A developmental stretch-and-fill process that optimises dendritic wiring

Ryan Rahy[1,2,3,,10,], Lothar Baltruschat[4,5,10], André Ferreira Castro[1,6,], Peter Jedlicka[2,3,7], Gaia Tavosanis[4,5,8,*,], and Hermann Cuntz[1,2,3,4,9,11,**,]

[1]Ernst Strüngmann Institute (ESI) for Neuroscience in cooperation with the Max Planck Society, Frankfurt-am-Main, 60528, Germany

[2]Computer-Based Modelling in the field of 3R Animal Protection, Faculty of Medicine, Justus Liebig University Giessen, Giessen, Germany

[3]Translational Neuroscience Network Giessen, Germany

[4]Max-Planck Institute of Biological Intelligence, 82152 Martinsried, Germany

[5]Center for Neurodegenerative Diseases (DZNE), Bonn, Germany

[6]Present address: School of Life Sciences, Technical University of Munich, Freising 85354, Germany

[7]Dr. Senckenberg Anatomy, Neuroscience Center, Goethe University, Frankfurt Main, Germany

[8]Present address: Institute of Developmental Biology / BioII, RWTH Aachen University, Germany

[9]Frankfurt Institute for Advanced Studies, Frankfurt-am-Main, 60438, Germany

[10]These authors contributed equally

[11]Lead contact

[*]Correspondence: gaia@devbiol.rwth-aachen.de

[**]Correspondence: hermann.neuro@gmail.com

Keywords

Dendrite growth, Space filling, Minimum spanning tree, Model, da neurons, *Drosophila*

In brief

We derive a detailed mathematical model that describes long-term time-lapse data of growing dendrites; it optimises total wiring and space-filling.

Highlights

- Fly neurons stretch and fill a given target area with precise scaling relations.

- We observe a sequence of two growth strategies.

- Each growth type implements optimal wiring which leads to optimal space filling.

- A model combining these programs captures the development of dendritic structures.

Summary

Circuit connectivity and computation depend on how dendrites branch and occupy space within neural tissue. While optimal wiring principles have long been known to constrain dendritic morphology and their scaling behaviour, the growth dynamics that produce such optimised structures remain unclear. Leveraging structural imaging across development, we identify two complementary growth strategies – inside-out versus outside-in – that together generate mature dendritic arbours. We formalise these dynamics in a mathematical model that captures the two growth modes and show that their interplay yields wiring-efficient, space-filling morphologies and class-specific developmental trajectories across species. This framework provides an algorithmic account of how local branching dynamics give rise to globally optimised architectures. By linking dendritic growth rules to functional design constraints, our theory offers a unifying description of dendritic differentiation and a basis for understanding how coverage and connectivity emerge during neural circuit formation.

Introduction   14

Dendritic branching defines the structural and functional identity of neurons by shaping   15
how they receive, integrate, and transmit information within neural circuits (Jan and Jan,   16
2003; London and Häusser, 2005; Poirazi and Papoutsi, 2020). Through complex arborisation   17
patterns, dendrites provide the receptive surface that underlies the brain's high degree of 18
connectivity, ultimately defining the connectome (Winding et al., 2023; Dorkenwald et al.,   19
2024; Schneider-Mizell et al., 2025; Piazza et al., 2025; Fritz et al., 2025). Class-type specific   20
morphologies emerge from coordinated influences of genetic programs and growth-promoting   21
signals that activate local interactions guiding branches into their appropriate spatial domains   22
(Parrish et al., 2007; Lefebvre et al., 2015; Barabási et al., 2025). Although these molecular and   23
cellular regulatory processes have been extensively characterised, the branching dynamics   24
that govern dendritic organisation remain poorly understood.   25

While several computational approaches can generate dendrite-like branching structures   26
(Ascoli et al., 2001; Luczak, 2006; Koene et al., 2009; Cuntz, 2015; Kanari et al., 2022; Kirchner et   27
al., 2025), morphological models based on optimal wiring (Cuntz et al., 2007, 2008; Budd et al.,   28
2010; Cuntz et al., 2010) are particularly well suited for studying dendrites and axons in their   29
role to implement circuit connectivity. Models derived from minimum spanning trees (MSTs)   30
(Kruskal, 1956; Prim, 1957) successfully reproduce key geometric and topological features of   31
dendritic architecture and provide a normative explanation for conserved scaling relationships   32
among neuronal size, wiring length, and connectivity (Cuntz et al., 2012). However, as MST-   33
based models describe what optimal wiring looks like, they leave open the question of how   34
branching dynamics in neurons achieve this during development.   35

Recent advances in time-lapse imaging now make it possible to observe dendritic growth as   36
a continuous process rather than a sequence of static morphologies (Sin et al., 2002; Lee et   37
al., 2013; Hogg et al., 2021). Quantitative live imaging captures branch extension, retraction,   38
and remodelling with sufficient temporal resolution to test computational models of arbour   39
formation (Mizrahi, 2007; Lee et al., 2013; Chalmers et al., 2016; Gonçalves et al., 2016; Radic   40
et al., 2017; Ferreira Castro et al., 2020; Stürner et al., 2022; Palavalli et al., 2021; Shree et al.,   41
2022; Rigaux et al., 2025). These studies reveal that class-specific dendritic arbours emerge   42
through sequential growth stages combining stochastic branch exploration with deterministic   43
refinement (Ferreira Castro et al., 2020; Palavalli et al., 2021). Despite these advances, imaging   44

alone cannot explain how local growth dynamics give rise to globally optimised architectures. 45

To uncover the growth rules that generate optimally wired dendritic morphologies, we 46
focused on class IV dendritic arborisation (da) neurons of *Drosophila* larvae, a well-established 47
model for dendrite morphogenesis (Jan and Jan, 2010; Grueber et al., 2002). These sensory 48
neurons form a highly branched, space-filling meshwork that tiles the larval body wall 49
through local self-avoidance and interactions with neighbouring arbours (Sugimura et al., 50
2007; Matthews et al., 2007; Soba et al., 2007; Shimono et al., 2010; Shree et al., 2022). Their 51
planar geometry, stereotyped architecture, and genetic accessibility enable direct quantification 52
of branch dynamics during growth, making them an ideal system for extracting general rules 53
of dendritic self-organisation. 54

To formalise dendritic maturation, we develop here a developmental morphological model 55
grounded in experimentally-observed growth dynamics after tracking all branches of class IV 56
da neurons (c4da) during development. The model reconstructs developmental trajectories 57
from quantitative morphometrics, incorporating two complementary growth programs: an 58
initial inside-out expansion driven by tip extension and a secondary outside-in phase that 59
fills space through new branch formation and stretching. This dual-process framework 60
reproduces class-specific morphologies and explains how neurons achieve optimal wiring 61
while occupying available territory. Validation of our model based on morphological statistics 62
using synaptic-resolution, densely reconstructed datasets confirms that the local growth 63
dynamics that we describe generalise across two- and three-dimensional environments and 64
across species. Together, these results establish a unified generative model of dendritic 65
development that links local growth rules to global wiring optimisation. 66

Results

Scaling behaviour of C4da neurons throughout larval development.

In order to dissect how optimal wiring is implemented in the growth of individual cells, we used live imaging to obtain time-lapse reconstructions of developing dendrites of the same C4da neurons throughout larval development (embryonic stage E – *red*, and instar larval stages L1 – *green*, L2 – *blue*, L3 – *black*, **Figure 1A**, see **Methods** "Time-lapse image acquisition"). Remarkably, large parts of the dendritic architecture were conserved over all stages of development while the overall shape was stretched, increasing the size in all directions (**Figures 1B and C**). Most importantly, individual dendritic branches from previous stages were easily identifiable and new branches grew in the space that became available through the extensive epidermal stretching.

We looked at different dendritic morphological parameters throughout the E–L3 larval stages to better quantify the geometrical aspects of the growth process. All branching statistics scaled up extensively with the developmental stages (**Figure 1D**) but were essentially invariant of the developmental stage once the differences in overall sizes were normalised out (**Figure 1E**), suggesting a general growth process conserving dendritic features.

The surface area S covered by the dendritic trees increased by a factor of $62\times$ between embryo (E) and third instar larva (L3) stages. The relation between the surface area S and the total dendrite cable length L during the L1–L3 stages was consistent with a strict power relation of exponent 0.78 ($L \sim S^{0.78}$; $R^2 = 0.97$). This implied that branches filled up newly available space more than in a simple stretching process ($L \sim S^{1/2}$, increasing the size without adding branches) but without conserving cable density ($L \sim S$, a linear relation with an exponent of 1) throughout development (**Figure 1F**, *left panel*). Similarly, the number of branch points B scaled with the surface area with an exponent of 0.66 ($B \sim S^{0.66}$; $R^2 = 0.84$), reflecting a branching process falling somewhere between a pure stretching process with no new branches ($B \sim S^0$, i.e. B does not change with S) and a process conserving branch point density ($B \sim S$; **Figure 1F**, *right panel*). The embryonic stage, on the other hand, showed markedly different scaling relations between these morphological parameters compared to L1–L3, suggesting that C4da neurons might exhibit a different initial growth process (**Figure 1F**), in line with

previous observations (Parrish et al., 2009). Overall, the analyses suggest a simple progression of growth with strong quantitative constraints for our mathematical theory.

Figure 1. Long-term time-lapse imaging of C4da neuron development.

A, Diagrams illustrating the imaging procedure throughout *Drosophila* larva developmental stages: embryo (E), larval instar (L) 1, 2 and 3. The larvae were kept in culture medium between the imaging steps. Times shown are AEL (after egg laying). **B**, Maximum intensity projections of image stacks from the same two neighbouring da dendrites (stars) for all four developmental stages. **C**, Digital reconstructions of the two neighbouring dendrites from **B** with small diameters clamped at $2\mu m$ for better visualisation. Triangular markers in L3 show axons, which are ignored in the rest of the study. **D–E**, Average branching statistics of reconstructions at the four developmental stages in absolute (**D**) and normalised (**E**) values. In all panels, the colours for the developmental stages are E (red), L1 (green), L2 (blue), L3 (black). **F**, Scaling behaviour of total length L and number of branch points B against the total surface S that the dendritic trees span. (See next page.)

Figure 1. (continued) Dashed lines show the best power law fits (linear fit in the log-log domain) for the data points from L1–L3 with respective power and r^2 values.

Dendritic developmental stretch-and-fill.   98

Beyond the quantification of the mathematical scaling relations, we dissected the C4da devel-   99
opmental growth process, leveraging our detailed time-lapse data to compare each time point   100
to the next. We tracked branch and termination points that were conserved in consecutive   101
reconstructions using a manual registration procedure (**Figure 2A**, see **Methods** "Time-lapse   102
analysis"). The analysis unveiled a surprisingly isometric stretching (**Figure 2B**) with a tight   103
linear slope between the stretching vector and the distance of points from the dendrite root   104
(**Figure 2C**). Interestingly, the overall origin of stretching did not lay precisely in the dendritic   105
root location (**Figure 2B**), consistent with a possible impact of superficial epithelium extension   106
on dendritic stretching during development (Jiang et al., 2014). The results from our manual   107
registration were all confirmed by an automated tracking of branches (see supplementary   108
Figure ??, Hermila et al., 2026, in preparation).   109

Having quantified the overall stretching of the dendritic spanning field, we scaled all the   110
dendritic reconstructions to the same surface area to focus on the changes inside. This was   111
done after applying a combination of small global alignment steps (a rotation with a consecu-   112
tive stretching in both spatial dimensions) to superimpose the obvious conserved branching   113
patterns. Our time-lapse registrations enabled us to then characterise the precise changes in   114
the branches that were not conserved. We quantified the branch dynamics and separated   115
them into: (1) retraction (branch elimination), (2) tip growth (branch elongation/shrinkage),   116
and (3) internal filling (new branch formation) (**Figure 2D**). Quantifying each of these types of   117
changes revealed that the amount of branch retraction was low and decreased throughout   118
development. In addition to that, while E to L1 showed more tip growth than internal filling,   119
this trend reversed for L1 to L3 that showed more internal filling than tip growth, once again   120
indicating a difference in developmental program between the embryonic and larval stages   121
(**Figure 2E**).   122

Figure 2. C4da neuron developmental stretching.
A, Time-lapse data analysis: Branch and termination points (coloured dots) in time 1 (*left*, here L2) are registered to the corresponding points in time 2 (*right*, here L3, same colours). The arrows on the left depict the movement of the respective topological points and illustrate the stretching of the entire structure between two acquisitions. **B**, Overall analysis for all pairs ($N = 47$). Arrows (every $25\mu m$) are binned averages only shown where data for at least 5 pairs was available. Dendrites were centred on their root (black dot). **C**, Analysis of stretching by comparing the arrow lengths (*top*, one dot per topological point with same colours as in **A**; *bottom*, topological points collected from all pairs with colours indicating larval instar at time 2 and their distance to the root). Greyed out points represent tips, linear fits (dashed orange lines) determine the slopes. **D–E**, Retraction (orange), tip growth (cyan), and internal filling (purple) processes observed in C4da cells, illustrated in **D** and quantified across all structures and phases in **E**. **F**, *Left*, diagram showing how θ is calculated as the distance one has to move away from the tree to cover $\sim 90\%$ of the spanning field of the dendrite, an inverse of space-filling-ness for neurons. *Right*, calculation of θ for all stages. Dashed orange line shows a smoothed Eilers estimate of the data. In panels **D–F**, all dendritic morphologies were scaled to cover a reference surface area of ($\frac{\sqrt{3}}{2} \cdot 1,000,000\mu m^2$, see **Figure ??**).

The considerable amount of internal filling and tip growth inside the dendritic spanning

area may be linked to the space-filling property of C4da neurons. We therefore introduce a [124] measure for the efficient coverage of available surface area S: θ, the maximum distance one [125] needs to move away from the dendrite to cover $\approx 90.69\%$ of the dendritic span. This measure [126] of space filling is inspired by considerations of circle packing on a hexagonal grid (**Figure** [127] **2F**, see **Methods** "Space-filling measure $1/\theta$" and supplementary **Figure ??**). Its inverse, $1/\theta$, [128] then quantifies how well the neuron fills the dendritic span (**Figure 2F**, *right*). In the current [129] case where all morphologies were stretched to cover the same surface area S, $1/\theta$ increased [130] steadily during larval instar growth L1–L3. [131]

Morphological model of dendrite development. [132]

Having quantified the simple progression of growth iterations and scaling of branching [133] statistics in the developmental dataset, we turn to morphological models of neurons to [134] analyse the growth through the lens of optimal wiring. We applied a minimum spanning [135] tree (MST) algorithm that optimises total cable length on our current C4da reconstructions [136] data (Cuntz et al., 2007; Prim, 1957, see Supplemental section "MST-based model reproducing [137] C4da development"). This showed that C4da dendrites followed optimal wiring principles [138] throughout development and were best matched with a trade-off parameter $bf = 0.225$ [139] between minimising total dendrite length L and path lengths PL along the tree from target [140] points toward the root ($wiring\ cost = L + bf \cdot PL$; **Figure ??**; see also Cuntz et al., 2010; Wen [141] and Chklovskii, 2008; Nanda et al., 2018). We next explored how the developmental process [142] leads to such optimally-wired trees from one stage to the next. [143]

Figure 3. Dendritic growth hypothesis.
Model incorporating inside-out growth (*top*) and outside-in growth (*bottom*), with new branches at every developmental step in *light blue* and *magenta* respectively compared to the conserved tree from previous developmental time points (in *grey*).

To this end, and based on our analysis of C4da development, we put forward a general $\quad$ 144
hypothesis of dendritic development, with two main processes in which dendrites spread $\quad$ 145
into the available space around them (**Figure 3**). Inside-out growth describes dendritic $\quad$ 146
changes that occur at the tips of the dendrites, with more tip retraction and elongation and $\quad$ 147
minimal interstitial branching; Outside-in growth describes a spreading pattern that fills out $\quad$ 148
the available spanning field with longer, relatively sparser primary dendrites before filling $\quad$ 149
the gaps in with interstitial branches. The resulting dendritic spreading pattern of a given $\quad$ 150
structure can involve both growth processes to varying degrees. $\quad$ 151

We formalised this hypothesis in an algorithm that builds on optimal wiring principles and $\quad$ 152
can generate different developmental trajectories. Developmental dynamics are determined $\quad$ 153
by a new parameter, the spreading pattern sp, which controls the timing of emergence of $\quad$ 154
different outgrowths (**Figure 4A**, *bottom panels*). The resulting temporal patterns can fall $\quad$ 155
anywhere on the continuous spectrum between the two growth types, with lower sp values $\quad$ 156
favouring inside-out growth and higher sp values favouring outside-in growth. Similar to $\quad$ 157
the aforementioned MST-based model, the branching pattern of the modelled structures is $\quad$ 158
determined by a balancing factor bf (**Figure 4A**, *top panel*). The resulting branching structure $\quad$ 159
is independent of the sp value, as can be seen in the distinct centripetal biases of the root angle $\quad$ 160
distributions (calculated as the distribution of angles between each dendritic segment and $\quad$ 161
its direct path to the soma Bird and Cuntz, 2019, **Figure 4B**, *bottom panel insets*) of the model $\quad$ 162
structures; this suggests consistent wiring optimality across sp values. In addition to that, the $\quad$ 163
scaling of the total dendritic length L with space filling $1/\theta$ was conserved throughout the $\quad$ 164
parameter space of our model (**Figure 4C**). $\quad$ 165

In order to generate realistic structures, an additional parameter k injects noise into the growth $\quad$ 166
process, subtly affecting the resulting structures along with their wiring cost optimality $\quad$ 167
and growth trajectories (**Figure ??**). **Figure ??** compares the growth dynamics and scaling $\quad$ 168
behaviours of these optimal-wiring-based morphological models. $\quad$ 169

Figure 4. growth_tree() algorithm is based on optimal wiring principles enabling distinct developmental trajectories along the continuous spectrum between the inside-out and outside-in growth type.
A, Effect of the main parameters on the output of the proposed algorithm modelling dendritic growth. *Top row*, Effect of changes in balancing factor (bf) for a set $sp = 0$. *Bottom panels*, Effects of changing the spreading pattern parameter sp for $bf = 0.225$ (*left*) and $bf = 0.9$ (*right*) on the developmental iterations of growth. **B**, Root angle distributions of the final morphologies from the bottom panels of **A**. **C**, Total length L vs space filling $1/\theta$ (see **Figure 2F**) during the developmental trajectories shown in the bottom panels of **A**.

Morphological model of C4da neurons throughout development

With this growth model, we simulated the complete *de novo* development of C4da dendrites in two steps in order to capture the change in growth program described earlier (**Figure 5A**). We first modelled the embryonic structure with a low $sp = 0.3$ then continued development until L3 with a higher $sp = 1$ to simulate the change from the inside-out growth program to outside-in. The bf parameter was unchanged from the wiring-optimal MST models (**Figure ??A**, *left*), and remained consistent throughout development. The synthesised L1 and L2 structures were selected as the intermediate model structures that length-matched the reconstructed structures.

Figure 5. Model of C4da development.
A, A two-stage model of the developmental trajectory of C4da cells (*top*), compared to the reconstructed structures scaled appropriately (*below*). In both modelling steps $bf = 0.225$, while sp was changed from 0.3 to 1 between the E and L1 stages. The shaded area behind each neuron shows the real dendritic span used to grow these neurons. Synthesised structures are coloured according to the sp value, following the colour scheme in **Figure 4A**. **B**, Scatter plots showing the scaling behaviour of real dendrites (in grey) compared to model dendrites (in pink), with the model trajectory from **A** shown in purple. Note that only structures for which all developmental phases were available were used.

Applying this model to all real C4da structures reproduced the scaling behaviour that real 179
dendrites underwent while optimising wiring and space filling throughout development 180
(**Figure 5B**). Modelling C4da cells in a single step without the switch in growth programs led 181
to structures with imbalanced branch lengths (see Supplemental section **??**). 182

Applying the growth model to other cell types 183

Our developmental growth process model reproduced the morphologies and the space-filling 184
characteristics of other example dendrites, both planar (fly C3da sensory neurons and lobula 185
plate tangential cells) and three-dimensional (rat dentate gyrus granule cells, fly second 186
order nociceptive neurons, and rat cortical layer 5 pyramidal cells), with bf and sp parameter 187
combinations that spanned different parts of the parameter space (**Figure 6**). The structures 188
predicted for these neurons were confirmed by statistical comparison to reconstructed neurons 189
(**Figure ??**). 190

Similar to C4da cells in **Figure 5A**, modelling the entire development of some other neuron 191
types required multiple steps (**Figurs 6A**). This included dendrites with growths having 192
differential spatial organisation, such as short terminal branch growth in C3da cells, and 193
growths with differential timing, such as oblique branch growth in L5 cortical pyramidal 194
neurons. 195

In order to further validate the model developmental trajectories, we compared the intermedi- 196
ate structures generated by our model to the earlier stages of the biological counterparts. This 197
was done using L1- and L3-stage structures of the same neurons obtained from two electron 198
microscopy datasets of the same fly nociceptive circuit (Gerhard et al., 2017). In this paradigm, 199
we first modelled the dendritic structures corresponding to L3 neurons, then analysed the 200
intermediate model structures to evaluate how well our trajectories captured the properties of 201
the reconstructed L1 structures. By length-matching the synthesised developmental steps to 202
the reconstructed L1 structures (**Figures 6**B, *first row*), we show that our model was able to cap- 203
ture both the individual structures and the trajectories that give rise to them. To evaluate the 204
overall performance across all models, we put together the artificial dendrite models at their 205
original locations in the connectome data and re-inserted the original axons and soma-linking 206
cables pertaining to the structures (**Figures 6**B, *second row*). Other than the expected difference 207
in tissue packing and stretching at the L1 stage, the identified synthesised structures matched 208

the real structures (**Figure S6**).

Figure 6. Modelling the development of additional cell types.

A, Growth model applied to a variety of neurons: *Orange*, Fly tangential cell; *pink*, Rat granule cell; *green*, Rat cortical pyramidal cell; *teal*, Fly C3da neuron. Synthesised structures from the model are shown in black and real cells in colour, with the grey shaded area marking the spanning area (derived from the reconstructed trees) in which these dendrites were grown. **B**, Intermediate structure prediction paradigm: *Top*, an example growth trajectory of a fly Basin A09a neuron (*lila*) reconstructed from EM data at larva instar stages L1 and L3. Synthesised L3 structure and its intermediate growth structures (*black*). *Bottom*, connectomes of all nociceptive neurons with the real axons and linking cables (in *black*) and modelled (large images) and real (insets) dendritic structures. Each colour represents a different dendrite.

Discussion

A central challenge in understanding neuronal development is to explain how complex dendritic arbours grow over time while maintaining functional organisation and wiring efficiency. The goal of this study was to obtain detailed time-lapse recordings of class IV dendritic arborisation (C4da) morphologies and comprehensively analyse developmental dendritic branching changes. We present a morphological model that captures the dendritic development constrained by that data as well as optimal wiring principles, and formulate a general hypothesis on dendritic growth.

From these analyses, several general principles of dendrite development emerge: (1) As the larva grows, C4da dendrites stretch with a remarkable conservation of branching structure and grow into new space as it becomes available; (2) Two alternative growth programs separated by what looks like a phase transition were observed throughout C4da development; (3) The two programs (inside-out vs outside-in) are required in the model to reproduce the data; (4) Both programs seem to guarantee optimal wiring throughout development; (5) Optimal wiring algorithms in general seem to guarantee that the dendritic span is optimally space-filled for a given dendritic cable length; (6) The growth model derived from C4da neurons applies widely to other neurons and explains existing data from cellular neuroanatomy.

From our *in vivo* time-lapse data we showed that the initial embryonic phase can be described as predominantly driven by external tip growth, consistent with previous observations (Shree et al., 2022). In contrast, the growth of the remaining developmental stages was better described by a regular and isometric stretching combined with internal in-filling (**Figure 2D,E**). This interpretation, of which we are confident due to the strong structural conservation between time points (see also our extensive appendix), is consistent with previous work showing a reorganisation during C4da development. One study found exocytotic machinery related to dendritic expansion shifting from being well-distributed throughout the entire dendrite during embryogenesis to becoming predominantly localised to primary (but not terminal) dendrites during larval development (Peng et al., 2015). Another study further showed that during development, C4da neurons first fill the empty space around them during embryonic development until they reach the hemisegment boundaries, then stretch loosely with the epithelium (Parrish et al., 2009). These changes in cellular mechanisms and extracellular contexts could reveal possible mechanistic or causal processes underlying the

change in dendritic spreading patterns that we report here. Relatedly, the timing of similar 241
changes in developmental programs coinciding with hatching has been observed in other 242
systems such as *C. elegans* (Nicosia et al., 2013). 243

Based on the C4da developmental trajectory, we derive a morphological model containing 244
two different growth programs. The inside-out program describes a growth type that is 245
driven by tip branching and growth (Hannezo et al., 2017) while the outside-in program 246
corresponds to more longer-ranging outgrowths followed by interstitial branching (Uçar 247
et al., 2021). This model introduces a new spreading pattern parameter sp that determines 248
the relative contribution of each growth type to development, which in turn determines the 249
temporal dynamics of dendritic outgrowth while preserving other dendritic features. We 250
were thus able to model a range of cell-type-specific differentiation trajectories for granule 251
cells (*c.f.* Gonçalves et al., 2016; Radic et al., 2017, *in vivo* and *in vitro*) and for fly nociceptive 252
neurons (Gerhard et al., 2017, *in vivo*). Our models also make predictions for structures and 253
trajectories with temporally- or spatially-specific growth (which could be driven by changes in 254
intrinsic or extrinsic growth cues) using multiple growth types. This includes the temporally 255
distinct multi-phasic growth of C4da cells which we dissect here extensively; the spatially 256
distinct short terminal branches (STBs) of class III cells, which have previously been shown to 257
have different underlying microtubule structures compared to the rest of the cell (Stürner et 258
al., 2022); and the branch compartmentalisation in pyramidal cells, where apical, basal, and 259
oblique branch growths are differentially affected by different neurotrophins and other factors 260
(Romand et al., 2011). 261

A key question arising from the observed developmental transitions is whether changes in 262
growth programs disrupt wiring optimality. Despite the change in growth programs, C4da 263
dendrites followed optimal wiring principles across all developmental stages, as shown using 264
simple MST models (**Figure ??**, Nanda et al., 2018). Optimal wiring follows from optimal 265
allocation of resources by balancing costs for total circuit wiring and signal propagation 266
speeds (Chklovskii, 2004; Stepanyants and Chklovskii, 2005; Cuntz et al., 2007; Wen and 267
Chklovskii, 2008; Ferreira Castro and Cardona, 2025). At the cellular level, such constraints 268
have been shown to have several implications: (1) They lead to precise scaling relationships 269
between measures of branching structure and connectivity, suggesting that different dendritic 270
morphologies follow similar underlying growth rules (Cuntz et al., 2012); (2) They predict 271
branching angle distributions (Kim et al., 2012; Bird and Cuntz, 2019); and (3), As shown 272

in this work, they necessarily lead to optimal space filling regardless of target distribution, meaning that the dendritic span is optimally filled for a given optimal dendritic length (**Figure 4A, ??C**). Both scaling relationships and angle distributions were independent of the growth programs (**Figure 4**), and fitted the biological data throughout development.

Taken together, the distinct growth programs observed *in vivo* appear to preserve similar functional outcomes. This suggests a form of degeneracy in neuronal growth, whereby different developmental trajectories converge on comparable wiring solutions (Albantakis et al., 2024). Incidentally, the inside-out growth is reminiscent of the algorithmic iterations behind the greedy MST implementation of optimal wiring (Cuntz et al., 2007, 2010) (**Figure ??B**). The outside-in growth in addition is somewhat reminiscent of rapidly exploring random trees (rrt*) from graph theory which were previously linked to space filling (Kuffner and LaValle, 2011).

We have previously proposed our scaling equations from Cuntz et al. (2012) as a signature for optimal wiring. Importantly, recent work has shown that optimal wiring alone is not necessary for the power relationships in those equations (Ouyang et al., 2025, *i.e.* $L \sim N^{2/3}$ in 3D or $L \sim N^{1/2}$ in 2D). However, the particular scaling relations are a a consequence of the developmental implementation of optimal wiring and are not trivially achieved by stochastic growth. This contrast highlights that scaling laws cannot be interpreted independently of explicit growth models. Only through precise modelling can the relevant scaling relationships be identified, allowing fine-grained predictions of growth changes across development.

A limitation of the present study is that the growth rules and optimisation principles we identify are inferred from morphological dynamics rather than directly linked to underlying molecular or biophysical mechanisms. While our time-lapse imaging constrains the timing and nature of developmental transitions, the temporal resolution is insufficient to resolve the fine-scale dynamics of branch elongation, retraction, and local interactions that likely mediate these processes (Chalmers et al., 2016; He and Cline, 2011). In addition, our modelling framework is phenomenological and focuses on wiring and space-filling constraints, without explicitly incorporating intracellular signalling, cytoskeletal dynamics, or activity-dependent plasticity. Future work could address these limitations by linking model parameters to mutant phenotypes or elements of structural plasticity (Tavosanis, 2012; Yalgin et al., 2015), and by directly probing branch–branch repulsion and molecular cues that establish dendritic territories (Hattori et al., 2009; Matthews et al., 2007; Parrish et al., 2007).

Here, we provide a framework that links dendritic growth dynamics, space filling, and wiring optimality during development. Considerations of simultaneous multi-dendritic development and the spread of neurons into existing tissue would help make predictions about neuron packing (*e.g.* Braitenberg, 2001), neuron avoidance (Soba et al., 2007), gap filling (*e.g.* during repair, Song et al., 2012), and extracellular signal interactions, and would allow the exploration of how spatial constraints affect optimal wiring and space filling. Ultimately, our model links neurite outgrowth and space filling with circuit formation (Schröter et al., 2017; Helmstaedter, 2026) in neural networks of arbitrary complexity, enabling simulations based on well-understood principles rather than vast collections of neuronal reconstructions (Markram et al., 2015).

Acknowledgments

We are grateful to M. Schölvinck, P. Soba, T. Stürner and A. Ziegler for comments on an earlier version of the manuscript and to C. Lossnitzer for helpful comments on the computer code. This work was supported by a BMBF grant (No. 01GQ1406 — Bernstein Award 2013), a DFG grant (CU 217/2-1), and by the German Federal Institute for Risk Assessment (Grant Agreement Number 60-0102-01.P636). The authors declare to have no competing financial interests.

Author contributions

R.R., L.B., G.T. and H.C. designed the study. L.B. and G.T. designed and performed the experiments. R.R., A.F.C, and H.C. analysed the data, designed the growth model and performed all simulations. R.R., L.B., A.F.C, P.J., G.T., and H.C. wrote the paper.

Methods

Fly handling

To label Class IV da (C4da) neurons specifically, we used a fly strain that carried a fusion of a fraction of the enhancer of the *ppk* gene and eGFP ($w^{1118}; ; ppk - eGFP$) (Grueber et al., 2003). This strain expresses eGFP in C4da neurons from embryonic stage 16 throughout larval stages. We concentrated on the dorsal ddaC neuron (Grueber et al., 2002). For embryo collection, young fertilised females were placed in population cages, supplemented with apple agar plates maintained in an incubator at $25°C$ and 60% relative humidity. To obtain developmentally synchronised embryos the agar plate was supplemented with fresh yeast paste to promote laying of retained eggs and replaced after one hour. Females were then allowed to lay eggs for $30min$. The collected eggs were placed back in the incubator and allowed to develop for $18h$, when they were harvested, dechorionated with bleach for $3.5min$ and rinsed with water. Not all embryos were dechorionated by this gentle treatment, and the chorion was then removed mechanically with a brush.

Time-lapse image acquisition

We developed a novel method to image the living larvae over several consecutive days causing as little interference as possible. Briefly, after each imaging session individuals were kept at $25°C$ and 60% relative humidity in a separate $500\mu l$ Eppendorf tube filled with $200\mu l$ standard cornmeal-agar fly food. To regain the test subject the fly food was dissolved in water and the larvae were localised under a binocular microscope (**Figure 1A**). To keep animals immobilised for imaging, but alive through the imaging sessions, we utilised three different physical immobilisation techniques at different stages. **Embryo**: We created a grid with 6 chambers of approximately $1mm \times 0.5mm \times 0.5mm$ on an object slide using adhesive strip. Dechorionated embryos were placed in the grid, covered with halocarbon oil 700 (Sigma) and oriented in such a way that the dorsal class IV ddaC neurons faced towards the coverslip. **L1 and L2 larvae**: Images of L1 and L2 larvae were acquired under a custom-made object slide (Dimitrova et al., 2008). The metal custom object slide fitted a round coverslip onto which the larva was placed, oriented and covered with halocarbon oil. A plastic net on a round plastic support was then

placed on top of the coverslip and fixed with screws to the metal slide. Detailed information ₃₅₄
about the design is available upon request. **L3 larvae**: L3 larvae were physically immobilised ₃₅₅
in halocarbon oil in-between an objective slide and a coverslip. Double-sided adhesive tape ₃₅₆
served as spacer. On average two out of five individuals survived through all imaging steps. ₃₅₇
During this procedure, development until the pupal stage was extended from typical five ₃₅₈
days at $25°C$ to about seven days in our preparation. Given the increased development time in ₃₅₉
these conditions from $\sim 120h$ to $\sim 168h$, embryo images were taken $18h - 19h\,AEL$, L1 images ₃₆₀
at $42h\,AEL$, L2 at $90h\,AEL$ and L3 images $138h\,AEL$. The measured ddaC dendrite size at $90h$ ₃₆₁
and at $138h$ corresponded to our previous measures of L2 and feeding L3 animals in control ₃₆₂
conditions, respectively. Images were acquired with a Leica TCS SP2 Confocal Microscope ₃₆₃
(`http://www.leica-microsystems.com`), maintaining minimal laser intensity to reduce ₃₆₄
tissue damage. To avoid issues due to larval movement, we minimised imaging time per ₃₆₅
$z-$stack and thus chose a relatively large $z-$step of $2\mu m$ combined with a larger pinhole size ₃₆₆
$(85-150\mu m)$ to obtain a voxel depth of $1.7-2.2\mu m$. For embryos, we used a $40\times$ oil immersion ₃₆₇
objective and for all the other stages a $20\times$ oil immersion objective. ₃₆₈

Data analysis ₃₆₉

All data analysis was performed in *Matlab* (`www.mathworks.com`) using our own software ₃₇₀
package, the *TREES toolbox* (`www.treestoolbox.org`, version 1.15 and interim version, ₃₇₁
Cuntz et al., 2010, 2011). A number of new TREES toolbox functions were custom-made ₃₇₂
for this study and will be incorporated in the existing *TREES toolbox* with the publication ₃₇₃
of this work: `span_tree`, `scaleS_tree`, `scaleV_tree`, `theta_tree`, `compare_tree`, `ui_comp_tree`, ₃₇₄
`theta_mc_tree`, `growth_tree`. See below for details on the individual functions. In the follow- ₃₇₅
ing, function names are in monospaced font with *TREES toolbox* functions having a `_tree` ₃₇₆
suffix. All data will be made available on `www.NeuroMorpho.Org` and all new *TREES toolbox* ₃₇₇
functions will be made available on `www.treestoolbox.org`. The entire code and data will ₃₇₈
be archived online on Zenodo. ₃₇₉

Anatomical data of C4da neurons
380

Image stacks from the confocal microscope were imported in the *TREES toolbox* and manual   381
reconstructions of all dendrites were performed individually (total amount of neurons $N =$ 382
168) using the dedicated reconstruction user interface `cgui_tree`. During the reconstruction   383
process we determined (1) adequate internode distances i.e. spatial resolution at which to   384
resample (`resample_tree`) the dendritic structures, (2) parameters for a diameter taper toward   385
the root (`quaddiameter_tree`) that best reproduced the overall diameter values of the real cells,   386
(3) soma length and soma diameter values to best map soma diameters (`soma_tree`) *post hoc*   387
on the existing reconstructions, and (4) region codes for dendrite, soma and axon. Axon nodes   388
were discarded for the remaining analysis.   389

Classical branching statistics
390

A large palette of branching statistics were collected for each dendrite reconstruction separately   391
using simple combinations of *TREES toolbox* functions: (1) **Total dendrite length** L was the   392
sum of all internode lengths (`len_tree`), (2) **number of branch points** B were calculated   393
as the sum of all nodes that are branch points (`B_tree`), (3) covered **surface area** S was   394
calculated with a novel custom written *TREES toolbox* function `span_tree` that counts $1 - bins$ 395
after morphological closing with a disc of $4\mu m$ radius of a binary matrix with $1 - bins$ 396
at tree node locations after resampling the trees to $1\mu m$ internode distances – similarly to   397
previous work on fly lobula plate tangential cells (Cuntz et al., 2008), (4) **Sholl analysis**   398
(Bird and Cuntz, 2019) that calculates the number of intersections of a tree with a growing   399
circle centered on the dendrite root was done by using the dedicated *TREES toolbox* function   400
`sholl_tree`, (5) normalised Sholl analysis was done by normalising the Sholl radii to $6\times$ the   401
mean Euclidean distance in the tree (`eucl_tree`) and normalising the number of intersections   402
with the integral over the Sholl intersections diagram, (6) **branch order** distributions were   403
calculated by identifying how many nodes and the overall dendritic cable length of their   404
connected segments in the tree had a given branch order (`BO_tree`), (7) normalised branch   405
orders were normalised to the next integer of $2.5\times$ the mean branch order in any given   406
tree and the cable distribution was normalised to the total length of dendrite, (8) **branch**   407
length distributions were obtained by combining two *TREES toolbox* functions that calculate   408

the path length from each node to the root (`Pvec_tree`) and that dissect the trees into their individual branches (`dissect_tree`), (9) normalised branch lengths were normalised to $3\times$ the mean branch length in the tree and branch length frequency was normalised to the integral over the distribution, (10) **compression** value distributions were calculated as the ratio between Euclidean distances (`eucl_tree`) and the path distances (`Pvec_tree`) to the root for all nodes, (11) the **Strahler order** distributions were calculated as the percentages of dendrite length per given Strahler order (`strahler_tree`) (Vormberg et al., 2017), (12) **branch angle** distributions were obtained by calculating the angles in the branching plane of each branch point (`angleB_tree`). Overall, we chose not to normalise to the maximum value but rather to a multiple of the mean because maximum values are more sensitive to noise.

Scaling of real trees

In order to compare real trees with a simple maximal space filling example, all reconstructions were scaled to the same surface area S as a reference diamond ($h \cdot 1,000,000 \mu m^2$, see supplementary **Figure ??** and supplementary section **'Mathematical calculations for maximal space filling')** by scaling the trees twice while measuring S (`span_tree`, see above) to make sure that target S is reached precisely in all cases (new *TREES toolbox* function `scaleS_tree`). Real trees in **Figures 2D-F, 5, supplementary ??** were scaled in such a manner.

Space-filling measure $1/\theta$

A custom-written new *TREES toolbox* function `theta_tree` was used to calculate how space filling a tree is. θ is the distance one has to move away from the tree to cover 90.69% of the spanning surface of the tree, in analogy to our maximal space filling mathematical framework derived from packing circles, see **Supplementary text**. Similar to the surface area S calculation, we calculated a binary matrix with $1 - bins$ after morphological closing with a disc of $4\mu m$ radius and used the *Matlab* function `bwdist` to calculate the distances from each bin within this spanning area to the closest bin occupied by the dendritic tree. Similar 'box-counting' methods have appeared in the literature (Wen et al., 2009; Shree et al., 2022), but the approach here allows the calculation of both single values of θ and the entire detailed distance distributions in histograms. The iterative growth process followed a slightly different but equivalent approach

using a Monte Carlo method since for 3D cases the binary matrices would have been very [437]
large (see section **'Iterative growth model'**). [438]

Time-lapse analysis [439]

Before registration of the branch and termination points, the morphologies of identified [440]
pairs of dendrites (the same cell in two consecutive time steps, $N = 47$ pairs) were rotated, [441]
translated and scaled separately in both dimensions to maximise the overlap of dendritic [442]
branches. Branch and termination points were then registered first manually and second [443]
using a fully-automated function (new *TREES toolbox* function `compare_tree`, see **Figure ??**). [444]

For the stretching analysis, the individual scaling steps were ignored and the vectors were [445]
drawn between branch and termination points of time points 1 and 2 for all pairs. [446]

We took advantage of both manual and automated of registration sets to track and and quantify [447]
the branching changes across the structures. Considering the branches formed between the [448]
soma and the tracked branching and termination points to be common between the structures [449]
in a given pair, we were able to isolate retracted branches (branches that appear in structure 1 [450]
and not in 2) and newly formed branches (branches that appear in structure 2 and not 1 and [451]
start from non-termination points). Branch elongation and shrinkage processes were identified [452]
as (1) the growth of untracked branches (similar to newly-found branches) from termination [453]
points and (2) the movement of termination points tracked with `compare_tree` from one [454]
time point to the next (since the fully-automated method stringently followed termination [455]
points, see **Figure ??**E). [456]

Dendritic minimum spanning trees [457]

Optimal wiring was implemented as previously described (Cuntz et al., 2007) with a minimum [458]
spanning tree algorithm connecting a given set of targets to minimise (1) total cable length [459]
and (2) path lengths from any target along the tree towards the root of the tree. This second [460]
cost was weighted by the balancing factor bf, the only parameter that the wiring algorithm [461]
takes: $wiring\ cost = L + bf \cdot PL$. This method was successfully used previously (Cuntz et al., [462]
2008, 2010) and all algorithms are available in the *TREES toolbox* (`www.treestoolbox.org`, [463]

`MST_tree`).   464

Targets were distributed uniformly in the surface area of each real dendrite, one at a time   465
(**Figure ??A**), and connected them with the minimum spanning tree algorithm. We generated   466
minimum spanning trees connecting as many targets as required to reproduce the number   467
of branch points B in the original tree and matched the resulting synthetic tree surface area   468
S each using the same scaling procedure used for real reconstructions (see above). We then   469
scanned the parameter bf of the model between 0 and 1 in steps of 0.025. An error function   470
comparing mean total length, branch order, compression and path length between synthetic   471
trees and their real counterparts yielded an optimal bf of 0.225. The corresponding synthetic   472
trees were then used with this bf value (**Figure 4D**). For visual purposes, all synthetic trees 473
were smoothed (`smooth_tree`) and a slight jitter was added (`jitter_tree`) that did not affect   474
the branching statistics.   475

Iterative growth model   476

The iterative growth model (new TREES function `growth_tree`) is an extension of the original   477
minimum spanning tree model implementing optimal wiring. It was fit to cover the two   478
developmental programs uncovered from reconstructions of dendrite development, and   479
the span of developmental trajectories between them. Given a surface area or volume in   480
which to grow a dendrite, this span is probed with randomly placed target points, with   481
every iteration of the algorithm growing towards one of these points to simulate a sequential   482
growth process. The number of target points is chosen by finding the lowest number that   483
allows the algorithm to reach the set termination condition before it runs out of points to   484
connect to. This is done by starting with (roughly) twice the number of branching points in   485
the reference structure, and decreasing that number until the algorithm stops prematurely.   486
Selecting the target point to grow to and where to connect it on the tree is based on optimising   487
two separate functions: minimising wiring cost, as described in the previous section, and   488
maximising space filling, which is based on how far a given target point is from the existing   489
tree structure. These two optimisation procedures are combined in a single scoring function   490
used to select the most optimal tree node to target point combination in a given iteration: 491
$score = ((1 - sp) * wiring\ cost) - (sp * node\ distance)$, where the scores pertaining to each   492
function are normalised to 1 and balanced based on the value of the spreading pattern (sp)   493

parameter. At $sp = 0$, the algorithm prioritises connections that minimise the wiring cost, while at $sp = 1$ the algorithm connects preferentially to points that are further away from the tree (while still minimising wiring costs as much as possible). An additional parameter k, which is internally saturated using $tanh(4k)$, adds uniform random noise to the scoring function. Following these parameters, the tree node – target point combination with the lowest calculated score is chosen and added to the tree. The new branch is then subdivided into $1\mu m$ segments and jittered slightly before the next iteration starts. The growth process stops when a termination condition is reached, which can either be running out of target points or stopping when a target total length L or number of termination points T is reached.

Synthesised morphologies of other planar and 3D dendrites

Five synthesised dendrites were grown from scratch using the growth_tree iterative growth model. Synthesised **Fly lobula plate tangential cells** (Cuntz et al., 2008) were obtained using $bf = 0.6$, $sp = 0.8$, $k = 0.4$, $radius = 800\mu m$; **Rat dentate gyrus granule cells** (Rihn and Claiborne, 1990) were obtained using $bf = 0.975$, $sp = 0.3$, $k = 0$, $radius = 400\mu m$; **Rat cortical layer 5 pyramidal cells** (Wittner and Miles, 2007) were obtained using $bf = 0.9$, $sp = 1$ $k = 0.5$, $radius = 1,000\mu m$; **Fly CIII dendritic arborisation sensory neurons** (Wittner and Miles, 2007) were obtained using $bf = 0.9$, $sp = 1$ $k = 0.5$, $radius = 1,000\mu m$.

Intermediate structure prediction

To validate our synthesised developmental trajectories, we tested our model's ability to predict the correct intermediate structures by comparing our model's iterations to earlier phases of real neurons. Two electron microscopy datasets containing overlapping neurons were used here: one from the fly L1 stage taken from the A1 segment, and another from the L3 stage/A3 segment. Six neuron types were isolated (A10a, A09a, A09c, A09l, A02m, A02n) from both datasets, and preprocessed by separating the dendrites, axons, and somatic linker cables. This paradigm then involved modeling each target L3 dendrite and combing through the iterations leading to it to find the L1-equivalent structures, which were defined as those closest length-wise to the real L1 dendrites.

Similar parameters were used to generate the structures for all six cell types: **A10a neurons** were obtained using $bf = 0.2$, $sp = 0$, $k = 0.5$, $radius = 80\mu m$; **A09a neurons** were obtained using $bf = 0.1$, $sp = 0$, $k = 0.6$, $radius = 80\mu m$; **A09c neurons** were obtained using $bf = 0.1$, $sp = 0.1$, $k = 0.6$, $radius = 80\mu m$; **A09l neurons** were obtained using $bf = 0.25$, $sp = 1$, $k = 0.6$, $radius = 200\mu m$; **A02m neurons** were obtained using $bf = 0.1$, $sp = 0$, $k = 0.5$, $radius = 120\mu m$; **A02n neurons** were obtained using $bf = 0.05$, $sp = 0.2$, $k = 0.6$, $radius = 80\mu m$.

The connectome model graphics in **Figure 6B** were generated by concatenating the modeled dendritic structures to the previously discarded axon and linker cable segments. In order to somewhat account for tissue stretching between the two stages, L1 structures were scaled down according to the overall difference in size between the real connectomes (using the horizontally protruding somata as reference).

References

Albantakis L, Bernard C, Brenner N, Marder E, Narayanan R (2024) The brain's best kept secret is its degenerate structure. *Journal of Neuroscience* 44:e1339242024.

Ascoli GA, Krichmar JL, Scorcioni R, Nasuto SJ, Senft SL (2001) Computer generation and quantitative morphometric analysis of virtual neurons. *Anatomy and Embryology* 204:283–301.

Barabási DL, Ferreira Castro A, Engert F (2025) Three systems of circuit formation: assembly, updating and tuning. *Nature Reviews Neuroscience* 26:232–243.

Bird AD, Cuntz H (2019) Dissecting Sholl analysis into its functional components. *Cell Reports* 27:3081–3096.e5.

Braitenberg V (2001) Brain size and number of neurons: an exercise in synthetic neuroanatomy. *Journal of Computational Neuroscience* 10:71–77.

Budd JML, Kovács K, Ferecskó AS, Buzás P, Eysel UT, Kisvárday ZF (2010) Neocortical axon arbors trade-off material and conduction delay conservation. *PLoS Computational Biology* 6:e1000711.

Chalmers K, Kita EM, Scott EK, Goodhill GJ (2016) Quantitative analysis of axonal branch dynamics in the developing nervous system. *PLOS Computational Biology* 12:e1004813.

Chklovskii DB (2004) Synaptic connectivity and neuronal morphology: two sides of the same coin. *Neuron* 43:609–617.

Cuntz H (2015) Modeling dendrite shape In *Dendrites. Oxford University Press, New York*, pp. 487–504.

Cuntz H, Forstner F, Borst A, Häusser M (2010) One rule to grow them all: A general theory of neuronal branching and its practical application. *PLOS Computational Biology* 6:e1000877.

Cuntz H, Forstner F, Borst A, Häusser M (2011) The TREES toolbox—probing the basis of axonal and dendritic branching. *Neuroinformatics* 9:91–96.

Cuntz H, Forstner F, Haag J, Borst A (2008) The morphological identity of insect dendrites. *PLoS Computational Biology* 4:e1000251.

Cuntz H, Haag J, Forstner F, Segev I, Borst A (2007) Robust coding of flow-field parameters by axo-axonal gap junctions between fly visual interneurons. *PNAS* 104:10229–10233.

Cuntz H, Mathy A, Häusser M (2012) A scaling law derived from optimal dendritic wiring. *PNAS* 109:11014–11018.

Dimitrova S, Reissaus A, Tavosanis G (2008) Slit and Robo regulate dendrite branching and elongation of space-filling neurons in Drosophila. *Developmental Biology* 324:18–30.

Dorkenwald S, Matsliah A, Sterling AR, Schlegel P, Yu Sc, McKellar CE, Lin A, Costa M, Eichler K, Yin Y, Silversmith W, Schneider-Mizell C, Jordan CS, Brittain D, Halageri A, Kuehner K, Ogedengbe O, Morey R, Gager J, Kruk K, Perlman E, Yang R, Deutsch D, Bland D, Sorek M, Lu R, Macrina T, Lee K, Bae JA, Mu S, Nehoran B, Mitchell E, Popovych S, Wu J, Jia Z, Castro MA, Kemnitz N, Ih D, Bates AS, Eckstein N, Funke J, Collman F, Bock DD, Jefferis GSXE, Seung HS, Murthy M (2024) Neuronal wiring diagram of an adult brain. *Nature* 634:124–138.

Ferreira Castro A, Baltruschat L, Stürner T, Bahrami A, Jedlicka P, Tavosanis G, Cuntz H (2020) Achieving functional neuronal dendrite structure through sequential stochastic growth and retraction. *eLife* 9:e60920.

Ferreira Castro A, Cardona A (2025) Invariant synaptic density across species.   575

Fritz I, Wang F, Chirif R, Malakasis N, Gjorgjieva J, Ferreira Castro A (2025) Synaptic density   576
    and relative connectivity conservation maintain circuit stability across development.   577

Gerhard S, Andrade I, Fetter RD, Cardona A, Schneider-Mizell CM (2017) Conserved neural   578
    circuit structure across Drosophila larval development revealed by comparative connec-   579
    tomics. *eLife* 6:e29089.   580

Gonçalves JT, Bloyd CW, Shtrahman M, Johnston ST, Schafer ST, Parylak SL, Tran T, Chang   581
    T, Gage FH (2016) In vivo imaging of dendritic pruning in dentate granule cells. *Nature*   582
    Neuroscience 19:788–791.   583

Grueber WB, Jan LY, Jan YN (2002) Tiling of the Drosophila epidermis by multidendritic   584
    sensory neurons. *Development* 129:2867–2878.   585

Grueber WB, Ye B, Moore AW, Jan LY, Jan YN (2003) Dendrites of distinct classes of   586
    Drosophila sensory neurons show different capacities for homotypic repulsion. *Current*   587
    Biology 13:618–626.   588

Hannezo E, Scheele CLGJ, Moad M, Drogo N, Heer R, Sampogna RV, van Rheenen J, Simons   589
    BD (2017) A unifying theory of branching morphogenesis. *Cell* 171:242–255.   590

Hattori D, Chen Y, Matthews BJ, Salwinski L, Sabatti C, Grueber WB, Zipursky SL (2009)   591
    Robust discrimination between self and non-self neurites requires thousands of Dscam1   592
    isoforms. *Nature* 461:644–648.   593

He HY, Cline HT (2011) Diadem X: automated 4 dimensional analysis of morphological data.   594
    Neuroinformatics 9:107–112.   595

Helmstaedter M (2026) Synaptic-resolution connectomics: towards large brains and connec-   596
    tomic screening. *Nature Reviews Neuroscience* 27:101–120.   597

Hermila B, Rahy R, Bird AD, Jedlicka P, Sümbül U, Cuntz H (2026) A systematic comparison   598
    of dendritic trees. *in preparation* .   599

Hogg PW, Coleman P, Toth TD, Haas K, Dellazizzo Toth T, Haas K (2021) Quantifying   600
    neuronal structural changes over time using dynamic morphometrics. *Trends in Neuro-*   601
    sciences 45:106–119.   602

Jan YN, Jan LY (2003) The control of dendrite development. *Neuron* 40:229–242.   603

Jan YN, Jan LY (2010) Branching out: mechanisms of dendritic arborization. *Nature Reviews*   604
Neuroscience 11:316–328.   605

Jiang N, Soba P, Parker E, Kim CC, Parrish JZ (2014) The microRNA bantam regulates a   606
developmental transition in epithelial cells that restricts sensory dendrite growth. *Develop-*   607
ment 141:2657–2668.   608

Kanari L, Dictus H, Chalimourda A, Arnaudon A, Van Geit W, Coste B, Shillcock J, Hess   609
K, Markram H (2022) Computational synthesis of cortical dendritic morphologies. *Cell*   610
Reports 39:110586.   611

Kim Y, Sinclair R, Chindapol N, Kaandorp JA, De Schutter E (2012) Geometric theory pre-   612
dicts bifurcations in minimal wiring cost trees in biology are flat. *PLoS Computational*   613
Biology 8:e1002474.   614

Kirchner JH, Euler L, Fritz I, Castro F, Gjorgjieva J (2025) Dendritic growth and synap-   615
tic organization from activity-independent cues and local activity-dependent plasticity.   616
eLife 12:e87527.   617

Koene RA, Tijms B, van Hees P, Postma F, De Ridder A, Ramakers GJA, Van Pelt J, Van Ooyen   618
A (2009) NETMORPH: A framework for the stochastic generation of large scale neuronal   619
networks with realistic neuron morphologies. *Neuroinformatics* 7:195–210.   620

Kruskal JB (1956) On the shortest spanning subtree of a graph and the traveling salesman   621
problem. *Proceedings of the American Mathematical Society* 7:48–50.   622

Kuffner JJ, LaValle SM (2011) Space-filling trees: A new perspective on incremental   623
search for motion planning. *IEEE International Conference on Intelligent Robots and Sys-*   624
tems pp. 2199–2206.   625

Lee PC, He HY, Lin Cy, Ching Yt, Cline HT (2013) Computer aided alignment and quantitative   626
4D structural plasticity analysis of neurons. *Neuroinformatics* 11:249–257.   627

Lefebvre JL, Sanes JR, Kay JN (2015) Development of dendritic form and function. *Annual*   628
Review of Cell and Developmental Biology 31:741–777.   629

London M, Häusser M (2005) Dendritic computation. *Annual Review of Neuroscience* 28:503–532.   630

Luczak A (2006) Spatial embedding of neuronal trees modeled by diffusive growth. *Journal of Neuroscience Methods* 157:132–141.

Markram H, Muller E, Ramaswamy S, Reimann MW, Abdellah M, Sanchez CA, Ailamaki A, Alonso-Nanclares L, Antille N, Arsever S, Kahou GAA, Berger TK, Bilgili A, Buncic N, Chalimourda A, Chindemi G, Courcol JD, Delalondre F, Delattre V, Druckmann S, Dumusc R, Dynes J, Eilemann S, Gal E, Gevaert ME, Ghobril JP, Gidon A, Graham JW, Gupta A, Haenel V, Hay E, Heinis T, Hernando JB, Hines M, Kanari L, Keller D, Kenyon J, Khazen G, Kim Y, King JG, Kisvarday Z, Kumbhar P, Lasserre S, Le Bé JV, Magalhães BRC, Merchán-Pérez A, Meystre J, Morrice BR, Muller J, Muñoz-Céspedes A, Muralidhar S, Muthurasa K, Nachbaur D, Newton TH, Nolte M, Ovcharenko A, Palacios J, Pastor L, Perin R, Ranjan R, Riachi I, Rodríguez JR, Riquelme JL, Rössert C, Sfyrakis K, Shi Y, Shillcock JC, Silberberg G, Silva R, Tauheed F, Telefont M, Toledo-Rodriguez M, Tränkler T, Van Geit W, Díaz JV, Walker R, Wang Y, Zaninetta SM, DeFelipe J, Hill SL, Segev I, Schürmann F (2015) Reconstruction and simulation of neocortical microcircuitry. *Cell* 163:456–492.

Matthews BJ, Kim ME, Flanagan JJ, Hattori D, Clemens JC, Zipursky SL, Grueber WB (2007) Dendrite self-avoidance is controlled by Dscam. *Cell* 129:593–604.

Mizrahi A (2007) Dendritic development and plasticity of adult-born neurons in the mouse olfactory bulb. *Nature Neuroscience* 10:444–452.

Nanda S, Chen H, Das R, Bhattacharjee S, Cuntz H, Torben-Nielsen B, Peng H, Cox DN, De Schutter E, Ascoli GA (2018) Design and implementation of multi-signal and time-varying neural reconstructions. *Scientific Data* 5:170207.

Nicosia V, Vértes PE, Schafer WR, Latora V, Bullmore ET (2013) Phase transition in the economically modeled growth of a cellular nervous system. *PNAS* 110:7880–7885.

Ouyang X, Sutradhar S, Trottier O, Shree S, Yu Q, Tu Y, Howard J (2025) Neurons exploit stochastic growth to rapidly and economically build dense dendritic arbors. *Nature Communications* 16:5903.

Palavalli A, Tizón-Escamilla N, Rupprecht JF, Lecuit T (2021) Deterministic and stochastic rules of branching govern dendrite morphogenesis of sensory neurons. *Current Biology* 31:459–472.

Parrish JZ, Emoto K, Kim MD, Jan YN (2007) Mechanisms that regulate establishment, maintenance, and remodeling of dendritic fields. *Annual Review of Neuroscience* 30:399–423.

Parrish JZ, Xu P, Kim CC, Jan LY, Jan YN (2009) The microRNA bantam functions in epithelial cells to regulate scaling growth of dendrite arbors in drosophila sensory neurons. *Neuron* 63:788–802.

Peng Y, Lee J, Rowland K, Wen Y, Hua H, Carlson N, Lavania S, Parrish JZ, Kim MD (2015) Regulation of dendrite growth and maintenance by exocytosis. *Journal of Cell Science* 128:4279–4292.

Piazza B, Barabási DL, Ferreira Castro A, Menichetti G, Barabási AL (2025) Physical Network Constraints Define the Lognormal Architecture of the Brain's Connectome.

Poirazi P, Papoutsi A (2020) Illuminating dendritic function with computational models. *Nature Reviews Neuroscience* 21:303–321.

Prim RC (1957) Shortest connection networks and some generalizations. *Bell System Technical Journal* 36:1389–1401.

Radic T, Jungenitz T, Singer M, Beining M, Cuntz H, Vlachos A, Deller T, Schwarzacher SW (2017) Time-lapse imaging reveals highly dynamic structural maturation of postnatally born dentate granule cells in organotypic entorhino-hippocampal slice cultures. *Scientific Reports* 7:43724.

Rigaux K, Ferreira Castro A, Kanari L (2025) Quantifying neuronal differentiation using temporal topological persistence.

Rihn LL, Claiborne BJ (1990) Dendritic growth and regression in rat dentate granule cells during late postnatal development. *Developmental Brain Research* 54:115–124.

Romand S, Wang Y, Toledo-Rodriguez M, Markram H (2011) Morphological development of thick-tufted layer v pyramidal cells in the rat somatosensory cortex. *Frontiers in Neuroanatomy* 5:5.

Schneider-Mizell CM, Bodor AL, Brittain D, Buchanan J, Bumbarger DJ, Elabbady L, Gamlin C, Kapner D, Kinn S, Mahalingam G, Seshamani S, Suckow S, Takeno M, Torres R, Yin W, Dorkenwald S, Bae JA, Castro MA, Halageri A, Jia Z, Jordan C, Kemnitz N, Lee K, Li K, Lu

R, Macrina T, Mitchell E, Mondal SS, Mu S, Nehoran B, Popovych S, Silversmith W, Turner NL, Wong W, Wu J, Reimer J, Tolias AS, Seung HS, Reid RC, Collman F, da Costa NM (2025) Inhibitory specificity from a connectomic census of mouse visual cortex. *Nature* 640:448–458.

Schröter M, Paulsen O, Bullmore ET (2017) Micro-connectomics: probing the organization of neuronal networks at the cellular scale. *Nature Reviews Neuroscience* 18:131–146.

Shimono K, Sugimura K, Kengaku M, Uemura T, Mochizuki A (2010) Computational modeling of dendritic tiling by diffusible extracellular suppressor. *Genes to Cells* 15:137–149.

Shree S, Sutradhar S, Trottier O, Tu Y, Liang X, Howard J (2022) Dynamic instability of dendrite tips generates the highly branched morphologies of sensory neurons. *Science Advances* 8:eabn0080.

Sin WC, Haas K, Ruthazer ES, Cline HT (2002) Dendrite growth increased by visual activity requires NMDA receptor and Rho GTPases. *Nature* 419:475–480.

Soba P, Zhu S, Emoto K, Younger S, Yang SJ, Yu HH, Lee T, Jan LY, Jan YN (2007) Drosophila sensory neurons require Dscam for dendritic self-avoidance and proper dendritic field organization. *Neuron* 54:403–416.

Song Y, Ori-McKenney KM, Zheng Y, Han C, Jan LY, Jan YNN (2012) Regeneration of Drosophila sensory neuron axons and dendrites is regulated by the Akt pathway involving Pten and microRNA bantam. *Genes & development* 26:1612–1625.

Stepanyants A, Chklovskii DB (2005) Neurogeometry and potential synaptic connectivity. *Trends in Neurosciences* 28:387–394.

Stürner T, Ferreira Castro A, Philipps M, Cuntz H, Tavosanis G (2022) The branching code: A model of actin-driven dendrite arborization. *Cell Reports* 39:110746.

Sugimura K, Shimono K, Uemura T, Mochizuki A (2007) Self-organizing mechanism for development of space-filling neuronal dendrites. *PLoS Computational Biology* 3:e212.

Tavosanis G (2012) Dendritic structural plasticity. *Developmental Neurobiology* 72:73–86.

Uçar MC, Kamenev D, Sunadome K, Fachet D, Lallemend F, Adameyko I, Hadjab S, Hannezo E (2021) Theory of branching morphogenesis by local interactions and global guidance. *Nature Communications* 12:6830.

Vormberg A, Effenberger F, Muellerleile J, Cuntz H (2017) Universal features of dendrites [716] through centripetal branch ordering. *PLOS Computational Biology* 13:e1005615. [717]

Wen Q, Chklovskii DB (2008) A cost-benefit analysis of neuronal morphology. *Journal of* [718] *Neurophysiology* 99:2320–2328. [719]

Wen Q, Stepanyants A, Elston GN, Grosberg AY, Chklovskii DB (2009) Maximization of the [720] connectivity repertoire as a statistical principle governing the shapes of dendritic arbors. [721] *PNAS* 106:12536–12541. [722]

Winding M, Pedigo BD, Barnes CL, Patsolic HG, Park Y, Kazimiers T, Fushiki A, Andrade IV, [723] Khandelwal A, Valdes-Aleman J, Li F, Randel N, Barsotti E, Correia A, Fetter RD, Hartenstein [724] V, Priebe CE, Vogelstein JT, Cardona A, Zlatic M (2023) The connectome of an insect brain. [725] *Science* 379:eadd9330. [726]

Wittner L, Miles R (2007) Factors defining a pacemaker region for synchrony in the hippocampus. *Journal of Physiology* 584:867–883. [727] [728]

Yalgin C, Ebrahimi S, Delandre C, Yoong LF, Akimoto S, Tran H, Amikura R, Spokony R, [729] Torben-Nielsen B, White KP, Moore AW (2015) Centrosomin represses dendrite branching [730] by orienting microtubule nucleation. *Nature Neuroscience* 18:1437–1445. [731]