**Fgf/Ets signalling in *Xenopus* ectoderm initiates neural induction and patterning in an**

**autonomous and paracrine manners**

Ikuko Hongo* and Harumasa Okamoto*[+]

Department of Life Science, Faculty of Science, Gakushuin University, 1-5-1 Mejiro,

Toshima-ku, Tokyo 171-8588, Japan

*Present address: Institute for Biomolecular Science, Gakushuin University, 1-5-1 Mejiro,

Toshima-ku, Tokyo 171-8588, Japan

[+]Author for correspondence (harumasa.okamoto@gakushuin.ac.jp)

**Running title**

Fgf/Ets in neural development

**Keywords**

Fgf, Ets, signalling, neural induction, AP patterning, morphogen

**Summary statement**

Fgf/Ets signalling in ectodermal cells is required to initiate the expression of both anterior and

posterior neural genes from the late blastula to gastrula stages, independent of anti-Bmp

signalling.

**ABSTRACT**

Fibroblast growth factor (Fgf) and anti-bone morphogenetic protein (Bmp) signals are derived

from the organiser of mesoderm origin and cooperate to promote *Xenopus* neural development

from the gastrula ectoderm. Using antisense oligos to Fgf2 and Fgf8 and dominant-negative Ets

transcription factors, we showed that the expression of Fgf2, Fgf8, and Ets in ectoderm cells is

essential to initiate neural induction both *in vivo* and *in vitro*. Our findings show that neural

induction is initiated primarily by autonomous signalling in ectoderm cells, rather than by

paracrine signalling from organiser cells. The signalling in ectoderm cells is transduced via the

Fgf/Ras/Mapk/Ets pathway, independent of Bmp signal inhibition via the Fgf/Ras/Mapk/Smad1

route, as indicated by earlier studies. Through the same pathway, Fgfs activated

position-specific neural genes dose-dependently along the anteroposterior axis in cultured

ectoderm cells. The expression of these genes coincides with the establishment of the activated

Ets gradient within the gastrula ectoderm. Organiser cells, being located posteriorly to the

ectoderm, secrete Fgfs as gastrulation proceeds, which among several candidate molecules

initially promote neural patterning of the induced neuroectoderm as morphogens.

**INTRODUCTION**

Neural development in amphibians is initiated by the conversion of the intrinsic epidermal fate

of the gastrula ectoderm to the anterior neural fate (i.e. neural induction). During subsequent

development, the induced neuroectoderm of anterior identity is progressively patterned along

the anteroposterior (AP) axis (i.e. neural patterning), yielding the fully differentiated central

nervous system with the definitive AP pattern. Neural induction appears to be elicited by an

instructive signal from the Spemann organiser towards the adjacent ectoderm during

gastrulation; fibroblast growth factor (Fgf) and anti-bone morphogenetic protein (Bmp) signals

are currently considered to mediate the action of the Spemann organiser (Stern, 2005).

Loss-of-function and gain-of-function experiments have revealed that the combined activation

of Fgf signalling and suppression of Bmp signalling in the ectoderm is necessary and sufficient

to initiate neural induction (Wilson and Edlund, 2001; Stern, 2005). Bmps can promote the

epidermal differentiation of ectoderm cells in an autocrine manner. The organiser cells inhibit

this Bmp activity by releasing Bmp antagonists, such as Noggin and Chordin, which sequester

Bmps from their receptors via direct binding, thereby suppressing the autonomous epidermal

differentiation of ectoderm cells (Muñoz-Sanjuán and Brivanlou, 2002). Although organiser

cells express Fgfs (Lea et al., 2009), their role in neural induction is undefined, since some

previous experiments suggested that Fgfs derived from ectoderm cells primarily contribute to

neural induction in an autonomous manner (Gruntz and Tacke, 1989; Sato and Sargent, 1989;

Hongo et al., 1999; Delaune et al., 2005; Kuroda et al., 2005).

It is argued that active Fgf signalling reinforces the inhibition of Bmp signalling via the

mitogen-activated protein kinase (Mapk)/Smad1 pathway (Pera et al., 2003; Kuroda et al.,

2005). Mapk, a downstream transducer of the Fgf/Ras pathway, suppresses the transcriptional

activity of Smad1, a downstream target of Bmp signalling, via the phosphorylation of inhibitory

sites in the Smad1 linker region, leading to the cytoplasmic retention of Smad1 (Kretzschmar et

al., 1997; Pera et al., 2003). This suggests that Fgf and anti-Bmp signals are integrated at the

Smad1 level, which leads to adequate suppression of Bmp signalling in ectoderm cells to

manifest their neural fate, which is the default state. However, some studies have suggested that

Fgf signalling also functions in a Bmp-independent manner, exerting additional effects on

ectoderm cells via pathways other than the Fgf/Ras/Mapk/Smad1 (Delaune et al., 2005; Marchal

et al., 2009), but the mechanism remains to be clarified. One of the most promising candidates

for such a pathway is the Fgf/Ras/Mapk/Ets pathway, whereby Fgfs can promote target gene

expression by phosphorylating Ets family transcription factors via Mapk. We have previously

shown that the Fgf/Ras/Mapk/Ets and anti-Bmp signalling pathways are integrated at the level

of transcriptional regulation of *cdx4* (*Xcad3*), a posterior neural marker gene that is expressed at

early-phase neural patterning (Haremaki et al., 2003). Activated Ets and Sox2, de-repressed via

the inhibition of Bmp/Smad1 signalling (Mizuseki et al., 1998), cooperatively bind to their

respective recognition sequences, activating *cdx4*. It is also notable that ascidians, which share

the last common ancestor with vertebrates in chordate evolution, initiate neural induction via the

Fgf/Ras/Mapk/Ets pathway (Bertrand et al., 2003; Miya and Nishida, 2003).

In the present study, we aimed to clarify the mode of action and the source of Fgf signals

that are responsible for initiating neural induction and patterning. For this purpose, we blocked

endogenous Ets activity in ectoderm cells using dominant-negative constructs of Ets, such as

ΔxEts1 and ΔhElk1, and depleted several Fgf members in ectoderm cells using

translation-blocking morpholino-modified antisense oligonucleotides (MOs). When we regard

the activation of *otx2* and *sox2* as proving neural induction definitively, our *in vivo* and *in vitro*

results provide evidence that the Ets transcriptional factor contributes to initiating neural

induction via the Mapk/Ets route directly at the level of gene regulation. This is independent of

Bmp signal inhibition at the level of Smad1 via the Mapk/Smad1 route. Since our *in vitro* results

indicate the autonomous activity of ectoderm cells for neural induction, and Fgf4 has been

suggested to function in neural induction (Marchal et al., 2009), we aimed to identify the

members among a family of Fgf proteins that are expressed in ectoderm cells, exerting effects

on them during neural induction. It turned out that Fgf2 and Fgf8 were primarily required for

neural induction instead of Fgf4. We further attempted to verify the autonomous activity of

ectoderm cells in Fgf signalling using isolated culture of ectoderm cells.

Finally, we asked about the role of Fgfs expressed in the organiser (Lea et al., 2009). There

is an expression gradient of activated Ets within the gastrula ectoderm, with increasing levels

towards the organiser, which is located posteriorly (Schohl and Fagotto, 2002). Here, we

showed that Fgf/Ets signalling could activate position-specific neural genes, such as *otx2*,

*hes7.1*, *foxb1*, and *cdx4*, dose-dependently along the AP axis in cultured ectoderm cells. The

expression of these genes in normal development coincides with the establishment of the

activated Ets gradient. This indicates that Fgfs derived from organiser cells contribute in a

paracrine manner to initiating neural patterning of the induced neuroectoderm along the AP axis.

**RESULTS**

**Plasmid construction for the expression of dominant-negative Ets transcription factors**

To investigate the functional role of Ets transcription factors in neural induction and patterning,

we prepared plasmids encoding several dominant-negative forms of Ets, as described in the

materials and methods (Fig. 1A; ΔxEts1, ΔxEtv1, and ΔhElk1). They lack the activation domain,

comprising mainly the DNA-binding Ets domain, thereby suppressing endogenous Ets

transcriptional activity by competing for the target genes (Wasylyk et al., 1994). Two of the

dominant-negative constructs were fused with the Engrailed repressor domain (EnR) to confirm

that their native forms function as transcriptional activators (Fig. 1A; ΔhElk1· EnR and EnR·

ΔxEtv1) (Conlon et al., 1996). ΔΔxEtv1 and EnR are used as negative control constructs.

**Ets transcriptional activity in ectoderm cells is required for anterior neural development**

When synthetic *Δxets1* RNA (80 pg/blastomere) or *Δhelk1* RNA (40 pg/blastomere) was injected

into four animal blastomeres at the 8-cell stage (Fig. 1B, upper left), morphological defects in

the anterior region were prominently visible at the tadpole stage (Fig. 1D, E) compared with

uninjected (Fig. 1C) or *ΔΔxetv1* RNA-injected (Fig. 1G; 80 pg/blastomere) controls. The defects

included substantial reductions in head mass and eye tissue. Another dominant-negative

construct, ΔxEtv1, was ineffective at the dose used (80 pg RNA/blastomere; Fig. 1F). However,

ΔxEtv1, once fused with EnR, could effectively suppress Ets transcriptional activity (Fig. S1).

Upon injection of *Δhelk1* RNA (40 pg/blastomere) into a single dorsal-animal blastomere

(Fig. 1B, lower left) to obtain a more targeted RNA distribution, the prominent defect was

mostly confined to the eye tissue on the injected side (Fig. 1I). In embryos injected in either

protocol, the axial structure appeared to have been minimally affected (Fig. 1D, E, I upper

panels). This suggests that the anterior defects were elicited via direct effects on the ectoderm

cells derived from RNA-injected blastomere(s) rather than secondarily, via neural-inducing

organiser cells that generate primarily axial tissues. The tadpoles exhibiting severe anterior

defects with a global RNA distribution of *Δxets1* or *Δhelk1* RNA (Fig. 1D, E) appeared to be

phenotypic copies of tadpoles injected with *Δxfgfr-4a* RNA (Fig. 1H; Hongo et al., 1999;

Hardcastle et al., 2000; Brunsdon and Isaacs, 2020) in a similar manner (Fig. 1B, right diagram).

These experiments suggest that signalling through the Fgf/Ras/Mapk/Ets pathway in ectoderm

cells is required for proper anterior neural development.

**Ets transcriptional activity in ectoderm cells is required to activate *sox2* and *otx2***

To ascertain whether Ets transcriptional activity was essential to initiate neural induction, we

injected *Δhelk1* RNA with lineage tracer *gfp* RNA into three dorsal-animal blastomeres at the

16-cell stage (Fig. 2A). We examined the expression of the two early neural genes *sox2*

(pan-neural marker) and *otx2* (anterior neural marker) using whole-mount *in situ* hybridisation.

Both *sox2* and *otx2* are regarded as definitive markers for neural induction because their

expression starts at the late blastula and early gastrula stages, respectively; the data for the

temporal expression profiles of neural marker genes employed in this and the following

experiments, except for *otx2* (Blitz and Cho, 1995), were based on previously published

RNA-Seq analysis (Session et al., 2016). Typical results for *otx2* expression are shown in Fig.

2B and 2C. Dorsal view of embryos double-stained for *otx2* and *gfp* RNAs (Fig. 2C, left panels)

showed a reduction in *otx2* expression in lineage-labelled dorsal ectoderm compared with

uninjected controls (Fig. 2B, left panels). *Otx2* is expressed not only in a prospective anterior

neural region of the outer ectoderm layer but also in the underlying mesoderm and endoderm

layers during gastrulation (Blitz and Cho, 1995). To verify the site of *otx2* suppression, we

bisected stained embryos sagittally through the dorsal–ventral mid-line (Fig. 2A, cut).

Cut-surface view (Fig. 2C, middle and right panels) revealed that the tracer *gfp* RNA (stained

red) was mostly confined to the ectoderm layer, and *otx2* expression was effectively suppressed

in this tracer-labelled region. To ascertain the specificity of ΔhElk1, we co-injected wild-type

*xets1* RNA to rescue *otx2* expression. Co-injection mostly recovered *otx2* expression,

confirming that the suppression of *otx2* expression could be attributed to decreased Ets

transcriptional activity in the ectoderm cells (Fig. 2D). We then examined the effects of ΔhElk1

on the expression of *sox2*, which encodes a transcriptional factor required to activate *otx2*

(Mizuseki et al., 1998).   The injection of *Δhelk1* RNA suppressed *sox2* expression, which was

rescued by the co-injection of wild-type *xets1* RNA (Fig. 2E). In contrast, the expression of

*chordin*, which marks the organiser tissue, was not affected by the injection of *Δhelk1* RNA (Fig.

2F). These results provide evidence that Ets transcriptional activity in ectoderm cells is essential

for initiating neural induction.

**Ectoderm cells require Ets transcriptional activity to respond to Fgf signals**

To confirm that ectoderm cells require Ets transcriptional activity for directly responding to Fgf

signals, we exploited their primary culture in a microculture system as described by Kengaku

and Okamoto (1995) (Fig. 3A). In the present study, the expression of a set of six

position-specific neural marker genes was analysed using quantitative RT-PCR (Fig. 3), which

differed from a set of genes used in the previous study. Ectoderm cells of the early gastrula were

prepared from a combination of *Δxets1* RNA-injected embryos and *ΔΔxetv1* RNA-injected

control embryos or a combination of *Δxfgfr-4a* RNA-injected embryos and uninjected control

embryos. In the control experiments (Fig. 3B, D), Fgf induced ectoderm cells to express the six

marker genes along the AP axis in a dose-dependent manner, similar to the previously described

set of neural marker genes. Quantified data showed that lower doses of Fgf elicited more

anterior marker genes, such as *bf1* (telencephalon marker), *rax* (eye marker), or *pax6*

(telencephalon to spinal cord marker). In comparison, higher doses elicited more posterior

marker genes, such as *en2* (midbrain-hindbrain boundary marker), *hoxc6* (whole spinal cord

marker), or *cdx4* (posterior spinal cord marker) (Fig. 3F). We noted that the anterior marker

genes were expressed in the absence of Fgf, but this default type of expression might be caused

by the dissociation procedure during the preparation of ectoderm cells for culture (Gruntz and

Tacke, 1989), which was necessarily employed in our culture system (Fig. 3A). The expression

of the anterior neural genes peaked at a lower dose range of around 0.1 ng/mL and then declined

as the dose increased (Fig. 3F and Fig. S2E, F, G). This characteristic of anterior neural genes

being suppressed by a higher dose of Fgfs may explain why a high level of Fgf signalling

notably suppressed anterior development, as previously reported (Cox and Hemmati-Brivanlou,

1995; Lamb and Harland, 1995; Pownall et al., 1996; Polevoy et al., 2019).

The overexpression of ΔxEts1, as well as ΔxFgfr-4a, suppressed the expression of all the

above-mentioned marker genes (Fig. 3C, E, G–R), indicating that ectoderm cells required Ets

transcriptional activity to activate neural genes in response to Fgf signalling. Notably, the

anterior neural genes were less efficiently suppressed by ΔxEts1 than by ΔxFgfr-4a. A possible

explanation for the incomplete suppression by ΔxEts1 is that it did not interfere with Smad1

function, while ΔxFgfr-4a did, since Ets transcription factors are downstream of the

Fgf/Ras/Mapk pathway (Wasylyk et al., 1998). In ΔxEts1-overexpressing ectoderm cells, Mapk

could boost the inhibition of Bmp signalling via Smad1 phosphorylation at a level sufficient to

activate anterior neural genes to a certain extent, possibly combined with residual weak

Mapk/Ets signalling. In contrast, ΔxEts1 efficiently suppressed the expression of posterior

neural genes elicited by the increased dose of Fgf, suggesting that the reinforcement of Bmp

signal inhibition via the Mapk/Smad1 route was not necessarily essential for posterior neural

gene expression as long as Fgf signalling levels were kept high.

**Fgf/Ets signalling activates *sox2* and *otx2* in ectoderm cells**

To establish the direct role of the Ets transcription factor in ectoderm cells in initiating neural

induction in response to Fgf signals, we analysed the expression of *sox2* and *otx2* in cultured

ectoderm cells with or without dominant-negative Ets expression. In this series of experiments,

we employed ΔhElk1·EnR and EnR·ΔxEtv1 (Fig. 1A) to ascertain their native functions as

transcriptional activators. EnR·ΔxEtv1 overexpression in ectoderm cells could effectively

suppress *pax6* expression (Fig. S1), even more strongly than ΔxEts1 overexpression could (Fig.

3I). In contrast, ΔhElk1·EnR had little effect on *pax6* expression. The difference in the

suppression efficiency among these three dominant-negative constructs may reflect the

difference in DNA-binding specificity of their Ets domain.

The overexpression of ΔhElk1·EnR or EnR·ΔxEtv1 could suppress the expression of both

*sox2* and *otx2* (Fig. 3S–W), although the degree of suppression was not prominent, being similar

to the degree of suppression of other anterior neural marker genes, which were expressed later

in neural development, by ΔxEts1 (Fig. 3G–I). The incomplete suppression of *sox2* and *otx2*

might be explained in the same way as that of the other anterior neural marker genes by ΔxEts1,

as noted before. These results imply that Fgf signalling in ectoderm cells contributed to neural

induction at the level of transcriptional activation of target genes via the Mapk/Ets route,

independent of Bmp signalling inhibition via the Mapk/Smad1 route.

**Fgf2 and Fgf8 expression in ectoderm cells promotes anterior neural development via the**

**Fgf/Ets pathway**

To search for Fgf ligands that promote neural induction and their source, we investigated the

phenotypic effects of reducing Fgf2, Fgf3, Fgf4, Fgf8, and Fgf9 expression in ectoderm cells

using respective translation-blocking antisense MOs. Transcripts encoding these Fgfs are

present in embryos during the late blastula and early gastrula stages based on previous

RNA-Seq data (Session et al., 2016). Among them, Fgf4 alone was examined and indicated to

contribute to neural induction (Marchal et al., 2009). Upon injection of each Fgf MO into a

single dorsal-animal blastomere at the 8-cell stage, however, we found that Fgf2 MO and Fgf8

MO were much more potent than the other MOs, including Fgf4 MO, in causing anterior head

defects, such as reductions in head mass and eye tissue on the injected side (Fig. 4A, C, E–G; H

for collected data). The morphological phenotypes elicited by Fgf2 MO and Fgf8 MO were

quite similar to those induced by overexpressed dominant-negative Ets (Fig. 1D, E, I) or

xFgfr-4a (Fig. 1H) in ectoderm cells in that they exhibited prominent anterior head defects,

ensuring that the overall axial structure remained well preserved. These apparent phenotypic

similarities imply that the Fgf/Ets pathway functions autonomously in ectoderm cells for

anterior neural development, given that Fgf MOs were targeted primarily to ectoderm cells. We

obtained several lines of evidence supporting the specificity of Fgf2 MO and Fgf8 MO. Most

embryos injected with 5mis-Fgf2 MO or 5mis-Fgf8 MO formed a normal anterior head

structure (Fig. 4B, D, H).   The inhibition of anterior head development in each Fgf2 and Fgf8

morphant was primarily rescued by the co-injection of either cognate *fgf2* or *fgf8* RNAs (Fig.

4I–N). Most significantly, co-injection of wild-type *Xenopus etv1* (*xetv1*) RNA could also

recover anterior head development (Fig. 4O, P), ruling out the possibility of non-specific side

effects of the MOs, such as deteriorative effects on embryonic morphogenesis via an innate

immune response (Gentsch et al., 2018; Paraiso et al., 2019).

**Autonomous Fgf/Ets signalling in ectoderm cells is required for neural induction**

Depleting Fgf2 or Fgf8 in ectoderm cells by injecting the respective antisense MO into a single

dorsal-animal blastomere at the 8-cell stage (Fig. 5A) reduced the expression of *otx2* and *sox2* in

the prospective neural region of the stage 11 gastrula on the injected side compared with

5mis-Fgf2 MO controls (Fig. 5B, C). In contrast, the expression of *chordin* in the organiser

region was not affected (Fig. 5D). These results provide evidence that Fgf2 and Fgf8 in

ectoderm cells function autonomously during neural induction. The suppression of *otx2* and

*sox2* expression in Fgf morphants could be efficiently rescued by the co-injection of the cognate

*fgf* RNA (Fig. 5E, F left panels) or wild-type *xetv1* RNA (Fig. 5E, F, right panels). The rescue

by a wild-type Ets protein strongly supports our claim that neural-inducing Fgfs act via the

Fgf/Ras/Mapk/Ets pathway.

We further verified the autonomous activity of ectoderm cells in Fgf signalling by culturing

them in isolation. Under Fgf-free culture conditions, we found that ectoderm cells expressed

*fgf8* in addition to anterior neural genes. The time course of *fgf8* activation was roughly parallel

with that of anterior neural genes, such as *pax6* and *rax*, whereas *otx2* expression decreased

gradually to a certain level during this period (Fig. 6A). As shown before (Fig. 3W, uninjected

control), *otx2* expression in cultured ectoderm cells was gradually suppressed as the added Fgf

ligand dose increased. Thus, the temporal expression patterns shown in Fig. 6A could be

explained by ectoderm cells synthesising and secreting Fgf8 gradually to either activate or

suppress anterior neural genes. When ectoderm cells were prepared from Fgf8 morphants, the

expression of *fgf8* and other neural genes, including *sox2*, was reduced compared to ectoderm

cells prepared from control embryos injected with 5mis-Fgf8 MO (Fig. 6B). The expression of

the same set of genes could also be suppressed by pharmacological inhibitors of the

Fgf/Ras/Mapk pathway, such as SU5402 (against Fgf receptor) and U0126 (against Mapk

kinase), albeit to a varying degree (Fig. 6C). Finally, the overexpression of EnR·ΔxEtv1 or

ΔhElk·EnR in ectoderm cells also reduced the expression of these genes (Fig. 6D). Collectively,

autocrine Fgf8 signalling in ectoderm cells contributed to neural induction via the

Fgf/Ras/Mapk/Ets pathway.

**Fgf/Ets signalling in ectoderm cells promotes neural patterning as an early morphogenic**

**factor**

To verify the role of Fgf/Ets signalling in initiating neural patterning, we examined the

expression profiles of position-specific neural marker genes of the early developmental stage in

ectoderm cells cultured with Fgf. They include *otx2* (forebrain marker), *hes7.1* (midbrain–

hindbrain boundary marker), *foxb1* (hindbrain marker), and *cdx4* (posterior spinal cord marker),

which are first activated around stage 10, peaking around stage 12 in normal development

(Session et al., 2016). Their expression patterns in whole embryos at stage 11, as revealed by *in*

*situ* hybridisation, are presented in Fig. 7A, with that of *sox2*. It should be noted that the

expression profiles of position-specific marker genes used earlier (Fig. 3F), except *cdx4*, did not

necessarily reflect the initial phase of neural patterning, since their expression began during the

mid-gastrula stage (around stage 12) and reached a peak during the early neurula stage (around

stage 15) in normal development.

Control experiments showed that Fgfs could induce ectoderm cells to express the earlier

marker genes along the AP axis in a dose-dependent manner (Fig. 7C, E; Fig. S2A, B, D),

similar to the later phase marker genes shown in Fig. 3F. The overexpression of EnR·ΔxEtv1 or

ΔhElk1·EnR reduced the expression of the marker genes to a varying degree. EnR·ΔxEtv1

effectively suppressed *hes7.1* expression but had little effect on *foxb1* and even enhanced *cdx4*

expression, whereas ΔhElk1·EnR effectively suppressed both (Fig. 7F–H). *Otx2* expression was

reduced by the two constructs to a similar extent (Fig. 3W). The varying degree of suppression

may be interpreted as the difference in DNA-binding specificities of the respective Ets domains

of xEtv1 and hElk1 to each marker gene, as noted before. Our data suggest that Fgf signalling in

ectoderm cells promoted neural patterning as early as the early gastrula stage in a

dose-dependent manner, which required Ets proteins as transcriptional activators.

**DISCUSSION**

**Fgf2 and Fgf8/Ets signalling in ectoderm cells is required to initiate neural induction**

By overexpressing a dominant-negative form of an Ets transcription factor (ΔhElk1), we

showed that endogenous Ets transcriptional activity in ectoderm cells is specifically required to

initiate neural induction, as revealed by the suppression of *sox2* and *otx2* expression and its

rescue by wild-type xEts1 in gastrula embryos (Fig. 2C, D, E). We also showed the requirement

of Ets transcriptional activity for cultured ectoderm cells to activate *sox2* and *otx2* (Fig. 3V, W).

Our previous studies, in which ΔxFgfr-4a was overexpressed in cultured ectoderm cells, showed

that Fgf receptor signalling is required for ectoderm cells to adopt a neural fate in response to

co-cultured intact organiser cells (Hongo et al., 1999). Furthermore, the morphological

phenotype of tadpoles elicited by ΔhElk1 or ΔxEts1 was quite similar to that by ΔxFgfr-4a (Fig.

1D, E, H). These findings indicate that the Fgf/Ras/Mapk/Ets pathway in ectoderm cells is

essential for initiating neural induction.

Numerous *fgf* transcripts are present in blastula and gastrula embryos, as shown by

RNA-Seq analysis (Suzuki et al., 2017). A report suggested that Fgf4 is a member of the Fgf

protein family that functioned in neural induction (Marchal et al., 2009), but the possibility of

contributions of other Fgf members remains unaddressed. We attempted to identify members of

the Fgf family exerting effects on ectoderm cells during neural induction. For this, we knocked

down Fgf activity in ectoderm cells by injecting several translation-blocking MOs into a single

8-cell-stage dorsal-animal blastomere. The depletion of Fgf2 and Fgf8 in ectoderm cells by

antisense MO most effectively suppressed *sox2* and *otx2* expression in gastrula embryos (Fig.

5B, C) and caused severe defects in the anterior head in tadpoles (Fig. 4A, C), similar to the

effects of the overexpression of dominant-negative Ets proteins (Fig. 1D, E; Fig. 2C, E middle

panels). The suppression of *otx2*, as well as anterior head development, in Fgf morphants was

rescued by xEtv1, a wild-type *Xenopus* Ets protein (Fig. 4O, P; Fig. 5E, F, right panels).

Collectively, these results provide evidence that Fgf2 and Fgf8 act via the Fgf/Ras/Mapk/Ets

pathway in ectoderm cells, independent of the Mapk/Smad1 branch, to initiate neural induction,

considering the expression of *sox2* and *otx2* as an indication of the start of neural induction.

**Fgf2 and Fgf8/Ets signalling acts autonomously in neural induction**

The microinjection of dominant-negative forms of Ets or Fgf MOs into animal blastomeres at

the 8- or 16-cell stage exhibited minimal effects on the functional activities of the organiser cells,

since the expression of *chordin* at the gastrula stage (Figs 2F, 5D), and subsequent formation of

the axial structures were barely affected (Fig. 1D, E; Fig. 4A, C). This indicates that

neural-inducing Fgfs function autonomously between ectoderm cells rather than in a paracrine

manner from the organiser cells. Consistently, some previous experiments suggested the

possible involvement of ectoderm cells in neural induction, which contribute in an autonomous

manner (Delaune et al., 2005; Kuroda et al., 2005). The idea is based on the observation that

ectoderm cells subjected to prolonged dissociation differentiate into neural cells of anterior

identity in the absence of organiser cells (Gruntz and Tacke, 1989; Sato and Sargent, 1989) and

that this autonomous type of *in vitro* neuralization is blocked by ΔxFgfr-4a overexpression

(Hongo et al., 1999).

In the microculture system we used, a dissociation procedure during the preparation of the

ectoderm cells was a prerequisite step, which caused the expression of pan-neural *sox2* and

anterior neural marker genes in the absence of Fgf (Fig. 3G, H, I, M, N, O, V, and W). We found

in this culture system of ectoderm cells that *fgf8* was also activated autonomously along with the

neural genes (Fig. 6A). In normal development, *fgf8* is activated as early as the late blastula

(stage 9), and maternally expressed *fgf2* is also present during the blastula and gastrula stages

(Suzuki et al., 2017). Our preliminary experiments showed that Fgf2 MO could suppress the

expression of *chordin* in the dorsal marginal region of the late blastula (Fig. S3), indicating that

Fgf2 signalling was indeed active during the late blastula stage. Maternal Fgf2 also possibly

triggered *fgf8* activation at this stage, and autonomous *fgf8* expression was reinforced by a

positive loop of Fgf signalling, as inferred from the suppression of *fgf8* in Fgf8 morphants (Fig.

6B). Our results correspond with an earlier report that phosphorylated Mapk, an indicator of

active Fgf signalling, is located in the dorsal-animal region during the blastula stage (Schohl and

Fagotto, 2002). These findings are consistent with the idea that autonomous Fgf signalling via

the Mapk/Ets route contributes to initiating neural induction, assuming the onset of *sox2*

expression at the blastula stage 9 as the onset of neural induction. Notably, autonomous action

of Chordin, Noggin, and Nordal3.1 within a group of cells in the dorsal-animal region is

required to initiate neural induction during the late blastula stage (Kuroda et al., 2004).

Consistently, activated (phosphorylated) Smad1 expression is maintained at a low level in this

period within the dorsal-animal region (Schohl and Fagotto, 2002). Thus, the integration of

autonomous positive-Fgf signalling and anti-Bmp signalling in ectoderm cells might be

essential for neural induction from the very start.

Besides Fgf2 and Fgf8, Fgf4 is involved in neural induction (Marchal et al., 2009), but *fgf4*

is activated around stage 10, with its transcript levels considerably lower than those of Fgf2 and

Fgf8 (Suzuki et al., 2017). However, comparison of dose-response profiles of these Fgfs for the

activation of neural genes indicated that Fgf4 was more potent than Fgf8 (Fig. S2), revealing the

possibility that Fgf4 expressed in the organiser region reinforced neural induction initiated by

Fgf2 and Fgf8 and subsequently contributed to neural patterning. It should also be noted that

*igf2*, encoding another candidate of neural inducer (Pera et al., 2003), is activated after stage 10

and may contribute to neural induction and patterning together with Fgf4.

**Fgf/Ets signalling is required for neural patterning as an early morphogenic factor**

In cultured isolated ectoderm cells, none of the posterior neural genes is activated, but Fgf can

induce them to express position-specific neural marker genes along the AP axis in a

dose-dependent manner, with lower doses eliciting more anterior genes and higher doses

eliciting more posterior genes (Kengaku and Okamoto, 1995). Here, we further showed that

Fgf/Ets signalling could dose-dependently activate a different set of position-specific markers,

such as *otx2*, *hes7.1*, *foxb1*, and *cdx4* (Fig. 7). Notably, their expression in normal development

starts around stage 10 and peaks around stage 12 (Session et al., 2016), coinciding with the

establishment of a gradient of the level of phosphorylated Ets within the gastrula ectoderm layer,

which increases towards the organiser, located posteriorly (Schohl and Fagotto, 2002). These

findings imply that Fgfs could function as early neural patterning morphogens in normal

development via the Fgf/Ras/Mapk/Ets pathway in the ectoderm layer. Indeed, the Fgf/Ets route

was shown to directly activate *cdx4* (Haremaki et al., 2003), which triggers the expression of

several *hox* genes, regulating posterior development during later developmental stages (Isaacs et

al., 1998).

Wnt/β-catenin signalling has been reported to regulate the AP neural patterning in *Xenopus*

(Kiecker and Niehrs, 2001). However, the gradient pattern of this signalling was not apparent

until mid-gastrula stage 11 (Schohl and Fagotto, 2002), suggesting that Wnt/β-catenin signalling

may reinforce and further promote the process of neural patterning. Retinoic acid (RA) is

another candidate for a neural patterning morphogen. A gradient of RA is established around the

stage 12 *Xenopus* neurula, exhibiting a peak at the hindbrain–spinal cord boundary (Pera et al.,

2014). Furthermore, RA-responsive enhancers are located at the 3′-region of the *hox* clusters

(Langston et al., 1997), which regulate AP patterning in the hindbrain and the spinal cord during

a later stage of neural patterning (Strate et al., 2009). Our results, in line with previous research,

indicated that Fgfs functioned as morphogens to provide a basis for neural patterning via the

Fgf/Ras/Mapk/Ets pathway.

**Source of Fgfs for neural patterning**

We attempted to ascertain how the neural patterning Fgfs are derived and contribute to the

establishment of the active Ets gradient. *Fgf8* was autonomously activated in cultured ectoderm

cells by a positive feedback loop (Fig. 6B). However, this autonomous *fgf8* expression was

suppressed by increasing the dose of Fgfs (Fig. S2H), suggesting that negative feedback

regulation of *fgf8* expression was also present in ectoderm cells. Our findings are consistent

with prevalent observations, including ours, that autonomous neuralization of ectoderm cells is

never accompanied by the expression of posterior neural genes (Gruntz and Tacke, 1989; Sato

and Sargent, 1989; Hongo et al., 1999; Muñoz-Sanjuán and Brivanlou, 2002). In contrast to the

expression of *fgf8* and anterior neural genes, the expression of posterior neural genes required

Fgfs in a dose-dependent manner along the AP axis; the higher dose elicited more posterior

genes (Fig. 3F and 7E). We previously showed that ectoderm cells, when co-cultured with

organiser cells, express position-specific posterior neural genes such as *egr2/krox20*,

*hoxc6/XlHbox1*, *hoxc9/XlHbox6*, and *cdx4/Xcad3* in an organiser cell number-dependent

manner, which was suppressed by ΔxFgfr-4a overexpression in ectoderm cells (Hongo et al.,

1999). These findings imply that, in normal development, organiser cells promote the patterning

of the neighbouring neuroectoderm along the AP axis by releasing Fgfs in a paracrine manner. A

gradient of active Fgf signalling along the AP axis is established in the neuroectoderm during

early (stage 10) to mid (stage 10.5) gastrula stages, with increasing levels towards the organiser

region (Schohl and Fagotto, 2002). When we re-examined this aspect, a gradient of P-Mapk

localisation along the AP axis in the prospective neural region of gastrula embryos was

confirmed (Fig. S4).

**Evolutionary aspects of Fgf/Ets signalling**

We wonder how far Fgf/Ras/Mapk/Ets signalling is conserved and plays an essential role in

neural development during animal evolution. The requirement of Fgf signals for neural

induction is conserved between several vertebrate species, such as zebrafish (Furthauer et al.,

2004), *Xenopus* (Hongo et al., 1999; Delaune et al., 2005), and chicken (Streit et al., 2000;

Wilson et al., 2000), but the employment of the Fgf/Ets pathway has only been shown for

*Xenopus* (this study). Notably, however, two species of ascidians, which share the last common

ancestor with vertebrates in chordate evolution, initiate neural induction via the

Fgf/Ras/Mapk/Ets pathway, as shown by *otx* expression (Bertrand et al., 2003 in *Ciona*

*intestinalis*; Miya and Nishida, 2003 in *Halocynthia roretzi*). Furthermore, in *Ciona*, the

regulatory element driving *otx* expression in the prospective neural cells was identified, which

directly responded to Fgf signalling. These findings suggest that the last common ancestor of

ascidians and vertebrates in the chordate lineage employs Fgf/Ets signalling for the process of

neural induction. It will be interesting to ask whether vertebrate species other than *Xenopus*

employ this signalling pathway.

The Fgf/Ras/Mapk/Ets signalling is also involved in neural patterning in *Ciona*, as revealed

by *six3* and *otx* expression (Gainous et al., 2015). Remarkably, even some protostomes, such as

annelids, arthropods, and onychophorans, express *six3* and *otx* in this order along the AP axis in

their anterior neuroectoderm (Steinmetz et al., 2010), as shown in *Xenopus* and *Ciona*, which

are classified as deuterostomes. Whether the expression pattern of *six3* and *otx* in protostomes is

regulated via Fgf/Ets signalling is yet to be addressed. However, in the planarian, another

protostome, Fgfr-related genes were involved in neural development (Cebria et al., 2002). It is

tempting to assume that the last common ancestor of bilaterians already acquired the Fgf/Ets

signalling pathway for neural development, although the original pathway might have been

modified or lost in some of its descendants.

**Concluding remarks**

Among a number of candidate signal molecules, Fgf2 and Fgf8 primarily contribute to initiating

neural induction. They function in ectoderm cells via the Mapk/Ets route directly at the level of

transcriptional regulation of the target genes. This is independent of Bmp signal inhibition via

the Mapk/Smad1 route, which was proposed to be the prevailing Fgf signalling route in neural

induction. Our findings satisfy the requirement of a different route for neural induction,

hypothesized from different lines of evidence. The Fgf signals for neural induction are derived

primarily from ectoderm cells themselves in an autonomous manner. This might conflict with

the leading idea that the organiser plays the primary role for both neural induction and

patterning. However, there is a possibility that the precursors of the organiser cells are present in

the dorsal marginal zone of the late blastula apart from regular ectoderm cells, and these

precursor cells release Fgfs, contributing to the earliest stage of neural induction in a paracrine

mode; the issue is yet to be addressed. Fgfs derived from the organiser at the gastrula stage

contribute to neural patterning in a paracrine manner. However, the Fgf members involved were

not identified, since loading MOs into the organiser cells blocked their differentiation,

disturbing the gastrulation process. Our work provides support for a shared use of Fgf/Ets

signalling for neural development during chordate evolution, but further work is needed to

ascertain its wider conservation during animal evolution.

**MATERIALS AND METHODS**

**Animal care**

*Xenopus laevis* embryos were obtained (Mitani and Okamoto, 1989) and staged (Nieuwkoop

and Farber, 1967) as previously described. The handling of animals was carried out in

accordance with nationally prescribed guidelines and the guidelines for animal experiments at

Gakushuin University.

**Plasmid construction**

Full or partially deleted coding sequences of *Xenopus*, *ets1* (*xets1*) and *etv1* (*xetv1*), and human

*elk1* (*helk1*) were subcloned into pSP64T as described previously (Haremaki et al., 2003);

*Δhelk1* RNA blocked Ets transcriptional activity more efficiently than *Δxets1* or *Δxetv1* RNA,

albeit being of human origin. *Δhelk1·enr* was generated by in-frame C-terminal fusion of the

*Drosophila engrailed* repressor region (EnR; Conlon *et al*., 1996) to *Δhelk1*, while *enr·Δxetv1*

was generated by in-frame N-terminal fusion of EnR to *Δxetv1*. *ΔΔxetv1* is deleted further in the

N-terminal region of *Δxetv1* to eliminate its DNA-binding capacity (Papoutsopoulou and

Janknecht, 2000), thereby serving as a negative control construct. Structural features of the Ets

proteins and their dominant-negative derivatives are shown in Fig. 1A.

**Antisense MOs**

Translation-blocking antisense MOs were obtained from Gene Tools LLC (Philomath, OR,

USA). These are listed in Supplementary information Table S1.

**Whole-mount *in situ* hybridisation**

*In situ* hybridisation experiments were performed following the previously described methods

(Harland, 1991; Sive *et al*., 1995). Digoxigenin-labelled antisense RNAs were prepared by the

transcription of respective plasmids carrying the target genes, which were linearised. Hybridised

digoxigenin-containing RNAs were visualised with anti-digoxigenin antibodies conjugated to

alkaline phosphatase and BM purple. For visualising the lineage tracer *gfp* RNA, embryos

stained with BM purple were collected in MEMFA (100 mM MOPS (pH 7.4), 2 mM EGTA, 1

mM $MgSO_4$, 3.7% formaldehyde) and rinsed twice with PBST (PBS + 0.1 % Tween 20) for 1h.

Embryos were then processed for *in situ* hybridisation to *gfp* RNA as described above, except

for the use of fluorescein-labelled antisense RNA, anti-fluorescein secondary antibody, and Fast

Red. In the experiments described in Fig. 2, stained embryos were bisected as described by

Sudou et al. (2012).

**Microinjection of RNAs and Fgf MOs**

Capped synthetic RNAs for microinjection were synthesized as described previously (Hongo et

al., 1999) and injected into animal blastomeres at the 8-cell, 16-cell, or 32-cell stage. Fgf MOs

with or without cognate RNA or wild-type *xetv1* RNA were injected into animal blastomeres at

the 8-cell stage. The cognate *fgf2* and *fgf8* RNAs were designed to mismatch with the respective

antisense MOs but conserved the same amino acid sequence as the wild-type RNAs. They were

synthesized from mutated plasmids that were constructed from wild-type *fgf2* and *fgf8* plasmids

via PCR, using 5′ forward primers with mutated cognate sequences (Table S2).

*Xenopus* **ectoderm cell microculture and quantitative RT-PCR**

Methods for culturing early gastrula cells were essentially as described previously (Mitani and

Okamoto, 1991; Kengaku and Okamoto, 1993; 1995). Animal cap fragments were dissected

from injected or uninjected embryos (stage 10), which were dissociated by incubation in $Ca^{2+}$-

and $Mg^{2+}$-deficient MBS containing 1% BSA at room temperature. The dispersed cells were

then suspended in standard MBS containing 1% BSA, and the desired number of cells were

inoculated onto Terasaki plates (150 ectoderm cells/well). After completion of reaggregation by

brief centrifugation, cells were incubated in the presence or absence of Fgfs at 22.5°C in

humidified air until control embryos reached stage 14 or 25 (Fig. 3A). In some experiments,

inhibitors of the Fgf/Ras/Mapk pathway, such as SU5402 (CAS 215543-92-3, Calbiochem) and

U0126 (CAS 109511-58-2, Calbiochem), were added with Fgfs. Recombinant Fgfs used were as

follows; bovine Fgf2 (bFgf) from Progen Biotechnik GmbH (Heidelberg,Germany), human

Fgf4 from PeproTech Inc (Cranbury, NJ, USA), and mouse Fgf8 from R&D Systems Inc

(Minneapolis, MN, USA). RT-PCR was performed as described previously (Hongo et al.,

1999). PCR products were separated on a 4% polyacrylamide gel, and the radioactivity of each

PCR product was estimated using a laser image analyser (Fujix BAS 2000, Fuji Film). When

isotopic measurement was not available, the intensity of the band stained in the gel was

estimated using ImageJ. PCR products were standardised against a co-amplified internal control,

*elongation factor 1α* (*ef1α*; Krieg et al., 1989). The primers used are listed in Table S1.

**Staining of activated Mapk (dpERK)**

Whole-mount dpERK staining was performed as previously described (Christian and Slack,

1999).

**ACKNOWLEDGEMENT**

We thank Professor Takashi Adachi-Yamada for critical reading of the manuscript.

**COMPETING INTERESTS**

The authors declare no competing or financial interests.

**FUNDING**

This work was supported in part by grant from MEXT*-Supported Program for the Private

University Research Branding Project 2016‑2020 (*Ministry of Education, Culture, Sports,

Science and Technology).

**REFERENCES**

**Bertrand, V., Hudson, C., Caillol, D., Popovici, C. and Lemaire, P.** (2003). Neural tissue in

ascidian embryos is induced by FGF9/16/20, acting via a combination of maternal GATA and

Ets transcription factors. *Cell* **115**, 615-627. doi: 10.1016/s0092-8674(03)00928-0

**Blitz, I. L. and Cho, K. W.** (1995). Anterior neurectoderm is progressively induced during

gastrulation: The role of the *Xenopus* homeobox gene *orthodenticle*. *Development* **121**,

993-1004.

**Brunsdon, H. and Isaacs, H. V.** (2020). A comparative analysis of fibroblast growth factor

receptor signalling during *Xenopus* development. *Biol. Cell* **112**, 1-13. doi:

10.1111/boc.201900089

**Cebria, F., Kobayashi, C., Umesono, Y. Nakazawa, M., Mineta, K., Ikeo, K., Gojobori, T.,**

**Itoh, M., Taira, M., Sanchez Alvarado, A. and Agata, K.** (2002). FGFR-related gene

nou-darake restricts brain tissues to the head region of planarians. *Nature* **419**, 620-624. doi:

10.1038/nature01042

**Christen, B. and Slack, J.M.** (1999) Spatial response to fibroblast growth factor signalling in

*Xenopus* embryos. *Development* **126**, 119-125.

**Conlon, F. L., Sedgwick, S. G., Weston, K. M. and Smith, J. C.** (1996). Inhibition of Xbra

transcription activation causes defects in mesodermal patterning and reveals autoregulation

of Xbra in dorsal mesoderm. *Development* **122**, 2427-2435.

**Cox, W. G. and Hemmati-Brivanlou, A.** (1995). Caudalization of neural fate by tissue

recombination and bFGF. *Development* **121**, 4349-4358.

**Delaune, E., Lemaire, P. and Kodjabachian, L.** (2005). Neural induction in *Xenopus* requires

early FGF signalling in addition to BMP inhibition. *Development* **132**, 299-310. doi:

10.1242/dev.01582

**Furthauer, M., van Celst, J., Thisse, C. and Thisse, B.** (2004). Fgf signalling controls the

dorsoventral patterning of the zebrafish embryo. *Development* **131**, 2853-2864. doi:

10.1242/dev.01156

**Gainous, T. B., Wagner, E. and Levine, M.** (2015). Diverse ETS transcription factors mediate

FGF signaling in the *Ciona* anterior neural plate. *Dev. Biol.* **399**, 218-225. doi:

10.1016/j.ydbio.2014.12.032

**Gentsch, G. E., Spruce, T., Monteiro, R. S., Owens, N. D. L., Martin, S. R. and Smith, J. C.**

(2018). Innate immune response and off-target mis-splicing are common

morpholino-induced side effects in *Xenopus*. *Dev. Cell* **44**, 597-610.e10. doi:

10.1016/j.devcel.2018.01.022

**Grunz, H. and Tacke, L.** (1989). Neural differentiation of *Xenopus laevis* ectoderm takes place

after disaggregation and delayed reaggregation without inducer. *Cell Differ. Dev.* **28**, 211-218.

doi: 10.1016/0922-3371(89)90006-3

**Hardcastle, Z., Chalmers, A. D. and Papalopulu, N.** (2000). FGF-8 stimulates neuronal

differentiation through FGFR-4a and interferes with mesoderm induction in Xenopus

embryos. *Curr. Biol.* **10**, 1511-1514. doi: 10.1016/S0960-9822(00)00825-3

**Haremaki, T., Tanaka, Y., Hongo, I., Yuge, M. and Okamoto, H.** (2003). Integration of

multiple signal transducing pathways on Fgf response elements of the *Xenopus caudal*

homologue *Xcad3*. *Development* **130**, 4907-4917. doi: 10.1242/dev.00718

**Harland, R. M.** (1991). In situ hybridization: An improved whole mount method for *Xenopus*

embryos. *Methods Cell Biol.* **36**, 685-695.

**Hongo, I., Kengaku, M. and Okamoto, H.** (1999). FGF signaling and the anterior neural

induction in *Xenopus*. *Dev. Biol.* **216**, 561-581. doi: 10.1006/dbio.1999.9515

**Isaacs, H. V., Pownall, M. E. and Slack, J. M.** (1998). Regulation of Hox gene expression and

posterior development by the *Xenopus* caudal homologue *Xcad3*. *EMBO J.* **17**, 3413-3427.

doi: 10.1093/emboj/17.12.3413

**Kengaku, M. and Okamoto, H.** (1993). Basic fibroblast growth factor induces differentiation

of neural tube and neural crest lineages of cultured ectoderm cells from *Xenopus* gastrula.

*Development* **119**, 1067-1078.

**Kengaku, M. and Okamoto, H.** (1995). bFGF as a possible morphogen for the anteroposterior

axis of the central nervous system in *Xenopus. Development* **121**, 3121-3130.

**Kiecker, C. and Niehrs, C.** (2001). A morphogen gradient of Wnt/β-catenin signalling regulates

anteroposterior neural patterning in *Xenopus*. *Development* **128**, 4189-4201.

**Kretzschmar, M., Doody, J. and Massague, J.** (1997). Opposing BMP and EGF signalling

pathways converge on the TGF-β family mediator Smad1. *Nature* **389**, 618-622. doi:

10.1038/39348

**Krieg, P. A., Varnum, S. M., Wormington, W. M. and Melton, D. A.** (1989). The mRNA

encoding elongation factor 1α (EF1α) is a major transcript at the mid-blastula transition in

*Xenopus. Dev. Biol.* **133**, 93-100. doi: 10.1016/0012-1606(89)90300-X

**Kuroda, H., Wessely, O. and Robertis, E. M.** (2004). Neural induction in *Xenopus*:

requirement for ectodermal and endomesodermal signals via Chordin, Noggin, β-Catenin,

and Cerberus. *PLoS Biol.* **2**, 623-634. doi: 10.1371/journal.pbio.0020092

**Kuroda, H., Fuentealba, L., Ikeda, A., Reversade, B. and De Robertis, E.** (2005). Default

neural induction: neuralization of dissociated *Xenopus* cells is mediated by Ras/MAPK

activation. *Genes Dev.* **19**, 1022-1027. doi: 10.1101/gad.1306605

**Lamb, T. M. and Harland, R. M.** (1995). Fibroblast growth factor is a direct neural inducer,

which combined with noggin generates anterior-posterior neural pattern. *Development* **121**,

3627-3636.

**Langston, A. W., Thompson, J. R. and Gudas, L. J.** (1997) Retinoic acid-responsive

enhancers located 3′ of the Hox A and Hox B homeobox gene clusters. Functional analysis. *J.*

*Biol. Chem.* **272**, 2167-2175. doi: 10.1074/jbc.272.4.2167

**Lea, R., Papalopulu, N., Amaya, E. and Dorey, K.** (2009). Temporal and spatial expression of

FGF ligands and receptors during *Xenopus* development. *Dev. Dyn.* **238**, 1467-1479. doi:

10.1002/dvdy.21913

**Marchal, L., Luxardi, G., Thome, V. and Kodjabachian, L.** (2009). BMP inhibition initiates

neural induction via FGF signaling and Zic genes. *Proc. Natl Acad. Sci*. USA **106**,

17437-17442. doi: 10.1073/pnas.0906352106

**Mitani, S. and Okamoto, H.** (1989). Embryonic development of *Xenopus* studied in a cell

culture system with tissue specific monoclonal antibodies. *Development* **105**, 53-59.

**Mitani, S. and Okamoto, H.** (1991). Inductive differentiation of two neural lineages

reconstituted in a microculture system from *Xenopus* early gastrula cells. *Development* **112**,

21-31.

**Miya, T. and Nishida, H.** (2003). An Ets transcription factor, HrEts, is target of FGF signaling

and involved in induction of notochord, mesenchyme, and brain in ascidian embryos. *Dev.*

*Biol*. **261**, 25-38. doi: 10.1016/s0012-1606(03)00246-x

**Mizuseki, K., Kishi, M., Matsui, M., Nakanishi, S. and Sasai, Y.** (1998). *Xenopus*

Zic-related-1 and Sox-2, two factors induced by chordin, have distinct activities in the

initiation of neural induction. *Development* **125**, 579-587.

**Muñoz-Sanjuán, I. and Brivanlou, A. H.** (2002). Neural induction, the default model and
embryonic stem cells. *Nat. Rev. Neurosci.* **3**, 271-280. doi: 10.1038/nrn786

**Nieuwkoop, P. D. and Faber, J.** (1967). *Normal Table of Xenopus laevis (Daudin).*
Amsterdam: North Holland, J. Faber Garland Publishing.

**Papoutsopoulou, S. and Janknecht, R.** (2000). Phosphorylation of ETS transcription factor
ER81 in a complex with its coactivators CREB-binding protein and p300. *Mol. Cell. Biol.* **20**,
7300-7310. doi: 10.1038/s41598-019-44685-3

**Paraiso, K. D., Blitz, I. L., Zhou, J. J. and Cho, K. W. Y.** (2019). Morpholinos do not elicit an
innate immune response during early *Xenopus* embryogenesis. *Dev. Cell* **49**, 643-650.e3. doi:
10.1016/j.devcel.2019.04.019

**Pera, E.M., Ikeda, A., Eivers. E. and De Robertis, M.** (2003). Integration of IGF, FGF, and
anti-BMP signals via Smad1 phosphorylation in neural induction. *Genes Dev.* **17**, 3023-3028.
doi: 10.1101/gad.1153603

**Pera, E.M., Acosta, H., Gouignard, N., Climent, M. and Arregi, I.** (2014). Active signals,
gradient formation and regional specificity in neural induction. *Exp. Cell Res.* **321**, 25-31.
doi: 10.1016/j.yexcr.2013.11.018

**Polevoy, H., Gutkovich, Y.E., Michaelov, A., Volovik, Y., Elkouby, Y.M. and Frank, D.**
(2019). New roles for Wnt and BMP signaling in neural anteroposterior patterning. *EMBO*
*Rep*. **20**, 1-13. doi: 10.15252/embr.201845842

**Pownall, M. E., Tucker, A. S., Slack, J. M. W. and Isaacs, H. V.** (1996). eFGF, Xcad3 and
Hox genes form a molecular pathway that establishes the anteroposterior axis in *Xenopus.*
*Development* **122**, 3881-3892.

**Sato, S. M. and Sargent, T. D.** (1989). Development of neural inducing capacity in dissociated
*Xenopus* embryos. *Dev. Biol.* **134**, 263-266. doi: 10.1016/0012-1606(89)90096-1

**Schohl, A. and Fagotto, F.** (2002). β-catenin, MAPK and Smad signaling during early *Xenopus*

development. *Development* **129**, 37-52.

**Session, A. M., Uno, Y., Kwon, T., Chapman, J. A., Toyoda, A., Takahashi, S., Fukui, A.,**

**Hikosaka, A., Suzuki, A., Kondo, M. et al.** (2016). Genome evolution in the allotetraploid

frog *Xenopus laevis*. *Nature* **538**, 336-343. doi: 10.1038/nature19840

**Sive, H. L., Grainger, R. M. and Harland, R. M.** (1995). *Early Development of Xenopus*

*laevis: Course Manual.* NY, USA: Cold Spring Harbor, Cold Spring Harbor Laboratory

Press.

**Steinmetz, P. R. H., Urbach, R., Posnien, N., Eriksson, J., Kostyuchenko, R. P., Brena, C.,**

**Guy, K., Akam, M., Bucher, G. and Arendt, D.** (2010). Six3 demarcates the anterior-most

developing brain region in bilaterian animals. *EvoDevo* **1**, 14. doi: 10.1186/2041-9139-1-14

**Stern, C.D.** (2005). Neural induction: old problem, new findings, yet more questions.

*Development* **132**, 2007-2021. doi: 10.1242/dev.01794

**Strate, I., Min, T. H., Iliev, D. and Pera, E. M.** (2009). Retinol dehydrogenase 10 is a

feedback regulator of retinoic acid signalling during axis formation and patterning of the

central nervous system, *Development* **136**, 461-472. doi: 10.1016/j.yexcr.2013.11.018

**Streit, A., Berliner, A. J., Papanayotou, C., Sirulnik, A. and Stern, C. D.** (2000). Initiation of

neural induction by FGF signalling before gastrulation. *Nature* **406**, 74-78. doi:

10.1038/35017617

**Sudou, N., Yamamoto, S., Ogino, H. and Taira, M.** (2012). Dynamic in vivo binding of

transcription factors to cis-regulatory modules of *cer* and *gsc* in the stepwise formation of the

Spemann-Mangold organizer. *Development* **139**, 1651-1661. doi: 10.1016/j.celrep.

2019.05.013

**Suzuki, A., Yoshida, H., van Heeringen, S.J., Takebayashi-Suzuki, K., Veenstra, G.J.C. and**

**Taira, M.** (2017). Genomic organization and modulation of gene expression of the TGF-β

and FGF pathways in the allotetraploid frog *Xenopus laevis*. *Dev. Biol.* **426**, 336-359. doi:

10.1016/j.ydbio.2016.09.016

**Wasylk, B., Hagman, J. and Gutierrez-Hartmann, A.** (1998). Ets transcription factors:

nuclear effectors of the Ras-MAP-kinase signaling pathway. *Trends Biochem. Sci.* **23**,

213-216. doi.org/10.1016/S0968-0004(98)01211-0

**Wasylk, C., Maira, S. M., Sobieszczuk, P. and Wasylk, B.** (1994). Reversion of Ras

transformed cells by Ets transdominant mutants. *Oncogene* **9**, 3665-3673.

**Wilson, S. I., Graziano, E., Harland, R., Jessell, T. M. and Edlund, T.** (2000). An early

requirement for FGF signalling in the acquisition of neural cell fate in the chick embryo.

*Curr. Biol.* **10**, 421-429. doi: 10.1016/S0960-9822(00)00431-0

**Wilson, S. I. and Edlund, T.** (2001). Neural induction: toward a unifying mechanism. *Nat.*

*Neurosci.* **4**, 1161-1168. doi: 10.1038/nn747

**FIGURE LEGENDS**

**Fig. 1. Suppression of anterior neural development *in vivo* by blocking Fgf/Ets signalling**

**in ectoderm cells.**

(A) Structural features of Ets proteins and their dominant-negative derivatives.

(B) Injection protocol.

(C) Lateral views of uninjected embryos at stage 35/36.

(D–J) Embryos were injected with synthetic RNA according to the respective protocol

illustrated in (B), reared until stage 35/36, and photographed. Lateral views of embryos

exhibiting typical phenotypes are shown. In (I) and (J), the lower panels are magnified

versions of the upper ones, highlighting the anterior head region. Scale bar = 1 mm. The

number of embryos exemplified in the photograph over the total number analysed is displayed

on each panel.

**Fig. 2. Suppression of *sox2* and *otx2* expression *in vivo* by blocking Ets transcriptional**

**activity in ectoderm cells.**

(A) Experimental procedure highlighting the injection and bisection protocols.

(B–F) Embryos were injected with *Δhelk1* RNA (45 pg/blastomere) plus the lineage tracer *gfp*

RNA (20 pg/blastomere). Controls were not injected. In (D) and (E; right panel), wild-type

*xets1* RNA (5.5 pg/blastomere) was co-injected. RNA *in situ* hybridisation was carried out on

stage 11 gastrula embryos. Scale bar = 0.25 mm. The number of embryos exemplified in the

photograph over the total number analysed is displayed on each panel.

**Fig. 3. Suppression of Fgf-induced activation of position-specific neural marker genes in**

**cultured ectoderm cells by blocking Ets transcriptional activity.**

(A) Experimental design for the primary culture of ectoderm cells. Cultured cells were

harvested for mRNA isolation when control embryos reached stage 25 (for B–E) or stage 14

(for S–U).

(B–E) PCR products of six position-specific neural marker gene mRNAs are indicated in (B),

each co-amplified with *ef1α* mRNA (internal control). Ectoderm cells were prepared from

embryos that had not been injected (D) or injected with *ΔΔxetv1* RNA (B), *Δxets1* RNA (C), or

*Δxfgfr-4a* RNA (E) (63 pg/dorsal-animal blastomere and 105 pg/ventral-animal blastomere for

each injection).

(F) A quantitative comparison of the dose-response profiles of marker transcript levels. Data

from control experiments (B) and (D) are combined. The ratio of the band intensities of the

markers that of *ef1α* plotted against Fgf2 doses on a semi-log graph. The percentage of the

maximum value of the ratio is presented in each profile.

(G–L) Data from (B) and (C) are quantified and presented as in (F).

(M–R) Data from (D) and (E) are quantified and presented as in (F).

(S–U) PCR products of *sox2* and *otx2* mRNA, each co-amplified with *ef1α* mRNA (internal

control). Ectoderm cells were prepared from embryos that had been injected with

*enr·Δxetv1*RNA (S; 40 pg/dorsal-animal blastomere and 50 pg/ventral-animal blastomere) or

*Δhelk1·enr* RNA (U; 4 pg/dorsal-animal blastomere and 5 pg/ventral-animal blastomere). For

(T), ectoderm cells were prepared from uninjected embryos.

**Fig. 4. Suppression of anterior head development *in vivo* by depleting Fgf2 or Fgf8 in**

**ectoderm cells.** Antisense morpholino oligo (MO) or 5mis-MO was injected into a single

dorsal-animal blastomere at the 8-cell stage. Injected embryos were cultured until stage 35/36.

Scale bar = 1 mm.

(A–D) Upper panels show a dorsal view of the anterior head of an injected embryo. Middle

panels show a lateral view of the whole body of two injected embryos. Lower panels show a

lateral view of the anterior head region of another injected embryo at a higher magnification.

Broken lines in the upper and lower panels represent the assumed mid-line of affected

embryos.

(E–G) Lateral views of the anterior head region of two injected embryos are shown.

(H) Numerical data were collected from the experiments exemplified in (A–G) and displayed

for comparison.

(I) Fgf2 MO was injected with no additives, (J) with cognate *fgf2* RNA, or (K) cognate *fgf8*

RNA.

(L) Fgf8 MO was injected with no additives, (M) with cognate *fgf2* RNA, or (N) cognate *fgf8*

RNA.

(O) Fgf2 MO or (P) Fgf8 MO was injected without (left panels) or with (right panels) wild-type

*xetv1* RNA. In (I–P), the number of embryos exemplified in the photograph over the total

number analysed is displayed on each panel.

**Fig. 5. Suppression of *otx2* and *sox2* expression *in vivo* by depleting Fgf2 or Fgf8 in**

**ectoderm cells.**

(A) Injection protocol. We poked a single dorsal-animal blastomere at three separate sites to

obtain a more global MO distribution in the progeny of the injected blastomere.

(B–D) The upper panel embryos were injected with Fgf2 MO (left panel) or 5mis-Fgf2 MO

(right panel) at a dose of 1.8 ng/blastomere, while embryos in the lower panels were injected

with Fgf8 MO (left panel) or 5mis-Fgf8 MO (right panel) at the same dose.

(E) Embryos in the left panels were injected with Fgf2 MO (2.4 ng/blastomere) with or without

cognate *fgf2* RNA (0.15 ng/blastomere), while embryos in the right panels were injected with

Fgf2 MO (2.4 ng/blastomere) with or without wild-type *xetv1* RNA (20 pg/blastomere).

(F) In the left panels, Fgf8 MO (2.4 ng/blastomere) was injected with or without cognate *fgf8*

RNA (0.065 pg/blastomere), while in the right panels, Fgf8 MO (2.4 ng/blastomere) was

injected with or without wild-type *xetv1* RNA (20 pg/blastomere). Scale bar = 0.25 mm. The

number of embryos exemplified in the photograph over the total number analysed is displayed

on each panel.

**Fig. 6. Fgf8/Ras/Mapk/Ets signalling in cultured ectoderm cells activates *fgf8*, *sox2*, and**

**anterior neural genes in an autocrine manner.**

(A) Time course of *fgf8* and anterior neural gene activation in cultured ectoderm cells. Ectoderm

cells were prepared from stage 10 gastrula embryos and cultured without Fgf. They were

harvested serially after the start of the culture and processed for RT-PCR. Relative amounts of

transcripts from each gene are plotted against the time of harvest.

(B–D) Suppression of autonomous *fgf8* and neural gene activation in ectoderm cells by

depleting Fgf8 (B) by adding inhibitors of the Fgf/Ras/Mapk pathway (C) or by blocking Ets

transcriptional activity (D). (B) Ectoderm cells were prepared from stage 10 embryos that had

been injected with Fgf8 MO or 5mis-Fgf8 MO (4.8 ng/dorsal-animal blastomere and 6.0

ng/ventral-animal blastomere) at the 8-cell stage, as described in Fig. 3A. (C) Ectoderm cells

were prepared from intact stage 10 embryos and cultured in the presence of SU5402 (400

μM) or U0126 (200 μM). (D) Ectoderm cells were prepared from stage 10 embryos that had

been injected with *enr·Δxetv1* RNA (40 pg/dorsal-animal blastomere and 50

pg/ventral-animal blastomere) or *Δhelk1·enr* RNA (4 pg/dorsal-animal blastomere and 5

pg/ventral-animal blastomere) at the 8-cell stage. After culturing without the addition of Fgf

until control sib-embryos reached stage 14, they were harvested and subjected to RT-PCR.

The band intensity was analysed using ImageJ. Relative amounts of transcripts from each

gene were compared.

**Fig. 7. Fgf/Ets signalling in cultured ectoderm cells activates position-specific early neural**

**marker genes in a dose-dependent manner.**

(A) Expression pattern of *sox2* and the position-specific early markers at stage 11. Scale bar =

0.25 mm.

(B–D) The experimental design was as described in Fig. 3A. *enr·Δxetv1* RNA and *Δhelk1·enr*

RNA were injected as in Fig 6D. Cultured cells were harvested when control embryos

reached stage 14. The band intensities were analysed using ImageJ.

(E) A quantitative comparison of the dose-response profiles of transcript levels of the four

markers. Data in (C) were quantified.

(F–H) Suppression of Fgf-induced activation of the position-specific early markers in cultured

ectoderm cells by ΔhElk1·EnR or EnR·ΔxEtv1. Data in (B), (C), and (D) were quantified.

**Fig.1**

833 **Fig.2**

**Fig.3**

| H | Injected Fgf MO (per blastomere) | Fgf2 MO (1.8 ng) | 5mis Fgf2 MO (2.5 ng) | Fgf3 MO (4.9 ng) | Fgf4 MO (3.7 ng) | Fgf8 MO (2.5 ng) | 5mis Fgf8 MO (3.7 ng) | Fgf9 MO (4.9 ng) |
|---|---|---|---|---|---|---|---|---|
| | Embryos with severe head defects | 25/29 (86%) | 1/30 (3%) | 2/37 (5%) | 4/33 (12%) | 17/20 (85%) | 2/28 (7%) | 2/27 (7%) |

Fig.4

**Fig.6**

**Fig.7**

43