# Proposed allosteric inhibitors bind to the ATP site of CK2α

Paul Brear[1*], Darby Ball[2], Katherine Stott[1], Sheena D'Arcy[2] and Marko Hyvönen[1*]

[1] Department of Biochemistry, University of Cambridge, UK

[2] Department of Chemistry and Biochemistry, The University of Texas at Dallas, USA

[*] Correspondence to Paul Brear (pdb47@cam.ac.uk) and Marko Hyvönen (mh256@cam.ac.uk)

**Keywords:** CK2α, inhibitor, ATP-competitive, kinase, biophysics

## Abstract

CK2α is a ubiquitous, well-studied protein kinase that is a target for small molecule inhibition, for treatment of cancers. While many different classes of ATP-competitive inhibitors have been described for CK2α, they tend to suffer from significant off-target activity and new approaches are needed. A series of inhibitors of CK2α has recently been described as allosteric, acting at a previously unidentified binding site. Given the similarity of these inhibitors to known ATP-competitive inhibitors, we have investigated these further. In our thorough structural and biophysical analyses, we have found no evidence that these inhibitors bind to the proposed allosteric site. Rather, we report crystal structures, competitive ITC and NMR, HDX mass spectrometry and chemoinformatic analyses that all point to these compounds binding in the ATP pocket. Our crystal structures however do show that the proposed allosteric site can bind ligands, just not those in the previously described series. Comparison of our results and experimental details with the data presented in the original report suggest several reasons for the disparity in our conclusions, the primary reason being non-specific inhibition by aggregation.

## Introduction

Potent inhibitors of kinases have existed for some time and the majority of these bind in the active site of the kinase, competing with the co-factor ATP[1,2]. Because of the significant conservation of the ATP binding site, kinase inhibitors often suffer from poor selectivity[3–7]. This poor selectivity is a disadvantage for several reasons. Firstly, promiscuous inhibitors tend to manifest off-target toxicity if used therapeutically and, to combat this, end up being used at suboptimal concentrations[8,9]. Secondly, they are poor chemical tools as they cannot provide unambiguous answers to target validation questions. In order to develop more selective inhibitors of kinases, a promising strategy is to target sites outside the highly conserved ATP pockets. This leads to increased selectivity as these sites are structurally not constrained by the shared property of binding to ATP and therefore tend to be less conserved among different kinases[10–13]. When developing inhibitors that bind outside the primary ATP site, it is vital that the new sites are thoroughly validated biochemically, biophysically, and structurally

to confirm the inhibitor binding mode unambiguously. Often inhibitors are identified as being

allosteric if they are shown to be non-competitive in biochemical assays. However, this data can be

misinterpreted, as non-specific inhibitors also often show similar non-competitive behaviour[14]. This

problem can be further compounded by moderate/weak affinity of the hits being investigated, as they

must be used at higher concentrations, which makes secondary binding more likely. Therefore,

apparent allosteric behaviour must be very carefully validated to ensure that vital time and resources

are not wasted on optimising these inhibitors further.

CK2α has a long history as a target for drug discovery, due to its ubiquitous role in multiple diseases

such as cancer and fibrosis[15–17]. CK2α is an unusual kinase in that it is constitutively active and does

not require phosphorylation for activation[18]. It is also a highly promiscuous kinase with hundreds of

recorded cellular substrates.[19] Inhibition of CK2α has been achieved mostly through an ATP-

competitive mechanism and a number of high affinity inhibitors of distinctly different chemotypes

are known and characterized structurally[17,20–27]. Despite significant effort, most inhibitors have poor

selectivity, although claims are often made to the contrary[28–32]. For example, the clinical trial

candidate CX-4945 inhibits at least ten other kinases with nanomolar affinity (see Table S1 in Brear

et al.[25]).

We have recently demonstrated a new approach to CK2α inhibition.  By using a cryptic so-called αD

pocket just below the active site as an anchoring point, we have been able to develop highly

selective inhibitors of CK2α[25]. The most potent of these is the most specific CK2α inhibitor reported

to date, CAM4066[25]. The fragment-based development of these inhibitors was supported by a

significant amount of structural data to identify the optimal αD site anchor moiety and subsequent

growth of the fragment to the ATP site[27]. In addition to active site targeting, another approach is the

inhibition of CK2α binding to its scaffolding partner CK2β. This method has the potential to generate

specific inhibitors that affect only a subset of CK2 substrates[24,33] and has led to some initial

encouraging success that requires further development[24,33,34].

Recently, a series of CK2α inhibitors (Series A, SI Fig. 1), proposed to bind in a novel site (Site A, SI

Fig. 2) just outside of the highly conserved ATP site, has been published[35,36]. The authors used

enzymatic assays, native mass spectrometry and competitive ligand-based NMR studies to

characterize the mode of inhibition. They interpret their data as supporting an allosteric mechanism.

While their data were presented as supporting non-ATP-competitive inhibition, there were several

reasons why we speculated that the mechanism of inhibition by these molecules was not necessarily

allosteric. In particular: the enzyme assays showed mixed mode inhibition; ATP analogues were

replaced in NMR studies; high affinity ligands showed unexpected signal in ligand-based NMR

experiments; and, mutations close to the ATP site affected inhibitory potential [35,36]. Our motivation

to investigate these inhibitors in more detail arose from the somewhat conflicting data presented,

[35,36] as well as the chemical properties of these new inhibitors, which are not unlike known ATP

competitive kinase inhibitors.

We have used several orthogonal biophysical and structural methods for the characterization of the

binding mode of these new CK2α inhibitors and we find no evidence for interaction with the

proposed allosteric site (Site A). We show, using crystal structures, competitive ITC and NMR,

hydrogen-deuterium exchange mass spectrometry (HDX) and computational analyses of the ligand

structures, that these molecules are clearly binding to the ATP site of the kinase. It thus seems likely

that most of the inhibitory effect, if not all, comes from traditional type I inhibition of the kinase

through orthosteric competition with ATP.

## Results

### Chemoinformatic analysis

The structures of the compounds presented in the original papers and their comparison to other

validated kinase and CK2α inhibitors raised concerns over the claim that these compounds had a

novel binding mode. The conserved scaffold of the compounds shares significant similarity to many

established and well validated kinase inhibitors that bind in the ATP site (Fig. 1)[27,37–63]. This scaffold

consists of a central thiazole ring that is connected directly to a phenyl group on one side and to a

benzoic acid moiety through an amine on the other side (Fig. 1A).  We have used two representative

compounds from Series A for further investigation, which we denote **1** and **2** (these were referred to

as **6** and **27** respectively in the original papers[35,36]). **1** was chosen as much of the validation of Series

A and Site A relied upon this compound[35]. **2** was chosen as it was the highest affinity inhibitor

reported in Series A.[36] A search of the Protein Data Bank (PDB, http://www.pdb.org) for ligands that

bind to serine/threonine kinases (EC=2.7.11.1) and contain a benzoic acid group, which is a critical

part of the inhibitors in question (Fig. 1B), identified 49 molecules. Of these, 46 bind in the ATP site

and the remaining three bind in interface sites that do not exist in CK2α (SI Table 1). Narrowing the

search by inclusion of the amine that links the benzoic acid to the thiazole ring identified 15 ligands

(SI Table 2). Of these ligands, all 15 bind in the ATP site. When a similar search was performed on

CK2α, 14 ligands were identified, all of which bound in the ATP site (SI Table 3). Four representative

structures of these compounds are shown in Fig. 1C. All of these inhibitors display the same binding

mode where the binding to CK2α is dominated by the interaction between the carboxylic acid and

the ζ-amino group of Lys68 (Fig. 1C). Further refining the search to include only compounds that also

bind in the hinge region, similar in structure to the thiazole-amine group of **1**, identified eight

inhibitors (SI Table 4). These compounds, including **CX4945**, an extensively validated ATP-

competitive CK2α inhibitor currently in clinical trials, interact with the hinge and Lys68[47]. Overlaying

the 2D structure of **1** with **CX4945** shows that **1** could replicate the binding mode of **CX4945** (SI Fig 2)

and bind in the ATP site with its carboxylic acid interacting with Lys68 and amine hydrogen bonding

with the hinge (Fig 1C). Indeed, modelling of **1** into the ATP site of CK2α  replicates the

crystallographic binding mode of **CX4945** (SI Fig 2). The carboxylic acid is interacting with Lys68 and

an ATP site water as predicted and the hydrophobic core of the molecule is sandwiched between

Met163 and Val66. The amine is not predicted to interact with the hinge. However, the benzyl group

is predicted to bind in a similar position to the benzyl chloride of **CX4945**.

Further evidence in support of our hypothesis that **1** may bind in the ATP site comes from the

analysis of a fragment-based X-ray crystallographic screen against CK2α[27] previously performed at

our lab. One purpose of fragment-based X-ray crystallographic screening is to use the fragments to

probe the target protein and identify binding hotspots[64]. From the screen of 354 fragments, 21 were

identified as binding to CK2α. All 21 bound in the ATP site and interacted with Lys68[27]. 19 of the

CK2α-binding fragments were structurally related to **1** and contained a carboxylic acid moiety that

mediated this interaction (SI Fig. 3). The preference of CK2α for binding ligands with a carboxylic acid

in this position is further supported by the serendipitous binding of acetate. Acetate ions can be

utilised as a very low molecular weight probe to identify binding sites for carboxylic acids[64]. In CK2α

structures deposited to the PDB, at least nine binding sites for acetate can be identified (SI Fig. 4).

Among these sites are the Lys68 hotspot in the ATP site, the CK2β interface and the substrate-

binding pocket. Notably, none of the nine sites overlaps with Site A, the proposed binding site of **1**.

When such a conserved binding mode is observed for an ATP site-binding pharmacophore, it seems

unlikely that inhibitors with significant structural similarity would not also bind to the ATP site. These

observations led us to initiate additional studies to investigate the binding mode of these inhibitors

using two compounds (**1** and **2**, Fig. 1A) from the original publications.

**Figure 1. Conserved pharmacophore binding in the ATP site of protein kinases** (A) The structures of **1** and **2**, denoted **6** and **27** respectively in the original studies. (B) A summary of the analysis of the established ligands that bind to Ser/Thr protein kinases (EC number 2.7.11.1) or CK2α in the ATP site. (C) The structures of the four CK2α inhibitors and their protein-bound crystal structures which contain a benzoic acid group that binds in the ATP site[27,37,46–55,38,56–63,39–45].

 X-ray crystallographic analysis

The original publication contained no structural data on the inhibitor binding mode. In fact, the

authors state that 'cocrystallisation attempts aimed at elucidating the non-ATP competitive binding

mode of 2-aminothiazole derivatives with CK2α were not successful'[35,36]. We have rectified this and

have determined structures of both **1** and **2** in complex with CK2α (Fig. 2, SI Table 5). These

structures were obtained by soaking the compounds into CK2α crystals at 10 mM for 16 hours. The

crystals diffracted at less than 2 Å resolution and unambiguous positive difference electron density

corresponding to both **1** and **2** can be clearly seen in the ATP site before the ligands were included

(Fig. 2A and 2B, SI Fig. 5). As we predicted, the carboxylic acid groups of both compounds interact

directly with Lys68 like the carboxylic acid of **CX4945** (Fig. 2C and 2D). The remainder of the

molecules however, each have their own binding nuances. The amine of the benzylamine of **1**

interacts very weakly with the hinge region backbone nitrogen of Val116 via a bridging water

molecule whereas the OH group of **2** interacts with His160, via a bridging water, which pulls the

amine of the benzylamine further away from the hinge region and prevents interactions with the

hinge (Fig. 2C and 2D). The 5-membered rings from **1** and **2** stack on top of Met163 which leads to

the naphthyl ring of **2** stacking on top of His160 at the entrance to the ATP site. The observed

binding mode agrees well with the binding mode of established ATP site inhibitors of CK2α, such as

**CX4945**. No density corresponding to either **1** or **2** was observed in the proposed allosteric binding

site (Site A, SI Fig. 5), despite the high concentration of the ligand (>1,000-fold excess over $K_D$) used

for soaking. The binding mode of **1** in the ATP site of CK2α observed in the crystal structure agrees

well with that predicted by the modelling study (SI Fig. XXXX).

**Figure 2. Crystal structures of compounds 1 and 2 bound to CK2α** (A) The Fo-Fc map (green, displayed at 3σ) in the ATP site and Site A before the ligand has been placed. Refined structure of **1** (green) has been superimposed on the ATP site (PDB: 6YPJ) and the predicted binding mode of **2** (transparent purple) is shown on Site A. (B) The Fo-Fc map (green, contoured at 3 σ) for the ATP site and Site A before the ligand has been placed. Refined structure of **2** (blue) has been superimposed on the ATP site (PDB: 6YPG) and the predicted binding mode of **2** (transparent purple, from supplementary material of reference 34) is shown on Site A. (C) **1** (green) bound to CK2α, (PDB: 6YPJ) with key interacting residues and H-bonding highlighted. (D) **2** bound to CK2α (PDB: 6YPG), with key interacting residues and H-bonding highlighted. The structure of **CX4945** (PDB:3PE1, yellow) has been superimposed on the structure of **1** and **2** in panels C and D. Electron densities of the final, refined structures are shown in SI Fig. 6.

To reduce the impact that different crystal forms and packing may have on the accessibility of

binding sites and bias in the resulting complex structure, we have soaked the inhibitors into two

different crystal forms of CK2α. One is obtained using an engineered form of CK2α (referred to as

CK2α KKK/AAA) in which lysines 74-76 are mutated to alanines to facilitate crystallization. These

lysines are in the vicinity of Site A and could affect binding of ligands to this site. However, these

lysines do not affect the ATP site and we show later that both **1** and **2** bind to CK2α KKK/AAA and to

wild type CK2α with the same affinity (see ITC measurements). The second crystal form (CK2α SF)

has no mutations in the lysine-rich loop but has a substrate peptide fused to the N-terminus of the

protein, aimed at facilitating crystallization. In both crystal forms, we observe electron density for **1**

and **2** only in the ATP site, and not in the proposed allosteric site (Site A, SI Fig. 5).

The two crystal forms of CK2α have been extensively used for the determination of CK2α:ligand

complexes (66 structures currently in the PDB). More than eight different ligand binding sites can be

accessed by soaking into these crystal forms (SI Fig. 7), including the proposed allosteric site for **1**

and **2**, Site A. We have observed both ADP and GDP bound to Site A in crystal structures of wild type

CK2α (PDB:6YPN) and in the KKK/AAA mutant (PDB:6YPK) (Fig. 3). The adenosine and guanosine

rings sit in the hydrophobic pocket identified in the previous modelling work[35] and the diphosphate

interacts with Arg80 that has been identified as an acetate-binding hotspot (SI Fig. 4). This confirms

that Site A is accessible to ligands in these crystal forms and soaking with **1** or **2** would be expected

to result in clear electron density in the site, should they bind there. Indeed, ADP can be seen

simultaneously in both the ATP site and in Site A (Fig. 3D, PDB: 6YPN). These observations

demonstrate that binding in Site A does not in itself prevent binding in the ATP site and that if **1** or **2**

did bind in both sites, we would expect to see density corresponding to it. Furthermore, **1** and **2**

would be expected to bind predominantly in the highest affinity site and this site would dominate

the inhibition.  These crystal structures prompted us to perform further experiments to

unambiguously establish the binding site in solution and verify the likely mode of inhibition for these

compounds.

**Figure 3.** Crystal structures of ADP and GDP binding to CK2α Site A**.** (A) GDP binding in Site A. The Fo-
Fc map (contoured at 3σ) centered on the proposed allosteric site. No density was observed in the
ATP site for GDP therefore the density in the ATP site was not shown for clarity. The CK2α KKK/AAA
construct was used for this structure (PDB: 6YPK). B) The structure of GDP bound to CK2α, with the
extensive H-bonding network shown by dotted lines (PDB: 6YPK). C) The proposed binding mode of **1**
(PDB: 6YPJ) superimposed on the structure of GDP bound to CK2α (6PYK). D) The structure of ADP
bound simultaneously in the ATP site and in Site A of WT CK2α (PDB: 6YPN).

Hydrogen-deuterium exchange mass spectrometry

Hydrogen-deuterium exchange (HDX) allows the localization of interaction sites and conformational
changes in proteins. We have analysed the binding of four different classes of ligands to CK2α (SI Fig.
8 and 9) using HDX. These ligands were: ADP and AMPPNP-Mg$^{2+}$, both of which bind across the

entire ATP site interacting with the hinge region and Lys68 (SI Fig. 8C and 8D);[60]  **3**, which is a

fragment that binds only on the right hand side of the ATP site, interacting with Lys68[60] (SI Fig. 8E);

**4**, which is an αD site binder (SI Fig. 8G)[26] and **6** which links the αD site and the ATP site (SI Fig. 8F)[27].

We recovered 192 shared peptides from each of these HDX experiments that redundantly span

almost the entire sequence of CK2α (SI Fig. 10). This allows for unbiased detection of ligand

interaction fingerprints across the whole of the protein. Binding can be identified by observing

changes in deuterium uptake between CK2α alone and a concurrently run ligand-bound sample (SI

Table 6). We show all changes greater than ±0.25 Da that have a p-value less than 0.01 in a Welch's

t-test (Fig. 4 and SI Fig. 11). Extended data for all peptides and timepoints are shown in the

Supplementary Information (SI Fig. 11 and a separate spreadsheet). For all the ligands, a decrease in

deuterium uptake is observed at late timepoints particularly for peptides containing residues 54-111

(sample peptides shown in SI Fig. 12). Since this change occurs with all the ligands, it appears to

result from general stabilization of the region termed the 'N-lobe', on ligand binding, rather than an

effect attributable to a specific site.

Addition of ADP and AMPPNP-$Mg^{2+}$ causes clear changes in deuterium uptake in the ATP site (Fig. 4).

The largest change is a decrease in deuterium uptake for peptides covering residues 39-55 and 121-

131. Residues 39-55 correspond to the P-loop that interacts directly with the phosphates of ADP,

while residues 121-131 are the αD loop that does not interact directly with ADP or AMPPNP-$Mg^{2+}$.

Protection occurs in the latter due to the interaction of the adenosine ring with the hinge region that

directly links to and stabilizes the αD loop. These observations are similar to those previously

observed when binding of ATP analogues to other kinases has been studied[65,66]. The observed

decreases in uptake are larger for AMPPNP-$Mg^{2+}$ than ADP-$Mg^{2+}$ because AMPPNP binds with a

higher affinity than ADP. To further probe interactions in a more localized area of the ATP site, we

used **3** that binds only at the right hand side of the ATP site (SI fig 8 E), coordinated to catalytic Lys68

(PDB: 5CSP)[60] (Fig. 4). With **3**, we see a decrease in deuterium uptake in the P-loop and no change in

the αD loop. The pattern of protection within the P-loop is also distinct, indicative of different

binding contacts being used by **3** compared to the ATP analogues.  We can thus distinguish binding

in the entire ATP site, including interaction with the hinge, from binding on just one side where

Lys68 is located.

To ensure we could detect different binding sites with our HDX experiments, we used another well

characterized ligand (**4**) that binds to the αD pocket only (SI Fig. 8G)[26]. This compound shows a

markedly different HDX fingerprint compared to the ATP analogues and **3** (Fig. 4). The largest

differences are seen in the αD loop, with binding of **4** resulting in increased uptake compared to

unbound CK2α. Increased uptake in the αD loop is opposite to the decrease seen upon ADP and

AMPPNP-Mg$^{2+}$ binding. This indicates that the binding of **4** in the αD pocket puts the loop in a more

flexible or open conformation amenable to exchange, as is seen in the crystal structure of CK2α in

complex with **4** (PDB: 5ORJ).  Finally, we characterised the HDX fingerprint for **6** that occupies both

the αD pocket and the ATP site. We observed a similar increase in uptake in the αD loop as with **4**

(Fig. 4). Compounds **4** and **6** also cause decreased uptake in the P-loop, with **6** showing more

protection than **4**, similar to the protection we observed with **3**.

All these fingerprints are consistent with the known binding modalities of these ligands to CK2α and

can be used to identify the binding modes of new compounds. We performed the same experiments

to characterise the binding of compound **1** to CK2α (Fig. 4). Addition of **1** to CK2α causes decreased

uptake in the P-loop with a pattern and magnitude of change similar to the addition of AMPPNP-

Mg$^{2+}$. However, no changes are observed in the αD pocket, as was observed with the addition of **3**.

These results are consistent with the binding mode seen in the crystal structures where **1** makes

significant hydrogen bonding just to Lys68 with only weak water-mediated interaction to the hinge

region. These data do not allow us to unequivocally eliminate binding at the proposed allosteric site.

However, if the allosteric binding mode were correct, two clear differences would be expected.

Firstly, a change in the uptake by residues 67-88, part of Site A, would be expected due to the main

interaction being with the acidic group of **1** (SI Fig. 8). We do not observe changes in uptake in this

region despite it being redundantly covered by eight peptides (SI Fig. 11). Secondly, no change in the

uptake by the P-loop would be expected since the modelling does not predict significant interactions

with that part of the protein. Interactions with the P-loop, such as that seen for **1**, are one of the

major features of ATP site binders, consistent with our structural characterization of **1** and

**CAM4066**.

**Figure 4.** HDX analysis of ligand binding to CK2α. Heatmaps show the change in deuterium uptake upon binding of different ligands to apo CK2α. Each panel, with the ligand listed on the top, shows the heatmaps for 10 s and 100 s time points for peptides corresponding to the P-loop and αD-loop, in comparison to unliganded CK2α. All depicted changes are greater than ±0.25 Da and have a p-value less than 0.01 in a Welch's t-test. The exact start and end residues for each peptide are shown on the left of the panels. Next to each heatmap are two graphs showing deuterium uptake for one peptide from each P-loop (top) and αD loop (bottom) for apo CK2α (black) and CK2α with ligand (grey). The Y-axis range is 80% of max D-uptake assuming the N-terminal residue undergoes 100%

back-exchange. Data have not been corrected for back-exchange. Error bars are ±2σ derived from

three technical replicates. A structural diagram in between the uptake plots has the P-loop and αD

loop coloured purple and green, respectively, and the bound ligand in surface rendering in yellow.

Full HDX data are shown in SI Fig. 11 and available in the supplementary spreadsheet.

Ligand-observed NMR analyses

To further explore the binding of the new compounds to CK2α, we have used various ligand-based

NMR methods, similar to the experiments in the original characterization of these molecules. We

were particularly concerned about the evidence for the simultaneous binding of **CX4945** and **1** due

to the presence of strong signals for **CX4945** in the STD NMR spectra. In order to observe a signal in

an STD spectrum, the saturation of protein resonances has to be transferred to the bound ligand,

which then dissociates to join the pool of bulk ligand in solution before it can be detected. This

means that the size of the signal for a given ligand is proportional to the amount of the ligand that

binds to *and* is released by the protein during the saturation step of the STD experiment. Therefore,

ligands with relatively fast association ($k_{on}$) and dissociation ($k_{off}$) rates give the largest signal.[67,68]

We have determined the off rate of **CX4945**, that binds to CK2α with low nanomolar to picomolar

affinity[47] , from CK2α to be 0.0039 $s^{-1}$ by SPR (SI Fig. 12). When ligands have $k_{off}$ rates <0.1 $s^{-1}$, the

saturation cannot be transferred effectively to bulk ligand in solution resulting in no observable STD

effect[68]. The experimentally determined off-rate for **CX4945** means that less than 1% of **CX4945**

would have dissociated after the 2-second saturation time used in the NMR pulse sequence. This

would mean that the concentration of saturated ligand in solution would be less than 0.1 μM. This

difference would not be observable by NMR.

To resolve these issues, we have repeated the STD NMR studies reported in the original publication

[35] and used two additional ligand-based NMR experiments, CPMG and Water-LOGSY[67],  for further

validation. Our STD experiment shows that **1** binds to CK2α, as revealed by detectable transferred $^1$H

intensities (Fig. 5A(i)). However, when the same experiment is run in the presence of **CX4945**, a

significant reduction of the STD signals from **1** are observed, indicating direct competition between

the two ligands (Fig. 5A(ii)). In the original paper, strong **CX4945**-derived signals are also clearly

visible. The signals - which are comparable in magnitude to the much weaker binding ligand **1** - could

instead arise from non-specific binding of **CX4945** to lower affinity sites on CK2α (as was shown by

native mass spectrometry[35]), precipitation of **CX4945** leading to false positive signals in the

experiment or binding of CX4945 to soluble aggregates of **1**.

Given the disparity in the STD NMR experiments between our results and those in the original

report, we have also used CPMG and Water-LOGSY experiments. These exploit different NMR

phenomena (relaxation rates, transfer of saturation) to observe the binding of the ligand to the

protein[67]. Using three different methods to observe the binding of the ligand reduces the chance of

experimental anomalies interfering with the outcome of the experiment. The Water-LOGSY

experiment shows clear binding of **1** to CK2α and complete inhibition of binding in the presence of

**CX4945** (Fig. 5B). Similarly, in the CPMG experiment, an increase in the signal from the aromatic

protons of **1** was observed in the presence of **CX4945** indicating a reduction in binding of **1** to CK2α

(Fig. 5C). In conclusion, all three ligand-observe NMR experiments show binding of **1** to CK2α and in

all three experiments binding is abolished by **CX4945**, suggesting direct competition between these

two ligands as they bind to the same site.

**A**

(i) +0 µM **CX4945**

(ii) +50 µM **CX4945**

**B**

(i) +0 µM **CX4945**

(ii) +50 µM **CX4945**

**C**

(i) +0 µM **CX4945**

(ii) +50 µM **CX4945**

0.5 mM **1** +7 µM CK2α
0.5 mM **1**

**Figure 5. Ligand-based NMR experiments to detect binding of 1 to CK2α and competition by CX4945**. All panels are overlays of spectra recorded in the presence (black) and absence (green) of CK2α. (A) Saturation Transfer Difference (STD) $^1$H spectra of **1** (i) in the absence of **CX4945**, and (ii) in the presence of 50 μM **CX4945**. (B) WaterLOGSY spectra of **1** (i) in the absence of **CX4945**, and (ii) in the presence of 50 μM **CX4945** (ii). (C) Relaxation-edited (CPMG) $^1$H spectra of **1** (i) in the absence of **CX4945**, and (ii) in the presence of 50 μM **CX4945** (ii). (NB: the signals at ca. 7.7 ppm are in fact two overlapping doublets and therefore small changes to either chemical shift results in an apparent change to the multiplet structure.)

To further confirm the validity of our results, we repeated the CPMG NMR study using the lower affinity ligands **3**, **5** and **CAM4066** as competitors (SI Fig. 13). **3** is a small, low affinity ligand in the ATP site and as such it is less likely to have an allosteric affect that might prevent binding at the adjacent Site A – an argument that could explain inhibition of binding to **1** in the presence of **CX4945**. Therefore, the reduction in the binding of **1** in the presence of **3** (SI Fig. 13B) is a more convincing proof that **1** is binding in the ATP site. **CAM4066** links the ATP site and the αD site and, as predicted, reduces the binding of **1** to CK2α (SI Fig. 13C). To rule out binding of **1** to the αD site, we have also used **5**, an optimized fragment with high affinity for the αD site. This should not prevent binding in the ATP site. Indeed, ligands similar to **5** have been shown to bind simultaneously with ATP site fragments by X-ray crystallography (PDB:6EHK)[26]. As predicted, and unlike with **CX4945**, **3** and **CAM4066**, the αD site ligand **5** has no effect on the binding of **1** to CK2α (SI Fig. 13D). Similar CPMG and STD NMR experiments were performed with the CK2α KKK/AAA mutant with the same outcomes (SI Fig. 14 and 15). These results further demonstrate that **1** binds in the ATP site.

Isothermal titration calorimetry

As a final confirmation, we have used isothermal titration calorimetry to determine affinities of **1** and **2** to CK2α and to evaluate the effect of competing ligands and CK2α mutations. This process has been successfully used to verify the binding site of novel CK2 inhibitors that bind in the αD site[26]. The affinities of compounds **1** and **2** for CK2α were determined to be 22 μM and 2.2 μM respectively (Fig. 6A and C). When the same experiments were performed in the presence of saturating concentrations of **CX4945**, no binding of either compound to CK2α was observed (Fig. 6B and 6D). This observation is in line with all our previous data and suggests that compounds **1** and **2** bind in the same site as **CX4945**. We also performed the experiment with **2** in the presence of **CAM4066,** which competes with ATP, but uses only a small benzoic acid moiety to interact with the ATP site, deriving most of its binding energy from the αD pocket. **CAM4066** abolished the binding of **2** to CK2α (SI Fig. 16). In the modelling of the binding of **1** and **2** reported previously[35,36], Lys74 was proposed to form a vital part of the binding site and to interact with the carboxylic acid of **1** and **2**. Therefore, mutations

to this residue should reduce the affinity of **1** and **2** for CK2α. To test this, we repeated the ITC

experiments using the construct CK2α KKK/AAA, in which three adjacent lysine residues (Lys74-76)

have been mutated to alanine. These mutations had a negligible effect on the affinity of **1** or **2** for

CK2α as we determined near identical affinities of 27 µM and 3.3 µM, respectively (SI Fig. 16). The

binding of **1** and **2** to CK2α KKK/AAA was also fully inhibited by the presence of **CX4945** and

**CAM4066** respectively (SI Fig. 16). These results are further evidence that the ATP site, not the

proposed allosteric site, is the primary binding site for **1** and **2**.

**Figure 6. ITC measurement of binding of 1 and 2 to CK2α.** (A) Titration of 1 into 10 µM CK2α

Titration of 2 into 10 µM CK2α. (B) Titration of **1** into 10 µM CK2α Titration of **2** into 10 µM CK2α in

the presence of **CX4945** (C) Titration of **2** into 20 µM CK2α. (D) Titration of **2** into 20 µM CK2α in the

presence of 100 µM **CX4945**.

Discussion

We present compelling and consistent evidence that **1** and **2** bind in the ATP site of CK2α, not in Site

A, as was previously proposed.[35,36] We see no evidence for binding in Site A. This is as expected,

based on our analysis of the structural data of similar kinase inhibitors. In the original study of Series

A,[36] five main pieces of evidence were used to verify a novel binding mode. This evidence included:

competitive native mass spectroscopy studies, ligand-based NMR, thermal shift analysis and

biochemical assays. While we were preparing this manuscript for publication, the Niefind group

published their analysis of the same series of compounds, which also showed binding to the ATP site

by X-ray crystallography.[69] While they have not done other biophysical analyses on the binding, they

do show by enzymatic assay that these inhibitors are ATP competitive, consistent with our data.

One of the cornerstones of the theory of allosteric inhibition was the observation that that **CX4945**

binds CK2α simultaneously with **1**, and that **1** competes with inositol hexaphosphate (IP$_6$, phytic

acid), shown by mass spectrometry. The data presented indicate that at least three molecules of **1**

bind to CK2α. In the presence of **CX4945** this is reduced to two binding sites, but no data were

presented to confirm the identity of these sites in which **1** was binding when it bound

simultaneously with **CX4945** (for example mutational studies combined with native mass

spectrometry). Some of these sites are expected to be non-specific, low affinity sites and the mass

spectrometric data does not confirm which is the primary binding site of **1.** The first evidence

presented for the location of the binding site of these compounds is the observation that in the

presence of **1** the complex between $IP_6$ and $CK2\alpha$ is not observed in the native mass spectra. This led

to the theory that **1** binds in the proposed site as a crystal structure was published by Lee et al

(PDB:3W8L)[70] of $IP_6$ apparently binding to $CK2\alpha$ in this site. However, the binding site of $IP_6$

presented in that crystal structure is more controversial than is discussed in the publication.[70] In this

structure, $IP_6$ is found at a crystal contact where half of the site is composed of residues from a

symmetry related $CK2\alpha$ molecule in the crystal lattice (SI Fig. 17). Therefore, $IP_6$ could in theory bind

to both these sites. From this crystal structure the binding site could be located around Lys74 or

His276. However, as the site is only fully formed in a crystal lattice (SI Fig. 17), $IP_6$ may not in actual

fact bind to $CK2\alpha$ at either of these sites in solution. $IP_6$ is also seen to bind weakly at a site centered

on His236 in several other crystal structures (PDB:5OQU, 5ORJ, 5MPJ, 5CVH)[26,27,60]. The native mass

spectrometric competition studies with **CX4945** or $IP_6$ do not therefore unambiguously confirm the

binding mode of **1**.

Another cornerstone of the allosteric site binding was the observation of the binding by ligand-based

NMR and simultaneous detection of **CX4945** binding, interpreted as lack of competition. As

discussed before, a high affinity ligand like **CX4945**, with slow dissociation rate should not yield

detectable signal in these experiments.  The binding of **CX4945** observed in those STD experiments is

however more likely to be due to an artefact in the experimental setup. This could be through a

number of mechanisms including aggregation of the ligand or binding to non-specific lower affinity

binding sites outside the ATP site.[71] These lower affinity sites would have higher on/off rates than

the ATP site and therefore show a larger signal in the STD spectrum. Non-specific and low affinity

sites can have a confusing influence on competition studies. This is a particular problem with the

NMR studies as ligand-observed NMR is optimized for studying low affinity sites and requires ligands

with rapid on/off rates in order for the signal (and therefore binding) to be observed. Therefore,

blocking the primary (high affinity) site may not remove the binding signal arising from other sites.

The competition studies in the original papers were done at 500 µM ligand concentration, well

beyond the observed affinities or $K_i$s of **1** and **2** and in a range where non-specific binding would be

more likely to occur.[35,36] The binding signals observed for **CX4945** in the STD spectra cast some doubt

on the conclusions drawn about the simultaneous binding of **1** and **CX4945.** This is because the

signals observed for **CX4945** cannot be due to binding in the ATP site as the dissociation rate for this

site is too slow. The major difference between our experimental system and the system reported in

the literature is the DMSO content. In brief, in the procedure reported here stocks of **1** and **2** were

made in $d_6$-DMSO and diluted 20-fold in the buffer. This was done to ensure the ligands were fully

soluble in the experiment and led to a final $d_6$-DMSO content of 5% (which was also used to provide

the field-frequency lock signal). When we first attempted to run the experiment without DMSO, we

observed precipitation of the ligands.  As aggregation can lead to false positives in STD experiments

it was decided to use DMSO in all our experiments.[71] However, in the original experiment $D_2O$ was

used for the lock and no DMSO was used to solubilise the ligand. The difference in the solvent

conditions may mean that **1** and **2** were closer to the limits of their solubility in the original

experiments and thus more likely to form aggregates. Aggregates are common causes of false

positives in biochemical assays and are often missed as they may not be easily visible to the naked

eye.  These aggregates may lead to the signals observed in the STD for **CX4945** and **1**.

The biochemical assays performed previously showed that the compounds exhibited mixed mode

non-competitive behavior and led to the understandable conclusion that they do not bind in the ATP

site. However, this is contradictory to the binding mode we see in the crystal structures.  A proposed

explanation for the non-competitive inhibition observed in the assay comes from the thermal

stability data also presented in the original paper.[35] The data show that the thermal stability of the

proteins decreases in the presence of **1** and **2**. This thermal de-stabilization could be due to a

number of different mechanisms caused by specific and non-specific effects. These mechanisms

include; breaking up of protein complexes by specific binding to the protein; binding to the unfolded

or partially unfolded protein; the formation of aggregates that bind the protein and unfold it and

specific binding that traps the protein in a less stable conformation[72–75].

It is impossible that the first mechanism is operating in this example as CK2α is present as a

monomer. The other mechanisms however may explain the conflicting results presented. All

proteins exist in an equilibrium between the native and unfolded form of the protein.[76] For the

majority of ligands, the ligand binds to the native form of the protein which stabilizes the folded

form leading to a positive thermal shift. If the ligand binds to the unfolded or partially unfolded form

of the protein and stabilize the unfolded form this could lead to a destabilization of the protein. This

would in turn lead to a decrease in the $T_m$. Binding to the unfolded protein can occur by two

mechanisms, the first mechanism is by the formation of colloidal aggregates of the small molecules,

onto which the protein would then be sequestered.[75] The aggregates are hydrophobic and

preferentially bind the unfolded or partially unfolded form of the protein where the hydrophobic

core is exposed. Therefore, a negative shift in the $T_m$ is observed. This is one of the most common

forms of non-specific inhibition that plague biochemical assays and would be seen in assays as

noncompetitive inhibition. The second mechanism occurs when the ligand monomers bind to a

partially unfolded or fully unfolded form of the protein stabilizing that form. It may be possible for

the binding to the partially unfolded form to be a specific form of inhibition.

Modelling experiments proposed Site A as the binding site for **1** and **2** (SI Fig. 8)[35]. We have

accumulated extensive evidence that these compounds bind in the ATP site. However, Site A as a

general ligand binding site has been verified by our structures of ADP and GDP bound in this

location.  This suggests that Site A could be utilized for the development of novel inhibitors of CK2α

and it also demonstrates that this site is accessible for ligand binding in our crystal forms. If this site

were to be successfully targeted by inhibitors, it would require extensive structural and biochemical

validation to confirm that compounds are really binding in the new site and not in one of the several

known small molecule binding sites in CK2α.

We believe that the data we have presented here leads to the conclusion that the compounds

represented by **1** and **2** bind to the ATP site and will inhibit CK2α via an orthosteric, ATP-competitive

mechanism. This diverse set of data pointing to **1** and **2** binding at the ATP site includes multiple

crystal structures, competitive ITC studies, competitive NMR studies, H/D exchange mass

spectrometry and computational analyses of the ligand structures. We also believe that the evidence

presented previously in support of an allosteric mode of inhibition for these compounds may have

been distorted by non-specific binding of the compounds to CK2α, which in turn affected the

interpretation of competition data with **CX4945.**

### Acknowledgements:
We would like to thank Dr Glyn Williams and Teodors Pantelejevs for their critical comments on the

manuscript. We are grateful for Diamond Light Source for access to beamlines i04, i04-1 and i24

(proposals: mx9537, mx14043, mx18548). We thank the X-ray Crystallographic and Biophysics

Facilities for access to instrumentation. We thank Apollo Therapeutics that funds our CK2α inhibitor

development project for allowing the use of the data on CX4945 binding kinetics for this publication.

The work in the D'Arcy group was funded by the NIH National Institute of General Medical Sciences

(GM133751) and start-up funds to S. D'Arcy.

### Structural coordinates in Protein Data Bank

477 6YPG (CK2α in complex with **2**), 6YPH (CK2α in complex with **2**), 6YPJ (CK2α in complex with **1**), 6YPK

478 (CK2α in complex with **GDP**), 6YPN (CK2α in complex with **ATP** ).

### Bibliography
(1)     Roskoski, R. Properties of FDA-Approved Small Molecule Protein Kinase Inhibitors.

*Pharmacological Research*. Academic Press June 2019, pp 19–50.

https://doi.org/10.1016/j.phrs.2019.03.006.

(2)   Wu, P.; Nielsen, T. E.; Clausen, M. H. FDA-Approved Small-Molecule Kinase Inhibitors. *Trends*
*Pharmacol. Sci.* **2015**, *36* (7), 422–439. https://doi.org/10.1016/j.tips.2015.04.005.

(3)   Smyth, L. A.; Collins, I. Measuring and Interpreting the Selectivity of Protein Kinase Inhibitors.
*J. Chem. Biol.* **2009**, *2* (3), 131–151. https://doi.org/10.1007/s12154-009-0023-9.

(4)   Davis, M. I.; Hunt, J. P.; Herrgard, S.; Ciceri, P.; Wodicka, L. M.; Pallares, G.; Hocker, M.;
Treiber, D. K.; Zarrinkar, P. P. Comprehensive Analysis of Kinase Inhibitor Selectivity. *Nat.*
*Biotechnol.* **2011**, *29* (11), 1046–1051. https://doi.org/10.1038/nbt.1990.

(5)   Drag, M.; Salvesen, G. S. Emerging Principles in Protease-Based Drug Discovery. *Nat. Rev.*
*Drug Discov.* **2010**, *9* (9), 690–701. https://doi.org/10.1038/nrd3053.

(6)   Bain, J.; Plater, L.; Elliott, M.; Shpiro, N.; Hastie, C. J.; Mclauchlan, H.; Klevernic, I.; Arthur, J. S.
C.; Alessi, D. R.; Cohen, P. The Selectivity of Protein Kinase Inhibitors: A Further Update.
*Biochem. J.* **2007**, *408* (3), 297–315. https://doi.org/10.1042/BJ20070797.

(7)   Bain, J.; McLauchlan, H.; Elliott, M.; Cohen, P. The Specificities of Protein Kinase Inhibitors: An
Update. *Biochem. J.* **2003**, *371* (1), 199–204. https://doi.org/10.1042/BJ20021535.

(8)   Tarcsay, Á.; Keserú, G. M. Contributions of Molecular Properties to Drug Promiscuity. *Journal*
*of Medicinal Chemistry*. American Chemical Society March 2013, pp 1789–1795.
https://doi.org/10.1021/jm301514n.

(9)   Bowes, J.; Brown, A. J.; Hamon, J.; Jarolimek, W.; Sridhar, A.; Waldron, G.; Whitebread, S.
Reducing Safety-Related Drug Attrition: The Use of in Vitro Pharmacological Profiling. *Nature*
*Reviews Drug Discovery*. Nat Rev Drug Discov December 2012, pp 909–922.
https://doi.org/10.1038/nrd3845.

(10)  Lu, X.; Smaill, J. B.; Ding, K. New Promise and Opportunities for Allosteric Kinase Inhibitors.
*Angew. Chemie Int. Ed.* **2019**. https://doi.org/10.1002/anie.201914525.

(11)  Lebakken, C. S.; Reichling, L. J.; Ellefson, J. M.; Riddle, S. M. Detection of Allosteric Kinase
Inhibitors by Displacement of Active Site Probes. *J. Biomol. Screen.* **2012**, *17* (6), 813–821.
https://doi.org/10.1177/1087057112439889.

(12)  Fasano, M.; Della Corte, C. M.; Califano, R.; Capuano, A.; Troiani, T.; Martinelli, E.; Ciardiello,
F.; Morgillo, F. Type III or Allosteric Kinase Inhibitors for the Treatment of Non-Small Cell Lung
Cancer. *Expert Opin. Investig. Drugs* **2014**, *23* (6), 809–821.

https://doi.org/10.1517/13543784.2014.902934.

(13) Wu, P.; Clausen, M. H.; Nielsen, T. E. Allosteric Small-Molecule Kinase Inhibitors.
514 *Pharmacology and Therapeutics*. Elsevier Inc. December 2015, pp 59–68.
515 https://doi.org/10.1016/j.pharmthera.2015.10.002.

(14) McGovern, S. L.; Shoichet, B. K. Kinase Inhibitors: Not Just for Kinases Anymore. *J. Med.*
517 *Chem.* **2003**, *46* (8), 1478–1483. https://doi.org/10.1021/jm020427b.

(15) Ahmad, K. A.; Wang, G.; Slaton, J.; Unger, G.; Ahmed, K. Targeting CK2 for Cancer Therapy.
519 *Anticancer. Drugs* **2005**, *16* (10), 1037–1043.

(16) Ruzzene, M.; Pinna, L. A. Addiction to Protein Kinase CK2: A Common Denominator of Diverse
521 Cancer Cells? *Biochim. Biophys. Acta - Proteins Proteomics* **2010**, *1804* (3), 499–504.
522 https://doi.org/10.1016/j.bbapap.2009.07.018.

(17) Panicker, R. C.; Chattopadhaya, S.; Coyne, A. G.; Srinivasan, R. Allosteric Small-Molecule
524 Serine/Threonine Kinase Inhibitors. *Adv. Exp. Med. Biol.* **2019**, *1163*, 253–278.
525 https://doi.org/10.1007/978-981-13-8719-7_11.

(18) Kim, H.; Choi, K.; Kang, H.; Lee, S. Y.; Chi, S. W.; Lee, M. S.; Song, J.; Im, D.; Choi, Y.; Cho, S.
527 Identification of a Novel Function of CX-4945 as a Splicing Regulator. *PLoS One* **2014**, *9* (4), 1–
528 8. https://doi.org/10.1371/journal.pone.0094978.

(19) Meggio, F.; Pinna, L. a. One-Thousand-and-One Substrates of Protein Kinase CK2? *FASEB J.*
530 **2003**, *17* (3), 349–368. https://doi.org/10.1096/fj.02-0473rev.

(20) Cozza, G.; Pinna, L. a; Moro, S. Protein Kinase CK2 Inhibitors: A Patent Review. *Expert Opin.*
532 *Ther. Pat.* **2012**, *22* (9), 1081–1097. https://doi.org/10.1517/13543776.2012.717615.

(21) Quotti Tubi, L.; Gurrieri, C.; Brancalion, A.; Bonaldi, L.; Bertorelle, R.; Manni, S.; Pavan, L.;
534 Lessi, F.; Zambello, R.; Trentin, L.; Adami, F.; Ruzzene, M.; Pinna, L. A.; Semenzato, G.; Piazza,
535 F. Inhibition of Protein Kinase CK2 with the Clinical-Grade Small ATP-Competitive Compound
536 CX-4945 or by RNA Interference Unveils Its Role in Acute Myeloid Leukemia Cell Survival, P53-
537 Dependent Apoptosis and Daunorubicin-Induced Cytotoxicity. *J. Hematol. Oncol.* **2013**, *6* (1),
538 78. https://doi.org/10.1186/1756-8722-6-78.

(22) Sarno, S.; Salvi, M.; Battistutta, R.; Zanotti, G.; Pinna, L. A. Features and Potentials of ATP-Site
Directed CK2 Inhibitors. In *Biochimica et Biophysica Acta - Proteins and Proteomics*; Biochim
Biophys Acta, 2005; Vol. 1754, pp 263–270. https://doi.org/10.1016/j.bbapap.2005.07.043.

(23)  Mazzorana, M.; Pinna, L. A.; Battistutta, R. A Structural Insight into CK2 Inhibition. *Mol. Cell.*
*Biochem.* **2008**, *316* (1–2), 57–62. https://doi.org/10.1007/s11010-008-9822-5.

(24)  Kufareva, I.; Bestgen, B.; Brear, P.; Prudent, R.; Laudet, B.; Moucadel, V.; Ettaoussi, M.; Sautel,
C. F.; Krimm, I.; Engel, M.; Filhol, O.; Borgne, M. Le; Lomberget, T.; Cochet, C.; Abagyan, R.
Discovery of Holoenzyme-Disrupting Chemicals as Substrate-Selective CK2 Inhibitors. *Sci. Rep.*
**2019**, *9* (1), 15893. https://doi.org/10.1038/s41598-019-52141-5.

(25)  Brear, P.; De Fusco, C.; Hadje Georgiou, K.; Francis-Newton, N. J.; Stubbs, C. J.; Sore, H. F.;
Venkitaraman, A. R.; Abell, C.; Spring, D. R.; Hyvönen, M. Specific Inhibition of CK2α from an
Anchor Outside the Active Site. *Chem. Sci.* **2016**, *7* (11), 6839–6845.
https://doi.org/10.1039/C6SC02335E.

(26)  Iegre, J.; Brear, P.; De Fusco, C.; Yoshida, M.; Mitchell, S. L.; Rossmann, M.; Carro, L.; Sore, H.
F.; Hyvönen, M.; Spring, D. R. Second-Generation CK2α Inhibitors Targeting the Ad Pocket.
*Chem. Sci.* **2018**, *9* (11), 3041–3049. https://doi.org/10.1039/c7sc05122k.

(27)  De Fusco, C.; Brear, P.; Iegre, J.; Georgiou, K. H.; Sore, H. F.; Hyvönen, M.; Spring, D. R. A
Fragment-Based Approach Leading to the Discovery of a Novel Binding Site and the Selective
CK2 Inhibitor CAM4066. *Bioorg. Med. Chem.* **2017**, *25* (13), 3471–3482.
https://doi.org/10.1016/j.bmc.2017.04.037.

(28)  Guerra, B.; Hochscherf, J.; Jensen, N. B.; Issinger, O.-G. Identification of a Novel Potent,
Selective and Cell Permeable Inhibitor of Protein Kinase CK2 from the NIH/NCI Diversity Set
Library. *Mol. Cell. Biochem.* **2015**. https://doi.org/10.1007/s11010-015-2433-z.

(29)  Prins, R. C.; Burke, R. T.; Tyner, J. W.; Druker, B. J.; Loriaux, M. M.; Spurgeon, S. E. CX-4945, a
Selective Inhibitor of Casein Kinase-2 (CK2), Exhibits Anti-Tumor Activity in Hematologic
Malignancies Including Enhanced Activity in Chronic Lymphocytic Leukemia When Combined
with Fludarabine and Inhibitors of the B-Cell Receptor Pathway. *Leukemia* **2013**, *27* (10),
2094–2096. https://doi.org/10.1038/leu.2013.228.

(30)  Dowling, J. E.; Alimzhanov, M.; Bao, L.; Chuaqui, C.; Denz, C. R.; Jenkins, E.; Larsen, N. A.;
Lyne, P. D.; Pontz, T.; Ye, Q.; Holdgate, G. A.; Snow, L.; O'Connell, N.; Ferguson, A. D. Potent
and Selective CK2 Kinase Inhibitors with Effects on Wnt Pathway Signaling in Vivo. *ACS Med.*
*Chem. Lett.* **2016**, *7* (3), 300–305. https://doi.org/10.1021/acsmedchemlett.5b00452.

(31)  Götz, C.; Gratz, A.; Kucklaender, U.; Jose, J. TF — A Novel Cell-Permeable and Selective
Inhibitor of Human Protein Kinase CK2 Induces Apoptosis in the Prostate Cancer Cell Line

LNCaP. *Biochim. Biophys. Acta - Gen. Subj.* **2012**, *1820* (7), 970–977.

https://doi.org/10.1016/j.bbagen.2012.02.009.

(32)  Pierre, F.; Chua, P. C.; O'Brien, S. E.; Siddiqui-Jain, A.; Bourbon, P.; Haddach, M.; Michaux, J.;

Nagasawa, J.; Schwaebe, M. K.; Stefan, E.; Vialettes, A.; Whitten, J. P.; Chen, T. K.; Darjania, L.;

Stansfield, R.; Bliesath, J.; Drygin, D.; Ho, C.; Omori, M.; Proffitt, C.; Streiner, N.; Rice, W. G.;

Ryckman, D. M.; Anderes, K. Pre-Clinical Characterization of CX-4945, a Potent and Selective

Small Molecule Inhibitor of CK2 for the Treatment of Cancer. *Mol. Cell. Biochem.* **2011**, *356*

(1–2), 37–43. https://doi.org/10.1007/s11010-011-0956-5.

(33)  Iegre, J.; Brear, P.; Baker, D. J.; Tan, Y. S.; Atkinson, E. L.; Sore, H. F.; O' Donovan, D. H.;

Verma, C. S.; Hyvönen, M.; Spring, D. R. Efficient Development of Stable and Highly

Functionalised Peptides Targeting the CK2α/CK2β Protein-Protein Interaction. *Chem. Sci.*

**2019**, *10* (19), 5056–5063. https://doi.org/10.1039/c9sc00798a.

(34)  Brear, P.; North, A.; Iegre, J.; Georgiou, K. H.; Lubin, A.; Carro, L.; Green, W.; Sore, H. F.;

Hyvönen, M.; Spring, D. R. Bioorganic & Medicinal Chemistry Novel Non-ATP Competitive

Small Molecules Targeting the CK2 α / β Interface. *Bioorg. Med. Chem.* **2018**, *26* (11), 3016–

3020. https://doi.org/10.1016/j.bmc.2018.05.011.

(35)  Bestgen, B.; Kufareva, I.; Seetoh, W.; Abell, C.; Hartmann, R. W.; Abagyan, R.; Le Borgne, M.;

Filhol, O.; Cochet, C.; Lomberget, T.; Engel, M. 2-Aminothiazole Derivatives as Selective

Allosteric Modulators of the Protein Kinase CK2. 2. Structure-Based Optimization and

Investigation of Effects Specific to the Allosteric Mode of Action. *J. Med. Chem.* **2019**, *62* (4),

1817–1836. https://doi.org/10.1021/acs.jmedchem.8b01765.

(36)  Bestgen, B.; Krimm, I.; Kufareva, I.; Kamal, A. A. M.; Seetoh, W. G.; Abell, C.; Hartmann, R. W.;

Abagyan, R.; Cochet, C.; Le Borgne, M.; Engel, M.; Lomberget, T. 2-Aminothiazole Derivatives

as Selective Allosteric Modulators of the Protein Kinase CK2. 1. Identification of an Allosteric

Binding Site. *J. Med. Chem.* **2019**, *62* (4), 1803–1816.

https://doi.org/10.1021/acs.jmedchem.8b01766.

(37)  Ngoei, K. R. W.; Langendorf, C. G.; Ling, N. X. Y.; Hoque, A.; Varghese, S.; Camerino, M. A.;

Walker, S. R.; Bozikis, Y. E.; Dite, T. A.; Ovens, A. J.; Smiles, W. J.; Jacobs, R.; Huang, H.; Parker,

M. W.; Scott, J. W.; Rider, M. H.; Foitzik, R. C.; Kemp, B. E.; Baell, J. B.; Oakhill, J. S. Structural

Determinants for Small-Molecule Activation of Skeletal Muscle AMPK Alpha 2 Beta 2 Gamma

1 by the Glucose Importagog SC4. *Cell Chem Biol* **2018**, *25*, 728-737.e9.

https://doi.org/10.2210/PDB6B1U/PDB.

(38)    Guerra, B.; Rasmussen, T. D.; Schnitzler, A.; Jensen, H. H.; Boldyreff, B. S.; Miyata, Y.;

606             Marcussen, N.; Niefind, K.; Issinger, O. G. Protein Kinase CK2 Inhibition Is Associated with the

607             Destabilization of HIF-1 Alpha in Human Cancer Cells. *Cancer Lett* **2015**, *356*, 751–761.

608             https://doi.org/10.2210/PDB4RLK/PDB.

609    (39)    Gazzard, L.; Appleton, B.; Chapman, K.; Chen, H.; Clark, K.; Drobnick, J.; Goodacre, S.;

610             Halladay, J.; Lyssikatos, J.; Schmidt, S.; Sideris, S.; Wiesmann, C.; Williams, K.; Wu, P.; Yen, I.;

611             Malek, S. Discovery of the 1,7-Diazacarbazole Class of Inhibitors of Checkpoint Kinase 1.

612             *Bioorg.Med.Chem.Lett.* **2014**, *24*, 5704–5709. https://doi.org/10.2210/PDB4QYE/PDB.

613    (40)    Kang, Y. N.; Stuckey, J. A.; Chang, P.; Russell, A. J. Crystal Structure of the CHK1. *TO BE Publ.*

614             https://doi.org/10.2210/PDB4FT0/PDB.

615    (41)    Lawrence, H. R.; Martin, M. P.; Luo, Y.; Pireddu, R.; Yang, H.; Gevariya, H.; Ozcan, S.; Zhu, J.-Y.;

616             Kendig, R.; Rodriguez, M.; Elias, R.; Cheng, J. Q.; Sebti, S. M.; Schonbrunn, E.; Lawrence, N. J.

617             Development of O-Chlorophenyl Substituted Pyrimidines as Exceptionally Potent Aurora

618             Kinase Inhibitors. *J. Med. Chem.* **2012**, *55* (17), 7392–7416.

619             https://doi.org/10.1021/jm300334d.

620    (42)    Xiao, B.; Sanders, M. J.; Carmena, D.; Bright, N. J.; Haire, L. F.; Underwood, E.; Patel, B. R.;

621             Heath, R. B.; Walker, P. A.; Hallen, S.; Giordanetto, F.; Martin, S. R.; Carling, D.; Gamblin, S. J.

622             Structural Basis of Ampk Regulation by Small Molecule Activators. *Nat.Commun.* **2013**, *4*,

623             3017. https://doi.org/10.2210/PDB4CFE/PDB.

624    (43)    Xue, Y.; Wan, P. T.; Hillertz, P.; Schweikart, F.; Zhao, Y.; Wissler, L.; Dekker, N. X-Ray Structural

625             Analysis of Tau-Tubulin Kinase 1 and Its Interactions with Small Molecular Inhibitors.

626             *ChemMedChem* **2013**, *8*, 1846. https://doi.org/10.2210/PDB4BTM/PDB.

627    (44)    Elkins, J. M.; Muniz, J. R. C.; Tan, I.; Leung, T.; Lafanechere, L.; Prudent, R.; Abdul Azeez, K.;

628             Szklarz, M.; Phillips, C.; Wang, J.; von Delft, F.; Bountra, C.; Edwards, A.; Knapp, S. Cdc42

629             Binding Protein Kinase Alpha (Mrck Alpha). *TO BE Publ.*

630             https://doi.org/10.2210/PDB4AW2/PDB.

631    (45)    Martin, M. P.; Zhu, J. Y.; Lawrence, H. R.; Pireddu, R.; Luo, Y.; Alam, R.; Ozcan, S.; Sebti, S. M.;

632             Lawrence, N. J.; Schönbrunn, E. A Novel Mechanism by Which Small Molecule Inhibitors

633             Induce the DFG Flip in Aurora A. *ACS Chem. Biol.* **2012**, *7* (4), 698–706.

634             https://doi.org/10.1021/cb200508b.

(46)    Battistutta, R.; Cozza, G.; Pierre, F.; Papinutto, E.; Lolli, G.; Sarno, S.; O'Brien, S. E.; Siddiqui-

Jain, A.; Haddach, M.; Anderes, K.; Ryckman, D. M.; Meggio, F.; Pinna, L. a. Unprecedented

Selectivity and Structural Determinants of a New Class of Protein Kinase CK2 Inhibitors in

Clinical Trials for the Treatment of Cancer. *Biochemistry* **2011**, *50* (39), 8478–8488.

https://doi.org/10.1021/bi2008382.

(47)  Ferguson, A. D.; Sheth, P. R.; Basso, A. D.; Paliwal, S.; Gray, K.; Fischmann, T. O.; Le, H. V.

Structural Basis of CX-4945 Binding to Human Protein Kinase CK2. *FEBS Lett.* **2011**, *585* (1),

104–110. https://doi.org/10.1016/j.febslet.2010.11.019.

(48)  Iegre, J.; Brear, P.; Baker, D. J.; Tan, Y. S.; Atkinson, E. L.; Sore, H. F.; O' Donovan, D. H.;

Verma, C. S.; Hyvonen, M.; Spring, D. R. Efficient Development of Stable and Highly

Functionalised Peptides Targeting the CK2 Alpha /CK2 Beta Protein-Protein Interaction. *Chem*

*Sci* **2019**, *10*, 5056–5063. https://doi.org/10.2210/PDB6Q38/PDB.

(49)  Filippakopoulos, P.; Rellos, P.; Fedorov, O.; Niesen, F.; Pike, A. C. W.; Pilka, E. S.; von Delft, F.;

Arrowsmith, C. H.; Edwards, A. M.; Weigelt, J.; Knapp, S. Crystal Structure of Human Death

Associated Protein Kinase 3 (DAPK3) in Complex with an Imidazo-Pyridazine Ligand. *TO BE*

*Publ.* https://doi.org/10.2210/PDB3BQR/PDB.

(50)  Pierce, A. C.; Jacobs, M.; Stuver-Moody, C. Docking Study Yields Four Novel Inhibitors of the

Protooncogene Pim-1 Kinase. *J.Med.Chem.* **2008**, *51*, 1972–1975.

https://doi.org/10.2210/PDB3BGZ/PDB.

(51)  Hou, Z.; Nakanishi, I.; Kinoshita, T.; Takei, Y.; Yasue, M.; Misu, R.; Suzuki, Y.; Nakamura, S.;

Kure, T.; Ohno, H.; Murata, K.; Kitaura, K.; Hirasawa, A.; Tsujimoto, G.; Oishi, S.; Fujii, N.

Structure-Based Design of Novel Potent Protein Kinase CK2 (CK2) Inhibitors with Phenyl-Azole

Scaffolds. *J. Med. Chem.* **2012**, *55* (6), 2899–2903. https://doi.org/10.1021/jm2015167.

(52)  Whelligan, D. K.; Solanki, S.; Taylor, D.; Thomson, D. W.; Cheung, K. M.; Boxall, K.; Mas-Droux,

C.; Barillari, C.; Burns, S.; Grummitt, C. G.; Collins, I.; Van Montfort, R. L.; Aherne, G. W.;

Bayliss, R.; Hoelder, S. Aminopyrazine Inhibitors Binding to an Unusual Inactive Conformation

of the Mitotic Kinase Nek2: Sar and Structural Characterization. *J.Med.Chem.* **2010**, *53*, 7682.

https://doi.org/10.2210/PDB2XK4/PDB.

(53)  Pike, A. C. W.; Savitsky, P.; Fedorov, O.; Krojer, T.; Ugochukwu, E.; von Delft, F.; Gileadi, O.;

Edwards, A.; Arrowsmith, C. H.; Weigelt, J.; Bountra, C.; Knapp, S. Structure of Human Serine-

Arginine-Rich Protein- Specific Kinase 2 (Srpk2) Bound to Purvalanol B. *TO BE Publ.*

https://doi.org/10.2210/PDB2X7G/PDB.

(54)    Dodson, C. A.; Kosmopoulou, M.; Richards, M. W.; Atrash, B.; Bavetsias, V.; Blagg, J.; Bayliss,
668            R. Crystal Structure of an Aurora-A Mutant That Mimics Aurora-B Bound to Mln8054: Insights
669            Into Selectivity and Drug Design. *Biochem.J.* **2010**, *427*, 19.
670            https://doi.org/10.2210/PDB2WTV/PDB.

671    (55)    Richards, M. W.; O'Regan, L.; Mas-Droux, C.; Blot, J. M.; Cheung, J.; Hoelder, S.; Fry, A. M.;
672            Bayliss, R. An Autoinhibitory Tyrosine Motif in the Cell-Cycle- Regulated Nek7 Kinase Is
673            Released Through Binding of Nek9. *Mol.Cell* **2009**, *36*, 560.
674            https://doi.org/10.2210/PDB2WQO/PDB.

675    (56)    Tong, Y.; Claiborne, A.; Stewart, K. D.; Park, C.; Kovar, P.; Chen, Z.; Credo, R. B.; Gu, W. Z.;
676            Gwaltney, S. L.; Judge, R. A.; Zhang, H.; Rosenberg, S. H.; Sham, H. L.; Sowin, T. J.; Lin, N. H.
677            Discovery of 1,4-Dihydroindeno[1,2-c]Pyrazoles as a Novel Class of Potent and Selective
678            Checkpoint Kinase 1 Inhibitors. *Bioorg.Med.Chem.* **2007**, *15*, 2759–2767.
679            https://doi.org/10.2210/PDB2E9O/PDB.

680    (57)    Pattanayek, R.; Egli, M. Protein-Protein Interactions in the Cyanobacterial Circadian Clock:
681            Structure of KaiA Dimer in Complex with C-Terminal KaiC Peptides at 2.8 Angstrom
682            Resolution. *Biochemistry* **2015**, *54*, 4575–4578. https://doi.org/10.2210/PDB5C5E/PDB.

683    (58)    Klaeger, S.; Heinzlmeir, S.; Wilhelm, M.; Polzer, H.; Vick, B.; Koenig, P. A.; Reinecke, M.;
684            Ruprecht, B.; Petzoldt, S.; Meng, C.; Zecha, J.; Reiter, K.; Qiao, H.; Helm, D.; Koch, H.; Schoof,
685            M.; Canevari, G.; Casale, E.; Depaolini, S. R.; Feuchtinger, A.; Wu, Z.; Schmidt, T.; Rueckert, L.;
686            Becker, W.; Huenges, J.; Garz, A. K.; Gohlke, B. O.; Zolg, D. P.; Kayser, G.; Vooder, T.;
687            Preissner, R.; Hahne, H.; Tonisson, N.; Kramer, K.; Gotze, K.; Bassermann, F.; Schlegl, J.;
688            Ehrlich, H. C.; Aiche, S.; Walch, A.; Greif, P. A.; Schneider, S.; Felder, E. R.; Ruland, J.; Medard,
689            G.; Jeremias, I.; Spiekermann, K.; Kuster, B. The Target Landscape of Clinical Kinase Drugs.
690            *Science (80-. ).* **2017**, *358*. https://doi.org/10.2210/PDB5MAF/PDB.

691    (59)    Niefind, K.; Bischoff, N.; Golub, A. G.; Bdzhola, V. G.; Balanda, A. O.; Prykhod'ko, A. O.;
692            Yarmoluk, S. M. Structural Hypervariability of the Two Human Protein Kinase CK2 Catalytic
693            Subunit Paralogs Revealed by Complex Structures with a Flavonol- and a Thieno[2,3-
694            d]Pyrimidine-Based Inhibitor. *Pharmaceuticals* **2017**, *10*.
695            https://doi.org/10.2210/PDB5M4F/PDB.

696    (60)    Brear, P.; De Fusco, C.; Hadje Georgiou, K.; Francis-Newton, N. J.; Stubbs, C. J.; Sore, H. F.;
697            Venkitaraman, A. R.; Abell, C.; Spring, D. R.; Hyvonen, M. Specific Inhibition of CK2 Alpha from
698            an Anchor Outside the Active Site. *Chem Sci* **2016**, *7*, 6839–6845.

699        https://doi.org/10.2210/PDB5CSP/PDB.

(61)    Ohno, H.; Minamiguchi, D.; Nakamura, S.; Shu, K.; Okazaki, S.; Honda, M.; Misu, R.; Moriwaki, H.; Nakanishi, S.; Oishi, S.; Kinoshita, T.; Nakanishi, I.; Fujii, N. Structure-Activity Relationship Study of 4-(Thiazol-5-Yl)Benzoic Acid Derivatives as Potent Protein Kinase CK2 Inhibitors. *Bioorg.Med.Chem.* **2016**, *24*, 1136–1141. https://doi.org/10.2210/PDB5B0X/PDB.

(62)    Boland, S.; Bourin, A.; Alen, J.; Geraets, J.; Schroeders, P.; Castermans, K.; Kindt, N.; Boumans, N.; Panitti, L.; Fransen, S.; Vanormelingen, J.; Stassen, J. M.; Leysen, D.; Defert, O. Design, Synthesis, and Biological Evaluation of Novel, Highly Active Soft ROCK Inhibitors. *J. Med. Chem.* **2015**, *58*, 4309–4324. https://doi.org/10.2210/PDB4WOT/PDB.

(63)    Guerra, B.; Bischoff, N.; Bdzhola, V. G.; Yarmoluk, S. M.; Issinger, O. G.; Golub, A. G.; Niefind, K. A Note of Caution on the Role of Halogen Bonds for Protein Kinase/Inhibitor Recognition Suggested by High- And Low-Salt CK2 Alpha Complex Structures. *Acs Chem.Biol.* **2015**, *10*, 1654–1660. https://doi.org/10.2210/PDB4UB7/PDB.

(64)    O'Reilly, M.; Cleasby, A.; Davies, T. G.; Hall, R. J.; Ludlow, R. F.; Murray, C. W.; Tisi, D.; Jhoti, H. Crystallographic Screening Using Ultra-Low-Molecular-Weight Ligands to Guide Drug Design. *Drug Discovery Today*. Elsevier Ltd May 2019, pp 1081–1086. https://doi.org/10.1016/j.drudis.2019.03.009.

(65)    Lee, T.; Hoofnagle, A. N.; Resing, K. A.; Ahn, N. G. Hydrogen Exchange Solvent Protection by an ATP Analogue Reveals Conformational Changes in ERK2 upon Activation. *J. Mol. Biol.* **2005**, *353* (3), 600–612. https://doi.org/10.1016/j.jmb.2005.08.029.

(66)    Andersen, M. D.; Shaffer, J.; Jennings, P. A.; Adams, J. A. Structural Characterization of Protein Kinase A as a Function of Nucleotide Binding. Hydrogen-Deuterium Exchange Studies Using Matrix-Assisted Laser Desorption Ionization-Time of Flight Mass Spectrometry Detection. *J. Biol. Chem.* **2001**, *276* (17), 14204–14211. https://doi.org/10.1074/jbc.M011543200.

(67)    Kleckner, I. R.; Foster, M. P. An Introduction to NMR-Based Approaches for Measuring Protein Dynamics. *Biochimica et Biophysica Acta - Proteins and Proteomics*. NIH Public Access August 2011, pp 942–968. https://doi.org/10.1016/j.bbapap.2010.10.012.

(68)    Viegas, A.; Manso, J.; Nobrega, F. L.; Cabrita, E. J. Saturation-Transfer Difference (STD) NMR: A Simple and Fast Method for Ligand Screening and Characterization of Protein Binding. *J. Chem. Educ.* **2011**, *88* (7), 990–994. https://doi.org/10.1021/ed101169t.

(69)    Lindenblatt, D.; Nickelsen, A.; Applegate, V.; Jose, J.; Niefind, K. Structural and Mechanistic

Basis of the Inhibitory Potency of Selected 2-Aminothiazole Compounds on Protein Kinase

CK2. *J. Med. Chem.* **2020**, acs.jmedchem.0c00587.

https://doi.org/10.1021/acs.jmedchem.0c00587.

(70)   Lee, W.-K.; Son, S. H.; Jin, B.-S.; Na, J.-H.; Kim, S.-Y.; Kim, K.-H.; Kim, E. E.; Yu, Y. G.; Lee, H. H.

Structural and Functional Insights into the Regulation Mechanism of CK2 by IP6 and the

Intrinsically Disordered Protein Nopp140. *Proc. Natl. Acad. Sci.* **2013**, *110* (48), 19360–19365.

https://doi.org/10.1073/pnas.1304670110.

(71)   Vom, A.; Headey, S.; Wang, G.; Capuano, B.; Yuriev, E.; Scanlon, M. J.; Simpson, J. S. Detection

and Prevention of Aggregation-Based False Positives in STD-NMR-Based Fragment Screening.

*Aust. J. Chem.* **2013**, *66* (12), 1518–1524. https://doi.org/10.1071/CH13286.

(72)   Feng, B. Y.; Shoichet, B. K. A Detergent-Based Assay for the Detection of Promiscuous

Inhibitors. *Nat. Protoc.* **2006**, *1* (2), 550–553. https://doi.org/10.1038/nprot.2006.77.

(73)   Shoichet, B. K. Screening in a Spirit Haunted World. *Drug Discov. Today* **2006**, *11* (13–14),

607–615. https://doi.org/10.1016/j.drudis.2006.05.014.

(74)   Coan, K. E. D.; Maltby, D. A.; Burlingame, A. L.; Shoichet, B. K. Promiscuous Aggregate-Based

Inhibitors Promote Enzyme Unfolding. *J. Med. Chem.* **2009**, *52* (7), 2067–2075.

https://doi.org/10.1021/jm801605r.

(75)   McGovern, S. L.; Helfand, B. T.; Feng, B.; Shoichet, B. K. A Specific Mechanism of Nonspecific

Inhibition. *J. Med. Chem.* **2003**, *46* (20), 4265–4272. https://doi.org/10.1021/jm030266r.

(76)   Walters, J.; Milam, S. L.; Clark, A. C. Chapter 1 Practical Approaches to Protein Folding and

Assembly. Spectroscopic Strategies in Thermodynamics and Kinetics. *Methods in Enzymology*.

NIH Public Access 2009, pp 1–39. https://doi.org/10.1016/S0076-6879(08)04201-8.

 Table of Content graphics