1

# CUT&RUN reveals unique positioning of pre-initiated

# RNA polymerase II in the steady state of transcription

Michi Miura* and Honglin Chen

State Key Laboratory for Emerging Infectious Diseases and

Department of Microbiology, The University of Hong Kong,

Hong Kong SAR, China

*Email: michi.miura.res@gmail.com

## Abstract

CUT&RUN is a powerful tool to study protein-DNA interactions *in vivo*. DNA fragments cleaved by the targeted micrococcal nuclease identify the footprints of DNA-binding proteins on the chromatin. We performed CUT&RUN on human lung carcinoma cell line A549 maintained in a multi-well cell culture plate to profile RNA polymerase II. Size-fractionation of aligned reads revealed that the collection of long (>270 bp) DNA fragments corresponded to the bimodal peak around the transcription start sites, as previously seen with chromatin immunoprecipitation. However, we found that shorter (<120 bp) fragments identify a well-defined peak localised at the transcription start sites. This distinct DNA footprint of shorter fragments, which constituted only about 5% of the total reads, indicates the unique positioning of the poised RNA polymerase II before transcription initiation, which has not been detected in the steady state of transcription by standard chromatin immunoprecipitation.

**Keywords:** CUT&RUN; protein-DNA interaction; RNA polymerase II; transcription initiation; promoter-proximal pausing; chromatin profiling

## Introduction

CUT&RUN (Cleavage Under Targets and Release Using Nuclease) (Skene & Henikoff, 2017) is a powerful tool to map protein-DNA interactions *in vivo*. CUT&RUN directs Protein A/G-fused micrococcal nuclease (pAG-MNase) (Meers, Bryson, Henikoff, & Henikoff, 2019) to the antibody-bound protein-DNA complex, with a subsequent release of the DNA fragments cleaved in the presence of calcium ions. There are two major advantages of CUT&RUN over conventional chromatin immunoprecipitation (ChIP). First, CUT&RUN does not require sonication for chromatin fragmentation. Physical shearing of chromatin by sonication does not occur randomly, which can skew signal detection; and excessive sonication may attenuate the epitope of the target protein (Marx, 2019). Second, CUT&RUN identifies the footprints of DNA-binding proteins on the chromatin, which revealed distinct binding configurations of transcription factors in yeast (Skene & Henikoff, 2017) and mammalian cells (Meers, Janssens, & Henikoff, 2019).

In this study, we performed CUT&RUN on the human lung carcinoma cell line A549 maintained in a multi-well cell culture plate (Figure 1A) to profile RNA polymerase II (Pol II) around the transcription start sites (TSS). We found that CUT&RUN could validate the bimodal signal around the TSS as previously detected using ChIP (Erickson, Sheridan, Cortazar, & Bentley, 2018). However, we also identified a distinct peak at the TSS from short DNA fragments released only by pAG-MNase. This finding indicates a transient positioning of the pre-initiated RNA polymerase II in the steady state of transcription, which has eluded detection by conventional ChIP (Erickson et al., 2018).

## Results

### Overall performance of "CUT&RUN on plate" for profiling Pol II

First, we tested the feasibility of performing CUT&RUN on adherent cells attached to a multi-well polystyrene cell culture plate with Pol II antibodies (Figure 1A). Pol II signals obtained by this method are shown in Figure 1B. The genomic region chr6 p22.2 was selected due to the presence of multiple histone genes. This location is ideal for efficient visual inspection of Pol II signals because of the compact size and close spacing of these genes. Our Pol II signals consistently mapped to the transcription units of histone genes with low background as previously reported (Meers, Bryson, et al., 2019; Skene & Henikoff, 2017) (Figure 1B). The signals along the gene body, in addition to 1.5 kb upstream of the TSS and downstream of the transcription end site (TES), were averaged over "singletons" (genes that are separated from other genes by >2 kb) (Figure 1C), showing consistent enrichment around the TSS. Therefore, through the successful capture of Pol II signals *in situ* with low background, the performance of our "C̲UT&R̲UN o̲n p̲late" (or CROP) protocol was shown to be comparable with the conventional CUT&RUN (Meers, Bryson, et al., 2019; Skene & Henikoff, 2017).

**Figure 1   Overall performance of "CUT&RUN on plate"**

(A) Protocol overview.  In conventional CUT&RUN (top row), cells are harvested and attached on Concanavalin A-magnetic beads.  Subsequent cell permeabilisation, antibody binding, and targeted pAG-MNase digestion are performed on these cells immobilised on the beads.  The magnetic beads are captured each time the buffers are changed.  "CUT&RUN on plate" (or CROP) (bottom row) circumvents the use of Concanavalin A-magnetic beads.  Adherent cells are anchored on the solid surface of the multi-well plate throughout the protocol, enabling rapid buffer change for multiple samples.  (B) Pol II signals in the histone gene cluster obtained by CROP.  The Pol II signal is expressed by the coverage normalised with the total read count.  (C) Metaplot analysis of the Pol II signal.  Pol II signal was averaged over genes that are at least 2 kb away from other genes.  The region between 1.5 kb upstream of the transcription start site (TSS) and 1.5 kb downstream of the transcription end site (TES) is presented.  The length of the gene body is scaled to 3 kb.

**Size-fractionation of aligned reads to reveal distinct footprints of Pol II-binding**

The size distribution of aligned paired-end reads showed multiple peaks (Figure 2A). We categorised the aligned reads into 5 subsets by length (Figure 2A), each of which is expected to reflect a unique footprint of the protein-DNA interaction (Meers, Janssens, et al., 2019; Skene & Henikoff, 2017). The size of DNA fractions was shown to reveal distinct signals of Pol II around the TSS (Figure 2B). The signal of the total reads (i.e. without size-fractionation) conformed to the characteristic bimodal peak as previously reported by ChIP (i.e. the peaks located at either ~150 bp upstream or downstream of the TSS) (Erickson et al., 2018). By contrast, the shorter fragments (<120 bp) constituted a single narrow peak at the TSS, and the longer fragments (>120 bp) accumulated adjacent to the TSS (Figure 2B). As expected, the bimodal signal of the longer fragments (>270 bp) is skewed towards the region downstream of the TSS (Figure 2B) (Erickson et al., 2018). To validate our findings, we analysed the TSS of individual genes. We confirmed that some genes showed an accumulation of longer (>270 bp) fragments on both sides of the TSS (Figure 2C), whereas some other genes had signals on only one side (Figure 2D and E). The signals of the shorter (<120 bp) and longer (>270 bp) fragments were mutually exclusive, showing two distinct footprints of Pol II adjacent to each other. The shorter fragments constituted only about 5% of the total reads (Figure 2A).

**Figure 2   Size-fractionation identifies distinct Pol II footprints**

(A) Size distribution of aligned reads (the insert size between the sequence adaptors).  The reads were categorised into five subsets by size, each of which is coded by a unique colour.  (B) Pol II signal (Millipore, 05-623; unphosphorylated and S5p) from each size fraction in ±1.5 kb around the transcription start site (TSS).  The signals in 13,452 annotated RefSeq singleton genes are presented in the heatmap; a *k*-mean clustering was performed on the total reads (coded in black) to enrich genes with Pol II signal (Cluster 1 and Cluster 2). The metaplot for each cluster is shown above the heatmap (Cluster 1, solid line; Cluster 2, dashed line).  (C-E) Pol II signal (Millipore, 05-623; unphosphorylated and S5p) in ±1 kb around the transcription start site of individual genes.  The vertical axis indicates the pile-up of aligned reads from the three size fractions.  The same colour codes are used as per panel (A).  Note that the coverage was normalised per million reads within each of the size fractions; the relative coverage within each plot is not proportional to the actual occurrence of these fractions.  (F) A schematic drawing of the Pol II footprints deduced by the alignment of size-fractionated reads.

Recent studies using ChIP-seq, as well as nascent transcript sequencing, have shown

that Pol II is paused at ~50 bp downstream of the TSS (Core, Waterfall, & Lis, 2008;

Erickson et al., 2018; Kwak, Fuda, Core, & Lis, 2013; Nojima et al., 2015; Shao &

Zeitlinger, 2017).  Our observations suggest two distinct binding configurations of Pol II

near the TSS (Figure 2F).  These configurations are: (1) poised Pol II at the TSS in the

pre-initiation state where nucleosomes are depleted, and (2) paused Pol II upstream or

downstream of the TSS where Pol II pausing is regulated by a higher-molecular complex

after transcription initiation.  The paused Pol II peaks were visible exclusively in the

longer (>120 bp) fragments adjacent to the TSS.  We reason that this is because Pol II

is associated with a higher-molecular complex at the paused site (Vos, Farnung, Urlaub,

& Cramer, 2018), which protects the chromatin from the pAG-MNase cleavage.  As a

result, pAG-MNase is unable to produce shorter (<120 bp) fragments associated with

the paused Pol II.  Composite signals were observed in our assay (Figure 2C to E), each

of which corresponds to one of the two distinct binding configurations as described.

These composite signals result from CUT&RUN taking a 'snapshot' of each gene from

hundreds of thousands of cells per assay at the TSS; a transcription start site

accommodates one molecule of Pol II at a time, either at the TSS (pre-initiation) or the

site of promoter-proximal pausing (post-initiation) (Gressel et al., 2017; Shao &

Zeitlinger, 2017).

**Impact of formaldehyde fixation on "CUT&RUN on plate"**

Formaldehyde fixation is often used to investigate transient protein interactions.  We

tested if formaldehyde fixation can be implemented in "CUT&RUN on plate".  Cells were

fixed with 1.5% formaldehyde in either pre- or post-cellular permeabilisation (see

Materials and Methods).  Although formaldehyde fixation reduced the detection of Pol II

signals, serine 5-phosphorylated (S5p) Pol II was shown to be enriched near the TSS

(Figure 3A).

**Figure 3   Impact of formaldehyde cell fixation on the S2p Pol II signals**

(A) S5p and S2p Pol II signal over a gene on chr10 (SIRT1) (S5p, ab5131; S2p, ab5095).  Two colour codes
indicate whether cells were fixed with formaldehyde (cyan, "No fix"; magenta, "Fixed").  Cells were fixed
following a brief cytoplasmic extraction ("Fixed" 1) or intact cells were fixed directly ("Fixed" 2).  The signals
were normalised with the yeast spike-in DNA.  (B) Metaplot analyses of S2p Pol II signal from 1.5 kb
upstream of the transcription start site (TSS) to 4.5 kb downstream of the transcription end site (TES).  The
signals were averaged over the singleton genes (n = 13,452).  The gene body is scaled to 4.5 kb in length.
(C) Two Pol II metrics measuring the ratio of signals in the gene body to TSS (left) and in the TES to TSS
(right).  These metrics were calculated for genes (n = 1,435) that were longer than 5 kb and aggregated in
Cluster 1 (Figure 2B).  The metrics of each gene are plotted for each combination of the assays and the
density of plots are represented by colour brightness (navy to yellow).  The data is presented in the
logarithmic scale (base = 10) for ease of visualisation.

We observed a profound effect of formaldehyde fixation on the serine 2-phosphorylated

(S2p) Pol II signal (Figure 3A).  The metaplot analysis showed that the average signal of

S2p RNA polymerase II over the singleton genes was more pronounced in the region

downstream of the TES when cells were fixed (Figure 3B).  This observation was

confirmed by signal quantification in the gene body or TES (relative to that in the TSS

respectively), for each singleton gene (Figure 3C, schematic drawing).  While these

metrics for each gene were in good concordance within the same fixation condition, we

found that the ratio of S2p Pol II signal in the gene body to TSS, or in the TES to TSS

was higher when the cells were fixed (Figure 3C, "No Fix" vs "Fixed").  We consider two

possible explanations of this observation: (1) fixation stabilises the elongating S2p RNA

polymerase II on the DNA, or more likely, (2) fixation preferentially reduces the detection

of Pol II near the TSS, thereby altering the ratio of signals in the TSS, gene body and

TES.

## Discussion

**Discussion**

In this study, we have devised "CUT&RUN on plate" (CROP) for profiling protein-DNA

interactions in adherent cells maintained in a multi-well cell culture plate. The current

CUT&RUN protocol (Hainer & Fazzio, 2019; Meers, Bryson, et al., 2019; Skene, Henikoff,

& Henikoff, 2018) relies on Concanavalin A-coated magnetic beads to immobilise the

cells before the targeted pAG-MNase cleavage. This immobilisation step introduces a

range of shortcomings associated with the use of Concanavalin A-beads. For example,

trypsinisation while harvesting the cells may alter the behaviour of DNA-binding

proteins and lower the binding affinity of cells to Concanavalin A. Concanavalin A also

stimulates certain types of cells. When the plasma membrane is removed to investigate

the nuclei, Concanavalin A does not bind efficiently to the nuclear membrane. Finally,

Concanavalin A-beads tend to aggregate when cells are attached. Our CROP protocol

described here overcomes these key problems associated with the use of Concanavalin-

A beads and may find widespread application.

Using CUT&RUN, we profiled RNA polymerase II near the transcription start sites. The

turnover of Pol II near the TSS is highly dynamic (Darzacq et al., 2007; Erickson et al.,

2018; Steurer et al., 2018). Since the duration of Pol II in the promoter-proximal

pausing state is about ten times longer than that of Pol II in the pre-initiation state

(Darzacq et al., 2007; Steurer et al., 2018), a majority of reads obtained by either ChIP

or CUT&RUN is from the paused site, approximately 50 bp downstream of the TSS.

Because ChIP fragments are typically hundreds of base pairs long, the rare signal of the

poised Pol II at the TSS is often obscured. While Erickson *et al.* had previously identified

the signal of the poised Pol II at the TSS by ChIP, this signal was only exposed following

the depletion of Pol II in hypotonic condition with the use of triptolide to block the

initiation (Erickson et al., 2018). Nascent transcript sequencing is, by its nature, unable

to capture the poised RNA polymerase II before transcription initiation. In contrast, we

accurately identified the positioning of poised Pol II in the physiological condition

through the recovery of its unique footprint (Figure 2F). Therefore, CUT&RUN is an

effective substitute for the conventional chromatin immunoprecipitation to investigate

the behaviour of RNA polymerase II at promoters.

## Materials and Methods

### Cells

Human lung carcinoma cell line A549 was maintained in Dulbecco's Modified Eagle Medium (Gibco, 11965-092) supplemented with 10% FBS (Gibco, 16140-071), 100 U/ml penicillin and 100 μg/ml streptomycin (Gibco, 15140-122).

### CUT&RUN in a cell culture plate

CUT&RUN was performed on a standard 24-well cell culture plate (TPP, 92012); cells were left attached to the bottom of the well throughout the protocol.

A549 was seeded in wells and grown to ~70 to 90% confluency (~0.5 million cells) at the time of an assay. Cells were washed with PBS and permeabilised with Perm buffer (0.1% Triton X-100 (Sigma, X100), 20 mM HEPES-KOH pH 7.5, 150 mM NaCl, 0.5 mM Spermidine (Sigma, S2626) and proteinase inhibitor (Roche, 04 693 132 001)) for 15 min at room temperature. After removing Perm buffer, cells were washed again with Perm buffer and incubated with an antibody (~0.7 μg) in 150 μl Perm buffer per well for one hour at room temperature. The antibodies used in this study were: mouse anti-RNA polymerase II (Millipore, 05-623; clone CTD4H8; 0.75 μg per well); rabbit anti-RNA polymerase II (phospho S2) (Abcam, ab5095; polyclonal; 0.75 μg per well); and rabbit anti-RNA polymerase II (phospho S5) (Abcam, ab5131; polyclonal; 0.675 μg per well). After incubation, unbound antibody was removed, and cells were washed twice with Perm buffer. Cells were then incubated with the recombinant pAG-MNase (Cell Signaling, 40366; 1:33 volume) in 150 μl Perm buffer per well for an hour at room temperature. Unbound pAG-MNase was removed, and cells were washed twice with

Perm buffer. During the second wash of this stage, the plate was placed on ice-cold

water. For pAG-MNase activation, 150 μl ice-cold Perm buffer containing 5 mM $CaCl_2$

was dispensed per well, and the plate was incubated on ice-cold water for 30 min. The

pAG-MNase digestion was halted by the addition of 50 μl 4× STOP solution (680 mM

NaCl, 40 mM EDTA (Sigma, E5134), 8mM EGTA (AG Scientific, E-2491), 100 μg/ml

RNase A (Invitrogen, 12091021) and 0.1% Triton X-100). The plate was incubated at

37°C for 30 min to facilitate the release of digested DNA fragments from the nucleus.

The supernatant was collected, and DNA was extracted using DNeasy Blood & Tissue

Kit (Qiagen, 69504).

In addition to the native "CUT&RUN on plate" described above, two protocols for cell

fixation with 1.5% formaldehyde were tested: (1) cells were fixed with 1.5% formaldehyde

for 10 min prior to the cell permeabilisation with Perm buffer; and (2) cells were fixed

following the permeabilisation (5 min in Perm buffer). The supernatant collected at the

end of the CUT&RUN protocol was supplemented with 1% SDS (final concentration) and

incubated at 65°C overnight for de-crosslinking. The DNA was purified with QIAquick

PCR Purification Kit (Qiagen, 28104).

**Pair-end sequencing and read alignment**

DNA libraries were constructed using NEBNext Ultra II DNA Library Prep Kit for Illumina

(NEB, E7645) with index primers (NEB, E7335 and E7710). Fragmented genomic DNA

from *S. cerevisiae* (Cell Signaling, 40366) was used (10 pg per sample) as spike-in.

DNA libraries were sequenced (150 bp × 2) with NovaSeq 6000 (Illumina), and the initial

50 bases of each read were aligned to a concatenation of human reference genome

(GRCh38) and *S. cerevisiae* (sacCer3) with a BWA algorithm (version 0.7.17) `bwa mem`

(Li & Durbin, 2009). Duplications of reads were marked with a Picard tool (version

2.18.9) `MarkDuplicates` (http://broadinstitute.github.io/picard). Only a minority of

the aligned reads (less than 0.151%) constituted the yeast genome, thereby confirming

the successful recovery of DNA fragments cleaved by pAG-MNase. A subset of aligned

reads was selected for the downstream analyses using Samtools (version 1.9) with the

following commands: SAM flag 3 (read paired; read mapped in proper pair) (`samtools`

`view -f 3`); mapping quality greater than 20 (`samtools view -q 20`); without tags

'XA:Z:' and 'SA:Z:' (`grep -v -e 'XA:Z:' -e 'SA:Z:'`); and aligned in either one of

the chromosomes chr1 to chr22, chrX or chrY (`samtools view chr1 chr2 ... chrY`).

**Data analyses and visualisation**

Aligned reads were fractionated by size (the 9th field of SAM) and stored in the BAM

format. Read coverage was computed and stored in the bigWig format with deepTools's

(version 3.1.3) `bamCoverage` (Ramirez et al., 2016), normalised per million reads. For

comparison of the Pol II signals across the fixation conditions (Figure 3A and B), read

coverage was computed with deepTools's (version 3.3.2) `bamCoverage` using scale

factors calculated with the number of spike-in reads mapped to the yeast genome. The

bigWig files were converted into the bedGraph format using `bigWigToBedGraph` (UCSC)

and the read coverage was visualised with the R package Sushi (version 1.24.0)

(Phanstiel, Boyle, Araya, & Snyder, 2014) in R (version 3.6.2). A matrix containing the

read coverage over a given region of genes was computed using deepTools's (version

3.3.2) `computeMatrix` with an annotation

GCA_000001405.15_GRCh38_full_analysis_set.refseq_annotation (NCBI). A subset of

genes that were separated from other genes by >2 kb was considered for this analysis;

such genes were extracted using bedtools's (version 2.26.0) (Quinlan & Hall, 2010)

`merge` and `intersect`. The matrix was visualised as a heatmap and a metaplot using

the R package EnrichedHeatmap (version 1.16.0) (Gu, Eils, Schlesner, & Ishaque, 2018).

**Signal quantification in the TSS, gene body and TES**

The regions for TSS, gene body and TES were defined: TSS, ±0.5 kb around the start

site of a gene; gene body, from 0.5 kb downstream of the start site to 1.5 kb upstream

of the end site; and TES, 1.5 kb upstream of the end site to 4.5 kb downstream of the

end site. For each of the TSS, gene body and TES regions, genome coordinates for the

singleton genes in the RefSeq annotation were stored in the BED format.

The BAM files containing the aligned paired reads were converted into the BED format

with bedtools's `bamtobed -bedpe` and the subsequent extraction of the 1st

(chromosome), 2nd, 6th (the start and end positions of an alignment, respectively), 7th,

8th and 9th fields.

For each singleton gene, paired reads aligned to the TSS, gene body or TES were counted

using bedtools's `intersect -F 0.5` (reads that overlapped by less than half of the

alignment length were rejected). The read counts in the gene body and TES were

normalised with the number of reads contained in the TSS.

## Acknowledgements

We thank Liew Jun Mun at University of Cambridge, for reviewing the manuscript to enhance the readability. We also thank Prof. Charles R. M. Bangham at Imperial College London, for reviewing the manuscript and critical comments. Illumina NovaSeq 6000 sequencing was performed at Centre for PanorOmic Sciences (CPOS) Genomics Core, LKS Faculty of Medicine, The University of Hong Kong. This study was supported in part by the Research Grants Council of the Hong Kong SAR (17119618, 17107019 and C7027-16G).

## Competing interests

The authors declare no competing interests.

# References

**References**

Core, L. J., Waterfall, J. J., & Lis, J. T. (2008). Nascent RNA sequencing reveals

widespread pausing and divergent initiation at human promoters. *Science,*

*322*(5909), 1845-1848. doi:10.1126/science.1162228

Darzacq, X., Shav-Tal, Y., de Turris, V., Brody, Y., Shenoy, S. M., Phair, R. D., & Singer,

R. H. (2007). In vivo dynamics of RNA polymerase II transcription. *Nat Struct Mol*

*Biol, 14*(9), 796-806. doi:10.1038/nsmb1280

Erickson, B., Sheridan, R. M., Cortazar, M., & Bentley, D. L. (2018). Dynamic turnover

of paused Pol II complexes at human promoters. *Genes Dev, 32*(17-18), 1215-

1225. doi:10.1101/gad.316810.118

Gressel, S., Schwalb, B., Decker, T. M., Qin, W., Leonhardt, H., Eick, D., & Cramer, P.

(2017). CDK9-dependent RNA polymerase II pausing controls transcription

initiation. *Elife, 6.* doi:10.7554/eLife.29736

Gu, Z., Eils, R., Schlesner, M., & Ishaque, N. (2018). EnrichedHeatmap: an

R/Bioconductor package for comprehensive visualization of genomic signal

associations. *BMC Genomics, 19*(1), 234. doi:10.1186/s12864-018-4625-x

Hainer, S. J., & Fazzio, T. G. (2019). High-Resolution Chromatin Profiling Using

CUT&RUN. *Curr Protoc Mol Biol, 126*(1), e85. doi:10.1002/cpmb.85

Kwak, H., Fuda, N. J., Core, L. J., & Lis, J. T. (2013). Precise maps of RNA polymerase

reveal how promoters direct initiation and pausing. *Science, 339*(6122), 950-953.

doi:10.1126/science.1229386

Li, H., & Durbin, R. (2009). Fast and accurate short read alignment with Burrows-

Wheeler transform. *Bioinformatics, 25*(14), 1754-1760.

doi:10.1093/bioinformatics/btp324

Marx, V. (2019). What to do about those immunoprecipitation blues. *Nat Methods, 16*(4),

289-292. doi:10.1038/s41592-019-0365-3

Meers, M. P., Bryson, T. D., Henikoff, J. G., & Henikoff, S. (2019). Improved CUT&RUN

chromatin profiling tools. *Elife, 8*. doi:10.7554/eLife.46314

Meers, M. P., Janssens, D. H., & Henikoff, S. (2019). Pioneer Factor-Nucleosome Binding

Events during Differentiation Are Motif Encoded. *Mol Cell, 75*(3), 562-575 e565.

doi:10.1016/j.molcel.2019.05.025

Nojima, T., Gomes, T., Grosso, A. R. F., Kimura, H., Dye, M. J., Dhir, S., . . . Proudfoot,

N. J. (2015). Mammalian NET-Seq Reveals Genome-wide Nascent Transcription

Coupled    to    RNA    Processing.    *Cell,    161*(3),    526-540.

doi:10.1016/j.cell.2015.03.027

Phanstiel, D. H., Boyle, A. P., Araya, C. L., & Snyder, M. P. (2014). Sushi.R: flexible,

quantitative and integrative genomic visualizations for publication-quality multi-

panel    figures.    *Bioinformatics,    30*(19),    2808-2810.

doi:10.1093/bioinformatics/btu379

Quinlan, A. R., & Hall, I. M. (2010). BEDTools: a flexible suite of utilities for comparing

genomic    features.    *Bioinformatics,    26*(6),    841-842.

doi:10.1093/bioinformatics/btq033

Ramirez, F., Ryan, D. P., Gruning, B., Bhardwaj, V., Kilpert, F., Richter, A. S., . . .

Manke, T. (2016). deepTools2: a next generation web server for deep-sequencing

data analysis. *Nucleic Acids Res, 44*(W1), W160-165. doi:10.1093/nar/gkw257

Shao, W., & Zeitlinger, J. (2017). Paused RNA polymerase II inhibits new transcriptional

initiation. *Nat Genet, 49*(7), 1045-1051. doi:10.1038/ng.3867

Skene, P. J., Henikoff, J. G., & Henikoff, S. (2018). Targeted in situ genome-wide
profiling with high efficiency for low cell numbers. *Nat Protoc, 13*(5), 1006-1019.
doi:10.1038/nprot.2018.015

Skene, P. J., & Henikoff, S. (2017). An efficient targeted nuclease strategy for high-
resolution mapping of DNA binding sites. *Elife, 6*. doi:10.7554/eLife.21856

Steurer, B., Janssens, R. C., Geverts, B., Geijer, M. E., Wienholz, F., Theil, A. F., . . .
Marteijn, J. A. (2018). Live-cell analysis of endogenous GFP-RPB1 uncovers rapid
turnover of initiating and promoter-paused RNA Polymerase II. *Proc Natl Acad Sci*
*U S A, 115*(19), E4368-E4376. doi:10.1073/pnas.1717920115

Vos, S. M., Farnung, L., Urlaub, H., & Cramer, P. (2018). Structure of paused
transcription complex Pol II-DSIF-NELF. *Nature, 560*(7720), 601-606.
doi:10.1038/s41586-018-0442-2