Multiple dimensions of social motivation in adult female degus

Navdeep K. Lidhar[1,2], Ayushi Thakur[1], Anna-Julia David[1], Kaori Takehara-Nishiuchi[1], Nathan Insel[3*]

[1] Department of Psychology, University of Toronto, Toronto, ON, Canada

[2] Department of Psychology, University of Toronto Mississauga, Mississauga, ON, Canada

[3] Department of Psychology, University of Montana, Missoula, MT, USA

*correspondence should be addressed to nathan.insel@umontana.edu, University of Montana, Skaggs Rm 143, 32 Campus Dr., Missoula, MT, USA

Abstract

In many species, social isolation increases subsequent interaction. This phenomenon may involve general drives, e.g. for social touch or companionship, as well as drives that are specific to particular peers, and which ultimately serve to re-establish expectations and relationships between the individuals. Female degus are known to be affiliative with multiple other individuals, including unrelated and unfamiliar conspecifics, offering an opportunity to study social motivation independent from exclusive pair-bonds or overt, same-sex competition. We attempted to disentangle factors driving peer interaction by examining reunion behavior across several social isolation and separation manipulations. High levels of interaction were observed between adult females who had been separated even without isolation, revealing a drive to re-establish relationships with specific peers. The content of separation-only reunions differed from isolation, with the latter involving more early-session interaction, higher levels of allogrooming before rear-sniffing, and a higher ratio of chitter vocalizations. To assess whether post-isolation behavior was related to stress, we examined reunions following a non-social (footshock) stressor. Like isolation, footshock increased early-session interactions, but did not increase allogrooming before rear-sniffing or chittering, as compared with controls. To test whether separation-only reunion behavior shared qualities with relationship formation, we also examined reunions of new (stranger) dyads. Strangers exhibited higher levels of interaction than cagemates, with particularly high levels of late-session rear-sniffing. Like separation-only reunions, strangers showed more non-chitter vocalizations and lower levels of allogrooming before rear-sniffing. Across experiments, an exploratory clustering method was used to identify vocalizations that differed between conditions. This yielded promising leads for future investigation, including a chaff-type syllable that may have been more common during relationship renewal. Overall, results are consistent with the hypothesis that female degu reunions are supported by both general and peer-stimulus specific drives, expressed through the structure of physical and vocal interactions over time.

keywords: motivation, ethology, relationship, reunion, rodents

Introduction

In many species, a period of social isolation can cause a subsequent increase in social activity (e.g., humans [1] and rats [2]). The drive to interact after social deprivation is reminiscent of traditional models of motivation, in which the energy for a behavior increases as more time passes since the behavior's expression [3,4]. But social interaction can take different forms and satisfy different needs, many of which are conflated during isolation. One distinction that can be made is between "stimulus-specific" drives between individuals that have not seen one-another for a period of time and more general drives for, for example, social touch, companionship, or a sense of social agency. These two domains—stimulus-specific and stimulus-general—are difficult to isolate experimentally. In pair-bonding species, for example, general social drives are largely exclusive to a single individual, so the expression of motivation for one, specific peer is conflated with social motivation more generally. Some insight on stimulus-specific social drives comes from research on greeting behavior between individuals that have been separate for extended periods of time. Such greetings are thought to help reestablish dyadic relationships and trust [5–8]; put another way, greetings reestablish expectations individuals have about one-other, including others' probable actions and reactions. Complete social isolation, meanwhile, deprives an animal of all forms of social activity. The motivational impact of isolation may be at least partly mediated by neuroendocrine stress responses, as isolation causes an increase in systemic glucocorticoids [9] and social interaction, in turn, is known reduce or "buffer" stress levels [10–15]. A number of physiological changes are known to take place during social isolation [16–23], some of which can be directly linked with stress, and others that may be more independent. An important question is what contribution stress, by itself, contributes to interactive behavior after isolation.

While multiple different drives likely contribute to peer interaction after isolation, it remains unclear how each differentially impacts the content or structure of social interactions. The primary goal of the present study is to help build a "lexicon" of social behaviors associated with specific motivational contexts that can later be used to infer motivational states in less controlled settings. We focus our examination on the degu (*Octodon degus*), a caviomorph rodent from Chile. Degus are an interesting species for studies of social interaction on account of their high sociality [24], dynamic group structures [25–28], as well as the parallel research that can be done in both field

and laboratory settings. Most importantly for the present experiments, adult female degus are highly interactive but do not show systematic preferences for familiar over unfamiliar peers [29]. This allows investigation of same-sex peer motivations without the confound of selective long-term bonds, meanwhile also minimizing aggressive behaviors that are typical between unfamiliar, same-sex adults of many rodent species. In native habitats, female degus are recognized for their constructive interactions, such as communal nursing, with both related and unrelated, same-sex individuals [26,30–33]. Degus also express a wide range of vocalizations in the human hearing range, and these have been manually classified based on spectral features and the social contexts in which they are used [34]. Degus therefore express a high dimensionality of social behavior that can be quantified to compare interactions between different experimental conditions.

We test four main hypotheses. Each experimental prediction assumes that more motivation results in more interactions during a 20 minute "reunion" session, and that different motivational drives are expressed with different interaction patterns in ways that generalize across dyads. Hypothesis 1: Given that adult degus show high sociality in their natural habitat and in prior laboratory research, motivation to interact increases the longer they go without socializing. This predicts that animals will spend more time interacting following longer (24 hr) periods of isolation compared with very little (1 min) isolation time, with intermediate levels of interaction following short-term (45 min) isolation. Hypothesis 2: Degus form peer-specific relationships and/or expectations that require renewal after a period of separation. This predicts that A) a degu will become more motivated to interact with a peer following an extended (24 hr) period of separation, as compared with a "baseline" 1 min of isolation, and that B) separation-without-isolation reunions will differ from those following isolation. Hypothesis 3: Stress at least partly mediates increased interaction levels following isolation. This predicts that A) a non-social (footshock) stressor will also increase female social interaction and, B) reunions following footshock will share behavioral patterns with those observed following isolation. Finally, Hypothesis 4: adult female degus are highly motivated to interact with strangers due to drives that help establish new expectations and relationships. This predicts that A) stranger reunions will involve levels of interaction at least as high as those between reunited cagemates and B) there will be similarities between interaction patterns observed between strangers and those between separated-but-not-isolated cagemate dyads.

We record degu behavior during 20 minute reunions following manipulations that include social
isolation, separating dyads without isolating them from other conspecifics, and exposure to an
external stressor. The results reveal several subtle but notable distinctions in how different forms
of motivation impact behavior.

Methods

Subjects

Thirty-eight female degus aged 6 to 19 months (mean = 10.2 ± 3.6 SD) were used for all
experiments. All degus came from our breeding colony at University of Toronto, descended from
animals from University of Michigan and an Ontario breeder. By the start of the study the colony
had been established in Toronto for approximately four years, and all degus used had been born
in-house. Degus reach developmental maturity by 3.5 months [35,36] and in the lab live up to 7 to
8 years [37]. We therefore consider the 6 to 19 month range as "young adult". While the study was
not designed to evaluate differences within this age range, a comparison of age against total
interactive time yielded a negative result: n = 17, r = -0.22, p = 0.39) so age was not otherwise
considered within the scope of the experiment. Prior to the start of experiments, degus were housed
either in pairs within 14" x 11" x 9.5" or, in the case of Experiment 2, in groups of four within 15"
x 19" x 8" polycarbonate cages. All were housed in a vivarium at the biological science facility at
the University of Toronto. Paired and group housing was usually with siblings, but in Experiment
1, three of the nine pairs were non-sibling (placed together sometime between weening and one
month prior to the experiment). Prior research has found that, in degus, familiarity more strongly
determines social interactions than kinship [38]; but effect sizes are described and compared for
sibling and non-sibling pairs in Results. the Degus were kept on a 12:12h light/dark cycle with
food and water available ad libitum. All procedures were conducted in accordance with the policies
of the biological facilities at the University of Toronto and all protocols were approved by the
University of Toronto Animal Care Committee.

Apparatuses

Behavioral recording took place in degus' homecages placed inside a 15''x19''x 24''plexiglass
chamber. The ceiling of the recording chamber was embedded with a Sony (Sony; Tokyo, Japan)
USB2 camera (video sampled at 30 Hz) and Sony lapel microphone (model number ECMCS3,
sampled at 44100 Hz). Data were collected using Bandicam multi-media recording software
(Bandisoft; Seoul, Korea). Social isolation was peformed by placing each degu into separate, 14"
x 11" x 9.5" chambers, either in the experiment room (for 1 and 45 min isolation periods) or the
vivarium (for 24 hr isolations). Footshocks used for the "acute stress" condition in Experiment 3
were delivered in a 18" × 17.5" × 15" conditioning chamber with a metal grid floor consisting of
5 mm thick stainless steel rods spaced 1 cm apart (Colbourne instruments; Holliston, USA).
145
146   *Protocol*.
147
Habituation: Prior to behavioral testing, degus were acclimated to the recording chamber. Degu
pairs were brought to the conditioning room and their homecage was placed into the soundproof
recording chamber. The pair spent 1 hour undisturbed in the chamber for each of 3 consecutive
days. Experimental protocols began immediately after acclimation except in the case of
Experiment 4, for which acclimation and prior experimentation took place at varying times prior
to the reunion.
154
Experiment 1: To test the effects of social isolation on interactive behavior, 9 pairs of female
cagemate degus (18 degus) were placed in separate cages for either 1 or 45 min in the recording
room, or 24 hr in the vivarium (the difference in holding room was logistical, as the shorter
intervals provided insufficient time for experimenters to shuttle both degus back to the vivarium,
and animals were required to be in the vivarium overnight). The isolation chambers were the same
size as the animals' homecages and were transparent and contained clean bedding. Notably,
although they were "isolated" in the sense of being unable to touch or interact with other
individuals, they could still see and hear other degus through the cage. Members of a dyad were
kept on different sides of the room to prevent direct communication between them. Following the
social isolation period, the original homecage was placed within the soundproof chamber for 20
minutes, during which time video and audio recordings were collected. The basic protocol is
illustrated in **Figure 1A**.

Experiment 2: To test the differences between social isolation and separation-without-isolation, each of 4 groups of 4 cagemate degus (16 degus) were first separated into two cages of only two degus (SEP). Following 24 hours, one member of each of the new cages was placed together in a clean 14" x 11" x 9.5" cage within in the recording chamber for a 20 min reunion session, after which the recorded pair was taken to the vivarium and kept together in that same cage for the subsequent 24 hours. The same was done with the remaining members of the two cages. Following 24 hours as reunited cagemates, the dyad was separated into two individual cages for a 1 minute period and then returned for a second, 20 minute reunion (CTL). Degus were then returned to separate cages and remained isolated for 24 hours, after which a final, 20 min reunion session was performed (ISO).

Experiment 3: To test the impact of stress on social interactions, the same degus from Experiment 1 were presented with two additional conditions. In one, cagemate dyads were separated into two different cages in two separate experiment rooms for a 45 minute period. During this time, following a 5 minute adaptation period, 8 sunflower seeds (a familiar, preferred food) were placed one at a time over a 2.5 minute period in each cage (RWD). When the 45 minute period had elapsed, they were placed together in their homecage, in the recording chamber for a 20 minute reunion session. In the other condition, cagemate dyads were separated into two separate rooms and placed in a conditioning chamber with a metal grid. Following a 5 minute baseline period, degus received 10, 1.0 mA footshocks lasting 2 s each over a 2.5 minute period (SHK). This shock level and pattern was based on a protocol we used previously to evoke contextual and observational fear learning [39], though to reduce total aversiveness we decreased the number of shocks by one-half. The 1 mA level offered a compromise for maximizing animal welfare while also eliciting a sufficient stress response (in rats, a series of 1.5 mA footshock stimulations evokes significantly higher levels and longer-lasting glucocorticoid responses compared with 0.5 mA [40]). After receiving footshock, degus were placed in separate cages for the remainder of the 45 minutes. To ensure that the reward and stress conditions did not contaminate subsequent exposures, degus were always tested with the neutral (NTL) 45 minute condition first, then RWD, and then SHK. The use of a consistent order could raise questions as to whether behavioral effects of SHK may be due to order effects of the exposures. The results ultimately showed that SHK differed more

198 from RWD than from the NTL condition in ways that were consistent with hypotheses but
199 inconsistent with an order effect.
200
Experiment 4: To investigate how interactive behavior was impacted by social novelty, potentially
associated with relationship establishment, degus were united with unfamiliar, stranger
conspecifics following 24 hours of isolation. Animals used for Experiment 4 were drawn from
those from Experiments 2 (12 degus) or were members of a separate set of animals that had
received pre-exposures and were subsequently used for a separate study (4 degus). Degus that had
been involved in another experiment had been pair-housed with a cagemate for at least one week
after the prior experiment had ended.
208
209 The sequence of reunion sessions for each dyad are illustrated in Table 1

Dyad	1-1min	1-45min	1-24hr	2-SEP	2-CTL	2-ISO	3-45min	3-RWD	3-SHK	4-STR
1	3	2	1				4	5	6	
2	1	3	2				4	5	6	
3	1	3	2				4	5	6	
4	1	2	3				4	5	6	
5	1	2	3				4	5	6	
6	1	2	3				4	5	6	
7	1	2	3				4	5	6	
8	1	2	3				4	5	6	
9	1	2	3				4	5	6	
10				1	2	3				
11				1	2	3				
12				1	2	3				4
13				1	2	3				4
14				1	2	3				4
15				1	2	3				4
16				1	2	3				4
17				1	2	3				4
18										1
19										1

210
211 **Table 1. Sequence of reunion exposures for each dyad.** Conditions are labeled as Experiment
212 number – condition. Numbers reflect the order that the dyad experienced the condition. Dyads 18
213 through 21 are listed as "7" because these animals had undergone testing in the Experiment 1 and
214 3 protocol but data were not used due to technical issues.
215

Behavioral analysis

Software: Analysis relied on several software packages. Scoring of physical behavior was
performed using BORIS (Behavioral Observation Interactive Research Software; Torino, Italy
[41]), which allows users to log events during video playback. Scoring of vocalization behavior
was performed using Audacity (Audacity v.2.1.2; Pittsburg, USA), as well as Sound Analysis Pro
(SAP), a vocalization detecting software developed for segmenting syllables and vocalization
structure in birdsong [42]. All other analyses were custom-written in MATLAB, as described
below and provided at https://github.com/ninsel/DeguReunionMotivation.

Physical interactions: Each video recording was manually scored by blind raters using BORIS.
Blinding was done by using coded filenames. They included four general types: *agonistic*
encounters (bites, characterized by face-to-body contact of one degu with the other making a
startled movement away; mounting, involving one degu placing forepaws on the other while the
other made avoidance movements; wrestling, characterized by rapid movements of both bodies
and usually accompanied by other aggressive actions; punching/boxing, with degus sometimes
rearing on hindlegs; body-slams, in which degus kicked with their hind legs, and any other
apparently aggressive encounter), *allogrooming/body sniffing* (one degu's nose touching and/or
interacting with the neck or body of the other), *rear-sniffing* (a.k.a. anogenital investigation, where
one degu's nose was near and directed toward the base of the tail of the other), and *face-to-face*
interactions (one degu's nose near and directed toward the nose or mouth of the other). Ethograms
were created based on a relatively minimal set of highly distinguishable behaviors that were
observed across most sessions, with a goal of maximizing simplicity and inter-rater reliability. To
assess inter-rater reliability, several sessions were scored by more than one rater, and Cronbach's
alpha was calculated to be 0.97 (very high). For each scored behavior, the identity of the animal
was also included. Often behaviors overlapped: e.g., one degu may initiate rear-sniffing and the
other may then reciprocate before the first behavior ended. In the case of face-to-face, the action
was necessarily mutual, but these were still labeled with the identity of the degu that initiated the
interaction. One potential problem with behaviors overlapping in time is that the sum of all
behaviors performed by one degu added to the sum of all behaviors performed by the other could,
in principle, exceed the total time of the session. Analyses therefore combined interaction time

from each degu by considering all time during which *either* one degu *or* the other was engaging in

the action (i.e., the union of time, rather than the combined total of both).

Comparisons of physical interactions between experimental conditions were analyzed using the

following three tests: 1) Total time dyads spent interacting, and total time engaging in different

interaction types. As noted above, interaction time was primarily analyzed as all time that either

one or the other degu of a dyad was engaging in an interactive behavior (body-sniffing, face-to-

face, etc.). 2) Temporal distribution of interactions across the session, primarily quantified and

illustrated by cumulative distributions of interaction over time. Cumulative distributions of

interactions were computed with a 2 s bin for each session (discussed below in *Statistics*). 3)

Temporal relationships between interaction types, quantified and illustrated by the cross

correlograms. Cross correlograms evaluated the time-lagged coincidence of one interaction

relative to another. To compute these, a histogram was generated: times preceding zero (left side

of the x-axis) showed the incidence of one ("target") action relative to when another ("reference")

action was initiated, while times following zero (right side of the x-axis) showed the incidence of

the target action relative to when the reference action had terminated. Actions that took place

simultaneously were therefore not included in the histogram. These histograms were then

normalized by the number of reference actions, thereby converting incidence into rates. For

interpretability, the histograms were further normalized by dividing by the proportion of time

during the session that the target action took place. In other words, values of "1" in the cross

correlation meant there was no observable temporal relationship between the two interaction types.

Importantly, this means that y-axis units were not correlations coefficients, but normalized rates.

To assess significance, all values from -10 to -1 s (before onset) and from 1 s to 10 s (after offset)

were averaged within each session, and a paired t-test was performed to assess whether values

were A) consistently higher than 1, and whether values were B) consistently higher than the

average of ten surrogate datasets, in which onset/offset times of the comparison action type were

randomly shuffled within a -5 to 5 minute interval. Only those cross correlations that met both

criteria at an alpha value of 0.05 were considered significant. In cases where comparisons were

being made between conditions, a paired or two-sample t-test was used. Issues with multiple

comparisons are addressed below under the *Statistics* subheading.

Vocal interactions: Vocalizations were scored by loading audio files into Audacity and using both
audio and visual spectrograms to identify vocalization start and end times. Blinding of vocalization
scoring used the same coded filenames as used for coding physical interactions. To reduce noise,
the Audacity "noise reduction" feature was used, which creates a filter based on a selected audio
baseline period (baseline was several seconds of audio recorded prior to degus placed in the
recording chamber). Experimenters scored vocalization by loading individual sound clips
generated by Audacity into SAP, where spectrograms and summary statistics (amplitude,
frequency modulation, duration) were compared with those presented by Long (2008; [34]). Inter-
rater reliability was estimated using Cronbach's alpha = 0.96 (very high). Comparisons of vocal
behavior between sessions relied on the following tests: 1) total vocalization rates (number of
syllables over session time), and rates of specific, manually scored vocalization types, and 2)
differences in the profile of vocalization types detected through unsupervised clustering of the
syllables (repeated-k-means).
291
We chose to use an unsupervised clustering method over a supervised algorithm out of interest for
a method that was complimentary to, rather than dependent on, manual-labeling. Unsupervised
methods have the potential of identifying syllable clusters based on features that human listeners
are not as well tuned to, thereby revealing syllable types that we would not have known to look
for. The sequence of steps involved in this analysis consisted of the following: 1) First, a set of 31
features was computed for each identified syllable. The first feature was syllable duration, while
the remaining thirty were dominant frequency, spectral dispersion, and amplitude across each of
ten deciles of time. These values were standardized across all syllables by converting each of the
31 features to a z-score relative to all other recorded vocalizations, and then weighted (each
multiplied by a scalar) such that syllable duration counted for 10 times each dominant frequency
value, 20 times each spectral dispersion value, and 40 times each amplitude value. The weightings
were chosen based on subjective assessment of which features should play the strongest roles in
syllable classification, an approach that has precedence in the field [43]. 2) To reduce the
dimensionality of the feature space, and also account for correlations between features (e.g.,
adjacent deciles are likely to be correlated), we converted features to principle components. 3)
Describing each syllable by its first 10 principle components (accounting for 99.5% of the
variance), we then entered all recorded syllables across all experiments into a k-means clustering

method. K-means clustering was used to subdivide the pool into 2 clusters, 3 clusters, 4, and on
up to 30 clusters (10 replicates each, with 20 repetitions each to assess cluster stability). 4) For
each syllable cluster, the number of cases in each session was identified. The proportional rate of
syllables in one condition could then be compared with another condition by computing an effect
size (Cohen's D) and a significance value (p value from a paired t-test when considering the same
animals, or two-sample t-test in Experiment 4 when comparing different dyads). In some cases,
more than one cluster might show a strong difference, with some being sub-divisions of others. To
identify only the *independent* groups of syllables that differed between conditions, only the
strongest cluster was considered, then all syllables from that cluster were removed and significance
testing between clusters was performed again, across multiple iterations until no significant
clusters were identified. This k-means clustering method was modified from a similar method used
in previous work from our group, which was designed to evaluate whether there were subdivisions
of distress/alarm calls [39]. The previous version differed in only minor respects; e.g., only four
features were included, so no dimension reduction was needed, and notable clusters were identified
with their p-values rather than effect sizes. Additionally, since there was expected to be fewer
categories, the earlier version did not repeated the process iteratively after removing syllables of
the strongest category.

One acknowledged risk of the repeated-k-means analysis is that if syllables are randomly
distributed across all sessions, one might expect to observe significant clusters roughly 5% of the
time, given an alpha of 0.05. Empirically this was not the case; however, taking this concern into
account, identification of clusters was treated as purely exploratory. For partial validation of the
results, we performed an empirical assessment of the individual clusters (including consistency of
the cluster across the 20 repetitions, as measured by the Rand index, and how the cluster mapped
onto manually-labeled vocalizations) and present figures of effect size distributions for qualitative
comparisons between experiments. All identified clusters were reported in Results, and to reduce
biases from multiple-comparisons across repetitions only the median significance values and effect
sizes across the 20 repetitions were reported.

Although behaviors were performed within a soundproof chamber, it was still possible that
vocalizations below a certain volume were undetected by out methods. To assess whether this

might be the case, vocalization syllables were segmented into 10 ms bins and the peak amplitude
was measured relative to the dominant frequency at that amplitude. This was first done across
syllables recorded in several, adult female degu "reunion" experiments which used the same
methods (24,535 syllables). An initial look at the data revealed that a substantial proportion of
syllables (approximately 43%) were not loud enough, or of high enough frequency, to exceed the
volume of the ambient noise at 400 Hz (the lower cutoff that was used to compute dominant
frequencies). Syllables were therefore separated into two populations: those with amplitude peaks
that exceeded a 400 Hz cutoff, and those that did not. Amplitudes and dominant frequencies of
those that reached threshold are illustrated in **Supplementary Figure 1A**; of these, only one small
grouping of low-amplitude, low-frequency syllables, almost exclusively labeled as chitters,
appeared to have a distribution with the lower-end cut-off from detection. Of those that did not
stand out from the noise, also most often labeled as chitters, the distribution also appeared to be
positively skewed (**Supplementary Figure 1B**). These data suggest that only a minority of
vocalizations failed to reach detection threshold, and most of these were likely to be "chitters."
354
Statistics
Parametric statistics are used in nearly all comparisons, including repeated measures ANOVAs
(used for all one-way tests), two-way ANOVAs (to evaluate interaction effects), and both paired
(for same-dyad comparisons) and two-sample (for between-dyad comparisons) t-tests. In the case
of manually scored vocalizations, variance across dyads was not distributed normally and
therefore a Wilcoxon rank sum test was used. Generally, sample sizes were low, and the high
probability of false negatives is addressed in discussion.
362
Many analyses addressed specific predictions in a given experiment; however, there were several
circumstances in which multiple comparisons were used in an exploratory fashion, raising the
possibility that false positives may arise by chance. The first of these is when testing whether
each individual behavior (agonistic, grooming/body-sniffing, rear-sniffing, face-to-face) is
affected by conditions within an experiment. In this case, we employ a Bonferroni correction
when assessing statistical significance (alpha $= 0.05/4 = 0.0125$). The second circumstance is
when looking at temporal interactions for each relationship between behaviors (e.g., rear-sniff
before grooming, rear-sniff after grooming, etc). Here, cross correlations had to reach two

criteria (changes from baseline and differences from a surrogate dataset) to be considered

significant. The analysis is performed with acknowledgment of a false positive rate of up to 5%,

though most comparisons elicited much lower p-values, and causal inferences focus on those

relationships that showed consistent and meaningful patterns between experiments. Finally,

multiple comparisons are used when assessing differences between conditions over different,

cumulative time intervals of the reunion session. In this case, a t-tests was used to compare each

of 600, 2 s intervals between conditions. This method was selected as an exploratory approach

that made no *a priori* assumptions on the distribution of behaviors over time, or how that may

vary between conditions (as a contrast to model-fitting, which would require would require either

overfitting a high number of parameters or assumptions about the distributions). To best address

the possibility of false positives, only extended intervals are reported, along with session times

when differences reach their peak, and the method is complement with a secondary, directed

analysis comparing the slopes of the cumulative distributions.

Results

Experiment I: Isolation time

We first sought to confirm that degus become increasingly motivated to interact following

increasing intervals of social isolation. Nine, adult female degu cagemate dyads were pre-exposed

to a behavioral recording chamber for one hour each day over three days and then, on sequential

days, isolated from one another in separate cages for 1 min, 45 min, and 24 hr (**Figure 1A**).

Following each isolation period, dyads were brought back together for 20 minutes in the recording

chamber while video and audio data were collected. Consistent with expectations, there was a

significant impact of isolation time on the time degus spent physically interacting (repeated-

measures ANOVA, $F_{2,16} = 4.93$, p = 0.021, **Figure 1B**). The effect of isolation was particularly

strong earlier in the session, generally peaking around 3 to 4 min (**Figure 1C**; using an alpha of

0.05, differences between 1 and 45 min conditions were observed for the entire interval between

78 and 422 s, peaking at 224 s: $t_8 = 4.79$, p = 0.0014; differences between 24 hr and 45 min observed

for the entire interval between 30 s and 1198 s, peaking at 282 s, $t_8 = 9.48$, p = 1.26×10^{-5}; and

differences between 24 hr and 1 min persisted throughout the session, significant by 20 s and

peaking at 194 s, $t_8 = 10.30$, $p = 6.81 \times 10^{-6}$). Interactions were parsed into four types: agonistic

(which included mounting), allogrooming/body sniffing, rear-sniffing (i.e., anogenital

investigation), and face-to-face. No interaction was observed between condition and interaction

type (2-way ANOVA, $F_{6,96} = 0.53$, $p = 0.78$). Examining each interaction type individually,

significant differences in interaction time (alpha $= 0.05/4 = 0.0125$, Bonferroni correction) were

observed for rear-sniffing ($F_{2,16} = 5.9$, $p = 0.012$) and face-to-face interactions ($F_{2,16} = 6.78$, $p =$

0.0074), with a trend also observed for grooming/body sniffing ($F_{2,16} = 3.58$, $p = 0.052$; **Figure**

1D). The time spent in agonistic behavior did not appear to vary across the conditions ($F_{2,16} = 0.23$,

$p = 0.80$).

When degus were not interacting with one-another, they were typically rearing and looking upward

along the walls or sniffing and sometimes digging into the chamber bedding. Self-grooming was

the only non-interactive behavior systematically scored, which significantly increase with

preceding isolation time ($F_{2,16} = 4.95$, $p = 0.021$). Although the study was not designed to compare

behavior in sibling versus non-sibling cagemates, there appeared to be no meaningful difference

in the effect size of the 24 hr versus 1 min isolation in non-siblings (n = 3, Cohen's D = 0.95)

compared with siblings (n = 6, Cohen's D = 0.82).

Figure 1. Effects of social isolation time on reunion behavior. **A)** Illustration of experimental design. Degus (n = 9 dyads) were separated from cagemates for 1 min, 45 min, or 24 hours then interactions were examined in a testing chamber. After being reunited for 24 hours the process was repeated. **B)** Proportion of time engaged in physical interactions (y axis) increased with the amount of preceding time of isolation (x-axis). **C)** Cumulative distributions of physical interactions over session time. Thicker lines are means, thin lines s.e.m. Higher levels of interaction in the 24 hr condition were seen within the first minute and remained high for the first 5 minutes. Inset shows full, 20 min of session time; although interaction time in the 24 hr condition remained higher, interactions in the 1 and 45 min conditions were more distributed across session time. **D)** Line graphs of interaction time across types of physical interaction; no interaction between type and condition was observed. **E)** Vocalization rates (y-axis) significantly increased across conditions (x-axis). **F)** Vocalization rates were consistently higher in the 24 hr condition across the 15 vocalization categories tested. **G)** Heatmap of identified syllable clusters with the strongest difference between the 24 hr and 1 min conditions, across k-means clusters repeated for 2-30 cluster divisions (columns), iterated after removing syllables from the best cluster (rows). Plot on

top is an errorbar showing variance of the first (top-row) across 20 repetitions. The hot-spot in the
heat map shows that when syllables were divided into 20 clusters, one cluster, consistent across
the 20 repetitions, showed a particularly high effect size and was significantly, across dyads, more
highly expressed in the 24 hr condition. **H)** Map of t-SNE projection of vocalization features. The
identified cluster on this projection is represented in blue dots. **I)** Proportion of manual label types
across syllables across all studies (grey) is plotted alongside the proportion of labels for the
identified cluster. Syllables in this cluster were predominately labeled as "chaffs", and almost
never as "chitters".
445

To assess the structure of behavior, we examined whether there were temporal relationships
between physical behaviors. We had no a priori hypotheses on how patterns might vary with
increasing isolation time, and anticipated that lower levels of interaction in short (1 min)
interaction conditions would impact the validity of a direct comparison between conditions. Initial
analyses therefore pooled all interactions across the 1 min, 45 min, and 24 reunion conditions to
identify a base-set of relationships, providing a framework for interpreting interactions observed
in subsequent experiments. Cross-correlograms were generated for each pair of interaction types
(e.g., face-to-face and rear-sniffing, rear-sniffing and allogrooming/body sniffing, etc), where
values on the negative side of the x-axis (time) represented the occurrence of an action type *before*
the onset of another "reference" type, and values on the positive side of the x-axis represented the
occurrence of an action type *following the offset* of the reference type. A significant temporal
relationship was determined if the cross correlogram values were A) consistently higher than
baseline (i.e., systematically higher than 1), and B) consistently higher than surrogate datasets, in
which onset/offset times of the comparison action were randomly shuffled within a -5 to 5 minute
interval (see Methods). In the initial analysis performed on Experiment 1 data Significant
relationships (described as comparison relative to reference) were identified for: 1)
allogrooming/body sniffing preceding rear-sniffing (above baseline: $t_{16} = 2.84$, p = 0.010, above
shuffled: $t_{16} = 4.12$, p = 5.0 x 10^{-4}), 2) agonistic preceding and following agonistic (above baseline:
$t_{16} = 2.49$, p = 0.024 and $t_{16} = 3.18$, p = 0.0059 respectively, above shuffled: $t_{16} = 2.59$, p = 0.020
and $t_{16} = 3.33$, p = 0.0042), 3) rear-sniffing following allogrooming/body sniffing (above baseline:
$t_{16} = 2.45$, p = 0.0024, above shuffled: $t_{16} = 5.68$, p = 0.0010), 4) allogrooming/body sniffing
following face-to-face (above baseline: $t_{16} = 3.40$, p = 0.0023, above shuffled: $t_{16} = 4.13$, p = 4.0
x 10^{-4}) 5) face-to-face preceding and following allogrooming/body sniffing (above baseline: $t_{16} =$
4.65, p = 1.0 x 10^{-4}, $t_{16} = 2.52$, p = 0.018 respectively, above shuffled: $t_{16} = 5.88$, p = 3.94 x 10-6,

$t_{16} = 4.16$, p = 3.0 x 10^{-4}). Averaged cross correlograms are presented in **Supplementary Figure**

2, with specific examples provided in subsequent sections.

As with physical interactions, isolation also appeared to increase vocalizing, measured by the

frequency of syllables over session time ($F_{2,16} = 4.18$, p = 0.034, **Figure 1E**). To unpack this result,

vocalization syllables were manually and automatically parsed into types. Manual classification

included 32 categories, guided by previous research on degu social vocalizations[34]. Of these,

only approximately 14 vocalization types were observed at least once, on average, per session—

for the purposes of analysis, all others were included within a 15^{th} "mixed" category. Each

vocalization type showed high variability between dyads and sessions, and no significant

interaction was observed between condition and vocalization type (2-way ANOVA, effect of

condition: $F_{2,360} = 3.04$, p = 0.049, effect of type: $F_{14,360} = 5.22$, p = 5.3 x 10^{-9}; condition x type,

$F_{28,360} = 0.38$, p = 1.00, **Figure 1F**).

Differences between manually-scored syllable categories were sometimes subtle, and there was

likely discrimination ambiguity between certain types (illustrated by the overlap of the different

classes in feature space, **Supplementary Figure 3**). We therefore also used an unsupervised

clustering method to assess whether clusters of syllables with shared features might be

differentially expressed between conditions. The repeated-k-means approach is detailed in

Methods. Briefly: syllables were divided into clusters using a k-means algorithm (2 to 29 clusters,

repeated 20 times, all across 10 iterations of removing the best cluster from the previous) and

syllable clusters that made-up a higher proportion of those expressed during one condition

compared with another were identified. For the purposes of this investigation, consistency across

dyads, was emphasized over magnitude of difference, and proportion of syllable type within a

session emphasized over total numbers. The method was considered an exploratory complement

to manual labeling that, aside from syllable start and end times, was unbiased by the experimenter.

Comparing the 24 hr to 1 min isolation conditions, only one cluster stood out as showing a

systematic (across dyad) difference (**Figure 1G & H**). This group of syllables was most precisely

isolated when the pool was divided into 20 clusters (median p across repetitions = 0.013, median

Cohen's D = 2.7; consistency of cluster across iterations was also high: average adjusted random

index = 0.96, where 1 is perfect cluster overlap). This syllable type was often manually scored as
a "chaff" (**Figure 1I**) and was distinguished by relatively high average dominant frequencies
around 6000 Hz, slightly lower-than-average entropy or frequency dispersion (i.e., more
"scratchy") and slightly higher than average amplitude. These vocalizations were not more
temporally associated with specific interactive behaviors than average vocalizations, although
prior work has suggested that chaffs may be more agonistic in nature [34].

To summarize the findings of Experiment 1, we first confirmed our prediction that adult, female
degu dyads interact with one-another more following increased isolation time. The increased
interaction did not significantly differ across interaction types, though notably there was no sign
of increases in agonistic behavior. Pooling conditions together, we found a number of significant
temporal interactions between interaction types, providing a starting point for more directed
analyses of behavioral structure in future experiments and studies. Vocalizing also increased with
longer isolation times. While differential increases across vocalizations were not observed with
increased isolation time, this could have been impacted by low power, and exploratory analyses
did identify one cluster of chaff-type syllables that might have been differentially increased in the
24 hr condition.

Experiment II: Isolation vs. Separation

The second experiment examined whether dyads interact differently when motivated by prior
social isolation, as compared with prior separation from one-another without being isolated (i.e.,
"general" versus "stimulus specific" social motivators). This was tested by comparing interactions
in reunited dyads following three manipulations (illustrated in **Figure 2A**): 1) Cages of four female
degus were split into two cages for 24 hours, thereby causing separation of two dyads (SEP); 2)
all individuals were socially isolated for 24 hours (ISO); and 3) all individuals were isolated for
only a 1 min interval (CTL).

Figure 2. Comparison of isolation versus separation on reunion behavior. **A)** Illustration of experimental design. Four cages of four (n = 8 dyads) were split into two cages for 24 hours, separated dyads were then united for reunion SEP tests. Subsequently these dyads were isolated for 1 min and then 24 hours for CTL and ISO tests respectively, reunited for 24 hrs between each. **B)** Proportion of time degus engaged in physical interactions was at least as high in SEP as ISO,

but lower in CTL. **C)** Cumulative distribution of physical interactions over session time (thicker
lines are means, thin lines s.e.m.). Although interactions in SEP were high over the 20 min,
following isolation degus were much more likely to interact during the first minute of the reunion
session. **D)** Interaction time for different types of physical interactions. There were no systematic
differences between SEP and ISO, though patterns hint at higher rear-sniffing and lower grooming
in SEP relative to ISO. **E)** Cross correlogram of levels of allogrooming/body-sniffing (traces)
relative to rear-sniffing. Negative time values are prior to rear-sniffing initiation, positive are after
termination of the behavior. Both ISO and the 24h condition from Experiment 1 showed higher
allogrooming before rear-sniffing compared with SEP. **F)** Cross correlogram of the levels of face-
to-face interactions relative to allogrooming/body sniffing. A significant increase in face-to-face
contact before allogrooming was observed for all conditions. **G)** Heatmap of identified syllable
clusters with the strongest difference between ISO and SEP, with increasing cluster divisions
across columns and iterations after removing syllables from the best cluster down rows. Plot on
top is an error bar showing the variance the first (top-row) across 20 repetitions. **H)** Map of t-SNE
projection of vocalization features. One cluster was identified of syllables that differentiated
between SEP and ISO conditions (blue points). **I)** Proportion of manual labeled types across all
syllables (gray trace) and for only those syllables in the identified cluster (blue trace) show that,
as with the cluster identified in Experiment 1, these were chaff-type syllables.
553
554

Degus interacted more in SEP than they did in CTL (repeated-measures ANOVA effect of
condition $F_{2,14} = 4.93$, p = 0.024; SEP vs. CTL, paired t-test, $t_7 = 3.94$, p = 0.0056; **Figure 2B**),
and there was no detectable difference between ISO and SEP ($t_7 = 0.79$, p = 0.45). Significant
differences also were not detected between ISO and SEP in any of the individual behaviors
(agonistic, body-sniffing, rear-sniffing, face-to-face; paired t-tests, p >> 0.0125). Importantly,
while there were no overall differences between these two conditions over the full 20 min period,
differences were observed in interaction levels during the start of the session. The differences were
most evident between 40 to 70 s, as illustrated in **Figure 2C** and identified quantitatively using a
series of paired t-tests, revealing a window of 2 s time bins between 42 and 66 s with a 95%
likelihood that the two curves differed. At 60 s, for example, ISO degus had interacted on average
20% of the time, while SEP degus had interacted an average of only about 5% (paired t-test, $t_7 =$
2.47, p = 0.043, Cowen's d = 1.16). There was a corresponding statistical trend for steeper initial
slopes in cumulative interaction behavior in the ISO condition compared with more evenly paced
interactive behavior in the SEP condition (paired t-test on the latency to half-amplitude, $t_{7 =} 2.10$,
p = 0.074). In summary, although interactive behavior between SEP and ISO conditions did not
show a net difference over the 20 min period, interactions tended to take place earlier following

isolation (ISO), while being more evenly distributed over the session following only separation

(SEP).

To help confirm that overall patterns were not due to the order that conditions were presented,

direct comparisons were made between interaction levels in Experiments 1 and 2. Interaction

levels in SEP (first exposure) were significantly higher than those of 1 min, Experiment 1

(including only dyads whose first exposure was 1 min isolation: two-sample t-test, $t_{14} = 2.35$, p =

0.034). Likewise, CTL showed significantly lower interaction levels than those following 24 hr

isolation in Experiment 1 (equivalent to ISO; only dyads with 24 hr as the second or final exposure

included; $t_{14} = 2.28$, p = 0.039).

By comparing the temporal relationships between physical interactions in ISO and SEP we found

a significant decrease in the proportion of rear-sniffing cases preceded by allogrooming/body-

sniffing (paired t-test, $t_6 = 3.71$, p = 0.010). Importantly, patterns in ISO replicated those observed

in the 24 hr condition of Experiment 1 (**Figure 2D**). In other words, while there were no apparent

systematic differences in the total levels of face-to-body contact between conditions, degus who

had been previously isolated were more prone to engage in allogrooming ("nuzzling") immediately

prior to rear-sniffing. This may reflect a tendency toward affiliation in animals that had been

previously deprived of social stimulation. All other temporal relationships were highly consistent

between ISO and SEP, and consistent with those observed in Experiment 1 (e.g., face-to-face

contact prior to allogrooming/body sniffing, **Figure 2E**).

As was observed in Experiment 1, vocalization rates were highly variable between sessions and

dyads. There were no significant differences across the three conditions (repeated-measures

ANOVA, $F_{2,14} = 2.27$, p = 0.14); however, it seems likely that lower vocalizing in CTL would be

observed with more statistical power (paired t-test between ISO and CTL, $t_7 = 2.30$, p = 0.055,

Cohen's D = 0.78). Given the low power and high variance, it is also unsurprising that we observed

no interaction between condition and vocalization type (2-way ANOVA, effect of condition: $F_{2,300}$

= 3.94, p = 0.020, effect of type: $F_{14,300} = 19.07$, p = 7.03×10^{-34}, condition x type: $F_{28,300} = 1.25$, p

= 0.19; **Supplementary Figure 4**). More generally, however, there appeared to be a higher ratio

of non-chitters to chitters in SEP compared with ISO (less than half the number of non-chitters to

chitters in ISO, and around twice the number of non-chitters to chitters in SEP; a non-parametric,
Wilcoxon rank-sum test was used because ratios did not distribute normally, $p = 0.0080$). This was
notable in that a similar but stronger pattern was observed in strangers (Experiment 4, below).

Our repeated-k-means clustering method revealed only one cluster with different proportional
expression in SEP versus ISO conditions (**Figure 2F & G**). The syllable type was most strongly
revealed when all syllables were subdivided into 8 clusters (d.f. = 7, median $p = 0.038$, Cohen's D
= 1.01, across 20 repetitions). Unexpectedly, while the cluster was highly overlapping with the
one identified in Experiment 1, and typically labeled as "chaffs", these syllables were expressed
more in SEP than ISO (**Figure 2H**). The syllables of the cluster identified in Experiment 1,
contained within this larger grouping, were themselves also more common in SEP than ISO (d.f.
= 7, $p = 0.037$). It is worth noting that in both cases—24 hr isolation in Expeirment 1 and ISO in
Experiment 2—this was the first time dyads had been reunited following 24 hrs apart. Comparing
ISO and CTL, several non-overlapping clusters reached criteria, but the effect magnitudes were
very low relative to the other clusters identified and did not lend themselves to straightforward
interpretation. Two of these clusters showed proportionally higher activity in ISO but were
distributed across several manually-labeled categories (e.g., chaffs, groans, chirps, tweets, whines;
d.f. = 7, median $p = 0.023$ and $p = 0.027$), another, of manually labeled tweets and chaffs, was
found to be proportionally more common in CTL (median $p = 0.027$).

In summary, Experiment II confirmed our prediction that adult female degus were highly
motivated to interact with one-another even after being only separated, and not isolated, for 24
hours, and also confirmed our prediction that the types of interactions among separated individuals
differed from those between isolated animals. SEP reunions included more non-chitter to chitter
vocalizations than ISO, and ISO elicited more allogrooming/body sniffing before rear-sniffing
(consistent with data from Experiment 1). Exploratory analyses identified a chaff-type vocalization
that may have been relatively increased in the separation compared with isolation condition.
Counterintuitively, this was effectively the same vocalization as was found to increase with 24 hr
isolation in Experiment 1, the one common factor between the two being that both Exp II ISO and
Exp I 24 hr conditions were the first time dyads had been reunited after extended (24 hrs) apart.

Experiment III: Acute Stressor vs. Reward

One factor that might have played a role in the differences between isolation and separation is the

stress associated with isolation [17–22]. To better understand the ways in which stress modulates

female social behavior, we examined behavior following a short period of exposure to an acute,

unpredictable stressor (a series of 10, uncued footshocks spaced over 2.5 minutes within a 45 min

isolation period; SHK), as compared against the effects of 45 minute isolation in a neutral

environment (NTL), and exposure to an unpredictable reward (un-cued delivery of sunflower seeds

over a 2.5 minute period within a 45 min isolation period; RWD; **Figure 3A**).

Figure 3. Effects of exposure to an acute stressor on reunion behavior. **A)** Experimental design.

Degus (n = 9 dyads) were isolated for a 45 min period during which they were exposed to

unexpected footshock (SHK), unexpected reward (sunflower seeds; RWD), or neither (NTL). **B)**
Proportion of time dyads spent interacting in each condition. **C)** Cumulative distribution of time
spent interacting across the session (thicker lines are means, thin lines standard error). Footshock
stressor increased social interaction earlier during the testing, though this behavior leveled off in
the latter half. **D)** Four line graphs depict the proportion of time degus spend interacting 10 min
into the testing session across each of the three conditions. There was a general pattern for higher
interaction across behavior types in SHK at the 10 min mark of the reunion. **E)** Vocalization rates
(syllables per second) across conditions. Vocalizations rates were highly variable between dyads.
F) Cumulative time spent vocalizing (y-axis) is plotted against session time (x-axis). Although
there was high variance between dyads, the general pattern was for vocalizations to begin after
several minutes of physical interactions, and to subsequently approximately follow the temporal
patterns of physical interactions. **G)** Vocalization rates are plotted across each of 15 syllable
categories. High variance made interpretation difficult, but generally rates of chitters seemed
particularly high in NTL relative to SHK and RWD. **H)** Two dimensional t-SNE projection of
syllable features, including those for all syllables (gray points) and those of the cluster of syllables
found to be expressed more frequently in SHK. **I)** Proportion of syllables manually labeled across
different category types, including all syllables (gray trace) and those syllables within the
identified cluster (blue trace). The data suggest that chirp- and tweet-type vocalizations may be
more common in degus recently exposed to a footshock stressor.

Physical interaction was affected by a preceding stressor primarily with respect to how the actions

were distributed over time. When taking into account the entire 20 minutes, differences between

SHK, NTL, and RWD conditions were hinted at by a statistical trend (repeated measures ANOVA,

$F_{2,16} = 3.93$, p = 0.074; **Figure 3B**), although these numbers were largely diluted by the high

variance of the neutral 45 min isolation condition, as a clear difference was observed in interaction

levels between SHK and RWD ($t_8 = 3.73$, p = 0.0058). Looking only at the first half of the session,

however, it was clear that the acute stressor did increase initial interactive behavior (**Figure 3C**).

Given an alpha of 0.05, there were extended stretches of time in which cumulative interaction time

differed between conditions, with differences being maximal between 4 to 6 min (SHK versus

NTL, 28 to 100 s and 146 to 958 s, maximum difference observed at 353, $t_8 = 5.30$, p = 7.2 x 10^{-}

^{4}; SHK versus RWD, from 42 s to the end of the session, maximum at 255, $t_8 = 4.10$, p = 0.0013).

Paralleling differences between ISO and SEP in Experiment 2, this was accompanied by a

statistical trend for steeper initial slopes in the cumulative interaction time (paired t-test on the

latency to half-amplitude, $t_8 = 2.15$, p = 0.064). There was no evidence that differences between

conditions differed between interaction types (2-way ANOVA, effect of condition: $F_{2,96} = 3.8$, p =

0.026, effect of type: $F_{3,96} = 10.13$, p = 7.4 x 10^{-6}, condition x type: $F_{6,96} = 0.48$, p = 0.82); however,

consistent with the notion that the stressor may have differentially increased affiliative compared

with agonistic or investigative behavior, during the first half session SHK sessions included much

higher levels of allogrooming/body sniffing and face-to-face contact compared with both NTL

(paired t-test, grooming: t_8 = 4.12, p = 0.0033, face to face: t_8 = 3.29, p = 0.011) and RWD

(grooming: t_8 = 3.30, p = 0.011, face-to-face: t_8 = 3.41, p = 0.0092; **Figure 3D**). No differences in

the temporal relationships between physical interactions were found between any condition (alpha

= 0.05), in all three cases levels of allogrooming-before-rear-sniffing appeared to be intermediate

between those observed in ISO/24 hr and SEP conditions.

Similar to Experiments 1 and 2, vocalizations were highly variable across sessions, and no

differences were observed between SHK, NTL, and RWD (repeated measures ANOVA on

vocalization time: $F_{2,16}$ = 1.24, p = 0.32; **Figure 3E & 3F**). Nor was there a clear differential effect

of condition on different types of manually-scored vocalizations (two-way ANOVA, effect of

condition: $F_{2,315}$ = 1.85, p = 0.16, effect of type: $F_{14,315}$ = 2.06, p = 0.014, condition x type: $F_{28,315}$

= 1.1, p = 0.34; **Figure 3G**). Repeated-k-means revealed one cluster of syllables differentially

increased in SHK relative to NTL (median p across iterations = 0.015, **Figure 3H**) and an almost

completely overlapping cluster that increased in SHK relative to RWD (median p = 0.033), while

no clusters met criterion when comparing NTL and RWD (alpha = 0.05). These syllables were

relatively short in duration (89±14 ms) and were often manually labeled as tweets or chirps (**Figure**

3I).

Summarizing Experiment 3, data confirmed the prediction that footshock stress would increase

interaction time, though unexpectedly this was only true for the first half of the session. Likewise,

the only common pattern between 24 hr isolation of Experiments I and II and SHK of Experiment

3 was the increased early-session interaction; neither a higher incidence of chitters nor increased

allogrooming/body-sniffing before rear-sniffing were observed in SHK. Exploratory analyses

identified only a potential increase in tweet/chirp type vocalizations in SHK, and no consistent

differences in vocalizations between NTL and RWD

Experiment IV: Stranger vs. Cagemate

 Results from Experiments 1 & 2 were consistent with the possibility that reunion behavior was

714   partly driven by relationship renewal. To further investigate behaviors associated with relationship

715   formation, we compared behavioral patterns in stranger dyads united after 24 hours of isolation

716   (STR) with the the 24 hour cagemate isolation data from Experiments 1 and 2 (CAG1, CAG2;

717   **Figure 4A**).

Figure 4. Comparison of social behavior between stranger and cagemate dyads. **A)** Illustration of protocol. Cagemates were isolated for 24 hour and then brought together with an unfamiliar conspecific. **B)** Proportion of time degus spent interacting. There was a statistical trend for stranger pairs to be more interactive than cagemates (STR = stranger, n = 8 dyads; CAG1 = cagemates Experiment 1, n = 9 dyads; CAG2 = cagemates Experiment 2, n = 8 dyads) **C)** Proportion of time dyads of each group engaged in different types of physical interaction. Although there was a pattern for higher levels of interaction between strangers across all types, a strong difference was only observed for rear-sniffing. **D)** Cumulative time dyads in each group spent engaged in rear-sniffing behavior (thicker lines are means, thin lines s.e.m.). Although rear-sniffing was common across all groups during the first several minutes, this behavior decreased among cagemates (CAG1 and CAG2, black and gray traces) but continued at high levels in strangers (STR, green trace). Inset shows cumulative interaction time across all interactive behaviors. **E)** Cross correlogram of levels of allogrooming/body-sniffing (traces) relative to rear-sniffing (see Figure 2E). Similar to SEP in Experiment 2, the STR group showed a lower incidence of allogrooming immediately before rear-sniffing compared with 24 hr isolated cagemates. **F)** Vocalization rates across groups. More vocalizations were observed in STR. G) Cumulative distributions of vocalization time across groups. After a quiet first few minutes, vocalizations were common in STR throughout the session. **H)** Vocalization rates across all 15, manually-labeled syllable categories examined in the present study. Higher vocalizing in strangers was observed across many types, although this was not as clear for chitters (the most common type of syllable observed between cagemates). **I)** Heatmap of identified syllable clusters with the strongest differences between STR and CAG groups (see Figures 1G and 2G). In contrast with prior experiments, the STR versus CAG difference showed multiple independent clusters with differential, proportional activity between the groups, as illustrated by the hot colors observed in each of the 3 top rows. **J)** t-SNE projection of vocalization features (see Figures 1H and 2H). Syllables from each of the three significant clusters are plotted (blue, green, and turquoise). The first two clusters are made-up of particularly long-duration syllables that cluster in a small island; these are most likely combinations of multiple overlapping syllables, notably far more common in stranger dyads than cagemates. **K)** Proportion of syllables manually labeled across different category types (as in Figures 1I, 2I, and 3I). Labeling of the long-duration clusters (blue and green) was variable, most likely because they comprised multiple overlapping syllables between the two individuals. The turquoise trace represented a group of syllables that was proportionally (relative to all syllables) higher in cagemates. These were typically labeled as chitters.

Stranger dyads interacted more than did cagemates (two-sample t-test, n = 8 stranger dyads, 17 pooled cagemate dyads, t_{23} = 2.29, p = 0.032; **Figure 4B**). The effect was largely driven by higher levels of rear-sniffing (anogenital investigation) between strangers (two-sample t-test, t_{23} = 3.55, p = 0.0017, Bonferroni corrected alpha = 0.05/4 = 0.0125; **Figure 4C**). Over the first several minutes of the session interaction levels between STR were on par with those of CAG1 and CAG2; however, while those of cagemates began to decrease and level off, those between strangers continued to remain high. This was particularly evident for rear-sniffing (**Figure 4D**).

Temporal relationships were also different between STR and CAG groups. Paralleling the effects of separation relative to isolation (Experiment 2), strangers were less likely to engage in putative allogrooming immediately before rear sniffing (two-sample t-test, $t_{23} = 2.02$, $p = 0.0075$, **Figure 4E**). This result is consistent with the idea that the allogrooming/body-sniffing taking place before anogenital investigation represents an affiliative gesture between animals that had been previously isolated.

Vocalization rates were very high between strangers, exceeding those between cagemates following 24 hours of isolation (two-sample t-test pooling cagemate groups, $t_{23} = 3.08$, $p = 0.0054$, **Figure 4F**). Higher levels of vocalizing between strangers was not uniform across syllable types (2-way ANOVA, effect of condition $F_{2,330} = 11.5$, $p = 1.5 \times 10^{-5}$, effect of type: $F_{14,330} = 5.33$, $p = 3.6 \times 10^{-9}$, interaction between condition and type: $F_{28,330} = 1.95$, $p = 3.5 \times 10^{-3}$). Among manually-scored vocalizations, increases were particularly pronounced for chaffs (two-sample t-test, pooling cagemate groups, $t_{23} = 2.59$, $p = 0.016$), chirps ($t_{23} = 2.61$, $p = 0.016$), grunts ($t_{23} = 2.88$, $p = 0.0085$), loud whistles ($t_{23} = 2.38$, $p = 0.026$), pips ($t_{23} = 2.21$, $p = 0.037$), squeals ($t_{23} = 2.88$, $p = 0.0084$), tweets ($t_{23} = 2.58$, $p = 0.017$), the mixed category ($t_{23} = 3.08$, $p = 0.0054$), and warbles ($t_{23} = 2.63$, $p = 0.015$); given the multiple comparisons No evidence of difference was observed for chitters, whistles, and whines ($p > 0.25$). More generally, STR exhibited higher non-chitter to chitter ratios (paralleling SEP and ISO differences in Experiment 2: Wilcoxon rank sum test, $p = 0.035$). These results were consistent with results of the repeated-k-means method. Three clusters of syllables were identified as being differentially expressed between STR and CAG, but the first two were part of a small cluster of very-long duration (>1 s) syllables that were likely, in fact, combinations of several syllables, and might have combined vocalizations from both individuals (these were often labeled as chaffs, loud whistles, tweets, and whines). The final cluster was comprised of chitters, found to be differentially expressed (relative to all syllables) in CAG. This is consistent with the idea that chitters represented affiliative interactions more prevalent between familiar individuals.

One question that could be raised is whether the differences between strangers and cagemates could be due to kinship differences, as most cagemates were also siblings, while strangers were

not born from the same breeding pairs. Since the experiment included only three, non-related

cagemate dyads, this could not be thoroughly tested. It is worth noting that non-sibling cagemates,

as with siblings, showed lower averages than strangers; however, the effect size for this difference

was lower for non-siblings (non-siblings, n = 3, Cohen's D = 0.51; siblings n = 17, Cohen's D =

0.94).

In summary, data from Experiment IV confirmed high levels of interaction between strangers, and

confirmed that these interactions took a different form than cagemates—including higher rear-

sniffing and very high levels of non-chitter vocalizations. Also similar to SEP, strangers did not

exhibit high levels of allogrooming/body-sniffing before rear-sniffing observed in cagemates who

had been previously isolated for 24 hrs. Exploratory analyses of vocalizations generally confirmed

the high degree of non-chitter vocalizing among strangers. Finally, when strangers were compared

with non-sibling cagemates effect sizes were reduced suggesting that some differences may have

been due to kinship.

Discussion

The present study was designed to disentangle factors contributing to social motivation in degus

after isolation by evaluating details of reunion behavior across different manipulations. A general

summary of observations is illustrated in **Figure 5**. Broken-down by experiment, they include the

following: **Experiment 1** found that adult female degus become more motivated to interact

following longer periods of social isolation. This is a novel result. Although it conforms to

observations that have been made in other non-solitary species, including humans [1] and the

traditional laboratory model of young adult male rats [2], work in some species has

counterintuitively found reduced interactive behavior during reunion (e.g., lion tamarins: [44], and

in some cases, marmosets: [45]). Notably, in the present experiment most dyads received the 24

hr condition last, creating a potential order confound. However, the difference between 1 min

versus 24 hr was reliable (replicated in Experiment 2) and unsurprising; moreover, in other

experiments we have found no evidence for increased interaction with repeated reunions

(unpublished data). **Experiment 2** found that separating degus of a dyad without isolating them

from other conspecifics also elicited high interaction levels. Multiple differences were observed

between reunions following isolation and separation: after isolation, interactions were distributed

earlier in the sessions, included more frequent allogrooming-to-rear-sniffing transitions, and
included a higher proportion of chitter vocalizations. Direct comparison of the effects of isolation
versus separation-without-isolation have rarely if ever been made between non-human adult,
same-sex peers, although results are consistent with other lines of research, including studies on
greetings and research opposite-sex, pair-bonded individuals, discussed below. The observed
patterns have implications for understanding behaviors dyads may use to relieve isolation stress
compared with those they may use to renew previous relationships. As with Experiment 1,
conditions proceeded in a particular sequence, though comparisons with the Experiment 1, 1 min
(first condition) and 24 hr (first reunion following an extended separation) reinforced results.
Finally, **Experiments 3 and 4** provided further evidence that "stress" and "relationship
establishment" are useful constructs for interpreting the behaviors: exposure to a footshock stressor
(Experiment 3), like isolation, elicited more early-session reunion interactions, and interactions
between strangers contained low levels of pre- rear-sniff allogrooming and relatively higher levels
of non-chitter vocalizations. Differences between footshock and isolation suggest stress was not
the only factor mediating motivation after isolation; e.g., the apparent nuzzling before rear-
sniffing, first identified in Experiment 1 (24 hr vs 1 min) but replicated in Experiment 2 (24 hr ISO
vs 1 min CTL), was highest between cagemates after 24 hr isolation, lowest in both strangers and
dyads separated without isolation, and apparently at intermediate levels in 45 min isolated animals
independent of the stress manipulation performed. The importance of these patterns for
understanding relationship renewal as compared with conspecific-general motivations like stress
buffering are discussed below.

Figure 5. Overview of observed differences between conditions. **A)** Amount of interaction over time (stylized illustration based on data). Both social isolation and footshock stress resulted in higher early-session interaction, while reunions with strangers and previously separated cagemates led to increased interaction over a more prolonged time period. **B)** Illustration of significant transitions between interaction types. An increase in allogrooming before rear-sniffing was primarily observed only in cagemates after isolation. **C)** General differences in vocalizing between groups. Strangers showed highest levels of vocalizations but these were primarily across non-chitters, while the chitter syllable rate in cagemates was very high. Non-isolated, previously separated cagemates showed slightly more resemblance to strangers.

Several factors associated with any study of natural behavior can impact results, including individual differences between animals, and biases or blind spots introduced by the ethograms used. Individuals likely differ in their social tendencies (i.e., "personalities"), and these differences may be further compounded for dyad combinations. Due to the relatively low sample sizes of the present experiments (8 or 9 dyads each), it is likely that effects that do not generalize across most dyads also failed to reach statistical significance, and therefore were not reported. The lower sample sizes may miss behaviors that vary between dyads, but can also be valuable for highlighting the "average behavioral patterns" common across most dyads. Another important factor that should be considered during interpretation is how the ethogram we used may have influenced our specific results. Selecting an ethogram for vocalizations was straightforward, as we were able to follow the extensive examinations of Long [34], and then complemented these with an automated clustering algorithm (our repeated-k-means approach). For physical interactions, behaviors were lumped into four types: agonistic (which included mounting as well as more aggressive behaviors like boxing, hindleg kicks, pushes, bites, etc), non-aggressive nose-to-body contact (allogrooming/body-sniffing), anogenital investigation (rear-sniffing), and face-to-face (which included both nose-nose and nose-mouth contact). It may be that visual inspection of face-to-body contact was insufficient for distinguishing between an affiliative behavior, allogrooming, known to play a role in consolation [46], and an information-seeking behavior, body sniffing. Ultimately, increasing sample sizes to reveal variable effects of the manipulations, and modifying our ethogram to tease-apart issues like grooming versus sniffing, would not have changed the overall conclusions that degus interact heavily with separated cagemates and these differ from stress-related interactions.

Several qualities about degus make this species uniquely suited for investigating reunion behaviors associated with peer relationships. Degus are highly social, potentially sensitive to isolation (previously studied in pups [24]), and there is converging evidence that females in particular are predisposed to make new relationships. Female degus show increased fitness with larger group sizes [47], but there is a high turnover of group membership over time [28] suggesting potential survival benefits to prosocial interactions with new individuals. Female degus are also known to share burrows with multiple, often genetically unrelated peers [25–27,30,31,33,48] (in some cases even cohabitating with unrelated species [48]). As a point of contrast, prairie voles reside with only familiar pair mates or family members [49] and tend to great strangers aggressively [50]. When the two species were directly compared using a partner preference test, prairie voles have been found to spend almost all of their time near or huddling with a familiar partner [51], while degus spend their time near and huddling with partners and strangers, exhibiting no average preference for either [29]. (Mice, meanwhile, tend not huddle with either partners or strangers, and spend much of their time in a non-social neutral chamber [51].) In the present study we confirm that degus interact heavily with strangers, but also show that similar drives may support cagemate interactions following a period of separation. These interactions were distributed across many minutes, suggesting they went beyond an ephemeral "greeting" and involved a prolonged period of re-establishing expectations. In the case of strangers, there was also a prolonged period of anogenital investigation, perhaps ensuring subsequent social recognition. One other domain that distinguished new and renewing relationships from conspecific-general motivation conditions was in vocal communication.

An intriguing feature of degus is their large repertoire of audible vocalizations [24], and the present study used both manually-scored and automatically-defined syllable categories to identify how vocalizations differed between conditions. Vocalizations were unexpectedly variable between sessions and dyads, far more than physical interactions. Also unexpectedly, while physical interactions began within 10 to 20 s of the reunion, vocal communication was not detected until later, within the first 1 to 2 min (e.g., **Figures 3F & 4G**). It seems unlikely that these patterns were only due to shifts from quieter, undetectable vocalizations, although the amplitude distributions of low-frequency syllables did suggest that some fell outside of the detection range (**Supplementary Figure 1B**). Instead, it may be that vocal communication is a "secondary" quality to their social

behavior, used only when an animal feels it's necessary to modulate another's behavior while patterning physical investigations and affiliative gestures. This would be consistent with the much higher levels of vocal communication observed in strangers, a situation in which individuals may feel more need to avoid misunderstandings. Consistent with this idea, when rats are devocalized by surgical removal of the larynx, their play behavior more readily escalates into fights [52]. The role of vocalizations for averting misunderstood intentions is much better studied in primates [53]. Grunting in baboons, for example, tends to be used in contexts that may help defuse tense situations, such as during potential conflict or when one female approaches a lower ranked female with an infant [54–57]. These patterns have been conceptualized in game theory terms, with repeated encounters requiring different kinds of information transmitted compared with single encounters [55]. In the present study, chaff-type vocalizations were found to be proportionally higher in relationship establishment/re-establishment conditions across three of the experiments. Long [34] reported chaffs as "[O]ften used when the subject is some distance away from a conspecific, but typically directed toward them. Used by males (7%) and females (93%) of all ages, usually during physical separation of observable conspecific (87.3%), but also when meeting an unfamiliar degu (4.5%)." They further specify the effect of chaffs as being approach, and the likely function as encouraging approach. The present work, which found chaffs to be more common in strangers than cagemates, suggests there more context about the state of an animal's relationship or recent experiences may be useful for understanding this call.

The term "social relationship" is often not used in rodent research, although has been defined and applied toward explaining animal behavior [58,59]. While the term is found only sparsely in the rodent literature, social relationships are studied in multiple forms, including dominance (e.g., mice: [60] rats: [61], marmots: [62], voles: [63], hamsters: [64,65], degus: [66]), pair bonds (e.g., voles: [67]), and same-sex peer affiliation (voles: [50]). Here we operationally define social relationships as the set of expectations, affiliative bonds, and intra-dyadic "roles" (e.g., dominance roles) that two individuals have for one-another. Research across a range of species has examined the importance of greetings for social bonds and dyadic expectations, although different species may show different emphases. As the temporal distance between two individuals increases, their uncertainty about one-another probably does as well, so greetings may be a means by which that learning can take place to mitigate uncertainty. In some cases this means a relief of tension from

potential agonistic or competitiveness interactions (e.g., male baboons [7], black and white colobus monkeys [68], and spider monkeys [5,69]) while in others this may mean a reaffirmation of affiliative and cooperative bonds (e.g., "high risk" greetings in spotted hyaenas [8] and male guinae baboons [6]). Greetings are less well studied in rodents, but there are a number of exceptions. In marmots, greetings seem to progress behaviorally in ways that arguably go beyond simple social recognition [70], and squirrel greetings vary considerably depending on dominance relationships [71]. Consistent with the present study, greetings between degus in the wild show strong differences depending on whether individuals are from the same or different group areas [48]. Where degu co-residents tended to show face-to-face and mounting behaviors, degus from different areas sometimes exhibiting fighting behaviors and chases. Although the present study does not explore the consequences of greeting behavior in previously-separated degus, it does demonstrate that degus are highly motivated to interact with previously separated individuals over a time period of tens of minutes, even in the absence of other stressors, including isolation. It seems reasonable to infer that the consequences of these interactions include re-establishment of expectations, bonds, and dyadic roles.

One goal of the present study was to examine how social motivation is expressed in adults of a species that does not exhibit pair bonds or same-sex partner preferences, but the results can be considered next to those from research in opposite-sex, pair-bonded animals. The effects of both separations and isolations between pair-bonded partners have been investigated in a number of pair-bonding species (e.g., titi monkeys [72], and both lion tamarins and marmosets [44]), but rarely have the two types of manipulations been directly compared. One exception is a study in Zebra finches, which show much more social interaction with a heterosexual partner after isolation than following separation only, particularly displaying more billing (touching of beaks) and allopreening (allogrooming) [73]. Similarly, marmosets show higher social proximity and potentially more grooming with a novel reproductive partner if they have been previously isolated compared with if they have been kept in social housing [74]. But many of the patterns observed between pair-bonded animals likely generalize to same-sex peers considered in the present work, including the multiple modalities which are used to maintain or re-establish bonds after time apart [75]. As already noted, vocalizations are a common characteristic of peer interactions in general [53], though it seems likely that duetting and courtship songs are more exclusive to either opposite-

sex or pair-bonded relationships. These few studies offer some hint at the differences and
similarities that may exist in separation versus isolation effects across species and types of
relationships, though much more work will be necessary to gain a more generalized understanding
the impact of lifespan, species, and social context.

Reunions between cagemates after both isolation and footshock resulted in more early-session
relative to late-session interactions. The similarities between the two conditions could, in principle,
be mediated by glucocorticoids resulting from hypothalamic-pituitary-adrenal (HPA) activation
[76]. Glucocorticoid responses associated with isolation have been reviewed thoroughly by
Hawley et al. [9] for primates, pigs, rats, hamsters, voles and mice (particularly well studied in
murine rodents [16–22,77,78]while those in response to footshock have primarily been studied in
mice (e.g., [79]) and rats (e.g., [40]). Responses following footshock may not be long-lasting, but
if the data in murines generalize to degus then we can expect that in the present experiment they
remained elevated across the 10 to 20 min interval between delivery of the footshocks and the start
of the reunion session. The increased social interaction may, in turn, have reduced or "buffered"
stress levels, a phenomenon that has been studied in a range of species [10–15], including even
cows and horses [80,81]).

While stress may account for similarities between reunions after isolation and footshock, isolation
was also distinct from footshock, including higher ratio of chitter vocalizations and more pre- rear-
sniff allogrooming. There are many possibilities for the differences. For example, vocalization
levels were highly variable following footshock stress, and it may be that degus may be less
inclined to make sound if there is a potential external threat in the environment. Another factor
may be differences in the levels, duration, or controllability of stress between the two conditions,
all of which can differentially impact physiology and behavior [76]. But perhaps the most
intriguing explanation for differences between social and non-social stressors is the possibility that
other mechanisms, including other signaling systems, are responsible for supporting social
motivation following social compared with non-social stress. One candidate is the nonapeptide
oxytocin, which has a long-standing reputation as a modulator of social information processing in
the brain [82–85]. Although the peptide is not traditionally thought of as a stress hormone, levels
of oxytocin have been found to increase not only during social stress, but also following physical

stressors, e.g., using restraint or other protocols [86–89] (though not cold exposure [86]). Oxytocin may, in turn, protect an animal against some of the negative behavioral and autonomic effects of social isolation [90]. Another class of signaling molecules that may contribute to social motivation following isolation are opioids [16,23,91] (although it has been argued these play a stronger role in primates [92]). Both oxytocin and endorphins have been linked to social touch, and the value of both molecules for buffering stress—and perhaps loneliness—may implicate these signaling systems in driving social behavior following isolation [23]. These signaling molecules likely interact with specific neurotransmitter systems to mediate behavioral changes. Dopamine neurons in the dorsal raphe nucleus exhibit synaptic changes following isolation and are necessary for increased social behavior following isolation [93]. Serotonin pathways likely also help mediate the motivational salience of social stimuli, as demonstrated not only in rodents, by the importance of serotonin release in the nucleus accumbens [94], but also in species as evolutionarily distant as octopuses [95]. Careful dissection of the contributions these different pathways and signaling systems make to social behavior will offer new insights into the multidimensionality of social motivation.

In summary, the present results provide new information on behaviors associated with relationship renewal and buffering of isolation stress in a highly social, gregarious rodent. These findings are a valuable lead into understanding how animals in general, and female degus in particular, establish and re-establish expectations about one-another after a period of time, as well as how they overcome the loneliness of isolation during reunion.

Acknowledgements

The authors would like to thank the animal care staff at University of Toronto and Parminder Mathaun for help with post-hoc behavioral scoring.

References

1. Cacioppo JT, Hawkley LC, Norman GJ, Berntson GG. Social isolation: Social isolation. Annals of the New York Academy of Sciences. 2011;1231: 17–22. doi:10.1111/j.1749-6632.2011.06028.x

2. Niesink RJ, van Ree JM. Short-term isolation increases social interactions of male rats: a parametric analysis. Physiol Behav. 1982;29: 819–825. doi:10.1016/0031-9384(82)90331-6

3. Lorenz TK. Ober die Bildung des Instinktbegriffes. : 12.

4. Hogan JA. Energy models of motivation: A reconsideration. Applied Animal Behaviour Science. 1997;53: 89–105. doi:10.1016/S0168-1591(96)01153-7

5. Dias PAD, Rodríguez Luna E, Canales Espinosa D. The functions of the "Greeting Ceremony" among male mantled howlers (Alouatta palliata) on Agaltepec Island, Mexico. American Journal of Primatology. 2008;70: 621–628. doi:10.1002/ajp.20535

6. Whitham JC, Maestripieri D. Primate Rituals: The Function of Greetings between Male Guinea Baboons. Ethology. 2003;109: 847–859. doi:10.1046/j.0179-1613.2003.00922.x

7. Smuts BB, Watanabe JM. Social relationships and ritualized greetings in adult male baboons (Papio cynocephalus anubis). International Journal of Primatology. 1990;11: 147–172. doi:10.1007/BF02192786

8. Smith JE, Powning KS, Dawes SE, Estrada JR, Hopper AL, Piotrowski SL, et al. Greetings promote cooperation and reinforce social bonds among spotted hyaenas. Animal Behaviour. 2011;81: 401–415. doi:10.1016/j.anbehav.2010.11.007

9. Hawkley LC, Cole SW, Capitanio JP, Norman GJ, Cacioppo JT. Effects of social isolation on glucocorticoid regulation in social mammals. Hormones and Behavior. 2012;62: 314–323. doi:10.1016/j.yhbeh.2012.05.011

10. Kikusui T, Winslow JT, Mori Y. Social buffering: relief from stress and anxiety. Phil Trans R Soc B. 2006;361: 2215–2228. doi:10.1098/rstb.2006.1941

11. Hennessy MB, Kaiser S, Sachser N. Social buffering of the stress response: diversity, mechanisms, and functions. Front Neuroendocrinol. 2009;30: 470–482. doi:10.1016/j.yfrne.2009.06.001

12. Kiyokawa Y, Hennessy MB. Comparative studies of social buffering: A consideration of approaches, terminology, and pitfalls. Neurosci Biobehav Rev. 2018;86: 131–141. doi:10.1016/j.neubiorev.2017.12.005

13.   Lieberwirth C, Wang Z. The neurobiology of pair bond formation, bond disruption, and social
buffering. Curr Opin Neurobiol. 2016;40: 8–13. doi:10.1016/j.conb.2016.05.006

14.   DeVries AC, Glasper ER, Detillion CE. Social modulation of stress responses. Physiol Behav.
2003;79: 399–407. doi:10.1016/s0031-9384(03)00152-5

15.   Hostinar CE, Sullivan RM, Gunnar MR. Psychobiological mechanisms underlying the social buffering
of the hypothalamic-pituitary-adrenocortical axis: a review of animal models and human studies
across development. Psychol Bull. 2014;140: 256–282. doi:10.1037/a0032671

16.   Panksepp J, Herman B, Conner R, Bishop P, Scott JP. The biology of social attachments: opiates
alleviate separation distress. Biol Psychiatry. 1978;13: 607–618.

17.   Ieraci A, Mallei A, Popoli M. Social Isolation Stress Induces Anxious-Depressive-Like Behavior and
Alterations of Neuroplasticity-Related Genes in Adult Male Mice. Neural Plasticity. 2016;2016: 1–
13. doi:10.1155/2016/6212983

18.   Berry A, Bellisario V, Capoccia S, Tirassa P, Calza A, Alleva E, et al. Social deprivation stress is a
triggering factor for the emergence of anxiety- and depression-like behaviours and leads to
reduced brain BDNF levels in C57BL/6J mice. Psychoneuroendocrinology. 2012;37: 762–772.
doi:10.1016/j.psyneuen.2011.09.007

19.   Djordjevic J, Djordjevic A, Adzic M, Radojcic MB. Effects of Chronic Social Isolation on Wistar Rat
Behavior and Brain Plasticity Markers. Neuropsychobiology. 2012;66: 112–119.
doi:10.1159/000338605

20.   Pibiri F, Nelson M, Guidotti A, Costa E, Pinna G. Decreased corticolimbic allopregnanolone
expression during social isolation enhances contextual fear: A model relevant for posttraumatic
stress disorder. Proc Natl Acad Sci USA. 2008;105: 5567–5572. doi:10.1073/pnas.0801853105

21.   Wallace DL, Han M-H, Graham DL, Green TA, Vialou V, Iñiguez SD, et al. CREB regulation of nucleus
accumbens excitability mediates social isolation-induced behavioral deficits. Nat Neurosci.
2009;12: 200–209. doi:10.1038/nn.2257

22.   Thorsell A, Slawecki CJ, El Khoury A, Mathe AA, Ehlers CL. The effects of social isolation on
neuropeptide Y levels, exploratory and anxiety-related behaviors in rats. Pharmacol Biochem
Behav. 2006;83: 28–34. doi:10.1016/j.pbb.2005.12.005

23.   Morrison I. Keep Calm and Cuddle on: Social Touch as a Stress Buffer. Adaptive Human Behavior
and Physiology. 2016;2: 344–362. doi:10.1007/s40750-016-0052-x

24.   Colonnello V, Iacobucci P, Fuchs T, Newberry RC, Panksepp J. Octodon degus. A useful animal
model for social-affective neuroscience research: basic description of separation distress, social
attachments and play. Neurosci Biobehav Rev. 2011;35: 1854–1863.
doi:10.1016/j.neubiorev.2011.03.014

25.   Davis GT, Vásquez RA, Poulin E, Oda E, Bazán-León EA, Ebensperger LA, et al. *Octodon degus* kin
and social structure. JMAMMAL. 2016;97: 361–372. doi:10.1093/jmammal/gyv182

26.   Ebensperger LuisA, Hurtado M, Soto-Gamboa M, Lacey EileenA, Chang AnnT. Communal nesting
and kinship in degus (Octodon degus). Naturwissenschaften. 2004;91. doi:10.1007/s00114-004-
0545-5

27.   Quirici V, Faugeron S, Hayes LD, Ebensperger LA. Absence of kin structure in a population of the
group-living rodent Octodon degus. Behavioral Ecology. 2011;22: 248–254.
doi:10.1093/beheco/arq196

28.   Ebensperger LA, Chesh AS, Castro RA, Tolhuysen LO, Quirici V, Burger JR, et al. Instability Rules
Social Groups in the Communal Breeder Rodent *Octodon degus*. Ethology. 2009;115: 540–554.
doi:10.1111/j.1439-0310.2009.01635.x

29.   Insel N, Shambaugh KL, Beery AK. Female degus show high sociality but no preference for familiar
peers. Behav Processes. 2020;174: 104102. doi:10.1016/j.beproc.2020.104102

30.   Ebensperger L, Veloso C, Wallem P. Do female degus communally nest and nurse their pups?
Journal of Ethology. 2002;20: 143–146. doi:10.1007/s10164-002-0063-x

31.   Ebensperger LA, Sobrero R, Quirici V, Castro RA, Tolhuysen LO, Vargas F, et al. Ecological drivers of
group living in two populations of the communally rearing rodent, Octodon degus. Behav Ecol
Sociobiol (Print). 2012;66: 261–274. doi:10.1007/s00265-011-1274-3

32.   Ebensperger LA, Villegas Á, Abades S, Hayes LD. Mean ecological conditions modulate the effects
of group living and communal rearing on offspring production and survival. Behavioral Ecology.
2014;25: 862–870. doi:10.1093/beheco/aru061

33.   Hayes LD, Chesh AS, Castro RA, Tolhuysen LO, Burger JR, Bhattacharjee J, et al. Fitness
consequences of group living in the degu Octodon degus, a plural breeder rodent with communal
care. Animal Behaviour. 2009;78: 131–139. doi:10.1016/j.anbehav.2009.03.022

34.   Long CV. Vocalisations of the degu Octodon degus, a social caviomorph rodent. Bioacoustics.
2007;16: 223–244.

35.   Mahoney MM, Rossi BV, Hagenauer MH, Lee TM. Characterization of the Estrous Cycle in Octodon
degus. Biology of Reproduction. 2011;84: 664–671. doi:10.1095/biolreprod.110.087403

36.   Hummer DL, Jechura TJ, Mahoney MM, Lee TM. Gonadal hormone effects on entrained and free-
running circadian activity rhythms in the developing diurnal rodent Octodon degus. Am J Physiol
Regul Integr Comp Physiol. 2007;292: R586-597. doi:10.1152/ajpregu.00043.2006

37.   Lee TM. Octodon degus: a diurnal, social, and long-lived rodent. ILAR J. 2004;45: 14–24.
doi:10.1093/ilar.45.1.14

38.   Villavicencio CP, Márquez IN, Quispe R, Vásquez RA. Familiarity and phenotypic similarity influence
kin discrimination in the social rodent Octodon degus. Animal Behaviour. 2009;78: 377–384.
doi:10.1016/j.anbehav.2009.04.026

39.   Lidhar NK, Insel N, Dong JY, Takehara-Nishiuchi K. Observational fear learning in degus is
correlated with temporal vocalization patterns. Behav Brain Res. 2017;332: 362–371.
doi:10.1016/j.bbr.2017.06.011

40.   Rabasa C, Muñoz-Abellán C, Daviu N, Nadal R, Armario A. Repeated exposure to immobilization or
two different footshock intensities reveals differential adaptation of the hypothalamic–pituitary–
adrenal axis. Physiology & Behavior. 2011;103: 125–133. doi:10.1016/j.physbeh.2011.02.022

41.   Friard O, Gamba M. BORIS : a free, versatile open-source event-logging software for video/audio
coding and live observations. Fitzjohn R, editor. Methods Ecol Evol. 2016;7: 1325–1330.
doi:10.1111/2041-210X.12584

42.   Tchernichovski O, Nottebohm F, Ho CE, Pesaran B, Mitra PP. A procedure for an automated
measurement of song similarity. Animal Behaviour. 2000;59: 1167–1176.
doi:10.1006/anbe.1999.1416

43.   Coffey KR, Marx RG, Neumaier JF. DeepSqueak: a deep learning-based system for detection and
analysis of ultrasonic vocalizations. Neuropsychopharmacol. 2019;44: 859–868.
doi:10.1038/s41386-018-0303-6

44.   Shepherd RE, French JA. Comparative analysis of sociality in lion tamarins (Leontopithecus rosalia)
and marmosets (Callithrix kuhli): Responses to separation from long-term pairmates. Journal of
Comparative Psychology. 1999;113: 24–32. doi:10.1037/0735-7036.113.1.24

45.   Taylor JH, Mustoe AC, Hochfelder B, French JA. Reunion behavior after social separation is
associated with enhanced HPA recovery in young marmoset monkeys. Psychoneuroendocrinology.
2015;57: 93–101. doi:10.1016/j.psyneuen.2015.03.019

46.   Burkett JP, Andari E, Johnson ZV, Curry DC, de Waal FBM, Young LJ. Oxytocin-dependent
consolation behavior in rodents. Science. 2016;351: 375–378. doi:10.1126/science.aac4785

47.   Ebensperger LA, Correa LA, León C, Ramírez-Estrada J, Abades S, Villegas Á, et al. The modulating
role of group stability on fitness effects of group size is different in females and males of a
communally rearing rodent. Dantzer B, editor. J Anim Ecol. 2016;85: 1502–1515.
doi:10.1111/1365-2656.12566

48.   Fulk GW. Notes on the Activity, Reproduction, and Social Behavior of Octodon degus. Journal of
Mammalogy. 1976;57: 495–505. doi:10.2307/1379298

49.   Getz LL, McGuire B, Pizzuto T, Hofmann JE, Frase B. Social Organization of the Prairie Vole
(Microtus ochrogaster). Journal of Mammalogy. 1993;74: 44–58. doi:10.2307/1381904

50.   Lee NS, Goodwin NL, Freitas KE, Beery AK. Affiliation, Aggression, and Selectivity of Peer
Relationships in Meadow and Prairie Voles. Front Behav Neurosci. 2019;13: 52.
doi:10.3389/fnbeh.2019.00052

51.   Beery AK, Christensen JD, Lee NS, Blandino KL. Specificity in Sociality: Mice and Prairie Voles
Exhibit Different Patterns of Peer Affiliation. Front Behav Neurosci. 2018;12: 50.
doi:10.3389/fnbeh.2018.00050

52.   Burke CJ, Kisko TM, Pellis SM, Euston DR. Avoiding escalation from play to aggression in adult male
rats: The role of ultrasonic calls. Behav Processes. 2017;144: 72–81.
doi:10.1016/j.beproc.2017.09.014

53.   Cheney DL, Seyfarth RM. Flexible usage and social function in primate vocalizations. Proc Natl Acad
Sci U S A. 2018;115: 1974–1979. doi:10.1073/pnas.1717572115

54.   Silk JB, Cheney DL, Seyfarth RM. The form and function of post-conflict interactions between
female baboons. Animal Behaviour. 1996;52: 259–268. doi:10.1006/anbe.1996.0171

55.   Silk JB, Kaldor E, Boyd R. Cheap talk when interests conflict. Animal Behaviour. 2000;59: 423–432.
doi:10.1006/anbe.1999.1312

56.   Silk JB, Seyfarth RM, Cheney DL. Strategic Use of Affiliative Vocalizations by Wild Female Baboons.
PLoS One. 2016;11: e0163978. doi:10.1371/journal.pone.0163978

57.   Cheney DL, Seyfarth RM, Silk JB. The role of grunts in reconciling opponents and facilitating
interactions among adult female baboons. Animal Behaviour. 1995;50: 249–257.
doi:10.1006/anbe.1995.0237

58.   Hinde RA. Interactions, Relationships and Social Structure. Man. 1976;11: 1. doi:10.2307/2800384

59.   Barrett L, Henzi SP. Constraints on Relationship Formation among Female Primates. Behaviour.
2002;139: 263–289.

60.   Wang F, Kessels HW, Hu H. The mouse that roared: neural mechanisms of social hierarchy. Trends
Neurosci. 2014;37: 674–682. doi:10.1016/j.tins.2014.07.005

61.   Blanchard DC, Blanchard RJ. Behavioral correlates of chronic dominance-subordination
relationships of male rats in a seminatural situation. Neuroscience & Biobehavioral Reviews.
1990;14: 455–462. doi:10.1016/S0149-7634(05)80068-5

62.   Huang B, Wey TW, Blumstein DT. Correlates and Consequences of Dominance in a Social Rodent:
Marmot Dominance. Ethology. 2011;117: 573–585. doi:10.1111/j.1439-0310.2011.01909.x

63.   Lanctot RB, Best LB. COMPARISON OF METHODS FOR DETERMINING DOMINANCE RANK IN MALE
AND FEMALE PRAIRIE VOLES (*MICROTUS OCHROGASTER*). Journal of Mammalogy. 2000;81: 734–
745. doi:10.1644/1545-1542(2000)081<0734:COMFDD>2.3.CO;2

64.   Drickamer LC, Vandenbergh JG. Predictors of social dominance in the adult female golden hamster
(Mesocricetus auratus). Animal Behaviour. 1973;21: 564–570. doi:10.1016/S0003-3472(73)80017-
X

65.   Drickamer LC, Vandenbergh JG, Colby DR. Predictors of dominance in the male golden hamster
(Mesocricetus auratus). Animal Behaviour. 1973;21: 557–563. doi:10.1016/S0003-3472(73)80016-
8

66. Correa LA, Frugone MJ, Soto-Gamboa M. Social dominance and behavioral consequences of
intrauterine position in female groups of the social rodent Octodon degus. Physiology & Behavior.
2013;119: 161–167. doi:10.1016/j.physbeh.2013.06.002

67. Young LJ, Wang Z. The neurobiology of pair bonding. Nat Neurosci. 2004;7: 1048–1054.
doi:10.1038/nn1327

68. Kutsukake N, Suetsugu N, Hasegawa T. Pattern, Distribution, and Function of Greeting Behavior
Among Black-and-White Colobus. International Journal of Primatology. 2006;27: 1271–1291.
doi:10.1007/s10764-006-9072-x

69. Aureli F, Schaffner CM. Aggression and conflict management at fusion in spider monkeys. Biol Lett.
2007;3: 147–149. doi:10.1098/rsbl.2007.0041

70. Barash DP. The social behaviour of the hoary marmot (Marmota caligata). Animal Behaviour.
1974;22: 256–261. doi:10.1016/S0003-3472(74)80077-1

71. Fisler GF. Agonistic Signals and Hierarchy Changes of Antelope Squirrels. Journal of Mammalogy.
1976;57: 94–102. doi:10.2307/1379514

72. Fernandez-Duque E, Mason WA, Mendoza SP. Effects of duration of separation on responses to
mates and strangers in the monogamous titi monkey (Callicebus moloch). Am J Primatol. 1997;43:
225–237. doi:10.1002/(SICI)1098-2345(1997)43:3<225::AID-AJP3>3.0.CO;2-Z

73. Remage-Healey L, Adkins-Regan E, Romero LM. Behavioral and adrenocortical responses to mate
separation and reunion in the zebra finch. Hormones and Behavior. 2003;43: 108–114.
doi:10.1016/S0018-506X(02)00012-0

74. Smith AS, Birnie AK, French JA. Social isolation affects partner-directed social behavior and cortisol
during pair formation in marmosets, Callithrix geoffroyi. Physiology & Behavior. 2011;104: 955–
961. doi:10.1016/j.physbeh.2011.06.014

75. Singletary B, Tecot S. Multimodal pair-bond maintenance: A review of signaling across modalities
in pair-bonded nonhuman primates. Am J Primatol. 2020;82. doi:10.1002/ajp.23105

76. Sapolsky RM. Why zebras don't get ulcers / Robert M. Sapolsky. 3rd ed. New York: Times Books;
2004.

77. Gamallo A, Villanua A, Trancho G, Fraile A. Stress adaptation and adrenal activity in isolated and
crowded rats. Physiology & Behavior. 1986;36: 217–221. doi:10.1016/0031-9384(86)90006-5

78. Hofer MA, Shair HN. Isolation distress in two-week-old rats: influence of home cage, social
companions, and prior experience with littermates. Dev Psychobiol. 1987;20: 465–476.
doi:10.1002/dev.420200410

79. Shanks N, Griffiths J, Zalcman S, Zacharko RM, Anisman H. Mouse strain differences in plasma
corticosterone following uncontrollable footshock. Pharmacology Biochemistry and Behavior.
1990;36: 515–519. doi:10.1016/0091-3057(90)90249-H

80. Coe CL, Franklin D, Smith ER, Levine S. Hormonal responses accompanying fear and agitation in the
squirrel monkey. Physiology & Behavior. 1982;29: 1051–1057. doi:10.1016/0031-9384(82)90297-9

81. Boissy A, Le Neindre P. Behavioral, cardiac and cortisol responses to brief peer separation and
reunion in cattle. Physiol Behav. 1997;61: 693–699. doi:10.1016/s0031-9384(96)00521-5

82. Goodson JL, Thompson RR. Nonapeptide mechanisms of social cognition, behavior and species-
specific social systems. Curr Opin Neurobiol. 2010;20: 784–794. doi:10.1016/j.conb.2010.08.020

83. Carter CS, Grippo AJ, Pournajafi-Nazarloo H, Ruscio MG, Porges SW. Oxytocin, vasopressin and
sociality. Prog Brain Res. 2008;170: 331–336. doi:10.1016/S0079-6123(08)00427-5

84. Carter CS. Oxytocin Pathways and the Evolution of Human Behavior. Annual Review of Psychology.
2014;65: 17–39. doi:10.1146/annurev-psych-010213-115110

85. Donaldson ZR, Young LJ. Oxytocin, vasopressin, and the neurogenetics of sociality. Science.
2008;322: 900–904. doi:10.1126/science.1158668

86. Gibbs DM. Dissociation of oxytocin, vasopressin and corticotropin secretion during different types
of stress. Life Sciences. 1984;35: 487–491. doi:10.1016/0024-3205(84)90241-8

87. Nishioka T, Anselmo-Franci JA, Li P, Callahan MF, Morris M. Stress increases oxytocin release
within the hypothalamic paraventricular nucleus. Brain Research. 1998;781: 57–61.
doi:10.1016/S0006-8993(97)01159-1

88. Taylor SE, Gonzaga GC, Klein LC, Hu P, Greendale GA, Seeman TE. Relation of Oxytocin to
Psychological Stress Responses and Hypothalamic-Pituitary-Adrenocortical Axis Activity in Older
Women: Psychosomatic Medicine. 2006;68: 238–245. doi:10.1097/01.psy.0000203242.95990.74

89. Grippo AJ, Gerena D, Huang J, Kumar N, Shah M, Ughreja R, et al. Social isolation induces
behavioral and neuroendocrine disturbances relevant to depression in female and male prairie
voles. Psychoneuroendocrinology. 2007;32: 966–980. doi:10.1016/j.psyneuen.2007.07.004

90. Grippo AJ, Trahanas DM, Zimmerman RR, Porges SW, Carter CS. Oxytocin protects against negative
behavioral and autonomic consequences of long-term social isolation. Psychoneuroendocrinology.
2009;34: 1542–1553. doi:10.1016/j.psyneuen.2009.05.017

91. Panksepp J, Herman BH, Vilberg T, Bishop P, DeEskinazi FG. Endogenous opioids and social
behavior. Neuroscience & Biobehavioral Reviews. 1980;4: 473–487. doi:10.1016/0149-
7634(80)90036-6

92. Machin AJ, Dunbar RIM. The brain opioid theory of social attachment: a review of the evidence.
Behav. 2011;148: 985–1025. doi:10.1163/000579511X596624

93. Matthews GA, Nieh EH, Vander Weele CM, Halbert SA, Pradhan RV, Yosafat AS, et al. Dorsal Raphe
Dopamine Neurons Represent the Experience of Social Isolation. Cell. 2016;164: 617–631.
doi:10.1016/j.cell.2015.12.040

94.   Dölen G, Darvishzadeh A, Huang KW, Malenka RC. Social reward requires coordinated activity of
1269   nucleus accumbens oxytocin and serotonin. Nature. 2013;501: 179–184. doi:10.1038/nature12518

1270   95.   Edsinger E, Dölen G. A Conserved Role for Serotonergic Neurotransmission in Mediating Social
1271   Behavior in Octopus. Current Biology. 2018;28: 3136-3142.e4. doi:10.1016/j.cub.2018.07.061

1272

1273   **Supplementary Materials**

1274

1275

1276

Supplementary Figure 1. Distribution of maximum amplitude and dominant pitch across all syllables. A)
Heatmap of 3D histogram of syllable amplitudes versus dominant pitch. Syllables fell into multiple
clusters, the cluster of particularly low amplitudes and pitch were almost exclusively chitters. B)
Histogram of clusters that did not have identifiable amplitude peaks above 400 Hz (red trace) and both
low pitch (blue trace) and high-pitch (black trace) syllables that did. Black dotted line shows
approximates the amplitude threshold for detecting syllables, although this would be more true for the
< 400 Hz syllables that fell within the ambient noise. The distribution of these syllables was asymmetric,
suggesting that some syllables fell below detection threshold.

1285

1286

1287

1288

1289   **Supplementary Figure 2.** Temporal relationships between physical behavior types. Each plot (blue
1290   traces) is a cross correlogram of normalized proportion of time degus spent using a comparison behavior
1291   (columns) relative to a reference behavior (rows). Times before zero are times before the reference
1292   behavior starts and times after zero are after the reference behavior ends. Gray traces show the cross
1293   correlograms of a surrogate dataset in which times of the comparison behavior are randomly jittered.

1294

1295

1296

1297   **Supplementary Figure 3**. Mapping of labeled syllables to a two-dimensional (t-SNE) projection. Although
1298   different vocalization types were distributed differently over multi-dimensional space, there was also a
1299   degree of overlap, inspiring the use of automated clustering methods. Axes are in arbitrary units.

1300

1301

1302

1303 Supplementary Figure 4. Rates of different vocalization types in the ISO (red), SEP (teal), and CTL
1304 conditions (black) conditions. On average, the ratio of non-chitter to chitter vocalizations was higher in
1305 SEP.

1306

1307

1308

1309

1310

1311

Supplementary Figure 4. Temporal relationships between interactions in STR (green), CTL1 (black), and CTL2 (grey) conditions of Experiment 4, organized as in Supplementary Figure 2. The only significant difference was observed in the higher levels of allogrooming/body-sniffing that preceded rear sniffing in both CAG1 and CAG2 conditions. Though not statistically significant, figure also shows apparently higher grooming/body-sniffing before and after agonistic interactions in strangers; this may have been due to either bites to the body, or overly-vigorous (painful) grooming labeled as agonistic in one or a few stranger dyads.

1319

A

interaction (y-axis) vs. time (x-axis)

isolation -> stranger
no isolation → separated cagemate
isolation -> cagemate
stress -> cagemate
no isolation -> cagemate

B

Face-to-face

Allogrooming/ body sniffing

(Isolation -> cagemates)

Agonistic (incl. mounting)

rear sniff (anogenital investigation)

C

chitters   non-chitters

Isolation -> strangers   lower   higher

no isolation -> separated cagemates

isolation -> cagemates   higher   lower