Age Influences on the Molecular Presentation of Tumours

Constance H. Li[1,2,3,4,5,6], Syed Haider[7], Paul C. Boutros[2,3,4,5,6,8,9]

[1]Ontario Institute for Cancer Research, Toronto, Ontario, Canada
[2] Department of Medical Biophysics, University of Toronto; Toronto, Ontario, Canada
[3] Department of Human Genetics, University of California, Los Angeles, CA, USA
[4] Department of Urology, University of California, Los Angeles, CA, USA
[5] Jonsson Comprehensive Cancer Center, University of California, Los Angeles, CA, USA
[6] Institute for Precision Health, University of California, Los Angeles, CA, USA
[7]The Breast Cancer Now Toby Robins Research Centre, The Institute of Cancer Research; London, United Kingdom
[8]Department of Pharmacology & Toxicology, University of Toronto; Toronto, Ontario, Canada
[9]Vector Institute for Artificial Intelligence, Toronto, Canada

Correspondence should be addressed to P.C.B. (PBoutros@mednet.ucla.edu)
12-109 CHS
10833 Le Conte Avenue
Los Angeles, CA, 90095
Phone: 310-794-7160

Abstract

Cancer is often called a disease of aging. There are numerous ways in which cancer epidemiology and behaviour change with the age of the patient. The molecular bases for these relationships remain largely underexplored. To characterize them, we analyzed age-associations in the nuclear and mitochondrial somatic mutational landscape of 20,033 tumours across 35 tumour-types. Age influences both the number of mutations in a tumour and their evolutionary timing. Specific mutational signatures are associated with age, reflecting differences in exogenous and endogenous oncogenic processes such as a greater influence of tobacco use in the tumours of younger patients, but more activity of DNA damage repair signatures in those of older patients. We find that known cancer driver genes are mutated in age-associated frequencies, and these alter the transcriptome and predict for clinical outcomes. These effects are most striking in brain cancers where alterations like *SUFU* loss and *ATRX* mutation are age-dependent prognostic biomarkers. Using three cancer datasets, we show that age shapes the somatic mutational landscape of cancer, with clinical implications.

Introduction

Cancer health disparities across different population stratifiers are common through a wide range of measures. These include differences in incidence rates, mortality rates, response to treatment, and survival between individuals of different sexes[1–6], races or ancestries[7–11] and ages[12–14], and these differences have been described across a range of tumour-types. Cancer disparities involving age are particularly well known. Aging is a leading risk factor for cancer, as it is associated with increased incidence of most tumour-types[9,15]. Older age is also typically associated with higher mortality and lower survival[16,17]. The links between older age and increased cancer burden are such that cancer is often described as a disease of aging[18,19].

However, there are many nuances in the relationship between aging and cancer. Pediatric cancers are an obvious exception, as cancers arising in children have different molecular and clinical characteristics[9,20–22]. Tumours arising in young adults (< 50 years of age) are often more aggressive: early onset tumours of the prostate[23], breast[24], pancreatic[25,26], colorectal[27], and soft tissue sarcomas[28] are diagnosed at higher stages and associated with lower survival. Molecular studies have described some striking differences in the mutational landscapes of early onset *vs.* later onset disease[29–31], suggesting differences in the underlying oncogenic processes driving cancer at different ages.

The mechanisms of how age shapes the clinical behaviour of cancers have been subject to intense study. Many factors and behaviours closely tied to aging have been implicated in observed epidemiological and clinical cancer health disparities. For example, higher age is associated with a greater burden of comorbidities such as diabetes and cardiovascular disease[32,33]. Higher prevalence of chronic disease, frailty and increased likelihood of adverse drug reactions also influence the choices of clinical interventions given to older cancer patients[34–36]. Nevertheless differences remain even after accounting for these factors[37]. Previous work associating somatic molecular changes with age suggest differences in overall tumour mutation burden[38], transcriptional profiles[39], and some mutational differences[29–31]. These studies have focused on single tumour-types, relatively small cohorts, or have only evaluated fractions of the whole-genome, leaving the landscape of age-associated cancer mutations largely unknown.

To fill this gap, we performed a pan-cancer, genome-wide study of age-associated molecular differences in 10,218 tumours of 23 tumour-types from The Cancer Genome Atlas (TCGA), 2,562 tumours of 30 tumour-types from the International Cancer Genome Consortium/The Cancer Genome Atlas Pan-cancer Analysis of Whole Genomes (PCAWG), and 7,259 tumours across 35 tumour-types from the AACR Genomics Evidence Neoplasia Information Exchange (GENIE) projects. We quantified age-associations in measures of mutation density, subclonal architecture, mutation timing,

74 mutational signatures and driver mutations in almost all tumour-types. These associations
75 remained even after adjusting for potential confounding factors such as sex and ancestry.
76 Many of these genomic age-associations were linked to clinical phenotypes: in particular,
77 we identified genomic alterations that were prognostic in specific age contexts,
78 suggesting the clinical utility of age-informed biomarkers.

Results

Age Associations in Mutation Density and Timing

We investigated TCGA, PCAWG and AACR GENIE datasets independently and performed pan-cancer analyses spanning all TCGA, all PCAWG, and all AACR GENIE tumours in separate analyses; these were supplemented with tumour-type-specific analyses. We used the recorded age at diagnosis (**Table 1**). We used a two-stage statistical approach: we first used univariate methods to identify putative age-associations, and then further modeled these putative hits with multivariate regression to evaluate age effects after adjusting for confounding factors. Our multivariate modeling accounted for a range of confounding variables for each tumour-type including sex and genetic ancestry. We modeled each genomic feature and tumour subtype based on available clinical data, *a priori* knowledge, variable collinearity and model convergence. Model and variable specifications, and results of association tests between model variables and age are presented in **Supplementary Table 1**. We perform multiple testing adjustment at both stages using the Benjamini-Hochberg false discovery rate procedure and these adjusted p-values are used throughout. Bonferroni-adjusted p-values provide similar support for our findings. Both Benjamini-Hochberg and Bonferroni, as well as unadjusted p-values are provided in supplementary materials.

The accumulation of mutations with age is a well-known phenomenon in both cancer and non-cancer cells[41–48]. To test the robustness of our statistical framework in detecting age-associated genomic events, we investigated age associations in two measures of mutation accumulation: single nucleotide variant (SNV) density and genome instability. Both SNV density and genome instability have clinical relevance as they are associated with poor outcome in some tumour-types[49–51] and response to immunotherapy in others[52,53]. We first identified univariate age-associations in SNV density using Spearman correlation. Putative age-associations identified at a false discovery rate (FDR) threshold of 10% were further analysed by multivariate linear regression (LNR) models to adjust for tumour-type-specific confounding effects (**Supplementary Table 1**). We have previously applied this statistical strategy successfully to identify sex-associated somatic mutational features[40,54].

As expected[41,42], we identified significant associations between age and the accumulation of SNVs across a range of tumour contexts (**Figure 1A**). There were pan-cancer positive correlations between age and SNV density in TCGA (pan-TCGA: $\rho = 0.31$, adjusted LNR $p = 4.1 \times 10^{-57}$) and PCAWG ($\rho = 0.43$, adjusted LNR $p = 1.6 \times 10^{-26}$) data. Using TCGA and PCAWG data, we estimate that SNV density increases at a rate of 0.077 mutations per megabase pair per year (**Table 2**, **Methods**). We also identified positive associations in 11 TCGA, 14 PCAWG, and six AACR GENIE tumour-types (**Figure 1A**). Of these, eight tumour-types showed consistent results in two of three datasets (**Supplementary Figure**

1, Supplementary Table 2) including prostate cancer (TCGA: ρ = 0.25, adjusted LNR p
= 0.015; PCAWG: ρ = 0.48, adjusted LNR p = 1.2×10^{-4}; **Figure 1B**). Estimates for per
year increase in mutation density are given in **Table 2** for tumour-types with consistent
evidence in at least two datasets.

We next asked whether there were differences in the timing of when these SNVs occurred
during tumour evolution and leveraged data describing the evolutionary history of
PCAWG tumours[55]. We first investigated polyclonality, or the number of cancer cell
populations detected in each tumour. Monoclonal tumours, or those where all tumour cells
are derived from one ancestral cell, are associated with better survival in several tumours
types[56–58]. While there were intriguing univariate associations between age and
polyclonality in non-Hodgkin lymphoma and prostate cancer, these were not significant
after multivariate adjustment (**Supplementary Figure 1, Supplementary Table 2**). We
then focused on polyclonal tumours and asked whether there were associations in
mutation timing: we investigated whether SNVs, indels or structural variants (SVs)
occurred more frequently as clonal mutations in the trunk or as subclonal ones in
branches.

We identified several significant associations between age and mutation timing. In pan-
PCAWG analysis, we found positive associations between age and proportion of clonal
SNVs (ρ = 0.20, adjusted LNR p = 1.4×10^{-3}, **Figure 1C**) and proportion of clonal indels (ρ
= 0.14, LNR p = 0.013, **Supplementary Table 2**). Age was also associated with
increasing clonal SNV proportion in two tumour-types: stomach cancer (Stomach-
AdenoCA: ρ =0.44, adjusted LNR p = 0.028), and medulloblastoma (CNS-Medullo: ρ =
0.34, adjusted LNR p = 2.5×10^{-3}, **Figure 1C**). Positive correlations in these tumour-types
indicate tumours arising in older individuals accumulated a greater fraction of SNVs
earlier in tumour evolution. In contrast, we identified the inverse trend in melanoma, where
tumours of younger patients accumulated more subclonal than clonal SNVs (ρ = -0.47,
adjusted LNR p = 7.8×10^{-3}). Differences in the proportion of clonal mutations suggest
differential mutation timing over the tumour evolution and could be caused by
mechanisms such as differences in mutational processes or driver mutation frequency.

We next focused on genome instability, a measure of copy number aberration (CNA)
burden and approximated by the percent of the genome altered by CNAs (PGA).
Analogous to SNV density measuring the burden of point mutations, PGA measures the
density of copy number alterations. We found that in pan-cancer analysis, PGA increased
with age in PCAWG (ρ = 0.19, adjusted LNR p = 0.022) and AACR GENIE (ρ = 0.041,
adjusted LNR p = 0.050) (**Figure 1D**) and estimate that PGA increases at 0.010% per
year (**Table 2**). We also identified positive correlations in six TCGA, three PCAWG, and
three AACR GENIE tumour-types. Again, prostate cancer showed consistent age-PGA
associations, this time in all three datasets (TCGA: ρ = 0.17, adjusted LNR p = 6.7×10^{-5};
PCAWG: ρ = 0.27, adjusted LNR p = 3.0×10^{-3}; AACR GENIE: ρ = 0.11, adjusted LNR p

= 0.050; **Figure 1E**). Age was associated with PGA in stomach cancer data in TCGA (ρ
= 0.11, adjusted LNR p = 0.011) and AACR GENIE (ρ = 0.38, adjusted LNR p = 0.041),
and while other age-PGA correlations were not statistically significant across multiple
datasets, they showed similar effect sizes (**Supplementary Figure 1**). Intriguingly, we
detected negative age-PGA associations in TCGA lung adenocarcinomas (**Figure 1D**),
and correspondingly negative associations in PCAWG (ρ = -0.13) and AACR GENIE lung
tumours (ρ = -0.099) (**Supplementary Figure 1**). Estimates for per year increase in PGA
for these tumour-types are given in **Table 2** for tumour-types with consistent evidence in
at least two datasets.

To better understand why lung cancers of younger patients might have higher genomic
instability, we investigated tobacco use. Tobacco exposure is a well-known risk factor in
lung cancers. Our multivariate lung cancer regression models do account for tobacco use,
but we supplemented our analysis with a more focused study of the relationship between
PGA, age and tobacco history. Age was associated with tobacco history (Kruskal-Wallis
p = 1.6×10^{-12}, **Figure 1F, top**), where current smokers were diagnosed with lung cancer
at younger ages than never and reformed smokers (difference in location = -6.0, 95% CI
= -8.0 − -4.0, Wilcoxon rank sum test p = 1.2×10^{-7}). PGA was also associated with
smoking history (Kruskal-Wallis p = 1.6×10^{-12}, **Figure 1F, bottom**) where tumours arising
in current smokers had higher PGA (difference in location = 5.0, 95% CI = 1.2 − 9.2,
Wilcoxon rank sum test p = 1.0×10^{-3}). We then examined the associations between age
and PGA within each tobacco use category and found no association in never and
reformed smokers ≥ 15 years (Spearman correlation p ≥ 0.1, **Figure 1G**). In contrast,
tumours of current and reformed smokers < 15 showed statistically significant negative
correlations, indicating the negative association between age and PGA are dependent on
current or recent tobacco use. These mutation density results demonstrate nuances in
the relationship between aging and the accumulation of mutations: there are differences
by when mutations occur throughout tumour evolution, and by how they occur through
different mutational processes.

Age-Associated Mutational Signatures
We continued exploring mutational processes through mutational signatures data.
Clinical data on exposures such as tobacco use history and frequency are self-reported
by the patient and may be inaccurate, mis-recorded, or not described. Moreover, clinical
data often do not include information on exposures such as second-hand smoke, which
is also a known cancer risk factor[59]. We leveraged the COSMIC mutational signatures,
which can be applied to deconvolve distinctive mutational patterns from genomic
sequencing[60]. Each mutational signature is thought to represent a specific oncogenic
process – for instance, single base signature (SBS) 4 represents tobacco exposure.
Using mutational signatures, we can not only detect the influence of oncogenic processes
within a tumour, but we can also quantify the number of mutations attributed to that

process and assess its relative activity compared with the activity of other mutational
processes active within the tumour.

Focusing on the tobacco smoking signature SBS4, we stratified TCGA lung
adenocarcinoma tumours into SBS4-postive (SBS4+) and -negative (SBS4-) groups.
SBS4+ tumours were detected more frequently in younger patients (Wilcox p = 1.1×10^{-5})
and have higher PGA (Wilcox p =1.0×10^{-3}, **Figure 2A left**). We also found that the number
of mutations attributed to SBS4 was negatively correlated with age (Spearman's ρ = -
0.12, p-value = 0.086) and positively correlated with PGA (Spearman's ρ = 0.31, p-value
= 2.9×10^{-6}, **Figure 2A centre**). These findings show that a greater burden of tobacco-
associated mutations occurs in lung tumours of younger patients and is associated with
increased genomic instability. When considering SBS4-attributed SNVs as a fraction of
total detected SNVs, we found that while age remained negatively correlated with SBS4
relative activity (Spearman's ρ = -0.16, p-value = 0.017), there was no association with
PGA (Spearman's ρ = 0.082, p-value = 0.23, **Figure 2A right**): while smoking-attributed
mutations account for a greater percentage of mutations in younger patients, this relative
activity is not associated with a change genomic instability. Also, when examining the
relationships between age and PGA separately in SBS4+ and SBS4- tumours, we found
significant negative correlations in both groups (**Figure 2B**), suggesting additional factors
beyond SBS4-attributed mutations may contribute to increased PGA in lung tumours
diagnosed in younger patients.

We repeated this analysis in TCGA lung squamous cell cancers and detected a similar
negative correlation between age and PGA in SBS4- tumours (Spearman's ρ = -0.17, p-
value = 5.8×10^{-3}, but no association in SBS+ tumours (Spearman's ρ = -0.0069, p-value
= 0.94, **Supplementary Figure 2**). SBS4 activity was also negatively associated with age
in PCAWG lung adenocarcinomas (Lung-AdenoCA: ρ = -0.50, adjusted LNR p = 0.025,
Supplementary Figure 2). Indeed, SBS4 and age were consistently negatively
associated across both subtypes of lung cancer and both datasets, though not all
associations were statistically significant after multiple testing adjustment. This supports
previous findings that tobacco has a larger tumorigenesis role in younger patients, with
tobacco-associated mutations contributing to a greater portion of the mutational
landscape of tumours derived from younger individuals[61].

The PCAWG project updated the COSMIC signatures v3 to a set of 49 single base
substitution , 11 doublet base substitution (DBS) and 17 small insertion and deletion (ID)
signatures[62]. We extended our analysis of SBS4 to all 77 mutational signatures in
PCAWG data and to SBS signatures in TCGA data. We did not investigate mutational
signatures in AACR GENIE data due to smaller mutation numbers resulting from the
limited genome coverage of the MSK-IMPACT panel. Similar to our analysis of SBS4, we
examined both the proportion of signature-positive tumours as well as relative mutation
activity. Previous studies of mutational signatures describe the correlations between age

and signature-attributed mutations but ignore the other aspects of signature detection and
relative activity. By comparing signature detection rates, we identify mutational processes
that are more likely to be active in younger *vs* older patients and vice versa. By analyzing
signature-attributed mutations as a proportion of total mutations per tumour, we can
derive information about that signature's contribution to the overall mutational spectrum.
For example, SBS1 is well-known as being 'clock-like' and its number of attributed
mutations increase with age[60,62]. However, because SBS1 is detected almost universally,
it is equally likely to occur in tumours of younger *vs* older patients; when analysed as a
proportion of total mutations, we find that the proportion of SBS1 mutations does not
change with age, suggesting that its relative activity is stable with age (Spearman's
correlation p > 0.1).

Across all 2,562 PCAWG tumours, we identified twelve mutational signatures with age-
associated detection frequency (**Figure 2C, left**) and ten with age-associated relative
signature activity (**Figure 2D, left**). For example, tumours arising in older patients were
more likely to be SBS3-positive (marginal log odds change = 0.0085, 95%CI = 0.0024-
0.015, adjusted LGR p = 0.075), but in these SBS-positive tumours, the proportion of
SBS3-attributed mutations decreased with age (ρ = -0.20, adjusted LNR p = 3.2×10^{-3}).
SBS3 mutations are thought to be caused by defective homologous recombination-based
DNA damage repair. These results imply that while tumours derived from older individuals
are more likely to harbour defective DNA damage repair, its relative impact on the burden
of SNVs is lower compared with tumours derived of younger individuals. A similar
relationship was seen for ID8, which is linked to defective non-homologous DNA end-
joining (marginal log odds change = 0.024, 95%CI = 0.020-0.028, adjusted LGR p = 3.4
x 10^{-3}; ρ = -0.099, adjusted LNR p = 3.7 x 10^{-5}) and ID1, associated with slippage during
DNA replication (marginal log odds change = 0.013, 95%CI = 0.0059-0.020, adjusted
LGR p = 0.018; ρ = -0.059, adjusted LNR p = 0.048). Like our results for SBS4, we
identified associations between age and other tobacco-related signatures DBS2 and ID3.
Conversely, tumours arising in older individuals were less likely to exhibit defective base
excision repair (SBS36). All mutations signatures findings are in **Supplementary Table**
2.

These pan-cancer differences persisted across individual tumour-types. We identified 23
age-associated signatures across eleven tumour-types, including six significant
signatures in melanoma. In this tumour-type, tumours arising in older patients were
preferentially SBS2-positive (marginal log odds change = 0.051, 95%CI = 0.013-0.095,
adjusted LGR p = 0.029, **Figure 2C**), attributed to APOBEC cytidine deaminase activity[63].
Melanomas arising in younger patients were more likely to be positive for signatures
related to UV damage (SBS 7a, 7b, 7d, **Figure 2C, Supplementary Table 2**). The
proportion of mutations attributed to UV damage was also higher in younger patients
(DBS1, ρ = -0.29, adjusted LNR p = 0.019, **Figure 2D**), while the proportion of mutations
attributed to slippage during DNA replication was higher in older patients (ID1, ρ = 0.27,

adjusted LNR p = 0.019, **Figure 2D**). These results suggest that melanomas in younger
patients more frequently involve UV exposure and damage, while melanomas in older
patients are more influenced by endogenous sources of mutation.

Leveraging data describing SBS signatures in TCGA data, we repeated this analysis to
identify age-associations in signatures derived from whole exome sequencing (WXS)
data. Across pan-TCGA tumours, we detected five signatures that occurred more
frequently in older individuals, and three that occurred more frequently in younger
individuals (**Figure 2C**). We also identified five signatures with higher relative activity in
younger patients (**Figure 2D**).

There was moderate agreement between TCGA and PCAWG findings: while the results
in one dataset never contradicted those of the other, some signatures were associated
with age exclusively in either TCGA or PCAWG data. Other signatures, such as SBS1
and SBS5 were associated with age in detection and relative activity across a range of
tumour-types in either dataset. There was complete agreement in only some signature
like SBS2 and SBS4. We hypothesized that this was due to differences in signature
detection rates between WXS and whole genome sequencing (WGS) data and compared
how frequently each signature was detected across all samples (**Figure 2E**). Signatures
with high agreement between datasets had similar detection rates, as observed for SBS2
(detection difference = 1.5%) and SBS4 (detection difference = 1.1%). Signatures where
findings did not replicate had vastly different detection rates, as was seen for SBS1
(detection difference = 7.2%) and SBS5 (detection difference = 10%). We further
examined this by comparing signatures data from non-PCAWG WGS and non-TCGA
WXS data. Differences in signature detection rates between PCAWG and TCGA data
were reflected in non-PCAWG WGS and non-TCGA WXS data (**Supplementary Figure
2**). We also looked specifically at identified age-associations and found high agreement
in data generated by the same sequencing strategy (**Supplementary Figure 2**). These
findings suggest high confidence in age-associations detected in both WGS and WXS
data, and that additional study is needed in independent WXS and WGS data to validate
TCGA- and PCAWG-specific findings.

CNA Differences Associated with Transcriptomic Changes
Global mutation characteristics such as genome instability are features of later stages in
a tumour's evolutionary history. In contrast, the early stages are often driven by
chromosome- or gene-specific events such as loss of specific chromosomes or mutation
of driver genes[55]. We therefore narrowed our focus to 87 cancer driver genes frequently
affected by copy number alterations as described by COSMIC[64]. We applied our statistical
framework to identify putative age-associated copy number driver gains and losses using
univariate logistic regression. Putative events identified with a false discovery rate
threshold of 10% were further analysed by multivariable logistic regression to account for
confounding factors. We applied these analyses to PCAWG, TCGA, and AACR GENIE

datasets separately to characterize pan-cancer and tumour-type-specific associations.
We assessed our statistical approach with p-value Q-Q plots as presented in
Supplementary Figure 4.

In pan-cancer analysis of TCGA data, we identified 23 driver genes that were more
frequently lost and eight driver genes that were more frequently gained in in tumours of
older individuals (**Figure 3A**). Age-associated loss of *FANCA* (marginal log odds change
= 0.015, 95%CI = 0.012-0.018, adjusted MLR p = 3.2×10^{-9}) was also statistically
significant in AACR GENIE data (marginal log odds change = 0.051, 95%CI = 0.036-
0.066, adjusted MLR p = 0.011). Other age-associated loss and gain events were
statistically significant in one dataset and corroborated by similar effects in at least one
other. For example, loss of *STK11* was significant in PCAWG (marginal log odds change
= 0.030, 95%CI = 0.023-0.037, adjusted MLR p = 0.014), and was positively associated
with age in TCGA (marginal log odds change = 0.014) and AACR GENIE (marginal log
odds change = 0.044) datasets without reaching our statistical significance thresholds
(**Supplementary Table 3-4**).

There were also age-associated CNAs in specific tumour-types: we detected age-
associated gains with evidence in at least two datasets in eight tumour-types (**Figure 3B**),
and losses in ten tumour-types (**Figure 3C**). Most of these associations were positive,
indicating they were more likely to occur in tumours of older patients. We found the most
evidence of age-associated drivers gains in ovarian cancer across all three datasets, such
as gain of *ALK* (marginal log odds changes: TCGA = 0.016; PCAWG = 0.070; AACR
GENIE = 0.071) and MDM4 (marginal log odds changes: TCGA = 0.023; PCAWG =
0.056; AACR GENIE = 0.071; **Supplementary Table 4**). There were also eight age-
associated driver losses in this tumour-type. All identified CNAs were more frequently
detected in tumours of older patients except for *TNFAIP3* loss, which occurred more
frequently in tumours of younger patients (marginal log odds changes: TCGA = -0.028;
PCAWG = -0.071; **Supplementary Table 3**). Other CNAs more likely to occur in tumours
of younger patients were detected in colorectal, lung, breast, colorectal, liver,
glioblastoma, and melanoma tumour-types (**Figure 3B-C**, **Supplementary Tables 3-4**).

We next asked whether age-associated CNAs might lead to downstream transcriptomic
changes by investigating TCGA tumour-matched mRNA abundance data. We used linear
models with age, copy number status, and the interaction between age and copy number
status as predictors. These terms tell us when mRNA abundance differs by age, when
the CNA event itself is significantly associated with mRNA abundance, and when the
effect of the CNA event on mRNA depends on age. We also adjusted for tumour purity
(as estimated by study pathologists) in all mRNA analyses. Of 58 age-associated CNAs
with mRNA data, we found 17 were significantly associated with changes in mRNA
abundance (**Figure 3D left, Supplementary Table 5**). For example, loss of *CDKN2A* in
glioblastoma, more frequent in tumours of older individuals, was associated with

decreased mRNA abundance (adjusted mRNA-CNA p = 1.3x10^{-3}). Intriguingly, *CDKN2A*
loss in sarcoma was not significantly associated with a change in mRNA on its own but
did have a significant interaction with age (**Figure 3D right**). There was a larger decrease
in *CDKN2A* mRNA in sarcomas derived of older individuals than younger individuals
(adjusted mRNA-CNA-age p = 0.024, **Figure 3E**). Thus, these age-associated CNAs not
only preferentially occur in tumours derived of individuals of a certain age group, they are
also associated with changes in mRNA abundance.

To investigate potential clinical significance of these age-associated CNAs, we performed
survival analysis to identify prognostic events. We used Cox Proportional-Hazards (Cox
PH) models with 5-year overall survival as the end point. Like our mRNA models, we used
predictors including copy number status, age and their interaction. In glioblastoma, age
itself is a known prognostic feature with older patients having poorer outcome (HR = 2.1,
95%CI = 1.7-2.6, Wald p = 1.4x10^{-13}). We found that loss of *SUFU* is itself not prognostic,
but incorporating age revealed that younger individuals with no *SUFU* loss have the best
outcome (HR = 0.42, 95%CI = 0.30 - 2.3, adjusted Wald p = 5.5x10^{-6}). Also, *SUFU* loss
stratifies younger individuals into two groups with distinct trajectories, but has no such
prognostic value in glioblastomas of older individuals (**Figure 3F**).

We repeated these mRNA and survival analyses for all TCGA tumour-types with age-
associated CNAs. Genes in these CNAs were associated with altered mRNA abundance
in six tumour-types. Five tumour-types also showed that age-associated CNAs can be
prognostic and that the prognostic value can also differ based on the age of the individual.
We were unable to repeat these analyses in PCAWG data due the small number of
patients with mRNA abundance and outcome data but present all mRNA and survival
analysis results in **Supplementary Table 5**.

SNVs Differences Associated with Functional Changes

Finally, we investigated gene-level SNVs for age-associations. In PCAWG analysis, we
used a predefined set of genomic driver and mitochondrial genes[65]. In TCGA analysis,
we focused on a set of 679 COSMIC driver genes[64] and applied a recurrence threshold
to filter out genes mutated in less than 1% of tumours. We used AACR GENIE data
generated using MSK-IMPACT targeted sequencing of up to 468 cancer genes, and we
applied a recurrence threshold of 1%. We included SNV density in our multivariate models
in addition to other confounding factors as previously described.

In pan-cancer analysis, we identified 102 age-associated genes in TCGA, nine age-
associated genes in AACR GENIE, and one in PCAWG data (**Supplementary Table 6**).
CREBBP-frequency was associated with age in both TCGA (marginal log odds change =
0.030, 95%CI = 0.024 – 0.040, adjusted LGR p = 0.049) and PCAWG (marginal log odds
change = 0.027, 95%CI = 0.0089 – 0.047, adjusted LGR p = 8.7x10^{-3}, **Figure 4A,**
Supplementary Table 6). In AACR GENIE, the positive association between *CREBBP*-
status and age was not significant after multiple testing correction (marginal log odds

change = 0.011, 95%CI = 0.0047 – 0.022, adjusted p > 0.1). *KDM6A* and *RBM10* were
more likely to have SNV in tumours derived of older patients in TCGA and AACR GENIE
but were not recurrently mutated in PCAWG data and not analysed in that dataset (**Figure
4A, Supplementary Table 6**). 35 other genes found to be associated with age in TCGA
data showed similar effect sizes in either PACWG or AACR GENIE data, but without
reaching statistical significance.

There were also tumour-type specific age-associations in SNV frequency in TCGA,
PCAWG and AACR GENIE. We identified three tumour-types with consistent and
significant age-associated SNVs in at least two datasets, and five tumour-types with
associations that were significant in one dataset and showed the same effect in one other
(**Figure 4B, Supplementary Table 6**). SNVs in *FOXA1* occurred more frequently in
breast and prostate tumours derived of older individuals. *SPOP* was also positively
associated with age in prostate tumours (PCAWG adjusted LGR p = 0.099, AACR GENIE
adjusted LGR p = 0.03). In melanoma, SNVs in *NF1* were more frequent in tumours of
older individuals, while *BRAF* SNVs were more frequent in tumours of younger
individuals. We also confirmed known associations between age and mutations in tumour
suppressors *IDH1-* and *ATRX-* in both high grade glioblastoma and lower grade gliomas,
IHD1- and *ATRX-* were more frequent in tumours derived of younger individuals (**Figure
4C**, **Supplementary Table 6**). Other age-associated SNVs include positive associations
in cervical and head and neck cancer, and negative associations in colorectal cancer
(**Supplementary Table 6**).

Like the nuclear genome, the mitochondrial genome is frequently mutated in cancer[66].
We leveraged mitochondrial SNV (mtSNV) data from PCAWG whole genome sequencing
and identified age associated mtSNVs in pan-cancer analysis and in ovarian cancer
(**Figure 4D**, **Supplementary Table 6**). All significant age-associations were mtSNVs that
occurred more frequently in tumours of older patients, even after controlling for the
number of mitochondria copies in each tumour. Implicated mitochondrial regions include
MT-CYB, which encodes a cytochrome b and the D-loop, a non-coding region that
controls replication and transcription[67,68] (**Figure 4E**). We also examined whether the
number of tumour mitochondria copies was associated with age by investigating the
foldchange compared with normal mitochondria copies. There was indeed a significant
association between mitochondria copy increase and age in pan-cancer analysis and
three tumour-types (**Figure 4F**, **Supplementary Table 6**). In these tumour contexts,
tumours of older patients gained more mitochondria copies than in tumours of younger
patients.

As with the age-associated CNAs, we evaluated the impact of SNVs on mRNA
abundance and survival in TCGA data. We identified significant associations between
age-associated SNVs and mRNA abundance for *ATRX* and *IDH1* in lower grade glioma
(**Supplementary Table 5**). Mutations in *ATRX* and *IDH1* were associated with lowered

mRNA abundance in both genes. There was also a significant interaction between age
and *IDH1*- frequency (adjusted p = 2.1 x 10^{-4}, **Figure 4G**) indicating an age-dependent
effect on mRNA abundance: mutated *IDH1* was associated with a greater mRNA
decrease in tumours arising in younger patients. Interestingly, this difference results from
a change in baseline *IDH1* mRNA: older patients have higher *IDH1* mRNA abundance
than younger, and mutated *IDH1* leads to equalised mRNA levels. *IDH1* encodes
isocitrate dehydrogenase 1, a component of the citric acid cycle: differences in its
baseline abundance may be due to differences in metabolism in younger and older
brains[69]. In contrast while there was no interaction between age and mutation status on
mRNA abundance (adjusted p = 0.36, **Figure 4G**), *ATRX*- and age were synergistically
associated with outcome, stratifying lower grade glioma patients into four groups (**Figure
4H**).

The lower grade glioma tumour-type is comprised of astrocytoma and oligodendroglioma
subtypes. *IDH1* mutation is intrinsically linked to glioma subtype as oligodendroglioma is
diagnosed based on the presence of both the 1p/19q co-deletion and mutation of either
IDH1 or *IDH2*[70]. While our multivariable models adjust for tumour subtype, we
investigated age-associations of *ATRX* and *IDH1* SNV mutation frequency in lower grade
gliomas in greater detail. We stratified the TCGA lower grade tumours into astrocytoma,
oligodendroglioma and oligoastrocytoma subtypes and repeated the SNV, mRNA and
survival analyses in each group. We found *TP53* and *ATRX* SNV mutations occurred
more frequently in tumours derived of younger individuals across all three subtypes
(Wilcox **Supplementary Figure 5**). *ATRX* SNV mutation frequency was associated with
decreased mRNA abundance in all three subtypes, and we also found a significant age-
ATRX interaction in astrocytoma, where *ATRX*- was associated with a greater decreased
in mRNA abundance in tumours of older individuals than younger (interaction p = 0.016,
Supplementary Figure 5). *IDH1* SNV mutation frequency was negatively associated with
age in astrocytoma (Wilcox p = 1.3x10^{-8}) and oligoastrocytoma (Wilcox p = 0.041), but
not oligodendroglioma (Wilcox p = 0.19), and we found significant age-*IDH1* interactions
in mRNA abundance analysis for oligodendroglioma (interaction p = 0.038) and
oligoastrocytoma (interaction p = 6.2x10^{-3}). Finally, we found significant age-dependent
associations of *ATRX* SNV status in astrocytoma but not oligodendroglioma or
oligoastrocytoma: ATRX SNVs were associated with improved survival in older patients,
but with worse survival in younger patients.

Discussion

Despite modest statistical power, suboptimal study designs and limited clinical annotation, we identified myriad age-associated differences in cancer genomes. Age-associated genomic features occur at the pan-cancer level and also across almost all individual tumour-types. Combined with similar reports of sex- and ancestry-associated differences in cancer genomes[40,71], these data reveal a set of host influences on the mutational characteristics of tumours. Indeed, a study in simultaneous preparation by Chatsirisupachai *et al.* describes corroborating evidence of age-associated differences in the genome and transcriptome, as well as age-specific differences in methylation and gene expression control[72]. Together, we find that characteristics of the tumour host appears to influence all aspects of the tumour molecular profile and that some of these lead to age-specific transcriptomic and clinical impacts. We note that throughout this study we report Bonferroni-adjusted p-values, rather than FDR-adjustment which is standard in our field. This change was required in peer-review, and provides our results significantly more statistical stringency that than common in genomic studies.

The mechanisms for these genomic associations are largely unknown. Our data suggest some endogenous or exogenous mutational processes preferentially occur in individuals of different age groups. Some of these mutational processes are related to aging-associated phenomena such as declining DNA damage repair[73,74], somatic mosaicism and the accumulation of mutations over time[60,75,76]. However, other processes related to immune surveillance, evolutionary selection, disease aetiology and epigenetics are also likely involved[77–79]. Pathogenic germline variants such as those in *BRCA1/2* or *TP53* also lead to earlier presentation of cancer. While our results remain unchanged on removing tumours with detected known pathogenic variants[80], it is likely there remains hereditary confounders that we have not accounted for.

In addition to such biological factors, lifestyle and socioeconomic considerations like diet[81] and microbiome composition[82] can continuously shape tumour evolution from its earlier steps. Many of these factors are deeply linked to not only an individual's age, but other fundamental characteristics over which we have limited control, such as ancestry or sex. For example, we found that tobacco exposure is closely linked to the negative correlation between age and PGA. It is possible that tobacco exposure leads to earlier presentation of mutation-dense lung cancers. However, it is also likely that there are other variables and interactions that influence the relationship between age and mutation density. Moreover cohort effects, where individuals born in one time period experience different risk exposures from those born in another, can greatly influence the somatic profile of tumours. Our analyses do not consider such cohort effects, and some described age-associations may instead be attributed to differences across time. A tumour's mutational history therefore reflects a complex interplay of biological, lifestyle and healthcare factors,

and we have little understanding of how these diverse processes interact to produce
molecular phenotypes.

Most tumour-types in our study showed some association between age and genomic
alterations. Prostate cancer showed persistent associations in all three datasets across
multiple measures of genomic alterations. These were all positive age-associations where
mutation density, driver CNA frequency, and SNV frequency of *SPOP* and *FOXA1* all
increased with age. The strong association of age with alterations in prostate cancer may
be related to its typically slow-growth and low mutation burden. Moreover, known
exogenous risk factors, including diet and endocrine disrupting chemicals are thought to
converge on hormone regulation, which then acts on the prostate[83]. In contrast, tumours
types with negative age-associations such as lung and liver cancers have exogenous risk
factors that impact cells through multiple indirect and direct mechanisms such as viral
infection, mutagenesis and inflammation[84].

The TCGA, PCAWG and AACR GENIE datasets sometimes identified different molecular
associations, highlighting the differences between the three datasets. TCGA and AACR
GENIE patients were largely North American while PCAWG had a greater international
component. While the ages represented in all three datasets were similar (**Table 1**), the
cohorts differ in other host and clinical characteristics. For instance, the representation of
ancestry groups is dissimilar, with many tumour-types differing vastly in ancestry
proportions (**Supplementary Table 1**). Furthermore, differences in sequencing targets
also contributed to variation in our results, most conspicuously in the detection rates of
some mutational signatures. We customized our analyses to take advantage of the
contrasting strengths of each dataset: WGS in PCAWG allowed us to interrogate a
greater breadth of mutation types, while the larger sample size and clinical annotation of
TCGA data improved statistical power and controls for confounders. The focused panel
used to generate AACR GENIE data facilitated the validation of age-associations in
cancer genes. Also, while we were able to identify more age-associations in TCGA data,
many of these findings were reflected in PCAWG and AACR GENIE data by similar
effects that did not reach our statistical significance threshold. More sequencing data
reflecting greater and more balanced diversity is needed to distinguish those age-
associations that are intrinsic to differences in biology, and those that are tied to
differences in lifestyle and geography.

Our findings have wide-reaching implications for both basic and translational cancer
research. Since cancer host characteristics like age, ancestry and sex widely shape the
somatic cancer landscape, we cannot consider discovery genomics complete they are
explicitly considered. Elderly individuals are underrepresented in cancer sequencing
studies and clinical trials[39,85,86]: better inclusion is needed to identify somatic changes
specific to older individuals and to leverage these changes to improve clinical care. In our
analysis, we found that some age-associated genomic differences associate with

transcriptional and clinical changes, but many do not – identifying the functional
consequences and mechanisms of these will be a long-term challenge. Finally, these
epidemiological factors should be considered and controlled for in personalized therapy
strategies. Indeed, every type of analysis from driver-discovery to biomarker-
development should explicitly test for and model the powerful influence of patient biology
and behaviour on tumour evolution.

Online Methods

Data acquisition & Processing

Genome-wide somatic copy-number, somatic mutation, and mRNA abundance profiles for the Cancer Genome Atlas (TCGA) datasets were downloaded from Broad GDAC Firehose (https://gdac.broadinstitute.org/), release 2016-01-28. For mRNA abundance, Illumina HiSeq rnaseqv2 level 3 RSEM normalised profiles were used. Genes with >75% of tumours having zero reads were removed from the respective dataset. GISTIC v2 (13) level 4 data was used for somatic copy-number analysis. mRNA abundance data were converted to $\log_2$ scale for subsequent analyses. Mutational profiles were based on TCGA-reported MutSig v2.0 calls. All pre-processing was performed in R statistical environment (v3.1.3). Genetic ancestry imputed by Carrot-Zhang[87] *et al*. was incorporated.

PCAWG whole genome sequencing data calls were downloaded from the PCAWG consortium through Synapse. All data pre-processing was performed by the consortium as described[88]. The individual datasets are available at Synapse (https://www.synapse.org/) and are denoted with synXXXXX accession numbers (listed under Synapse ID); all these datasets are also mirrored at https://dcc.icgc.org. Tumour histological classifications were reviewed and assigned by the PCAWG Pathology and Clinical Correlates Working Group (annotation version 9; https://www.synapse.org/#!Synapse:syn10389158, https://www.synapse.org/#!Synapse:syn10389164). Ancestry imputation was performed using an ADMIXTURE23-like algorithm based on germline SNP profiles determined by whole-genome sequencing of the reference sample (https://www.synapse.org/#!Synapse:syn4877977). The consensus somatic SNV and indel (https://www.synapse.org/#!Synapse:syn7357330) file covers 2778 whitelisted samples from 2583 donors. Driver events were called by the PCAWG Drivers and Functional Interpretation Group (https://www.synapse.org/#!Synapse:syn11639581). Consensus CNA calls from the PCAWG Structural Variation Working Group were downloaded in VCF format (https://www.synapse.org/#!Synapse:syn8042988). Subclonal reconstruction was performed by the PCAWG Evolution and Heterogeneity Working Group (https://www.synapse.org/#!Synapse:syn8532460). SigProfiler mutation signatures were determined by the PCAWG Mutation Signatures and Processes Working Group for single base substitution (https://www.synapse.org/#!Synapse:syn11738669), doublet base substitution (https://www.synapse.org/#!Synapse:syn11738667) and indel (https://www.synapse.org/#!Synapse:syn11738668) signatures. Signatures data for TCGA, non-PCAWG WGS and non-TCGA WXS samples were downloaded from Synapse (https://www.synapse.org/#!Synapse:syn11804040).

Controlled access AACR GENIE data was downloaded from AACR Project GENIE for the
MSK project[89]. (https://portal.gdc.cancer.gov/projects/GENIE-MSK). AACR GENIE MSK
data was generated by the custom hybridization-based capture panel MSK-IMPACT,
which includes up to 410 genes. Preprocessed gene-wise SNV and CNA calls were
obtained through the GDC Data Portal. We filtered 6,841 metastatic tumours and
matched on tumour-types included in the TCGA and PCAWG datasets (**Supplementary
Table 1**) for better comparison across datasets. The final pan-cancer AACR GENIE
cohort consisted of 7,259 tumours across 35 tumour-types.

We used TCGA data describing 10,212 distinct TCGA tumour samples across 23 tumour-
types and 2,562 distinct PCAWG samples across 29 tumour-types. Tumour-types with no
age annotation were excluded from analysis. Age is treated as a continuous variable for
both TCGA and PCAWG analyses. We matched tumour-types between datasets as
described in **Supplementary Table 1**.

General Statistical Framework
For each genomic feature of interest, we used univariate tests first followed by false
discovery rate (FDR) adjustment to identify putative age-associations of interest ($q < 0.1$).
We used two-sided non-parametric univariate tests to minimize assumptions on the data.
For putative age-associations, we then follow up the univariate analysis with multivariate
modeling to account for potential confounders using bespoke models for each tumour-
type.

Model variables for each tumour context are presented in **Supplementary Table 1** and
were included based on availability of data (<15% missing), sufficient variability (at least
two levels) and collinearity (as assessed by variance inflation factor). Discrete data was
modeled using logistic regression (LGR). Continuous data was first transformed using the
Box-Cox family and modeled using linear regression (LNR). The Box-Cox family of
transformations is a formalized method to select a power transformation to better
approximate a normal-like distribution and stabilize variance. We used the Yeo-Johnson
extension to the Box-Cox transformation that allows for zeros and negative values[90].

FDR adjustment was performed for p-values for the age variable significance estimate
and an FDR threshold of 10% was used to determine statistical significance. More detail
is provided for each analysis below. A summary of all results is presented in
Supplementary Table 1. We present 95% confidence intervals for all tests.

Mutation Density
Performed for both TCGA, PCAWG and AACR GENIE data. Overall mutation prevalence
per patient was calculated as the sum of SNVs across all genes on the autosomes and
scaled to mutations/Mbp. Coding mutation prevalence only considers the coding regions
of the genome, and noncoding prevalence only considers the noncoding regions. TCGA
mutation density reflects coding mutation prevalence. AACR GENIE mutation density

reflects the targeted sequences of the 341-, 410- or 468-gene MSK-IMPACT panel used
for each sample. Mutation density was compared age using Spearman correlation for
both pan-cancer and tumour-type specific analysis. Comparisons with univariate q-values
meeting an FDR threshold of 10% were further analyzed using linear regression to adjust
for tumour subtype-specific variables. Mutation density analysis was performed
separately for each mutation context, with pan-cancer and tumour subtype p-values
adjusted together. Full mutation density results are in **Supplementary Table 2.**

Per year increase in SNV density was estimated by combining TCGA, PCAWG and
AACR GENIE data. We fit a linear regression model with formula SNV density ~ age +
project and took the coefficient estimate for age as the per year increase in SNV density
value.

Genome instability
Performed for TCGA, PCAWG and AACR GENIE data. Genome instability was calculated
as the percentage of the genome affected by copy number alterations. The number of
base pairs for each CNA segment was summed to obtain a total number of base pairs
altered per patient. The total number of base pairs was divided by the number of assayed
bases to obtain the percentage of the genome altered (PGA). Genome instability was
compared using Spearman correlation for both pan-cancer and tumour-type specific
analysis. Comparisons with univariate q-values meeting an FDR threshold of 10% were
further analyzed using linear regression to adjust for tumour subtype-specific variables.
Genome instability analysis was performed separately for each mutation context, with
pan-cancer and tumour subtype p-values adjusted together. Full mutation density results
are in **Supplementary Table 2.** Per year increase in PGA was estimated similarly to the
estimation for SNV density: we fit a linear regression model with formula PGA ~ age +
project and took the coefficient estimate for age as the per year increase in PGA.

Clonal structure and mutation timing analysis
Performed for PCAGW data only. Subclonal structure data was binarized from number of
subclonal clusters per tumour to monoclonal (one cluster) or polyclonal (more than one
cluster). Putative age-associations were identified using univariate logistic regression and
putative associations were further analysed using multivariate logistic regression. A
multivariate q-value threshold of 0.1 was used to determine statistically significant age-
associated clonal structure.

Mutation timing data classified SNVs, indels and SVs into clonal (truncal) or subclonal
groups. The proportion of truncal variants was calculated for each mutation type
($\frac{Number\ truncal\ SNVs}{total\ SNVs}$, *etc.*) to obtain proportions of truncal SNVs, indels and SVs for each
tumour. These proportions were compared using Spearman correlation. Univariate p-
values were FDR adjusted to identify putatively age-associated mutation timing. Linear
regression was used to adjust for confounding factors and a multivariate q-value threshold

of 0.1 was used to determine statistically significant age-a mutation timing. The mutation
timing analysis was performed separately for SNVs, indels and SVs. All results for clonal
structure and mutation timing analyses are in **Supplementary Table 2**.

Mutational Signatures analysis
Performed for TCGA and PCAWG data. For each signature, we compared the proportion
of tumours with any mutations attributed to the signatures ("signature-positive") using
logistic regression to identify univariate age-associations. Signatures with putative age-
associations were further analysed using multivariable logistic regression.

We also compared relative signature activity using the proportions of mutations attributed
to each signature. The numbers of mutations per signature were divided by total number
of mutations for each tumour to obtain the proportion of mutations attributed to the
signature. Spearman correlation was used. Putative age-associated signatures were
further analysed using multivariable linear regression after Box-cox adjustment.

Signatures that were not detected in a tumour subtype was omitted from analysis for that
tumour subtype. All results for clonal structure and mutation timing analyses are in
Supplementary Table 2.

Genome-spanning CNA analysis
Performed for TCGA, PCAWG and AACR GENIE data. The copy number profiles for
COSMIC driver CNA genes were extracted. Copy number calls were collapsed to ternary
(loss, neutral, gain) representation by combining loss groups (mono-allelic and bi-allelic)
and gain groups (low and high). Logistic regression was used to identify univariate age-
associated CNAs. After identifying candidate pan-cancer univariately significant genes,
multivariate logistic regression was used to adjust ternary CNA data for tumour-type-
specific variables. The genome-spanning analysis was performed separately for losses
and gains for each tumour subtype. All CNA results are in **Supplementary Tables 3-4**.

Driver SNV Analysis
Performed for TCGA, PCAWG and AACR GENIE data. We focused on driver events
described by the PCAWG consortium[65] and by COSMIC[64]. For TCGA and AACR GENIE,
we also applied a 1% recurrence filter. Driver mutation data was binarized to indicate
presence or absence of the driver event in each patient. Proportions of mutated genes
were compared using univariate logistic regression. A q-value threshold of 0.1 was used
to select genes for further multivariate analysis using binary logistic regression. SNV
density was included in all models. FDR correction was again applied and genes with
significant age terms were extracted from the models (q-value < 0.1). Driver event
analysis was performed separately for pan-cancer analysis and for each tumour subtype.
All SNV and driver event analysis results are in **Supplementary Table 6**.

mRNA functional analysis

Performed for TCGA data. Genes in bins altered by age-associated CNAs and SNVs after multivariate adjustment were further investigated to determine functional consequences. Tumour purity was included in all mRNA models. Tumours with available mRNA abundance data were matched to those used in CNA analysis. For each gene affected by an age-associated loss, its mRNA abundance was modeled against age, copy number loss status, an age-copy number loss interaction term, and tumour purity. The interaction term was used to identify genes with age-associated mRNA changes. FDR adjusted p-values and fold-changes were extracted for visualization. A q-value threshold of 0.1 was used for statistical significance. For genes affected by age-associated gains, the same procedure was applied using copy number gains. mRNA modeling results for age-associated CNAs and SNVs are in **Supplementary Tables 5-6**.

Survival analysis

Performed for TCGA data. Genes found to have significant (FDR threshold of 10%) age-associated CNAs and SNVs were also analyzed using Cox proportional hazards modelling after checking proportional hazards assumption. Cox proportional hazard regression models incorporating age, CNA/SNV status, and an age-CNA/SNV group interaction were fit for overall survival after checking the proportional hazards assumption. Age was treated as a continuous variable for modeling, but median dichotomized into 'low age' and 'high age' groups for visualization. FDR-adjusted interaction p-values and $\log_2$ hazard ratios were extracted for visualization. A q-value threshold of 0.1 was used to identify genes with sex-influenced survival. Survival modeling results for age-associated CNAs and SNVs are in **Supplementary Tables 5 and 6**.

Statistical Analysis & Data Visualization Software

All statistical analyses and data visualization were performed in the R statistical environment (v3.2.1) using the BPG[91] (v5.9.8) and Survival (v2.44-1.1) packages, and with Inkscape (v0.92.3).

References

1. Cook, M. B. *et al.* Sex disparities in cancer incidence by period and age. *Cancer Epidemiol. Biomark. Prev. Publ. Am. Assoc. Cancer Res. Cosponsored Am. Soc. Prev. Oncol.* **18**, 1174–1182 (2009).

2. Edgren, G., Liang, L., Adami, H.-O. & Chang, E. T. Enigmatic sex disparities in cancer incidence. *Eur. J. Epidemiol.* **27**, 187–196 (2012).

3. Elsaleh, H. *et al.* Association of tumour site and sex with survival benefit from adjuvant chemotherapy in colorectal cancer. *Lancet Lond. Engl.* **355**, 1745–1750 (2000).

4. Cook, M. B., McGlynn, K. A., Devesa, S. S., Freedman, N. D. & Anderson, W. F. Sex Disparities in Cancer Mortality and Survival. *Cancer Epidemiol. Biomarkers Prev.* **20**, 1629–1637 (2011).

5. Scelo, G., Li, P., Chanudet, E. & Muller, D. C. Variability of Sex Disparities in Cancer Incidence over 30 Years: The Striking Case of Kidney Cancer. *Eur. Urol. Focus* **4**, 586–590 (2018).

6. Zheng, D. *et al.* Sexual dimorphism in the incidence of human cancers. *BMC Cancer* **19**, 684 (2019).

7. Nipp, R. *et al.* Disparities in cancer outcomes across age, sex, and race/ethnicity among patients with pancreatic cancer. *Cancer Med.* **7**, 525–535 (2018).

8. Zhang, W., Edwards, A., Flemington, E. K. & Zhang, K. Racial disparities in patient survival and tumor mutation burden, and the association between tumor mutation burden and cancer incidence rate. *Sci. Rep.* **7**, 13639 (2017).

9. Siegel, R. L., Miller, K. D. & Jemal, A. Cancer statistics, 2020. *CA. Cancer J. Clin.* **70**, 7–30 (2020).

10. Torre, L. A. *et al.* Cancer statistics for Asian Americans, Native Hawaiians, and Pacific

Islanders, 2016: Converging incidence in males and females: Cancer Statistics for

Asian Americans, Native Hawaiians, and Pacific Islanders, 2016. *CA. Cancer J. Clin.*

66, 182–202 (2016).

11. DeSantis, C. E., Miller, K. D., Goding Sauer, A., Jemal, A. & Siegel, R. L. Cancer

statistics for African Americans, 2019. *CA. Cancer J. Clin.* **69**, 211–233 (2019).

12. Cook, P. J., Doll, R. & Fellingham, S. A. A mathematical model for the age distribution

of cancer in man. *Int. J. Cancer* **4**, 93–112 (1969).

13. de Magalhães, J. P. How ageing processes influence cancer. *Nat. Rev. Cancer* **13**,

357–365 (2013).

14. Harding, C., Pompei, F. & Wilson, R. Peak and decline in cancer incidence, mortality,

and prevalence at old ages. *Cancer* **118**, 1371–1386 (2012).

15. White, M. C. *et al.* Age and Cancer Risk. *Am. J. Prev. Med.* **46**, S7–S15 (2014).

16. DePinho, R. A. The age of cancer. *Nature* **408**, 248 (2000).

17. Armitage, P. & Doll, R. The age distribution of cancer and a multi-stage theory of

carcinogenesis. *Br. J. Cancer* **8**, 1–12 (1954).

18. Ershler, W. B. Cancer: a disease of the elderly. *J. Support. Oncol.* **1**, 5–10 (2003).

19. Aunan, J. R., Cho, W. C. & Søreide, K. The Biology of Aging and Cancer: A Brief

Overview of Shared and Divergent Molecular Hallmarks. *Aging Dis.* **8**, 628 (2017).

20. Steliarova-Foucher, E. *et al.* International incidence of childhood cancer, 2001–10: a

population-based registry study. *Lancet Oncol.* **18**, 719–731 (2017).

21. the St. Jude Children's Research Hospital–Washington University Pediatric Cancer
Genome Project. Whole-genome sequencing identifies genetic alterations in pediatric
low-grade gliomas. *Nat. Genet.* **45**, 602–612 (2013).

22. the St. Jude Children's Research Hospital–Washington University Pediatric Cancer
Genome Project. The genomic landscape of diffuse intrinsic pontine glioma and
pediatric non-brainstem high-grade glioma. *Nat. Genet.* **46**, 444–450 (2014).

23. Trama, A. *et al.* Survival of European adolescents and young adults diagnosed with
cancer in 2000–07: population-based data from EUROCARE-5. *Lancet Oncol.* **17**,
896–906 (2016).

24. Anders, C. K. *et al.* Young Age at Diagnosis Correlates With Worse Prognosis and
Defines a Subset of Breast Cancers With Shared Patterns of Gene Expression. *J.*
Clin. Oncol. **26**, 3324–3330 (2008).

25. Tingstedt, B., Weitkämper, C. & Andersson, R. Early onset pancreatic cancer: a
controlled trial. *Ann. Gastroenterol.* **24**, 206–212 (2011).

26. Piciucchi, M. *et al.* Early onset pancreatic cancer: Risk factors, presentation and
outcome. *Pancreatology* **15**, 151–155 (2015).

27. Katz, M. *et al.* The Effect of Race/Ethnicity on the Age of Colon Cancer Diagnosis. *J.*
Health Disparities Res. Pract. **6**, 62–69 (2013).

28. Tricoli, J. V. *et al.* Biologic and clinical characteristics of adolescent and young adult
cancers: Acute lymphoblastic leukemia, colorectal cancer, breast cancer, melanoma,
and sarcoma: Biology of AYA Cancers. *Cancer* **122**, 1017–1028 (2016).

29. Willauer, A. N. *et al.* Clinical and molecular characterization of early-onset colorectal
cancer. *Cancer* **125**, 2002–2010 (2019).

30. Lieu, C. H. *et al.* Comprehensive Genomic Landscapes in Early and Later Onset

Colorectal Cancer. *Clin. Cancer Res.* **25**, 5852–5858 (2019).

31. Parsons, D. W. *et al.* An Integrated Genomic Analysis of Human Glioblastoma

Multiforme. *Science* **321**, 1807–1812 (2008).

32. Williams, G. R. *et al.* Comorbidity in older adults with cancer. *J. Geriatr. Oncol.* **7**, 249–

257 (2016).

33. Jørgensen, T. L., Hallas, J., Friis, S. & Herrstedt, J. Comorbidity in elderly cancer

patients in relation to overall and cancer-specific mortality. *Br. J. Cancer* **106**, 1353–

1360 (2012).

34. Balducci, L., Colloca, G., Cesari, M. & Gambassi, G. Assessment and treatment of

elderly patients with cancer. *Surg. Oncol.* **19**, 117–123 (2010).

35. Given, B. & Given, C. W. Older adults and cancer treatment. *Cancer* **113**, 3505–3511

(2008).

36. Chen, R. C., Royce, T. J., Extermann, M. & Reeve, B. B. Impact of Age and

Comorbidity on Treatment and Outcomes in Elderly Cancer Patients. *Semin. Radiat.*

Oncol. **22**, 265–271 (2012).

37. Andaya, A. A. *et al.* Race and Colon Cancer Survival in an Equal-Access Health Care

System. *Cancer Epidemiol. Biomarkers Prev.* **22**, 1030–1036 (2013).

38. Milholland, B., Auton, A., Suh, Y. & Vijg, J. Age-related somatic mutations in the

cancer genome. *Oncotarget* **6**, 24627–24635 (2015).

39. Wahl, D. R. *et al.* Pan-Cancer Analysis of Genomic Sequencing Among the Elderly.

Int. J. Radiat. Oncol. **98**, 726–732 (2017).

40. Li, C. H., Haider, S., Shiah, Y.-J., Thai, K. & Boutros, P. C. Sex Differences in Cancer

Driver Genes and Biomarkers. *Cancer Res.* **78**, 5527 (2018).

41. Chalmers, Z. R. *et al.* Analysis of 100,000 human cancer genomes reveals the

landscape of tumor mutational burden. *Genome Med.* **9**, (2017).

42. Martincorena, I. *et al.* Somatic mutant clones colonize the human esophagus with

age. *Science* **362**, 911–917 (2018).

43. Martincorena, I. *et al.* High burden and pervasive positive selection of somatic

mutations in normal human skin. *Science* **348**, 880–886 (2015).

44. Lee-Six, H. *et al.* The landscape of somatic mutation in normal colorectal epithelial

cells. *Nature* **574**, 532–537 (2019).

45. Yokoyama, A. *et al.* Age-related remodelling of oesophageal epithelia by mutated

cancer drivers. *Nature* **565**, 312–317 (2019).

46. Suda, K. *et al.* Clonal Expansion and Diversification of Cancer-Associated Mutations

in Endometriosis and Normal Endometrium. *Cell Rep.* **24**, 1777–1789 (2018).

47. Jaiswal, S. *et al.* Age-Related Clonal Hematopoiesis Associated with Adverse

Outcomes. *N. Engl. J. Med.* **371**, 2488–2498 (2014).

48. Welch, J. S. *et al.* The Origin and Evolution of Mutations in Acute Myeloid Leukemia.

Cell **150**, 264–278 (2012).

49. Vollan, H. K. M. *et al.* A tumor DNA complex aberration index is an independent

predictor of survival in breast and ovarian cancer. *Mol. Oncol.* **9**, 115–127 (2015).

50. Lalonde, E. *et al.* Tumour genomic and microenvironmental heterogeneity for

integrated prediction of 5-year biochemical recurrence of prostate cancer: a

retrospective cohort study. *Lancet Oncol.* **15**, 1521–1532 (2014).

51. Hieronymus, H. *et al.* Copy number alteration burden predicts prostate cancer

relapse. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 11139–11144 (2014).

52. Rizvi, N. A. *et al.* Mutational landscape determines sensitivity to PD-1 blockade in

non–small cell lung cancer. *Science* **348**, 124–128 (2015).

53. Samstein, R. M. *et al.* Tumor mutational load predicts survival after immunotherapy

across multiple cancer types. *Nat. Genet.* **51**, 202–206 (2019).

54. PCAWG Tumour Subtypes and Clinical Translation *et al.* Sex differences in oncogenic

mutational processes. *Nat. Commun.* **11**, 4330 (2020).

55. PCAWG Evolution & Heterogeneity Working Group *et al.* The evolutionary history of

2,658 cancers. *Nature* **578**, 122–128 (2020).

56. Espiritu, S. M. G. *et al.* The Evolutionary Landscape of Localized Prostate Cancers

Drives Clinical Aggression. *Cell* **173**, 1003-1013.e15 (2018).

57. Landau, D. A. *et al.* Evolution and Impact of Subclonal Mutations in Chronic

Lymphocytic Leukemia. *Cell* **152**, 714–726 (2013).

58. Shah, S. P. *et al.* The clonal and mutational evolution spectrum of primary triple-

negative breast cancers. *Nature* **486**, 395–399 (2012).

59. Asomaning, K. *et al.* Second hand smoke, age of exposure and lung cancer risk. *Lung*

Cancer **61**, 13–20 (2008).

60. Alexandrov, L. B. *et al.* Signatures of mutational processes in human cancer. *Nature*

500, 415–421 (2013).

61. Alexandrov, L. B. *et al.* Mutational signatures associated with tobacco smoking in

human cancer. *Science* **354**, 618–622 (2016).

62. PCAWG Mutational Signatures Working Group *et al.* The repertoire of mutational

signatures in human cancer. *Nature* **578**, 94–101 (2020).

63. Alexandrov, L. *et al.* The Repertoire of Mutational Signatures in Human Cancer.

(2018) doi:10.1101/322859.

64. Tate, J. G. *et al.* COSMIC: the Catalogue Of Somatic Mutations In Cancer. *Nucleic*

Acids Res. **47**, D941–D947 (2019).

65. PCAWG Drivers and Functional Interpretation Working Group *et al.* Analyses of non-

coding somatic drivers in 2,658 cancer whole genomes. *Nature* **578**, 102–111 (2020).

66. Stewart, J. B. *et al.* Simultaneous DNA and RNA Mapping of Somatic Mitochondrial

Mutations across Diverse Human Cancers. *PLOS Genet.* **11**, e1005333 (2015).

67. Hopkins, J. F. *et al.* Mitochondrial mutations drive prostate cancer aggression. *Nat.*

Commun. **8**, 656 (2017).

68. Ye, C. *et al.* Mutations in the mitochondrial DNA D-loop region and breast cancer risk.

Breast Cancer Res. Treat. **119**, 431–436 (2010).

69. Nugent, S. *et al.* Brain glucose and acetoacetate metabolism: a comparison of young

and older adults. *Neurobiol. Aging* **35**, 1386–1395 (2014).

70. van den Bent, M. J., Smits, M., Kros, J. M. & Chang, S. M. Diffuse Infiltrating

Oligodendroglioma and Astrocytoma. *J. Clin. Oncol.* **35**, 2394–2401 (2017).

71. Li, C. H. *et al. Sex Differences in Oncogenic Mutational Processes.*

http://biorxiv.org/lookup/doi/10.1101/528968 (2019) doi:10.1101/528968.

72. Chatsirisupachai, K., Lesluyes, T., Paraoan, L., Van Loo, P. & de Magalhães, J. P. *An*

Integrative Analysis of the Age-Associated Genomic, Transcriptomic and Epigenetic

Landscape across Cancers. http://biorxiv.org/lookup/doi/10.1101/2020.08.25.266403

(2020) doi:10.1101/2020.08.25.266403.

73. Maynard, S., Fang, E. F., Scheibye-Knudsen, M., Croteau, D. L. & Bohr, V. A. DNA

Damage, DNA Repair, Aging, and Neurodegeneration. *Cold Spring Harb. Perspect.*

Med. **5**, a025130 (2015).

74. Hoeijmakers, J. H. J. DNA Damage, Aging, and Cancer. *N. Engl. J. Med.* **361**, 1475–

1485 (2009).

75. Fernández, L. C., Torres, M. & Real, F. X. Somatic mosaicism: on the road to cancer.

Nat. Rev. Cancer **16**, 43–55 (2016).

76. Tomasetti, C., Li, L. & Vogelstein, B. Stem cell divisions, somatic mutations, cancer

etiology, and cancer prevention. *Science* **355**, 1330–1334 (2017).

77. Özdemir, B. C. & Dotto, G.-P. Racial Differences in Cancer Susceptibility and Survival:

More Than the Color of the Skin? *Trends Cancer* **3**, 181–197 (2017).

78. Cramer, D. W. & Finn, O. J. Epidemiologic perspective on immune-surveillance in

cancer. *Curr. Opin. Immunol.* **23**, 265–271 (2011).

79. Vick, A. D. & Burris, H. H. Epigenetics and Health Disparities. *Curr. Epidemiol. Rep.*

4, 31–37 (2017).

80. Huang, K. *et al.* Pathogenic Germline Variants in 10,389 Adult Cancers. *Cell* **173**, 355-

370.e14 (2018).

81. Research, W. C. R. F. I. for C. Diet, nutrition, physical activity and cancer: a global

perspective. *Contin. Update Proj. Expert Rep.* (2018).

82. Gopalakrishnan, V., Helmink, B. A., Spencer, C. N., Reuben, A. & Wargo, J. A. The

Influence of the Gut Microbiome on Cancer, Immunity, and Cancer Immunotherapy.

Cancer Cell **33**, 570–580 (2018).

83. Bostwick, D. G. *et al.* Human prostate cancer risk factors. *Cancer* **101**, 2371–2490

(2004).

84. Irigaray, P. & Belpomme, D. Basic properties and molecular mechanisms of

exogenous chemical carcinogens. *Carcinogenesis* **31**, 135–148 (2010).

85. Talarico, L., Chen, G. & Pazdur, R. Enrollment of Elderly Patients in Clinical Trials for

Cancer Drug Registration: A 7-Year Experience by the US Food and Drug

Administration. *J. Clin. Oncol.* **22**, 4626–4631 (2004).

86. Ruiter, R., Burggraaf, J. & Rissmann, R. Under-representation of elderly in clinical

trials: An analysis of the initial approval documents in the Food and Drug

Administration database. *Br. J. Clin. Pharmacol.* **85**, 838–844 (2019).

87. Carrot-Zhang, J. *et al.* Comprehensive Analysis of Genetic Ancestry and Its Molecular

Correlates in Cancer. *Cancer Cell* **37**, 639-654.e6 (2020).

88. The ICGC/TCGA Pan-Cancer Analysis of Whole Genomes Consortium. Pan-cancer

analysis of whole genomes. *Nature* **578**, 82–93 (2020).

89. The AACR Project GENIE Consortium. AACR Project GENIE: Powering Precision

Medicine through an International Consortium. *Cancer Discov.* **7**, 818–831 (2017).

90. Yeo, I.-K. & Johnson, R. A. A New Family of Power Transformations to Improve

Normality or Symmetry. *Biometrika* **87**, 954–959 (2000).

91. P'ng, C. *et al.* BPG: Seamless, automated and interactive visualization of scientific

data. *BMC Bioinformatics* **20**, (2019).

Acknowledgments

The authors thank all the members of the Boutros lab for insightful discussion. We also thank João Pedro de Magalhães, Kasit Chatsirisupachai, Peter Van Loo, and the de Magalhães lab for collaborative discussion. This work was supported by the Discovery Frontiers: Advancing Big Data Science in Genomics Research program, which is jointly funded by the Natural Sciences and Engineering Research Council (NSERC) of Canada, the Canadian Institutes of Health Research (CIHR), Genome Canada and the Canada Foundation for Innovation (CFI). P.C.B. was supported by a Terry Fox Research Institute New Investigator Award and a CIHR New Investigator Award. This work was supported by an NSERC Discovery grant and by Canadian Institutes of Health Research, grant #SVB-145586, to PCB. The results described here are in part based upon data generated by the TCGA Research Network: http://cancergenome.nih.gov/. This work was supported by the NIH/NCI under award number P30CA016042 and by an operating grant from the National Cancer Institute Early Detection Research Network (U01CA214194).

Author Contributions

CHL and PCB initiated the project. CHL, and SH analyzed data. PCB supervised research. CHL and PCB wrote the first draft of the manuscript, which all authors edited and approved.

Figures & Figure Legends

Figure 1

Figure 1 | Mutation density and timing are associated with age at diagnosis.
Summary of associations between age and **(A)** SNV density and **(D)** PGA in TCGA,
PCAWG, and AACR GENIE tumours. The dot size and colour show the Spearman
correlation, and background shading indicate adjusted multivariate p-value. Only tumour-
types with at least univariately significant associations are shown. Associations between
(B) SNV density and **(E)** PGA with age in prostate cancer across the three datasets.
Univariate Spearman correlation, adjusted correlation p-value and adjusted multivariate
p-values shown. **(C)** Correlations between age and proportion of SNVs occurring in the
truncal clone in four PCAWG tumour contexts. **(F)** In TCGA lung adenocarcinoma, age
and PGA are associated with smoking history, and **(G)** the negative association between
PGA and age remains significant in current smokers and recently reformed smokers
(<=15 years). Dot colours show smoking history groups as indicated in **(F)**.

Figure 2

Figure 2 | Age-associated mutational signatures suggest differences in underlying mutational processes.

(A) In TCGA lung adenocarcinoma, age (top plots) and PGA (bottom plots) are associated
with the tobacco-associated signature SBS4 (left). The absolute number of SBS4-
attributed mutations is also associated with age and PGA (middle). The relative proportion
of SBS4-attibuted mutations is negatively associated with age and has no significant
relationship with PGA (right). **(B)** The negative association between PGA and age
remains significant in both SBS4+ and SBS4- groups. Yellow and blue dots indicate
SBS4- and SBS4+ tumours, respectively. **(C)** Summary of associations between age and
the proportion of signature-positive tumours, where dot size shows the marginal log odds
from logistic regression and background shading show adjusted multivariate p-values.
PCAWG data is on left and TCGA on right. **(D)** Similarly, the summary of associations
between age and relative signature activity, with dot size showing Spearman correlations
and background indicating adjusted linear regression p-values. **(E)** Comparison of
PCAWG and TCGA signature detection frequency. Filled in and open circles indicate
comparisons where the differences are statistically significant (proportion test q < 0.05)
and not, respectively. Proposed SBS signature aetiologies are as indicated. Proposed
DBS and ID aetiologies are: DBS1: UV, DBS2: tobacco, DBS5: platinum chemotherapy,
DBS7: defective MMR, ID1: slippage during DNA replication, ID2: slippage during DNA
replication, ID3: tobacco, ID6: defective HR, ID8: NHEJ, ID13: UV

Figure 3

Figure 3 | Age-associations in copy number drivers are associated with functional changes in mRNA and survival.

Summary of all detected age-associated driver CNAs for **(A)** pan-cancer and **(B-C)** tumour-type specific analyses across three datasets. Dot size shows the magnitude of the association as the difference in proportion and the background shading shows adjusted multivariate p-values. Right covariate indicates copy number gain drivers in red and loss drivers in blue. **(D)** Age-associated CNAs lead to differential mRNA abundance with respect to the CNA itself (left), age (middle), as well as age-specific effects of the CNA (right). **(E)** *CDKN2A* mRNA abundance changes between copy number loss (blue) or no loss (black) and broken down by age for low *vs.* high age. Tukey boxplots are shown with the box indicating quartiles and the whiskers drawn at the lowest and highest points within 1.5 interquartile range of the lower and upper quartiles, respectively. Adjusted p-values are indicated. **(F)** *SUFU* loss interacts with age to further stratify patient prognosis. The adjusted p-value for the copy number loss-age interaction term is shown.

Figure 4

age-associated prognostic biomarker in lower grade glioma.

(A) Summary of age- associated nuclear driver SNVs for **(A)** pan-cancer, **(B)** tumour-type specific analyses across three datasets. Dot size shows the magnitude of the association as the difference in proportion and the background shading shows adjusted multivariate p-values. Left covariate in **(B)** indicates relevant tumour-type. **(C)** TCGA lower grade glioma age-associations in driver mutation frequency with adjusted multivariate p-values, marginal log odds changes for 10-year age increment, and age of tumours compared between those with (red) and without (grey) the mutation. **(D)** Summary of age-associated mitochondrial SNVs in PCAWG data with specific examples from ovarian cancer shown in **(E)**. **(F)** Mitochondrial copy number foldchange is also associated with age in four tumour contexts. **(G)** In TCGA lower grade glioma: mRNA abundance changes for *IDH1* and *ATRX* when the gene is mutated (red) or not (black) compared by median-dichotomized age. Adjusted SNV-age interaction p-values are shown. **(H)** *ATRX* mutation interacts with age to stratify patient prognosis into four groups. Log-odds p-value is shown. Tukey boxplots are shown with the box indicating quartiles and the whiskers drawn at the lowest and highest points within 1.5 interquartile range of the lower and upper quartiles, respectively.