Bayesian inference of population prevalence

Robin A. A. Ince[1], Jim W. Kay[2], Philippe G. Schyns[1,3]

[1] Institute of Neuroscience and Psychology, University of Glasgow
[2] Department of Statistics, University of Glasgow
[3] School of Psychology, University of Glasgow

Abstract

Within neuroscience, psychology and neuroimaging, the most frequently used statistical approach is null-hypothesis significance testing (NHST) of the population mean. An alternative approach is to perform NHST within individual participants and then infer, from the proportion of participants showing an effect, the prevalence of that effect in the population. We propose a novel Bayesian method to estimate such population prevalence that offers several advantages over population mean NHST. This method provides a population-level inference that is currently missing from study designs with small participant numbers, such as in traditional psychophysics and in precision imaging. It delivers a quantitative estimate with associated uncertainty instead of reducing an experiment to a binary inference on a population mean. Bayesian prevalence is widely applicable to a broad range of studies in neuroscience, psychology and neuroimaging. Its emphasis on detecting effects within individual participants could also help address replicability issues in these fields.

Introduction

Within neuroscience, psychology and neuroimaging, the common experimental paradigm is to run an experiment on a sample of participants and then infer and quantify any effect of the experimental manipulation in the population from which the sample was drawn. For example, in a psychology experiment, a particular set of stimuli (e.g. visual or auditory stimuli) might be presented to a sample of human participants, who are asked to categorize the stimuli or to perform some other task. Each participant repeats the procedure several times with different stimuli (experimental trials), and their responses and reaction times are recorded. In a neuroimaging experiment, the same procedure is employed with neuroimaging signals recorded in addition to behavioural responses. The researcher analyses these responses to infer something about brain function in the population from which the participants were drawn.

In this standard experimental paradigm, the implicit goal is usually to determine the presence of a causal relationship between the experimental manipulation and the response of interest. For example, between a stimulus property, and the neural activity in a particular brain region, as reflected through non-invasive neuroimaging signals, or between neural activity and behaviour. A properly controlled experimental design in which other extraneous factors are held constant (i.e. a randomised control trial) enables a causal interpretation of a correlational relationship (Angrist and Pischke, 2014; Pearl, 2009). We use tools from statistics to evaluate the measured effect, and to ensure that we are not being fooled by randomness— that what we observe does not result from random fluctuations that we could expect to see by chance even if there was no effect. This is often formalised as Null Hypothesis Significance Testing (NHST). We reject the null hypothesis when the probability, if the null hypothesis was true, of observing an effect as large as that actually observed is less than some prespecified value (often *p=0.05*). Simply stated, we would be unlikely to obtain the observed effect due to chance, if the null hypothesis was true.

Researchers performing such studies usually wish to infer something about the population from which the experimental participants are selected (Holmes and Friston, 1998), rather than about the specific sample of participants that were examined (as in a case study). Importantly, any statistical inference from a sample to the population requires a model of the effect in the population. The ubiquitous approach used in psychology and neuroimaging is to model the effect in the population with a normal distribution and perform inference on the mean of this model: the population mean (see Methods). For example, the null hypothesis is often that the population mean is zero, and the probability of the observed sample data under this null is computed, taking the variance across the sample as an estimate of the variance across the population. However, an alternative and equally valid question to ask is how typical is an effect in the population (Friston et al., 1999a). In this approach, we infer an effect in each individual of the sample, and from that infer the *prevalence* of the effect in the population—i.e. the proportion of the population that would show the effect, if tested (Allefeld et al., 2016; Donhauser et al., 2018; Friston et al., 1999b; Rosenblatt et al., 2014). The results obtained using these two approaches, population mean versus population prevalence, can differ, particularly when effects are heterogenous across participants.

Here, we argue that in many experimental applications in psychology and neuroscience, the individual participant is the natural replication unit of interest (Little and Smith, 2018; Nachev et al., 2019; Smith and Little, 2018; Thiebaut de Schotten and Shallice, 2017). This is because many aspects of cognition, and the neural mechanisms underlying them, are likely to be heterogenous across individuals. Therefore, we should seek to quantify effects within individual participants, and ensure that our results can be reliably distinguished from chance within individual participants. We argue that with this shift in perspective towards experimental assessment within individual participants, our statistical focus at the population level could also shift from NHST of the population mean to estimating the population prevalence: the proportion of individuals in the population who would show a true positive above-chance effect in a specific experiment. Although we focus here on a broad class of experiments in psychology and neuroimaging that feature human participants and non-invasive recording modalities, the arguments we present are general and apply equally well to other experimental model organisms or sampling units.

To support this shift in perspective, we present a simple but novel Bayesian method to estimate population prevalence based on the results of within-participant NHST, including prevalence differences between groups of participants or between tests on the same participants. This approach can also be applied without explicit within-participant NHST to estimate prevalence of different effect sizes, giving a new view on what can be learned about a population from an experimental sample. We suggest that applying Bayesian population prevalence in studies that are sufficiently powered within individual participants could address many of the recent issues raised about replicability in psychology and neuroimaging research (Benjamin et al., 2018; Ioannidis, 2005). This novel approach provides a population prevalence estimate with associated uncertainty and therefore avoids reducing an entire experiment to a binary NHST inference on a population mean (McShane et al., 2019).

Results

Figure 1: *Population vs individual inference*. For each simulation, we sample $N = 50$ individual participant mean effects from a normal distribution with population mean μ (**A,B**: $\mu = 0$; **C,D**: $\mu = 1$) and between-participant standard deviation $\sigma_b = 2$. Within each participant, T trials (**A,C**: $T = 20$; **B,D**: $T = 500$) are drawn from a normal distribution with the participant-specific mean and a common within-participant standard deviation $\sigma_w = 10$ (Baker et al. 2019). Orange and blue indicate, respectively, exceeding or not exceeding a $p = 0.05$ threshold for a t-test at the population level (on the within-participant means, population normal density curves) or at the individual participant level (individual sample means +/- s.e.m.). **E**: Bayesian posterior distributions of population prevalence of true positive results for the 4 simulated data sets (A-D). Circles show Bayesian maximum a posteriori estimates. Thick and thin horizontal lines indicate 50% and 96% highest posterior density intervals, respectively. MAP [96% HPDI] values are shown in the legend.

Population vs within-participant statistical tests

To illustrate, we simulate data from the standard hierarchical population model that underlies inference of a population mean effect based on the normal distribution (Friston, 2007; Holmes and Friston, 1998; Penny and Holmes, 2007) (see Methods). Figure 1 illustrates the results of four simulations that differ in the true population mean, μ (A, B: $\mu = 0$; C, D, $\mu = 1$) and in the number of trials performed per participant (A, C: 20 trials, B, D: 500 trials).

For each simulation, we performed inference based on a standard two-sided, one-sample t-test against zero at two levels. First, we applied the standard summary statistic approach: we

took the mean across trials for each participant and performed the second-level population t-test on these per-participant mean values. This provides an inference on the population mean, taking into account between-participant variation. This is equivalent to inference on the full hierarchical model in a design where participants are separable (Holmes and Friston, 1998; Penny and Holmes, 2007; Penny et al., 2003) (see Methods). The modelled population distribution is plotted as a solid curve, coloured according to the result of the population-mean inference (orange for significant population mean, blue for non-significant population mean). Second, we performed inference within each participant, applying the t-test on within-participant samples, separately for each participant. The sample mean +/- s.e.m. is plotted for each participant (orange for significant participants, blue for non-significant participants).

The population mean inference correctly fails to reject the null hypothesis for Figure 1 panels A and B (ground truth $\mu = 0$), and correctly rejects the null in panels C and D (ground truth $\mu = 1$). But consider carefully Figure 1B and C. With 500 trials in panel B, 32/50 participants (orange markers) show a significant within-participant result. The probability of this happening by chance, if there was no effect in any members of the population, can be calculated from the cumulative density function of the binomial distribution. In this case, it is tiny—for a test with false positive rate $\alpha = 0.05$, and no effect in any individual, $p < 2.2 \times 10^{-16}$ (below 64-bit floating-point precision). Compare that to $p = 0.008$ for the population t-test in panel C. Thus, the panel B results provide much stronger statistical evidence of a non-zero effect *at the population level*; the observed results are very unlikely if the proportion of individuals in the population that would show an effect in this test is zero. This would be ignored in analyses based only on the population mean. Considering inference at the individual level, panel C results (11/50 significant) have $p = 4.9 \times 10^{-6}$ if the proportion of the population with the effect was zero, i.e. there was no effect within any individuals. Thus, even panel C, which reports simulating an experiment with only 20 trials per participant, provides stronger evidence for a population effect from the within-participant perspective than from the population-mean perspective.

Obviously, these two different p-values are calculated from two different null hypotheses. The population t-test tests the null hypothesis that the population mean is zero, assuming that individual means follow a normal distribution:

$$H_0: \mu_{pop} = 0$$

while the p-value for the number of significant within-participant tests comes from the null hypothesis that there is no effect in any individual in the population, termed the *global null* (Allefeld et al., 2016; Donhauser et al., 2018; Nichols et al., 2005):

$$H_0: \mu_i = 0 \text{ for all } i$$

where i indexes all members of the population. These are testing different questions, but importantly both are framed at the population level and both provide a population-level inference. We agree that it is important to align "the meaning of the quantitative inference with the meaning of the qualitative hypothesis we're interested in evaluating" (Yarkoni, 2019). We suggest that often, if the goal of the analysis is to infer the presence of a causal relationship within individuals in the population, the within-participant perspective may be more appropriate. Clearly, the global null itself is quite a blunt tool, as it would be untrue mathematically if even one in a million participants showed an effect. The goal of the

prevalence methods we present here is to quantify within-participant effects at the population level in a more meaningful and graded way.

We emphasise that this is the simplest possible illustrative example, intended to demonstrate the different perspective of within-participant vs. population mean inference. This example is derived from the standard null model implicit in almost every random effects inference employed in psychology and neuroimaging studies. Although we have only considered the global null, the simulations show that the within-participant perspective can give a very different view of the evidence for a population-level effect provided by a specific dataset. We should also emphasise that the simulated data values within individuals are considered to represent the *effect* of an experimental manipulation (e.g. the difference in a neuroimaging response between stimuli of different classes). Therefore, the results illustrated in Figure 1B represent strong evidence of a non-zero effect within many participants, albeit one that seems to be approximately balanced across participants between positive and negative directions. We do not suggest that such two-sided effects are particularly common but wish to highlight how they are implicitly assumed in traditional population mean inference models. In fact, if the researcher believes all effects in an experiment are positive, then the standard null model illustrated here is inappropriate, and the traditional population mean t-test is actually providing a fixed-effects analysis based on the global null rather than the random-effects interpretation of generalization to the population that is normally applied (Allefeld et al., 2016). However, the prevalence method we develop below is not limited to two-sided effects, we could just as well use a one-sided t-test in these examples.

In fact, the point we make here is applicable to any within-participant statistical test of an experimental manipulation. For example, we could consider a within-participant d-prime, a coefficient or contrast from a linear model (e.g. a General Linear Model of fMRI data), a cross-validated out-of-sample predictive correlation from a high-dimensional stimulus encoding model (e.g. a model predicting auditory cortex MEG responses to continuous speech stimuli), a rank correlation of dissimilarity matrices in a Representational Similarity Analysis, or parameters of computational models of decision making (e.g. the Diffusion Drift Model). Equivalence tests (Lakens, 2017) could be applied within each participant and prevalence estimation applied to effects that are significantly smaller than the SESOI. In all of these cases, the distinction between performing inference on the population mean vs. within individual participants still holds. We argue that performing NHST at the individual participant level is preferable for conceptual reasons in psychology and neuroimaging and also for practical reasons related to the replicability crisis (see Discussion).

Estimating population prevalence

The *p*-values under the global null are obtained from the cumulative density function of the binomial distribution, based on a within-participant false positive rate $\alpha = 0.05$. However, we can also model the number of above-threshold individuals in a sample when the population prevalence of true positive (significant) test results is γ (see Methods). Consider a within-participant test with a false-positive rate α. In this case, the distribution of the number of significant test results follows a binomial distribution with success probability $\theta = (1 - \gamma)\alpha + \gamma$. Here, we present a Bayesian approach to estimate the population prevalence of true positive test results, γ, from this binomial model of within-participant testing. Alternatively, frequentist inference approaches can be used (Allefeld et al., 2016; Donhauser

et al., 2018; Friston et al., 1999b). Note that we are not estimating the population prevalence of the ground truth binary status of the tested effect. We could only obtain a lower bound on this prevalence, because there might be individuals with an effect too small to detect with our test. Therefore, throughout we focus on the prevalence of true positive test results, i.e. the proportion of the population we would expect to give a true positive test result given the specific experimental test procedure considered.

The Bayesian approach provides a full posterior distribution for γ, from which we can obtain the *maximum a posteriori* (MAP) estimate, together with measures of uncertainty—e.g. highest posterior density intervals (HPDIs) or lower bound quantiles. Figure 1E shows the Bayesian posteriors, MAPs and HPDIs for the 4 simulated data sets in Figure 1A-D. Even though there is no population mean effect in Figure 1B, the MAP estimate of the prevalence is 0.62 (96% HPDI: [0.47 0.76]). Given the data, the probability that the population prevalence is greater than 47% is higher than 0.96. Based on this result, we would consider it highly likely that >47% of the population would show a true positive effect in the same experiment with 500 trials.

Figure 2 illustrates how Bayesian prevalence inference scales with numbers of participants and trials. Figure 2A-C suggests that there are benefits to having larger numbers of participants for the Bayesian prevalence metrics (including, decrease in variance of obtained MAP and HPDI width, increase in prevalence lower bound). However, above around 50 participants, these benefits become less pronounced. Figures 2E shows that inferred prevalence is mostly sensitive to the number of trials per participant (horizontal contours), and is invariant to the number of participants (although variance decreases with increasing N, as in Figure 2A,C,F). By comparison, t-test power (Figure 2D) is mostly sensitive to the number of participants (vertical contours) and is largely invariant to the number of trials above around 100 trials per participant (Baker et al., 2019). In sum, compared to the population mean t-test, prevalence exhibits greater sensitivity to the number of trials obtained per participant, and less sensitivity to the number of participants.

Estimating differences in prevalence

Often the scientific question of interest might involve comparing an effect between groups of participants, or between different experimental conditions in the same set of participants. For example, in the first case, a researcher might be interested in looking for differences in a behavioural or neuroimaging effect between patients and healthy controls, or between participants from two different cultures. In the second case, a researcher might be interested in determining the effect of an intervention on a particular measured effect, testing the same set of participants on two occasions, once with an intervention and once in a control condition. From the population-mean perspective these questions would often be addressed with a two-sample unpaired t-test for the former case, and a two-sample paired t-test for the latter. From the prevalence perspective, the question would be whether the prevalence of true positive results differs between two sampled populations (in the first case) or between two experimental tests (in the second case). We therefore provide additional functions (see Methods) to directly estimate the difference in prevalence of true positive test results for these two comparisons, which we term between-group (same test applied to two different samples) and within-group (two different tests applied to a single sample).

To estimate the difference in prevalence of true positive results for a given test between two samples from separate populations (e.g. patients vs healthy controls), the input data required is the count of positive results and the total number of participants in each group. We illustrate this with a simulated dataset. We specify the true prevalence in the two populations as $\gamma_1 = 0.75$ and $\gamma_2 = 0.25$ respectively, and draw a random sample based on the respective binomial distribution for the parameters θ_i (see Methods). We simulate $N_1 = 60$ participants in the first group and $N_2 = 40$ participants in the second group. The results of one such draw gives $k_1 = 45, k_2 = 11$ positive tests in each group, respectively. In this case, the MAP [96% HPDI] prevalence difference $\gamma_1 - \gamma_2$, calculated from these four values (k_1, k_2, N_1, N_2), is 0.49 [0.29 0.67], which closely matches the ground truth (0.5). Figure 3A-B shows how the between-group posterior prevalence difference estimates scale with the number of participants for three different simulations.

To estimate the difference in prevalence of true positive results between two different tests applied to the same sample of participants, the input parameters are the number of participants significant in both tests, the number significant only in each of the two tests, and the total number of participants. We simulate two tests applied to a group of $N = 50$ participants. Each test detects a certain property with false positive rate $\alpha = 0.05$. The ground truth prevalence of true positives for the two tests are $\gamma_1 = 0.5$ and $\gamma_2 = 0.25$, respectively, and the correlation between positive results is $\rho_{12} = 0.2$ (i.e. participants who test positive on one test are more likely to also test positive on the other test). The results of one random draw from this model gives: (see Methods) $k_{11} = 8$ participants with a

significant result in both tests; $k_{10} = 19$ participants with a significant result in the first test but not the second; and $k_{01} = 5$ participants with a significant result in the second but not the first. In this case, the MAP [96% HPDI] prevalence difference $\gamma_1 - \gamma_2$, calculated from these four values ($k_{11}, k_{10}, k_{01}, N$), is 0.28 [0.08 0.46], which again matches the ground truth (0.25). Figure 3C-D shows how the within-group posterior prevalence difference estimates scale with the number of participants for three different ground truth situations, given as $[\gamma_1 \; \gamma_2] \; \rho_{12}$.

Both these approaches are implemented using Monte Carlo methods, and the functions return posterior samples (Gelman, 2014), which can be used to calculate other quantities, such as the posterior probability that one test or group has a higher prevalence than the other. The posterior log odds in favour of this hypothesis can be computed by applying the logit function to the proportion of posterior samples in favour of a hypothesis. In the between-group example above, the hypothesis $\gamma_1 > \gamma_2$ has a proportion 0.9999987 of posterior samples in favour, corresponding to log posterior odds of 13.5. In the above within-group comparison, the hypothesis $\gamma_1 > \gamma_2$ has a proportion 0.9979451 of posterior samples in favour, corresponding to log posterior odds of 6.2 (each computed from 10 million posterior samples).

Figure 3: *Bayesian inference of difference of prevalence*. **A,B**: We consider two independent groups of participants with population prevalence of true positives $[\gamma_1, \gamma_2]$ of [25% 25%] (blue), [25% 50%] (red) and [25% 75%] (yellow). We show how **A**: the Bayesian MAP estimate, and **B**: 96% HPDI width, of the estimated between-group prevalence difference $\gamma_1 - \gamma_2$ scale with the number of participants. **C,D**: We consider two tests applied to the same sample of participants. Here, each simulation is parameterised by the population prevalence of true positives for the two tests, $[\gamma_1, \gamma_2]$, as well as ρ_{12}, the correlation between the (binary) test results across the population. We show this for [50% 50%] with $\rho_{12} = 0.2$ (blue), [50% 0%] with $\rho_{12} = 0$ (red), and [75% 50%] with $\rho_{12} = -0.2$ (yellow). We show how **C**: the Bayesian MAP estimate, and **D**: 96% HPDI width, of the estimated within-group prevalence difference $\gamma_1 - \gamma_2$ scale with the number of participants.

Prevalence as a function of effect size

In the above, we focussed on performing explicit statistical inference within each participant. A possible criticism of this approach is that the within-participant binarization of a continuous effect size can lose information. If the null distribution is the same for each participant, then the within-participant inference involves comparing each participant effect size, E_p, to a common statistical threshold $\hat{E}$. The prevalence estimation described above can therefore be interpreted as estimating the population prevalence of participants for which $E_p > \hat{E}$. In the NHST case, $\hat{E}$ is chosen so that $P(E > \hat{E}) = \alpha$, (usually $\alpha = 0.05$) but we can generally consider the prevalence of participants with effects exceeding any value $\hat{E}$. We therefore estimate the prevalence of $E_p > \hat{E}$ as a function of $\hat{E}$. This can reveal if a data set provides evidence for population prevalence of a sub-threshold within-participant effect, as well as showing how population prevalence decreases for larger effects. If desired, a frequentist approach to control the error rate can be applied by using the method of maximum statistics to correct over the multiple prevalence inferences. Figure 4 demonstrates this approach for the simulated systems of Figure 1, showing prevalence results for both right-sided, $E_p > \hat{E}$, and left-sided, $E_p < \hat{E}$, effect size thresholds. Note that this approach requires the null distribution to be the same for each participant, and requires the false positive rate α to be calculated for each effect size considered. It reveals everything we can learn about the population prevalence of different effect sizes from our dataset, exploring beyond the binarization of the within-participant inference.

Figure 4: *One-sided prevalence as a function of effect size.* We consider the same simulated systems shown in Figure 1, showing both right-tailed ($E_p > \hat{E}$) and left-tailed ($E_p < \hat{E}$) prevalence as a function of effect size. Orange lines show the effect size corresponding to the two-sided $\alpha = 0.05$ within-participant test, as used in Figure 1. Dashed lines show the effect size corresponding to the ground truth of the simulation. **A,B:** $\mu_{pop} = 0$, **C,D:** $\mu_{pop} = 1$. **A,C:** $k = 20$ trials, **B,D:** $k = 500$ trials. Black line shows MAP, shaded region shows 96% HPDI.

How to apply Bayesian prevalence in practice

As in the simulation of Figure 1, a typical population mean inference is often framed as a two-level summary statistic procedure. At the first level, the effect is quantified within each participant (e.g. a difference in mean response between two conditions). At the second level, the population mean is inferred under the assumption that the effect is normally distributed in the population (i.e. based on the mean and standard deviation of the measured effect across participants). Bayesian prevalence is similarly framed as a two-level procedure. At the first level, a statistical test is applied within each participant, the result of which can be binarized via a within-participant NHST (e.g. using a parametric test, as in our simulation, or alternatively using non-parametric permutation methods, but independently for each participant), or via an arbitrary effect size threshold $\hat{E}$. At the second level, the binary results from the first level (i.e. the counts of significant participants) are the input to the Bayesian population prevalence computation. To accompany this paper, we provide code in Matlab, Python and R to visualise the full posterior distribution of the population prevalence, as well as extract properties, such as the MAP point estimate and HPDIs. We also provide functions to provide Bayesian estimates of the difference in prevalence between two mutually exclusive participant groups to the same test (between-group prevalence difference), as well as the difference in prevalence between two different tests applied to a single sample of participants (within-group prevalence difference). We suggest reporting population prevalence inference results as the MAP estimate together with one or more HPDI's (e.g. with probability 0.5 or 0.96, see Methods).

It is important to stress that the second-level prevalence inference does not impose any requirements on the first level within-participant tests, other than that each test should have the same false positive rate of α (see Methods). It is not required, for example, that each participant have the same number of trials or degrees of freedom. The within-participant test can be parametric (e.g. a t-test) or non-parametric (e.g. based on permutation testing). It can be a single statistical test, or an inference over multiple tests (e.g. a neuroimaging effect within a certain brain region), provided that the family-wise error rate is controlled at α (e.g. by using permutation tests with the method of maximum statistics).

Discussion

The fields of psychology and neuroimaging are currently dominated by an approach in which the primary focus of statistical analysis is the population mean. Here, we propose a conceptual shift in perspective – from estimating population means to estimating the prevalence of effects detected within individuals. We provide a simple but novel Bayesian method to estimate the population prevalence of true positives from any experimental test. We propose that within-participant inference, combined with prevalence estimation, may in some cases better match a qualitative hypothesis regarding the existence of an effect within individuals, relative to a binary inferential result that the population mean differs from zero. Our method can easily be applied to almost any statistical evaluation of any experiment, provided a NHST can be performed at the individual participant level. The simulations presented here can also be used for simple power analyses when designing studies from this perspective.

Together, this conceptual shift and novel practical method support an alternative perspective for statistics in which the individual participant is the most relevant experimental unit to consider for the purposes of replication (Nachev et al., 2019; Smith and Little, 2018; Thiebaut de Schotten and Shallice, 2017). From this perspective, power should be calculated for effects within individual participants. This perspective gives a very different view of the strength of evidence provided by a dataset and of the importance of sample size (for both participants and trials) compared to the more common population mean perspective (Baker et al., 2019). For example, the simulation of 50 participants with 20 trials in Figure 1C has $p = 0.008$ for a group mean different from zero, a result that is as surprising, under the null hypothesis, as observing 7 heads in a row from tosses of a fair coin (Rafi and Greenland, 2020). This is weaker evidence than just 2 out of 5 participants showing an effect at $\alpha = 0.05$ ($p = 0.0012$ under the global null, or about as surprising as 10 heads in a row). 3 out of 5 significant participants corresponds to $p = 0.00003$ under the global null (as surprising as 15 heads in a row); this is substantially stronger evidence for a population-level effect than that provided by the population mean inference in Figure 1D (from 50 participants, even with 500 trials). However, in the current scientific climate, the weaker result obtained from the larger sample size would commonly be viewed as providing more satisfactory evidence by most readers and reviewers. We would like to highlight this pervasive misunderstanding – that larger participant numbers automatically imply better evidence at the population level. The crux of our argument is that most studies focus on the difference between panels A and C (a small difference in population mean, ignoring the implications of large between-participant variance in studies that are underpowered for within-participant effects). By contrast, moving towards the situation shown in Fig 1B and D (increased power within individual participants) would provide both improved replicability and greater insight into the size and prevalence of effects in the population. Ensuring that studies are sufficiently powered to obtain reliable effect size estimates within individual participants has several advantages. More accurate characterisation of within-participant effect sizes could improve studies relating individual differences in experimental effects to other external factors. Comparisons between groups can look beyond means to consider other differences in the distribution of effect sizes (Rousselet et al., 2017).

Bayesian prevalence produces a quantitative prevalence estimate with associated uncertainty, rather than reducing an entire experiment to a single, binary population NHST result. This addresses many of the problems noted with the NHST framework (Amrhein et al., 2019; McShane et al., 2019) and also reduces the risk of questionable research practices, such as *p*-hacking (Forstmeier et al., 2017). Each participant serves as a replication since the same statistical analysis is performed on separate sets of data. We have also shown how to estimate the prevalence of different effect size thresholds, which avoids focussing on a single within-participant dichotomisation. Importantly, estimated prevalence is only one approach to evaluating the strength of evidence provided by a dataset, and it should be assessed alongside the quality of the experimental design, within-participant power, and the effect sizes within individuals. Prevalence and population mean estimation are not mutually exclusive; ideally one could report within-participant effect sizes and estimated population prevalence together with estimation and inference of the population mean.

The practice of collecting enough data to perform within-participant inference is not a new idea – much of traditional psychophysics employs this approach (Smith and Little, 2018). We have employed this technique with EEG (Schyns et al., 2011; Smith et al., 2006) and MEG (Ince et al., 2015; Zhan et al., 2019), and in fMRI it is becoming more common to collect large quantities of data for fitting high-dimensional, cross-validated machine learning models within individual participants (Huth et al., 2016; Stansbury et al., 2013). Recently, this practice has also been adopted in the resting-state fMRI field where it is termed dense sampling or precision imaging (Braga and Buckner, 2017; Gordon et al., 2017; Laumann et al., 2015; Poldrack, 2017). One benefit of Bayesian prevalence is to obtain a population level inference from such studies, even when the number of participants is small.

Note that while similar points regarding within-participant inference have been made elsewhere (Nachev et al., 2019; Smith and Little, 2018; Thiebaut de Schotten and Shallice, 2017), such results are typically considered to form a case study, without an inferential link that allows generalization to the population (Neuroscience, 2020). The methods presented here address this concern by providing an inferential bridge from within-participant statistics to the population level. We have focussed on human participants in typical psychology or neuroimaging experiments, but the proposed methods can be applied to infer the population prevalence of effects of other types of experimental units. For example, our approach could be directly applied to identified single-unit neurons in electrophysiological recordings. Typically, only a subset of the identified neurons respond to a specific stimulus or experimental manipulation. Bayesian prevalence could be applied to estimate the proportion of neurons that respond in a particular way in the population of neurons in a particular brain region. Our between-group comparison method could be used to formally compare the population prevalence of a certain type of responsive neuron between different brain areas or between different species. Thus, although it is common in electrophysiology to have individual neurons as the replication unit and to perform inference at that level, the inferential bridge to the population that Bayesian prevalence provides offers a new perspective on the results of such studies.

The prevalence approach, however, is not without certain limitations. One criticism is that the demonstration of within-participant effects, which this approach requires, sets a much higher bar of evidence. It might be impractical to reach sufficient within-participant power in

some experimental designs. However, many statisticians have explicitly argued that the replicability crisis results from standards of evidence for publication being too weak (Benjamin et al., 2018). If so, the within-participant approach should lead to increased replicability. Indeed, it is common in neuroimaging studies to have no consistent pattern discernible in the effect size maps of individual participants, and yet such studies report a significant population mean effect in a focal region. In our view, this is problematic if our ultimate goal is to relate neuroimaging results to cognitive functions within individuals (Smith and Little, 2018). By contrast, as our simulations show (Figure 1B), strong evidence for a modulation can be evident in the absence of a population mean effect when the effect is heterogeneous across the population.

It is natural to expect considerable heterogeneity to exist across populations for a wide range of experimental tasks. In fact, the normal distribution that underlies most inferences on the population mean implies such heterogeneity (Figure 1B). It has recently been suggested that researchers should define a smallest effect size of interest (SESOI) (Lakens, 2017) and should consider this when calculating statistical power. We suggest that the SESOI should also be considered at the individual level, and explicitly related to the population variance obtained from the hierarchical mixed effects model. If the population variance is large, then this implies that individuals exist in the population who have effects that are larger than the individual SESOI. This possibility is almost always ignored in applications of such modelling, which usually focus only on inference on the population mean. In clinical studies of rare diseases, patient numbers are often limited, and heterogeneity can be higher among these patient populations relative to the healthy population. If an experiment is sufficiently powered within individual participants, then Bayesian prevalence provides a statistical statement about the population of patients with the disease, even from a small sample and without assuming a normal distribution for the population.

Furthermore, the common assumption that effect sizes in the population follow a normal distribution is strong, although seldom justified (Lakens et al., 2018). For example, information processing, decision making, risk taking, and other strategies vary within the healthy population and across clinical and sub-clinical groups (Nachev et al., 2019; Smith and Little, 2018). In neuroimaging studies, there are related issues of variability in anatomical and functional alignment. To address these issues, results are often smoothed within individuals before performing population mean inference. However, many new experimental developments, such as high-resolution 7T fMRI to image cortical layers and high-density intracranial recordings in humans, pose increasing difficulties in terms of aligning data across participants. A major advantage of the prevalence approach is that we can perform within-participant inference corrected for multiple comparisons, and then estimate the population prevalence of these effects without requiring them to be precisely aligned (in space, time frequency etc.) across participants. For example, one might report that 24/30 participants showed an EEG alpha band power effect between 500 ms and 1000 ms post stimulus, which implies a population prevalence MAP of 0.79 (96% HPDI [0.61 0.91]), without requiring these individual effects to occur at precisely the same time points in those 24 participants. Similarly, if within-participant inference is corrected for multiple comparisons, one can infer prevalence of, say, an effect in a certain layer of primary visual cortex, without requiring precise alignment of the anatomical location of the effect between participants.

In fMRI concerns have recently been raised with common statistical approaches, many of which might be ameliorated by adopting the prevalence perspective. For example, Botvinik-Nezer et al. (Botvinik-Nezer et al., 2020) reported large variability in the inferential results that were obtained when different groups of researchers tested the same hypotheses on the same dataset. However, they note broad agreement between the population-level effect size maps of different pipelines, suggesting it is the final binary population inference that is inconsistent. It would be interesting to compare the consistency of graded prevalence results based on within-participant inference between different analysis pipelines in a similar way. Elliott et al. (Elliott et al., 2020) considered the within-participant test-retest reliability of common fMRI designs. They noted that reliable population mean effects can arise from unreliable within-participant measurements, and that this is the case in many common experimental designs that focus on statistical power from the perspective of the population mean. They argue that such designs are therefore problematic for individual-differences research. Focussing on within-participant power, as we suggest here, may increase test-retest reliability and enable more robust individual-difference applications.

Of course, there are many cases where the population mean is indeed the primary interest; in such cases, estimating and inferring on the population mean using hierarchical models is the most appropriate analysis, when possible. Linear mixed-effect models can also be interrogated in different ways to investigate the question of prevalence, for example, by examining the variances of the random-slopes for each participant, or explicitly computing prevalence of different effect sizes from the normal distribution fit to the population. It is possible to extend Bayesian hierarchical models to explicitly account for different sub-groups of participants (Bartlema et al., 2014; Haaf and Rouder, 2019). However, these approaches currently are not widely adopted, cannot easily be applied to non-linear or high-dimensional analysis methods common in neuroimaging, and add both mathematical and computational complexity compared to the second-level Bayesian prevalence method we present here, which is straightforward to apply to any first-level, within-participant analysis. Further, for many complex computational techniques – from modelling learning behaviour and decision making, to neuroimaging analysis techniques such as Representational Similarity Analysis, or high-dimensional encoding models (Haxby et al., 2014) – it is currently not possible to employ a multi-level linear modelling approach down to the trial level, due to the complexity of the non-linear analysis functions and models employed. It is also important to note that if the effect of interest is an unsigned information measure such as mutual information (Ince et al., 2017) or classification accuracy (Haxby et al., 2014) then the t-test based inference on population mean does not actually provide a random-effects inference that generalises to the population, but is equivalent to rejection of the global null showing only that there exist some subjects with an effect, i.e. a fixed-effect analysis (Allefeld et al., 2016). In such cases, Bayesian prevalence estimates may provide a stronger and more descriptive population result than the population mean NHST.

Bayesian prevalence has a broad range of applicability – spanning dense sampling studies with high within-participant power, to more traditional sampling models (more participants, fewer trials, e.g. Figure 1C). It is applicable to any behavioural study, including detailed computational models of behaviour, provided that model comparison or inference on model parameters can be performed within individuals. In neuroimaging, Bayesian prevalence can be applied to any imaging modality (EEG, MEG, fMRI, intracranial EEG), to individual neurons

within a brain region (to infer the proportion of responsive neurons), and with any statistical approach, including non-linear and multivariate analysis methods. The crucial requirement is an adequately powered experiment to detect effects within individual participants (or other units of interest, e.g. neurons). We argue that ensuring experiments are adequately powered to detect effects within individuals would have a wide range of advantages, including strengthening the robustness and replicability of reported results.

Conclusions

While the problems that underlie the replication crisis are being increasingly recognised, there is currently no consensus as to alternative statistical approaches that are needed to address the underlying problems. Here, we propose that shifting our focus to quantifying and inferring effects within individuals addresses many of the pressing concerns recently highlighted in psychology and neuroimaging (Amrhein et al., 2019; Benjamin et al., 2018; Forstmeier et al., 2017; Ioannidis, 2005; McShane et al., 2019). We present a Bayesian approach to estimating population prevalence which is broadly applicable, as it places no assumptions on the within-participant tests nor on the distribution of effects in the population. Further, prevalence does not require a Bayesian treatment; frequentist inference approaches can be used instead. The crucial point is to shift our perspective to first evaluate effects within individual participants, who represent the natural replication unit for studies of human behaviour.

Methods

Simulations from hierarchical population model
The data shown in Figure 1 were simulated from the standard hierarchical model:

$$y_{ij} \sim N(\mu_i, \sigma_w^2)$$
$$\mu_i \sim N(\mu_{pop}, \sigma_b^2)$$

Where y_{ij} denotes the measurement made on the j^{th} trial (out of t) of the i^{th} participant (out of n). μ_i represents the mean of each individual participant, σ_w represents a common within-participant standard deviation over trials, σ_b represents the standard deviation between participants, and μ_{pop} represents the overall population mean. This can be written as

$$y_{ij} = \mu_{pop} + \eta_{ij} + \epsilon_i$$

where $\eta_{ij} \sim N(0, \sigma_w^2)$, and $\epsilon_i \sim N(0, \sigma_b^2)$. Note that under this model, the distribution of the within participant means is $N(\mu_{pop}, \sigma_b^2 + \frac{1}{t}\sigma_w^2)$.

Binomial model of population prevalence of the ground truth state of sampled units
We consider a population of experimental units (for example, human participants, or individual neurons) which are of two types: those that have a particular binary effect or property, and those that do not. We consider the population prevalence of the ground trust state of each unit γ_{gt}, which is the proportion of the population from which the sample was drawn that have the effect ($0 < \gamma_{gt} < 1$). If the true status of each individual unit could be directly observed, then the sample could be modelled with a binomial distribution with probability parameter γ_{gt}. However, we cannot directly observe the true status of each unit. Instead, we apply to each unit a statistical test following the NHST framework, which outputs a binary result (which we term here positive vs negative). This test has a false positive rate α, and sensitivity β. Thus, the probability that a randomly selected unit from the population that does not possess the defined effect but will produce a positive test result is α, while the probability that a randomly selected unit that does possess the defined effect will produce a positive test result is β. Under the assumption that the units are independent and α and β are constant across units, the number of positive tests k in a sample of size n can be modelled as a binomial distribution with parameter θ (Donhauser et al., 2018; Friston et al., 1999a, 1999b; Rogan and Gladen, 1978):

$$P(X = k|\theta) = \binom{n}{k}\theta^k(1-\theta)^{n-k}$$
$$\theta = (1 - \gamma_{gt})\alpha + \gamma_{gt}\beta$$

Binomial model of population prevalence of true positives for a given test procedure
A major issue with the above approach is that it requires the sensitivity β to be specified and constant across units. β is the probability of a significant result given that the individual has an effect, and differs as a function of the ground truth effect size. In general, we do not assume that all members of the population have the same effect size, so it is not possible to specify a common β for all individuals.

Therefore, rather than modelling the prevalence of ground truth effects, we consider estimation of the prevalence of true positive test results for the particular test procedure we employ, γ_{tp}:

$$P(X = k|\theta) = \binom{n}{k}\theta^k(1-\theta)^{n-k}$$
$$\theta = (1 - \gamma_{tp})\alpha + \gamma_{tp}$$

In this case, the only assumption is that test procedure has the same false positive rate α for every individual, which is easily satisfied by most common parametric and non-parametric statistical tests. Note that this is equivalent to estimating ground truth prevalence with a test with $\beta = 1$. In general, a test with lower sensitivity allows inference of a higher prevalence for an observed k, because some of the observed negative results will be missed true-positive results. Therefore, prevalence of true positives obtained with $\beta = 1$ is a conservative lower bound on the prevalence of ground truth state (Allefeld et al., 2016; Donhauser et al., 2018; Friston et al., 1999b).

Throughout this paper we consider the prevalence of true positives for a given test procedure, rather than the prevalence of ground truth state and so omit the subscript tp. This quantifies the proportion of the population that would be expected to provide a true positive test result (i.e. have a non-null effect that would be detected by the experimental test procedure considered). It is simply not possible to infer anything at the population level about effects too weak to detect with the performed experiment.

Frequentist estimation of and inference on population prevalence

Various frequentist approaches can be used with the above binomial model of statistical testing. First, the maximum likelihood of the population prevalence of true positives can be obtained as:

$$\hat{\gamma} = \frac{k/n - \alpha}{1 - \alpha}$$

Standard bootstrap techniques (Efron and Tibshirani, 1994) can give percentile bootstrap confidence intervals as an indication of uncertainty in this estimate. We can also explicitly test various null hypotheses at the population level. For example, we can test a compound null hypothesis $\gamma < 0.5$, termed the majority null (Allefeld et al., 2016; Donhauser et al., 2018). This is chosen with the idea that a prevalence of >50% supports a claim that the effect is *typical* in the population. Other explicit compound nulls of this form can also be tested (e.g. that $\gamma < 0.25$ or $\gamma < 0.75$). Alternatively, it is possible to infer a lower bound on the population prevalence, by finding the largest value γ_c, such that $p(X > k \mid \gamma < \gamma_c) < 0.05$ (Allefeld et al., 2016; Donhauser et al., 2018). This inferred lower bound provides a more graded output than a binary significance result of testing against a specific compound null (i.e. the continuous value γ_c).

Bayesian estimation of population prevalence

We apply standard Bayesian techniques to estimate the population prevalence parameter of this model (Gelman, 2014). Assuming a Beta prior distribution for θ with shape parameters r, s, together with a Binomial likelihood function, the posterior distribution for θ is given by a Beta distribution with parameters $(k + r, n - k + s)$, truncated to the interval $[\alpha, 1]$, where k is the number of participants showing an above-threshold effect out of n tested. In the

examples shown here, we use a uniform prior (beta with shape parameters $r = s = 1$), as in the general case, there is no prior information regarding θ. This implies a uniform prior also for γ, so, a priori, we consider any value of population prevalence equally likely. While we default to the uniform prior, the code supports any beta distribution as a prior. Alternative priors could be implemented via Markov chain Monte Carlo methods (Gelman, 2014) together with the models described here. Note that similar Bayesian approaches have been applied in the field of epidemiology, where sometimes multiple complementary diagnostic tests for a disease are applied with or without a gold standard diagnosis in a subset of the sampled units (Berkvens et al., 2006; Enøe et al., 2000; Joseph et al., 1995).

Under the uniform prior, the Bayesian maximum a posteriori (MAP) estimate for prevalence proportion of true positives, γ, is available analytically and is equivalent to the maximum likelihood estimate:

$$\gamma_{map} = \frac{k/n - \alpha}{1 - \alpha}$$

Following (McElreath, 2016), we present 96% Highest Posterior Density Intervals (HPDIs) here to emphasise the arbitrary nature of this value and to reduce the temptation to interpret the interval as a frequentist p=0.05 inference.

Bayesian estimation of the prevalence difference between two independent groups

We consider here a situation where the same test is applied to units sampled from two different populations. In addition to the prevalence of true positives within each population, we wish to directly estimate the difference in prevalence between the two populations. We denote the prevalence within each population as γ_1, γ_2, respectively. We sample n_1, n_2 participants at random from each population and record k_1, k_2, the number of significant within-participant tests in each sample. Assuming independent uniform priors on the prevalences, and associated θ_i variables as above with:

$$\theta_i = (1 - \gamma_i)\alpha + \gamma_i$$

then the posterior distribution for (θ_1, θ_2) is given by the product of two truncated beta distributions, with parameters $(k_i + 1, n_i - k_i + 1)$ respectively, both truncated to the interval $[\alpha, 1]$. The prevalence difference can be obtained as:

$$\gamma_1 - \gamma_2 = (\theta_1 - \theta_2)/(1 - \alpha)$$

For non-truncated beta distributions, an analytic exact result is available (Pham-Gia and Turkkan, 1993). This result could be extended to provide an exact distribution for the prevalence difference, but the mathematical expressions involved are quite complex. We find it simpler to employ Monte Carlo methods, which can provide as close an approximation to the exact answer as desired. Here we use Monte Carlo methods to draw samples from the posterior for (θ_1, θ_2), obtaining samples for the prevalence difference with the above expression. We use these samples to numerically compute the MAP and HPDIs.

Bayesian estimation of the prevalence difference of two different tests within the same sample of participants

In this situation, we consider that two different test procedures are applied to a single sample of n units. We assume both tests have the same value of α, and define:

$$\theta_i = (1 - \gamma_i)\alpha + \gamma_i = \alpha + (1 - \alpha)\gamma_i$$

Here θ_i is the probability that a randomly selected unit from the population will show a positive result on the i^{th} test, and γ_i is the population prevalence of true positives associated with the i^{th} test. Now, each unit provides one of four mutually exclusive results based on the combination of binary results from the two tests. k_{11} represents the number of units that have a positive result in both tests, k_{10}, k_{01} represent the number of units that have a positive result only in the first or second test, respectively, and k_{00} is the number of units that do not show a positive result in either test. So $\sum_{i,j} k_{ij} = n$. We can define analogous variables $\boldsymbol{\theta} = \{\theta_{ij}\}$, representing the population proportion for each of the four combined test outcomes. Note that $\theta_{ij} > 0$ and $\sum_{i,j} \theta_{ij} = 1$. The marginal success probabilities θ_i can be expressed as:

$$\theta_1 = \theta_{11} + \theta_{10}, \quad \theta_2 = \theta_{11} + \theta_{01}$$

and so

$$\gamma_1 - \gamma_2 = (\theta_{10} - \theta_{01})/(1 - \alpha)$$

The marginal probabilities θ_i are subject to the constraints

$$\alpha < \theta_i < 1$$

and so

$$\alpha < \theta_{11} + \theta_{10} < 1, \quad \alpha < \theta_{11} + \theta_{01} < 1$$

Assuming a uniform prior, and a multinomial distribution for the k_{ij}, the posterior of $\boldsymbol{\theta}$ is a truncated Dirichlet distribution with parameters $m_{ij} = k_{ij} + 1$ subject to the constraints above (which are analogous to the truncation of the beta posterior distribution in the case of a single test). We use a Monte Carlo approach to draw samples from this posterior following a modified stick-breaking process.

- Draw a sample θ_{11} from a $Beta(m_{11}, m_{10} + m_{01} + m_{00})$ distribution truncated to the interval $[0,1]$.
- Draw a sample z_{10} from a $Beta(m_{10}, m_{01} + m_{00})$ distribution truncated to the interval $\left[\max\left(\frac{\alpha - \theta_{11}}{1 - \theta_{11}}, 0\right), \frac{1 - \theta_{11}}{1 - \theta_{11}}\right]$. Set $\theta_{10} = (1 - \theta_{11})z_{10}$.
- Draw a sample z_{01} from a $Beta(m_{01}, m_{00})$ distribution truncated to the interval $\left[\max\left(\frac{\alpha - \theta_{11}}{1 - \theta_{11} - \theta_{10}}, 0\right), \min\left(\frac{1 - \theta_{11}}{1 - \theta_{11} - \theta_{10}}, 1\right)\right]$. Set $\theta_{01} = (1 - \theta_{11} - \theta_{10})z_{01}$.
- Set $\theta_{00} = 1 - \theta_{11} - \theta_{10} - \theta_{01}$.
- Then $(\theta_{11}, \theta_{01}, \theta_{10}, \theta_{00})$ is a draw from the required truncated Dirichlet distribution, and $(\theta_{10} - \theta_{01})/(1 - \alpha)$ is a draw from the posterior distribution of the prevalence difference.

We use these samples to numerically compute properties like the MAP estimate and HPDIs.

To specify a ground truth to simulate data from two tests applied to the same participants (Figure 3), we require γ_1 and γ_2, the population prevalences of the two tested effects, together with ρ_{12}, the correlation between the presence of the two effects across the population. From this we can calculate γ_{11}, the proportion of true positive results to both tests as:

$$\gamma_{11} = \gamma_1\gamma_2 + \rho_{12}\sqrt{\gamma_1(1-\gamma_1)\gamma_2(1-\gamma_2)}$$

Similarly, we can define γ_{ij} representing the population proportions corresponding to the other test result configurations. Then we can generate multinomial samples using the parameters θ_{ij} computed as:

$$\theta_{11} = \gamma_{11} + \alpha\gamma_{10} + \alpha\gamma_{01} + \alpha^2\gamma_{00}$$
$$\theta_{10} = \alpha + (1-\alpha)\gamma_1 - \theta_{11}$$
$$\theta_{01} = \alpha + (1-\alpha)\gamma_2 - \theta_{11}$$
$$\theta_{00} = 1 - \theta_{11} - \theta_{10} - \theta_{01}$$

Further mathematical details of the Bayesian estimation of prevalence and prevalence differences are given in supplemental notes available with the code at https://github.com/robince/bayesian-prevalence.

Prevalence as a function of effect size threshold

Estimating the prevalence of $E_p > \hat{E}$ proceeds as for prevalence inference based on within-participant NHST. One additional step is the need to calculate α, the false positive rate under the null hypothesis of no effect, for each threshold value $\hat{E}$. This is simply the probability of $E_p > \hat{E}$ under the null hypothesis. In the examples shown here, we calculate this from the cumulative distribution function of the appropriate t-distribution, but for other tests this could also be estimated non-parametrically. A number of $\hat{E}$ values are selected, linearly spaced over the observed range of the sample. For each of these values the count of the number of participants satisfying the inequality, and the α value corresponding to the inequality, are used to obtain the Bayesian posterior for prevalence of true positives. Note that this can be applied to either tail $E_p > \hat{E}$, or $E_p < \hat{E}$.

Data Availability

Scripts implementing the simulations and creating the figures from the manuscript are available at https://github.com/robince/bayesian-prevalence

Code Availability

To accompany this paper, we provide functions in Matlab, Python and R to calculate the Bayesian prevalence posterior density (e.g. to plot the full posterior distribution), the MAP estimate of the population prevalence, HPDI intervals of the posterior and lower bound quantiles of the posterior, as well as prevalence differences between samples or between tests within a sample. We also provide example scripts that produce posterior plots as in Figure 1E. See https://github.com/robince/bayesian-prevalence

Acknowledgements

We thank Carsten Allefeld for useful discussions. RAAI was supported by the Wellcome Trust [214120/Z/18/Z]. PGS was supported by EPSRC [MURI 1720461] and the Wellcome Trust [107802]. PGS is a Royal Society Wolfson Fellow [RSWF\R3\183002].

References

Allefeld, C., Görgen, K., and Haynes, J.-D. (2016). Valid population inference for information-based imaging: From the second-level t-test to prevalence inference. NeuroImage *141*, 378–392.

Amrhein, V., Greenland, S., and McShane, B. (2019). Retire statistical significance. Nature *567*, 305–307.

Angrist, J.D., and Pischke, J.-S. (2014). Mastering 'Metrics: The Path from Cause to Effect (Princeton ; Oxford: Princeton University Press).

Baker, D.H., Vilidaite, G., Lygo, F.A., Smith, A.K., Flack, T.R., Gouws, A.D., and Andrews, T.J. (2019). Power contours: optimising sample size and precision in experimental psychology and human neuroscience. ArXiv:1902.06122 [q-Bio, Stat].

Bartlema, A., Lee, M., Wetzels, R., and Vanpaemel, W. (2014). A Bayesian hierarchical mixture approach to individual differences: Case studies in selective attention and representation in category learning. Journal of Mathematical Psychology *59*, 132–150.

Benjamin, D.J., Berger, J.O., Johannesson, M., Nosek, B.A., Wagenmakers, E.-J., Berk, R., Bollen, K.A., Brembs, B., Brown, L., Camerer, C., et al. (2018). Redefine statistical significance. Nature Human Behaviour *2*, 6.

Berkvens, D., Speybroeck, N., Praet, N., Adel, A., and Lesaffre, E. (2006). Estimating Disease Prevalence in a Bayesian Framework Using Probabilistic Constraints: Epidemiology *17*, 145–153.

Botvinik-Nezer, R., Holzmeister, F., Camerer, C.F., Dreber, A., Huber, J., Johannesson, M., Kirchler, M., Iwanir, R., Mumford, J.A., Adcock, R.A., et al. (2020). Variability in the analysis of a single neuroimaging dataset by many teams. Nature 1–7.

Braga, R.M., and Buckner, R.L. (2017). Parallel Interdigitated Distributed Networks within the Individual Estimated by Intrinsic Functional Connectivity. Neuron *95*, 457-471.e5.

Donhauser, P.W., Florin, E., and Baillet, S. (2018). Imaging of neural oscillations with embedded inferential and group prevalence statistics. PLOS Computational Biology *14*, e1005990.

Efron, B., and Tibshirani, R.J. (1994). An introduction to the bootstrap (CRC press).

Elliott, M.L., Knodt, A.R., Ireland, D., Morris, M.L., Poulton, R., Ramrakha, S., Sison, M.L., Moffitt, T.E., Caspi, A., and Hariri, A.R. (2020). What Is the Test-Retest Reliability of Common Task-Functional MRI Measures? New Empirical Evidence and a Meta-Analysis: Psychological Science.

Enøe, C., Georgiadis, M.P., and Johnson, W.O. (2000). Estimation of sensitivity and specificity of diagnostic tests and disease prevalence when the true disease state is unknown. Preventive Veterinary Medicine *45*, 61–81.

Forstmeier, W., Wagenmakers, E.-J., and Parker, T.H. (2017). Detecting and avoiding likely false-positive findings – a practical guide. Biological Reviews *92*, 1941–1968.

Friston, K.J. (2007). Statistical parametric mapping: the analysis of functional brain images (Academic Press).

Friston, K.J., Holmes, A.P., and Worsley, K.J. (1999a). How Many Subjects Constitute a Study? NeuroImage *10*, 1–5.

Friston, K.J., Holmes, A.P., Price, C.J., Büchel, C., and Worsley, K.J. (1999b). Multisubject fMRI Studies and Conjunction Analyses. NeuroImage *10*, 385–396.

Gelman, A. (2014). Bayesian data analysis (Boca Raton: CRC Press).

Gordon, E.M., Laumann, T.O., Gilmore, A.W., Newbold, D.J., Greene, D.J., Berg, J.J., Ortega, M., Hoyt-Drazen, C., Gratton, C., Sun, H., et al. (2017). Precision Functional Mapping of Individual Human Brains. Neuron *95*, 791-807.e7.

Haaf, J.M., and Rouder, J.N. (2019). Some do and some don't? Accounting for variability of individual difference structures. Psychon Bull Rev *26*, 772–789.

Haxby, J.V., Connolly, A.C., and Guntupalli, J.S. (2014). Decoding Neural Representational Spaces Using Multivariate Pattern Analysis. Annual Review of Neuroscience *37*, 435–456.

Holmes, A., and Friston, K. (1998). Generalisability, random effects and population inference. Neuroimage *7*.

Huth, A.G., de Heer, W.A., Griffiths, T.L., Theunissen, F.E., and Gallant, J.L. (2016). Natural speech reveals the semantic maps that tile human cerebral cortex. Nature *532*, 453–458.

Ince, R.A.A., van Rijsbergen, N., Thut, G., Rousselet, G.A., Gross, J., Panzeri, S., and Schyns, P.G. (2015). Tracing the Flow of Perceptual Features in an Algorithmic Brain Network. Scientific Reports *5*, 17681.

Ince, R.A.A., Giordano, B.L., Kayser, C., Rousselet, G.A., Gross, J., and Schyns, P.G. (2017). A statistical framework for neuroimaging data analysis based on mutual information estimated via a gaussian copula. Hum. Brain Mapp. *38*, 1541–1573.

Ioannidis, J.P.A. (2005). Why Most Published Research Findings Are False. PLOS Medicine *2*, e124.

Joseph, L., Gyorkos, T.W., and Coupal, L. (1995). Bayesian Estimation of Disease Prevalence and the Parameters of Diagnostic Tests in the Absence of a Gold Standard. Am J Epidemiol *141*, 263–272.

Lakens, D. (2017). Equivalence Tests: A Practical Primer for t Tests, Correlations, and Meta-Analyses. Social Psychological and Personality Science *8*, 355–362.

Lakens, D., Adolfi, F.G., Albers, C.J., Anvari, F., Apps, M.A.J., Argamon, S.E., Baguley, T., Becker, R.B., Benning, S.D., Bradford, D.E., et al. (2018). Justify your alpha. Nature Human Behaviour *2*, 168.

Laumann, T.O., Gordon, E.M., Adeyemo, B., Snyder, A.Z., Joo, S.J., Chen, M.-Y., Gilmore, A.W., McDermott, K.B., Nelson, S.M., Dosenbach, N.U.F., et al. (2015). Functional System and Areal Organization of a Highly Sampled Individual Human Brain. Neuron *87*, 657–670.

Little, D.R., and Smith, P.L. (2018). Replication is already mainstream: Lessons from small-N designs. Behavioral and Brain Sciences *41*.

McElreath, R. (2016). Statistical rethinking: a Bayesian course with examples in R and Stan (Boca Raton: CRC Press/Taylor & Francis Group).

McShane, B.B., Gal, D., Gelman, A., Robert, C., and Tackett, J.L. (2019). Abandon Statistical Significance. The American Statistician *73*, 235–245.

Nachev, P., Rees, G., and Frackowiak, R. (2019). Lost in translation. F1000Research *7*, 620.

Neuroscience, S. for (2020). Consideration of Sample Size in Neuroscience Studies. J. Neurosci. *40*, 4076–4077.

Nichols, T., Brett, M., Andersson, J., Wager, T., and Poline, J.-B. (2005). Valid conjunction inference with the minimum statistic. NeuroImage *25*, 653–660.

Pearl, J. (2009). Causality (Cambridge: Cambridge University Press).

Penny, W., and Holmes, A. (2007). Random effects analysis. Statistical Parametric Mapping: The Analysis of Functional Brain Images 156–165.

Penny, W., Holmes, A., and Friston, K. (2003). Random effects analysis. Human Brain Function 843–850.

Pham-Gia, T., and Turkkan, N. (1993). Bayesian analysis of the difference of two proportions. Communications in Statistics - Theory and Methods *22*, 1755–1771.

Poldrack, R.A. (2017). Precision Neuroscience: Dense Sampling of Individual Brains. Neuron *95*, 727–729.

Rafi, Z., and Greenland, S. (2020). Semantic and Cognitive Tools to Aid Statistical Science: Replace Confidence and Significance by Compatibility and Surprise. ArXiv:1909.08579 [q-Bio, Stat].

Rogan, W.J., and Gladen, B. (1978). ESTIMATING PREVALENCE FROM THE RESULTS OF A SCREENING TEST. Am J Epidemiol *107*, 71–76.

Rosenblatt, J.D., Vink, M., and Benjamini, Y. (2014). Revisiting multi-subject random effects in fMRI: Advocating prevalence estimation. NeuroImage *84*, 113–121.

Rousselet, G.A., Pernet, C.R., and Wilcox, R.R. (2017). Beyond differences in means: robust graphical methods to compare two groups in neuroscience. European Journal of Neuroscience *46*, 1738–1748.

Schyns, P.G., Thut, G., and Gross, J. (2011). Cracking the Code of Oscillatory Activity. PLoS Biol *9*, e1001064.

Smith, P.L., and Little, D.R. (2018). Small is beautiful: In defense of the small-N design. Psychon Bull Rev *25*, 2083–2101.

Smith, M.L., Gosselin, F., and Schyns, P.G. (2006). Perceptual Moments of Conscious Visual Experience Inferred from Oscillatory Brain Activity. PNAS *103*, 5626–5631.

Stansbury, D.E., Naselaris, T., and Gallant, J.L. (2013). Natural Scene Statistics Account for the Representation of Scene Categories in Human Visual Cortex. Neuron *79*, 1025–1034.

Thiebaut de Schotten, M., and Shallice, T. (2017). Identical, similar or different? Is a single brain model sufficient? Cortex *86*, 172–175.

Yarkoni, T. (2019). The Generalizability Crisis (PsyArXiv).

Zhan, J., Ince, R.A.A., van Rijsbergen, N., and Schyns, P.G. (2019). Dynamic Construction of Reduced Representations in the Brain for Perceptual Decision Behavior. Current Biology *29*, 319-326.e4.