The maintenance of polygenic sex determination depends on the dominance of fitness

effects which are predictive of the role of sexual antagonism

Richard P. Meisel

Department of Biology and Biochemistry, University of Houston, Houston, TX 77204

Running title: Polygenic sex determination

Keywords: sex chromosomes, house fly, sexual conflict, recessive, additive, overdominant

Corresponding author:

Richard P. Meisel

Science & Research Building 2

3455 Cullen Blvd., Suite 342

Houston, Texas 77204-5001

1-713-743-3607

rpmeisel@uh.edu

Acknowledgments

This work was funded by the National Science Foundation (DEB-1845686). This manuscript

was improved by comments from members of the Meisel lab, editors, and multiple anonymous

reviewers.

Abstract

In species with polygenic sex determination, multiple male- and/or female-determining loci on different proto-sex chromosomes segregate as polymorphisms within populations. The extent to which these polymorphisms are at stable equilibria is not yet resolved. Previous work demonstrated that polygenic sex determination is most likely to be maintained as a stable polymorphism when the proto-sex chromosomes have opposite (sexually antagonistic) fitness effects in males and females. However, these models usually consider polygenic sex determination systems with only two proto-sex chromosomes, or they do not broadly consider the dominance of the alleles under selection. To address these shortcomings, I used forward population genetic simulations to identify selection pressures that can maintain polygenic sex determination under different dominance scenarios in a system with more than two proto-sex chromosomes (modeled after the house fly). I found that overdominant fitness effects of male-determining proto-Y chromosomes are more likely to maintain polygenic sex determination than dominant, recessive, or additive fitness effects. Dominance also affects the predicted role of sexual antagonism in maintaining polymorphism. Dominant fitness effects that maintain polygenic sex determiantion, for example, are predicted to have sexually antagonistic multi-chromosomal genotypes. However, the individual proto-sex chromosomes are not expected to have sexually antagonstic effects, demonstrating that sexual antagonism can be an emergent property of the multi-chromosome genotype. Overdominant fitness effects, in contrast, are expected to maintain polygenic sex determination with proto-Y chromosomes that have sexually antagonstic effects even though the multi-chromosome genotypes do not have signatures of sexual antagonism, My results therefore demonstrate that the dominance of fitness effects has consequences for both the likelihood that polygenic sex determination will be

41   maintained as well as the role sexually antagonistic selection is expected to play in its

42   maintenance.

Introduction

Sex determination is the developmental process by which a genetic signal or environmental cue initiates sexually dimorphic gene regulatory pathways to produce phenotypically different males and females (Bachtrog *et al.* 2014). The master regulators of sex determination evolve fast, often differing between closely related species and sometimes even polymorphic within populations (Bull 1983; Wilkins 1995; Beukeboom and Perrin 2014). Master regulators can be found on sex chromosomes, and the evolutionary transitions of sex determiners can drive turnover of the sex chromosomes (Abbott *et al.* 2017). These evolutionary transitions include a period of polygenic sex determination (PSD) during which multiple master sex determining loci on different chromosomes segregate within natural populations (Moore and Roberts 2013). Understanding the population dynamics of PSD is informative of the factors responsible for the evolutionary divergence of sex determination pathways and sex chromosomes.

PSD has been observed in many animal and plant species (Moore and Roberts 2013; Beukeboom and Perrin 2014; Bachtrog *et al.* 2014). Most population genetic models predict that PSD will be an unstable intermediate between monogenic equilibria (Rice 1986; van Doorn and Kirkpatrick 2007). Models that do allow for stable PSD often predict that opposing selection pressures in males and females (i.e., sexually antagonistic selection) may be important for maintaining the polymorphism (Bull 1983; Rice 1986; van Doorn and Kirkpatrick 2007; Meisel *et al.* 2016). It is also possible that variable selection pressures across heterogeneous environments could maintain PSD (Bateman and Anholt 2017).

Most previous theoretical treatments of the selection pressures that maintain PSD considered specific cases with a small number of sex chromosomes segregating within a population. For example, in the platyfish, *Xiphophorus maculatus*, a single chromosome can be

an X, Y, or Z (Kallman 1973). Orzack et al. (1980) showed that this polymorphism is maintained

because heterogametic females (WY or WX) and males (XY) have higher fitness than their

homogametic counterparts (XX females or YY males). In addition, van Doorn and Kirkpatrick

(2007, 2010) found that sexually antagonistic selection could maintain a polymorphism in which

one chromosome segregates as an XY pair and another chromosome is either an XY or ZW

pair. However, these models do not capture the full diversity of PSD, which can include more

than two XY or ZW pairs segregating within a single population (Moore and Roberts 2013).

Previous models also did not broadly consider how the maintenance of PSD depends on

the dominance of fitness effects. The dominance of sexually antagonistic alleles affects their

ability to segregate as stable polymorphisms (Kidwell *et al.* 1977; Connallon and Chenoweth

2019), suggesting that dominance may be important for the maintenance of PSD under sexually

antagonistic selection. Dominance also differentially affects the evolutionary fate of X-linked and

autosomal alleles that experience sex-specific selection pressures (Rice 1984; Charlesworth *et*

al. 1987; Fry 2010; Connallon *et al.* 2012). In a species with PSD, there are proto-sex

chromosomes that behave neither like autosomes nor conventional heteromorphic sex

chromosomes; proto-sex chromosomes are diploid in both sexes (like autosomes) but have

sex-biased modes of inheritance (like conventional sex chromosomes). This is superficially

similar to freely recombining regions on sex chromosomes (Otto *et al.* 2011), except that in PSD

a proto-Y (or proto-W) chromosome can be directly carried by females (males) without X-Y (or

Z-W) recombination. Therefore, the expected effects of dominance and sexual antagonism on

the evolution of PSD cannot be determined from existing models of autosomal or sex-linked

alleles.

The house fly, *Musca domestica*, is a well-suited organism for investigating the

maintenance of complex PSD because it has a highly polymorphic sex determination system

(Hamm *et al.* 2015). The *M. domestica male determiner* (*Mdmd*) can be found on at least four of

the six house fly chromosomes in natural populations (Sharma *et al.* 2017). *Mdmd* causes the

house fly ortholog of *transformer* (*Md-tra*) to be spliced into a non-functional isoform, which

leads to male-specific splicing and expression of downstream target genes, thereby producing

fertile males (Hediger *et al.* 2010). In the absence of *Mdmd*, *Md-tra* is spliced into a

female-determining transcript that regulates the splicing of downstream targets, which promote

the development of female morphological and behavioral traits (Hediger *et al.* 2004; Meier *et al.*

2013).

There are two common male-determining proto-Y chromosomes and one

female-determining proto-W chromosome in house fly populations. *Mdmd* is most commonly

found on the Y (Y^M) and third (III^M) chromosomes in natural populations (Hamm *et al.* 2015).

Both Y^M and III^M are young proto-Y chromosomes that arose recently and are nearly identical in

gene content to their homologous proto-X chromosomes, known as X and III (Meisel *et al.*

2017). In addition, there is a dominant allele of *Md-tra* that is found in many house fly

populations (*Md-tra^D*), which is resistant to *Mdmd* and causes female development regardless of

whether there are copies of *Mdmd* in the genotype (McDonald *et al.* 1978; Hediger *et al.* 2010).

Md-tra is found on the fourth chromosome, and thus fourth chromosomes carrying *Md-tra^D*

(which I will refer to as IV^F) are W chromosomes. All sequences of *Md-tra^D* from around the

world are identical (Scott *et al.* 2014), suggesting that *Md-tra^D* arose recently and IV^F is a young

proto-W chromosome. In populations where Y^M, III^M, and *Md-tra^D* all segregate, there are 18

possible sex chromosome genotypes (Table 1).

There is evidence that natural selection maintains PSD in house fly populations. First,

the proto-Y chromosomes have remained at stable frequencies within house fly populations

over decades (Kozielska *et al.* 2008; Meisel *et al.* 2016). Second, Y^M and III^M are distributed

along north-south clines on multiple continents (Denholm *et al.* 1986; Tomita and Wada 1989;

Hamm *et al.* 2005; Kozielska *et al.* 2008), and temperature is the best predictor of their

frequencies (Feldmeyer *et al.* 2008). This suggests that heterogenous (temperature-dependent)

selection pressures maintain the Y^M and III^M chromosomes across populations. The stable

frequencies within populations, yet divergent frequencies across populations, suggest that there

are specific fitness effects associated with the house fly proto-Y chromosomes within each

population. Third, males carrying multiple proto-Y chromosomes (e.g., both Y^M and III^M, or two

copies of III^M) can be found in some populations (e.g., Hamm and Scott 2009). The frequency of

Md-tra^D is positively correlated with the frequency of these multi-Y males across populations,

suggesting that selection for balanced sex ratios may further maintain PSD (Meisel *et al.* 2016).

The rich body of observational data in house fly makes it a well-suited system around

which to develop population genetic models to determine how PSD can be maintained in natural

populations. For example, Kozielska et al. (2006) showed that, even in the complex house fly

PSD system, sex ratios are not expected to deviate substantially from 1:1 male:female. In

addition, despite no experimental evidence for sexually antagonistic effects of any house fly

proto-sex chromosomes, another population genetic model demonstrated that the frequencies

of the proto-sex chromosomes in natural populations are consistent with sexual antagonism

(Meisel *et al.* 2016). However, that model assumed that the proto-Y chromosomes (Y^M and III^M)

have additive fitness effects, which may not be true if, for example, beneficial mutations are

recessive or dominant (Orr 2010). In addition, Y chromosomes are expected to carry recessive

deleterious alleles (Charlesworth and Charlesworth 2000), which can further violate the

assumption of additive fitness effects. To address the shortcomings of previous models, I used

population genetic simulations to investigate whether sex chromosomes with non-additive

fitness effects can maintain the complex house fly PSD system. I specifically tested if these

selection pressures can produce proto-sex chromosome frequencies similar to those observed

in natural populations of house fly, and I determined the conditions under which sexually

antagonistic effects of those proto-sex chromosomes are expected.

Table 1. House fly sex chromosome genotypes.

	genotype			
	X or Y^M	III or III^M	IV or IV^F	sex
f_1	X/X	III/III	IV/IV	female
f_2	X/X	III/III	IV/IV^F	female
f_3	X/X	III/III^M	IV/IV^F	female
f_4	X/X	III^M/III^M	IV/IV^F	female
f_5	X/Y^M	III/III	IV/IV^F	female
f_6	X/Y^M	III/III^M	IV/IV^F	female
f_7	X/Y^M	III^M/III^M	IV/IV^F	female
f_8	Y^M/Y^M	III/III	IV/IV^F	female
f_9	Y^M/Y^M	III/III^M	IV/IV^F	female
f_{10}	Y^M/Y^M	III^M/III^M	IV/IV^F	female
m_1	X/X	III/III^M	IV/IV	male
m_2	X/X	III^M/III^M	IV/IV	male
m_3	X/Y^M	III/III	IV/IV	male
m_4	X/Y^M	III/III^M	IV/IV	male
m_5	X/Y^M	III^M/III^M	IV/IV	male
m_6	Y^M/Y^M	III/III	IV/IV	male
m_7	Y^M/Y^M	III/III^M	IV/IV	male
m_8	Y^M/Y^M	III^M/III^M	IV/IV	male

Materials and Methods

General overview

I used a simulation approach to identify fitness effects of two proto-Y chromosomes (Y^M and III^M) and one proto-W chromosome (IV^F) that maintain PSD under four different dominance scenarios. To do so, I first randomly assigned fitness values to each of the single chromosome genotypes in males and females separately (Figure 1A), assuming either additive, dominant, recessive, or overdominant (i.e., heterozygote advantage) fitness effects of the proto-Y chromosomes (Y^M and III^M). I then calculated the fitness of each of the 18 multi-chromosome genotypes based on the single chromosome fitness effects (Table 1). I next determined the frequency of each genotype and proto-sex chromosome after 1,000 generations of random mating with the assigned fitness values (Figure 1A). I did this using simulations for one million different fitness values and four possible starting genotype frequencies for each dominance model. I retained fitness effects in which all three proto-sex chromosomes remained polymorphic after 1,000 generations, and from those I selected the 1,000 fitness effects that produced proto-sex frequencies closest to those observed in natural populations (Figure 1A). In doing so, I am essentially applying an approximate Bayesian computation approach to fit selection pressures to natural populations (Beaumont *et al.* 2002). I also evaluated how well those fitness effects maintain PSD in a finite population (Figure 1B).

I used simulations to test if fitness effects maintain PSD because the model parameters were too complex to analytically solve for equilibrium conditions. In my simulations, I tested if PSD persists for 1,000 generations, which may not reflect an equilibrium state. In the Results, I

show how this approach can be problematic if proto-Y chromosomes are on a path to fixation or

loss but have not yet reached fixation at 1,000 generations. I do this by simulating 1,000,000

generations for a subset of the fitness effects.

Figure 1. Steps in forward population genetic simulations with **(A)** infinite or **(B)** finite populations. The
best-fitting arrays are those with the lowest mean squared error (MSE) comparing simulated proto-sex
chromosome frequencies to those observed in each natural population.

Fitness effects and genotypic fitness

To generate the fitness of the 18 possible multi-chromosome genotypes (Table 1), I first

assigned fitness effects to each single chromosome genotype (e.g., for the X and Y^M

chromosomes there are three possible genotypes: X/X, X/Y^M, and Y^M/Y^M). The single

chromosome genotype is analogous to a single locus genotype in other population genetic

models (e.g., Kidwell *et al.* 1977). I generated the single chromosome genotype fitness values

by assigning a sex-specific fitness effect, s_{ij}, to each j proto-Y or proto-W chromosome (Y^M,

III^M, or IV^F) in each sex i. The fitness effect of IV^F was only assigned for females because the

female-determining proto-W chromosome (carrying *Md-tra^D*) cannot be found in male genotypes

(Table 1). Thus, there are five fitness effects: Y^M in males ($i=\mathrm{male}, j=\mathrm{Y^M}$), Y^M in females (

$i=\mathrm{female}, j=\mathrm{Y^M}$), III^M in males ($i=\mathrm{male}, j=\mathrm{III^M}$), III^M in females ($i=\mathrm{female}, j=\mathrm{III^M}$),

and IV^F in females ($i=\mathrm{female}, j=\mathrm{IV^F}$). Each s_{ij} was drawn from a uniform distribution

between -1 and 1. Positive fitness effects ($s_{ij}>0$) mean that the proto-Y or proto-W

chromosome increases fitness (i.e., it is beneficial), and negative values ($s_{ij}<0$) mean that it

decreases fitness (i.e., it is deleterious).

The fitness effects of the proto-Y and proto-W chromosome were then used to calculate

the fitness of each single chromosome genotype, using one of four dominance scenarios. The

calculations depend on whether the proto-Y chromosome has beneficial ($s_{ij}>0$) or deleterious

($s_{ij}<0$) effects, which ensures that the maximum genotypic fitness for a given chromosome

and sex is equal to 1 (Table 2). First, in the additive model, the sex-specific single chromosome

fitness values are defined such that the heterozygous genotype (X/Y^M or III/III^M) is intermediate

between the two homozygous genotypes. Second, when the proto-Y chromosomes have

dominant fitness effects, the single chromosome fitness of the heterozygote is equal to the

proto-Y homozygote. Third, when the effects of the proto-Y chromosomes are recessive, the

single chromosome fitness of the heterozygote is equal to the proto-X homozygote and different

from the proto-Y homozygote (Table 2). Fourth, the Y^M and III^M proto-Y chromosomes have

overdominant effects in males, but are additive in females. In the overdominant model, the

fitness effect of recessive deleterious proto-Y alleles is equal in magnitude to the fitness benefit

of dominant advantageous proto-Y alleles. This simplifying assumption may not be biologically

realistic, and future work could examine the implications of this assumption. In all four models,

there is a single fitness effect associated with carrying a copy of the IV^F proto-W chromosome in

females; only females can carry IV^F and it is impossible to be homozygous for IV^F (Table 1). In

all four dominance models, the fitness of each of the multi-chromosomal genotypes (Table 1) is

calculated as the product of the three relevant sex-specific single chromosome genotype fitness

values.

Table 2. Single proto-Y/proto-X chromosome genotype fitness values with different dominance scenarios.

	XX	XY	YY
additive			
$s_{ij}>0$	$1-s_{ij}$	$1-0.5s_{ij}$	1
$s_{ij}<0$	1	$1+0.5s_{ij}$	$1+s_{ij}$
dominant			
$s_{ij}>0$	$1-s_{ij}$	1	1
$s_{ij}<0$	1	$1+s_{ij}$	$1+s_{ij}$
recessive			
$s_{ij}>0$	$1-s_{ij}$	$1-s_{ij}$	1
$s_{ij}<0$	1	1	$1+s_{ij}$
overdominant (male only)			
$s_{ij}>0$	$1-s_{ij}$	1	$1-s_{ij}$
$s_{ij}<0$	$1+s_{ij}$	1	$1+s_{ij}$

Three genotypes are shown (XX, XY, and YY). The Y chromosome could either be Y^M or III^M, and the X
chromosome could either be X or III.

I performed a normalization to ensure that the the maximum multi-chromosome

genotype fitness in each sex is equal to 1. This is necessary because not all multi-chromosome

genotypes are possible for both males and females, which means that the maximum product of

all single chromosome genotypes can be <1 even though the maximum single chromosome

genotype fitness values are set to 1. To ensure that the maximum female multi-chromosome

genotype is 1, I divided each female multi-chromosome genotype fitness by the maximum

female multi-chromosome genotype fitness. I did the same for males.

All four dominance scenarios (additive, dominant, recessive, or overdominant) attribute a

single sex-specific fitness effect to each proto-Y and proto-W chromosome, whether it is Y^M, III^M,

or IV^F. I am therefore assuming that all copies of each proto-sex chromosome in the population

carry an identical suite of beneficial and deleterious alleles (i.e., all copies of III^M are identical, all

copies of Y^M are identical, and all copies of IV^F are identical). In addition, all models assume

complete linkage between any allele(s) under selection and the *Mdmd* locus on the Y^M and III^M

chromosomes. Similarly, I assume complete linkage between the female-determining *Md-tra^D*

locus and any alleles on the IV^F chromosome. Complete linkage may be achieved if, for

example, chromosomal inversions suppress recombination between the proto-Y and proto-X (or

proto-Z and proto-W) chromosomes in heterozygotes (Bergero and Charlesworth 2009; Wright

et al. 2016). Free recombination in sex chromosome homozygotes would not affect genetic

variation on a sex chromosome because all copies of each chromosome are assumed to carry

the same alleles.

Simulations with infinite population size

I used forward simulations to determine how the fitness effects of the proto-sex

chromosomes affect their frequencies in populations. These simulations were performed with

non-overlapping generations and random mating, assuming a population of infinite size

(simulations with finite population sizes are described later). Each generation of a simulation

consists of two discrete steps (Figure 1A). First, the frequency of each of the 18 genotypes is

multiplied by its corresponding fitness value, which models differential survival across

genotypes. Second, recursion equations developed by Hamm (2008) and previously

implemented by Meisel *et al.* (2016) are used to model the production of progeny by random

mating (Supplemental Table S1). After 1,000 generations of selection and random mating, the

frequency of each genotype and proto-sex chromosome was calculated. This process was

performed for 1,000,000 fitness arrays for each dominance scenario (additive, dominant,

recessive, and overdominant) and four possible initial genotype frequencies (16,000,000 total

simulations).

I considered four possible initial genotype frequencies for each simulation. The first three

initial frequencies are based on the observed frequencies of Y^M, III^M, and *Md-tra^D* from three

North American populations (Meisel *et al.* 2016). These populations were sampled in Chino, CA

(Meisel *et al.* 2016), Wake County, NC (Hamm and Scott 2008), and Chemung County, NY

(Scott *et al.* 2013). Initial genotype frequencies were calculated based on the observed

frequencies of Y^M, III^M, and *Md-tra^D*, assuming random mating (Meisel *et al.* 2016). These three

populations were chosen because Y^M, III^M, and *Md-tra^D* have all remained at a frequency >1%

across multiple years of sampling (Hamm *et al.* 2005; Hamm and Scott 2008; Meisel *et al.*

2016). I used frequencies from actual populations as the initial frequencies because I am

specifically interested in how selection can maintain PSD at the frequencies observed in natural

populations. The fourth starting values consist of all 18 genotypes at the same initial frequency

(i.e., 1/18 for each genotype). This allows me to evaluate how selection pressures can drive the

proto-sex chromosomes to the frequencies observed in natural populations if they start at

arbitrary values that deviate from the observed frequencies.

To determine whether the fitness effects maintain PSD, I first selected fitness arrays that

produced frequencies of each proto-sex chromosome (Y^M, III^M, IV^F, and their homologous

chromosomes, X, III, and IV) that are all >0.1% after 1,000 generations. This criterion ensures

that all three chromosomes are polymorphic with both alleles at a frequency that is measurable

(i.e., 1/1,000) given the sampling schemes used in previous studies of natural house fly

populations (Hamm *et al.* 2005; Hamm and Scott 2008; Meisel *et al.* 2016). I refer to genotype

fitness arrays with all proto-sex chromosomes present at a frequency >0.1% as maintaining

PSD, although not necessarily at the frequencies observed in natural populations.

From the genotype fitness arrays that maintain PSD (i.e., all proto-sex chromosomes

>0.1%), I selected those that produce proto-sex chromosome frequencies most similar to the

frequencies observed in natural populations. To do so, I calculated the mean squared error

(MSE) between the frequencies of Y^M, III^M, and IV^F after 1,000 generations in a simulation (S)

and the observed frequencies (O) in a natural population (CA, NC, or NY):

269
$$\mathrm{MSE} = \frac{1}{3}[(O_{Y^M} - S_{Y^M})^2 + (O_{III^M} - S_{III^M})^2 + (O_{tra^D} - S_{tra^D})^2].$$

When using simulations that started with proto-sex chromosome frequencies observed in a

population (e.g., CA), I only used MSE to compare with observed frequencies from that same

population. When simulations started with equal frequencies of all 18 genotypes, I compared

simulated chromosomes frequencies with the observed frequencies in each of the three

populations (CA, NC, and NY).

I retained the 1,000 fitness arrays that produce proto-sex chromsome frequencies with

the lowest MSE for a given dominance model, population, and starting genotype frequency. I

refer to each of these 1,000 fitness arrays as the best-fitting arrays for each population.

Simulations for the 1,000 best-fitting arrays were then run for 1,000,000 generations and

compared to the 1,000 generation simulations to assess the stability of chromosomal

frequencies. I also selected 1,000 random fitness arrays and 1,000 fitness arrays that maintain

PSD to perform 1,000,000 generation simulations.

Simulations with finite population sizes

I assessed how well the 1,000 best-fitting fitness arrays for each population, starting

frequency, and dominance model maintain PSD when population sizes are finite. Finite

population sizes introduce stochasticity to the change in allele frequencies across generations

(i.e., genetic drift), in contrast to the deterministic effects of constant fitness values in a

population of infinite size (Hartl and Clark 2007). It is not possible to calculate the probability of

fixation for alleles in a finite population when there is complex sex-linked inheritance, as in the

house fly PSD system (Meisel *et al.* 2016). To overcome this limitation, I estimated the

probability of fixation using simulations that modeled finite populations by including multinomial

sampling each generation.

I simulated finite populations by adding one additional step to the simulations described

above in order to model a population size (N) of 10,000 individuals (Figure 1B). This population

size was chosen because it is small enough to capture the effects of drift within the time scale of

my simulations. Each simulation was started with the frequencies of the proto-sex

chromosomes observed in the population where the fitness array was identified as best-fitting.

In each generation, following multiplication by the fitness array (i.e., natural selection), I used

multinomial sampling to calculate the new frequencies of all 18 genotypes. Multinomial sampling

for all genotype was performed with 10,000 trials (i.e., $N=10^4$) and a probability of success equal

to the frequency of each genotype after selection. The resulting array of 18 genotype

frequencies was then divided by the sum of all values to ensure that the genotype frequencies

sum to 1. This array of genotype frequencies (after selection and multinomial sampling) was

next used as input into the same recursion equations described above to calculate genotype

frequencies after random mating (Supplemental Table S1). The process (selection, multinomial

sampling, and random mating) was repeated for 1,000 generations. I simulated 100 replicate

finite populations for each fitness array. I then used the frequency with which fixation and loss

occur in those 100 replicates as an estimate of the probability of fixation and loss of proto-sex

chromosomes within 1,000 generations.

I determined a null expectation for fixation or loss of proto-sex chromosomes in finite

populations by using simulations without selection (i.e., genetic drift and no fitness differences

across genotypes). To those ends, I performed simulations with 10,000 individuals for 1,000

generations, as was done in the simulations with selection and drift. I performed 1,000 replicate

"drift-only" simulations with starting values at the observed frequencies of the proto-sex

chromosomes in each population (CA, NC, and NY). These drift-only simulations included the

same steps as the simulations with both selection and drift, except that there is no differential

survival (i.e., natural selection) step in the drift-only simulation. After 1,000 generations, I

calculated the frequency of each proto-Y and proto-W chromosome (Y^M, III^M, and IV^F), as well

as the frequency with which each chromosome was lost, across the 1,000 simulations.

Results

I used forward population genetic simulations to assess how natural selection can

maintain PSD in a species with two proto-Y chromosomes (Y^M and III^M) and one proto-W

chromosome (IV^F), modeled after the house fly (Hamm *et al.* 2015). Simulations were performed

assuming either additive, dominant, recessive, or overdominant fitness effects of the Y^M and III^M

proto-Y chromosomes. Simulations were started with genotype frequencies based on proto-sex

chromosome counts in each of three natural populations (CA, NC, or NY) or equal frequencies

of all 18 genotypes. The general trends are the same regardless of the starting genotype

frequencies, and I only present the results of simulations starting with equal frequencies in the

main text (other results are presented in the Supplemental Material).

Dominance of fitness effects and the maintenance of polygenic sex determination

The probability that PSD is maintained (with a frequency of each proto-sex chromosome

>0.1% after 1,000 generations) is affected by the dominance of the fitness effects of the Y^M and

III^M proto-Y chromosomes (Supplemental Figures S1A, S2A). When fitness effects of proto-Y

chromosomes are additive, <5% of fitness arrays maintain PSD. In contrast, when fitness

effects of the proto-Y chromosomes are recessive, ~20% of fitness arrays maintain PSD.

Overdominant fitness effects are the most likely to maintain PSD, with >40% of fitness arrays

maintaining proto-sex chromosomes at a frequency >0.1% after 1,000 generations. When

additive, recessive, or overdominant (but not dominant) fitness effects maintain PSD, there is a

broad range of frequencies at which the proto-Y chromosomes (Y^M and III^M) are maintained after

1,000 generations (Figure 2; Supplemental Figure S3). Randomly chosen fitness effects, in

comparison, tend to produce one of three proto-Y chromosome frequency classes: 0 (loss of a

proto-Y, because a different XY system has taken over), 1 (fixation of a proto-Y chromosome,

corresponding to a monogenic ZW system), and 0.25 (the expectation for a monogenic XY

system).

Most dominant fitness effects that appear to maintain PSD are paths to the eventual

fixation of Y^M and III^M. More than 25% of the dominant fitness effects that I considered maintain

PSD for 1,000 generations (Supplemental Figures S1A, S2A), but most of those produce very

high frequencies of the Y^M and III^M chromosomes (Figure 2B; Supplemental Figure S3). A

proto-Y chromosome with a dominant fitness effect segregating at very high frequency is

consistent with a beneficial effect that masks the recessive variant (in this case the homologous

proto-X), allowing the proto-X chromosome to persist at a low frequency for a long time (Hartl

and Clark 2007). To test this prediction, I ran simulations for 1,000,000 generations with the

best-fitting selection pressures, and I considered a proto-sex chromosome to reach "fixation" or

"loss" if it achieves a frequency >99.9% or <0.1%, respectively (i.e., the same threshold for

maintaining PSD after 1,000 generations). Consistent with the prediction, >50% of dominant

fitness effects that maintain PSD for 1,000 generations eventually cause the Y^M or III^M proto-Y

chromosomes to reach fixation (frequency >99.9%) within 1,000,000 generations, regardless of

the starting genotype frequencies (Supplemental Figures S4–S7). In contrast, ≤1% of additive,

recessive, or overdominant fitness effects of proto-Y chromosomes that maintain PSD for 1,000

generations eventually result in fixation of a proto-Y chromosome after 1,000,000 generations.

Despite the eventual fixation of proto-Y chromosomes in many parameterizations of dominant

fitness effects, there are a subset of dominant fitness effects that maintain Y^M and III^M at lower

frequencies, resembling those observed in natural populations (Figure 2B). For these dominant

fitness effects that produce proto-sex chromosome frequencies similar natural populations,

fixation of proto-Y chromosomes never occurs after 1,000,000 generations (Supplemental

Figures S4–S7). Therefore, dominant fitness effects can maintain PSD at equilibrium, although

most dominant fitness effects result in fixation or loss of a proto-Y chromosome.

Selecting only the 1,000 fitness effects that produce proto-sex chromosome frequencies

most similar to natural populations further affects the relationships between dominance and

proto-sex chromosome frequencies. For example, proto-sex chromosome frequencies are more

similar to those in natural populations when fitness effects of the proto-Y chromosomes are

dominant, recessive, or overdominant, as opposed to additive fitness effects (Figure 2;

Supplemental Figures S1, S2, S3). The poorer fit of additive models arises because III^M and IV^F

segregate at lower frequencies in the CA, NC, and NY populations than they do in the

best-fitting simulations with additive fitness effects (Figure 2; Supplemental Figure S3).

Figure 2. Smoothed histograms show the frequency of each proto-sex chromosome (Y^M, III^M, and IV^F)
after 1,000 generations in simulations using either 1,000 random fitness arrays (random), fitness arrays
that maintain PSD (polygenic), or the 1,000 best-fitting fitness arrays for each population (CA, NC, or NY).
Simulations were started with equal frequencies of all genotypes. The vertical line within each histogram
shows the median. Red dots show the observed proto-sex chromosome frequencies in each natural
population. Fitness arrays were calculated assuming either **(A)** additive, **(B)** dominant, **(C)** recessive, or
(D) overdominant fitness effects of the proto-Y chromosomes.

The low frequency of the III^M chromosome in the CA, NC, and NY populations has

consequences for the ability of natural selection to maintain the polymorphism. As above, I

define loss of the III^M chromosome (or fixation of the III chromosome) as a frequency below

0.1% (or above 99.9% for III). Additive fitness effects that maintain the III^M chromosome at

frequencies similar to those in CA, NC, or NY for 1,000 generations often (9–17% of the time)

result in the loss of III^M within 1,000,000 generations (Supplemental Figures S8, S9). In contrast,

<7% of dominant, recessive, or overdominant fitness effects that maintain III^M at frequencies

observed in natural populations for 1,000 generations lead to the eventual loss of III^M within

1,000,000 generations. III^M is especially likely to be lost within 1,000,000 generations when it is

segregating at a lower frequency after 1,000 generations, regardless of the dominance of fitness

effects (Supplemental Figures S8, S9). The Y^M chromosome, in comparison, segregates at a

higher frequency that III^M all three populations (Figure 2B). Y^M is lost in ≤2% of 1,000,000

generation simulations when I apply fitness effects that maintain proto-sex chromosomes at

frequencies similar to those observed in natural populations, regardless of the dominance of

fitness effects (Supplemental Figure S10). I therefore conclude that low frequency proto-sex

chromosomes, such as III^M, are difficult to maintain as polymorphisms at equilibrium under the

model I have applied here.

The majority of fitness effects that maintain PSD cause IV^F to segregate at a frequency

of 25%, regardless of the dominance of the proto-Y chromosomes (Figure 2; Supplemental

Figure S3). A frequency of 25% is the expectation for a W chromosome in a randomly mating

population with monogenic ZW sex determination. Therefore, in a system with 2 proto-Y

chromosomes and one proto-W, PSD is expected to be maintained by the (near) fixation of the

ZW female genotype. In contrast, IV^F makes up 2–9% of all fourth chromosomes in the CA, NC,

and NY populations (Hamm and Scott 2008; Scott *et al.* 2013; Meisel *et al.* 2016). A subset of

fitness arrays in my simulations maintain IV^F at <25% for all dominance models, including those

that produce proto-sex chromosome frequencies most similar to those observed in natural

populations (Figure 2; Supplemental Figure S3). It is thus possible for PSD to be maintained

with a single proto-W chromosome at a frequency <25%. However, even with those best fitting

selection parameters, the frequency of IV^F tends to be greater than observed in natural

populations (Figure 2).

Selection pressures can maintain polygenic sex determination in finite populations

I next used simulations to examine how well natural selection maintains PSD when

population sizes are finite ($N=10^4$ individuals). I calculated the proportion of simulations (out of

100) in which a given proto-sex chromosome reached fixation for each fitness array to estimate

of the probability of fixation (P_{fix}). The P_{fix} value for a given proto-Y or proto-W chromosome

(Y[M], III[M], or IV[F]) is equal to the probability of loss (P_{loss}) of the homologous proto-X or proto-Z

(X, III, or IV), and *vice versa*. I used Fisher's exact test to determine if the number of fixations of

a proto-sex chromosome is greater or less in a finite population of the same size and without

selection (i.e., genetic drift only).

When a proto-sex chromosome is at low frequency, selection pressures that maintain

PSD decrease P_{fix} and P_{loss} in finite populations, relative to populations without selection.

For example, III[M] is rare in all three populations, and it is the most likely proto-sex chromosome

to be lost when population size is finite (Supplemental Figures S11, S12). In turn, selection

pressures that maintain PSD result in a decreased P_{fix} of chromosome III relative to drift alone

(Figure 3; Supplemental Figure S13), which is equivalent to decreasing P_{loss} of III[M]. The IV[F]

proto-W chromosome is also at a low frequency in all three populations, and the maintenance of

the IV[F] polymorphism (i.e., the standard IV chromosome does not fix) is substantially greater

when there is selection acting in a finite population (Figure 3; Supplemental Figure S13).

Therefore, when loss via drift is likely because a proto-sex chromosome is at low frequency

(such as III[M] and IV[F]), selection pressures that maintain PSD decrease the probability of loss.

Figure 3. Bar plots show the number of fitness arrays that result in a deficiency (gray) or excess (white)
of fixations of each proto-sex chromosome (X, Y^M, III, III^M, or IV) relative to simulations in which there are
no fitness differences across proto-sex chromosomes (i.e., drift only). The fitness arrays are the 1,000
best-fitting fitness values for each population (CA, NC, or NY) when simulations were started with equal
frequencies of all genotypes. Fitness effects of the proto-Y chromosomes are either **(A)** additive, **(B)**
dominant, **(C)** recessive, or **(D)** overdominant.

A different effect is observed for sex chromosomes segregating at higher frequencies.

The X and Y^M chromosomes rarely fix in finite populations with selection pressures that should

maintain the polymorphism (Supplemental Figures S11, S12). Y^M is at a higher frequency than

III^M and IV^F across all three populations (Meisel *et al.* 2016), and the high frequency of Y^M is

likely responsible for its low P_{loss} within 1,000 generations. However, for a small fraction of the

best-fitting arrays (<10%), P_{fix} for both the X and Y^M chromosomes is greater with selection

and drift than under drift alone (Figure 3). The slightly elevated P_{fix} of the X and Y^M

chromosomes with selection is observed for all four dominance models and regardless of how

the best-fitting arrays are identified (Supplemental Figure S13). Therefore, when fixation or loss

is unlikely via drift (because the proto-Y chromosome is at high frequency), selection pressures

that maintain PSD do not decrease P_{fix} or P_{loss} of a proto-sex chromosome in a finite

population.

Sexual antagonism and the maintenance of polygenic sex determination

I next inspected if there is a negative correlation between male and female fitness when

PSD is maintained. A negative intersexual fitness correlation is a hallmark of sexually

antagonistic genetic variation (Rice and Chippindale 2001; Chippindale *et al.* 2001;

Bonduriansky and Chenoweth 2009). Each of the eight male house fly proto-sex chromosome

genotypes has a corresponding female genotype that differs only because females carry a copy

of the IV^F proto-W chromosome and males do not (Table 1). There are two additional female

genotypes (X/X; III/III; IV/IV and X/X; III/III; IV^F/IV) that do not have a corresponding male

genotype because they do not have a proto-Y chromosome. To test for sexually antagonistic

multi-chromosome genotypes, I calculated Spearman's rank order correlation of fitness between

males and females (ρ_{MF}) for the eight pairs of male and female genotypes in each fitness

array.

Additive fitness effects that maintain PSD (i.e., all proto-sex chromosomes at frequency

>0.1%) tend to have $\rho_{MF} < 0$ (Figure 4A; Supplemental Figure S14). However, when

proto-sex chromosome frequencies resemble those in natural populations and fitness effects

are additive, ρ_{MF} can be either positive or negative. There is also no consistent signal of

$\rho_{MF} < 0$ for random fitness effects, regardless of dominance (Figure 4; Supplemental Figure

S14), demonstrating that $\rho_{MF} < 0$ is not an intrinsic property of the model. In comparison,

$\rho_{MF} < 0$ when I previously applied a model with additive fitness effects both within and across

chromosomes (Meisel *et al.* 2016). In the model presented here, multi-chromosome genotype

fitness is calculated as the product of single chromosome genotype fitness values, rather than

by summing across chromosomes. Therefore, there is some evidence that additive fitness

effects that maintain PSD result in $\rho_{MF} < 0$, but it depends on specifics of how the model is

parameterized.

Figure 4. Smoothed histograms show distributions of intersexual fitness correlations (ρ_{MF}) for
genotype fitness arrays used in simulations started with equal frequencies of all genotypes. Correlations
are shown for the 1,000 best-fitting genotypic fitness arrays for the CA, NC, and NY populations.
Correlations are also shown for all fitness arrays that maintain PSD (polygenic), as well as 1,000 random
fitness arrays. Dashed vertical lines show a correlation of 0, and solid vertical lines within histograms
479   show the median.

When fitness effects are dominant, $\rho_{MF} < 0$ when proto-sex chromosome frequencies

resemble those in natural populations. In contrast, ρ_{MF} can be positive or negative for

dominant fitness effects that maintain PSD regardless of the proto-sex chromosome frequency

(Figure 4B; Supplemental Figure S14B). However, as noted above, many of the cases where

dominant fitness effects appear to maintain PSD are instead on a path to fixation that is not yet

reached at 1,000 generations. Indeed, $\rho_{MF} < 0$ when I only sample the dominant fitness

effects that maintain PSD for 1,000,000 generations, and not for the dominant fitness effects

that eventually reach fixation within 1,000,000 generations (Supplemental Figure S15).

Therefore, assuming 1,000,000 generations adequately approximates the equilibrium state,

$\rho_{MF} < 0$ when dominant fitness effects maintain PSD.

When fitness effects are overdominant or recessive, ρ_{MF} does not provide evidence for

sexually antagonistic selection. First, there is no clear pattern of $\rho_{MF} < 0$ when fitness effects

are recessive (Figure 4C; Supplemental Figure S14C). Second, overdominant fitness effects

that produce the best-fitting genotype fitness arrays nearly all have $\rho_{MF} > 0$ (Figure 4D;

Supplemental Figure S14D), but this result is difficult to interpret because overdominant effects

produce a concave relationship between male genotype and fitness (i.e., heterozygous males

are defined as the most fit). Therefore, a simple rank order correlation (i.e., ρ_{MF}) does not

adequately capture the relationship between male and female fitness when the proto-Y

chromosomes have overdominant effects in males.

Negative intersexual fitness correlations are not caused by sexually antagonistic proto-Y

chromosomes

The negative ρ_{MF} that I observe for additive and dominant fitness effects that maintain

PSD (Figure 4; Supplemental Figure S14) are suggestive of sexually antagonistic selection. A

potential cause of $\rho_{MF} < 0$ is sexually antagonistic effects of the proto-Y chromosomes if, for

example, Y^M and III^M carry male-beneficial and female-detrimental alleles (Fisher 1931; Rice

1987). In contrast to this hypothesis, I do not observe a consistent signal of male-beneficial and

female-detrimental proto-Y chromosomes (Figure 5; Supplemental Figure S16).

Figure 5. Smoothed histograms show the distributions of fitness effects of each proto-sex chromosome in
each sex for 1,000 random fitness arrays (random), fitness arrays that maintain PSD (polygenic), or the
1,000 best-fitting fitness arrays for each population (CA, NC, or NY). Simulations were started with equal
frequencies of all genotypes. The vertical line within each histogram shows the median, and dashed lines
in each panel are at fitness value of 0. Fitness arrays were calculated assuming either (**A**) additive, (**B**)
dominant, (**C**) recessive, or (**D**) overdominant fitness effects of the proto-Y chromosomes.

Additive fitness effects that maintain PSD often have female-beneficial and

male-detrimental effects of the proto-Y chromosomes (Figure 5A; Supplemental Figure S16A).

Female-beneficial and male-detrimental Y chromosomes are opposite of the expected effects

under most models (Rice 1987, 1996). However, some previous population genetic modeling

has suggested the possibility of "feminization" of the Y chromosome (Cavoto *et al.* 2018). Within

individual additive parameterizations that maintain PSD, the proto-Y chromosomes do not

consistently confer a greater fitness benefit to either males or females (Supplemental Figures

S17, S18).

There is some evidence that Y^M is male-beneficial and female-detrimental when

dominant fitness effects maintain PSD (Figure 5B; Supplemental Figure S16B). Additionally,

both Y^M and III^M often confer a greater fitness benefit to males than females when dominant

fitness effects maintain PSD (Supplemental Figures S17, S18, S19). However, III^M does not

tend to have male-beneficial and female-detrimental effects in the models with dominant fitness

effects (Figure 5B; Supplemental Figure S16B). Therefore, when dominant fitness effects

maintain PSD the multi-chromosome genotypes tend to have sexually antagonistic fitness

effects (Figure 4; Supplemental Figures S14, S15), but the individual chromosomes are not

necessarily sexually antagonistic themselves.

Overdominant fitness effects that maintain proto-sex chromosomes at frequencies

similar to those observed in natural populations have the strongest evidence of male-beneficial

and female-detrimental proto-Y chromosomes (Figure 5D; Supplemental Figures S16D, S17D,

S18D). The overdominant model is constrained to require male-beneficial effects of the proto-Y

chromosomes in heterozygotes, but female fitness effects can be beneficial or deleterious.

When proto-sex chromosomes are at frequencies similar to those in the CA, NC, or NY

populations, the proto-Y chromosomes are almost always female-deterious. These sexually

antagonistic effects of the Y^M and III^M proto-Y chromosomes in the overdominant models does

not cause negative intersexual correlations of multi-chromosomal genotype fitness values

(Figure 4; Supplemental Figure S14) because of the concave relationship between male

genotype and fitness (i.e., heterozygous males are defined as the most fit).

Female-beneficial effects of the proto-W chromosome

The IV^F proto-W chromosome is nearly always beneficial to females in fitness arrays that

maintain PSD (Figure 5). I observe the female-beneficial effect of IV^F in all cases when PSD is

maintained, regardless of the dominance of fitness effects of the proto-Y chromosomes,

whether or not proto-sex chromosomes frequencies are similar to those in natural populations,

and independently of the genotype frequencies used to start the simulations (Figure 5;

Supplemental Figure S16). There also tends to be a positive correlation between the fitness

effect and frequency of IV^F when proto-Y chromosomes have additive, dominant, recessive, or

overdominant effects, regardless of the starting genotype frequencies used in the simulations

(Supplemental Figures S20, S21). A positive correlation indicates that the more

female-beneficial the IV^F chromosome, the higher its frequency. The only exception to this rule

is when dominant fitness effects maintain PSD at frequencies not similar to those observed in

natural populations; however, as described above, these cases are on a path to fixation and

therefore not at equilibrium

Discussion

I used forward simulations of a complex PSD system—with two proto-X/Y pairs and one

proto-Z/W, modeled after the house fly—to investigate how the dominance of alleles under

selection affects the maintenance of PSD. I found that the ability of natural selection to maintain

PSD does indeed depend on the dominance of fitness effects. Specifically, PSD is more likely to

be maintained if proto-Y chromosomes have overdominant fitness effects, and least likely if the

fitness effects are additive (Supplemental Figures S1, S2). Dominance also affects how similar

proto-sex chromosome frequencies are to those in natural populations (Figure 2; Supplemental

Figure S3). Moreover, the extent of sexually antagonism depends on the dominance of fitness

effects: dominant effects of proto-Y chromosomes result in the strongest signatures of sexually

antagonistic multi-chromosomal genotypes (Figure 4; Supplemental Figures S14, S15),

whereas overdominant fitness effects produce the strongest signatures of sexually antagonistic

proto-Y chromosomes (Figure 5; Supplemental Figure S16). Despite these differences across

dominance models, selection pressures that maintain PSD are all effective at preventing the

fixation or loss of proto-sex chromosomes via genetic drift, regardless of the dominance of

fitness effects (Figure 3; Supplemental Figures S11–S13).

Sexually antagonistic genotypes can maintain polygenic sex determination without

sexually antagonistic proto-Y chromosomes

Sexual antagonism has previously been identified as a key component of the

maintenance of PSD within populations (Bull 1983; Rice 1986; van Doorn and Kirkpatrick 2007;

Meisel *et al.* 2016). A negative inter-sexual correlation of fitness across genotypes is a hallmark

of sexually antagonistic selection (Rice and Chippindale 2001; Chippindale *et al.* 2001;

Bonduriansky and Chenoweth 2009). I find strong evidence for $\rho_{MF} < 0$ when dominant fitness

effects maintain PSD (Figure 4B; Supplemental Figure S15). There is also some evidence that

additive fitness effects that maintain PSD tend to have negative ρ_{MF}, although it is dependent

on the frequency at which the proto-sex chromosomes are maintained (Figure 4A;

Supplemental Figure S14A). However, neither dominant nor additive fitness effects that

maintain PSD have male-beneficial and female-detrimental effects of all individual proto-Y

chromosomes (Figure 5; Supplemental Figure S16). Therefore, the sexually antagonistic

multi-chromosomal genotypes that maintain PSD are emergent properties of cumulative fitness

effects across chromosomes. This is evidence for a decoupling between single chromosome

sexually antagonistic effects and sexual antagonism across multi-chromosome genotypes. It

also suggests that testing for sexually antagonistic effects of individual proto-sex chromosomes

may not be a sufficient assay of the role sexual antagonism plays in the maintenance of

complex, multi-chromosomal PSD.

Overdominance and the maintenance of polygenic sex determination

Overdominant fitness effects of proto-Y chromosomes in males are more likely to

maintain PSD than any other dominance scenario that I tested. Nearly half of the 4,000,000

overdominant fitness effects that I tested maintain all three proto-sex chromosomes at a

frequency >0.1% for at least 1,000 generations (Supplemental Figures S1A, S2A). In addition,

nearly all of those overdominant fitness effects that maintain PSD for 1,000 generations

continue to do so for at least 1,000,000 generations (Supplemental Figures S4–S7), suggesting

they are indeed stable polymorphisms at equilibrium. The frequencies with which overdominant

fitness effects can maintain multiple polymorphic proto-sex chromosomes covers the full range

of possible values, from close to 0 to nearly 1 (Figure 2D; Supplemental Figure S3D).

The extent of overdominant genetic variation—and balancing selection more

generally—has been a topic of considerable debate in population genetics (Crow 1987; Delph

and Kelly 2014; Fijarczyk and Babik 2015), with a few classic examples of heterozygote

advantage (e.g., Dobzhansky 1947; Allison 1956). Contemporary evidence for overdominant

fitness effects in natural populations is mixed, including some work that suggests

overdominance is rare (Andrés *et al.* 2009; Sellis *et al.* 2011, 2016; Hedrick 2012; Bitarello *et al.*

2018). Notably, in platyfish, which has a single chromosome PSD system, heterozygous males

(XY) and females (WY or WX) have higher fitness than homozygotes (Orzack *et al.* 1980).

Regardless of the prevalence of overdominance in natural populations, it is one of the most

straightforward mechanisms by which polymorphisms can be maintained (Hartl and Clark 2007).

In my overdominant model, I assume that the proto-Y chromosomes carry male-beneficial

additive or dominant alleles (Rice 1984, 1992), along with recessive deleterious alleles that

have opposing effects equal in magnitude to the beneficial alleles (Charlesworth and

Charlesworth 2000). Testing if these assumptions are biologically realistic would require

measuring the magnitude and dominance of beneficial and deleterious alleles on proto-Y

chromosomes.

When overdominant fitness effects maintain proto-sex chromosomes at frequencies

most similar to those observed in natural populations, I find that they do so with

female-deterious effects of the proto-Y chromosomes (Figure 5D; Supplemental Figure S16D).

This is the strongest evidence I observe for sexually antagonistic fitness effects of individual

proto-Y chromosomes (as opposed to multi-chromosomal genotypes) in any dominance

scenario. It is also consistent with previous work demonstrating that overdominance in one sex

can maintain genetic variants at individual loci even if there is directional selection against one

allele in the other sex (Kidwell *et al.* 1977). In contrast to the additive and dominant fitness

effects, there is no evidence for $\rho_{MF} < 0$ when overdominant fitness effects maintain PSD

(Figure 4; Supplemental Figure S14). This provides additional evidence for a decoupling

between single chromosome sexually antagonistic effects and sexual antagonism across

multi-chromosome genotypes.

Sexually antagonistic proto-Y chromosomes require there to be segregating sexually

antagonistic genetic variation. Male-beneficial and female-deleterious alleles are likely to

accumulate on Y chromosomes because of their male-limited inheritance (Rice 1984). It is not

clear, however, if that prediction applies when the Y chromosome can be transmitted through

females in a complex PSD system such as in house fly. A more appropriate point of comparison

for PSD systems may be autosomal and X-linked sexually antagonistic genetic variation, of

which there is evidence for a substantial amount (Rice 1992; Chippindale *et al.* 2001; Calsbeek

and Sinervo 2004; Fedorka and Mousseau 2004; Foerster *et al.* 2007; Brommer *et al.* 2007;

Innocenti and Morrow 2010).

My results are also consistent with theory that suggests sexually antagonistic selection is

important for the early evolution of sex chromosomes, as well as empirical studies that identified

sexually antagonistic variants on young sex chromosomes (Lindholm and Breden 2002; van

Doorn and Kirkpatrick 2007, 2010; Roberts *et al.* 2009). It remains to be determined, however,

whether young sex chromosomes are enriched for sexually antagonistic variants, if those

variants are maintained by balancing selection, and if the sexually antagonistic alleles contribute

to the maintenance of PSD. Investigating these problems could follow the approaches

previously developed to test if balancing selection maintains sexually antagonistic alleles

throughout the genome (e.g., Dutoit *et al.* 2018; Ruzicka *et al.* 2019).

The dominance of sexually antagonistic alleles

The dominance of fitness effects is an important parameter affecting the ability sexually

antagonistic selection to maintain genetic variation (Connallon and Chenoweth 2019). For

example, a sexually antagonistic allele is more likely to be maintained as a polymorphism if the

sex-specific deleterious effects are recessive (Kidwell *et al.* 1977). In contrast, when fitness

effects are dominant, stronger selection pressures are required to maintain the polymorphism

(Kidwell *et al.* 1977). My work presented here contributes to the special case where the alleles

under selection are on proto-sex chromosomes (as opposed to autosomes or heteromorphic

sex chromosomes). Consistent with Kidwell *et al.* (1977), I find that dominant fitness effects that

maintain PSD do indeed have strong selection in favor of or against proto-Y chromosomes in

one sex (Figure 5; Supplemental Figure S16).

In most of the models I consider here, the dominance of fitness effects of the proto-Y

chromosomes is the same in both sexes. However, the dominance of fitness effects may be

sex-specific (Barson *et al.* 2015; Grieshop and Arnqvist 2018). Such dominance reversals could

be important for the maintenance of sexually antagonistic alleles (Spencer and Priest 2016;

Connallon and Chenoweth 2019), including at sex-linked loci (Fry 2010). Therefore, it is

possible that sex-specific dominance could create a broader range of fitness values that

maintain PSD via sexually antagonistic selection. Future work could consider this possibility by

including sex-specific dominance as an additional randomly chosen variable within my modeling

framework.

Recombination on the proto-sex chromosomes and Y-linked alleles

My model makes assumptions about recombination on the sex chromosomes that may

affect my conclusions about the maintenance of PSD. Specifically, I assume that there is no

recombination between the male- or female-determining locus on each proto-Y or proto-W

chromosome and the allele(s) under selection. X-Y and Z-W recombination may be suppressed

by chromosomal inversions that create tight genetic linkage between the sex-determining locus

and sexually antagonistic alleles (Bergero and Charlesworth 2009; Wright *et al.* 2016). While

there is no direct evidence for inversions or other recombination suppressors on the house fly

sex chromosomes, male meiosis in flies is thought to occur without crossing over (Gethmann

1988). A lack of crossing over in males would prevent X-Y recombination in males without the

need for sex-chromosome-specific modifiers. However, there is evidence for male

recombination in house fly (Feldmeyer *et al.* 2010), which opens up the possibility for X-Y

recombination. Furthermore, in the absence of Y-linked suppressors, X-Y recombination could

also occur in the multiple female genotypes that carry a proto-Y chromosome (Table 1).

There are at least three other consequences of recombination on the sex chromosomes

that could affect how sex-specific selection maintains PSD. First, my model assumes that all

copies of each proto-Y and proto-W chromosome have identical fitness effects. This assumption

is consistent with population genetics theory that predicts Y-linked variants cannot segregate as

protected polymorphisms when there is no X-Y recombination (Clark 1987). In contrast to this

prediction, there are numerous examples of Y-linked polymorphisms with phenotypic effects in

Drosophila, where the X and Y do not recombine (Clark 1990; Zhang *et al.* 2000; Lemos *et al.*

2008, 2010; Griffin *et al.* 2015; Brown *et al.* 2020). Relevant to the house fly, these examples

include Y-linked variation across heterogeneous environments over the species' geographic

range (Rohmer *et al.* 2004). Future work could incorporate variation in fitness effects of each

proto-Y and proto-W chromosome into my model to assess how segregating polymorphisms on

the proto-sex chromosomes affect the maintenance of PSD.

A second consequence of sex chromosome recombination could affect how

overdominant fitness effects maintain PSD. I assume that overdominance arises from proto-Y

chromosomes that carry recessive deleterious alleles. This assumption is based on the

prediction that suppressed X-Y recombination leads to the "degeneration" of the Y chromosome

via accumulation of (recessive) deleterious alleles because of Muller's ratchet and hitchhiking

(Charlesworth and Charlesworth 2000; Bachtrog 2013). In contrast, X-Y recombination in males

and/or females could help purge recessive deleterious mutations from the Y^M and III^M

chromosomes (Muller 1964; Felsenstein 1974). Moreover, there are multiple genotypes that are

homozygous for a proto-Y chromosome (Table 1), which would expose recessive deleterious

mutations to selection. Selection in these proto-Y chromosome homozygotes would further

remove recessive deleterious alleles from populations (Charlesworth *et al.* 1990; Barrett and

Charlesworth 1991). Therefore, while overdominant fitness effects are likely to maintain PSD,

proto-Y chromosomes may not possess the necessary alleles for such selection pressures to

act upon. However, overdominance can also occur by other mechanisms besides degeneration

of the Y chromosome and independent of accumulation of recessive deleterious Y-linked alleles

(Sellis *et al.* 2011; Connallon and Clark 2014).

Third, X-Y recombination would introduce X-linked alleles onto the proto-Y

chromosomes. Male-limited inheritance of a Y chromosome is expected to lead to the

accumulation of male-beneficial alleles in Y-linked genes, even if they have female-deleterious

effects (Rice 1987, 1996). Conversely, female-biased inheritance of the X chromosome is

expected to result in stronger selection on X-linked alleles in females if those alleles are partially

dominant or segregating at moderate frequencies (Rice 1984; Charlesworth *et al.* 1987; Orr and

Betancourt 2001), potentially feminizing the X chromosome. If there is X-Y recombination, those

feminized X-linked alleles would be continuously introduced onto the Y chromosome, possibly

feminizing the Y as well (Cavoto *et al.* 2018). This could provide a biological mechanism that

allows for female-beneficial and male-detrimental effects of the proto-Y chromosomes that I

predict when additive fitness effects maintain PSD (Figure 5A; Supplemental Figure S16A).

Proto-sex chromosomes are reminiscent of pseudoautosomal regions, with some

notable differences

Proto-sex chromosomes have some superficial similarities to pseudoautosomal regions

(PARs) of heteromorphic sex chromosomes. The PAR is a region within which there is X-Y (or

Z-W) recombination, even if recombination is suppressed between the remainder of the X and Y

(Otto *et al.* 2011). Notably, PARs are capable of maintaining genetic variation, including sexually

antagonistic alleles, under broader conditions than autosomes or even non-recombining regions

of sex chromosomes (Jordan and Charlesworth 2012). There is evidence for sexually

antagonistic alleles in PARs from plants (Delph *et al.* 2010; Spigler *et al.* 2011) and fish (Tripathi

et al. 2009; Kitano *et al.* 2009), but it is not yet clear whether PARs are generally enriched for

sexually antagonistic alleles. Furthermore, the unique evolutionary dynamics of PARs is

generally limited to loci close to the boundary with the non-recombining region of the sex

chromosomes (Charlesworth *et al.* 2014).

Despite the superficial similarities, there are important differences that make PARs

imperfect analogs to proto-sex chromosomes. First, when there is an epistatic dominant female

determining proto-W chromosome (as in house fly), the proto-Y chromosome itself can be

transmitted through females. In contrast, the Y PAR cannot ever be carried by females, but

ancestrally Y-linked PAR alleles can recombine onto the X PAR and thus be carried by females.

Moreover, males can be homozygous for the proto-Y chromosome, which is superficially similar

to a male carrying an X PAR that has an ancestrally Y-linked allele. However, as above, this

ancestral Y allele on the X PAR must recombine onto the X PAR, whereas, when there is PSD,

homozygosity for the proto-Y occurs without X-Y recombination. Additional theoretical analyses

are required to directly compare the invasion dynamics, fixation probabilities, and regions of

stable polymorphisms across PARs and proto-sex chromosomes.

Low frequency proto-sex chromosomes are unlikely to be maintained by selection

My analysis suggests that it is difficult for selection to maintain proto-sex chromosomes

at low frequencies within populations. The III^M proto-Y is the lowest frequency proto-sex

chromosome I consider here, segregating at ≤2% in all three populations (Hamm *et al.* 2005;

Meisel *et al.* 2016). Selection pressures that maintain III^M at a frequency similar to those

observed in the populations for 1,000 generations allow for the loss of III^M within 1,000,000

generations (Supplemental Figures S8, S9). Additive fitness effects are especially likely to allow

for the loss of III^M. In addition, when PSD is maintained with proto-sex chromosomes at

frequencies similar to natural populations, III^M tends to be at a higher frequency in my

simulations than in any of the actual populations (Figure 2; Supplemental Figure S3). III^M is also

frequently lost when population size is finite, even when selection pressures exist that should

maintain the polymorphism (Supplemental Figures S11, S12). IV^F is also found at a low

frequency (2–9% of all fourth chromosomes in the natural populations); it is similarly found at a

higher frequency in the simulations than in the actual populations (Figure 2; Supplemental

Figure S3), and it is often lost in finite populations with selection (Supplemental Figures S11,

S12).

My results suggest that factors in addition to natural selection within populations may be

required for the maintenance of low frequency proto-sex chromosomes. Variation in proto-sex

chromosome frequencies across house fly populations are suggestive of what those factors

may be. For example, spatially variable, temporally fluctuating, and other heterogeneous

selection pressures can contribute to the maintenance of genetic variation across populations

(Levene 1953; Hedrick 2006), including maintaining PSD (Bateman and Anholt 2017). The

house fly Y^M and III^M proto-Y chromosomes are distributed along north-south clines on multiple

continents (Denholm *et al.* 1986; Tomita and Wada 1989; Hamm *et al.* 2005; Kozielska *et al.*

2008), suggesting spatially heterogeneous selection pressures may be important for maintaining

PSD across house fly populations. Migration between these populations would continuously

introduce rare proto-Y and proto-W chromosomes, potentially overwhelming selection pressures

that would otherwise allow for the loss of those proto-sex chromosomes within populations

(Lenormand 2002; Tigano and Friesen 2016). Future work should directly model the effect of

migration across populations with different selection pressures on the maintenance of PSD

within populations.

Generalizing to even more complex polygenic sex determination systems

I anticipate that many of the general patterns I observe here would be similar in other

complex PSD systems with at least three different proto-sex chromosomes. For example, in a

system with more than two proto-Y chromosomes, I expect to observe similar effects of

dominance on the maintenance of PSD and sexual antagonism as in the system I modeled with

two proto-Y chromosomes. In addition, if there are multiple proto-W chromosomes and a single

(epistatic) proto-Y (i.e., the opposite of the house fly system), I would expect similar patterns as

I observe in the house fly system, but with the sexes reversed. These expectations could be

tested in future work. It is not clear, however, what would be expected if there were both multiple

proto-W and multiple proto-Y chromosomes in a population. It is worth noting that polygenic

systems with many sex-determining loci are expected to be evolutionarily unstable (Rice 1986).

Zebrafish may represent a promising model organism for combining empirical and theoretical

approaches to make additional progress in understanding the maintenance of many sex

determining loci (Liew *et al.* 2012; Wilson *et al.* 2014).

Conclusions

My results demonstrate that the maintenance of PSD depends on the dominance of

fitness effects, which are further predictive of the sexually antagonistic effects of proto-sex

chromosomes and genotypes. Notably, overdominant fitness effects are most likely to maintain

PSD, but that conclusion requires at least one of two important assumptions: genetic variation

with overdominant fitness effects must exist in natural populations, or proto-Y chromosomes

must carry recessive deleterious alleles (in addition to additive or dominant beneficial alleles).

My results also suggest that sexual antagonistic multi-chromosomal genotypes should be most

prominent when dominant fitness effects (and to a lesser extent additive fitness effects) maintain

PSD. This sexual antagonism is an emergent property of multi-chromosomal genotypes, and

not intrinsic to the fitness effects of all individual proto-Y chromosomes. In contrast, when

overdominant fitness effects maintain PSD, the proto-Y chromosomes tend to be

male-beneficial and female-detrimental, but the multi-chromosomal genotypes do not have

opposing fitness effects in males and females. Lastly, future work is needed to evaluate how

X-Y (and Z-W) recombination, heterogeneous selection pressures, and migration across demes

affect the ability of sex-specific selection pressures to maintain complex PSD, including in

systems with other combinations of proto-sex chromosomes.

 Supplemental Material

Supplemental Figure S1. Smoothed
histograms show the distributions of mean
squared error (MSE) values for fitness arrays
that maintain PSD (i.e., all proto-sex
chromosomes at a frequency >0.1%). MSE is
calculated based on the frequencies of Y^M, III^M,
and IV^F found in either CA, NC, or NY.
Simulations were started with equal frequencies
of all genotypes. Fitness arrays are based on
additive, dominant, recessive, or overdominant
fitness effects of Y^M and III^M. **(A)** Distributions of
MSE are shown for all fitness arrays that
maintain PSD, with the percent of those arrays
that maintain PSD shown in each panel.
(B) Distributions of MSE are shown for the 1,000
best-fitting arrays with the lowest MSE.

Supplemental Figure S2. Smoothed
histograms show the distributions of mean
squared error (MSE) values for fitness arrays
that maintain PSD (i.e., all proto-sex
chromosomes at a frequency >0.1%). MSE is
calculated based on the frequencies of Y^M, III^M,
and IV^F found in either CA, NC, or NY.
Simulations were started with genotype
frequencies observed in natural populations.
Fitness arrays are based on additive, dominant,
recessive, or overdominant fitness effects of Y^M
and III^M. **(A)** Distributions of MSE are shown for
all fitness arrays that maintain PSD, with the
percent of those arrays that maintain PSD
shown in each panel. **(B)** Distributions of MSE
are shown for the 1,000 best-fitting arrays with
the lowest MSE.

Supplemental Figure S3. Smoothed histograms show the frequency of each proto-sex chromosome (Y^M,

III^M, and IV^F) after 1,000 generations in simulations using either 1,000 random fitness arrays (random),

fitness arrays that maintain PSD (polygenic), or the 1,000 best-fitting fitness arrays for each population

(CA, NC, or NY). Simulations were started with genotype frequencies observed in each population. The

vertical line within each histogram shows the median. Red dots show the observed proto-sex

chromosome frequencies in each natural population. Fitness arrays were calculated assuming either

(**A**) additive, (**B**) dominant, (**C**) recessive, or (**D**) overdominant fitness effects of the proto-Y

chromosomes.

Supplemental Figure S4. The frequency of fixation of the Y^M proto-Y chromosome after 1,000
generations (white) and 1,000,000 generations (gray) is shown for the 1,000 best-fitting genotype fitness
arrays (best-fit), 1,000 fitness arrays that maintain PSD (polygenic), and 1,000 randomly chosen fitness
arrays (random) in each population (CA, NC, or NY). Best-fitting fitness values and fitness values that
maintain PSD were identified using simulation in which all genotypes started at equal frequencies. Fitness
effects of each proto-Y chromosome were either additive, dominant, recessive, or overdominant.

Supplemental Figure S5. The frequency of fixation of the Y^M proto-Y chromosome after 1,000
generations (white) and 1,000,000 generations (gray) is shown for the 1,000 best-fitting genotype fitness
arrays (best-fit), 1,000 fitness arrays that maintain PSD (polygenic), and 1,000 randomly chosen fitness
arrays (random) in each population (CA, NC, or NY). Best-fitting fitness values and fitness values that
maintain PSD were identified using simulation in which all genotypes started at the frequencies observed
in each population. Fitness effects of each proto-Y chromosome were either additive, dominant,
842   recessive, or overdominant.

Supplemental Figure S6. The frequency of fixation of the III^M proto-Y chromosome after 1,000
generations (white) and 1,000,000 generations (gray) is shown for the 1,000 best-fitting genotype fitness
arrays (best-fit), 1,000 fitness arrays that maintain PSD (polygenic), and 1,000 randomly chosen fitness
arrays (random) in each population (CA, NC, or NY). Best-fitting fitness values and fitness values that
maintain PSD were identified using simulation in which all genotypes started at equal frequencies. Fitness
effects of each proto-Y chromosome were either additive, dominant, recessive, or overdominant.

Supplemental Figure S7. The frequency of fixation of the III^M proto-Y chromosome after 1,000
generations (white) and 1,000,000 generations (gray) is shown for the 1,000 best-fitting genotype fitness
arrays (best-fit), 1,000 fitness arrays that maintain PSD (polygenic), and 1,000 randomly chosen fitness
arrays (random) in each population (CA, NC, or NY). Best-fitting fitness values and fitness values that
maintain PSD were identified using simulation in which all genotypes started at the frequencies observed
in each population. Fitness effects of each proto-Y chromosome were either additive, dominant,
recessive, or overdominant.

Supplemental Figure S8. Histograms of the frequency of the III^M proto-Y chromosome after 1,000
generations are shown for the 1,000 fitness arrays that produce proto-sex chromosomes most similar to
those observed in each natural population (CA, NC, or NY). Fitness arrays are divided into those in which
the III^M chromosome eventually reaches a frequency <0.1% after 1,000,000 generations (lost) and those
in which the III^M chromosome is maintained at a frequency >0.1% (no). Fractions in each panel indicate
how often the III^M chromosome was lost after 1,000,000 generations in the 1,000 best-fitting arrays.
Simulations were started with equal frequencies of all genotypes, and fitness effects of proto-Y
863   chromosomes were either additive, dominant, recessive, or overdominant.

Supplemental Figure S9. Histograms of the frequency of the III^M proto-Y chromosome after 1,000

generations are shown for the 1,000 fitness arrays that produce proto-sex chromosomes most similar to

those observed in each natural population (CA, NC, or NY). Fitness arrays are divided into those in which

the III^M chromosome eventually reaches a frequency <0.1% after 1,000,000 generations (lost) and those

in which the III^M chromosome is maintained at a frequency >0.1% (no). Fractions in each panel indicate

how often the III^M chromosome was lost after 1,000,000 generations in the 1,000 best-fitting arrays.

Simulations were started with genotypes frequencies observed in each population, and fitness effects of

proto-Y chromosomes were either additive, dominant, recessive, or overdominant.

872 **Supplemental Figure S10.** Bar graphs show the frequency with which the Y^M chromosome is lost after
873 1,000,000 generation using fitness effects that maintain proto-sex chromosome frequencies similar to
874 those observed in each natural population (CA, NC, or NY). The best-fitting arrays were identified with
875 simulations starting with equal frequencies of all genotypes (equal starting freqs) or frequencies observed
876 in each population (location start freqs). Genotype fitness values were calculating assuming either
877 additive, dominant, recessive, or overdominant fitness effects.

Supplemental Figure S11. Histograms show the distributions of frequency of fixation or loss each
proto-Y or proto-W chromosome in each population when $N=10^4$. Numbers in each panel (n) indicate the
number of fitness arrays with at least one fixation in 100 simulations. The x-axis is the frequency of
fixations in 100 simulations for those arrays with at least one fixation. Simulations were performed with
the 1,000 best-fitting selection pressures for each population (CA, NC, or NY) that were identified from
simulations started with equal frequencies of all genotypes, using either **(A)** additive, **(B)** dominant,
(C) recessive, or **(D)** overdominant fitness effects of the proto-Y chromosomes.

Supplemental Figure S12. Histograms show the distributions of frequency of fixation or loss each
proto-Y or proto-W chromosome in each population when $N=10^4$. Numbers in each panel (n) indicate the
number of fitness arrays with at least one fixation in 100 simulations. The x-axis is the frequency of
fixations in 100 simulations for those arrays with at least one fixation. Simulations were performed with
the 1,000 best-fitting selection pressures for each population (CA, NC, or NY) that were identified from
simulations started with the observed frequencies of each genotype in each population, using either
(A) additive, **(B)** dominant, **(C)** recessive, or **(D)** overdominant fitness effects of the proto-Y
chromosomes.

Supplemental Figure S13. Bar plots show the number of fitness arrays that result in a deficiency (gray)
or excess (white) of fixations of each proto-sex chromosome (X, Y^M, III, III^M, or IV) relative to simulations
in which there are no fitness differences across proto-sex chromosomes (i.e., drift only). The fitness
arrays are the 1,000 best-fitting fitness values for each population (CA, NC, or NY) when simulations
were started with genotype frequencies observed in natural populations. Fitness effects of the proto-Y
chromosomes are either **(A)** additive, **(B)** dominant, **(C)** recessive, or **(D)** overdominant.

Supplemental Figure S14. Smoothed histograms show intersexual fitness correlations (ρ_{MF}) for the

1,000 best-fitting genotypic fitness arrays for the CA, NC, and NY populations (best), all fitness arrays

that maintain PSD (polygenic), and 1,000 random fitness arrays. Simulations were started with genotype

frequencies observed in natural populations. Dashed vertical lines show $\rho_{MF} = 0$, and solid vertical

lines within histograms show the median. Fitness arrays were calculated assuming either **(A)** additive,

(B) dominant, **(C)** recessive, or **(D)** overdominant fitness effects of the proto-Y chromosomes.

Supplemental Figure S15. Smoothed histograms show intersexual fitness correlations (ρ_{MF}) for the
1,000 best-fitting genotypic fitness arrays based on dominant fitness effects of the proto-Y chromosomes
for the CA, NC, and NY populations after 1,000,000 generations. Fitness arrays that maintain PSD for
1,000 generations are divided into those the maintain PSD for 1,00,000 generations (polygenic) and
those that allow at least one proto-sex chromosome to reach fixation (fixed). Simulations were started
with either equal frequencies of all genotypes (left) or genotype frequencies observed in natural
populations (right). Dashed vertical lines show $\rho_{MF} = 0$, and solid vertical lines within histograms show
912   the median.

Supplemental Figure S16. Smoothed histograms show the distributions of fitness effects of each
proto-sex chromosome in each sex for 1,000 random fitness arrays (random), fitness arrays that maintain
PSD (polygenic), or the 1,000 best-fitting fitness arrays for each population (CA, NC, or NY). Simulations
were started with genotype frequencies observed in natural populations. The vertical line within each
histogram shows the median, and dashed lines in each panel are at fitness value of 0. Fitness arrays
were calculated assuming either (**A**) additive, (**B**) dominant, (**C**) recessive, or (**D**) overdominant fitness
effects of the proto-Y chromosomes.

Supplemental Figure S17. Smoothed histograms show the distributions of the difference in the fitness
effect of each proto-Y chromosome (Y^M or III^M) between males and females within individual
parameterizations of the model. If the fitness effect is greater in males, the values are positive. If the
fitness effect is greater in females, the values are negative. Fitness effects of each proto-Y chromosome
come from the 1,000 random fitness arrays (random), fitness arrays that maintain PSD (polygenic), or the
1,000 best-fitting fitness arrays for each population (CA, NC, or NY). Simulations were started with equal
frequencies of all genotypes. The vertical line within each histogram shows the median, and dashed lines
in each panel are a difference of 0 (i.e., equal fitness effects in males and females). Genotype fitness
arrays were calculated assuming either (**A**) additive, (**B**) dominant, (**C**) recessive, or (**D**) overdominant
929   fitness effects of the proto-Y chromosomes.

Supplemental Figure S18. Smoothed histograms show the distributions of the difference in the fitness
effect of each proto-Y chromosome (Y^M or III^M) between males and females within individual
parameterizations of the model. If the fitness effect is greater in males, the values are positive. If the
fitness effect is greater in females, the values are negative. Fitness effects of each proto-Y chromosome
come from the 1,000 random fitness arrays (random), fitness arrays that maintain PSD (polygenic), or the
1,000 best-fitting fitness arrays for each population (CA, NC, or NY). Simulations were started with
genotype frequencies observed in natural populations. The vertical line within each histogram shows the
median, and dashed lines in each panel are a difference of 0 (i.e., equal fitness effects in males and
females). Genotype fitness arrays were calculated assuming either (**A**) additive, **(B)** dominant, **(C)**
939   recessive, or **(D)** overdominant fitness effects of the proto-Y chromosomes.

Supplemental Figure S19. Smoothed histograms show the distributions of the difference in the fitness
effect of each proto-Y chromosome (Y^M or III^M) between males and females within individual
parameterizations of the model. If the fitness effect is greater in males, the values are positive. If the
fitness effect is greater in females, the values are negative. Simulations were performed using 1,000
fitness arrays based on dominant fitness effects of the proto-Y chromosomes that maintain PSD for 1,000
generations. Fitness arrays that maintain PSD for 1,000 generations are divided into those in which the
proto-Y chromosome reaches fixation within 1,00,000 generations (fixed) and those in which it does not
reach fixation (no) in each population (CA, NC, or NY). Simulations were started with either equal
frequencies of all genotypes (left) or genotype frequencies observed in natural populations (right). The
vertical line within each histogram shows the median, and dashed lines in each panel are a difference of
0 (i.e., equal fitness effects in males and females).

Supplemental Figure S20. Bar plots show Spearman's rank order correlation between the frequency the IV^F proto-W chromosome across simulated populations and its fitness effect in the populations. Simulated populations include those that best-fit proto-sex chromosome frequencies in natural populations (CA, NC, or NY), those that maintain PSD (polygenic), and 1,000 random populations (random). Simulations were started with equal frequencies of all genotypes. Genotype fitness in populations was calculated assuming **(A)** additive, **(B)** dominant, **(C)** recessive, or **(D)** overdominant effects of the proto-Y chromosomes.

Supplemental Figure S21. Bar plots show Spearman's rank order correlation between the frequency the IV^F proto-W chromosome across simulated populations and its fitness effect in the populations. Simulated populations include those that best-fit proto-sex chromosome frequencies in natural populations (CA, NC, or NY), those that maintain PSD (polygenic), and 1,000 random populations (random). Simulations were started with the observed frequencies of each genotype in each population. Genotype fitness in populations was calculated assuming **(A)** additive, **(B)** dominant, **(C)** recessive, or **(D)** overdominant effects of the proto-Y chromosomes.

 Supplemental Table S1. Recursion equations to simulate random mating.

genotype	frequency	frequency after random mating
X/X; III/III; IV/IV	f_1	$f_1 m_1/2 + f_1 m_3/2 + f_1 m_4/4$ $+ f_2 m_1/4 + f_2 m_3/4 + f_2 m_4/8$ $+ f_3 m_1/4 + f_3 m_3/8 + f_3 m_4/16$ $+ f_5 m_1/8 + f_5 m_3/8 + f_5 m_4/16$ $+ f_6 m_1/16 + f_6 m_3/16 + f_6 m_4/32$
X/X; III/III; IV/IV$^{\mathrm{F}}$	f_2	$f_2 m_1/4 + f_2 m_3/4 + f_2 m_4/8$ $+ f_3 m_1/8 + f_3 m_3/8 + f_3 m_4/16$ $+ f_5 m_1/8 + f_5 m_3/8 + f_5 m_4/16$ $+ f_6 m_1/16 + f_6 m_3/16 + f_6 m_4/32$
X/X; III/III$^{\mathrm{M}}$; IV/IV$^{\mathrm{F}}$	f_3	$f_2 m_1/4 + f_2 m_2/2 + f_2 m_4/8 + f_2 m_5/4$ $+ f_3 m_1/4 + f_3 m_2/4 + f_3 m_3/8$ $+ f_3 m_4/8 + f_3 m_5/8$ $+ f_4 m_1/4 + f_4 m_3/4 + f_4 m_4/8$ $+ f_5 m_1/8 + f_5 m_2/4 + f_5 m_4/16 + f_5 m_5/8$ $+ f_6 m_1/8 + f_6 m_2/8 + f_6 m_3/16$ $+ f_6 m_4/16 + f_6 m_5/16$ $+ f_7 m_1/8 + f_7 m_3/8 + f_7 m_4/16$
X/X; III$^{\mathrm{M}}$/III$^{\mathrm{M}}$; IV/IV$^{\mathrm{F}}$	f_4	$f_3 m_1/8 + f_3 m_2/4 + f_3 m_4/16 + f_3 m_5/8$ $+ f_4 m_1/4 + f_4 m_2/2 + f_4 m_4/8 + f_4 m_5/4$ $+ f_6 m_1/16 + f_6 m_2/8 + f_6 m_4/32 + f_6 m_5/16$ $+ f_7 m_1/8 + f_7 m_2/4 + f_7 m_4/16 + f_7 m_5/8$
X/Y$^{\mathrm{M}}$; III/III; IV/IV$^{\mathrm{F}}$	f_5	$f_2 m_3/4 + f_2 m_4/8 + f_2 m_6/2 + f_2 m_7/4$ $+ f_3 m_3/8 + f_3 m_4/16 + f_3 m_6/4 + f_3 m_7/8$ $+ f_5 m_1/8 + f_5 m_3/4 + f_5 m_4/8$ $+ f_5 m_6/4 + f_5 m_7/8$ $+ f_6 m_1/16 + f_6 m_3/8 + f_6 m_4/16$ $+ f_6 m_6/8 + f_6 m_7/16$ $+ f_8 m_1/4 + f_8 m_3/4 + f_8 m_4/8$ $+ f_9 m_1/8 + f_9 m_3/8 + f_9 m_4/16$

X/Y$^\mathrm{M}$; III/III$^\mathrm{M}$; IV/IV$^\mathrm{F}$	f_6	$f_2m_4/8 + f_2m_5/4 + f_2m_7/4 + f_2m_8/2$

$$f_2m_4/8 + f_2m_5/4 + f_2m_7/4 + f_2m_8/2$$
$$+ f_3m_3/8 + f_3m_4/8 + f_3m_5/8$$
$$+ f_3m_6/4 + f_3m_7/4 + f_3m_8/4$$
$$+ f_4m_3/4 + f_4m_4/8 + f_4m_6/2 + f_4m_7/4$$
$$+ f_5m_1/8 + f_5m_2/4 + f_5m_4/8$$
$$+ f_5m_5/4 + f_5m_7/8 + f_5m_8/4$$
$$+ f_6m_1/8 + f_6m_2/8 + f_6m_3/8 + f_6m_4/8$$
$$+ f_6m_5/8 + f_6m_6/8 + f_6m_7/8 + f_6m_8/8$$
$$+ f_7m_1/8 + f_7m_3/4 + f_7m_4/8$$
$$+ f_7m_6/4 + f_7m_7/8$$
$$+ f_8m_1/4 + f_8m_2/2 + f_8m_4/8 + f_8m_5/4$$
$$+ f_9m_1/4 + f_9m_2/4 + f_9m_3/8$$
$$+ f_9m_4/8 + f_9m_5/8$$
$$+ f_{10}m_1/4 + f_{10}m_3/4 + f_{10}m_4/8$$

X/Y$^\mathrm{M}$; III$^\mathrm{M}$/III$^\mathrm{M}$; IV/IV$^\mathrm{F}$ f_7

$$f_3m_4/16 + f_3m_5/8 + f_3m_7/8 + f_3m_8/4$$
$$+ f_4m_4/8 + f_4m_5/4 + f_4m_7/4 + f_4m_8/2$$
$$+ f_6m_1/16 + f_6m_2/8 + f_6m_4/16$$
$$+ f_6m_5/8 + f_6m_7/16 + f_6m_8/8$$
$$+ f_7m_1/8 + f_7m_2/4 + f_7m_4/8$$
$$+ f_7m_5/4 + f_7m_7/8 + f_7m_8/4$$
$$+ f_9m_1/8 + f_9m_2/4 + f_9m_4/16 + f_9m_5/8$$
$$+ f_{10}m_1/4 + f_{10}m_2/2 + f_{10}m_4/8 + f_{10}m_5/4$$

Y$^\mathrm{M}$/Y$^\mathrm{M}$; III/III; IV/IV$^\mathrm{F}$ f_8

$$f_5m_3/8 + f_5m_4/16 + f_5m_6/4 + f_5m_7/8$$
$$+ f_6m_3/16 + f_6m_4/32 + f_6m_6/8 + f_6m_7/16$$
$$+ f_8m_3/4 + f_8m_4/8 + f_8m_6/2 + f_8m_7/4$$
$$+ f_9m_3/8 + f_9m_4/16 + f_9m_6/4 + f_9m_7/8$$

Y$^\mathrm{M}$/Y$^\mathrm{M}$; III/III$^\mathrm{M}$; IV/IV$^\mathrm{F}$ f_9

$$f_5m_4/16 + f_5m_5/8 + f_5m_7/8 + f_5m_8/4$$
$$+ f_6m_3/16 + f_6m_4/16 + f_6m_5/16$$
$$+ f_6m_6/8 + f_6m_7/8 + f_6m_8/8$$
$$+ f_7m_3/8 + f_7m_4/16 + f_7m_6/4 + f_7m_7/8$$
$$+ f_8m_4/8 + f_8m_5/4 + f_8m_7/4 + f_8m_8/2$$
$$+ f_9m_3/8 + f_9m_4/8 + f_9m_5/8$$
$$+ f_9m_6/4 + f_9m_7/4 + f_9m_8/4$$
$$+ f_{10}m_3/4 + f_{10}m_4/8 + f_{10}m_6/2 + f_{10}m_7/4$$

Y^M/Y^M; III^M/III^M; IV/IV^F	f_{10}	$f_6 m_4/32 + f_6 m_5/16 + f_6 m_7/16 + f_6 m_8/8$ $+ f_7 m_4/16 + f_7 m_5/8 + f_7 m_7/8 + f_7 m_8/4$ $+ f_9 m_4/16 + f_9 m_5/8 + f_9 m_7/8 + f_9 m_8/4$ $+ f_{10} m_4/8 + f_{10} m_5/4 + f_{10} m_7/4 + f_{10} m_8/2$
X/X; III/III^M; IV/IV	m_1	$f_1 m_1/2 + f_1 m_2/1 + f_1 m_4/4 + f_1 m_5/2$ $+ f_2 m_1/4 + f_2 m_2/2 + f_2 m_4/8 + f_2 m_5/4$ $+ f_3 m_1/8 + f_3 m_2/4 + f_3 m_3/8$ $+ f_3 m_4/8 + f_3 m_5/8$ $+ f_4 m_1/4 + f_4 m_3/4 + f_4 m_4/8$ $+ f_5 m_1/8 + f_5 m_2/4 + f_5 m_4/16 + f_5 m_5/8$ $+ f_6 m_1/8 + f_6 m_2/8 + f_6 m_3/16$ $+ f_6 m_4/16 + f_6 m_5/16$ $+ f_7 m_1/8 + f_7 m_3/8 + f_7 m_4/16$
X/X; III^M/III^M; IV/IV	m_2	$f_3 m_1/8 + f_3 m_2/4 + f_3 m_4/16 + f_3 m_5/8$ $+ f_4 m_1/4 + f_4 m_2/2 + f_4 m_4/8 + f_4 m_5/4$ $+ f_6 m_1/16 + f_6 m_2/8 + f_6 m_4/32 + f_6 m_5/16$ $+ f_7 m_1/8 + f_7 m_2/4 + f_7 m_4/16 + f_7 m_5/8$
X/Y^M; III/III; IV/IV	m_3	$f_1 m_3/2 + f_1 m_4/4 + f_1 m_6/1 + f_1 m_7/2$ $+ f_2 m_3/4 + f_2 m_4/8 + f_2 m_6/2 + f_2 m_7/4$ $+ f_3 m_3/8 + f_3 m_4/16 + f_3 m_6/4 + f_3 m_7/8$ $+ f_5 m_1/8 + f_5 m_3/4 + f_5 m_4/8$ $+ f_5 m_6/4 + f_5 m_7/8$ $+ f_6 m_1/16 + f_6 m_3/8 + f_6 m_4/16$ $+ f_6 m_6/8 + f_6 m_7/16$ $+ f_8 m_1/4 + f_8 m_3/4 + f_8 m_4/8$ $+ f_9 m_1/8 + f_9 m_3/8 + f_9 m_4/16$ $+ f_5 m_6/4 + f_5 m_7/8$ $+ f_6 m_1/16 + f_6 m_3/8 + f_6 m_4/16$ $+ f_6 m_6/8 + f_6 m_7/16$ $+ f_8 m_1/4 + f_8 m_3/4 + f_8 m_4/8$ $+ f_9 m_1/8 + f_9 m_3/8 + f_9 m_4/16$

X/Y$^{\text{M}}$; III/III$^{\text{M}}$; IV/IV	m_4	$f_1m_4/4 + f_1m_5/2 + f_1m_7/2 + f_1m_8/1$
		$+f_2m_4/8 + f_2m_5/4 + f_2m_7/4 + f_2m_8/2$
		$+f_3m_3/8 + f_3m_4/8 + f_3m_5/8$
		$+f_3m_6/4 + f_3m_7/4 + f_3m_8/4$
		$+f_4m_3/4 + f_4m_4/8 + f_4m_6/2 + f_4m_7/4$
		$+f_5m_1/8 + f_5m_2/4 + f_5m_4/8$
		$+f_5m_5/4 + f_5m_7/8 + f_5m_8/4$
		$+f_6m_1/8 + f_6m_2/8 + f_6m_3/8 + f_6m_4/8$
		$+f_6m_5/8 + f_6m_6/8 + f_6m_7/8 + f_6m_8/8$
		$+f_7m_1/8 + f_7m_3/4 + f_7m_4/8$
		$+f_7m_6/4 + f_7m_7/8$
		$+f_8m_1/4 + f_8m_2/2 + f_8m_4/8 + f_8m_5/4$
		$+f_9m_1/4 + f_9m_2/4 + f_9m_3/8$
		$+f_9m_4/8 + f_9m_5/8$
		$+f_{10}m_1/4 + f_{10}m_3/4 + f_{10}m_4/8$
X/Y$^{\text{M}}$; III$^{\text{M}}$/III$^{\text{M}}$; IV/IV	m_5	$f_3m_4/16 + f_3m_5/8 + f_3m_7/8 + f_3m_8/4$
		$+f_4m_4/8 + f_4m_5/4 + f_4m_7/4 + f_4m_8/2$
		$+f_6m_1/16 + f_6m_2/8 + f_6m_4/16$
		$+f_6m_5/8 + f_6m_7/16 + f_6m_8/8$
		$+f_7m_1/8 + f_7m_2/4 + f_7m_4/8$
		$+f_7m_5/4 + f_7m_7/8 + f_7m_8/4$
		$+f_9m_1/8 + f_9m_2/4 + f_9m_4/16 + f_9m_5/8$
		$+f_{10}m_1/4 + f_{10}m_2/2 + f_{10}m_4/8 + f_{10}m_5/4$
Y$^{\text{M}}$/Y$^{\text{M}}$; III/III; IV/IV	m_6	$f_5m_3/8 + f_5m_4/16 + f_5m_6/4 + f_5m_7/8$
		$+ f_6m_3/16 + f_6m_4/32 + f_6m_6/8 + f_6m_7/16$
		$+ f_8m_3/4 + f_8m_4/8 + f_8m_6/2 + f_8m_7/4$
		$+ f_9m_3/8 + f_9m_4/16 + f_9m_6/4 + f_9m_7/8$

Y^M/Y^M; III/III^M; IV/IV	m_7	$f_5 m_4/16 + f_5 m_5/8 + f_5 m_7/8 + f_5 m_8/4$
		$+ f_6 m_3/16 + f_6 m_4/16 + f_6 m_5/16$
		$+ f_6 m_6/8 + f_6 m_7/8 + f_6 m_8/8$
		$+ f_7 m_3/8 + f_7 m_4/16 + f_7 m_6/4 + f_7 m_7/8$
		$+ f_8 m_4/8 + f_8 m_5/4 + f_8 m_7/4 + f_8 m_8/2$
		$+ f_9 m_3/8 + f_9 m_4/8 + f_9 m_5/8$
		$+ f_9 m_6/4 + f_9 m_7/4 + f_9 m_8/4$
		$+ f_{10} m_3/4 + f_{10} m_4/8 + f_{10} m_6/2 + f_{10} m_7/4$
Y^M/Y^M; III^M/III^M; IV/IV	m_8	$f_6 m_4/32 + f_6 m_5/16 + f_6 m_7/16 + f_6 m_8/8$
		$+ f_7 m_4/16 + f_7 m_5/8 + f_7 m_7/8 + f_7 m_8/4$
		$+ f_9 m_4/16 + f_9 m_5/8 + f_9 m_7/8 + f_9 m_8/4$
		$+ f_{10} m_4/8 + f_{10} m_5/4 + f_{10} m_7/4 + f_{10} m_8/2$

References cited

Abbott, J. K., A. K. Nordén, and B. Hansson, 2017 Sex chromosome evolution: historical insights and future perspectives. Proc. R. Soc. B. 284: 20162806.

Allison, A. C., 1956 The sickle-cell and haemoglobin C genes in some African populations. Ann. Hum. Genet. 21: 67–89.

Andrés, A. M., M. J. Hubisz, A. Indap, D. G. Torgerson, J. D. Degenhardt *et al.*, 2009 Targets of balancing selection in the human genome. Mol. Biol. Evol. 26: 2755–2764.

Bachtrog, D., 2013 Y-chromosome evolution: emerging insights into processes of Y-chromosome degeneration. Nat. Rev. Genet. 14: 113–124.

Bachtrog, D., J. E. Mank, C. L. Peichel, M. Kirkpatrick, S. P. Otto *et al.*, 2014 Sex determination: why so many ways of doing it? PLoS Biol. 12: e1001899.

Barrett, S. C., and D. Charlesworth, 1991 Effects of a change in the level of inbreeding on the genetic load. Nature 352: 522–524.

Barson, N. J., T. Aykanat, K. Hindar, M. Baranski, G. H. Bolstad *et al.*, 2015 Sex-dependent dominance at a single locus maintains variation in age at maturity in salmon. Nature 528: 405–408.

Bateman, A. W., and B. R. Anholt, 2017 Maintenance of polygenic sex determination in a fluctuating environment: an individual-based model. J. Evol. Biol. 30: 915–925.

Beaumont, M. A., W. Zhang, and D. J. Balding, 2002 Approximate Bayesian computation in population genetics. Genetics 162: 2025–2035.

Bergero, R., and D. Charlesworth, 2009 The evolution of restricted recombination in sex chromosomes. Trends Ecol. Evol. 24: 94–102.

Beukeboom, L. W., and N. Perrin, 2014 *The Evolution of Sex Determination*. Oxford University

989  Press.

990 Bitarello, B. D., C. de Filippo, J. C. Teixeira, J. M. Schmidt, P. Kleinert *et al.*, 2018 Signatures of

991  long-term balancing selection in human genomes. Genome Biol. Evol. 10: 939–955.

992 Bonduriansky, R., and S. F. Chenoweth, 2009 Intralocus sexual conflict. Trends Ecol. Evol. 24:

993  280–288.

994 Brommer, J. E., M. Kirkpatrick, A. Qvarnström, and L. Gustafsson, 2007 The intersexual genetic

995  correlation for lifetime fitness in the wild and its implications for sexual selection. PLoS One

996  2: e744.

997 Brown, E. J., A. H. Nguyen, and D. Bachtrog, 2020 The Y chromosome may contribute to

998  sex-specific aging in *Drosophila*. Nat Ecol Evol 4: 853–862.

999 Bull, J. J., 1983 *Evolution of Sex Determining Mechanisms*. Benjamin/Cummings.

1000 Calsbeek, R., and B. Sinervo, 2004 Within-clutch variation in offspring sex determined by

1001  differences in sire body size: cryptic mate choice in the wild. J. Evol. Biol. 17: 464–470.

1002 Cavoto, E., S. Neuenschwander, J. Goudet, and N. Perrin, 2018 Sex-antagonistic genes, XY

1003  recombination and feminized Y chromosomes. J. Evol. Biol. 31: 416–427.

1004 Charlesworth, B., and D. Charlesworth, 2000 The degeneration of Y chromosomes. Philos.

1005  Trans. R. Soc. Lond. B Biol. Sci. 355: 1563–1572.

1006 Charlesworth, B., J. A. Coyne, and N. H. Barton, 1987 The relative rates of evolution of sex

1007  chromosomes and autosomes. Am. Nat. 130: 113–146.

1008 Charlesworth, B., C. Y. Jordan, and D. Charlesworth, 2014 The evolutionary dynamics of

1009  sexually antagonistic mutations in pseudoautosomal regions of sex chromosomes.

1010  Evolution 68: 1339–1350.

1011 Charlesworth, D., M. T. Morgan, and B. Charlesworth, 1990 Inbreeding depression, genetic

1012  load, and the evolution of outcrossing rates in a multilocus system with no linkage.

Evolution 44: 1469–1489.

Chippindale, A. K., J. R. Gibson, and W. R. Rice, 2001 Negative genetic correlation for adult
fitness between sexes reveals ontogenetic conflict in *Drosophila*. Proc. Natl. Acad. Sci. U.
S. A. 98: 1671–1675.

Clark, A. G., 1990 Two tests of Y chromosomal variation in male fertility of *Drosophila melanogaster*. Genetics 125: 527–534.

Clark, A. G., 1987 Variation in Y chromosome segregation in natural populations of *Drosophila melanogaster*. Genetics 115: 143–151.

Connallon, T., and S. F. Chenoweth, 2019 Dominance reversals and the maintenance of genetic variation for fitness. PLoS Biol. 17: e3000118.

Connallon, T., and A. G. Clark, 2014 Balancing selection in species with separate sexes: insights from Fisher's geometric model. Genetics 197: 991–1006.

Connallon, T., N. D. Singh, and A. G. Clark, 2012 Impact of genetic architecture on the relative rates of X versus autosomal adaptive substitution. Mol. Biol. Evol. 29: 1933–1942.

Crow, J. F., 1987 Muller, Dobzhansky, and overdominance. J. Hist. Biol. 20: 351–380.

Delph, L. F., A. M. Arntz, C. Scotti-Saintagne, and I. Scotti, 2010 The genomic architecture of sexual dimorphism in the dioecious plant *Silene latifolia*. Evolution 64: 2873–2886.

Delph, L. F., and J. K. Kelly, 2014 On the importance of balancing selection in plants. New Phytol. 201: 45–56.

Denholm, I., M. G. Franco, P. G. Rubini, and M. Vecchi, 1986 Geographical variation in house-fly (*Musca domestica* L.) sex determinants within the British Isles. Genet. Res. 47: 19–27.

Dobzhansky, T., 1947 Genetics of natural populations. XIV. A response of certain gene arrangements in the third chromosome of *Drosophila pseudoobscura* to natural selection.

Genetics 32: 142–160.

van Doorn, G. S., and M. Kirkpatrick, 2010 Transitions between male and female heterogamety

caused by sex-antagonistic selection. Genetics 186: 629–645.

van Doorn, G. S., and M. Kirkpatrick, 2007 Turnover of sex chromosomes induced by sexual

conflict. Nature 449: 909–912.

Dutoit, L., C. F. Mugal, P. Bolívar, M. Wang, K. Nadachowska-Brzyska *et al.*, 2018 Sex-biased

gene expression, sexual antagonism and levels of genetic diversity in the collared

flycatcher (*Ficedula albicollis*) genome. Mol. Ecol. 27: 3572–3581.

Fedorka, K. M., and T. A. Mousseau, 2004 Female mating bias results in conflicting sex-specific

offspring fitness. Nature 429: 65–67.

Feldmeyer, B., M. Kozielska, B. Kuijper, F. J. Weissing, L. W. Beukeboom *et al.*, 2008 Climatic

variation and the geographical distribution of sex-determining mechanisms in the housefly.

Evol. Ecol. Res. 10: 797–809.

Feldmeyer, B., I. Pen, and L. W. Beukeboom, 2010 A microsatellite marker linkage map of the

housefly, *Musca domestica*: evidence for male recombination. Insect Mol. Biol. 19:

575–581.

Felsenstein, J., 1974 The evolutionary advantage of recombination. Genetics 78: 737–756.

Fijarczyk, A., and W. Babik, 2015 Detecting balancing selection in genomes: limits and

prospects. Mol. Ecol. 24: 3529–3545.

Fisher, R. A., 1931 The evolution of dominance. Biol. Rev. Camb. Philos. Soc. 6: 345–368.

Foerster, K., T. Coulson, B. C. Sheldon, J. M. Pemberton, T. H. Clutton-Brock *et al.*, 2007

Sexually antagonistic genetic variation for fitness in red deer. Nature 447: 1107–1110.

Fry, J. D., 2010 The genomic location of sexually antagonistic variation: some cautionary

comments. Evolution 64: 1510–1516.

Gethmann, R. C., 1988 Crossing over in males of higher Diptera (Brachycera). J. Hered. 79:

344–350.

Grieshop, K., and G. Arnqvist, 2018 Sex-specific dominance reversal of genetic variation for

fitness. PLoS Biol. 16: e2006810.

Griffin, R. M., D. Le Gall, H. Schielzeth, and U. Friberg, 2015 Within-population Y-linked genetic

variation for lifespan in *Drosophila melanogaster*. J. Evol. Biol. 28: 1940–1947.

Hamm, R. L., 2008 Exploring the population genetics of the house fly sex determining genes, M

and F [Ph.D.]: Cornell University.

Hamm, R. L., R. P. Meisel, and J. G. Scott, 2015 The evolving puzzle of autosomal versus

Y-linked male determination in *Musca domestica*. G3 5: 371–384.

Hamm, R. L., and J. G. Scott, 2009 A high frequency of male determining factors in male *Musca*

domestica (Diptera: Muscidae) from Ipswich, Australia. J. Med. Entomol. 46: 169–172.

Hamm, R. L., and J. G. Scott, 2008 Changes in the frequency of Y^M versus III^M in the housefly,

Musca domestica L., under field and laboratory conditions. Genet. Res. 90: 493–498.

Hamm, R. L., T. Shono, and J. G. Scott, 2005 A cline in frequency of autosomal males is not

associated with insecticide resistance in house fly (Diptera: Muscidae). J. Econ. Entomol.

98: 171–176.

Hartl, D. L., and A. G. Clark, 2007 *Principles of Population Genetics*. Sinauer.

Hediger, M., G. Burghardt, C. Siegenthaler, N. Buser, D. Hilfiker-Kleiner *et al.*, 2004 Sex

determination in *Drosophila melanogaster* and *Musca domestica* converges at the level of

the terminal regulator *doublesex*. Dev. Genes Evol. 214: 29–42.

Hediger, M., C. Henggeler, N. Meier, R. Perez, G. Saccone *et al.*, 2010 Molecular

characterization of the key switch *F* provides a basis for understanding the rapid divergence

of the sex-determining pathway in the housefly. Genetics 184: 155–170.

Hedrick, P. W., 2006 Genetic polymorphism in heterogeneous environments: the age of

genomics. Annu. Rev. Ecol. Evol. Syst. 37: 67–93.

Hedrick, P. W., 2012 What is the evidence for heterozygote advantage selection? Trends Ecol.

Evol. 27: 698–704.

Innocenti, P., and E. H. Morrow, 2010 The sexually antagonistic genes of *Drosophila*

melanogaster. PLoS Biol. 8: e1000335.

Jordan, C. Y., and D. Charlesworth, 2012 The potential for sexually antagonistic polymorphism

in different genome regions. Evolution 66: 505–516.

Kallman, K. D., 1973 The sex-determining mechanism of the platyfish, *Xiphophorus maculatus*,

pp. 19–28 in *Genetics and Mutagenesis of Fish*, Springer Berlin Heidelberg.

Kidwell, J. F., M. T. Clegg, F. M. Stewart, and T. Prout, 1977 Regions of stable equilibria for

models of differential selection in the two sexes under random mating. Genetics 85:

171–183.

Kitano, J., J. A. Ross, S. Mori, M. Kume, F. C. Jones *et al.*, 2009 A role for a neo-sex

chromosome in stickleback speciation. Nature 461: 1079–1083.

Kozielska, M., B. Feldmeyer, I. Pen, F. J. Weissing, and L. W. Beukeboom, 2008 Are autosomal

sex-determining factors of the housefly (*Musca domestica*) spreading north? Genet. Res.

90: 157–165.

Kozielska, M., I. Pen, L. W. Beukeboom, and F. J. Weissing, 2006 Sex ratio selection and

multi-factorial sex determination in the housefly: a dynamic model. J. Evol. Biol. 19:

879–888.

Lemos, B., L. O. Araripe, and D. L. Hartl, 2008 Polymorphic Y chromosomes harbor cryptic

variation with manifold functional consequences. Science 319: 91–93.

Lemos, B., A. T. Branco, and D. L. Hartl, 2010 Epigenetic effects of polymorphic Y

chromosomes modulate chromatin components, immune response, and sexual conflict.

Proc. Natl. Acad. Sci. U. S. A. 107: 15826–15831.

Lenormand, T., 2002 Gene flow and the limits to natural selection. Trends Ecol. Evol. 17:

183–189.

Levene, H., 1953 Genetic equilibrium when more than one ecological niche is available. Am.

Nat. 87: 331–333.

Liew, W. C., R. Bartfai, Z. Lim, R. Sreenivasan, K. R. Siegfried *et al.*, 2012 Polygenic sex

determination system in zebrafish. PLoS One 7: e34397.

Lindholm, A., and F. Breden, 2002 Sex chromosomes and sexual selection in poeciliid fishes.

Am. Nat. 160 Suppl 6: S214–24.

McDonald, I. c., P. Evenson, C. A. Nickel, and O. A. Johnson, 1978 House fly genetics: isolation

of a female determining factor on chromosome 4. Ann. Entomol. Soc. Am. 71: 692–694.

Meier, N., S. C. Käppeli, M. Hediger Niessen, J.-C. Billeter, S. F. Goodwin *et al.*, 2013 Genetic

control of courtship behavior in the housefly: evidence for a conserved bifurcation of the

sex-determining pathway. PLoS One 8: e62476.

Meisel, R. P., T. Davey, J. H. Son, A. C. Gerry, T. Shono *et al.*, 2016 Is multifactorial sex

determination in the house fly, *Musca domestica* (L.), stable over time? J. Hered. 107:

615–625.

Meisel, R. P., C. A. Gonzales, and H. Luu, 2017 The house fly Y Chromosome is young and

minimally differentiated from its ancient X Chromosome partner. Genome Res. 27:

1417–1426.

Moore, E. C., and R. B. Roberts, 2013 Polygenic sex determination. Curr. Biol. 23: R510–2.

Muller, H. J., 1964 The relation of recombination to mutational advance. Mutat. Res. 106: 2–9.

Orr, H. A., 2010 The population genetics of beneficial mutations. Philos. Trans. R. Soc. Lond. B

Biol. Sci. 365: 1195–1201.

Orr, H. A., and A. J. Betancourt, 2001 Haldane's sieve and adaptation from the standing genetic

variation. Genetics 157: 875–884.

Orzack, S. H., J. J. Sohn, K. D. Kallman, S. A. Levin, and R. Johnston, 1980 Maintenance of the

three sex chromosome polymorphism in the platyfish *Xiphophorus maculatus*. Evolution 34:

663–672.

Otto, S. P., J. R. Pannell, C. L. Peichel, T.-L. Ashman, D. Charlesworth *et al.*, 2011 About PAR:

the distinct evolutionary dynamics of the pseudoautosomal region. Trends Genet. 27:

358–367.

Rice, W. R., 1996 Evolution of the Y sex chromosome in animals. Bioscience 46: 331–343.

Rice, W. R., 1986 On the instability of polygenic sex determination: the effect of sex-specific

selection. Evolution 40: 633–639.

Rice, W. R., 1984 Sex chromosomes and the evolution of sexual dimorphism. Evolution 38:

735–742.

Rice, W. R., 1992 Sexually antagonistic genes: experimental evidence. Science 256:

1436–1439.

Rice, W. R., 1987 The accumulation of sexually antagonistic genes as a selective agent

promoting the evolution of reduced recombination between primitive sex chromosomes.

Evolution 41: 911–914.

Rice, W. R., and A. K. Chippindale, 2001 Intersexual ontogenetic conflict. J. Evol. Biol. 14:

685–693.

Roberts, R. B., J. R. Ser, and T. D. Kocher, 2009 Sexual conflict resolved by invasion of a novel

sex determiner in Lake Malawi cichlid fishes. Science 326: 998–1001.

Rohmer, C., J. R. David, B. Moreteau, and D. Joly, 2004 Heat induced male sterility in

Drosophila melanogaster: adaptive genetic variations among geographic populations and

role of the Y chromosome. J. Exp. Biol. 207: 2735–2743.

Ruzicka, F., M. S. Hill, T. M. Pennell, I. Flis, F. C. Ingleby *et al.*, 2019 Genome-wide sexually

antagonistic variants reveal long-standing constraints on sexual dimorphism in fruit flies.

PLoS Biol. 17: e3000244.

Scott, J. G., C. A. Leichter, F. D. Rinkevich, S. A. Harris, C. Su *et al.*, 2013 Insecticide resistance

in house flies from the United States: resistance levels and frequency of pyrethroid

resistance alleles. Pestic. Biochem. Physiol. 107: 377–384.

Scott, J. G., W. C. Warren, L. W. Beukeboom, D. Bopp, A. G. Clark *et al.*, 2014 Genome of the

house fly, *Musca domestica* L., a global vector of diseases with adaptations to a septic

environment. Genome Biol. 15: 466.

Sellis, D., B. J. Callahan, D. A. Petrov, and P. W. Messer, 2011 Heterozygote advantage as a

natural consequence of adaptation in diploids. Proc. Natl. Acad. Sci. U. S. A. 108:

20666–20671.

Sellis, D., D. J. Kvitek, B. Dunn, G. Sherlock, and D. A. Petrov, 2016 Heterozygote advantage is

a common outcome of adaptation in *Saccharomyces cerevisiae*. Genetics 203: 1401–1413.

Sharma, A., S. D. Heinze, Y. Wu, T. Kohlbrenner, I. Morilla *et al.*, 2017 Male sex in houseflies is

determined by *Mdmd*, a paralog of the generic splice factor gene *CWC22*. Science 356:

642–645.

Spencer, H. G., and N. K. Priest, 2016 The Evolution of Sex-Specific Dominance in Response to

Sexually Antagonistic Selection. Am. Nat. 187: 658–666.

Spigler, R. B., K. S. Lewers, and T.-L. Ashman, 2011 Genetic architecture of sexual dimorphism

in a subdioecious plant with a proto-sex chromosome. Evolution 65: 1114–1126.

Tigano, A., and V. L. Friesen, 2016 Genomics of local adaptation with gene flow. Mol. Ecol. 25:

2144–2164.

Tomita, T., and Y. Wada, 1989 Multifactorial sex determination in natural populations of the

    housefly (*Musca domestica*) in Japan. Jpn. J. Genet. 64: 373–382.

Tripathi, N., M. Hoffmann, E.-M. Willing, C. Lanz, D. Weigel *et al.*, 2009 Genetic linkage map of

    the guppy, Poecilia reticulata, and quantitative trait loci analysis of male size and colour

    variation. Proc. Biol. Sci. 276: 2195–2208.

Wilkins, A. S., 1995 Moving up the hierarchy: a hypothesis on the evolution of a genetic sex

    determination pathway. Bioessays 17: 71–77.

Wilson, C. A., S. K. High, B. M. McCluskey, A. Amores, Y.-L. Yan *et al.*, 2014 Wild sex in

    zebrafish: loss of the natural sex determinant in domesticated strains. Genetics 198:

    1291–1308.

Wright, A. E., R. Dean, F. Zimmer, and J. E. Mank, 2016 How to make a sex chromosome. Nat.

    Commun. 7: 12087.

Zhang, P., B. Timakov, R. L. Stankiewicz, and I. Y. Turgut, 2000 A trans-activator on the

    Drosophila Y chromosome regulates gene expression in the male germ line. Genetica 109:

    141–150.