1   Title: Urban living influences the reproductive success of Darwin's finches in the Galápagos Islands

3   JOHANNA A. HARVEY[1,2*], KILEY CHERNICKY[1], SHELBY R. SIMONS[1],

4   TAYLOR B. VERRETT[1], JAIME A. CHAVES[3,4], and SARAH A. KNUTIE[1,5]

[1]Department of Ecology and Evolutionary Biology, University of Connecticut, Storrs, CT, USA

[2]Current address: Division of Invertebrate Zoology, American Museum of Natural History, New York,

NY, 10024, USA

[3]Colegio de Ciencias Biológicas y Ambientales, Campus Cumbayá, Universidad San Francisco de Quito,

Diego de Robles y Av. Interoceánica, Cumbayá, Quito, Ecuador

[4]Department of Biology, San Francisco State University, 1600 Holloway Ave, San Francisco, CA, 94132,

USA

[5]Institute for Systems Genomics, University of Connecticut, Storrs, CT, 06269, USA.

*Corresponding author: johannaaharvey@gmail.com

Abstract

1. Urbanization is expanding worldwide and can have major consequences for organisms, anthropogenic factors can reduce the fitness of animals but may also have benefits, such as consistent human food availability. Understanding these trade-offs is critically important in environments with unreliable annual natural food availability, such as the Galápagos Islands where urbanization is rapidly increasing. For example, during dry climatic condition years, the reproductive success of bird species, such as Darwin's finches, is low because low precipitation reduces food availability. Urban areas in the Galápagos provide supplemental human food to finches, which could improve their reproductive success during years with low natural food availability. However, urban finches might face trade-offs, as the incorporation of anthropogenic debris (e.g. string, hair, plastic) into their nests can increase mortality.

2. In our study, we determined the effect of urbanization on the reproductive effort and success of small ground finches (*Geospiza fuliginosa*; a species of Darwin's finch) during a dry year on San Cristóbal Island, Galápagos Islands. We also documented the abundance of anthropogenic debris incorporated in to nests.

3. We quantified nest building, egg laying, hatching, and fledging of small ground finches in an urban and non-urban area. We also qualified the type of anthropogenic debris in finch nests and the quantified the percent of debris comprising total nest mass. We determined whether incorporating these materials into the nest directly led to entanglement- or ingestion-related mortalities.

4. Overall, urban finches built more nests, laid more eggs, and produced more fledglings than non-urban finches. However, every nest in the urban area contained anthropogenic-related material, which resulted in entanglement- or ingestion-related mortalities in 18% of nests with nestlings. Non-urban nests did not contain any anthropogenic-related material.

5. Our study showed that urban living has trade-offs during drier climatic conditions: urban birds have overall higher reproductive success than non-urban birds, but urban birds can also suffer a

negative consequence by using anthropogenic-related material for nesting. These results suggest

that despite the potential cost of urban living, finches benefit overall from urban living and

urbanization may buffer the effects of limited resource availability in the Galápagos Islands.

Keywords: climate change, dry year, entanglement, Galápagos Islands, plastic, trash, urban ecology

1. Introduction

Urbanization is increasing worldwide with fewer pristine places remaining untouched by humans

(Vitousek, 1997). Urbanization can directly change the physical structure of the ecosystem through the

creation of roads and buildings and introducing artificial light, pollution, and noise (Fernández-Juricic,

2002; Dominoni, Quetting, & Partecke, 2013; Herrera-Dueñas et al., 2017). Consequently, native fauna

can suffer reduced fitness or extirpation in response to urbanization-related stressors, such as

environmental change, increased predation, limited natural food availability, and increased disease and

parasites (Blair, 1996; Lepczyk, Mertig, & Liu, 2004; Bailly et al., 2016; Johnson & Munshi-South,

2017). However, urban living can also benefit organisms by reducing the natural predation risk and

increasing alternative resource availability, such as human food sources and habitat structures (Gering &

Blair, 1999; Lowry, Lill, & Wong, 2013; Møller et al., 2015). The effect of anthropogenic materials on

seabirds has been well examined (Roman et al., 2019); however, the impacts on passerines, particularly

on reproductive success in urban areas, have not been well assessed. The effect of urbanization on birds

can vary, but include earlier lay dates and lower reproductive success in urban versus non-urban areas

(Chamberlain et al., 2009; Sepp et al., 2018). Urban food availability has been suggested as a principal

factor driving the variation of demographic responses across passerines (Chamberlain et al., 2009). While

species diversity can decline in urban areas (Kark et al., 2007), urban areas still sustain a number of native

species (Aronson et al., 2014); this duality presents an opportunity to better understand the trade-offs

experienced by a species in response to urbanization.

Determining the effects of urbanization on islands is especially important given that islands host

higher human population densities, as compared to the mainland, and harbor 20% of all terrestrial plant

and vertebrate species diversity (Kier et al., 2009; Courchamp et al., 2014). Furthermore, island endemic

species, particularly specialist species, can be highly sensitive to natural and anthropogenic perturbations

(Buckley & Jetz, 2007) due to their small population sizes, low immigration, and associated genetic

factors, such as low genetic diversity (Benning et al., 2002). However, the effect of urbanization on island

species can vary. For example, native Caribbean island reptiles can persist in urban environments, but in

lower numbers (Jesse et al., 2018). In contrast, urban adapted dark-eyed juncos have longer breeding

seasons, which result in higher reproductive success (Yeh & Price, 2004). These few studies suggest that

native species can differ in their responses to urbanization. As island species face extinction threats on

many fronts, examining trade-offs for urban animals on islands could provide insight into their ability to

respond to anthropogenic change or help inform management and conservation of the species.

The Galápagos Islands of Ecuador has experienced recent urbanization due in large part to growth

in ecotourism. Since the 1990s, Galápagos tourism has increased by an average of 9.4% per year, with

current estimates of nearly 225,000 visiting tourists each year. The resident human population has

increased by an average of 6.4% per year since the early 1990's, reaching 25,244 in 2015 (Epler, 2007;

Walsh & Mena, 2016). The recent human population growth and the incipient urbanization of the

Galápagos islands provides an ideal "laboratory" to determine the effects of human activity on endemic

animals. For example, recent studies have shown that Darwin's finches in urban areas prefer non-natural

food compared to finches in non-urban areas (De León et al., 2018), which consequently resulted in

changes to their microbiota (Knutie, Chaves, & Gotanda, 2019), epigenetics (McNew et al., 2017), and

morphology (Hendry et al., 2006; De Léon et al., 2011).

The Galápagos also face natural stressors, such as highly variable climatic conditions. The islands

have a hot, wetter season from approximately January to May, and a cool, drier season from

approximately June to December (P. R. Grant & Boag, 1980). The conditions during these seasons

depend on the Inter-Tropical Convergence Zone (ITCZ) and the periodically irregular El Niño Southern

Oscillation (ENSO) (Trueman & d'Ozouville, 2010). El Niño events can often result in wetter seasons

with high primary productivity and therefore high food resources for the finches, whereas La Niña events

are characterized by drier seasons with limited primary productivity and food resources (P. R. Grant &

Boag, 1980; Trueman & d'Ozouville, 2010). Consequently, low reproductive fitness has been a consistent

documented pattern reported in Darwin's finches across the Galapagos islands in dry La Niña years, with

this effect being more pronounced in the arid coastal zones (Boag & Grant, P.R., 1981; Gibbs & Grant,

P.R., 1987; Grant & Grant, 1989, 1999; Koop, LeBohec, & Clayton, 2013). Low reproductive success in

response to dry years has also been found in other island land birds, such as Galápagos mockingbirds

(Curry & Grant, P.R., 1989; McNew et al., 2019). To date, studies have not examined the influence of

urbanization on the reproductive fitness of Darwin's finches during dry conditions, Therefore, the

Galápagos islands present a unique opportunity to examine the effects of growing, yet incipient,

urbanization in a landscape where climate could be further exacerbating the positive or negative effects of

urbanization on an endemic species.

In our study, we examined the effect of urbanization on the reproductive effort and success of

small ground finches (*Geospiza fuliginosa*; a species of Darwin's finch) during a year with dry climate

conditions. First, we determined whether reproductive effort (i.e. nests built, eggs laid, hatchlings) and

success (i.e. young fledged) of small ground finches differed between urban and non-urban areas by

tracking the survival of nests from construction to egg laying, hatching, nestling survival, and confirmed

fledging of young. During years with dry conditions, the reproductive effort and success of Darwin's

finches is lower than years with wet conditions, which has been linked to reduced natural food availability

(Boag & Grant, P.R., 1981; Koop, LeBohec, & Clayton, 2013). Because urban areas are supplemented

with additional human food resources (De León et al., 2018), finch reproductive effort and success is

predicted to increase in urban areas compared to non-urban areas. However, finches incorporate human-

related material into their nest (Knutie et al., 2014), which is readily available in urban areas and can

result in death due to entanglement (Townsend & Barker, 2014; Jagiello et al., 2018; Jiguet et al., 2019,

Theodosopoulos & Gotanda, 2019). Therefore, although urban finches are predicted to have higher

overall reproductive success, urban finches likely face a trade-off related to anthropogenic debris in their

nests: urban finches are predicted to incorporate more anthropogenic debris into their nests and suffer

higher mortality related to this debris (i.e. entanglement, ingestion) than non-urban finches.

2. Materials and Methods

2.1 Study system

We conducted our study between February – May 2018 (during the breeding season) in the arid

lowland climatic zone of San Cristóbal (557 km^2) in the Galápagos Islands. Breeding for ground finches

is initiated by heavy rainfall events and continued breeding is dependent on continued rainfall (Boag &

Grant, P.R., 1984; S. Kleindorfer, 2007). Rainfall on San Cristóbal is highly variable, with interannual

variation alternating between high and low rainfall (P. R. Grant & Boag, 1980).

We quantified nest building, egg laying, hatching, and fledging of small ground finches in an

urban and non-urban area. The urban area was in the capital city of Puerto Baquerizo Moreno (hereon,

urban area), which is the second largest city in the Galápagos archipelago with a human population of

7,199 (INEC, 2016). The urban area primarily consists of impermeable concrete or stone surfaces and

human built structures in altered landscapes. Our urban study area measured 0.79 km^2 (~1.2 km by 0.62

km) and included tourist and residential zones (Fig. 1C). The search area within the urban area was

determined due to suitability of habitat, i.e. presence of trees for nesting and accessibility, permission to

search and access granted to private properties. The non-urban area was in the Jardin de Opuntias (hereon,

non-urban area), which is a Galápagos National Park site located 4.5 km southeast of the urban area

consisting of vegetated natural habitats with no unnatural impermeable surfaces present. Our non-urban

study area measured 0.21 km^2 and covered 1.4 km of the main trail and 0.15 km to each side (Fig. 1D).

The search area is larger in the urban area than the non-urban area due to spatial mismatch and differences

in environmental structure, which can result in urban patches devoid of suitable nesting areas. Search

efforts, via person-hours (total number of hours searched per person for each day), were tracked for each

study area to normalize search efforts. Our search area was primarily, but not exclusively centered, by the

presence of the arboreal cactus, *Opuntia megasperma*, which are one of the preferred nesting locations of

small ground finches. Cacti are rare on San Cristóbal, likely due to destruction by introduced mammals in

the 1800s, but are found in high abundance within the Jardin de Opuntias (Phillips, Wiedenfeld, & Snell,

2012; Dvorak et al., 2019). Small ground finch nests are commonly found in cacti as well as trees such as

matazarno (*Piscidia cathagenensi*) and Galápagos acacia (*Acacia rorudiana*). The non-urban area

receives very low human visitation: locals occasionally, but rarely, visit the site to access the beach. Cacti

are also frequently cultivated in urban areas, and therefore are found in garden beds, planters, city parks,

and the main boardwalk. Urban finches also nest in native and non-native trees and occasionally nest in

human built structures, such as gutters and building signs.

Fig. 1. (A) Map of the major islands of the Galápagos archipelago and inset map of (B) San Cristóbal

Island with sampling areas noted (black dots) for the urban area (Puerto Baqueizo Moreno) and the non-

urban area (Jardin de Opuntias). Satellite maps of the (C) urban area and (D) the non-urban area showing

nests which were builds only (grey circle), nests with eggs that failed (grey square), nests with nestlings

that failed (dark grey triangle), and nests with nestlings that fledged (black circle) across each sampling

site with the search area delineated by the black polygon border. Map data Google Maps Imagery © 2019

and the GADM database (Hijmans et al., 2014).

2.2 Locating nest sites and data collection

164   In each urban and non-urban area, we searched intensively for nests and for small ground finches

exhibiting nest-building behaviors, including vocalization and behavioral cues. The field sites were

searched nearly every other day for evidence of nest-building activity by small ground finches. We

followed all nests accessible with the use of a 10-foot (~3 meter) ladder. Once found, nests were checked

every other day until eggs were found in the nest. Observations were made primarily through binoculars

to minimize nest disturbance, and secondarily through a small camera (Contour LLC, Provo, USA)

attached to an extendable pole when the nest was not attended by adults. Once the eggs hatched, we

followed the survival of nestlings and banded them with a unique color band combination when they were

7-8 days of age (hatch date = day 0). Successful fledging was confirmed by identifying individual birds

once they left the nest, as in previous studies (Knutie et al., 2016). After nestling birds fledged or died, the

nest was collected and placed in a sealed plastic bag. Each nest was carefully dissected to separate natural

and anthropogenic materials, after which each material type was weighed (g). Anthropogenic materials

were then qualitatively identified (composition of material, possible source, and color of materials) when

possible. All detected nest failures and mortalities were documented, and causation was determined when

possible. Materials associated with mortality via ingestion or entanglement were also identified and

documented.

180   Our study resulted from a single year of sampling, and no building or breeding individuals in our

study had been previously banded. We did not observe any dispersal of banded birds across urban and

non-urban areas. The flight distance between urban and non-urban sampling areas is eight km, and

previous studies have not found dispersal in small ground finches to occur across habitats (Sonia

Kleindorfer et al., 2006). Therefore, it is unlikely that small ground finches disperse between the urban

and non-urban area of this study at significant rates.

2.3 Statistical analyses

All data were analyzed in Rstudio v1.2 (R Core Team, 2012). We calculated total person-hours

by multiplying the number of hours searched by the number of people searching for each day at both

sites (urban and non-urban respectively) across all days of the survey period. We tested survey effort

using an independent t-test on total person-hours per day at each site after examining data for

homogeneity of variance using a Fligner-Killeen test and Q-Q plots for assessment of normality. For

effect size, we calculated Cohen's D (Cohen, 2013) for percent of nestlings fledged per nest across

sampling areas and percent nest mass of anthropogenic debris across sampling areas, respectively.

Cohen's D is a standardized measure of difference in means divided by the pooled standard deviation,

and it was calculated using the lsr package (Navarro & Foxcroft,).

We used generalized linear models (GLMs) using the glm function in the lme4 package (Bates et

al., 2015) and ANOVAs using the car package (Fox & Weisberg, 2011). We first verified that data met

assumptions of models by checking for overdispersion and underdispersion. We used GLMs with a

binomial error structure to determine whether location (urban and non-urban) affected the binary

presence or absence of nests with eggs, nests with nestlings, and trash in nest material. Additionally, we

used binomial logistic regressions, which are a special case of GLM to determine the effect of location

on hatching and nestling survival trials, independently. The response variable is a matrix of trials, which

are successes and failures, where the number of eggs hatched and nestling fledged per nest represent

successes while eggs not hatched and nestling mortalities represent failures, respectively. We also used a

binomial logistic regression to determine the effect of location along with the proportion of

anthropogenic debris in the nest on nestling survival, where nest survival was measured as successes and

failures.

We used a Mayfield logistic regression, an extension of the Mayfield method, which examines

nest survival data over the incubation and nestling periods and incorporates explanatory variables in a

logistic regression (Hazler, 2004). Nest fates (success or failure) are modeled over the number of

exposure days where each exposure day is a trial. The explanatory variables are covariates of location

(urban and non-urban) and proportion of anthropogenic debris in nests which are also weighted by the

number of exposure days (Hazler, 2004). Exposure days were calculated using the median of the last day

the nest was known active to the last check day when the nest outcome or fate was known (Mayfield,

1961).

3. Results

Nest searching was conducted over 49 days in the urban area with an average of 12.51 ± 4.21

person-hours search day. Nest searching was conducted over 40 days in the non-urban area with an

average of 12.23 ± 7.5 person-hours per search day. The number of person-hours spent searching did not

significantly vary across the urban and non-urban study areas (Independent t-test, t = -0.22, $df = 75.51$, P

= 0.82).

The first urban finch nest with eggs was found on February 10, 2018, whereas the first non-urban

finch nest with eggs was found on March 7, 2018 (Fig. 2). The last urban nest with eggs was found on

April 14, 2018 and the last non-urban finch nest with eggs was found on April 13, 2018, resulting in a

breeding season of 68 days in the urban area and 37 days in the non-urban area. Small ground finches

built 29 nests in the urban area and 29 nests in the non-urban area (Fig. 1, Table 1). The urban area had

more nests with eggs (n = 25 nests with eggs out of 29 nests built) than in the non-urban area (n = 12

nests with eggs out of 29 nests built) (GLM, $\chi^2 = 13.28$, $df = 1$, $P < 0.0001$).

Fig. 2. Histograms of reproductive activity noting frequency of nests with (A) eggs, (B) nestlings, and (C) fledgling (where fledging occurred that week) across each week of the study period in non-urban (grey) and urban (black) sampling areas.

233   **Table 1.** Reproductive success of small ground finches in non-urban and urban areas.

Variable	Non-urban	Urban
No. built	29	29
No. with eggs	12/29 (41.3%)	25/29 (86.2%)
No. with nestlings	6/12 (50.0%)	17/25(68%)
No. with at least one fledgling	2/6 (33.3%)	15/17 (88.2%)
No. with anthropogenic materials	0/10 (0%)	22/22 (100%)

235
236
The urban area had more nests with hatchlings (n = 16) from the total nests built (n = 29)

compared to the non-urban area (n = six nests with hatchlings from 29 nests built) (GLM, χ^2 = 9.0, df =

1, P = 0.003). Of the 25 urban nests with eggs, eight failed at the egg stage. Three nests were damaged

during likely predation events, with no remains found and the nest entrance destroyed. Two nests were

found abandoned with eggs intact and cold. One nest had only one egg and was found predated by ants.

One nest was found with a single egg ejected and the remaining eggs found cold in the unattended nest.

One nest was found empty with no evidence of predation or cause of failure determined. Of the 12 non-

urban nests with eggs, six failed at the egg stage. One nest had clear signs of predation with a broken egg

shell found outside of the nest. One egg nest was found empty with no evidence to explain its failure.

The remaining four failed egg nests were abandoned, with cold eggs found in the unattended nest.

Overall survival, from egg stage and fledging, was higher in the urban compared to the non-

urban area: urban nests had a higher proportion of nests with eggs that hatched (χ^2 = 4.34, df = 1, P =

0.04) and nestlings that fledged (χ^2 = 14.35, *df* = 1, *P* = 0.0002, *Cohen's D* = 1.48) than the non-urban

nests. Only two urban nest failures occurred during the nestling stage. One nest was found with a six-day

old dead nestling hanging from ingested hair that was woven into the nest (Fig. 3A) and the two other

nestlings missing from the nest. One nest was found empty with no apparent cause for mortality or

nestling remains found. For three out of the four nest failures during the nestling stage in the non-urban

area, nestlings were missing with no clear signs of depredation. These mortalities occurred post-hatching

at days six, seven, and ten days old. One non-urban nest had signs of depredation, with partial nestling

remains (i.e. limbs and skull) found near the nest.

Fig. 3. Documented small ground finch mortalities due to ingestion/entanglement of anthropogenic nest

materials: a. 6-day old nestling ingestion/entanglement with plastic and human hair, b. 12-day old nestling

entanglement with synthetic string, c. adult female entanglement during nest building with human hair).

Fig. 4. Identification of anthropogenic materials dissected (a. shredded plastic tarp strands, b. cigarette fibers, c. fibers/thread, d. fiberglass from fishing boats, e. caution tape, f. paper shreds, g. human hair, h. shredded twine, i. synthetic stuffing) from a single *G. fuliginosa* urban nest which were sorted by material type and color (central figure, labeled with lower case) with potential urban source materials shown (below and above, labeled with upper case).

Anthropogenic debris was not found in non-urban areas, whereas all nests in the urban area

contained anthropogenic debris (*Cohen's D* = 2.140, Table 1, Fig. 4). The percent of anthropogenic

debris out of total nest mass varied from 3.1% to 22.7% (Supporting Information). Within the urban

area fledging survival, examining the number of nestlings fledged versus nestlings failed, declined with

increasing proportion of anthropogenic material comprising total nest mass (GLM, χ^2 = 13.80, df = 1, P =

0.0002). When we jointly model the covariates of location, urban and non-urban, with the proportion of

nest trash to determine the effect on daily nestling survival, using a Mayfield logistic regression, we

found both covariates to be significant (β_1= -1.169, P <0.0001, β_2= -4.109, P = 0.0391).

The most common materials found across nests were synthetic strings and fibers and synthetic

stuffing followed by plastic (Supporting Information, Fig. 4). The color of anthropogenic nest debris

most collected was white (present in all nests), followed by blue/purple and green. In the urban area, we

found and documented four cases of nest entanglement-associated mortalities, which include three

nestlings from separate nests and one adult female, directly related to anthropogenic debris: 1) one

nestling was found hanging from ingested hair that was woven into the nest, with its leg also tangled in

free hanging plastic string (Fig. 3A), 2) one nestling was hanging from ingested synthetic string (Fig.

3B), 3) one nestling (near fledging) was found with its leg entangled in plastic string, and 4) one adult

female was found hanging, strangulated, from human hair during active nest building (Fig. 3C).

4. Discussion

Low reproductive success in dry years is an established pattern in Darwin's finches. The effect of

dry years on the reproductive success of urban Darwin's finches, however, has not been previously

addressed. In this study, we found, that consistent with the majority of urban rural comparisons

(Chamberlain et al., 2009) urban small ground finches' reproductive effort, egg laying, began earlier then

in their rural congeners. However, reproductive effort as well as reproductive success was significantly

higher, in urban finches compared to non-urban finches: the urban area resulted in higher numbers of

nests with eggs (Fig. 2A), nests with nestlings (Fig. 2B), and fledging success (Fig. 2C). All urban finches

were found to incorporate anthropogenic debris (3.1% to 22.7% of total nest mass) into their nests while

non-urban nests had no anthropogenic debris incorporated. Critically, mortalities due to anthropogenic

nest debris entanglement were recorded across four urban nests, affecting 18% of nests with nestlings and

one female in active nest building, and shown to be a cost associated with nesting in urban areas. These

results suggest that despite anthropogenic debris related mortalities, small ground finches derived an

overall reproductive benefit from urban habituation during a dry year, perhaps associated with higher

reproductive output due to earlier initiation of the breeding season and higher offspring survival, which

may be attributable to the high urban food availability.

The later initiation of breeding and reproductive failures documented in non-urban finches are

likely associated with limited precipitation and dry La Niña conditions as seen previously (Boag & Grant,

1981, 1984; Koop, LeBohec, & Clayton, 2013). Rainfall and the associated increase in primary

productivity is known to initiate and sustain finch breeding (Gibbs & Grant, P.R., 1987; Zhang et al.,

2019; MEI.v2 Retreived 21 December 2019)(Boag & Grant, P.R., 1984). Indicating enough rain occurred

to trigger breeding but perhaps not enough natural food resources were available to sustain breeding

efforts. During the 2018 breeding season, we recorded a high percent of nest abandonments in the non-

urban area: 33% of finch nests (four of 12 nests) were abandoned and failed with eggs. The lower limit of

rainfall required to sustain finch breeding is unknown and may also be impacted by rainfall the previous

year and the associated carryover effects on vegetation and invertebrate communities. The previous year,

2017, was a milder La Niña year (Zhang et al., 2019), providing some support for the premise of

carryover of low primary productivity further affecting the availability of resources in 2018. The

differences in overall reproductive output of urban and non-urban small ground finches in our 2018 study

might be related to the higher food resource availability in the urban area (Lochmiller & Deerenberg,

2000), and low primary and secondary production associated with low precipitation typical of a La Niña

year in the non-urban area. Urban food resources are independent of climatic variability, whereas natural

food resources (seeds and insects) are both lower in dry years (low precipitation, drought). Accordingly,

Galápagos land nesting birds have shown reduced breeding success in dry years (Grant & Grant, 1993;

Koop, LeBohec, & Clayton, 2013; McNew et al., 2019). Non-urban finch fitness, in terms of both

breeding success and adult survival, is dictated by precipitation patterns and the resulting food availability

(Gibbs & Grant, P.R., 1987; Koop, LeBohec, & Clayton, 2013; Grant & Grant, 2014). Therefore, future

studies should examine whether the urban breeding success exceeds non-urban success in wet years and

across years in order to understand the long-term demographic patterns of urban finches. Urban and non-

urban birds seem to face distinct and at least partially non-overlapping stressors; non-urban birds are more

impacted by climatic variables, and resulting food availability, and urban birds are being impacted by

anthropogenic stressors, such as anthropogenic debris and possibly low nutritional food quality.

While human-based food availability in urban environments can benefit adult birds, nestlings

require a higher protein diet (Boag, 1987). Natural high-protein food sources, such as arthropods,

demonstrate declines with increasing urbanization (Shochat et al., 2004). Additionally, the lower quality

of food in urban environments has been seen to negatively affect nestling and juvenile growth in other

urban bird species (Pierotti & Annett, 2001; Seress et al., 2012, 2020). Even if urban food quality is lower

nutritionally than natural food sources, the quantity and consistent availability of the resources can

provide short-term benefits for survival of urban finch nestlings. However, poor nestling diet can directly

impact morphological metrics later in life (Boag, 1987), indicating possible lasting changes in urban finch

adults. Urban finches were observed foraging on a variety of human foods (i.e. chips, rice, bread, corn),

but also seen foraging on refuse (i.e. out of garbage bins, rotting organic material in street) (pers. obs.

JAH, TBV). While previous studies have examined the diet of finches in urban vs. non-urban areas (De

León et al., 2018), an examination is still needed of the resulting effect of low quality diet on urban finch

on long term trends in demography. Furthermore, future studies are needed to determine the long-term

effects of urban nestling diet on adult morphology and overall fitness. The marked variation between

urban and non-urban breeding success found here may have also been compounded by climatic factors

which impact resource availability and these differences, such as parental provisioning rates and diet

analysis, must be taken into account in future studies.

We found that 100% of urban nests contained anthropogenic debris, whereas no anthropogenic

debris was recovered from non-urban nests. These results were likely due to debris availability: abundant

anthropogenic debris was available in the urban area and only two pieces of debris were seen at the non-

urban area during the field season. Furthermore, 24% of urban nests were associated with debris-related

mortalities and the proportion of anthropogenic debris comprising the total nest mass impacted nestling

survival. Other studies have observed mortalities due to both entanglement and ingestion of

anthropogenic debris in nestling and adult land birds (Mee et al., 2007; Henry, Wey, & Balança, 2011;

Theodosopoulos & Gotanda, 2019). However, few studies have examined the effect of debris on

reproductive success in land birds (Jagiello et al., 2018) and only one previous study has examined

passerines (Townsend & Barker, 2014). Several types of common anthropogenic debris (e.g., plastic

string, human hair, and string fibers) are frequently used by urban finches in nest building and pose a

higher risk, as seen in the documented entanglement mortalities. The incorporation of anthropogenic

debris into nests presents a clear cost to birds nesting in urban habitats; however, despite suffering

mortalities due to entanglement, urban nestlings demonstrated higher survival than non-urban nestlings.

Our study was conducted in an area with intermediate, albeit increasing, urbanization, relative to

the other human-inhabited Galápagos Islands. We do not know if a higher degree of urbanization would

yield the same beneficial result for finches. A number of species succeed and are abundant at intermediate

levels of urbanization and disturbance (Ausprey & Rodewald, 2011; Stracey & Robinson, 2012; Perrier et

al., 2018). However, species could reach thresholds at which they no longer benefit from increased

urbanization because of further changes in habitat structure which result in a loss of resources (water,

food resources, perches, nesting sites) (Blair, 1996; Lee et al., 2004); a threshold of urbanization that is

detrimental for Darwin's finches might not have been reached yet. In addition to direct anthropogenic

threats, Darwin's finches in both urban and non-urban areas are also affected by an introduced nest

parasite, *Philornis downsi*, which can affect their reproductive success (Knutie et al., 2016; Sonia

Kleindorfer & Dudaniec, 2016; McNew & Clayton, 2018). We did not consider possible effects of *P.*

downsi in our study because we lacked the power to detect an effect of the parasite (due to the low

reproductive success in the non-urban area). Future studies could examine whether the costs and benefits

of urbanization for the finches vary across islands with different levels of human population sizes, as well

as study the interactions between urbanization and *P. downsi* in limiting reproductive success of Darwin's

finches.

Acknowledgements

We thank Karla Vasco for her assistance and logistical support. We would like to thank the Galápagos

Science Center and the Galápagos National Park for support. We also thank Kiyoko Gotanda, Sabrina

McNew and members of the Knutie lab (Alyssa Addesso, Lauren Albert, Anna Sjodin, Grace Vaziri, and

Mackenzie Watkins) for comments on the manuscript. Galápagos photos credited to SAK, JAH, SRS, and

TBV. Additional photos credited to Monica Hinkle and Santeri Viinamäk.

Supporting Information

Table of qualitative data for nest material dissections is available as supplemental document.

Author contributions: JAH and SAK conceived the study, conducted the data analyses, and wrote the

manuscript; JAH, SAK, KC, SRS, TBV collected data; JAC provided logistical support. All authors

revised and approved the manuscript.

Data availability statement: All raw data are available on Figshare (doi provided upon acceptance).

ORCID

Johanna A. Harvey https://orcid.org/0000-0003-4504-6777

Jaime A. Chaves https://orcid.org/0000-0003-0679-2603

Sarah A. Knutie https://orcid.org/0000-0001-6423-9561

Funding: The work was supported by start-up funds and a Research Excellence Program Grant from the

University of Connecticut and National Science Foundation Grant (DEB-1949858) to SAK and an

American Ornithological Society Hesse Research Award to JAH. JAH was supported by an American

Museum of Natural History Gerstner Scholar Fellowship. We the authors declare we have no direct

financial benefits that could result from publication.

Competing interests: The authors declare that they have no conflict of interest.

Ethical statement: All applicable international, national, and/or institutional guidelines for the care and

use of animals were followed. All bird handling and work was conducted according to approved

University of Connecticut IACUC (Institutional Animal Care and Use Committee) protocols (No. A17-

044). Our work was done under GNP permits PC 03-18 and Genetic Access permit MAE-DNB-CM-

2016-0041.

References

Aronson, M. F. J., La Sorte, F. A., Nilon, C. H., Katti, M., Goddard, M. A., Lepczyk, C. A., Warren, P.

S., Williams, N. S. G., Cilliers, S., Clarkson, B., Dobbs, C., Dolan, R., Hedblom, M., Klotz, S.,

Kooijmans, J. L., Kühn, I., Macgregor-Fors, I., Mcdonnell, M., Mörtberg, U., Pyšek, P., Siebert, S.,

Sushinsky, J., Werner, P., & Winter, M. (2014). A global analysis of the impacts of urbanization on

bird and plant diversity reveals key anthropogenic drivers. *Proc. R. Soc. B Biol. Sci.* **281**.

http//doi.org/10.1098/rspb.2013.3330

Ausprey, I. J., & Rodewald, A. D. (2011). Postfledging Survivorship and Habitat Selection Across A

Rural-To-Urban Landscape Gradient. *Auk* **128**, 293–302. https://doi.org/10.1525/auk.2011.10158

Bailly, J., Scheifler, R., Berthe, S., Clément-Demange, V. A., Leblond, M., Pasteur, B., & Faivre, B.

(2016). From eggs to fledging: Negative impact of urban habitat on reproduction in two tit species.

J. Ornithol. **157**, 377–392. https://doi.org/10.1007/s10336-015-1293-3

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using

lme4. *J. Stat. Softw.* **67**, 251–264. http://doi.org/10.18637/jss.v067.i01

Benning, T. L., LaPointe, D., Atkinson, C. T., & Vitousek, P. M. (2002). Interactions of climate change

with biological invasions and land use in the Hawaiian Islands: Modeling the fate of endemic birds

using a geographic information system. *Proc. Natl. Acad. Sci.* **99**, 14246–14249.

https://doi.org/10.1073/pnas.162372399

Blair, R. B. (1996). Land use and avian species diversity alng an urban gradient. *Ecol. Appl.* **6**, 506–519.

http://doi.org/10.2307/2269387

Boag, P. T. (1987). Effects of Nestling Diet on Growth and Adult Size of Zebra Finches (Poephila

guttata). *Auk* **104**, 155–166. https://doi.org/10.1093/auk/104.2.155

Boag, P. T., & Grant, P. R. (1981). Intense Natural Selection in a Population of Darwin's Finches

(Geospizinae) in the Galapagos. *Science (80-.)*. **214**, 82–85.

http://doi.org/10.1126/science.214.4516.82

Boag, P. T., & Grant, P. R. (1984). Darwin's Finches (*Geospiza*) On Isla Daphne Major, Galapagos :

Breeding and Feeding Ecology in a Climatically Variable Environment. *America (NY)*. **54**, 463–489.

https://doi.org/10.2307/1942596

Buckley, L. B., & Jetz, W. (2007). Insularity and the determinants of lizard population density. *Ecol. Lett.*

10, 481–489. http://doi.org/10.1111/j.1461-0248.2007.01042.x

Chamberlain, D. E., Cannon, A. R., Toms, M. P., Leech, D. I., Hatchwell, B. J., & Gaston, K. J. (2009).

Avian productivity in urban landscapes: a review and meta-analysis. *Ibis (Lond. 1859)*. **151**, 1–18.

https://doi.org/10.1111/j.1474-919X.2008.00899.x

Cohen, J. (2013). Statistical Power Analysis for the Behavioral Sciences (2nd ed.). USA: Lawrence

Erlbaum Associate Publishers.

Courchamp, F., Hoffmann, B. D., Russell, J. C., Leclerc, C., & Bellard, C. (2014). Climate change, sea-

level rise, and conservation: keeping island biodiversity afloat. *Trends Ecol. Evol.* **29**, 127–130.

https://doi.org/10.1016/j.tree.2014.01.001

Curry, R. L., & Grant, P. R. (1989). Demography of the Cooperatively Breeding Galapagos Mockingbird,

Nesomimus parvulus, in a Climatically Variable Environment. *J. Anim. Ecol.* **58**, 441.

http://doi.org/10.2307/4841

De Léon, F. L., Raeymaekers, J. A. M., Bermingham, E., Podos, J., Herrel, A., & Hendry, A. P. (2011).

Exploring possible human influences on the evolution of Darwin's finches. *Evolution (N. Y)*. **65**,

2258–2272. https://doi.org/10.1111/j.1558-5646.2011.01297.x

De León, L. F., Sharpe, D. M. T., Gotanda, K. M., Raeymaekers, J. A. M., Chaves, J. A., Hendry, A. P.,

& Podos, J. (2018). Urbanization erodes niche segregation in Darwin's finches. *Evol. Appl.* 1–15.

https://doi.org/10.1111/eva.12721

Dominoni, D., Quetting, M., & Partecke, J. (2013). Artificial light at night advances avian reproductive

physiology. *Proc. R. Soc. B Biol. Sci.* **280**. https://doi.org/10.1098/rspb.2012.3017

Dvorak, M., Fessl, B., Nemeth, E., Anchundia, D., Cotín, J., Schulze, C. H., Tapia, W., & Wendelin, B.

(2019). Survival and extinction of breeding landbirds on San Cristóbal, a highly degraded island in

the Galápagos. *Bird Conserv. Int.* 1–15. http://doi.org/10.1017/S0959270919000285

Epler, B. (2007). Tourism, the Economy, Population Growth, and Conservation in Galapagos. Galapagos

Islands, Ecuador. https://www.galapagos.org/wp-content/uploads/2012/01/TourismReport2.pdf

Fernández-Juricic, E. (2002). Can human disturbance promote nestedness? A case study with breeding

birds in urban habitat fragments. *Oecologia* **131**, 269–278. https://doi.org/10.1007/s00442-002-

0883-y

Fox, J., & Weisberg, S. (2011). An R Companion to Applied Regression (2nd ed). Thousand Oaks, CA,

USA: Sage.

Gering, J. C., & Blair, R. B. (1999). Predation on artificial bird nest along an urban gradient: predatory

risk or relaxation in urban enviroments? *Ecography.* **22**, 532–541. www.jstor.org/stable/3683146

Gibbs, H. L., & Grant, P. R. (1987). Ecological Consequences of an Exceptionally Strong El Nino Event

on Darwin's Finches. *Ecology* **68**, 1735–1746. http://doi.org/ 10.2307/1939865

Grant, B. R., & Grant, P. R. (1993). Evolution of Darwin's finches caused by a rare climatic event. *Proc.*

R. Soc. London. Ser. B Biol. Sci. **251**, 111–117. https://doi.org/10.1098/rspb.1993.0016

Grant, P. R., & Boag, P. T. (1980). Rainfall on the Galápagos and the Demography of Darwin's Finches.

Auk Ornithol. Adv. **97**, 227–244. http://doi.org/10.1093/auk/97.2.227

Grant, P. R., & Grant, B. R. (1989). Evolutionary dynamics of a natural population: the large cactus finch

of the Galapagos. Chicago, USA: University of Chicago Press.

Grant, P. R., & Grant, B. R. (1999). Effects of the 1998 El Niño on Darwin's Finches on Daphne. *Not.*

Galápagos **60**, 29–30. http://aquaticcommons.org/id/eprint/7468

Grant, P. R., & Grant, B. R. (2014). Synergism of natural selection and introgression in the origin of a

new species. *Am. Nat.* **183**, 671–681. http://doi.org/10.1086/675496

Hazler, K. R. (2004). Mayfield Logistic Regression: A Practical Approach for Analysis of Nest Survival.

Auk **121**, 707–716. https://doi.org/10.1093/auk/121.3.707

Hendry, A. P., Grant, P. R., Grant, B. R., Ford, H. A., Brewer, M. J., & Podos, J. (2006). Possible human

impacts on adaptive radiation: beak size bimodality in Darwin's finches. *Proc. Biol. Sci.* **273**, 1887–

1894. http://doi.org/10.1098/rspb.2006.3534

Henry, P.-Y., Wey, G., & Balança, G. (2011). Rubber Band Ingestion by a Rubbish Dump Dweller, the

White Stork (Ciconia ciconia). *Waterbirds* **34**, 504–508. https://doi.org/10.1675/063.034.0414

Herrera-Dueñas, A., Pineda-Pampliega, J., Antonio-García, M. T., & Aguirre, J. I. (2017). The influence

of urban environments on oxidative stress balance: A case study on the house sparrow in the Iberian

Peninsula. *Front. Ecol. Evol.* **5**, 1–10. https://doi.org/10.3389/fevo.2017.00106

Hijmans, R., Kapoor, J., Wieczorek, J., Garcia, N., Maunahan, A., Rala, A., & Mandel, A. (2014).

GADM Database of Global Administrative Areas (V 3.6). https://gadm.org

INEC. (2016). Análisis de resultados definitivos censo de población y vivienda Galápagos 2015. Quito,

Ecudaor. https://www.ecuadorencifras.gob.ec/censo-de-poblacion-y-vivienda-galapagos/

Jagiello, Z. A., Dylewski, Ł., Winiarska, D., Zolnierowicz, K. M., & Tobolka, M. (2018). Factors

determining the occurrence of anthropogenic materials in nests of the white stork Ciconia ciconia.

Environ. Sci. Pollut. Res. **25**, 14726–14733. https://doi.org/10.1007/s11356-018-1626-x

Jesse, W. A. M., Behm, J. E., Helmus, M. R., & Ellers, J. (2018). Human land use promotes the

abundance and diversity of exotic species on Caribbean islands. *Glob. Chang. Biol.* **24**, 4784–4796.

http://doi.org/ 0.1111/gcb.14334

Jiguet, F., Sunnen, L., Prévot, A. C., & Princé, K. (2019). Urban pigeons loosing toes due to human

activities. *Biol. Conserv.* **240**, 108241. https://doi.org/10.1016/j.biocon.2019.108241

Johnson, M. T. J., & Munshi-South, J. (2017). Evolution of life in urban environments. *Science* **358**,

eaam8327. http://doi.org/10.1126/science.aam8327

Kark, S., Iwaniuk, A., Schalimtzek, A., & Banker, E. (2007). Living in the city : can anyone become an

'urban exploiter'? *J. Biogeogr.* 638–651. https://doi.org/10.1111/j.1365-2699.2006.01638.x

Kier, G., Kreft, H., Lee, T. M., Jetz, W., Ibisch, P. L., Nowicki, C., Mutke, J., & Barthlott, W. (2009). A

global assessment of endemism and species richness across island and mainland regions. *Proc. Natl.*

Acad. Sci. **106**, 9322–9327. https://doi.org/10.1073/pnas.0810306106

Kleindorfer, S. (2007). The ecology of clutch size variation in Darwin's Small Ground Finch Geospiza

fuliginosa: Comparison between lowland and highland habitats. *Ibis (Lond. 1859).* **149**, 730–741.

https://doi.org/10.1111/j.1474-919X.2007.00694.x

Kleindorfer, Sonia, Chapman, T. W., WInkler, H., & Sulloway, F. J. (2006). Adaptive divergence in

contiguous populations of Darwin's Small Ground Finch (Geospiza fuliginosa). *Evol. Ecol. Res.* **8**,

357–372.

https://dspace2.flinders.edu.au/xmlui/bitstream/handle/2328/9243/2006002287.pdf?sequence=1

Kleindorfer, Sonia, & Dudaniec, R. Y. (2016). Host-parasite ecology, behavior and genetics: a review of

the introduced fly parasite Philornis downsi and its Darwin's finch hosts. *BMC Zool.* **1**, 1.

https://doi.org/10.1186/s40850-016-0003-9

Knutie, S. A., Chaves, J. A., & Gotanda, K. M. (2019). Human activity can influence the gut microbiota

of Darwin's finches in the Galapagos Islands. *Mol. Ecol.* **9,**2441-2450.

https://doi.org/10.1111/mec.15088

Knutie, S. A., McNew, S. M., Bartlow, A. W., Vargas, D. A., & Clayton, D. H. (2014). Darwin's finches

combat introduced nest parasites with fumigated cotton. *Curr. Biol.* **24**, R355–R356. http://doi.org/

10.1016/j.cub.2014.03.058

Knutie, S. A., Owen, J. P., McNew, S. M., Bartlow, A. W., Arriero, E., Herman, J. M., DiBlasi, E.,

Thompson, M., Koop, J. A. H., & Clayton, D. H. (2016). Galápagos mockingbirds tolerate

introduced parasites that affect Darwin's finches. *Ecology* **97**, 940–950. https://doi.org/10.1890/15-

0119.1

Koop, J. A. H., LeBohec, C., & Clayton, D. (2013). Dry year does not reduce invasive parasitic fly

prevalence or abundance in Darwin's finch nests. *Reports Parasitol.* **3**, 11. http://doi.org/

10.2147/RIP.S48435

Lee, P.-F., Ding, T.-S., Hsu, F.-H., & Geng, S. (2004). Breeding bird species richness in Taiwan:

distribution on gradients of elevation, primary productivity and urbanization. *J. Biogeogr.* **31**, 307–

314. https://doi.org/10.1046/j.0305-0270.2003.00988.x

Lepczyk, C. A., Mertig, A. G., & Liu, J. (2004). Landowners and cat predation across rural-to-urban

landscapes. *Biol. Conserv.* **115**, 191–201. https://doi.org/10.1016/S0006-3207(03)00107-1

Lochmiller, R. L., & Deerenberg, C. (2000). Trade-offs in evolutionary immunology: just what is the cost

of immunity? *Oikos* **88**, 87–98. https://doi.org/10.1034/j.1600-0706.2000.880110.x

Lowry, H., Lill, A., & Wong, B. B. M. (2013). Behavioural responses of wildlife to urban environments.

Biol. Rev. **88**, 537–549. https://doi.org/10.1111/brv.12012

Mayfield, H. (1961). Nesting Success Calculated from Exposure. *Wilson Bull.* **73**, 255–261.

McNew, S. M., Beck, D., Sadler-Riggleman, I., Knutie, S. A., Koop, J. A. H., Clayton, D. H., & Skinner,

M. K. (2017). Epigenetic variation between urban and rural populations of Darwin's finches. *BMC*

Evol. Biol. **17**, 183. https://doi.org/10.1186/s12862-017-1025-9

McNew, S. M., & Clayton, D. H. (2018). Alien Invasion: Biology of *Philornis* Flies Highlighting

Philornis downsi, an Introduced Parasite of Galápagos Birds. *Annu. Rev. Entomol.* **63**, 369–387.

https://doi.org/10.1146/annurev-ento-020117-043103

McNew, S. M., Knutie, S. A., Goodman, G. B., Theodosopoulos, A., Saulsberry, A., Yépez R., J., Bush,

S. E., & Clayton, D. H. (2019). Annual environmental variation influences host tolerance to

parasites. *Proc. R. Soc. B Biol. Sci.* **286**, 20190049. https://doi.org/10.1098/rspb.2019.0049

Mee, A., Rideout, B. A., Hamber, J. A., Todd, J. N., Austin, G., Clark, M., & Wallace, M. P. (2007). Junk

ingestion and nestling mortality in a reintroduced population of California Condors Gymnogyps

californianus. *Bird Conserv. Int.* **17**, 119–130. https://doi.org/10.1017/S095927090700069X

Møller, A. P., Díaz, M., Flensted-Jensen, E., Grim, T., Ibáñez-Álamo, J. D., Jokimäki, J., Mänd, R.,

Markó, G., & Tryjanowski, P. (2015). Urbanized birds have superior establishment success in novel

environments. *Oecologia* **178**, 943–950. https://doi.org/10.1007/s00442-015-3268-8

Navarro, D. J., & Foxcroft, D. R. (2019). Learning statistics with Jamovi: a tutorial for psychology

students and other beginners (Version 0.70). Vol. 53. http://doi.org/ 10.24384/hgc3-7p15

Perrier, C., Lozano del Campo, A., Szulkin, M., Demeyrier, V., Gregoire, A., & Charmantier, A. (2018).

Great tits and the city: Distribution of genomic diversity and gene–environment associations along

an urbanization gradient. *Evol. Appl.* **11**, 593–613. https://doi.org/10.1111/eva.12580

Phillips, R. B., Wiedenfeld, D. A., & Snell, H. L. (2012). Current status of alien vertebrates in the

Galápagos Islands: Invasion history, distribution, and potential impacts. *Biol. Invasions* **14**, 461–

480. https://doi.org/10.1007/s10530-011-0090-z

Pierotti, R., & Annett, C. (2001). The ecology of Western Gulls in habitats varying in degree of urban

influence. In *Avian Ecol. Conserv. an Urban. World*. pp. 307–329. Boston, MA: Springer.

http://doi.org/10.1007/978-1-4615-1531-9_15

R Core Team. (2012). R: A Language and Environment for Statistical Computing. R Foundation for

Statistical Computing. (L. Ranjard, F. Poly, & J. C. Lata, Eds.). Vienna, Austria.

Roman, L., Hardesty, B. D., Hindell, M. A., & Wilcox, C. (2019). A quantitative analysis linking seabird

mortality and marine debris ingestion. *Sci. Rep.* **9**, 3202. https://doi.org/10.1038/s41598-018-36585-

9

Sepp, T., McGraw, K. J., Kaasik, A., & Giraudeau, M. (2018). A review of urban impacts on avian life-

history evolution: Does city living lead to slower pace of life? *Glob. Chang. Biol.* **24**, 1452–1469.

https://doi.org/10.1111/gcb.13969

Seress, G., Bókony, V., Pipoly, I., Szép, T., Nagy, K., & Liker, A. (2012). Urbanization, nestling growth

and reproductive success in a moderately declining house sparrow population. *J. Avian Biol.* **43**,

403–414. https://doi.org/10.1111/j.1600-048X.2012.05527.x

Seress, G., Sándor, K., Evans, K. L., & Liker, A. (2020). Food availability limits avian reproduction in

the city: An experimental study on great tits Parus major. *J. Anim. Ecol.* **89**, 1570–1580.

https://doi.org/10.1111/1365-2656.13211

Shochat, E., Stefanov, W. L., Whitehouse, M. E. A., & Faeth, S. H. (2004). Urbanization and spider

diversity: Influenences of human modification of habitat structure and productivity. *Ecol. Appl.* **14**,

268–280. https://doi.org/10.1007/978-0-387-73412-5_30

Stracey, C. M., & Robinson, S. K. (2012). Are urban habitats ecological traps for a native songbird?

Season-long productivity, apparent survival, and site fidelity in urban and rural habitats. *J. Avian*

Biol. **43**, 50–60. https://doi.org/10.1111/j.1600-048X.2011.05520.x

Theodosopoulos, A. N., & Gotanda, K. M. (2019). Death of a Darwin's Finch: a consequence of human-

made debris? *Wilson J. Ornithol.* **130**, 1023. https://doi.org/10.1676/1559-4491.130.4.1023

Townsend, A. K., & Barker, C. M. (2014). Plastic and the nest entanglement of urban and agricultural

crows. *PLoS One* **9**, 1–5. https://doi.org/10.1371/journal.pone.0088006

Trueman, M., & d'Ozouville, N. (2010). Characterizing the Galapagos terrestrial climate in the face of

global climate change. *Galapagos Res.* **67**, 26–37.

Vitousek, P. M. (1997). Human Domination of Earth's Ecosystems. *Science.* **277**, 494–499.

http://doi.org/10.1126/science.277.5325.494

Walsh, S. J., & Mena, C. F. (2016). Interactions of social, terrestrial, and marine sub-systems in the

Galapagos Islands, Ecuador. *Proc. Natl. Acad. Sci.* **113**, 14536–14543.

https://doi.org/10.1073/pnas.1604990113

Yeh, P. J., & Price, T. D. (2004). Adaptive Phenotypic Plasticity and the Successful Colonization of a

Novel Environment. *Am. Nat.* **164**, 531–542. http://doi.org/10.1086/423825

Zhang, T., Hoell, A., Perlwitz, J., Eischeid, J., Murray, D., Hoerling, M., & Hamill, T. M. (2019).

Towards Probabilistic Multivariate ENSO Monitoring. *Geophys. Res. Lett.* **46**, 10532–10540.

https://doi.org/10.1029/2019GL083946