Speciation and the developmental alarm clock

Asher D. Cutter [1] & Joanna D. Bundus [2]

[1]Department of Ecology & Evolutionary Biology, University of Toronto

asher.cutter@utoronto.ca

[2]Department of Integrative Biology, University of Wisconsin – Madison

bundus@wisc.edu

Abstract

New species arise as the genomes of populations diverge. The developmental 'alarm clock' of speciation sounds off when sufficient divergence in genetic control of development leads hybrid individuals to infertility or inviability, the world awoken to the dawn of new species with intrinsic post-zygotic reproductive isolation. Some developmental stages will be more prone to hybrid dysfunction due to how molecular evolution interacts with the ontogenetic timing of gene expression. Considering the ontogeny of hybrid incompatibilities provides a profitable connection between 'evo-devo' and speciation genetics to better link macroevolutionary pattern, microevolutionary process, and molecular mechanisms. Here we explore speciation alongside development, emphasizing their mutual dependence on genetic network features, fitness landscapes, and developmental system drift. We assess models for how ontogenetic timing of reproductive isolation can be predictable. Experiments and theory within this synthetic perspective can help identify new rules of speciation as well as rules in the molecular evolution of development.

Impact Statement

Integrating speciation genetics with ontogeny can identify predictable rules in the molecular evolution of developmental pathways and in the accumulation of reproductive isolation as genomes diverge.

Introduction

"*Are certain developmental processes especially likely to be disrupted in hybrids? This question has been surprisingly neglected given that hybrid defects provide a rare window on those developmental processes and pathways that diverge rapidly between taxa.*" – Coyne and Orr (2004), p.309

Distinct species are those separate collections of genomes that, if you were to put them together in the same cells, the mixture would create a broken organism. Development in the hybrid individuals would go awry in such a way as to cause inviability, infertility, or a phenotypic mismatch to ecology or mating interactions that compromises further reproductive success. Given common descent from an ancestor, how could evolution produce such disastrous phenotypic consequences of genetic change? Darwin recognized this dilemma (Darwin 1859), in that natural selection will oppose changes that confer a net fitness cost. But evolution can circumvent this problem through interactions between multiple genetic factors, as intuited by W. Bateson, T. Dobzhansky and H.J. Muller and made explicit in the genetic mechanism of Dobzhansky-Muller incompatibility (DMI) for post-zygotic disruption in hybrids (Box 1A,B). Put simply: when evolution substitutes an allele at one locus, it makes no guarantee that this new derived genetic background will be compatible with allele substitutions occurring in other populations at other loci. Such inter-locus incompatibilities could involve two, or three, or many more interacting loci to create a DMI to genetically enforce species boundaries.

In this way, the mutational substitutions that accumulate in one population are indifferent to the substitutions that accumulate in other populations, but only so long as the two populations do not share alleles with one another through gene flow. When the populations of incipient species do intermingle genetically, then developmental programs in hybrid individuals must confront whether alleles derived from such 'out-of-sight, out-of-mind' evolution interact in a way that allows ontogeny to proceed normally (Box 1A,B). Incompatible combinations of alleles in hybrid individuals at two or more loci can be thought of as negative epistasis or as a kind of non-linear genetic perturbation. Put this way, the genetics of speciation sound eerily similar to notions of cryptic genetic variation (Wade et al. 1997, Ledón-Rettig et al. 2014, Paaby and

Rockman 2014) and developmental system drift in the literature on the evolution of development

(True and Haag 2001, Pavlicev and Wagner 2012, Tulchinsky et al. 2014a). The regulatory

pathways that govern development are an important component of how and whether divergence

will lead to genetic incompatibility and speciation (Porter and Johnson 2002). Our aim here is to

draw these connections more explicitly, to place speciation in a firmer developmental context

(Sucena and Stern 2000, Johnson and Porter 2001), and to emphasize the relevance of the

evolution of reproductive isolation to problems in developmental biology.

Distinguishing detectable versus undetectable differences in phenotype between species is key to

thinking about divergence in the genes and genetic networks that underlie the developmental

pathways that create an organism. Detectable differences in the development of traits between

species, often due to adaptive divergence, clearly involve genetic changes. The converse,

however, is not true: conservation of phenotype does not imply conservation of genetic controls.

For example, stabilizing selection on expression levels leads to similar expression levels between

species, despite evidence of widespread compensatory effects of molecular evolution to both *cis-*

and *trans*-acting regulators of gene expression (Mack and Nachman 2017). This idea of genetic

change despite phenotypic stasis is known in evo-devo as developmental system drift (DSD), and

in population genetics as evolution along a ridge in a fitness landscape (Box 1C,D). Adaptive

evolution at the molecular level also can contribute to DSD for a given trait, particularly if the

genetic changes affect fitness through pleiotropic effects with phenotypic consequences for only

a subset of traits or from subsequent compensatory evolution (Box 1C).

When molecular evolutionary differences between species involve two or more loci, they may

not interact properly in hybrid individuals that have copies of both genomes. This can happen

regardless of whether those differences induce detectable phenotypic differences that distinguish

the species and regardless of whether selection or genetic drift caused their fixation. Each

mutational substitution that distinguishes species in protein coding sequence or gene regulatory

control thus has some probability of contributing to the formation of a DMI in hybrid

individuals. The more substitutions, the more chances for incompatibilities to create hybrid

dysfunction at some point in development (Orr 1995). In this way, inter-species hybrids can

reveal genetic divergence in the control of even those developmental programs that yield

seemingly-equivalent phenotypic outputs. Hybrids can also reveal the incidence and role of

different kinds of changes, such as *cis-* vs *trans-*acting regulators of gene expression (Box 2)

(Wittkopp et al. 2004, Mack et al. 2016).

Divergence in developmental genetic programs and intrinsic post-zygotic reproductive isolation

in the speciation process are thus close conceptual brethren, despite their largely separate

research traditions. But how do features of genetic networks and evolutionary forces intersect to

create such profound developmental genetic divergence revealed as DMIs in hybrids? Are some

stages of ontogeny predisposed to genetic architectures and evolutionary pressures to be more

likely to yield dysfunctional development in hybrid organisms? Applying our timepiece

metaphor: what gears and springs in the genetic clockwork will set the developmental alarm

clock to ring at one time versus another during ontogeny to signal speciation? The more

molecular evolutionary change, the more likely are genetic incompatibilities to manifest in inter-

species hybrids. We must therefore set our goal to define the factors that make some genes

evolve more quickly than others. The answers will help us to determine ways in which the

molecular evolution that underpins development is predictable and, consequently, in what ways

the genetics of speciation is predictable.

To address this goal, here we first consider how molecular evolution is influenced by the

properties of genes and genetic networks, such as pleiotropy, modularity, robustness, and

cis/trans regulation. These are the "gears and springs" in the genetic architecture of

development. We then explore how genetic architecture may sensitize some phases of ontogeny

disproportionately to disruptive effects of misexpression in genetic networks. By integrating

genetic architecture and ontogenetic timing, we arrive at distinct predictions for how molecular

evolution and hybrid dysfunction manifest across developmental time. Finally, we summarize

the literature on these issues for three case study systems (*Caenorhabditis* nematodes,

Drosophila fruit flies, and anuran *Bufo* and *Xenopus*). In the present state of the field, we find

that few general answers emerge from these factors considered in isolation, motivating deeper

attention from theory and empirical study.

Speciation and development: divergence in genetic networks

As we explore how the evolution of development intersects with speciation, it is valuable to consider some key aspects of genetic architecture from the perspective of multi-gene networks that, in turn, control organismal development (Johnson and Porter 2007, Palmer and Feldman 2009). Here we focus on how pleiotropy, network modularity, and robustness can influence the molecular evolution of coding sequences and non-coding regulatory elements. These links will help ground our expectations for incorporating ontogenetic time into our thinking to then consider predictions for molecular evolution and the production of incompatibilities in genetic networks of hybrid individuals.

Pleiotropic roles and effects

The mapping of genotype on phenotype and fitness leads us to predict that evolution will proceed along genetic lines of least resistance (Schluter 1996). Genetic "resistance" to evolutionary change is affected by the mutability and covariation of traits within and across developmental stages, in addition to natural selection. That is, the net rate of evolution integrates the likelihood and accumulation of mutational input (genetic variance) with the consequences of genetic variants for the ensemble of traits that comprise an organism (genetic covariances between traits) and fitness (natural selection). From the perspective of developmental biologists, this means that genetic hot-spots of evolutionary change over the long term ought to favor mutations with minimally-pleiotropic effects (Carroll 2005, Stern and Orgogozo 2008), a form of developmental bias. As more molecular evolutionary change occurs, genetic incompatibilities become more likely to form from dysfunctional interactions between diverged sequences in inter-species hybrids. Our goal, then, is to enumerate the factors that facilitate molecular evolution to make some genes evolve more quickly than others. The structure of genes and gene regulatory networks provide clues to predict what these factors are (Garfield et al. 2013). These clues about molecular evolution can help us in thinking how to connect the temporal dynamics of developmental genetic networks to genetic incompatibilities between species.

The extent of pleiotropic effects induced by mutation to a gene is a major determinant of the likelihood of evolutionary change to that gene (Carroll 2005, Stern and Orgogozo 2008). This logic follows from the idea that most traits experience stabilizing selection on short timescales, and so the effects of mutation on most traits will be detrimental and offset any fitness benefit conferred to a particular trait by a mutation. On longer timescales, conserved traits are thought to track a moving but bounded fitness landscape (Estes and Arnold 2007), which can facilitate compensatory evolution among the collection of interacting genes of the genetic network to result in sequence divergence and reproductive isolation between populations (Haag and Molla 2005, Tulchinsky et al. 2014a, Tulchinsky et al. 2014b). Consequently, we might expect that more rapid evolution should occur in genes with fewer pleiotropic roles or for classes of changes that perturb fewer of a gene's roles. The position of a gene within gene regulatory networks defines the pleiotropic roles it plays in development. Genes in more central and highly connected network positions can influence a greater fraction of the genome, and so changes to those genes have the potential to induce greater pleiotropic effects (Promislow 2004, Hahn and Kern 2005, He and Zhang 2006, Kahali et al. 2009). However, the precise relationship between centrality and pleiotropy remains complex (Siegal et al. 2007). And, despite the idea that network 'kernels' may perdure through evolution (Erwin and Davidson 2009), the molecular interactions among kernel components may nevertheless coevolve in ways that could generate DMIs (Haag 2007, Tulchinsky et al. 2014a). Genes that are expressed at high levels also are more likely to have a large number of interaction partners and therefore to occupy central network positions (Bloom and Adami 2003).

Coding sequence changes can affect protein activity everywhere they get expressed, and so ought to have greater potential for pleiotropic effects than regulatory changes (Wray et al. 2003, Carroll 2005, Stern and Orgogozo 2008). Alterations of *cis*-regulatory elements, in contrast, generally influence only a subset of a gene's expression, and so will exert the fewest pleiotropic effects relative to coding sequence and *trans*-regulatory changes. Consequently, *trans*-regulatory evolution and coding sequence divergence ought generally to be slower for genes with high centrality and connectivity in genetic networks. We should expect such genes to have especially high ratios of *cis*:*trans* regulatory divergence, provided that those genes also have reasonably complex arrays of *cis*-regulatory elements controlling their expression that can facilitate

divergence. Because *trans*-acting regulation tends to show greater condition-dependence than

does *cis*-regulation (Smith and Kruglyak 2008, Tirosh et al. 2009), however, pleiotropy-based

predictions for their evolution may be tempered by the fact that varying conditions experienced

by organisms could limit the net negative pleiotropic effects of *trans*-regulatory changes.

Moreover, *trans*-regulatory evolution may often be coupled to subsequent *cis*-regulatory

compensatory evolution that ameliorates negative pleiotropic effects of a population having fixed

a beneficial *trans*-regulatory change. Such compensatory evolution involving both *cis*- and *trans*-

acting regulatory evolution may be an especially important contributor to DMI formation

(Landry et al. 2005, Ortíz-Barrientos et al. 2007, Takahasi et al. 2011, Mack et al. 2016).

Experiments show that *trans*-acting regulatory factors contribute disproportionately to

segregating genetic variation for gene expression levels within species, whereas *cis*-acting

regulatory differences disproportionately underlie fixed genetic differences in expression

between species (Wittkopp and Kalay 2012). The loci controlling DMIs also can be polymorphic

or fixed (Cutter 2012), and so to the extent that regulatory evolution is responsible for creating

DMIs, we ought to expect polymorphic DMIs to often be controlled by *trans*-acting factors.

Because *trans*-regulatory mutations tend to arise more readily due to larger mutational target size

in the genome, and to be more recessive, than *cis*-acting mutations (Landry et al. 2007, Gruber et

al. 2012), they are expected to be a common and persistent kind of polymorphism within

populations.

Modularity of genetic network architecture

Modularity of genetic networks reduces the scope for genes to have highly pleiotropic roles

(Wagner and Altenberg 1996). By constraining the neighborhood of partner interactions, greater

modularity limits the pleiotropic effects of changes to gene expression or function for any given

gene. Like tissue-specific expression in the spatial modularity of gene networks (Larracuente et

al. 2008, Packer et al. 2019), we can also consider temporal modularity for those portions of gene

regulatory networks that show stage-specific expression. Modularity of genetic network structure

will increase over ontogeny as cells differentiate and tissues establish autonomous regulatory

programs (Packer et al. 2019). Some transient phases of development are thought to reverse this

217 trend, however, as when tissues integrate during gastrulation or reorganize during

218 metamorphosis (Chin-Sang and Chisholm 2000). Consequently, if pleiotropy at the scale of the

219 whole organism constrains evolution of individual genes, then we might expect faster molecular

220 evolution for genes that experience highly modular genetic network structures due to spatially-

221 or temporally-restricted expression. This logic provides one way to frame the 'early

222 conservation' and 'hourglass' models in the evolution of development (Figure 1).

224 Despite a trend of increasing genetic network modularity from single-celled zygote to adult,

225 some genes will experience the modularity more than others at any given place or time in

226 ontogeny. Genes with greater breadth of expression across tissues, and genes with greater

227 temporal persistence of expression, will experience more scope to contribute to different genetic

228 networks and their phenotypic outputs. Consequently, genes most likely to produce pleiotropic

229 effects when genetically perturbed are those expressed at high levels across many tissues

230 throughout a long span of developmental time (high expression breadth and low temporal

231 expression bias; (Larracuente et al. 2008, Coronado-Zamora et al. 2019)). Even a modest role in

232 any given tissue or at any single stage of development, however, may compound when integrated

233 over space and time.

235 The appropriate weighting of the importance of spatial versus temporal expression breadth and

236 modularity, however, is not entirely obvious. Therefore, empirically, it will be valuable to

237 partition the ontogenetic trajectories of genes with similar expression breadths across tissues, or,

238 complementarily, partition the spatial profiles of expression for genes with similar ontogenetic

239 expression dynamics (Box 3). The transcriptome analysis of distinct cell lineages, as conducted

240 for *C. elegans* precursors of endoderm, mesoderm and ectoderm (Hashimshony et al. 2014) or

241 for single-cell transcriptomes throughout embryogenesis (Hashimshony et al. 2012, Tintori et al.

242 2016, Packer et al. 2019), provides one intriguing scheme for approaching this issue. Another

243 approach could incorporate developmental time into transcriptome analysis of serial-sectioned

244 samples to access 4-dimensional expression profiles through developmental space and time

245 (Ebbing et al. 2018).

247 ***Robustness to genetic change in gene regulatory networks***

Selection on traits within a species often is stabilizing, favoring a particular phenotypic value and

meaning that factors that lead to alternative phenotypic values will have lower fitness. A

genetically 'robust' phenotype is insensitive to genetic perturbations, which permits an organism

to produce the same phenotype and to maintain fitness despite mutational disruption to a gene or

genetic network that contributes to the trait's developmental program (Felix and Wagner 2008).

That is, a greater fraction of mutations to genes in more robust networks have effectively neutral

effects (Ohta 2011). A gene network's robustness therefore ought to influence the mutations that

can accumulate, which in turn affects rates of molecular evolutionary divergence and the

manifestation of genetic incompatibilities in hybrids. Differences across gene networks in

genetic robustness could arise from differential selection on robustness itself (adaptive

robustness or canalization) or could simply arise as a byproduct of differences in their epistatic

genetic architectures (Hermisson and Wagner 2004). Genes and genetic networks that are more

robust to genetic perturbation can be thought of as larger phenotypic capacitors (Paaby and Testa

2018), letting more cryptic genetic divergence in developmental controls accumulate to be

revealed upon the formation of inter-species hybrids. This effect is analogous to how specific

environmental circumstances or genetic backgrounds can expose so-called cryptic genetic

variation or conditionally-neutral variants within a species (Ledón-Rettig et al. 2014, Paaby and

Rockman 2014).

As a consequence, traits with greater genetic robustness ought to accumulate more cryptic

genetic variation within a species that can then can contribute to greater developmental system

drift between species as substitutions accrue (Felix and Wagner 2008). This process will, in turn,

lead separate populations to travel independently along a fitness ridge in the landscape of

genotype space (Box 1D) (Felix and Wagner 2008, Fragata et al. 2019). More robust networks

are more likely to contain more cryptic genetic variation, but the presence of cryptic genetic

variation alone is insufficient to conclude whether or not selection directly favored greater

robustness, and characterizing the robustness of genetic networks empirically can be difficult.

The selective regime that shapes the fitness landscape will influence the speed at which

developmental system drift can evolve for any given trait (Tulchinsky et al. 2014b). DSD is

especially likely to result from directional selection on traits, perhaps facilitated by a role for

cryptic genetic variation in adaptation to environmental shifts (Felix and Wagner 2008), and

from selection that drives molecular co-evolution among genetic elements (Tulchinsky et al.

2014b).

Phenotypes with a larger underlying genetic network are expected to be more robust to

perturbation (Fragata et al. 2019), implying that they should more readily tolerate the

accumulation of genetic changes. Genes holding more central positions in genetic networks also

appear to be associated with greater robustness upon perturbation (Proulx et al. 2007), though

this effect may depend more on expression level than connectivity per se (Siegal et al. 2007).

This means that changes to any given gene in a network that is shared with other networks, and

that confers a beneficial effect to that other network, are less likely to manifest negative

pleiotropic effects for those genetic networks that are larger (Pavlicev and Wagner 2012).

Consequently, cryptic variation and DSD may be disproportionately prevalent in developmental

programs comprised of larger genetic networks. Genetic networks with low modularity, by

definition, have all constituent genes interacting, and networks with low modularity indeed are

most robust to genetic perturbation (Tran and Kwon 2013).

Recall, however, that we argued how such low modularity conditions maximizes the pleiotropic

roles of genes, which should impede rather than tolerate molecular evolution. These effects thus

appear to present a contradiction. The resolution may lie in assumptions about network size, such

that the implications of low modularity for pleiotropy may be secondary to robustness for large

genetic networks. Analysis of networks in high-resolution time series of expression showing

distinct tissues with distinct transcriptomic profiles that vary in size may help in testing this idea

(Packer et al. 2019). The resolution may also depend on whether we consider coding or

regulatory sequence evolution, as coding sequence evolution does not appear to depend strongly

on how connected the gene is in the genetic network (Jordan et al. 2003, Batada et al. 2006). The

way that connections define the network output also can be crucial. Simple connectivity metrics

may just poorly summarize functional and evolutionary properties of networks, making the

contradiction more apparent than real (Siegal et al. 2007). For example, genetic networks with

switch-like behavior may be more capable of accumulating cryptic variation, as appears to occur

disproportionately among genes with early zygotic expression in sea urchin embryogenesis

(Garfield et al. 2013).

Ontogenetic stages that have an especially high incidence of mutationally-robust phenotypes

within species should thus be especially prone to experience developmental system drift. Such

DSD can get revealed in the novel genomic environment of inter-species hybrids, taking the

form of DMIs. A counter-argument to this idea is that mutationally-robust traits might also be

predisposed to being robust to the genomic perturbation experienced in hybrid individuals, and

therefore be less likely to show dysfunction. The logic of this counter-argument might be more

pertinent at the very early stages of divergence between populations, becoming less and less

applicable as substitutions accrue in the speciation process. It will be interesting for future

research to resolve this question. In general, it remains unclear how network robustness within

species versus between species might scale differently with the number of genetic changes.

Measurements of mutational variance for traits shared across ontogenetic stages provide another

possible way to test for stage-dependent changes in mutational robustness, for example, using

mutation accumulation experiments that, so far, point to this possibility (Farhadifar et al. 2016,

Zalts and Yanai 2017). It will be interesting to see in future studies whether the genetic

robustness of traits truly can predict the incidence of cryptic genetic variation, developmental

system drift, and the propensity to reveal DMIs in hybrids.

Degeneracy via partially-redundant genetic network pathways provides another way that

phenotypes could be robust to genetic perturbation, potentially allowing DSD to accumulate

without yielding DMIs in F1 hybrids. Indeed, genes associated with redundant networks evolve

more quickly (Wagner 2000, Kitami and Nadeau 2002). Robustness mediated by distributed,

rather than redundant (Felix and Wagner 2008), genetic networks may thus more readily

experience DSD in a way that would foster DMIs. Moreover, translational buffering of gene

expression leads to the inference that protein production shows much less misexpression than do

mRNA transcripts in F1 hybrids (Leducq et al. 2012, Khan et al. 2013, Artieri and Fraser 2014,

McManus et al. 2014). And, it is fitness as a phenotype that is the ultimate readout of robustness.

Consequently, genetic networks important for development may be less disrupted in hybrids than

transcriptome analyses might otherwise suggest. Overall, discerning the relative incidence of

distinct mechanisms conferring phenotypic robustness will be important in defining whether

different genetic networks and different stages of development will be more or less likely to

contribute to post-zygotic reproductive isolation as divergence between species accumulates.

Evolvability of distinct genetic components

Mode of selection: Traits and sequences with greater propensity to diverge are said to be more

evolvable (Wagner and Zhang 2011). One way to detect evolvability is when traits and

sequences diverge as a result of natural selection, because adaptive divergence in phenotypes

between species makes it easy to conclude that there must have been changes to the underlying

genetic networks. Such selection can involve multiple changes on an adaptive path, for example,

if first a large-effect *trans*-regulatory mutation gets fixed and is followed by subsequent

compensatory *cis*-regulatory substitutions that ameliorate suboptimal pleiotropic effects of the

initial *trans*-acting substitution (Box 1C) (Goncalves et al. 2012); similar logic can also apply to

coding sequence changes (Clark et al. 2009). Moreover, positive selection and multi-locus

antagonistic coevolution will drive molecular evolution that is much more rapid than will genetic

drift. Repeated adaptive evolution of orthologous genes in disparate lineages is implicated as a

general feature of some types of phenotypic change, including aggregate effects of multiple

mutations affecting the same locus (Stern and Orgogozo 2008, Martin and Orgogozo 2013). The

genes contributing to such adaptive divergence may thus be especially likely to contribute to

post-zygotic DMIs (Presgraves 2010b), in addition to pre-mating, gametic, or ecological

reproductive isolation barriers.

Alternately, some genes may evolve faster than others simply because they experience weaker

purifying selection or greater mutational target size. The evolutionary accumulation of changes

in this way is rate-limited by the input of mutation and the speed of genetic drift, and so will be

faster in species with small population sizes.

Haldane's rule as developmentally predictable evolution: One developmentally-predictable

rule in evolution says something about sex: Haldane's rule, in which the heterogametic sex is

more likely to suffer inviability or sterility in inter-species hybrids (Haldane 1922, Delph and

Demuth 2016). Dominance theory provides one explanation for this pattern: incompatibility loci

linked to sex chromosomes will reveal themselves in F1 hybrids disproportionately for the sex

that has only one copy of a given sex chromosome (Turelli and Orr 1995). This tells us that

genomic compartmentalization of how traits are genetically encoded also is important for the

implications of molecular evolution; when genes contributing to a given developmental process

are biased in genomic location, it may confer greater or lesser tendency to yield DMIs in hybrids.

When males are heterogametic (X/Y or X/O sex-determination), other factors may also

contribute to Haldane's rule (Delph and Demuth 2016). Genes that control male-biased traits

may evolve especially rapidly due to sexual selection or mutational biases ('faster male' theory),

or gene regulatory networks controlling male traits may be unusually sensitive to genetic

perturbation in hybrids ('fragile males'), or genes linked the X-chromosome may evolve

especially fast ('faster X') (Charlesworth et al. 1987, Wu and Davis 1993). Consequently, X-

linked genes may show distinct patterns of misexpression in hybrids (Moehring et al. 2007,

Turner et al. 2014, Civetta 2016, Sanchez-Ramirez et al. 2020). Developmental pathways related

to spermatogenesis seem especially sensitive to disruption in hybrids, perhaps being predisposed

to disproportionate compensatory *cis-trans* regulatory coevolution (Mack et al. 2016, Mack and

Nachman 2017). Sex-limited genetic networks also may differ for males versus females in size,

location of genomic encoding, or predominant mechanism of regulatory control, and so influence

the relative accumulation of cryptic genetic variation and developmental system drift.

Coding vs regulatory evolution: Both coding sequences and regulatory sequences diverge

between species, despite the fact that most selection on each of them is expected to be purifying

(Casillas et al. 2007). The rate of evolution for a coding sequence and its *cis*-regulatory regions,

however, correlate only weakly (Castillo-Davis et al. 2004, Liao and Zhang 2006, Tirosh and

Barkai 2008). This observation implies that the strength and mode of selection affecting

mutations to protein structure tells us little about the strength and mode of selection on mutations

affecting expression, and vice versa. Consequently, molecular evolution associated with

ontogenetic timing may show distinct patterns for coding and regulatory sequences, and so also

yield different implications for when DMIs manifest over ontogeny.

The type of regulatory change may be key, however, in understanding the relation between
regulatory and coding sequence divergence (i.e. evolvability), as well as to the likelihood of
contributing to a DMI. For example, genes showing evidence of *trans*-regulatory divergence
appear to evolve slower in their coding sequences than do genes with *cis*-regulatory divergence
(Goncalves et al. 2012). And, DMIs due to misregulation generally involve non-compensatory
evolution of both *cis*- and *trans*-regulators of expression (Ortíz-Barrientos et al. 2007, Mack and
Nachman 2017). Specifically, regulatory divergence involving reinforcing *cis+trans* changes
(thought to result from directional selection within species) may actually be less likely to create
DMIs than changes with opposing *cis-trans* effects (thought to result from stabilizing selection
within species) (Mack et al. 2016) (but see (Tulchinsky et al. 2014b)). Consequently, positive
selection affecting regulatory versus coding sequences may differ in their propensity to yield
DMIs.
414
Might the functional role of genes also represent an axis of predictability to factors that instigate
DMIs in inter-species hybrids? The relatively few known 'speciation genes' in animals give little
clue to whether particular molecular functions may be predisposed to involvement in inter-
species incompatibilities (Blackman 2016). At a coarse level, because DMIs require interaction,
we should expect sequences that affect interactions between DNA, RNA, and protein products to
be more prevalent among DMI loci than, say, enzymes that interact predominantly with
metabolites. From the perspective of gene regulatory networks, transcription factors experience
faster sequence evolution than other genes in the genome (Gilad et al. 2006, Haerty et al. 2008).
piRNA genes also turnover rapidly and are implicated in hybrid dysfunction (Assis and
Kondrashov 2009, Bagijn et al. 2012, Kelleher et al. 2012), along with other classes of regulatory
endogenous small RNAs (Li et al. 2016). Genes that act cell non-autonomously, as for secreted
proteins or diffusible signaling molecules, may tend to evolve slower if they also tend to exhibit
greater expression breadth and lower modularity, unless they are predisposed to co-evolutionary
dynamics. Reproductive isolation between species need not only involve classic regulators of
development, however, as attested by incompatibilities that commonly seem to involve
chromosome segregation and cyto-nuclear interactions (Blackman 2016, Lima et al. 2019).
Despite the fundamental role of the mitochondrial genome in energy metabolism, mitochondrial
genes experience adaptive molecular evolution (Bazin et al. 2006) and mito-nuclear

incompatibilities can provide important reproductive barriers between species (Hill 2015, Lima

et al. 2019). To the extent that some developmental stages may be more sensitive to the

disruption of chromosome segregation or mitochondrial function, such as phases of heightened

cell division or metabolism, such developmentally-tangential genetic pathways might

nevertheless contribute to ontogenetic patterns of hybrid dysfunction.

Gene duplication and gene origination: Gene duplication is a powerful factor in the evolution

of phenotypic novelty (Kaessmann 2010). Moreover, regulatory or structural

subfunctionalization in the evolution of gene copies following duplication could lead to

dysfunction in hybrids and so contribute to a DMI (Mack and Nachman 2017). Similarly, a DMI

could result from 'divergent resolution' in the loss of alternate copies in different species for

functionally equivalent gene duplicates (Lynch and Force 2000). The de novo origin of new

genes also can instigate species differences in gene network structure (Neme and Tautz 2014), as

could divergent use of alternative splice forms of a gene (Ortíz-Barrientos et al. 2007). Thus, the

phenotypic evolvability promoted by gene duplication and gene origination also confers the

potential to induce reproductive isolation as gene copies evolve through distinct trajectories in

different evolutionary lineages.

Ontogenetic timing in genetic network and molecular evolution

Gene expression profiles and genetic network architecture are dynamic over the course of

ontogeny. This dynamism suggests that molecular evolution could differ for the distinct subsets

of genes associated with different stages of development (Box 3). Importantly, this molecular

divergence may or may not correspond to divergence in organismal phenotypes (True and Haag

2001), despite the common emphasis on how changes to gene regulatory networks alter the

development of phenotypes (Erwin and Davidson 2009). Stages especially prone to rapid

molecular evolution, of coding sequences or regulatory elements, may also translate into a

greater incidence of DMIs. Ontogenetic predispositions toward sequence evolution may

therefore confer predictable ontogenetic detection of reproductive isolation between species,

letting us know when to listen most carefully for the developmental alarm clock of speciation. In

our search of the literature, we identified 147 studies involving 106 species that quantified

transcriptome, proteome, or related molecular data across multiple developmental stages (Table

1, Supplementary file 1). We found 40% of studies to focus on embryogenesis only. Over 60%

of the studies included a comparative analysis of expression divergence between species (52%)

or of DNA sequence divergence (13%). Integrating such studies with developmental time

courses of hybrid dysfunction would provide a powerful collection of empirical tests for how

incompatibilities in genetic networks arise through ontogeny. In the meantime, ideas from both

development and population genetics lead to several partially-distinct predictions for the

evolution of genes expressed differentially across ontogeny, which we enumerate below (Figure

1). We conclude that a conceptual gap is the lack of a modeling framework that integrates these

disparate perspectives on the role of ontogenetic timing in molecular evolution.

Early conservation model

The early conservation model ("von Baer's third law") derives from the fact that time is

unidirectional, so changes early in development may cascade catastrophically as cell division and

differentiation proceeds. Consequently, genes expressed early in development would experience

stronger purifying selection, with slower molecular and phenotypic evolution early in ontogeny

(Kalinka and Tomancak 2012). Some perspectives on gene regulatory network structure follow

the spirit of this view, as well (Erwin and Davidson 2009). This idea would also be consistent

with genes and genetic networks expressed in early development having lower robustness,

modularity, and evolvability, and with greater pleiotropy when perturbed (Figure 1). The early

conservation model is mostly applied to embryogenesis, with some empirical support from

vertebrates based on gene expression divergence (Roux and Robinson-Rechavi 2008, Irie and

Kuratani 2014). In principle, however, the logic of this model applies to the entirety of ontogeny.

If the likelihood of DMIs scales with rate of evolution, then we ought to expect post-zygotic

reproductive isolation to be more likely to manifest later in development and that species at

earlier stages of divergence would manifest defects later in development.

A caveat about inferring the developmental timing of hybrid dysfunction is that early-acting

incompatibilities may preclude detection of DMIs that would otherwise be revealed later (see

uniform chance model, below). Disproportionate observation of early-acting developmental

defects in hybrids thus may not imply that molecular evolution disproportionately accrues for

genes in early-acting developmental programs. This 'pull of the early' represents a general

challenge in characterizing the profile across ontogeny of disrupted genetic networks that confer

post-zygotic reproductive isolation. There are at least four ways to potentially address this issue

empirically: i) assess distinct species comparisons from different phylogenetic depths, ii) exploit

partial penetrance of F1 hybrid dysfunction (Bundus et al. 2015), iii) use early-acting hybrid-

rescue genotypes to test for late-acting hybrid dysfunction (cf. *Hmr* in *Drosophila* (Hutter et al.

1990)), or iv) focus on misregulation of gene expression over ontogeny for species with

relatively weak post-zygotic isolation (i.e. without catastrophic effects in early life stages).

Uniform chance model

We suggest that the 'uniform chance' model represents a null model for the manifestation of

hybrid incompatibilities over the course of ontogeny. If molecular evolution is unbiased with

respect to the timing of expression of genes that have diverged between species, then every point

in development can be considered to have an equal chance of expressing a DMI to yield hybrid

disruption of a tissue type or termination of development (Figure 1). This could arise from each

stage adapting continuously to the 'habitat' it experiences distinctively from other stages,

whether inside an egg (or uterus), juvenile environmental circumstances, or post-metamorphosis

adulthood. Consequently, species pairs with greater overall divergence would be more likely to

have hybrids that terminate development earlier in ontogeny; the probability that the earliest

developmental stage avoids the effects of DMIs declines exponentially (p^n with n DMIs each

with probability p of not occurring in the earliest stage). This null model therefore predicts a

negative relationship of the genetic distance between species and the terminal stage to which

hybrids develop, and predicts that later stages will have more tissue types exhibiting dysfunction

in hybrids. Note that these predictions overlap with several other models that we describe, but do

not depend on differential selection pressures, pleiotropy, or network features for genes

expressed at different times in development. This model, however, predicts no association of

molecular evolutionary rates for genes (or the mode of selection) with the developmental timing

of their expression.

Mutation accumulation model of aging

The mutation accumulation model of aging and senescence predicts that genes and genetic networks expressed earlier in development will evolve more slowly (Promislow and Tatar 1998, Partridge 2001). Here the logic and developmental timing, however, is different from the early conservation model: the reproductive value of individuals declines following the onset of reproductive maturity, meaning that purifying selection is weaker against deleterious mutations that affect adult phenotypes relative to embryonic and juvenile phenotypes (Medawar 1952, Flatt and Schmidt 2009). This perspective presumes similarly strong purifying selection for all genes expressed prior to maturity, and makes no specific prediction about positive selection across ontogeny (Figure 1). Not explicitly formulated by the mutation accumulation model, however, is whether the strong early-life selection would indirectly favor mutational robustness to embryonic and juvenile developmental programs. Such robustness could facilitate greater molecular evolution of early-expressed genes than otherwise anticipated, which also could facilitate the accumulation of developmental system drift.

Antagonistic pleiotropy model of aging

Gain in fitness from traits that increase survival or reproduction early in life will experience disproportionate selection pressure. Consequently, positive selection may more fiercely favor beneficial mutations to genes expressed early in life, irrespective of negative pleiotropic consequences on fitness-related traits expressed later on. This idea is the essence of the antagonistic pleiotropy model for the evolution of aging and senescence (Williams 1957, Partridge 2001, Flatt and Schmidt 2009). In terms of molecular evolution, it ought to yield a signature of excess positive selection on genes first expressed at pre-reproductive stages (Figure 1). However, subsequent selection may lead to compensatory evolution for genes expressed late in life to ameliorate negative pleiotropic effects. This sequential evolutionary series of events could produce developmental system drift, especially for late-life traits. Regulatory divergence between species often reflects compensatory changes in *cis* and *trans*, with *trans*-acting changes

being more likely to exert pleiotropic effects and arising at a higher rate (Gruber et al. 2012). It

will be interesting to determine whether the developmental timing of *cis* versus *trans* changes are

biased over ontogeny in a way that would be consistent with the antagonistic pleiotropy model,

i.e., disproportionate *cis*-regulatory changes that modulate late-acting expression.

Hourglass model

The hourglass model is a pattern in search of a mechanism that derives from classic phenotypic

observations of a 'phylotypic stage' in mid-embryogenesis, a point of greatest morphological

similarity across organisms. Profiles of gene expression divergence between species, a molecular

phenotype, also are often interpreted to be consistent with this pattern (Raff 1996, Kalinka and

Tomancak 2012). In terms of DNA sequence evolution, it is less obvious why genes expressed in

mid-embryogenesis would exhibit slowest rates of sequence evolution, whether due to stronger

purifying selection or less frequent adaptive divergence (Figure 1). Ontogenetic trends in genetic

network architecture may provide some intuition, as it is proposed that the phylotypic stage may

coincide with developmental periods of intense integration of distinct cell lineages, often around

gastrulation (Levin et al. 2012). Consequently, mid-embryogenesis may experience a shift

toward fewer and larger genetic networks (low modularity) that leads to greater pleiotropic

effects when perturbed. Changes in network structure related to switch-like or threshold traits

also could contribute (Garfield et al. 2013). If such a genetic network architecture in mid-

embryogenesis also confers greater within-species genetic robustness, then it may be especially

prone to DSD and the production of DMIs and hybrid dysfunction in crosses between species. If,

instead, it represents a point of low robustness to genetic perturbation, then DSD from genes

expressed at earlier stages may induce a tipping point of dysfunction that manifests in hybrids

during such a phase of integration.

The potential for distinctive properties in the 'waist of the hourglass' has drawn most scrutiny by

researchers, but an alternative hypothesis might suppose that it is the earlier points in

development that are unusual and require special explanation. This possibility has recently

received greater theoretical and empirical attention (Demuth and Wade 2007, Dapper and Wade

2016, Zalts and Yanai 2017, Coronado-Zamora et al. 2019). For example, theory predicts more

rapid accumulation of mutations to genes acting in the earliest genetic networks that derive from

maternal resources, leading to their unusually fast molecular evolution with the potential to drive

DSD and DMIs (Demuth and Wade 2007). Faster-than-anticipated molecular evolution at the

very earliest stages of development might also arise from especially high robustness of genetic

networks involving maternally-deposited gene products, or from co-evolutionary dynamics

associated with parent-offspring or other genomic conflicts (Brandvain and Haig 2005, Crespi

and Nosil 2013). In such scenarios, the point in time that coincides with 'waist of the hourglass'

might simply represent the onset of conditions consistent with the early conservation model.

Sex-biased selection

Selection on genes with sex-limited expression can lead to their more rapid evolution. This faster

evolution can arise in two ways: sexual selection/conflict and weaker purifying selection. Sexual

selection and sexual conflict that occurs within and between the sexes for reproductive adults can

promote rapid phenotypic and molecular evolution via positive selection, especially in

reproduction- and gamete-related traits (Swanson and Vacquier 2002, Mank 2017, Rowe et al.

2018). Thus, the timing of expression for genes that are part of the developmental programs that

build such traits in late juvenile and adult stages would be expected to be most strongly

impacted. This ontogenetic timing is similar to the mutation accumulation model, except that it

should reflect adaptive evolution rather than weaker purifying selection and that it deals

primarily with the subset of traits and genes associated with sexual interactions between

individuals (Figure 1). Genes controlling gamete traits may be especially prone to rapid

molecular evolution (Swanson and Vacquier 2002), with the effects on hybrids potentially

reflected in the earliest-stage zygotes. Speciation genetics research has incorporated this idea of

rapid evolution of sex-limited genes into the 'faster male' theory for Haldane's rule and into the

explanation for why hybrid sterility often appears to arise earlier in the speciation process than

does hybrid inviability (Coyne and Orr 1989, Wu and Davis 1993, Ortíz-Barrientos et al. 2007,

Presgraves 2010a).

In addition to the possibility of more prevalent positive selection on genes with sex-limited

expression, they may also experience weaker purifying selection (Dapper and Wade 2016).

Weaker purifying selection means greater accumulation of divergence between species. This

weaker efficacy of selection arises because half of the individuals in the population, one sex or

the other, do not express the gene and so mask the effects of new detrimental mutations. This

logic applies to both sex-limited gene expression for adult traits, as well as for maternally-

provisioned transcripts delivered to eggs (Demuth and Wade 2007, Dapper and Wade 2016).

Rapid evolution of maternal-provisioning genes may lead to incompatibilities when they interact

with zygotically-expressed genes in inter-species hybrids, potentially elevating the incidence of

mid-embryonic hybrid dysfunction independently of any temporal changes in genetic network

modularity or robustness. Genetic incompatibilities involving uniparentally-inherited genetic

factors also may lead to predictable asymmetries in reproductive isolation between species pairs

(Turelli and Moyle 2007), including roles for genes encoded on sex chromosomes and

mitochondrial genomes (Bolnick et al. 2008, Barreto et al. 2018, Cutter 2018).

Towards predictable rules in the evolution of development in speciation

To decipher how predictable molecular evolution and post-zygotic reproductive isolation might

be, it is important to consider the ontogenetic context of organisms to identify trends in the

tempo of gene expression, cell division, and differentiation (Table 1). The kind, number, and

molecular evolutionary implications of genetic interactions also likely are sensitive to classes of

embryogenesis programs (e.g., syncytial with late cellularization as in *Drosophila* and other

insects, totipotent versus highly cell-autonomous cell lineage development as in *C. elegans*,

nourishment by minimal vs very large yolk vs maternal tissues as seen in many insects, birds,

and mammals) (Church et al. 2019). While here we emphasize animal systems, similar

consideration of the evolutionary genetics of development and speciation for plant systems may

also reveal valuable insights (Rieseberg and Blackman 2010, Bedinger et al. 2011, Baack et al.

2015). Here we distill in abbreviated form what is known for a few concrete motivating example

systems (*Caenorhabditis* nematodes, *Drosophila* insects, *Bufo* toads) to frame these issues about

how ontogenetic features link to molecular evolution and hybrid dysfunction. Surprisingly, few

broad conclusions can be drawn from even these well-studied systems, and general principles

await concerted research efforts. Nevertheless, these study systems present clear promise for

future research to disentangle the constellation of causal contributing factors that link

microevolutionary mechanisms to developmental programs and macroevolutionary patterns.

C. elegans ontogenetic profiles of gene expression, molecular evolution and hybrid

dysfunction

Taking *C. elegans* development as a point of reference, embryonic cell numbers grow

approximately exponentially until ~520 cells upon which cell count increases relatively slowly to

the 959 somatic and ~2000 germline cells that comprise the adult hermaphrodite animal

(Giurumescu et al. 2012) (Figure 2A). Cell lineages derived from 5 blastomeres exhibit distinct

transcriptome profiles in ontogenetic timecourses (Hashimshony et al. 2014), with single-cell

transcriptome sequencing of embryos in a time series showing even finer resolution (Packer et al.

2019). The number of genes expressed increases over most of embryogenesis, with relatively

similar numbers of genes expressed at stages post-hatching (Boeck et al. 2016) (Figure 2B). The

change in identity of expressed genes, however, is greatest early in embryogenesis (primarily due

to down-regulation) and late in embryogenesis (primarily due to up-regulation) (Boeck et al.

2016) (Figure 2C). Gene connectivity peaks in early embryogenesis, declining until a spike in

adulthood (Liu and Robinson-Rechavi 2018a). Cellular development in embryos is conserved

across species, being nearly indistinguishable at least to the 350-cell stage (Zhao et al. 2008,

Memar et al. 2018). In hybrid crosses of most species pairs, however, embryos arrest around

gastrulation (Baird and Yen 2000, Baird and Seibert 2013), near the time that expression

divergence appears minimized across *Caenorhabditis* (Levin et al. 2012) and that expression-

weighted coding sequence divergence is lowest (Cutter et al. 2019). Coding sequences with

fastest molecular evolution show peak expression very early in embryogenesis or toward

adulthood (Cutter and Ward 2005, Cutter et al. 2019) (Box 3). Hybrids of *C. briggsae* and *C.*

nigoni show high embryonic inviability, but those genetically-identical individuals that hatch

successfully exhibit little larval mortality (Bundus et al. 2015). Gene misexpression is

widespread in adults of both sexes for these hybrids (Sanchez-Ramirez et al. 2020) and

misregulation of small-RNAs in hybrid genetic backgrounds is implicated in spermatogenic

dysfunction (Li et al. 2016). Together, these observations suggest the possibility that

developmental system drift may accrue more readily and generate hybrid incompatibility

disproportionately at stages showing greatest selective constraint on gene expression levels and

coding sequences.

Drosophila divergence in sequences and transcriptomes over ontogeny

Coding sequences expressed across *D. melanogaster* development show slowest evolution in

mid- to late-embryogenesis (Davis et al. 2005, Coronado-Zamora et al. 2019). The faster

sequence evolution of maternally-deposited transcripts in early embryogenesis is due

disproportionately to non-adaptive divergence whereas it is adaptive divergence that is

disproportionately implicated in faster sequence evolution of genes expressed post-

embryonically (Coronado-Zamora et al. 2019) (Figure 2E). Gene connectivity similarly shows a

peak in early embryogenesis and in adulthood (Liu and Robinson-Rechavi 2018a).

Transcriptome divergence across species, however, is low throughout most of embryogenesis

and highest after reproductive maturity; within embryogenesis, however, expression divergence

is greatest at the earliest stages (Kalinka et al. 2010, Liu and Robinson-Rechavi 2018a). Sterility

tends to arise before inviability in hybrids of a given phylogenetic distance in *Drosophila* (Coyne

and Orr 1997), and hybrid misexpression among spermatogenesis genes is much more

pronounced in adults than in late-stage larvae (Moehring et al. 2007). Genes with greater tissue-

specificity show faster coding sequence evolution (Larracuente et al. 2008). Hybrid

misexpression less often involves genes implicated in transcriptional regulation than expected

(Moehring et al. 2007), and defects of chromosome condensation in mitosis, nucleoporins, and

transcriptional regulators of selfish elements have been implicated in the inviability of hybrid

larvae (Orr et al. 1997, Barbash et al. 2003, Tang and Presgraves 2009, Satyaki et al. 2014). It

will be interesting for future work to assess whether those embryonic stages that show greatest

conservation in expression and coding sequence evolution might show disproportionate hybrid

dysfunction, as hinted from experiments in *Caenorhabditis*. The syncytial nature of *Drosophila*

embryos through the ~6,000-cell stage, however, may lead to distinct genetic network

architecture and expectations in the timing of selective pressures on gene products deposited

maternally versus expressed zygotically across embryogenesis.

Bufo ontogenetic profiles of hybrid dysfunction

The developmental timing of hybrid inviability in frogs and toads enjoys much richer literature

than for many other systems. Hybrid developmental data exist for several genera, including *Hyla*

(Fouquette 1960, Mecham 1960, 1965, Kuramoto 1984, Kawamura et al. 1990), *Pseudacris*

(Mecham 1965), *Rana* (Kuramoto 1974, Frost and Platz 1983, Sumida et al. 2003), and most

notably *Bufo* (Blair 1972, Malone and Fontenot 2008). The developmental stages of hybrid

inviability are described for over 600 inter-species *Bufo* cross combinations, with late-stage

hybrid dysfunction being most prevalent for species pairs that are less genetically divergent

(Figure 2). By contrast, genome and transcriptome information is rare, due to the large genome

sizes of most frogs and toads. One key exception among anurans is for the model species

Xenopus laevis and *X. tropicalis*: comparative transcriptome analysis across their embryonic

development showed changes in expression profiles of many genes despite a background of

conservation in expression for most genes (Yanai et al. 2011). Interestingly, earlier stages of

embryogenesis showed greater divergence in gene expression (Figure 2). Post-embryonic

development was not analyzed, however, making it unclear what profile of gene expression

divergence describes all of ontogeny. It will be valuable in future work to link directly the

ontogeny of hybrid dysfunction to the degree of divergence in transcriptomes and sequences for

genes expressed differentially across development.

Conclusions

A general understanding of genetic mechanisms in the speciation process integrates the

ontogenetic timing of gene expression and gene action in developmental programs. Predictable

ontogenetic trends in the molecular evolution of proteins and their regulation will introduce

predictability into how genetic networks diverge and how Dobzhansky-Muller incompatibilities

manifest dysfunctional phenotypes in hybrids. In this underexploited way, the genetics of post-

zygotic isolation in the speciation process dovetails with research programs in evolutionary

developmental genetics. The diversity of theoretical perspectives that contribute predictions to

ontogenetic patterns of molecular evolution are, however, at best, incompletely integrated with

conceptions about genetic architectures (pleiotropy, modularity, robustness). We still have only a

rudimentary understanding of how the genetic clockwork might set the ontogenetic timing of this

developmental alarm clock in the dawn of new species. Consequently, it is a challenge to extract

consensus on patterns and predictions about sequence evolution and DMI incidence over the

course of ontogeny.

Key features of an ontogenetic view of molecular evolution include the pleiotropic roles of genes

and the pleiotropic effects of genetic perturbation, as well as the dynamism of genetic network

modularity over development, to influence the evolvability of genes and the robustness of

phenotypic outputs. A growing body of empirical literature is documenting the dynamics of gene

expression and molecular evolution over developmental time. What is missing in the nascent

state of the field is an integrated set of theoretical expectations for how these features can

produce emergent trends of genome evolution and of dysfunctional genetic networks in the

divergent set of genomes of hybrid individuals. Empirical tests of existing theoretical

predictions, from models that focus primarily on only a subset of ontogeny, will benefit from

exploring the relative influence of adaptive molecular evolution and purifying selection on genes

that vary in expression over all ontogeny. Answering the neglected question -- "Are certain

developmental processes especially likely to be disrupted in hybrids?" -- offers the promise to

identify new rules of speciation as well as rules in the molecular evolution of development.

Acknowledgements

This research was supported by a Discovery Grant from the Natural Sciences and Engineering

Research Council (NSERC) of Canada to A.D.C. We are grateful for the constructive comments

by reviewers on an earlier draft of this manuscript.

Literature cited

Artieri, C. G., and H. B. Fraser. 2014. Evolution at two levels of gene expression in yeast. Genome Research **24**:411-421.

Assis, R., and A. S. Kondrashov. 2009. Rapid repetitive element-mediated expansion of piRNA clusters in mammalian evolution. Proceedings of the National Academy of Sciences **106**:7079.

Baack, E., M. C. Melo, L. H. Rieseberg, and D. Ortiz-Barrientos. 2015. The origins of reproductive isolation in plants. New Phytologist **207**:968-984.

Bagijn, M. P., L. D. Goldstein, A. Sapetschnig, E.-M. Weick, S. Bouasker, N. J. Lehrbach, M. J. Simard, and E. A. Miska. 2012. Function, targets, and evolution of *Caenorhabditis elegans* piRNAs. Science **337**:574.

Baird, S. E., and S. R. Seibert. 2013. Reproductive isolation in the Elegans-Group of *Caenorhabditis*. Natural Science **5**:18-25.

Baird, S. E., and W.-C. Yen. 2000. Reproductive isolation in *Caenorhabditis*: terminal phenotypes of hybrid embryos. Evolution and Development **2**:9-15.

Barbash, D. A., D. F. Siino, A. M. Tarone, and J. Roote. 2003. A rapidly evolving MYB-related protein causes species isolation in *Drosophila*. Proceedings of the National Academy of Sciences of the United States of America **100**:5302-5307.

Barreto, F. S., E. T. Watson, T. G. Lima, C. S. Willett, S. Edmands, W. Li, and R. S. Burton. 2018. Genomic signatures of mitonuclear coevolution across populations of Tigriopus californicus. Nature Ecology & Evolution **2**:1250-1257.

Batada, N. N., L. D. Hurst, and M. Tyers. 2006. Evolutionary and physiological importance of hub proteins. PLoS Computational Biology **2**:e88.

Bazin, E., S. Glemin, and N. Galtier. 2006. Population size does not influence mitochondrial genetic diversity in animals. Science **312**:570-572.

Bedinger, P. A., R. T. Chetelat, B. McClure, L. C. Moyle, J. K. Rose, S. M. Stack, E. van der Knaap, Y. S. Baek, G. Lopez-Casado, P. A. Covey, A. Kumar, W. Li, R. Nunez, F. Cruz-Garcia, and S. Royer. 2011. Interspecific reproductive barriers in the tomato clade: opportunities to decipher mechanisms of reproductive isolation. Sex Plant Reprod **24**:171-187.

Blackman, B. K. 2016. Speciation genes. Pages 166-175 *in* R. M. Kliman, editor. Encyclopedia of Evolutionary Biology. Academic Press, Oxford.

Blair, W. F. 1972. Evolution in the genus *Bufo*. University of Texas Press.

Bloom, J. D., and C. Adami. 2003. Apparent dependence of protein evolutionary rate on number of interactions is linked to biases in protein-protein interactions data sets. BMC Evolutionary Biology **3**:1-10.

Boeck, M. E., C. Huynh, L. Gevirtzman, O. A. Thompson, G. Wang, D. M. Kasper, V. Reinke, L. W. Hillier, and R. H. Waterston. 2016. The time-resolved transcriptome of *C. elegans*. Genome Research **26**:1441-1450.

Bolnick, D. I., M. Turelli, H. Lopez-Fernandez, P. C. Wainwright, and T. J. Near. 2008. Accelerated mitochondrial evolution and "Darwin's corollary": asymmetric viability of reciprocal F1 hybrids in Centrarchid fishes. Genetics **178**:1037-1048.

Brandvain, Y., and D. Haig. 2005. Divergent mating systems and parental conflict as a barrier to hybridization in flowering plants. American Naturalist **166**:330-338.

Bundus, J. D., R. Alaei, and A. D. Cutter. 2015. Gametic selection, developmental trajectories and extrinsic heterogeneity in Haldane's rule. Evolution **69**:2005-2017.

Carroll, S. B. 2005. Evolution at two levels: on genes and form. PLoS Biology **3**:e245.

811   Casillas, S., A. Barbadilla, and C. M. Bergman. 2007. Purifying selection maintains highly conserved
812   noncoding sequences in Drosophila. Molecular Biology and Evolution **24**:2222-2234.
813   Castillo-Davis, C. I., D. L. Hartl, and G. Achaz. 2004. *cis*-Regulatory and protein evolution in orthologous
814   and duplicate genes. Genome Research **14**:1530-1536.
815   Charlesworth, B., J. A. Coyne, and N. H. Barton. 1987. The relative rates of evolution of sex-
816   chromosomes and autosomes. American Naturalist **130**:113-146.
817   Chin-Sang, I. D., and A. D. Chisholm. 2000. Form of the worm:: genetics of epidermal morphogenesis in
818   C. elegans. Trends in Genetics **16**:544-551.
819   Church, S. H., S. Donoughe, B. A. S. de Medeiros, and C. G. Extavour. 2019. Insect egg size and shape
820   evolve with ecology but not developmental rate. Nature **571**:58-62.
821   Civetta, A. 2016. Misregulation of gene expression and sterility in interspecies hybrids: causal links and
822   alternative hypotheses. Journal of Molecular Evolution **82**:176-182.
823   Clark, N. L., J. Gasper, M. Sekino, S. A. Springer, C. F. Aquadro, and W. J. Swanson. 2009. Coevolution of
824   interacting fertilization proteins. PLoS Genetics **5**:e1000570.
825   Coronado-Zamora, M., I. Salvador-Martínez, D. Castellano, A. Barbadilla, and I. Salazar-Ciudad. 2019.
826   Adaptation and conservation throughout the *Drosophila melanogaster* life-cycle. Genome
827   Biology and Evolution:10.1093/gbe/evz1086.
828   Coyne, J. A., and H. A. Orr. 1989. Patterns of speciation in Drosophila. Evolution **43**:362-381.
829   Coyne, J. A., and H. A. Orr. 1997. "Patterns of speciation in Drosophila" revisited. Evolution **51**:295-303.
830   Coyne, J. A., and H. A. Orr. 2004. Speciation. Sinauer, Sunderland, MA.
831   Crespi, B., and P. Nosil. 2013. Conflictual speciation: species formation via genomic conflict. Trends in
832   Ecology & Evolution **28**:48-57.
833   Cutter, A. D. 2012. The polymorphic prelude to Bateson-Dobzhansky-Muller incompatibilities. Trends in
834   Ecology & Evolution **27**:209-218.
835   Cutter, A. D. 2018. X exceptionalism in Caenorhabditis speciation. Molecular Ecology **27**:3925-3934.
836   Cutter, A. D., R. H. Garrett, S. Mark, W. Wang, and L. Sun. 2019. Molecular evolution across
837   developmental time reveals rapid divergence in early embryogenesis. Evolution Letters **3**:359-
838   373.
839   Cutter, A. D., and S. Ward. 2005. Sexual and temporal dynamics of molecular evolution in *C. elegans*
840   development. Molecular Biology and Evolution **22**:178-188.
841   Dapper, A. L., and M. J. Wade. 2016. The evolution of sperm competition genes: The effect of mating
842   system on levels of genetic variation within and between species. Evolution **70**:502-511.
843   Davis, J. C., O. Brandman, and D. A. Petrov. 2005. Protein evolution in the context of Drosophila
844   development. Journal of Molecular Evolution **60**:774-785.
845   Delph, L. F., and J. P. Demuth. 2016. Haldane's rule: genetic bases and their empirical support. Journal of
846   Heredity **107**:383-391.
847   Demuth, J. P., and M. J. Wade. 2007. Maternal expression increases the rate of bicoid evolution by
848   relaxing selective constraint. Genetica **129**:37-43.
849   Ebbing, A., Á. Vértesy, M. C. Betist, B. Spanjaard, J. P. Junker, E. Berezikov, A. van Oudenaarden, and H.
850   C. Korswagen. 2018. Spatial transcriptomics of *C. elegans* males and hermaphrodites identifies
851   sex-specific differences in gene expression patterns. Developmental Cell **47**:801-813.e806.
852   Ehrenreich, I. M., N. Torabi, Y. Jia, J. Kent, S. Martis, J. A. Shapiro, D. Gresham, A. A. Caudy, and L.
853   Kruglyak. 2010. Dissection of genetically complex traits with extremely large pools of yeast
854   segregants. Nature **464**:1039-U1101.
855   Erwin, D. H., and E. H. Davidson. 2009. The evolution of hierarchical gene regulatory networks. Nature
856   Reviews Genetics **10**:141-148.

Estes, S., and S. J. Arnold. 2007. Resolving the paradox of stasis: Models with stabilizing selection explain
evolutionary divergence on all timescales. American Naturalist **169**:227-244.
Farhadifar, R., J. M. Ponciano, E. C. Andersen, D. J. Needleman, and C. F. Baer. 2016. Mutation is a
sufficient and robust predictor of genetic variation for mitotic spindle traits in *Caenorhabditis*
elegans. Genetics **203**:1859.
Felix, M. A., and A. Wagner. 2008. Robustness and evolution: concepts, insights and challenges from a
developmental model system. Heredity **100**:132-140.
Flatt, T., and P. S. Schmidt. 2009. Integrating evolutionary and molecular genetics of aging. Biochimica et
Biophysica Acta (BBA) - General Subjects **1790**:951-962.
Fouquette, M. J. 1960. Isolating mechanisms in three sympatric treefrogs in the canal zone. Evolution
14:484-497.
Fragata, I., A. Blanckaert, M. A. Dias Louro, D. A. Liberles, and C. Bank. 2019. Evolution in the light of
fitness landscape theory. Trends in Ecology & Evolution **34**:69-82.
Frost, J. S., and J. E. Platz. 1983. Comparative assessment of modes of reproductive isolation among four
species of leopard frogs (*Rana pipiens* complex). Evolution **37**:66-78.
Garfield, D. A., D. E. Runcie, C. C. Babbitt, R. Haygood, W. J. Nielsen, and G. A. Wray. 2013. The impact of
gene expression variation on the robustness and evolvability of a developmental gene
regulatory network. PLoS Biology **11**:e1001696.
Gavrilets, S. 2003. Models of speciation: what have we learned in 40 years? Evolution **57**:2197-2215.
Gilad, Y., A. Oshlack, G. K. Smyth, T. P. Speed, and K. P. White. 2006. Expression profiling in primates
reveals a rapid evolution of human transcription factors. Nature **440**:242-245.
Giurumescu, C. A., S. Kang, T. A. Planchon, E. Betzig, J. Bloomekatz, D. Yelon, P. Cosman, and A. D.
Chisholm. 2012. Quantitative semi-automated analysis of morphogenesis with single-cell
resolution in complex embryos. Development **139**:4271-4279.
Goncalves, A., S. Leigh-Brown, D. Thybert, K. Stefflova, E. Turro, P. Flicek, A. Brazma, D. T. Odom, and J.
C. Marioni. 2012. Extensive compensatory cis-trans regulation in the evolution of mouse gene
expression. Genome Research **22**:2376-2384.
Gruber, J. D., K. Vogel, G. Kalay, and P. J. Wittkopp. 2012. Contrasting properties of gene-specific
regulatory, coding, and copy number mutations in *Saccharomyces cerevisiae*: frequency, effects,
and dominance. PLoS Genetics **8**:e1002497.
Guerrero, R. F., C. D. Muir, S. Josway, and L. C. Moyle. 2017. Pervasive antagonistic interactions among
hybrid incompatibility loci. PLoS Genetics **13**:e1006817.
Haag, E. S. 2007. Compensatory vs. pseudocompensatory evolution in molecular and developmental
interactions. Genetica **129**:45-55.
Haag, E. S., and M. N. Molla. 2005. Compensatory evolution of interacting gene products through
multifunctional intermediates. Evolution **59**:1620-1632.
Haerty, W., C. Artieri, N. Khezri, R. S. Singh, and B. P. Gupta. 2008. Comparative analysis of function and
interaction of transcription factors in nematodes: extensive conservation of orthology coupled
to rapid sequence evolution. BMC Genomics **9**:399.
Hahn, M. W., and A. D. Kern. 2005. Comparative genomics of centrality and essentiality in three
eukaryotic protein-interaction networks. Molecular Biology and Evolution **22**:803-806.
Haldane, J. B. S. 1922. Sex ratio and unisexual sterility in hybrid animals. Journal of Genetics **12**:101-109.
Hashimshony, T., M. Feder, M. Levin, B. K. Hall, and I. Yanai. 2014. Spatiotemporal transcriptomics
reveals the evolutionary history of the endoderm germ layer. Nature **519**:219.
Hashimshony, T., F. Wagner, N. Sher, and I. Yanai. 2012. CEL-Seq: single-cell RNA-Seq by multiplexed
linear amplification. Cell Reports **2**:666-673.
He, X., and J. Zhang. 2006. Toward a molecular understanding of pleiotropy. Genetics **173**:1885-1891.

Hermisson, J., and G. P. Wagner. 2004. The population genetic theory of hidden variation and genetic
robustness. Genetics **168**:2271.
Hill, G. E. 2015. Mitonuclear ecology. Molecular Biology and Evolution **32**:1917-1927.
Hutter, P., J. Roote, and M. Ashburner. 1990. A genetic basis for the inviability of hybrids between sibling
species of *Drosophila*. Genetics **124**:909.
Irie, N., and S. Kuratani. 2014. The developmental hourglass model: a predictor of the basic body plan?
Development **141**:4649-4655.
Johnson, N. A., and A. H. Porter. 2001. Toward a new synthesis: Population genetics and evolutionary
developmental biology. Genetica **112-113**:45-58.
Johnson, N. A., and A. H. Porter. 2007. Evolution of branched regulatory genetic pathways: directional
selection on pleiotropic loci accelerates developmental system drift. Genetica **129**:57-70.
Jordan, I. K., Y. I. Wolf, and E. V. Koonin. 2003. No simple dependence between protein evolution rate
and the number of protein-protein interactions: only the most prolific interactors tend to evolve
slowly. BMC Evolutionary Biology **3**:1.
Kaessmann, H. 2010. Origins, evolution, and phenotypic impact of new genes. Genome Research
20:1313-1326.
Kahali, B., S. Ahmad, and T. C. Ghosh. 2009. Exploring the evolutionary rate differences of party hub and
date hub proteins in *Saccharomyces cerevisiae* protein–protein interaction network. Gene
429:18-22.
Kalinka, A. T., and P. Tomancak. 2012. The evolution of early animal embryos: conservation or
divergence? Trends in Ecology & Evolution **27**:385-393.
Kalinka, A. T., K. M. Varga, D. T. Gerrard, S. Preibisch, D. L. Corcoran, J. Jarrells, U. Ohler, C. M. Bergman,
and P. Tomancak. 2010. Gene expression divergence recapitulates the developmental hourglass
model. Nature **468**:811-U102.
Kawamura, T., M. Nishioka, and H. Ueda. 1990. Reproductive isolation in treefrogs distributed in Japan,
Korea, Europe and America. Scientific Reports of the Laboratory Amphibian Biology of Hiroshima
University **10**:255-293.
Kelleher, E. S., N. B. Edelman, and D. A. Barbash. 2012. *Drosophila* interspecific hybrids phenocopy
piRNA-pathway mutants. PLoS Biology **10**:e1001428.
Khan, Z., M. J. Ford, D. A. Cusanovich, A. Mitrano, J. K. Pritchard, and Y. Gilad. 2013. Primate transcript
and protein expression levels evolve under compensatory selection pressures. Science
342:1100.
Kim, B., B. Suo, and Scott W. Emmons. 2016. Gene function prediction based on developmental
transcriptomes of the two sexes in *C. elegans*. Cell Reports **17**:917-928.
Kitami, T., and J. H. Nadeau. 2002. Biochemical networking contributes more to genetic buffering in
human and mouse metabolic pathways than does gene duplication. Nature Genetics **32**:191-
194.
Kuramoto, M. 1974. Experimental hybridization between brown frogs of Taiwan, Ryukyu Islands and
Japan. Copeia:815-822.
Kuramoto, M. 1984. Systematic implications of hybridization experiments with some Eurasian treefrogs
(genus *Hyla*). Copeia:609-616.
Landry, C. R., B. Lemos, S. A. Rifkin, W. J. Dickinson, and D. L. Hartl. 2007. Genetic properties influencing
the evolvability of gene expression. Science **317**:118.
Landry, C. R., P. J. Wittkopp, C. H. Taubes, J. M. Ranz, A. G. Clark, and D. L. Hartl. 2005. Compensatory
cis-trans evolution and the dysregulation of gene expression in interspecific hybrids of
Drosophila. Genetics **171**:1813-1822.

Larracuente, A. M., T. B. Sackton, A. J. Greenberg, A. Wong, N. D. Singh, D. Sturgill, Y. Zhang, B. Oliver,
and A. G. Clark. 2008. Evolution of protein-coding genes in Drosophila. Trends in Genetics
24:114-123.
Ledón-Rettig, C. C., D. W. Pfennig, A. J. Chunco, and I. Dworkin. 2014. Cryptic genetic variation in natural
populations: a predictive framework. Integrative and Comparative Biology **54**:783-793.
Leducq, J.-B., G. Charron, G. Diss, I. Gagnon-Arsenault, A. K. Dubé, and C. R. Landry. 2012. Evidence for
the Robustness of Protein Complexes to Inter-Species Hybridization. PLoS Genetics **8**:e1003161.
Levin, M., T. Hashimshony, F. Wagner, and I. Yanai. 2012. Developmental milestones punctuate gene
expression in the *Caenorhabditis* embryo. Developmental Cell **22**:1101-1108.
Li, R., X. Ren, Y. Bi, V. W. S. Ho, C.-L. Hsieh, A. Young, Z. Zhang, T. Lin, Y. Zhao, L. Miao, P. Sarkies, and Z.
Zhao. 2016. Specific down-regulation of spermatogenesis genes targeted by 22G RNAs in hybrid
sterile males associated with an X-Chromosome introgression. Genome Research **26**:1219-1232.
Liao, B.-Y., and J. Zhang. 2006. Evolutionary conservation of expression profiles between human and
mouse orthologous genes. Molecular Biology and Evolution **23**:530-540.
Lima, T. G., R. S. Burton, and C. S. Willett. 2019. Genomic scans reveal multiple mito-nuclear
incompatibilities in population crosses of the copepod *Tigriopus californicus*. Evolution **73**:609-
620.
Liu, J., and M. Robinson-Rechavi. 2018a. Adaptive evolution of animal proteins over development:
support for the Darwin selection opportunity hypothesis of evo-devo. Molecular Biology and
Evolution **35**:2862-2872.
Liu, J., and M. Robinson-Rechavi. 2018b. Developmental constraints on genome evolution in four
bilaterian model species. Genome Biology and Evolution **10**:2266-2277.
Lynch, M., and A. G. Force. 2000. The origin of interspecific genomic incompatibility via gene duplication.
American Naturalist **156**:590-605.
Mack, K. L., P. Campbell, and M. W. Nachman. 2016. Gene regulation and speciation in house mice.
Genome Research **26**:451-461.
Mack, K. L., and M. W. Nachman. 2017. Gene regulation and speciation. Trends in Genetics **33**:68-80.
Malone, J. H., and B. E. Fontenot. 2008. Patterns of reproductive isolation in toads. PLoS ONE **3**:e3900.
Mank, J. E. 2017. Population genetics of sexual conflict in the genomic era. Nature Reviews Genetics
18:721.
Martin, A., and V. Orgogozo. 2013. The loci of repeated evolution: a catalog of the genetic hotspots of
phenotypic variation. Evolution **67**:1235-1250.
Masly, J. P., and D. C. Presgraves. 2007. High-resolution genome-wide dissection of the two rules of
speciation in Drosophila. Public Library of Science Biology **5**:e243.
McManus, C. J., J. D. Coolon, M. O. Duff, J. Eipper-Mains, B. R. Graveley, and P. J. Wittkopp. 2010.
Regulatory divergence in Drosophila revealed by mRNA-seq. Genome Research **20**:816-825.
McManus, C. J., G. E. May, P. Spealman, and A. Shteyman. 2014. Ribosome profiling reveals post-
transcriptional buffering of divergent gene expression in yeast. Genome Research **24**:422-430.
Mecham, J. S. 1960. Introgressive hybridization between two southeastern treefrogs. Evolution **14**:445-
457.
Mecham, J. S. 1965. Genetic relationships and reproductive isolation in southeastern frogs of genera
Pseudacris and *Hyla*. American Midland Naturalist **74**:269-&.
Medawar, P. B. 1952. An Unsolved Problem of Biology. H.K. Lewis & Co., London.
Memar, N., S. Schiemann, C. Hennig, D. Findeis, B. Conradt, and R. Schnabel. 2018. Twenty million years
of evolution: The embryogenesis of four *Caenorhabditis* species are indistinguishable despite
extensive genome divergence. Developmental Biology **447**:182-199.

Moehring, A. J., K. C. Teeter, and M. A. F. Noor. 2007. Genome-wide patterns of expression in *Drosophila*
pure species and hybrid males. II. Examination of multiple-species hybridizations, platforms, and
life cycle stages. Molecular Biology and Evolution **24**:137-145.
Neme, R., and D. Tautz. 2014. Evolution: dynamics of de novo gene emergence. Current Biology
24:R238-R240.
Ohta, T. 2011. Near-neutrality, robustness, and epigenetics. Genome Biology and Evolution **3**:1034-
1038.
Orr, H. A. 1995. The population genetics of speciation: the evolution of hybrid incompatibilities. Genetics
139:1805-1813.
Orr, H. A., L. D. Madden, J. A. Coyne, R. Goodwin, and R. S. Hawley. 1997. The developmental genetics of
hybrid inviability: a mitotic defect in *Drosophila* hybrids. Genetics **145**:1031.
Ortíz-Barrientos, D., B. A. Counterman, and M. A. F. Noor. 2007. Gene expression divergence and the
origin of hybrid dysfunctions. Genetica **129**:71-81.
Paaby, A. B., and M. V. Rockman. 2014. Cryptic genetic variation: evolution's hidden substrate. Nature
Reviews Genetics **15**:247-258.
Paaby, A. B., and N. D. Testa. 2018. Developmental plasticity and evolution. Pages 1-14 *in* L. Nuno de la
Rosa and G. Müller, editors. Evolutionary Developmental Biology: A Reference Guide. Springer
International Publishing, Cham.
Packer, J. S., Q. Zhu, C. Huynh, P. Sivaramakrishnan, E. Preston, H. Dueck, D. Stefanik, K. Tan, C. Trapnell,
J. Kim, R. H. Waterston, and J. I. Murray. 2019. A lineage-resolved molecular atlas of *C. elegans*
embryogenesis at single-cell resolution. Science **365**:eaax1971.
Palmer, M. E., and M. W. Feldman. 2009. Dynamics of hybrid incompatibility in gene networks in a
constant environemnt. Evolution **63**:418-431.
Partridge, L. 2001. Evolutionary theories of ageing applied to long-lived organisms. Experimental
Gerontology **36**:641-650.
Pavlicev, M., and G. P. Wagner. 2012. A model of developmental evolution: selection, pleiotropy and
compensation. Trends in Ecology & Evolution **27**:316-322.
Porter, A. H., and N. A. Johnson. 2002. Speciation despite gene flow when developmental pathways
evolve. Evolution **56**:2103-2111.
Presgraves, D. C. 2010a. Darwin and the origin of interspecific genetic incompatibilities. American
Naturalist **176 Suppl 1**:S45-S60.
Presgraves, D. C. 2010b. The molecular evolutionary basis of species formation. Nature Reviews
Genetics **11**:175-180.
Promislow, D. E. L. 2004. Protein networks, pleiotropy and the evolution of senescence. Proceedings of
the Royal Society of London. Series B: Biological Sciences **271**:1225-1234.
Promislow, D. E. L., and M. Tatar. 1998. Mutation and senescence: where genetics and demography
meet. Genetica **102/103**:299-314.
Proulx, S. R., S. Nuzhdin, and D. E. L. Promislow. 2007. Direct Selection on Genetic Robustness Revealed
in the Yeast Transcriptome. PLoS ONE **2**:e911.
Quint, M., H.-G. Drost, A. Gabel, K. K. Ullrich, M. Bönn, and I. Grosse. 2012. A transcriptomic hourglass in
plant embryogenesis. Nature **490**:98-101.
Raff, R. A. 1996. The Shape of Life: Genes, Development, and the Evolution of Animal Form. University of
Chicago Press, Chicago.
Rieseberg, L. H., and B. K. Blackman. 2010. Speciation genes in plants. Annals of Botany **106**:439-455.
Roux, J., and M. Robinson-Rechavi. 2008. Developmental constraints on vertebrate genome evolution.
PLoS Genetics **4**:e1000311.

Rowe, L., S. F. Chenoweth, and A. F. Agrawal. 2018. The genomics of sexual conflict. The American
Naturalist **192**:274-286.
Sanchez-Ramirez, S., J. G. Weiss, C. G. Thomas, and A. D. Cutter. 2020. Sex-specific and sex-chromosome
regulatory evolution underlie widespread misregulation of inter-species hybrid transcriptomes.
bioRxiv:10.1101/2020.1105.1104.076505.
Satyaki, P. R. V., T. N. Cuykendall, K. H. C. Wei, N. J. Brideau, H. Kwak, S. Aruna, P. M. Ferree, S. Ji, and D.
A. Barbash. 2014. The *Hmr* and *Lhr* hybrid incompatibility genes suppress a broad range of
heterochromatic repeats. PLoS Genetics **10**:e1004240.
Schluter, D. 1996. Adaptive radiation along genetic lines of least resistance. Evolution **50**:1766-1774.
Siegal, M. L., D. E. L. Promislow, and A. Bergman. 2007. Functional and evolutionary inference in gene
networks: does topology matter? Genetica **129**:83-103.
Signor, S. A., and S. V. Nuzhdin. 2018. The evolution of gene expression in cis and trans. Trends in
Genetics **34**:532-544.
Smith, E. N., and L. Kruglyak. 2008. Gene–environment interaction in yeast gene expression. PLoS
Biology **6**:e83.
Stern, D. L., and V. Orgogozo. 2008. The loci of evolution: how predictable is genetic evolution?
Evolution **62**:2155-2177.
Sucena, É., and D. L. Stern. 2000. Divergence of larval morphology between *Drosophila sechellia* and its
sibling species caused by cis-regulatory evolution of *ovo/shaven-baby*. Proceedings of the
National Academy of Sciences USA **97**:4530.
Sumida, M., H. Ueda, and M. Nishioka. 2003. Reproductive isolating mechanisms and molecular
phylogenetic relationships among palearctic and oriental brown frogs. Zoological Science
20:567-580.
Swanson, W. J., and V. D. Vacquier. 2002. The rapid evolution of reproductive proteins. Nature Reviews
Genetics **3**:137-144.
Takahasi, K. R., T. Matsuo, and T. Takano-Shimizu-Kouno. 2011. Two types of cis-trans compensation in
the evolution of transcriptional regulation. Proceedings of the National Academy of Sciences
USA **108**:15276-15281.
Tang, S., and D. C. Presgraves. 2009. Evolution of the Drosophila nuclear pore complex results in multiple
hybrid incompatibilities. Science **323**:779-782.
Tintori, Sophia C., E. Osborne Nishimura, P. Golden, Jason D. Lieb, and B. Goldstein. 2016. A
transcriptional lineage of the early *C. elegans* embryo. Developmental Cell **38**:430-444.
Tirosh, I., and N. Barkai. 2008. Evolution of gene sequence and gene expression are not correlated in
yeast. Trends in Genetics **24**:109-113.
Tirosh, I., S. Reikhav, A. A. Levy, and N. Barkai. 2009. A yeast hybrid provides insight into the evolution of
gene expression regulation. Science **324**:659.
Tran, T.-D., and Y.-K. Kwon. 2013. The relationship between modularity and robustness in signalling
networks. Journal of The Royal Society Interface **10**:20130771.
True, J. R., and E. S. Haag. 2001. Developmental system drift and flexibility in evolutionary trajectories.
Evolution and Development **3**:109-119.
Tulchinsky, A. Y., N. A. Johnson, and A. H. Porter. 2014a. Hybrid incompatibility despite pleiotropic
constraint in a sequence-based bioenergetic model of transcription factor binding. Genetics
198:1645-1654.
Tulchinsky, A. Y., N. A. Johnson, W. B. Watt, and A. H. Porter. 2014b. Hybrid incompatibility arises in a
sequence-based bioenergetic model of transcription factor binding. Genetics **198**:1155-1166.
Turelli, M., and L. C. Moyle. 2007. Asymmetric postmating isolation: Darwin's corollary to Haldane's rule.
Genetics **176**:1059-1088.

Turelli, M., and H. A. Orr. 1995. The dominance theory of Haldane's rule. Genetics **140**:389-402.
Turner, L. M., M. A. White, D. Tautz, and B. A. Payseur. 2014. Genomic networks of hybrid sterility. PLoS
1091   Genetics **10**:e1004162.
1092   Wade, M. J., N. A. Johnson, R. Jones, V. Siguel, and M. McNaughton. 1997. Genetic variation segregating
1093   in natural populations of *Tribolium castaneum* affecting traits observed in hybrids with *T.*
1094   *freemani*. Genetics **147**:1235-1247.
1095   Wagner, A. 2000. Robustness against mutations in genetic networks of yeast. Nature Genetics **24**:355-
1096   361.
1097   Wagner, G. P., and L. Altenberg. 1996. Complex adaptations and the evolution of evolvability. Evolution
1098   **50**:967-976.
1099   Wagner, G. P., and J. Z. Zhang. 2011. The pleiotropic structure of the genotype-phenotype map: the
1100   evolvability of complex organisms. Nature Reviews Genetics **12**:204-213.
1101   Wang, Z., X. Sun, Y. Zhao, X. Guo, H. Jiang, H. Li, and Z. Gu. 2015. Evolution of gene regulation during
1102   transcription and translation. Genome Biology and Evolution **7**:1155-1167.
1103   Williams, G. C. 1957. Pleiotropy, natural selection, and the evolution of senescence. Evolution **11**:398-
1104   411.
1105   Wittkopp, P. J., B. K. Haerum, and A. G. Clark. 2004. Evolutionary changes in cis and trans gene
1106   regulation. Nature **430**:85.
1107   Wittkopp, P. J., and G. Kalay. 2012. Cis-regulatory elements: molecular mechanisms and evolutionary
1108   processes underlying divergence. Nature Reviews Genetics **13**:59-69.
1109   Wray, G. A., M. W. Hahn, E. Abouheif, J. P. Balhoff, M. Pizer, M. V. Rockman, and L. A. Romano. 2003.
1110   The evolution of transcriptional regulation in eukaryotes. Molecular Biology and Evolution
1111   **20**:1377-1419.
1112   Wu, C. I., and A. W. Davis. 1993. Evolution of postmating reproductive isolation: the composite nature of
1113   Haldane rule and its genetic bases. American Naturalist **142**:187-212.
1114   Yanai, I., L. Peshkin, P. Jorgensen, and Marc W. Kirschner. 2011. Mapping gene expression in two
1115   *Xenopus* species: evolutionary constraints and developmental flexibility. Developmental Cell
1116   **20**:483-496.
1117   Zalts, H., and I. Yanai. 2017. Developmental constraints shape the evolution of the nematode mid-
1118   developmental transition. Nature Ecology & Evolution **1**:0113.
1119   Zhao, Z., T. J. Boyle, Z. Bao, J. I. Murray, B. Mericle, and R. H. Waterston. 2008. Comparative analysis of
1120   embryonic cell lineage between *Caenorhabditis briggsae* and *Caenorhabditis elegans*.
1121   Developmental Biology **314**:93-99.

1122

Box 1. Visualizing, conceptualizing, and modeling Dobzhansky-Muller incompatibilities.
Dobzhansky-Muller Incompatibilities (DMIs) can be viewed as disrupted gene networks and as valleys (or holes) in fitness landscapes (Gavrilets 2003, Fragata et al. 2019). Distinct biological species comprise separate groups of individuals that fail to successfully interbreed, though 'good' biological species may still yield hybrid F1 and F2 offspring, so long as they suffer clear fitness deficits (Coyne and Orr 2004). Of special interest for development are genetically intrinsic post-zygotic species barriers that manifest as DMIs in hybrid individuals and do not depend on extrinsic circumstances, reflecting disruptive changes to developmental programs. DMIs between populations can evolve from a single common ancestral population through the independent substitution of two or more mutations distinguishing the descendant lineages. The substitutions may fix due to positive selection or genetic drift. This DM model also encapsulates the essence of DSD: independent evolution in distinct lineages that causes divergence in genetic architectures while retaining within-lineage fitness (True and Haag 2001).

Species trees in (A) show the history of substitutions for loci in two genetic networks that differ in molecular evolution (case on left with e.g. fewer genes, stronger purifying selection, less adaptation, greater modularity, or lower pleiotropic effects) and the potential for DMIs (solid purple lines = potential DMI for derived-derived substitutions between species, dashed lines = potential derived-ancestral DMIs; lowercase = ancestral alleles, uppercase = derived substitutions unique to one lineage). The number of potential DMIs (purple) scales faster than

linear with number of substitutions (red and blue hashes); faster evolving genetic networks may
be more likely to experience this 'snowball effect' of reproductive isolation (Orr 1995). DSD
arises when the outward phenotype remains constant despite molecular divergence between
descendant species. Panel (B) shows how two loci (a and b) that diverge can potentially create a
DMI upon formation of F1 hybrids between descendant species. Panel (C) illustrates with a
Fisher's geometric model visualization, for two traits with a shared genetic architecture, how
adaptive evolution with respect to one trait (Trait 1) can generate DSD in another (Trait 2).
Concentric circles represent lines of equal fitness; filled dots (black = ancestor, red and blue =
descendant species) indicate genotypes (letters as in A) that evolve via three substitutions
(arrows) toward the fitness optimum at the center. Note the DSD in Trait 2 due to no net
phenotypic difference relative to the ancestor at the end of the adaptive walk for both species,
despite underlying genetic changes. Panel (D) shows evolution along ridges of equal fitness in a
fitness landscape comprised of a genetic architecture with many genes. Genotypic paths evolve
independently in different species (ancestral black to derived red and blue species), similarly to
DSD, such that hybrids between them (purple) occupy a portion of genotype space with low
fitness ('holes').

Box 2. Developmental disruption and regulatory inference in inter-species hybrids.
1160

Analysis of inter-species hybrids can reveal genome-wide mechanisms of gene regulatory
change. (A) Comparison of allele-specific expression ratios in F1 hybrids to expression ratios for
orthologous genes in parental individuals allows inference of expression changes due to local
cis-acting regulatory differences, distant *trans*-acting changes, or distinct ways that both *cis*- and
trans-regulatory divergence can jointly influence gene expression. (B) Another type of
expression comparison between F1 hybrid individuals and parental species can characterize the
dominance in allele-specific expression, with misexpression in hybrids reflecting
disproportionately high (overdominant) or low (underdominant) gene expression. (C) The
inference of misexpression in hybrids at the transcriptional level may be buffered at the
translational level, leading to more severe misexpression of the transcriptome than proteome.
Other approaches to using inter-species hybrids for deciphering the genetics of divergence in
development include quantitative trait locus (QTL) mapping. Inter-species QTL analyses can
incorporate screens with deletion libraries (Masly and Presgraves 2007), recombinant inbred line
(RIL) panels (Bedinger et al. 2011), near isogenic line (NIL) or introgression line panels
(Guerrero et al. 2017), or multigeneration selection approaches such as X-QTL mapping
(Ehrenreich et al. 2010).

1177
1178 Panels A and B redrawn from data for *Drosophila* flies in McManus et al. (2010). Panel C data
1179 for *Saccharomyces* yeast from Wang et al. (2015).
1180
1181

Box 3. Measuring molecular divergence across ontogeny.

Incorporating ontogeny into molecular evolution requires developmental time series of gene expression, which necessarily involves more experimental effort and sophistication of analysis than studies using a single developmental timepoint. This idea can be explored in several ways. First, one may quantify divergence in gene expression as a phenotypic output that closely maps to genotype. Studies in diverse organisms have compared transcriptome profiles over ontogeny with this approach (Table 1). Second, the contrast of gene expression levels between two parental species and their F1 hybrids provides a way to infer whether changes in expression result from *cis-* or *trans*-regulatory evolution (Box 1) (Wittkopp et al. 2004, Signor and Nuzhdin 2018). This approach has challenges and opportunities: environmental conditions also can influence transcriptome expression and the balance of *cis*:*trans* controls of the set of expressed genes (Tirosh et al. 2009), and it may be difficult to deconvolve maternal from zygotic *trans*-effects in F1 embryos. The technique also has not yet been applied in a developmentally dynamic way, making it ripe for future studies. Third, rates of coding sequence evolution (dN/dS or K_A/K_S) for genes expressed differentially across development provide a means of assessing selection on the encoded protein sequences used in genetic networks. A simple way of testing for trends in purifying selection and positive selection across ontogeny is to compare the average dN/dS value for the set of genes with peak expression at a given developmental timepoint (Cutter and Ward 2005, Davis et al. 2005) or an expression-weighted mean dN/dS value for all genes expressed at a given timepoint (Quint et al. 2012, Liu and Robinson-Rechavi 2018b). When a multi-species phylogeny is used, or if polymorphism data are incorporated in a McDonald-Kreitman testing framework, then positive selection may be distinguished from relaxed purifying selection (Liu and Robinson-Rechavi 2018a). Genes also can be grouped according to patterns of co-expression to analyze for consistent differences in rates of molecular evolution among them or among parameters of mathematical functions that describe the temporal profiles of expression (Coronado-Zamora et al. 2019, Cutter et al. 2019). Spatial definition of expression can enhance such approaches using tissue- and sex-specificity (Kim et al. 2016), restricted cell-lineages (Hashimshony et al. 2014), or tomographic expression profiling (Ebbing et al. 2018). While mRNA expression is most commonly accessed, all of these approaches can be extended to protein expression; protein levels appear to differ less between species than do transcriptomes (Leducq et al. 2012, Khan et al. 2013). Quantifying chromatin misregulation in a developmental time-course also could help discern the influence of epigenetic factors in hybrid dysfunction.

Panels (A) for *C. elegans* redrawn from (Cutter et al. 2019) and (B) for *Drosophila melanogaster* from (Coronado-Zamora et al. 2019) illustrate sets of genes with shared profiles of expression

1218   across development (non-linear timepoint scaling), and their rates of coding sequence evolution

1219   (*D. melanogaster* analysis excludes non-dynamic constitutive genes, gray for *C. elegans*; non-

1220   synonymous substitution rate K_A, adaptive non-synonymous substitution rate ω_a, non-adaptive

1221   non-synonymous substitution rate ω_{na}).

1222 **Tables**
1223
1224 **Table 1.** Studies in the literature that characterize expression profiles across development
1225 (Supplementary file 1).
1226

	Number of studies (%)		Number of species (%)	
Developmental stage analyzed				
embryogenesis only	58	39.5%	45	42.5%
non-embryo stages	37	25.2%	17	16.0%
all ontogeny	51	34.7%	44	41.5%
Type of molecular time series data				
transcriptome	128	87.1%	106	100.0%
proteome	7	4.8%	5	4.7%
other	14	9.5%	12	11.3%
Inter-species divergence feature				
gene expression	32	21.8%	55	51.9%
DNA sequence	5	3.4%	14	13.2%
non-comparative	99	67.3%	42	39.6%
Taxonomic group				
Angiosperm	16	10.9%		
Arthropoda	26	17.7%		
Chordata	55	37.4%		
Nematoda	10	6.8%		
other group only	30	20.4%		
multiple groups	10	6.8%		

1227

1228

Figures

Figure 1. Predictions and hypotheses in the evolution of ontogeny and reproductive isolation.
(A-C) Models from population genetics and evo-devo suppose that some modes of natural
selection may be more potent at particular life stages, as described in the main text. However, not
all models make clear predictions about the ontogenetic dynamism of selection for all selection
modes (purifying vs adaptive vs neutral) or for all stages of development. (D-E) Differential
incidence of selection across development may be mediated by genetic architecture in terms of
the pleiotropic effects of genetic changes and how that translates into the robustness of fitness-
related traits. When fitness is disproportionately robust to changes to genes expressed at a given
stage, then that stage will be more likely to accumulation cryptic genetic variation (CV) within
species, divergence between species as developmental system drift (DSD), and to result in
production of Dobzhansky-Muller incompatibilities (DMIs) in inter-species hybrid individuals.
1243
1244

1245

Figure 2. Developmental dynamics in *C. elegans* nematodes and in *Bufo* and *Xenopus*
amphibians. (A) *C. elegans* cell counts grow exponentially in early embryogenesis, before
slowing later. Redrawn with data from (Giurumescu et al. 2012) and wormatlas.org. (B) Gene
expression changes dynamically over ontogeny in terms of number of genes expressed and the
incidence of up- and down-regulated genes over time. Redrawn from (Boeck et al. 2016). (C)
Ontogenetic timing in the accumulation of reproductive isolation with genetic divergence for
Bufo toads. Hybrid individuals between more closely-related species develop to later stages than
do hybrids from distantly-related species pairs. Redrawn from (Malone and Fontenot 2008). (D-
E) *Xenopus* gene expression level differentiation decreases over developmental time (yolk
comprises ~½ of embryo volume; maternal-zygotic transition at stage ~8; yolk consumption
begins around gastrulation). Redrawn from (Yanai et al. 2011).

1257 **Supplementary file 1.**
1258 Summary of literature survey for studies analyzing ontogenetic profiles of transcriptome or
1259 proteome expression.