Dental diversity in early chondrichthyans and the multiple origins of shedding teeth

Richard P. Dearden[1] and Sam Giles[2,3,4]

[1]CR2P, Centre de Recherche en Paléontologie–Paris, Muséum national d'Histoire naturelle, Sorbonne Université, Centre National de la Recherche Scientifique, CP 38, 57 rue Cuvier, F75231 Paris cedex 05, France.

[2]School of Geography, Earth and Environmental Sciences, University of Birmingham, Birmingham, UK

[3]Department of Earth Sciences, Natural History Museum, Cromwell Road, London SW7 5BD, UK

[4]Department of Earth Sciences, University of Oxford, South Parks Road, Oxford, OX1 3AN, UK.

Abstract

The teeth of sharks famously form a series of parallel, continuously replacing files borne directly on the mandibular cartilages. In contrast, bony fishes possess site-specific shedding dentition borne on dermal plates. Understanding how these disparate systems evolved is challenging, not least because of poorly understood relationships and the profusion of morphologically and terminologically diverse bones, splints and whorls seen in the earliest chondrichthyans. Here we use tomographic methods to investigate the nature of mandibular structures in several early branching 'acanthodian'-grade stem-chondrichthyans. We characterise the gnathal plates of ischnacanthids as growing bones, and draw similarities between early chondrichthyan and stem gnathostome teeth and jaws. We further build the case for *Acanthodopsis*, a Carboniferous taxon, as an acanthodid, and show that, unexpectedly, its teeth are borne directly on the mandibular cartilage. Mandibular splints are formed from dermal bone and appear to be an acanthodid synapomorphy. The development of a unidirectionally growing dentition may be a feature of the chondrichthyan total-group. More generally, ischnacanthid and stem gnathostome gnathal plates share a common construction and are likely homologous, and shedding teeth evolved twice in gnathostomes.

Keywords (3-6): dentitions, early vertebrates, acanthodians, chondrichthyans, tooth evolution, Palaeozoic

1. Background

The structure and position of teeth and jaws are one of the major anatomical distinctions between osteichthyans (bony fishes: ray-finned fishes, lobe-finned fishes, and tetrapods) and chondrichthyans (cartilaginous fishes: sharks, rays, and chimaeras) (1). In osteichthyans, teeth are partially resorbed, shed, and replaced in position on dermal bones lateral to and overlying endoskeletal jaw cartilages. These form inner (dermal: coronoids, dermopalatines; endoskeletal: Meckel's cartilage, palatoquadrate) and outer (dermal: dentary, maxilla, premaxilla) dental arcades. In chondrichthyans, teeth grow, shed, and are replaced in parallel rows of labially-directed series directly on the jaw cartilages (endoskeletal: Meckel's cartilage, palatoquadrate). These two conditions are difficult to reconcile. Their origins can be observed in the morphologies of Palaeozoic gnathostome fossils, which suggest that the last common ancestor of jawed fishes (gnathostomes), as well as crownward stem gnathostomes (a paraphyletic assemblage referred to as 'placoderms'), possessed non-shedding teeth fused to the underlying dermal jaw bone (2–4).

The advent of micro-computed tomography has led to a renewed interest in tooth evolution and development in 'placoderms' (3–5) and osteichthyans (6–10). However, 'acanthodians', the earliest-branching members of the chondrichthyan total-group, have received comparatively little study (but see: (11–13)), despite a bewildering array of dermal oral structures (e.g. (11,14–18)).

Here we provide new tomographic data on teeth and jaws in several early-diverging stem chondrichthyans. We place this in the context of what we know

62 about early chondrichthyan relationships, and discuss its implications for the

63 early evolution of gnathostome dentitions.

2. Materials and Methods

All specimens studied here are housed at the Natural History Museum,

London (NHM), and comprise: an isolated *Taemasacanthus erroli* jaw

(NHMUK PV P33706); an isolated *Atopacanthus* sp. jaw (NHMUK PV

P.10978); an isolated *Acanthodopsis* sp. jaw (NHMUK PV P.10383); and a

partial head of *Acanthodes* sp. (NHMUK PV P.8065). Full descriptions, as well

as details of an additional isolated *Ischnacanthus* sp. jaw (NHMUK PV

P.40124) are given in the supplement.

CT scanning of two specimens took place at the Imaging and Analysis Centre,

NHMUK, using a Metris X-Tek HMX ST 225 with the following settings:

Taemasacanthus erroli: 3142 projections, 130 kV, 131 μA, 0.1 mm copper

filter, voxel size 17.3 μm; *Atopacanthus*: 3142 projections, 130 kV, 154 μA,

0.1 mm copper filter, voxel size 19.508/07/2020 18:05:00 μm.

CT scanning of three specimens took place at the Bristol University

Department of Life Sciences using a Nikon XT H 225 ST with the following

settings: *Acanthodopsis*: 3142 projections, 180 kV, 92 μA, no filter, voxel size

22.6 μm; *Acanthodes*: 3142 projections, 215 kV, 165 μA, 0.1 mm tin filter,

voxel size 44.9 μm; *Ischnacanthus*: 3142 projections, 222 kV, 105 μA, 0.5 mm

copper filter, voxel size 24.6 μm.

Reconstructed tomographic datasets were segmented in Mimics v.19

(biomedical.materialise.com/mimics; Materialise, Leuven, Belgium). Images of

the resulting models were then generated using Blender (blender.org).

88 **3. Results**

The left lower jaw of *Taemasacanthus* (previously described in Long (19): fig.

4A-C) comprises a tooth-bearing dermal gnathal plate (also referred to as a

dentigerous jaw bone: Burrow (15)) with a concave ventral surface overlying a

partially-ossified Meckel's cartilage (figure 1a,b, figure S1). The *Atopacanthus*

specimen examined here belongs to either a right lower or left upper jaw

(figure 1d,e, figure S2), and only the gnathal plate is preserved. The jaw

bones of both taxa have a broadly similar construction, which also

corresponds to that of *Ischnacanthus* (figure S3). The largest component of

the dentigerous jaw bone is the gnathal plate. Internally, this plate is highly

vascularised with interlinked antero-posteriorly polarised canals (figure 1c,f).

Three rows of teeth are borne on the biting edge of the gnathal plate: a lateral,

medial, and lingual row. Teeth within the lateral and lingual rows are fused to

the jaw but histologically separate from the underlying gnathal plate. The

medial row lies on the mesial ridge, formed by the occlusal corner of the

gnathal plate, and comprises small disorganised cusps (one row in

Taemasacanthus, two rows in *Atopacanthus*). All mesial teeth are

vascularised in *Atopacanthus*, but only the posteriormost ones are in

Taemasacanthus. The lateral and lingual rows of teeth in both taxa are much

larger, are ridged, and comprise a vascular base topped with an avascular

crown (figure 1a,c,f). Both grow by the addition of new teeth anteriorly, as

evidenced by anterior teeth partially overlying posterior ones, with cusps

becoming progressively larger in the direction of growth. The sole exception to

this is the penultimate lingual tooth in *Taemasacanthus* (figure 1a), which

partially overlies and is thus younger than the tooth anterior to it. The lateral

tooth row in both taxa lies on the dorso-lateral (or ventro-lateral) surface of the

gnathal plate. Its teeth are laterally unornamented and continuous with the

lateral surface of the dermal bone, connected to one another via antero-

posterior lateral ridges. The lingual tooth row grows on the lingual side of the

dermal plate, curving away from the occlusal surface anteriorly. These teeth

appear to sit on a bony lingual plate, with a distinct histology from the main

gnathal plate, something particularly obvious in *Atopacanthus* (figure S2).

The right lower jaw in *Acanthodopsis* comprises a tooth-bearing Meckel's

cartilage and a mandibular splint (figure 1g,i,j, figure S4). The Meckel's

cartilage is similar in form to that of *Acanthodes* (figure 1k–m, figure S5)

(15,20), with an identical articular cotylus and symphyseal fossa, and is

mineralised as a thick shell of perichondral bone which would have contained

cartilage in life. Unlike in *Acanthodes*, the Meckel's cartilage in *Acanthodopsis*

is mineralised along its entire length. A row of ten monocuspid, triangular

teeth runs along the dorsal surface of the Meckel's cartilage. The largest tooth

is in the middle of the jaw, with teeth becoming smaller and more closely set

anteriorly and posteriorly; they are slightly lingually convex, with a smooth (but

possibly weathered) lateral face and a longitudinally striated lingual face.

Rather than being tooth-shaped extensions of perichondral bone (15) these

are histologically distinct from Meckel's cartilage, formed from a thick outer

shell and a vascular inner pulp, with no obvious pulp cavity (figure 1i,j).

Although indistinct, growth lines suggest that the largest tooth is the oldest,

with younger teeth added both anteriorly and posteriorly. The mandibular

splint in *Acanthodopsis* is an unornamented, slightly sinusoidal bone that fits

into a groove on the ventro-lateral part of Meckel's cartilage, extending almost

its entire length. Internally, the element is pierced by a series of thin,

longitudinally oriented canals (figure 1i). This, combined with the fact that no

other endochondral mineralisation is preserved in either Meckelian element,

suggest that it is formed from dermal bone (21), rather than endochondral

tissue (15). In all respects the mandibular splint in *Acanthodes* (figure 1l,m,

figure S5) is almost identical to that of *Acanthodopsis* (20)*.

4. Discussion

Our new data show conclusively that the gnathal plates (also referred to as

dentigerous jaw bonens) of ischnacanthids were growing bones with new

teeth added in an anterior direction, as supposed by Ørvig (22) based on

directional wear. These teeth were fused to, but distinct from, the underlying

gnathal plate, which presumably grew with the rest of the jaw. This mode of

growth is comparable to that of stem-gnathostome arthrodire gnathal plates in

Compagopiscis and an unnamed buchanosteid, in which teeth are added in

multiple directions onto a growing basal dermal bone (3,5,23). This also

appears to be the case in the stem gnathostome acanthothoracid *Romundina*,

although its exact mode of growth is disputed (4,24–26). This organisation of

dental elements may be a plesiomorphic condition, shared with certain stem-

group gnathostomes (figure 2). Unlike these taxa however, the tooth growth in

ischnacanthids is unidirectional. This is more comparable to what is seen in

'acanthodian' tooth whorls (12,13), where non-shedding cusps are added

unidirectionally onto a bony base. This unidirectional mode of tooth growth

may be a chondrichthyan synapomorphy (figure 2).

The row of monocuspid dermal teeth borne directly on the Meckelian element

of *Acanthodopsis* is unlike any other known chondrichthyan (with the possible

exception of *Pucapampella*: (27)), and distinct from that seen in any other

known gnathostome. Although in the past *Acanthodopsis* has been

considered to have dermal dentigerous jaw bones (15,22), our CT data show

conclusively that the main body of the jaw is endoskeletal in origin. A

symphyseal fossa and *Acanthodes*-like mandibular splint further support

Burrow's (15) assertion that *Acanthodopsis* is closely related to

acanthodiforms, rather than ischnacanthiforms. However, beyond this its

dental morphology is difficult to interpret. The teeth are comparable to the

tooth whorls of more crownward chondrichthyans in that they are borne

directly on the Meckelian element and are apparently oriented perpendicular

to the direction of the jaw bone. In this sense they could be interpreted as a

non-growing tooth whorl with a single generation. However, this interpretation

is confounded by the fact that *Acanthodes* is completely toothless, and

phylogenetic analyses recover it as nested within a larger clade of toothless

acanthodiforms inferred to be filter-feeders, the earliest members of which

existed in the Early Devonian (28). If this phylogeny is correct, and the teeth

of *Acanthodopsis* are homologous with tooth whorls, it would demand at least

two convergent losses of teeth in this clade (i.e. in deeper-diverging

acanthodiforms and *Acanthodes*). Alternatively, teeth may simply be

unobserved in some acanthodiform taxa due to their small size: teeth were

recently found in the supposedly edentulous filter feeder *Gladbachus*

adentatus (28).

Mandibular splints (variously termed dentohyoids, extramandibular spines,

splenials, and mandibular bones (29)) have been reported in a range of

'acanthodian'-grade taxa including acanthodids (29,30), mesacanthids

(31,32), cheiracanthids (21), ischnacanthids (33,34), and diplacanthids

(35,36). Their small size and unclear association with other bones of the

mandibular arch has made them difficult to characterise, of unclear homology,

and prone to being mixed up with other mandibular elements; the supposed

mandibular splint in diplacanthids has for example been shown to be an

"occlusal bone" (18). Our characterisation of the mandibular splint in

Acanthodopsis and *Acanthodes* as a dermal bone with a distinctive shape

allows reassessment of these mandibular bones in other taxa. The

morphology in other acanthodids appears likely to match that in the two taxa

we describe in that they have a slightly sinusoidal shape, for example in other

species of *Acanthodes* (20,29,37), *Halimacanthodes* (30), and *Howittacanthus*

(38). In *Ischnacanthus* (figure S3), the ventral margin of Meckel's cartilage is

reinforced and laterally flattened; we suggest this is also likely to be the case

in other ischnacanthids with "mandibular splints" (33). This may also be the

case in mesacanthids, in which the mandibular splint is not convincingly

separate from the Meckel's cartilage or branchiostegal plates, for example in

Promesacanthus (31). The condition is uncertain in cheiracanthids: a

mandibular splint is absent in *Cheiracanthus* and *Homalacanthus*, and while

Protogonacanthus is described as having a mandibular splint (21) its

morphology looks more similar to the reinforced ventral margin in

ischnacanthids. Thus it is possible that a separate "true" mandibular splint, as

seen in *Acanthodopsis* and *Acanthodes*, is an acanthodid synapomorphy. Its

similarity to the ventral branchiostegal rays in *Acanthodes* suggest that it may

simply be part of this series that has been co-opted to support the jaw.

Our new data on 'acanthodian' jaw elements feed into an emerging picture of

stem chondrichthyan evolution. A unidirectional mode of tooth growth appears

to be a chondrichthyan synapomorphy (with a possible reversal in

Acanthodopsis; figure 2). While fine-scale relationships remain poorly

understood, phylogenetic analyses increasingly recover a stemward grade of

ischnacanthiforms, acanthodiforms, and diplacanthiforms (28), with a

climatiid-grade more proximate to the chondrichthyan crown. Consequently, a

dentition consisting entirely of tooth whorls—which extend along the entire jaw

length—as well as absence of dermal gnathal plate, is restricted to climatiids

and more crownward taxa (figure 2), although possibly homologous dentitions

may be seen in ischnacanthid tooth whorls, *Acanthodopsis*, and

Latviacanthus. Tooth shedding is restricted to the node proximate to the

chondrichthyan crown. Some uncertainty remains, however, largely due to the

uncertain position of edentulous taxa such as *Lupopsyrus* and

Kathemacanthus, and the peculiar site-specific dentition apparently present in

Pucapampella (27).

Our new data also have a bearing on the broader question of jaw and tooth

evolution in gnathostomes. Ischnacanthid gnathal plates (also referred to as

dentigerous jaw bones) appear homologous with the gnathal plates of stem-

gnathostomes: they are positionally, structurally and histologically similar,

being vascularised dermal bones overlying the mandibular cartilage, with non-

shedding tooth cusps added onto an underlying dermal plate. These gnathal

plates have been homologised with the inner dental arcade of osteichthyans

(39–41), and more recently with the outer dental arcade (42,43). In either

case, assuming homology between gnathal plates and either the inner or

outer dental arcade, this is a rare example of a macromeric skeletal structure

preserved on all three branches incident to the gnathostome crown node.

Our reassessment of early chondrichthyan dentitions also presents an

opportunity to reconsider the homology of tooth whorls. The rows of denticles

on the marginal jaw bones of the stem-osteichthyans *Lophosteus* and

Andreolepis have been said to recall the tooth "families" of chondrichthyans

(6). However, given that multiple rows of tooth whorls are absent in the

deepest-diverging chondrichthyans, and that the parasymphyseal tooth whorls

that are present grow from the lingual side of the jaw cartilage as in living

chondrichthyans (6), we consider any resemblance superficial (9).Similarly,

ischnacanthids cheek scales have been argued to be incipient tooth whorls

formed from cheek denticles (11,44). However, given the phylogenetic

position of ischnacanthids, and the propensity of dermal ornament bordering

the mouth to resemble dentition (e.g. *Mimipiscis*, *Ptomacanthus* (45,46)), we

consider these structures removed from hypotheses of early tooth evolution.

Parasymphyseal tooth whorls in which the teeth replace via resorption and

anterior rotation (10) are also known in osteichthyans (e.g. in *Onychodus*

(10,47)), but these are phylogenetically and structurally removed from

chondrichthyan tooth whorls and unlikely to be homologous. However, it is

difficult to assess whether non-shedding parasymphyseal whorls are

homologous due to the unclear condition in psarolepids (variably interpreted

as stem sarcopterygians or stem osteichthyans (48–52)), in which a whorl is

inferred (48,52) but is yet to be described. Finally, the interposition of many

non-shedding stem-chondrichthyan taxa between shedding chondrichthyans

and shedding osteichthyans confirms that a shedding dentition evolved twice,

in two different ways, in crown-gnathostomes (7,10). The teeth of extant

chondrichthyans, borne directly on endoskeletal mandibular cartilages, are

positionally distinct from both the inner and outer dental arcades of

osteichthyans, where teeth are borne on dermal bones.

Acknowledgements

We thank E. Bernard and Z. Johanson (NHMUK) for assistance with

specimen access, V. Fernandez and B. Clark (both NHMUK) and T. Davies

(University of Bristol) for assistance with CT scanning, and M. Brazeau

(Imperial College London), B. Davidson and C. Burrow (Queensland Museum)

for discussion. M. Colfer (University of Oxford) assisted with segmentation of

Mimics files.

Funding

This work was supported by a Junior Research Fellowship, Christ Church,

Oxford, and a Royal Society Dorothy Hodgkin Research Fellowship, both to

S.G.. R.P.D. is supported by a Paris Île-de-France Region – DIM "Matériaux

anciens et patrimoniaux" grant (PHARE projet).

Ethics. This research is based exclusively on specimens from natural history

collections.

Data accessibility. Raw data (.vol or .tiff stacks), Mimics files, and 3D PLY

files for each specimen are deposited in Dropbox (see links in supplementary

file) and will be permanently deposited in Dryad upon manuscript acceptance.

Authors' contributions. S.G. conceived the project and selected specimens.

S.G. and R.P.D. carried out CT scanning. R.P.D. segmented the specimens,

made Blender renders and constructed figures with input from S.G. S.G. and

R.P.D. drafted the manuscript. Both authors revised and edited the

manuscript, approved the final version and agree to be accountable for all

aspects of the work.

Competing interests. We have no competing interests.

5. References

1.   Tucker AS, Fraser GJ. Evolution and developmental diversity of tooth

regeneration. Semin Cell Dev Biol. 2014;25–26:71–80.

2.   Ørvig T. Histologic Studies of Ostracoderms, Placoderms and Fossil

Elasmobranchs: 3. Structure and growth of the gnathalia of certain

arthrodires'. Zool Scr. 1980;9(1–4):141–159.

3.   Rücklin M, Donoghue PCJ, Johanson Z, Trinajstic K, Marone F,

Stampanoni M. Development of teeth and jaws in the earliest jawed

vertebrates. Nature. 2012;491.

4.   Rücklin M, Donoghue PCJ. *Romundina* and the evolutionary origin of

teeth. Biol Lett. 2015;11:0–4.

5.   Hu Y, Young GC, Burrow C, Zhu Y, Lu J. High resolution XCT scanning

reveals complex morphology of gnathal elements in an Early Devonian

arthrodire. Palaeoworld. 2018;

6.   Botella H, Blom H, Dorka M, Ahlberg PE, Janvier P. Jaws and teeth of

the earliest bony fishes. Nature. 2007;448:583–6.

7.   Chen D, Blom H, Sanchez S, Tafforeau P, Märss T, Ahlberg PE.

Development of cyclic shedding teeth from semi-shedding teeth : the

inner dental arcade of the stem osteichthyan Lophosteus. 2017;

8.   Chen D, Blom H, Sanchez S, Tafforeau P, Ahlberg PE. The stem

osteichthyan *Andreolepis* and the origin of tooth replacement. Nature .

2016; Available from: http://dx.doi.org/10.1038/nature19812

9.   Cunningham J a., Rücklin M, Blom H, Botella H, Donoghue PCJ. Testing

models of dental development in the earliest bony vertebrates,

Andreolepis and *Lophosteus*. Biol Lett. 2012;8(5):833–7.

10.   Doeland M, Couzens AMC, Donoghue PCJ, Rücklin M. Tooth

replacement in early sarcopterygians. R Soc Open Sci. 2019;6.

11.   Maisey JG, Turner S, Naylor GJP, Miller RF. Dental patterning in the

earliest sharks: Implications for tooth evolution. J Morphol. 2014

May;275(5):586–96.

12.   Qu Q, Sanchez S, Blom H, Tafforeau P, Ahlberg PE. Scales and Tooth

Whorls of Ancient Fishes Challenge Distinction between External and

Oral "Teeth." PLoS ONE. 2013;8(8).

13.   Rücklin M, Giles S, Janvier P, Donoghue PCJ. Teeth before jaws?

Comparative analysis of the structure and development of the external

and internal scales in the extinct jawless vertebrate *Loganellia scotica*.

Evol Dev. 2011;13:523–32.

14.   Brazeau MD. A revision of the anatomy of the Early Devonian jawed

vertebrate *Ptomacanthus anglicus* Miles. Palaeontology. 2012;55:355–

367.

15.   Burrow CJ. Acanthodian fishes with dentigerous jaw bones : the

Ischnacanthiformes and Acanthodopsis. Foss Strata. 2004;(50):8–22.

16. Burrow CJ, Blaauwen JD, Newman MJ, Davidson RG. The diplacanthid

fishes (Acanthodii , Diplacanthiformes , Diplacanthidae) from the Middle

Devonian of Scotland. Palaeontol Electron. 2016;(February):1–83.

17. Hanke GF, Davis SP, Wilson MVH. New species of the acanthodian

genus *Tetanopsyrus* from northern Canada, and comments on related

taxa. J Vertebr Paleontol. 2001;21(4):740–753.

18. Newman MJ, Davidson RG, Blaauwen JL Den, Burrow CJ. The Early

Devonian Acanthodian *Uraniacanthus curtus* (Powrie , 1870) n . comb .

from the Midland Valley of Scotland. Geodiversitas. 2012;34(4):739–59.

19. Long JA. New ischnacanthid acanthodians from the Early Devonian of

Australia, with comments on acanthodian interrelationships. Zool J Linn

Soc. 1986;87:321–339.

20. Miles RS. Relationships of acanthodians. In: Greenwood P, Miles RS,

Patterson C, editors. Interrelationships of fishes. London: Zoological

Journal of the Linnean Society; 1973. p. 63–103.

21. Miles RS. The acanthodian fishes of the Devonian Plattenkalk of the

Paffrath Trough in the Rhineland. Ark För Zool. 1964;18(9):147–94.

22. Ørvig T. Acanthodian dentition and its bearing on the relationships of the

group. Palaeontogr Abt A. 1973;143:119–50.

23. Hu Y, Lu J, Young GC. New findings in a 400 million-year- old Devonian

placoderm shed light on jaw structure and function in basal

gnathostomes. Sci Rep. 2017;7:1–12.

24. Burrow CJ, Hu Y, Young GC. Placoderms and the evolutionary origin of

teeth: a comment on Rücklin and Donoghue (2015). Biol Lett. 2016;12.

25. Rücklin M, Donoghue PCJ. Reply to "placoderms and the evolutionary

origin of teeth": Burrow et al. (2016). Biol Lett. 2016;12.

26. Smith MM, Clark B, Goujet D, Johanson Z. Evolutionary origins of teeth

in jawed vertebrates: conflicting data from acanthothoracid dental plates

('Placodermi'). Palaentology. 2017;60(6):829–36.

27. Maisey JG, Janvier P, Pradel A, Denton JSS, Bronson A, Miller R, et al.

Doliodus and Pucapampellids Contrasting Perspectives on Stem

Chondrichthyan Morphology. In: Johanson Z, Underwood CJ, Richter M,

editors. Evolution and Development of Fishes. Cambridge: Cambridge

University Press; 2019. p. 87–109.

28. Coates MI, Finarelli JA, Sansom IJ, Andreev PS, Criswell KE, Tietjen K,

et al. An early chondrichthyan and the evolutionary assembly of a shark

body plan. Proc R Soc B Biol Sci. 2018;285(1870).

29. Zidek J. Kansas Hamilton Quarry (Upper Pennsylvanian) *Acanthodes*,

with remarks on the previously reported North American occurrences of

the genus. Univ Kans Palaeontol Contrib. 1976;83.

30. Burrow CJ, Trinajstic K, Long JA. First acanthodian from the Upper

Devonian (Frasnian) Gogo Formation , Western Australia. Hist Biol.

2012;24(4):349–57.

31. Hanke GF. *Promesacanthus eppleri* n. gen., n. sp., a mesacanthid

(Acanthodii, Acanthodiformes) from the Lower Devonian of northern

Canada. Geodiversitas. 2008;30(2):287–302.

32. Watson DMS. The Acanthodian Fishes. Philos Trans R Soc Lond B Biol

Sci. 1937 Sep 9;228(549):49–146.

33. Ørvig T. Some new acanthodian material from the lower Devonian of

Europe. Zool J Linn Soc. 1967;47(1966):131–53.

34. Jessen H. Weitere fischreste aus dem Oberen Plattenkalk der Bergisch-

Gladbach Paffrather Mulde (Oberdevon, Rheinisches Schiefergebirge).

Palaeontogr Abt A. 1973;143:159–87.

35. Young GC, Burrow CJ. Diplacanthid acanthodians from the Aztec

Siltstone (late Middle Devonian) of southern Victoria Land, Antarctica.

Foss Strata. 2004;(50).

36. Hanke GF, Davis SP. Redescription of the acanthodian *Gladiobranchus*

probaton Bernacsek &Dineley, 1977, and comments on diplacanthid

relationships. Geodiversitas. 2008;30(2):303–30.

37. Beznosov P. A redescription of the Early Carboniferous acanthodian

Acanthodes lopatini Rohon, 1889. Acta Zool. 2009;90(May):183–93.

38. Long JA. A new Late Devonian acanthodian fish from Mt. Howitt, Victoria,

Australia, with remarks on acanthodian biogeography. Proc R Soc Vic.

1986;98:1–17.

39. Stensiö E. Elasmobranchiomorphi; Placodermata, Arthrodires. In: Piveteau J, editor. Traité de paleontologie. Paris: Masson et Cie.; 1969. p. 71–692.

40. Young GC. The relationships of placoderm fishes. Zool J Linn Soc. 1986 Sep 15;88(1):1–57.

41. Zhu M, Yu X, Ahlberg PE, Choo B, Lu J, Qiao T, et al. A Silurian placoderm with osteichthyan-like marginal jaw bones. Nature. 2013 Oct 10;502(7470):188–93.

42. Zhu M, Ahlberg PE, Pan Z, Zhu Y, Qiao T, Zhao W. A new Silurian maxillate placoderm illuminates jaw evolution. Science. 2016;

43. Zhu Y, Ahlberg PE, Zhu M. The evolution of vertebrate dermal jaw bones in the light of maxillate placoderms. In: Johanson Z, Underwood CJ, Richter M, editors. Evolution and Development of Fishes. Cambridge: Cambridge University Press; 2019. p. 71–86.

44. Blais SA, MacKenzie LA, Wilson MVH. Tooth-like scales in Early Devonian eugnathostomes and the outside-in hypothesis for the origins of teeth in vertebrates. J Vertebr Paleontol. 2011 Nov;31(6):1189–99.

45. Gardiner BG. The relationships of the palaeoniscid fishes, a review based on new specimens of *Mimia* and *Moythomasia* from the Upper Devonian of Western Australia. Bull Br Mus Nat Hist Geol. 1984;37(4):175–418.

46.   Miles RS. Articulated acanthodian fishes from the Old Red Sandstone of

England, with a review of the structure and evolution of the acanthodian

shoulder girdle. Bull Br Mus Nat Hist Geol. 1973;24(2).

47.   Andrews SM, Long JA, Ahlberg PE, Barwick RE, Campbell K. The

structure of the sarcopterygian *Onychodus jandemarrai* n. sp. from Gogo,

Western Australia: with a functional interpretation of the skeleton. Trans

R Soc Edinb Earth Sci. 2006;96:197–307.

48.   Zhu M, Yu X, Janvier P. A primitive fossil fish sheds light on the origin of

bony fishes. Nature. 1999;6004(1978):607–610.

49.   Qu Q, Haitina T, Zhu M, Ahlberg PE. New genomic and fossil data

illuminate the origin of enamel. Nature. 2015;526:108–111.

50. Choo B, Zhu M, Qu Q, Yu X, Jia L, Zhao W. A new osteichthyan from the

late Silurian of Yunnan, China. Plos One. 2017;12(3):e0170929.

51. Lu J, Giles S, Friedman M, Zhu M. A new stem sarcopterygian illuminates

patterns of character evolution in early bony fishes. Nat Commun 2017;8.

http://dx.doi.org/10.1038/s41467-017-01801-z

52. Zhu M, Zhao W, Jia L, Lu J, Qiao T, Qu Q. The oldest articulated

osteichthyan reveals mosaic gnathostome characters. Nature.

2009;458(7237):469–74.

53. Dearden RP, Stockey C, Brazeau MD. The pharynx of the stem-

chondrichthyan *Ptomacanthus* and the early evolution of the

gnathostome gill skeleton. Nat Commun. 2019;10(1):1–7.

54. Kamminga P, De Bruin PW, Geleijns J, Brazeau MD. X-ray computed

tomography library of shark anatomy and lower jaw surface models. Sci Data.

2017;in press:1–6.

Figures

Figure 1. Tomographic models of jaw elements in acanthodian-grade stem-

chondrichthyans: (a-c) the left lower jaw of *Taemasacanthus erroli* NHMUK

PV P.33706 in (a) medial view, (b) lateral view, and (c) a reconstructed

tomogram showing a sagittal section through the lingual tooth row; (d-f) a

gnathal plate of *Atopacanthus* sp. NHMUK PV P.10978 in (d) medial view, (e)

lateral view, and (f) a reconstructed tomogram showing a sagittal section

through the lingual tooth row; (g-j) the right lower jaw of *Acanthodopsis* sp.

NHMUK PV P.10383 in (g) lateral view, (h) medial view, (i,j) reconstructed

tomograms showing (i) a sagittal section through the entire jaw and (j) a

transverse section through the jaw; (k-m) the lower jaws of *Acanthodes* sp.

NHMUK PV P.8085 in (k) ventral view against the digital cast of the surface

with (l,m) reconstructed tomograms showing (l) a coronal section through the

specimen, and (m) a transverse section through a lower jaw. Abbreviations:

Mand. splint., Mandibular splint; Meck. cart, Meckel's cartilage; L., Left; R.,

Right.

Multidirectional
tooth addition

buchanosteid

Compagopiscis

Entelognathus

Teeth
lost

Andreolepis

Tooth
shedding

crown-ost.

Teeth on
dermal
plate
on Meck.
cart.

Ischnacanthus

Taemasacanthus

Atopacanthus

Teeth
lost

Diplacanthus

Teeth
lost

Acanthodes

Monodirectional
tooth addition

Mandibular
splint

Acanthodopsis

Tooth add.
bidirectional

Teeth on
Meck. cart.

Ptomacanthus

Doliodus

Teeth on
Meck. cart.

crown-chond

Tooth whorls
comprise
entire dentition

Tooth
shedding

Figure 2. Hand-drawn cladogram showing relationships of total-group

gnathostomes pertinent to tooth evolution and schematics of their lower jaw

morphologies. Anterior to right and all specimens in lingual view except for

Compagopiscis, *Entelognathus*, and *Andreolepis*. Black arrow indicates

position of initial tooth growth, blue graded arrow indicates direction of

subsequent growth. Colour scheme: grey, Meckel's cartilage; dark blue,

gnathal plates and attached teeth; green, mandibular splints; light blue, teeth

placed directly on endoskeleton; white, dermal bones of uncertain homology

and attached teeth. Abbreviations: Add., addition; Meck. cart, Meckel's

cartilage. Images redrawn from (3,5,8,11,16,32,41,45,53,54).

Supplementary material (separate pdf)

Supplementary information for this manuscript is included as a single

separate file. It includes supplementary figures 1-5, supplementary text

comprising a full description of the elements scanned and a review of

acanthodian jaw elements, and supplementary references. It also contains

links to 3D models (as plys), mimics files, and tomographic data (as volume

files or TIFFs) for all the data used in this study.