Diverse stem-chondrichthyan oral structures and evidence for an independently acquired acanthodid dentition

Richard P. Dearden[1] and Sam Giles[2,3*]

[1]CR2P, Centre de Recherche en Paléontologie–Paris, Muséum national d'Histoire naturelle, Sorbonne Université, Centre National de la Recherche Scientifique, CP 38, 57 rue Cuvier, F75231 Paris cedex 05, France.

[2]School of Geography, Earth and Environmental Sciences, University of Birmingham, Birmingham, UK

[3]Department of Earth Sciences, Natural History Museum, Cromwell Road, London SW7 5BD, UK

*Corresponding author

<u>Abstract</u>

The teeth of sharks famously form a series of transversely-organised files with conveyor-belt replacement that are borne directly on the jaw cartilages, in contrast to the dermal plate-borne dentition of bony fishes that undergoes site-specific replacement. A major obstacle in understanding how this system evolved is the poorly understood relationships of the earliest chondrichthyans and the profusion of morphologically and terminologically diverse bones, cartilages, splints and whorls that they possess. Here we use tomographic methods to investigate mandibular structures in several early branching 'acanthodian'-grade stem-chondrichthyans. We show that the dentigerous jaw bones of disparate genera of ischnacanthids are united by a common construction, being growing bones with non-shedding dentition. Mandibular splints, which support the ventro-lateral edge of the Meckel's cartilage in some taxa, are formed from dermal bone and may be an acanthodid synapomorphy. We demonstrate that the teeth of *Acanthodopsis* are borne directly on the mandibular cartilage and that this taxon is deeply nested within an edentulous radiation, representing an unexpected independent origin of teeth. Many or even all of the range of unusual oral structures may be apomorphic, but they should nonetheless be considered when building hypotheses of tooth and jaw evolution, both in chondrichthyans and more broadly.

<u>Keywords (3-6):</u> dentitions, early vertebrates, acanthodians, chondrichthyans, tooth evolution, Palaeozoic

1. Introduction

The structure and position of teeth and jaws are amongst the major anatomical distinctions between crown osteichthyans (bony fishes: ray-finned fishes, lobe-finned fishes, and tetrapods) and crown chondrichthyans (cartilaginous fishes: sharks, rays, and chimaeras) [1]. In osteichthyans, teeth are partially resorbed at their base, shed, and replaced in position on dermal bones lateral to and overlying endoskeletal jaw cartilages as part of outer and inner dental arcades. In crownward chondrichthyans, teeth grow, shed, and are replaced in parallel rows of labiolingually-directed series directly on the jaw cartilages. The origins of these dental structures can be traced back to Palaeozoic taxa, which suggest that the last common ancestor of jawed fishes (gnathostomes), as well as crownward stem gnathostomes (a paraphyletic assemblage referred to as 'placoderms'), possessed non-shedding teeth fused to the underlying dermal jaw bone [2–7]. However, oral structures in many Palaeozoic gnathostomes remain poorly characterised, and as a result their relevance to the evolution of teeth is unclear.

The advent of micro-computed tomography has led to a renewed interest in tooth evolution and development in Palaeozoic gnathostomes. These have mostly focussed on stem-gnathostome 'placoderms' [2,5,6,8] and osteichthyans [9–14] and have revealed an unexpected range of morphologies. Stem-group gnathostomes have non-shedding dentitions, which may be arranged radially [5,8] or in parallel rows [6], borne on an underlying dermal bone. The homology of the dermal jaw bones in stem-group gnathostomes to the inner and outer dental arcades of crown

gnathostomes is uncertain [6,15,16]. Meanwhile, many Palaeozoic

osteichthyans possessed shedding dentitions comparable to more recent

taxa, although some stem osteichthyans have dental structures such as

symphyseal tooth whorls [14] and marginal cusps organised into rows

[9,10,11,12,13], which are more broadly comparable in the gnathostome total

group. Early-branching members of the chondrichthyan total-group (including

'acanthodians'), have received less attention, with only a handful of taxa

described using CT data [7,17-20]. This is despite a remarkable array of

dermal oral structures across the assemblage: various early chondrichthyans

possess tooth whorls [17,21,22], gracile or molariform teeth not organised into

whorls [18,23,24], dermal plates of differing constrictions with and without

teeth [22,23,25-28], or may lack dermal mandibular structures entirely [29]. A

variety of extramandibular 'dentitions' and other oral structures are also

known [29,30].

Teeth arranged into files are widespread in chondrichthyans both living and

extinct. In the larger of the two constituent chondrichthyan clades

(elasmobranchs: sharks and rays), teeth are continuously replaced in

generative series [1]. They grow on the inner margin of the jaw, move through

a labiolingual file (figure 1a), and are shed at the labial jaw margin [1]. In

holocephalans (chimaeras and relatives), the dentition is modified to two

upper pairs and one lower pair of non-shedding, hypermineralised toothplates

[31]. The elasmobranch-like condition of labiolingual files of teeth is seen in

both stem-group elasmobranchs [32] and stem-group holocephalans [33],

implying that this condition is plesiomorphic for chondrichthyans. Many

Palaeozoic chondrichthyans possessed tooth whorls, where the tooth file

comprises multiple cusps fused onto a common base. Tooth whorls are

common in taxa in the chondrichthyan stem-group, and may form the entire

dentition (e.g. *Ptomacanthus* [21]; *Doliodus* [17]) or be present at and/or

restricted to the symphysis (e.g. ischnacanthids: [20,22]). It is unclear whether

individual teeth were shed from tooth whorls borne on mandibular rami, or

whether the whorls themselves were shed [17]. Some probable crown-group

chondrichthyans had a further condition in which teeth did not share a

common base but post-functional teeth were retained at the labial margin of

the jaw [34]. This, alongside the prevalence of tooth whorls with fused teeth in

stem-chondrichthyans, indicates that tooth shedding was acquired later in the

chondrichthyan total-group.

In addition, a ream of stem-group chondrichthyans have unusual dentitions

that do not conform to a file-like arrangement, and which are less commonly

considered in hypotheses of tooth evolution (figure 1b-f). The most well-

characterised of these are dentigerous jaw bones: large, tooth-bearing dermal

plates which sit on both the upper and lower jaw cartilages in 'ischnacanthid'

stem-group chondrichthyans [22]. Dentigerous jaw bones bear one or more

rows of antero-posteriorly aligned teeth, the cusps of which can be quite

morphologically and presumably functionally variable [25]. The tooth rows

have been shown to grow via anterior addition of new cusps in one

taxonomically unidentified specimen, but construction of the bone beyond the

tooth rows, as well as the array of different morphologies, remain poorly

characterised [7,26,27]. Other stem-group chondrichthyans, such as

Gladbachus [24], *Pucapampella* [18] and *Acanthodopsis* [23] have teeth that

are neither part of dentigerous jaw bones nor arranged into files. Diplacanthid

stem-group chondrichthyans [27,28] bear smooth, toothless dermal plates on

their lower jaws. Numerous stem-group chondrichthyans lack teeth altogether,

and the lower jaws of many of these taxa bear poorly-characterised

mandibular splints [30,37] which have also been identified in some

diplacanthids and ischnacanthids [38,39].

Further confounding this diversity of oral structures are competing and

unsettled hypotheses of relationships for early chondrichthyans. While

'acanthodians' are now established as stem-group chondrichthyans [24,32,33,

40-44], there is limited certainty over the monophyly of 'acanthodian'

subgroups. There is some evidence that an assemblage of diplacanthiform,

ischnacanthiform, and acanthodiform taxa may form a clade or grade

subtending the remainder of the chondrichthyan total-group [24]. 'Climatiid'

acanthodians, which have overlapping character complements with more

shark-like chondrichthyans [18], tend to be recovered in a more crownward

position (although see [35]). Beyond this, however, different phylogenetic

analyses present very different schemes of relationships, often with low

support. These conflicting patterns of relationships present major obstacles to

understanding patterns evolution for many of the dental structures seen in

early chondrichthyans, although likelihood-based methods provide a possible

approach [7].

Here we use computed tomography to image the teeth, jaws and associated

oral structures of several early-diverging stem-group chondrichthyans with the

aim of more broadly sampling the diversity of their oral structures. We aim to

characterise the anatomy of several different ischnacanthid dentigerous jaw

bones of different constructions, the mandibular splint of acanthodids, and the

‘teeth’ of *Acanthodopsis*. We contextualise our new data within a wider review

of stem-group chondrichthyan oral structures and identify those that represent

synapomorphies, while also discussing challenges in reconstructing dental

evolutionary histories and homologies.

2. Materials and Methods

2.1 Taxa examined

All specimens studied here are housed at the Natural History Museum,

London (NHMUK), and comprise: an isolated jaw of *Taemasacanthus erroli*

(NHMUK PV P.33706); an isolated jaw of *Atopacanthus* sp. (NHMUK PV

P.10978); an isolated jaw of *Acanthodopsis* sp. (NHMUK PV P.10383); a

partial head of *Acanthodes* sp. (NHMUK PV P.8065); and an isolated jaw of

Ischnacanthus sp. (NHMUK PV P.40124).

Taemasacanthus erroli is known from eight isolated jaws from the Emsian

(Lower Devonian) Murrumbidgee Group in New South Wales, Australia [45].

Other species, also based on isolated jaw bones, have been assigned to the

genus [46,47], but no articulated animals are known. *Taemasacanthus* is

understood to be an ischnacanthid on the basis of its dentigerous jaw bone

[45]. NHMUK PV P.33706, described here (figure 2), is a right lower jaw, and

comprises two main parts: a dermal dentigerous jaw bone and the articular

ossification of Meckel’s cartilage. The external morphology of this specimen of

Taemasacanthus has been fully described [45] but is briefly redescribed here

to contextualise our new information. An additional ventral fragment of the

articular (previously figured [45]) has become detached from the rest of the

ossification and was not included in the CT scan.

Atopacanthus is known throughout the Middle-Upper Devonian [50]. The type

species, *Atopacanthus dentatus*, is known from several dentigerous jaw

bones from near Hamburg, New York and is presumed to be an ischnacanthid

[23,49]. The sole articulated specimen attributed to *Atopacanthus* sp., from

the Upper Devonian of the Rhineland [50,51], has since been referred to

Serradentus [52] and so the genus is known only from disarticulated remains.

The specimen described here, NHMUK PV P.10978 (figure 3), is a

dentigerous jaw bone collected from Elgin, Scotland, and is Late Devonian in

age. It was originally labelled as a possible dipnoan toothplate before later

being referred to *Atopacanthus*, and its morphology conforms with that of

other specimens described as *Atopacanthus*. It is not associated with any

endoskeletal material, and it is impossible to tell whether it is from a right

lower jaw or left upper jaw. For ease of comparison with other specimens, we

describe its morphology as if it were a part of the lower jaw.

Ischnacanthus is the best known ischnacanthid 'acanthodian', represented by

numerous articulated specimens of Lochkovian (Lower Devonian) age from

the Midland Valley in Scotland [22]. The material described here (NHMUK PV

P.40124; figure 4) is an isolated left lower jaw from the Lower Devonian

(Lochkovian) Midland Valley in Tealing, Forfarshire [53]. It is fairly complete,

but parts of the dorsal and anterior margins have been lost to the counterpart

(which is preserved, but was not CT scanned), and the whole jaw is laterally

flattened. It comprises a dentigerous jaw bone and Meckel's cartilage.

Acanthodopsis is known from the Carboniferous of the UK and Australia.

Acanthodopsis has been previously considered an ischnacanthid on the basis

of its "dentigerous jaw bones", but in terms of its skeletal anatomy it is more

similar to acanthodids [23,45]. The material described here, NHMUK PV

P.10383 from the Northumberland Coal Measures (figure 5), comprises a

laterally flattened lower right jaw, consisting of a Meckel's cartilage with teeth

and a mandibular splint.

Acanthodes is the latest occurring genus of 'acanthodian' found as articulated

body fossils from the Mississippian (Carboniferous) into the Lower Permian

[31]. It is the only genus of 'acanthodian'-grade animal known from

extensively preserved endoskeleton, seen in specimens of *Acanthodes*

confusus from Lebach, Germany [54]. The material described here (NHMUK

PV P.8065) comprises part of the ventral half of the head of a specimen from

the Knowles Ironstone of Staffordshire (figure 6). As the dorsal margins of the

jaw bones are obscured within the rock, it was originally referred to

'*Acanthodopsis* or *Acanthodes*'. As CT scanning shows that dentition is

absent, we can confirm it to be *Acanthodes* sp. Most of the left jaw is

preserved, and of the right jaw only the mandibular splint is preserved, as are

some of the lower branchiostegal ray series and isolated dermal gill rakers.

Scattered parts of the rest of the head endoskeleton are also present,

including parts of ceratobranchial and a hyomandibular.

2.2 CT scanning

Full details of scanning parameters are given in supplementary table 1. The

voxel size for each scan are as follows: *Acanthodes*, 44.9 *µ*m; *Acanthodopsis*,

22.6 µm; *Atopacanthus*, 19.51 *µ*m; *Ischnacanthus*, 24.6 *µ*m; *Taemasacanthus*

erroli, 17.3 *µ*m.

Reconstructed tomographic datasets were segmented in Mimics v.19

(biomedical.materialise.com/mimics; Materialise, Leuven, Belgium) and

images were generated using Blender (blender.org).

2.3 Phylogenetic analyses

Our dataset is based on Dearden *et al*. [44], the most recently published

phylogenetic dataset specifically tackling stem-group chondrichthyan

relationships. We have added one taxon and four characters and made minor

modifications to some codes (full details given in the supplementary text). We

performed a parsimony analysis in TNT [55] with the following settings: a

parsimony ratchet with 10,000 iterations, holding 100 trees per iteration, with

TBR branch swapping. Galeaspida was set as the outgroup and the following

constraint applied: (Galeaspida(Osteostraci(Mandibulate Gnathostomes))).

We also performed a Bayesian analysis, with Galeaspida set as the outgroup,

mandibulate gnathostomes constrained as monophyletic, and the following

settings. We used a uniform prior with the Mkv model and gamma-distributed

rates; searched for 10,000,000 generations, sampling every 1,000

generations; and calculated the majority rule consensus tree with a relative

burn-in of 25%.

3. Results

3.1 *Taemasacanthus*

The dentigerous jaw bone of *Taemasacanthus* is approximately half the full

depth of the jaw and is sinusoidal in dorsal view (figure 2). A circular ridge,

previously suggested to be for a labial cartilage attachment [45] but more

likely the attachment site for a ligament, is present on the lateral surface

approximately ¼ of the way along its length (figure 2b,c). Posteriorly, the bone

curves laterally and broadens to wrap around the articular. The lateral

expansion is larger than the medial expansion, and both are rounded

posteriorly. Ventrally, a groove formed by the posterior confluence of these

two processes runs underneath the entire length of the dentigerous jaw bone,

and would have overlain Meckel's cartilage (figure 2c). The dentigerous jaw

bone is approximately trapezoid in cross section, and bears three rows of

teeth (figure 2a,b,d,g,h). Histologically the underlying dermal plate comprises

heavily vascularised dermal bone, similar to that observed in thin sections of

other ischnacanthid dentigerous jaw bones [7,22,56], with a relatively thin

layer of less vascularised bone around the plate margins (figure 2g; figure

S1a-d). The vascularisation comprises an interlinked network of tubules,

which are strongly anteroposteriorly polarised in the tooth-bearing section of

the bone. Vascular channels occasionally open onto the surface of the bone,

particularly the ventral groove. On the lingual surface of the bone, a change in

orientation of the vascularisation (from anteroposterior to more random)

indicates the presence of a separate plate-like unit of growth (Figure S1d).

The more posterior, wrapped part of the bone has a radial arrangement of

vasculature suggesting that the bone grew posteriorly and ventrally as well as

anteriorly as the underlying endoskeletal jaw grew. The circular ridge is

formed of avascular bone, but is otherwise a similar tissue to that forming the

outer margin of the dermal plate (figure 2f).

Three rows of teeth are borne on the biting edge of the dermal plate, all

starting approximately at the level of the mesial ridge: a lateral, medial, and

lingual row (figure 2a,b,d). Teeth of the medial row are far smaller and less

distinct than those of the lateral and lingual rows. Teeth within the lateral and

lingual rows are fused to the jaw, but the base of the tooth is marked by an

increase in the density of random and dorsally-oriented vascular canals

(figure 2f; figure S1a-c). The medial row lies on the mesial ridge and

comprises a single row of small disorganised cusps that are continuous with

the underlying dermal plate and do not appear to overlap with each other.

Vasculature in the medial row is only visible in the more posterior cusps and

does not seem related to size (figure S1f). The lateral and lingual rows of

teeth are much larger and ridged, with a vascular base topped with a mostly

avascular crown. The younger, larger, more anterior teeth of both rows have

extensively vascularised crowns. The smaller, more posterior cusps have

less—or even no—vascularisation, indicating that the crown was infilled in

older teeth. The basal vasculature has occasional connections with the

vasculature of the underlying dermal plate. The teeth seemingly lack a

continuous enameloid covering (figure 2f), contra recent reports of enameloid

in an ischnacanthid [7], although we caution that this may be due to the

resolution of our dataset. Both tooth rows grow by the addition of new teeth

onto the anterior end of the row, as evidenced by anterior teeth partially

overlying posterior ones, and cusps becoming progressively larger in the

direction of growth. The sole exception to this is in the lingual tooth row,

where, in what is probably a pathology, the eighth cusp of the lingual row is

incomplete and is either damaged or its growth has been aborted (figure 2a).

Both the damaged eighth cusp and the undamaged tenth cusp are overgrown

by the youngest cusp in the row. This, the ninth cusp, is oriented noticeably

more medially than other cusps and may have disrupted the growth of the

smaller tenth cusp, although the underlying bone appears unaffected. The

vasculature of the ninth cusp appears to be isolated from the surrounding

vasculature, while the vasculature of the underlying eighth cusp opens into a

large, central hollow. The lateral tooth row comprises around twelve cusps,

and its teeth are laterally unornamented and continuous with the lateral

surface of the dermal bone, connected to one another via antero-posterior

lateral ridges (figure 2b). The lingual side of each cusp is rounded, and

ornamented with a number of ridges, which become longer and progressively

more tuberculated on more anterior cusps. The lingual tooth row comprises

ten cusps, which curve away from the occlusal surface anteriorly.

Only the posteriormost portion of the Meckel's cartilage, the articular

ossification, is preserved (figure 2). It is formed from a sheath of perichondral

bone and would have been filled with cartilage in life. Some spongy texture is

apparent on the interior surface of the perichondral bone. A shallow groove on

the posterior surface does not appear to continue ventrally, making it unlikely

to have accommodated a mandibular splint as previously suggested [45].

Articulation with the palatoquadrate appear to be via an open, oval, fossa [45]

(figure 2d). The tissue forming this is notably ill-formed, and appears to lack a

solid perichondral covering; as Burrow [22] suggests it seems likely to be a

articular process like that in other ischnacanthids [22,25] which is broken.

3.2 *Atopacanthus*

The dentigerous jaw bone is robust, trapezoid in cross-section, but flattened laterally and taller proportionate to its length compared to *Taemasacanthus* (figure 3). It is slightly medially convex. The anterior fifth of the preserved bone (it is broken both anteriorly and posteriorly) is toothless and tapers slightly. A narrow, shallow groove to accommodate the mandibular cartilage runs along its ventral surface (figure 3c). The histology is similar to *Taemasacanthus*, with heavily vascularised dermal bone surrounded by a less vascular layer, but the vascular tubules are even more strongly polarised in an antero-dorsal direction (figure 3e; figure S1e-g). Again, the vasculature principally opens into the ventral groove. Towards the surface, bone vascularisation is less dense, and not polarised. The lingual face of the dentigerous jaw bone supports a distinct thin, lingual, tooth-bearing plate (figure 3a,d,e), which is comparable in position and vascularisation to the plate-like region in *Taemasacanthus* (figure S1d). This plate is still heavily vascularised, but tubes are polarised dorso-lingually. On the outer perimeter of the lateral face of the main bone, the vasculature is oriented obliquely.

As in *Taemasacanthus*, lateral, mesial and lingual tooth rows are borne on the dorsal surface of the underlying dermal plate (figure 3a,b,d). The medial ridge bears two disorganised rows of cusps along its anterior half, with the posterior half being smooth. All cusps are vascular, and are histologically continuous with the medial ridge. The lateral tooth row comprises eight cusps, which become progressively larger anteriorly, and their lateral surfaces are continuous with the outside of the dermal plate (figure 3b). Their lingual surfaces are rounded and ornamented with untuberculated ridges. The lingual

tooth row comprises ten main cusps, which curve medially across the

dentigerous jaw bone. Two additional small cusps are present near the

posterior margin of the dermal plate, ventral to the main lingual tooth row,

which are closest in appearance to the teeth of the medial ridge (figure 3a,d).

The lingual tooth row lies on top of a lingual plate, which is apposited onto the

lingual surface of the main dermal plate (figure 3a,d,e) . As in the lateral row

and in *Taemasacanthus*, cusps become larger anteriorly, and are ornamented

with ridges. The histology of the teeth of the lateral and lingual rows

comprises a vascular base topped with an avascular cap lacking enameloid,

with the vascular canals oriented distinctly from the underlying dermal and

lingual plates (figure 3e). As in *Taemasacanthus*, younger teeth are more

heavily vascularised with an extensively vascularised crown, and the

anteriormost tooth still posseses a clear pulp cavity (figure S1e-g), suggesting

that teeth vasculature became infilled with age. Teeth in both the lateral and

lingual rows were added anteriorly, with anterior cusps partly overlying their

posterior fellows.

3.3 *Ischnacanthus*

Only the anterior part of the dentigerous jaw bone is preserved in the part

(figure 4), although the mould of the posterior region is visible in outline. The

underlying dermal plate is much shallower than in *Taemasacanthus* and

Atopacanthus. A lateral tooth row and a medial ridge are present. The lateral

tooth row preserves four cusps, the third of which is exposed on the surface

and therefore incomplete dorsally. The cusps are linked by a cuspidate ridge

along their lateral faces. Relative size and age are difficult to determine due to

the mode of preservation, but the anteriormost cusp is the largest, and cusp

overlap indicates that teeth were added anteriorly. Although the ventralmost

parts of the dermal plate are missing, the ventral margin of the teeth is

marked by a noticeable shift in density and orientation of the vascular canals

(figure 4d). The tissue forming the teeth is similar to *Taemasacanthus* and

Atopacanthus, with a vascularised base and an avascular crown apparently

lacking enameloid [22]. The internal vasculature of the bone in *Ischnacanthus*

is also longitudinally polarised and connected, although less well-visualised in

our scan data (figure S1h,i).

The large Meckel's cartilage is near-complete and preserved as a single

ossification (figure 4). It is curved posteriorly and tapers anteriorly. The

dentigerous jaw bone is borne on its dorso-lingual surface. A laterally-directed

articular condyle is present at the posterior extent, and a shallow groove

extends ventral to the condyle. The majority of the Meckel's cartilage is

formed of globular calcified cartilage. Parts of its lateral surface, as well as its

ventral, anterior, and posterior extents, are covered by a thin, densely

mineralised tissue that appears to be perichondral bone [22]. This tissue

thickens ventrally and posteriorly and is fractured. The perichondral sheath is

avascular, but has fractured in such a way that cracks and voids artificially

resemble the vasculature of the dentigerous jaw bone. A thickened ridge

along the posteroventral and posterior margin is continuous with the

perichondral rind that extends onto the lateral surface, but externally gives the

appearance of a separate ossification (figure 4bc,). In section this is closely

comparable to the so called "mandibular bone" that Ørvig [39] described in

Xylacanthus and is probably responsible for accounts of mandibular splints in

ischnacanthids.

3.4 *Acanthodopsis*

The lower jaw in *Acanthodopsis* comprises a tooth-bearing Meckel's cartilage

and a mandibular splint (figure 5). The Meckel's cartilage is long and thin and

similar in form to that of *Acanthodes* (figure 5; 23,54], with an identical

articular cotylus and marked preglenoid process. It tapers anteriorly,

terminating in a small, cup-shaped anterior symphyseal fossa (figure 5c,e).

The Meckel's cartilage is formed from a shell of what we infer to be

perichondral bone [23], which has collapsed and cracked under compression

and appears to be unmineralized internally (figure 5g,h), although some

mineralisation appears to be present in the jaw articulation. Unlike

Acanthodes confusus, it is perichondrally mineralised along its entire length

(figure 5), rather than in separate articular and mentomandibular sections.

Ten monocuspid, triangular teeth form a row along the dorsal surface of the

Meckel's cartilage (figure 5). The largest tooth is in the middle of the jaw, with

teeth becoming smaller and more closely set anteriorly and posteriorly; they

are slightly lingually convex, each with a smooth (but possibly weathered)

lateral face and a longitudinally striated lingual face. Previous descriptions of

the teeth [23] were undecided as to their tissue makeup. The tissue

comprising them comprises distinct inner and outer layers. The outer layer is

thick and covers the outside surface of each tooth but does not close

ventrally: unlike the perichondral surface of the cartilage it is not crushed and

the surface appears intact. The inner layer has a spongy texture, which may

reflect internal vasculature: no obvious pulp canals are present. Although the

contrast between the teeth and jaw bone is subtle, they can be differentiated

in that the internal tissues of the teeth have a spongy texture, whereas the

perichondral bone is solid. We infer the teeth to be dermal due to this

histological distinction from the Meckel's cartilage as well as their gross

structure, which shows a separation from the Meckel's cartilage, and their

ornamentation. We consider it likely that these tissues are dentinous but more

detailed study is needed to establish their identity. The direction of growth is

difficult to infer. In successive tomograms viewed in sequence, the largest

tooth appears to be overlapped by the anterior and posterior teeth, possibly

making it the oldest. This contrasts to the order of growth in *Taemasacanthus*,

Atopacanthus and *Ischnacanthus*, where teeth are added anteriorly. However,

as with the identity of the tissue, we express caution at this interpretation.

The mandibular splint in *Acanthodopsis* is an unornamented, slightly

sinusoidal bone that fits into a groove on the ventro-lateral part of Meckel's

cartilage, extending almost its entire length (figure 5a,b). This groove was

likely originally much shallower, and its depth has been exaggerated by lateral

flattening of the specimen. The tissue forming the splint is solid, and is

organised into multiple concentric lamellae likely representing lines of arrested

growth. It is pierced by a series of thin, longitudinally oriented canals (figure

5g,h). This tissue is distinct from that forming Meckel's cartilage, in particular

being denser and better organised, and can be interpreted as dermal bone,

especially given the ornamentation of the mandibular splints of some

acanthodids [57]. Burrow [23] reported that the mandibular splint in

Acanthodopsis was formed from cartilage and bone based on thin sections,

but the cartilage identified in these reports may have been the Meckel's

cartilage above the bone instead (CJB pers. comm. July 2020).

3.2 *Acanthodes*

The left lower jaw comprises ossified articular and mentomandibular parts of

Meckel's cartilage, as well as a mandibular splint (figure 6). Mineralised parts

of the Meckel's cartilage are formed from thick perichondral bone, and are

slightly laterally crushed (figure. 6a-d). The articular is as previously described

[54]. The mentomandibular has a distinct cup-like symphyseal fossa at its

anterior tip, forming part of the mandibular symphysis (figure 6c). The

mandibular splint is unornamented, slightly sinusoidal in shape, and ellipsoid

in cross-section. It sits in a groove in the lateral faces of the mentomandibular

and articular. Internally it is solid and vascularised by sparse long, thin canals

running its length (figure 6e,f), as in *Acanthodopsis*. A single tooth-like cusp

sitting in (although separate from) a cushion-shaped base is probably a

branchial or hyoid raker (figure 6b,e).

3.6 *Phylogenetic results*

Our parsimony analysis recovered 26101 most parsimonious trees with a

length of 704 steps. The strict consensus of these results (figures 7a, S2) is

consistent with other recent analyses in finding all 'acanthodians' to be stem-

group chondrichthyans. We recover "Acanthodii" *sensu* Coates *et al.* [24] as a

clade subtending the remainder of the chondrichthyan total group, with

Euthacanthus as the sister group to the "Acanthodii". However,

acanthodiforms (i.e. cheiracanthids, mesacanthids, and acanthodids) are

paraphyletic. Ischnacanthids plus diplacanthids form a clade, but

ischnacanthids themselves are paraphyletic, and diplacanthids are a clade to

the exclusion of *Tetanopsyrus*. Remaining stem chondrichthyan taxa,

including climatiids, *Gladbachus* and *Doliodus*, are recovered in a polytomy

along with a monophyletic chondrichthyan crown group. Support values aside

from the chondrichthyan total group and crown group nodes are typically low.

The Bayesian majority rule consensus tree (figure 7b, S3) is broadly

consistent with the parsimony strict consensus tree, although "Acanthodii" is

instead recovered as a polytomy subtending all more crownwards

chondrichthyans. *Acanthodopsis* is recovered as the sister-taxon to

Acanthodes in both analyses.

4. Discussion

4.1 Stem-chondrichthyan oral structures

Our new data show that the dentigerous jaw bones of all ischnacanthids,

including articulated taxa such as *Ischnacanthus* and those only known from

isolated jaws (e.g. atopacanthids and taemasacanthids), were united by a

common construction. These follow the model of tooth growth hypothesized

by Ørvig [26] and demonstrated by Rucklin *et al.* [7] on the basis of directional

wear and overlapping cusps. These teeth were fused to, but distinct from, the

underlying bone, which grew with the endoskeletal component of the jaw.

Based on the vasculature of *Taemasacanthus* the dermal bone grew radially

from a point posterior to the tooth row. This condition and the positions of

tooth rows relative to the underlying dermal plate is common to the three

different morphologies of dentigerous jaw bone that we describe and we infer

it to have been a common feature of ischnacanthid dentigerous jaw bones

The presence of an out of sequence tooth in *Taemasacanthus*, where the

youngest tooth has partially overgrown a cusp anteriorly (figure 2), suggests

that non-sequential growth was possible in the otherwise ordered tooth rows,

likely in response to a pathology. Dentigerous jaw bones are broadly

comparable with the condition in stem-gnathostomes in the sense that non-

shedding teeth are ankylosed to and growing on a basal bone. However,

phylogenetic topologies supporting homology between these conditions are in

limited supply (figures 7,8), and is not upheld in our topology or by more

detailed analysis [7].

Although dentigerous jaw bones are typically contrasted with tooth whorls [7],

we suggest that dentigerous jaw bones can be usefully interpreted by

comparison to antero-posteriorly "stretched out" tooth whorls. Teeth and

tooth-like structures are added directionally across the gnathostome total-

group and this may be a plesiomorphic feature of gnathostome dentitions [6].

However, tooth rows on dentigerous jaw bones are more comparable to

whorls than to these other structures in that tooth files are located in a specific

position on an underlying dermal plate, growing in a single direction. This

stands in contrast to single-directional, but haphazardly arranged, tooth files

reported in stem-gnathostomes and early osteichthyan marginal jaw bones

[6,9]. Notably, symphyseal tooth whorls and whorl-like cheek scales in some

ischnacanthids could suggest common patterning mechanisms affecting

dermal structures in and around the mouth of early chondrichthyans [58]. This

organisation tooth files could be apomorphic for chondrichthyans if

osteichthyan tooth whorls are optimised as homoplasious [7], and could

potentially be a character uniting chondrichthyan dentitions more inclusively

than the presence of tooth whorls.

The row of monocuspid 'teeth' borne directly on the Meckelian element of

Acanthodopsis is unlike that of any other known chondrichthyan or

gnathostome. Furthermore, the presence of teeth in a Carboniferous taxon

deeply nested within an edentulous radiation (figure 7), the oldest of which are

Early Devonian in age, strongly suggests that this represents an independent

acquisition of dentition. Previous studies of the jaw of *Acanthodopsis* have

interpreted it either as a dentigerous jaw bone [26] or as a perichondrally

ossified Meckelian bone with 'teeth' [23]. Our data confirm the latter view and

show that the teeth in *Acanthodopsis* are histologically distinct from the

underlying perichondral bone and so are presumably dermal ossifications

attached to its surface. Our phylogenetic analysis supports the view of Long

[45] and Burrow [23] that *Acanthodopsis* is closely related to *Acanthodes*, with

the presence of teeth representing the only difference between the genera. A

possible morphological comparison to these teeth lies in the branchial and

hyoid rakers found in acanthodiform fishes like *Acanthodes*, *Cheiracanthus*,

and *Homalacanthus* [31,59]. As with the teeth of *Acanthodopsis*, these rakers

are conical, sometimes striated ([61] fig. 3), and decrease in size from the

centre outwards (Dearden *pers obvs*). An alternative to homology between

the mandibular dentition in *Acanthodopsis* and other gnathostomes, which is

difficult to reconcile with their phylogenetic relationships, may be the co-option

of hyoid and branchial rakers to form a novel "dentition" on the mandibular

arch.

4.2 Distribution of oral structures in the chondrichthyan stem-group

Teeth and other oral structures in stem-chondrichthyans take on a broad

variety of forms, raising questions about their homology and the primitive or

derived condition for different features. Here we briefly review this diversity

and map key characters onto our phylogeny.

A diverse array of **teeth** (light blue icons, figures 7 and 8) are present in the

majority of Palaeozoic chondrichthyans, including as tooth whorls, but are

also remarkable for the breadth of taxa in which they are absent. Acanthodids

(e.g. *Acanthodes*: figure 6, ref. [54]) are completely toothless, except for

Acanthodopsis, as are the likely related mesacanthids (e.g. *Promesacanthus*

[37]) and cheiracanthids (e.g. *Cheiracanthus* [59]). In this latter group, tooth-

like hyoid rakers have sometimes been mistaken for teeth [61]. Teeth are also

absent in diplacanthids [28], with the possible exception of *Tetanopsyrus* (see

"Occlusal plates" section below) [29]. A number of toothless taxa with

otherwise diverse anatomies, including *Obtusacanthus*, *Lupopsyrus*,

Euthacanthus, *Brachyacanthus*, and *Kathemacanthus* [62-65], are often

resolved as more closely related to the crown-group (figures 7,8). Given the

small size and two-dimensional preservation of some of these taxa, it is

possible that teeth are present but reduced and so far undetected, as in

Gladbachus adentatus [24]. Teeth are inferred to be homologous across

gnathostomes [7], demanding numerous independent losses of teeth in stem

chondrichthyans (figures 7 and 8).

Tooth whorls (dark green icons, figures 7 and 8) are tooth files with a fused

bony base, which grow by the lingual addition of new cusps along a single

axis [20]. Tooth whorls are understood to be the evolutionary precursor to

modern chondrichthyan tooth families, which have a similar morphology but

lack a common bony base [66]. Tooth whorls are present in a range of stem-

group chondrichthyans with otherwise dissimilar anatomies, including those

with dentigerous jaw bones (e.g. *Ischnacanthus* [22]), densely tesserate head

skeletons (e.g. *Climatius* [67]), and more conventionally shark-like taxa (e.g.

Doliodus [17]). They have also been described in the acanthodiform-like

Latviacanthus [68], although as this is based on x-ray plates these may be

mischaracterised hyoid rakers as in *Homalacanthus* [59]. Tooth whorls are

also present in some stem-group holocephalans such as iniopterygians [69],

although these are likely to be secondarily derived given their phylogenetic

remoteness from the chondrichthyan stem-group. Some osteichthyan taxa

also possess tooth whorls, but here the tooth crowns are shed via resorption

of the tooth base, a mechanism not present in statodont chondrichthyan tooth

whorls [14].There is variation in their distribution on the jaw: in ischnacanthids

and osteichthyans, tooth whorls are few in number and limited to the

symphysis [22], whereas in more crownward chondrichthyans they are

arrayed along the length of the jaw and comprise the entire dentition (gold and

yellow-green icons, figures 7 and 8). The distribution of tooth whorls across

osteichthyans and within chondrichthyans is complex, with whorl-bearing taxa

often nested within whorl-less radiations. Probabilistic ancestral state

reconstruction indicates that tooth whorls evolved independently multiple

times both within chondrichthyans and across gnathostomes [7]. Within

chondrichthyans, different phylogenetic topologies have quite different

implications for the gain and loss of tooth whorls and their distribution across

the group (figure 8).

Some chondrichthyans have **teeth that are not organised into files** but

which lie directly on the jaw cartilage (coral icons [in part; this icon also

captures teeth arranged in files that are borne on the jaw cartilages], figures 7

and 8). This condition is present in *Acanthodopsis* (figure 5), *Pucapampella*,

and *Gladbachus* [18,24], although expressed in different ways. In

Acanthodopsis, teeth are triangular in profile and diminish in size anteriorly

and posteriorly (figure 5). In *Pucapampella*, teeth form a single row along the

jaw but show a variety of sizes, shapes, and spacings [18]. In *Gladbachus*,

teeth are much reduced and individually separate, although possibly aligned

linguo-labially [24]. However, these taxa are scattered across the tree, and

their tooth morphologies can be radically different, suggesting that teeth that

lie directly on the jaw cartilage and are not organised into files evolved

multiple times independently. Although *Gladbachus* and *Pucapampella* are

recovered as a clade in our analyses, this is contrary to most other recent

findings (figure 8c,d) and we view this result with extreme caution.

Dentition cones are tooth-like cones with smaller denticles attached. They

are only known in three partially-articulated ischnacanthids (*Zemylacanthus*

(*Poracanthodes*), *Acritolepis*, and *Serradentus* [52,70,71]), and are absent in

the better characterised *Ischnacanthus* and relatives [22,58]. The lack of fully

articulated fossils bearing dentition cones leaves open the possibility that they

represent a displaced part of the branchial apparatus rather than oral

structures [50], and are perhaps comparable to gill and hyoid rakers in

cheiracanthids and acanthodids [31,59]. Whether oral or branchial in origin,

their presence may unite the subset of ischnacanthids that possess them,

although this has not been tested in a phylogenetic context.

Tooth-like scales are present along the oral margin of some stem-group

chondrichthyans, and include part of the cheek squamation, the 'lip' and the

ventral rostral area. They are best characterised in *Ischnacanthus*-like

'acanthodians' [32] in which they show a variety of morphologies, and in life

may have helped with grasping prey. Some of these scales are strikingly

similar in organisation to tooth whorls (which are also present within the gape

of the same animals), comprising a file of denticles oriented towards the

mouth [22,32]. Specialised tooth-like scales have also been identified along

the margin of the mouth in *Obtusacanthus* [63]. More generally, tooth-like

denticles are common along the oral margin of the tooth row in early

osteichthyans [14,72]. Although potentially interesting from a developmental

perspective, they seem unlikely to carry any phylogenetic signal.

Dentigerous jaw bones are tooth-bearing dermal jaw bones present in the

upper and lower jaws of a number of stem-chondrichthyan taxa [23].

Articulated fossils bearing dentigerous jaw bones include *Zemlyacanthus,*

Nerepisacanthus, and *Serradentus* [44,52,73]. By far the best anatomically

characterised taxa with dentigerous jaw bones are *Ischnacanthus* and similar

taxa [22,25]. There are few anatomical characters to group taxa possessing

dentigerous jaw bones, but all have a complement of oral structures including

some combination of symphyseal tooth whorls, dentition cones, and tooth-like

cheek scales. Dentigerous jaw bones themselves display anatomical diversity,

for example relating to the structure of the bone, the number and morphology

of tooth rows [23], and variance in dentition shapes likely linked to diet [25]. In

our phylogeny, taxa with dentigerous jaw bones (i.e. ischnacanthids) are

recovered in a polytomy, in a broader grouping of 'acanthodians' with dermal

mouth plates (dark purple icons, figures 7 and 8). Dermal jaw bones, both

edentulous and tooth-bearing, are also present in 'placoderms' and

osteichthyans, but few phylogenetic results support their homology with those

of ischnacanthids (Fig. 7,8).

Occlusal plates are a pair of smooth dermal plates in the gapes of some

stem-chondrichthyans (light purple icons, figures 7 and 8). Their detailed

anatomy is poorly characterised and in the past they have become

terminologically and anatomically confused with the mandibular splint [30].

Occlusal plates are present in *Diplacanthus*, *Rhadinacanthus*, *Milesacanthus*,

Uraniacanthus, *Culmacanthus*, and *Tetanopsyrus* [28-30,74–76]. At least

some of these taxa have other common morphologies (i.e. similar body

shapes, scapular processes with posterior lamina, large postorbital scales,

deep, striated dorsal fin spine insertions), and on this basis they are grouped

into the diplacanthids [28]. There is some variation in the morphology of

occlusal plates. In all taxa but *Tetanopsyrus* [29,77], they are only present in

the lower jaws. *Tetanopsyrus* may also have tooth-like denticles along the

inner surface of the plates, although this is only known from an isolated

Meckel's cartilage associated with a complete *Tetanopsyrus* specimen [29]

and its attribution is therefore uncertain. In *Uraniacanthus* and *Culmacanthus*,

a dorsal process is present [74,78]. We recover diplacanthids as monophyletic

in our Bayesian analysis (figure 7, S3), but paraphyletic with respect to

Tetanopsyrus in our parsimony analysis, and occlusal plates appear to be a

character uniting diplacanthids (figures 7, 8). *Tetanopsyrus*, with its upper and

lower plates, may represent a link between the occlusal plates of

diplacanthids and dermal jaw bones of ischnacanthids.

A **mandibular splint** (variously termed dentohyoid, extramandibular spine,

splenial, or mandibular bone) is a slightly sinusoidal dermal bone that

underlies the Meckel's cartilage ventrolaterally (grey icons, figures 7 and 8).

Unlike the other structures discussed here, it did not lie within the gape, and

likely reinforced the lower jaw. Mandibular splints are present in *Acanthodes*,

Acanthodopsis, *Halimacanthodes*, *Howittacanthus*, and *Protogonacanthus*

[55,57,79,80]. They have also been incorrectly identified in a variety of other

taxa. Mandibular splints in mesacanthids [35,62] are more similar in size to

gular plates, and may represent displaced elements of this series. Reports in

diplacanthids [74,76] are better interpreted as occlusal plates [30]. Although a

mandibular splint has been identified in the putative cheiracanthid

Protogonacanthus [57], the taxon in question is likely not a cheiracanthid but

an acanthodid [59]. Finally, as we show, descriptions of mandibular splints in

ischnacanthids [39,50] instead represent a reinforced ventral margin of the

endoskeletal mandible. Mandibular splints in acanthodids are very

conservative in form, although may be ornamented as in *Acanthodes sulcatus*

[57]. Its similarity to the ventral branchiostegal rays in *Acanthodes* (figure

6a,b), which are also dermal, tubular, and slighty sinusoidal, suggests that it

may be part of this series that has been co-opted to support the jaw. Our

phylogeny suggests that a mandibular splint unites *Acanthodes* and

Acanthodopsis but either evolved convergently in *Halimacanthodes* (figure. 7)

or was lost in *Homalacanthus*; we consider it most likely that it unites

acanthodids to the exclusion of other stem-chondrichthyans, and that this

distribution is a result of undersampling acanthodids and their characters in

our phylogeny.

4.3 The evolution of chondrichthyan teeth

Although phylogenetic topologies for early chondrichthyans are poorly

resolved and often suggest conflicting hypotheses, there are some signals

that may provide insight into the evolution of a modern shark-like dentition.

The placement of 'climatiid' acanthodians in a relatively crownward position

on the chondrichthyan stem [24,42-44] (Fig. 7,8a-c) suggests that taxa with a

dentition entirely formed from tooth whorls share a last common ancestor to

the exclusion of other stem-group chondrichthyan taxa. Not all topologies

support this hypothesis, however: Frey *et al.* [35] recover climatiids as remote

from the chondrichthyan crown node, implying that an extensively whorl-

based dentition borne on the jaw cartilages either developed independently in

the crownward lineage or was lost in the Acanthodii *sensu* Coates *et al.* This

phylogeny is based on a more limited selection of stem-group chondrichthyan

taxa, which may have had an influence on reconstructed patterns of character

evolution. Either scenario still invokes multiple episodes of secondary tooth

loss (e.g. *Lupoposyrus*) and divergences from a whorl-like tooth anatomy

(*Gladbachus*, *Pucapampella*). In our phylogeny, tooth-shedding is restricted to

the crown-node. Generative tooth series are present in stem-group

elasmobranchs (e.g. *Phoebodus* [34]) and stem-group holocephalans with

both shark-like (e.g. *Ferromirum* [35]) and more chimaeroid-like (e.g.

Debeerius [81]) forms. However, it is unclear how widespread the non-

shedding condition described in some sharks with cladodont teeth [36] is.

Either way, this suggests a 'two-step' development of the stereotypical

chondrichthyan dentition, with an initial shift towards tooth whorls borne

exclusively on the jaw cartilages, followed by the eventual loss of fused bases

and concomitant development of tooth shedding in crown-group

chondrichthyans. However, significant phylogenetic uncertainty persists, and

this scenario warrants further testing as hypotheses of relationship stabilise.

Despite this, the interposition of multiple lineages of non-shedding stem-

chondrichthyan taxa between shedding chondrichthyans and shedding

osteichthyans confirms that a shedding dentition evolved twice, in two

different ways, in crown-gnathostomes [6,7,11,14].

In stark contrast to the clade comprising chondrichthyans with tooth whorls,

the clade or grade including diplacanthids, acanthodids, and

ischnacanthiforms exhibits a diverse array of oral structures, none of which

seem to persist beyond the end of the Palaeozoic [24]. In phylogenetic

analyses, this grade is consistently recovered at the base of the

chondrichthyan total-group, with the exception of Frey *et al.* [35], who recover

it in a more crownward position. A number of likely apomorphic oral

morphologies are present within this clade, including diplacanthid occlusal

plates, ischnacanthid dentigerous jaw bones and toothless acanthodids with

mandibular splints. Many of these morphologies are known from the Late

Silurian and Devonian, approximately contemporaneously to 'acanthodians'

with tooth whorls and the unusual dentitions in more shark-like taxa (e.g.

Pucapampella, *Gladbachus*. Novel oral morphologies have been linked to a

period of inferred rapid gnathostome evolution [7,43]. Furthermore, in the

Devonian small-bodied chondrichthyans were significant in freshwater

nektonic faunas [24]: diverse oral structures seem likely to have accompanied

their radiation into these niches. While the latest surviving lineage of the

'acanthodian' grade was the remarkably morphologically conservative

acanthodids [79], *Acanthodopsis* shows that experimentation with novel oral

apparatus in stem-group chondrichthyans continued well into the Late

Carboniferous.

733 **4. Summary**

'Acanthodian' stem-group chondrichthyans display a diverse array of oral and

dental morphologies, including an apparently independent origin of teeth deep

within an edentulous clade. However, interpreting patterns of tooth evolution

is complicated by conflicting and unresolved phylogenetic hypotheses, both

for the chondrichthyan (figure 7; figures S1,2) and gnathostome stem-group

(e.g. [6,43]). CT- and synchrotron-based investigations seem likely to provide

the anatomical information necessary to resolve these instabilities, and

likelihood-based methods provide another potential way of overcoming

uncertainties [7]. In the meantime, the proliferation of different tree shapes in

conjunction with generally low support values means that morphologies

should be considered across multiple potential topologies. This illustrates the

challenges of drawing broad scale conclusions for gnathostome tooth

evolution on the basis of unstable relationships or tentatively placed taxa.

748 **Acknowledgements**

We thank E. Bernard and Z. Johanson (NHMUK) for assistance with

specimen access, V. Fernandez and B. Clark (both NHMUK) and T. Davies

(University of Bristol) for assistance with CT scanning. M. Brazeau (Imperial

College London), B. Davidson, and C. Burrow (Queensland Museum) and P.

Ahlberg (Uppsala University) contributed to helpful discussion. M. Colfer

(University of Oxford) carried out preliminary segmentation and interpretation

of the Mimics files. Four anonymous reviewers provided helpful comments,

and we also thank two anonymous reviewers for their comments on an earlier

version of this manuscript.

Funding

This work was supported by a Junior Research Fellowship, Christ Church,

Oxford, and a Royal Society Dorothy Hodgkin Research Fellowship, both to

S.G.. R.P.D. was supported by a Paris Île-de-France Région – DIM

"Matériaux anciens et patrimoniaux" grant (PHARE project).

Ethics. This research is based exclusively on specimens from natural history

collections.

Data accessibility. Raw data (.vol or .tiff stacks), Mimics files, and 3D PLY

files for each specimen are deposited in Zenodo (10.5281/zenodo.5238205).

Authors' contributions. S.G. conceived the project and selected specimens.

S.G. and R.P.D. carried out CT scanning. R.P.D. segmented the specimens,

made Blender renders and constructed figures with input from S.G. S.G. and

R.P.D. drafted the manuscript. Both authors revised and edited the

manuscript, approved the final version and agree to be accountable for all

aspects of the work.

Competing interests. We have no competing interests.

5. References

1. Tucker AS, Fraser GJ. 2014 Evolution and developmental diversity of tooth regeneration. *Semin. Cell Dev. Biol.* **25–26**, 71–80. (doi:10.1016/j.semcdb.2013.12.013)

2. Jobbins M, Rücklin M, Argyriou T, Klug C. 2021 A large Middle Devonian eubrachythoracid 'placoderm' (Arthrodira) jaw from northern Gondwana. *Swiss J. Palaeontol.* **140**, 2. (doi:10.1186/s13358-020-00212-w)

3. Ørvig T. 1980 Histologic Studies of Ostracoderms, Placoderms and Fossil Elasmobranchs: 3. Structure and growth of the gnathalia of certain arthrodires'. *Zool. Scr.* **9**, 141–159. (doi:10.1111/j.1463-6409.1980.tb00660.x)

4. Rücklin M, Donoghue PCJ. 2015 *Romundina* and the evolutionary origin of teeth. *Biol. Lett.* **11**, 0–4.

5. Rücklin M, Donoghue PCJ, Johanson Z, Trinajstic K, Marone F, Stampanoni M. 2012 Development of teeth and jaws in the earliest jawed vertebrates. *Nature* (doi:10.1038/nature11555)

6. Vaškaninová V, Chen D, Tafforeau P, Johanson Z, Ekrt B, Blom H, Ahlberg PE. 2020 Marginal dentition and multiple dermal jawbones as the ancestral condition of jawed vertebrates. *Science* **369**, 211–216. (doi:10.1126/science.aaz9431)

7. Rücklin M, King B, Cunningham JA, Johanson Z, Marone F, Donoghue PCJ. 2021 Acanthodian dental development and the origin of gnathostome dentitions. *Nat. Ecol. Evol.* , 14. (doi: 10.1038/s41559-021-01458-4)

8. Hu YZ, Young GC, Burrow CJ, an Zhu Y, Lu J. 2018 High resolution XCT scanning reveals complex morphology of gnathal elements in an Early Devonian arthrodire. *Palaeoworld* (doi:10.1016/j.palwor.2018.12.003)

9. Botella H, Blom H, Dorka M, Ahlberg PE, Janvier P. 2007 Jaws and teeth of the earliest bony fishes. *Nature* **448**, 583–586. (doi:10.1038/nature05989)

10. Chen D, Blom H, Sanchez S, Tafforeau P, Ahlberg PE. 2016 The stem osteichthyan Andreolepis and the origin of tooth replacement. *Nature* (doi:10.1038/nature19812)

11. Chen D, Blom H, Sanchez S, Tafforeau P, Märss T, Ahlberg PE. 2017 Development of cyclic shedding teeth from semi-shedding teeth : the inner dental arcade of the stem osteichthyan *Lophosteus R. Soc. open sci.* **4**: 161084.

12. Chen D, Blom H, Sanchez S, Tafforeau P, Märss T, Ahlberg PE. 2020 The developmental relationship between teeth and dermal odontodes in the

most primitive bony fish Lophosteus. *eLife* **9**, e60985.
(doi:10.7554/eLife.60985)

13. Cunningham JA, Rücklin M, Blom H, Botella H, Donoghue PCJ. 2012
Testing models of dental development in the earliest bony vertebrates,
Andreolepis and Lophosteus. *Biol. Lett.* **8**, 833–837.
(doi:10.1098/rsbl.2012.0357)

14. Doeland M, Couzens AMC, Donoghue PCJ, Rücklin M. 2019 Tooth
replacement in early sarcopterygians. *R. Soc. Open Sci.* **6**.

15. Zhu M, Ahlberg P E,Pan, Z, Zhu Y, Qiao T, Zhao W, Jia L, Lu J. 2016 A
Silurian maxillate placoderm illuminates jaw evolution. *Science* **354** 6310
334-336.

16 Zhu Y, Ahlberg, P A, Zhu M. 2018 The evolution of vertebrate dermal jaw
bones in the light of maxillate placoderms. In *Evolution and Development*
of Fishes (Ed. Zerina Johanson) 71-86.

17. Maisey JG, Turner S, Naylor GJP, Miller RF. 2014 Dental patterning in the
earliest sharks: Implications for tooth evolution. *J. Morphol.* **275**, 586–96.
(doi:10.1002/jmor.20242)

18. Maisey JG, Janvier P, Pradel A, Denton JSS, Bronson A, Miller RF,
Burrow CJ. 2019 *Doliodus* and pucapampellids: contrasting perspectives
on stem chondrichthyan morphology. In *Evolution and Development of*
Fishes (eds Z Johanson, C Underwood, M Richter), pp. 87–109.
Cambridge: Cambridge University Press.

19. Qu Q, Sanchez S, Blom H, Tafforeau P, Ahlberg PE. 2013 Scales and
Tooth Whorls of Ancient Fishes Challenge Distinction between External
and Oral 'Teeth'. *PLoS ONE* **8**. (doi:10.1371/journal.pone.0071890)

20. Rücklin M, Giles S, Janvier P, Donoghue PCJ. 2011 Teeth before jaws?
Comparative analysis of the structure and development of the external
and internal scales in the extinct jawless vertebrate Loganellia scotica.
Evol. Dev. **13**, 523–532. (doi:10.1111/j.1525-142X.2011.00508.x)

21. Miles RS. 1973 Articulated acanthodian fishes from the Old Red
Sandstone of England, with a review of the structure and evolution of the
acanthodian shoulder girdle. *Bulletin of the British Museum (Natural*
History). Geology. **24**, 111–213.

22. Burrow CJ, Newman MJ, den Blaauwen J, Jones R, Davidson RG. 2018
The Early Devonian ischnacanthiform acanthodian *Ischnacanthus gracilis*
(Egerton, 1861) from the Midland Valley of Scotland. *Acta Geol. Pol.* **68**,
335–362. (doi: 10.1515/agp-2018-0008)

23. Burrow CJ. 2004 Acanthodian fishes with dentigerous jaw bones : the
Ischnacanthiformes and *Acanthodopsis*. *Foss. Strata* , 8–22.

857
24. Coates MI, Finarelli JA, Sansom IJ, Andreev PS, Criswell KE, Tietjen K,
Rivers ML, Riviere PJL, Coates MI. 2018 An early chondrichthyan and the
evolutionary assembly of a shark body plan. *Proc. R. Soc. B Biol. Sci.* **285**.

25. Blais SA, Hermus CR, Wilson MVH. 2015 Four new Early Devonian
ischnacanthid acanthodians from the Mackenzie Mountains, Northwest
Territories, Canada: an early experiment in dental diversity. *J. Vertebr.*
Paleontol. **35**, 37–41. (doi:10.1080/02724634.2014.948546)

26. Ørvig T. 1973 Acanthodian dentition and its bearing on the relationships of
the group. *Palaeontogr. Abt. A* **143**, 119–150.

27. Smith, MM. 2003 Vertebrate dentitions at the origin of jaws: when and how
pattern evolved. *Evolution & Development.* **5**(4) 394-413 (doi:
10.1046/j.1525-142X.2003.03047.x)

28. Burrow CJ, Blaauwen JD, Newman MJ, Davidson RG. 2016 The
diplacanthid fishes (Acanthodii, Diplacanthiformes, Diplacanthidae) from
the Middle Devonian of Scotland. *Palaeontol. Electron.* , 1–83.

29. Hanke GF, Davis SP, Wilson MVH. 2001 New species of the acanthodian
genus *Tetanopsyrus* from northern Canada, and comments on related
taxa. *J. Vertebr. Paleontol.* **21**, 740–753.

30. Newman MJ, Davidson RG, Blaauwen JLD, Burrow CJ. 2012 The Early
Devonian Acanthodian *Uraniacanthus curtus* (Powrie , 1870) n . comb .
from the Midland Valley of Scotland. *Geodiversitas* **34**, 739–759.

31. Zidek J. 1976 Kansas Hamilton Quarry (Upper Pennsylvanian)
Acanthodes, with remarks on the previously reported North American
occurrences of the Genus. *Univ. Kans. Palaeontol. Contrib.* , 1–41.

32. Blais SA, MacKenzie LA, Wilson MVH. 2011 Tooth-like scales in Early
Devonian eugnathostomes and the outside-in hypothesis for the origins of
teeth in vertebrates. *J. Vertebr. Paleontol.* **31**, 1189–1199.
(doi:10.1080/02724634.2011.607992)

33. Meredith Smith M, Underwood C, Goral T, Healy C, Johanson Z. 2019
Growth and mineralogy in dental plates of the holocephalan *Harriotta*
raleighana (Chondrichthyes): Novel dentine and conserved patterning
combine to create a unique chondrichthyan dentition. *Zoological Letters* **5**,
1–30. (doi:10.1186/s40851-019-0125-3)

34. Frey L, Coates MI, Ginter M, Hairapetian V, Rücklin M, Jerjen I, Klug C.
2019 The early elasmobranch *Phoebodus* :phylogenetic relationships,
ecomorphology and a new time-scale for shark evolution. *Proceedings of*
the Royal Society B: Biological Sciences **286**. (doi:
10.1098/rspb.2019.1336)

35. Frey L, Coates MI, Tietjen K, Rücklin M, Klug C. 2020 A symmoriiform
from the Late Devonian of Morocco demonstrates a derived jaw function in
ancient chondrichthyans. *Commun. Biol.* **3**, 681. (doi:10.1038/s42003-020-
01394-2)

36. Williams ME. 2001 Tooth retention in cladodont sharks: with a comparison
between primitive grasping and swallowing, and modern cutting and
gouging feeding mechanisms. *Journal of Vertebrate Paleontology* **21**,
214–226. (doi:10.1671/0272)

37. Hanke GF. 2008 *Promesacanthus eppleri n. gen., n. sp*., a mesacanthid
(Acanthodii, Acanthodiformes) from the Lower Devonian of northern
Canada. *Geodiversitas* **30**, 287–302.

38. Hanke GF, Davis SP. 2008 Redescription of the acanthodian
Gladiobranchus probaton Bernacsek &Dineley, 1977, and comments on
diplacanthid relationships. *Geodiversitas* **30**, 303–330.

39. Ørvig T. 1967 Some new acanthodian material from the lower Devonian of
Europe. *Zool. J. Linn. Soc.* **47**, 131–153.

40. Zhu M *et al.* 2013 A Silurian placoderm with osteichthyan-like marginal jaw
bones. *Nature* **502**, 188–93. (doi:10.1038/nature12617)

41. Giles S, Friedman M, Brazeau MD. 2015 Osteichthyan-like cranial
conditions in an Early Devonian stem gnathostome. *Nature* **520**, 82–175.
(doi:10.1038/nature14065)

42. Chevrinais M, Sire J-Y, Cloutier R. 2017 From body scale ontogeny to
species ontogeny: Histological and morphological assessment of the Late
Devonian acanthodian *Triazeugacanthus affinis* from Miguasha, Canada.
PLOS ONE **12**, e0174655. (doi:10.1371/journal.pone.0174655)

43. King B, Qiao T, Lee MSY, Zhu M, Long JA. 2016 Bayesian Morphological
Clock Methods Resurrect Placoderm Monophyly and Reveal Rapid Early
Evolution in Jawed Vertebrates. *Syst. Biol.* **66**, 499–516.
(doi:10.1093/sysbio/syw107)

44. Dearden RP, Stockey C, Brazeau MD. 2019 The pharynx of the stem-
chondrichthyan *Ptomacanthus* and the early evolution of the gnathostome
gill skeleton. *Nat. Commun.* **10**, 1–7. (doi:10.1038/s41467-019-10032-3)

45. Long JA. 1986 New ischnacanthid acanthodians from the Early Devonian
of Australia, with comments on acanthodian interrelationships. *Zool. J.
Linn. Soc.* **87**, 321–339.

46. Lindley ID. 2000 Acanthodian fish remains from the Lower Devonian
Cavan Bluff Limestone (Murrumbidgee Group), Taemas district, New
South Wales. *Alcheringa* **24**, 11–35.

47. Lindley ID. 2002 Lower Devonian ischnacanthid fish (Gnathostomata:
Acanthodii) from the Taemas Limestone, Lake Burrinjuck, New South

Wales. *Alcheringa Australas. J. Palaeontol.* **25**, 269–291.
(doi:10.1080/03115510108527804)

48. Burrow CJ. 2004 A Redescription of *Atopacanthus dentatus* Hussakof and
Bryant , 1918 (Acanthodii , Ischnacanthidae). *Jvp* **24**, 257–267.
(doi:10.1671/1928)

49. Hussakof L, Bryant WL. 1918 Catalog of the fossil fishes in the Museum of
Buffalo Society of Natural Sciences. *Bull. Buffalo Soc. Nat. Sci.* **12**, 5–178.

50. Jessen H. 1973 Weitere fischreste aus dem Oberen Plattenkalk der
Bergisch-Gladbach Paffrather Mulde (Oberdevon, Rheinisches
Schiefergeb.pdf. *Palaeontogr. Abt. A* **1–6**, 159–187.

51. Denison R. 1979 *Handbook of Palaeoichthyology: Acanthodii*. Verlag Dr.
Friedrich Pfeil.

52. Newman MJ, Burrow CJ, Blaauwen JLD. 2019 A new species of
ischnacanthiform acanthodian from the Givetian of Mimerdalen , Svalbard.
Nor. J. Geol. **99**, 1–13.

53. Dineley DL, Metcalf S J. 1999 *Fossil Fishes of Great Britain*.
Peterborough: Joint Nature Conservation Committee.

54. Miles RS. 1973 Relationships of acanthodians. In *Interrelationships of*
fishes (eds P Greenwood, RS Miles, C Patterson), pp. 63–103. London:
Zoological Journal of the Linnean Society.

55. Goloboff PA, Catalano SA. 2016 TNT version 1.5, including a full
implementation of phylogenetic morphometrics. *Cladistics* **32**, 221–238.

56. Gross W. 1971 Downtonische und Dittonische acanthodier-reste des
Ostseegebietes. *Palaeontogr. Abt. A* **136**, 1–82.

57. Miles RS. 1966 The acanthodian fishes of the Devonian Plattenkalk of the
Paffrath Trough in the Rhineland with an appendix containing a
classification of the Acanthodii and a revision of the genus
Homalacanthus. *Ark. För Zool.* **18**, 147–194.

58. Blais SA, MacKenzie LA, Wilson MVH. 2011 Tooth-like scales in Early
Devonian eugnathostomes and the outside-in hypothesis for the origins of
teeth in vertebrates. *J. Vertebr. Paleontol.* **31**, 1189–1199.
(doi:10.1080/02724634.2011.607992)

59. Burrow CJ, Blaauwen JD, Newman M. 2020 A redescription of the three
longest-known species of the acanthodian *Cheiracanthus* from the Middle
Devonian of Scotland. *Palaeontol. Electron.* **23**, 1–43.

60. Zidek J. 1985 Growth in *Acanthodes* (Acanthodii: Pisces) data and
implications. *Palaontologische Z.* **59**, 147–166.

61. Schultze H-P. 1972 *Homalacanthus*, ein oberdevonishcer Acanthodier mit
Haifisch-ähnlichen Zähnen. *Neues Jahrb. Für Geol. Paläontol.* , 315–320.

62. Watson DMS. 1937 The Acanthodian Fishes. *Philos. Trans. R. Soc. Lond.*
B. Biol. Sci. **228**, 49–146. (doi:10.2307/1436828)

63. Hanke GF, Wilson MVH. 2004 New teleostome fishes and acanthodian
systematics. In *Recent Advances in the Origin and Early Radiation of*
Vertebrates (eds G Arratia, MVH Wilson), pp. 189–216. München,
Germany.

64. Newman MJ, Burrow CJ, Blaauwen JLD, Davidson RG. 2014 The Early
Devonian acanthodian *Euthacanthus macnicoli* Powrie, 1864 from the
Midland Valley of Scotland. *Geodiversitas* **36**, 321.
(doi:10.5252/g2014n3a1)

65. Hanke GF, Wilson MVH. 2010 The putative stem-group chondrichthyans
Kathemacanthus and *Seretolepis* from the Lower Devonian MOTH
locality, Mackenzie Mountains, Canada. In *Phylogeny and*
Paleobiogeography of Fossil Fishes (eds DK Elliott, JG Maisey, X Yu, D
Miao), pp. 159–182. München, Germany: Verlag Dr. Friedrich Pfeil.

66. Smith MM, Coates MI. 1998 Evolutionary origins of the vertebrate
dentition: phylogenetic patterns and developmental evolution. *Eur. J. Oral*
Sci. **106**, 482–500

67. Burrow CJ, Davidson RG, Den Blaauwen JL, Newman MJ. 2015 Revision
of *Climatius reticulatus* Agassiz, 1844 (Acanthodii, Climatiidae), from the
Lower Devonian of Scotland, based on new histological and morphological
data. *J. Vertebr. Paleontol.* **35**, e913421.
(doi:10.1080/02724634.2014.913421)

68. Schultze H, Zidek J. 1982 Ein primitiver acanthodier (pisces) aus dem
unterdevon lettlands. *Paläontologische Z.* **56**, 95–105.
(doi:10.1007/BF02988788)

69. Zangerl R. 1981 *Handbook of Paleoichthyology, Chondrichthyes I,*
Paleozoic Elasmobranchii. Stuttgart: Gustav Fischer Verlag.

70. Valiukevičius J. 1992 First articulated *Poracanthodes* from the Lower
Devonian of Severnaya Zemlya. In *Fossil Fishes as Living Animals*, pp.
193–213.

71. Valiukevičius J. 2003 Devonian acanthodians from Severnaya Zemlya
Archipelago (Russia). **25**, 131–204.

72. Giles S, Coates MI, Garwood RJ, Brazeau MD, Atwood R, Johanson Z,
Friedman M. 2015 Endoskeletal structure in *Cheirolepis* (Osteichthyes,
Actinopterygii), An early ray-finned fish. *Palaeontology* **58**.
(doi:10.1111/pala.12182)

73. Burrow CJ, Rudkin D. 2014 Oldest near-complete acanthodian: The first
vertebrate from the silurian bertie formation Konservat-Lagerstätte,
Ontario. *PLoS ONE* **9**, e104171. (doi:10.1371/journal.pone.0104171)

74. Burrow CJ, Young GC. 2012 New Information on *Culmacanthus*
(Acanthodii : Diplacanthiformes) from the ?Early – Middle Devonian of
Southeastern Australia. *Proc. Linn. Soc. New South Wales* **134**, 21–29.

75. Hanke GF, Davis SP. 2008 Redescription of the acanthodian
Gladiobranchus probaton Bernacsek &Dineley, 1977, and comments on
diplacanthid relationships. *Geodiversitas* **30**, 303–330.

76. Young GC, Burrow CJ. 2004 Diplacanthid acanthodians from the Aztec
Siltstone (late Middle Devonian) of southern Victoria Land, Antarctica.
Foss. Strata **50**, 23-43

77. Gagnier P, Hanke GF, Wilson MVH. 1999 *Tetanopsyrus lindoei* gen. et sp.
nov., an Early Devonian acanthodian from the Northwest Territories,
Canada. *Acta Geol. Pol.* **49**, 81–96.

78. Hanke GF, Davis SP. 2008 Redescription of the acanthodian
Gladiobranchus probaton Bernacsek &Dineley, 1977, and comments on
diplacanthid relationships. *Geodiversitas* **30**, 303–330.

79. Burrow CJ, Trinajstic K, Long JA. 2012 First acanthodian from the Upper
Devonian (Frasnian) Gogo Formation , Western Australia. *Hist. Biol.* **24**,
349–357. (doi:10.1080/08912963.2012.660150)

80. Long JA. 1986 A new Late Devonian acanthodian fish from Mt. Howitt,
Victoria, Australia, with remarks on acanthodian biogeography. *Proc. R.*
Soc. Vic. **98**, 1–17.

81. Grogan ED, Lund R. 2000 *Debeerius ellefseni* (Fam. Nov., Gen. Nov.,
Spec. Nov.), an Autodiastylic Chondrichthyan From the Mississippian Bear
Gulch Limestone of Montana (USA), the Relationships of the
Chondrichthyes, and Comments on Gnathostome Evolution. *Journal of*
Morphology. **243:** 219-245.

82. Kamminga P, De Bruin PW, Geleijns J, Brazeau MD. 2017 X-ray
computed tomography library of shark anatomy and lower jaw surface
models. *Scientific Data*, 1–6.

Figure captions

Figure 1. Summary figure showing lower jaws of six chondrichthyans, and major structures discussed in the text. (a) the extant elasmobranch *Isurus* in medial view and with tooth series in lateral view, drawn from 3D data [82]; (b) the stem-chondrichthyan *Ptomacanthus* in medial view reconstructed from [44] and with tooth whorl from [21] (c) the ischnacanthid stem-chondrichthyan *Erymnacanthus* in medial view redrawn from [25] (d) the acanthodid stem-chondrichthyan *Acanthodes* in lateral view redrawn from [54] (e) the ?acanthodid stem-chondrichthyan *Acanthodopsis* in lateral view redrawn from Burrow [23]; (f) the diplacanthid stem-chondrichthyan *Diplacanthus* in medial view reconstructed from Watson *et al.*[61] and Burrow *et al.* [28] Not to scale.

1056

1057

Figure 2. Tomographic model of the left lower jaw of *Taemasacanthus erroli*

NHMUK PV P.33706 in (*a*) medial view, (*b*) lateral view, (*c*) ventral view, (*d*)

dorsal view, (*e*) posterior view, and reconstructed tomograms showing (*f*) a

transverse section through the circular ridge, (*g*) a transverse section through

the lower jaw, (*h*) a sagittal section through the lingual tooth row and (*i*)-(*j*)

progressively medial sagittal sections through the aberrant youngest cusp and

neighbouring cusps. Teeth in panels (*a*), (*b*) and (*d*) are coloured separately

from the dentigerous jaw bone. Arrow indicates direction of anterior, and top

right inset shows location of rendered jaw components. Abbreviations: art,

articular (Meckel's cartilage); art.con, articular 'condyle'; circ.ri, circular ridge;

djb, dentigerous jaw bone; lat.t.r., lateral tooth row; ling.t.r, lingual tooth row;

mes.ri, mesial ridge; mes.t.r, mesial tooth row; post.gr, posterior groove;

young.t, out-of-order youngest tooth; vent.gr, ventral groove; vas.can,

vascular canals. Scale bar = 5 mm.

1072

1073

Figure 3. Tomographic model of a dentigerous jaw bone of *Atopacanthus* sp.

NHMUK PV P.10978 in (*a*) medial view, (*b*) lateral view, (*c*) ventral view, (*d*)

dorsal view and (*e*) a reconstructed tomogram showing a sagittal section

through the lingual tooth row. Arrow indicates direction of anterior, and top

right inset shows possible locations of rendered jaw components. Teeth in

panels (*a*), (*b*) and (*d*) are coloured separately from the dentigerous jaw bone.

Abbreviations: add.c, additional cusps; djb, dentigerous jaw bone; lat.t.r.,

1082   lateral tooth row; ling.pl, lingual plate; ling.t.r, lingual tooth row; mes.ri, mesial

1083   ridge; mes.t.r, mesial tooth row; vent.gr, ventral groove; vas.can, vascular

1084   canals. Scale bar = 5 mm in (*a*)-(*e*) and (*h*)-(*j*), 2 mm in (*f*)-(*h*).

1085

1086

1087

Figure 4. Tomographic model of the left lower jaw of *Ischnacanthus* sp.

NHMUK PV P.40124 in (*a*) lateral view, (*b*) medial view, and (*c*), (*d*), (*e*)

reconstructed tomograms showing successively deeper sagittal sections.

Arrow indicates direction of anterior, and top right inset shows location of

rendered jaw components. Abbreviations: art.proc, articular process; djb

dentigerous jaw bone; lat.t.r., lateral tooth row; Meck.c, Meckel's cartilage;

mes.ri, mesial ridge; per, perichondral bone; rid, ridge; vas.can, vascular

canal. Scale bar = 5 mm.

1096

1097

1098

Figure 5. Tomographic model of the right lower jaw of *Acanthodopsis* sp.

NHMUK PV P.10383 in lateral view with (*a*) and without (*b*) mandibular splint,

(*c*) medial view, (*d*) dorsal view, (*e*) anteromedial view, (*f*) posterior view, and

reconstructed tomograms showing (*g*) a sagittal section through the entire jaw

and (*h*) a transverse section through the jaw. Arrow indicates direction of

anterior, and top right inset shows location of rendered jaw components.

Abbreviations: art.cot, articular cotylus; gr.mand.spl, groove for mandibular

splint; inn.lay, inner layer; mand.add.fo, mandibular adductor fossa; mand.spl,

mandibular splint; Meck.c, Meckel's cartilage; out.lay, outer layer; pgl.pr,

preglenoid process; sym.fos, symphyseal fossa; 't', 'teeth'. Scale bar = 5 mm

in (*a*)-(*g*), 2 mm in (*h*).

1112

Figure 6. The lower jaws of *Acanthodes* sp. NHMUK PV P.8065 in (*a*) dorsal

view against the matrix, (*b*) in ventral view superimposed on a digital mould of

the matrix's surface, the left lower jaw isolated in (*c*) medial and (*d*) lateral

view, and reconstructed tomograms showing (*e*) a coronal section through the

specimen, and (*f*) a transverse section through a lower jaw. Arrow indicates

direction of anterior, and top right inset shows location of rendered jaw

components. Abbreviations: art, articular (Meckel's cartilage); br,

branchiostegal rays; mand.spl, mandibular splint; Meck.c, Meckel's cartilage;

ment, mentomandibular (Meckel's cartilage); pgl.pr, preglenoid process; rak,

gill raker; sym.fos, symphyseal fossa; vis.ar, visceral arch fragments. Scale

bar = 5 mm in (*a*)-(*e*), 2mm in (*f*).

1124

Figure 7. Phylogenetic relationships of early chondrichthyans and distribution

of oral structures. Strict consensus of 26101 most parsimonious trees on left

and Bayesian analysis on right, with some non-chondrichthyan taxa excluded

(full tree with support values in figures S2 and S3). Numbers at nodes on right

represent Bayesian posterior probabilities. Character distribtion is based on

coding in the data matrix as follows: teeth, character (c.) 82; teeth ankylosed,

c.93; tooth whorls fused, c. 85; whorls at symphysis and whorls entire

dentition, c.88; teeth on jaw cartilages and dermal jaw plate, c.94 ; occlusal

plate, c.269 ; mandibular splint, c.268. Filled boxes indicate presence of

1135   feature; 'x' indicates feature absent; '?' indicates uncertainty; '-'indicates

1136   inapplicability.

1137

Figure 8. Summaries of four contrasting recent phylogenetic schemes of early

chondrichthyans, with distribution of oral characters. (a) King *et al.* 2016 [43],

(b) Chevrinais *et al.* 2017 [42], (c) Coates *et al.* 2018 [24], (d) Frey *et al.* 2020

1142   [35]. Icons and character numbers as in Fig. 7 Acanthodiforms includes taxa

1143   both with and without mandibular splints.

1144

Supplementary material (separate pdf)

Supplementary information for this manuscript is included as two files. The

first includes notes on the phylogenetic analysis (taxon addition, character

addition and coding changes) supplementary table 1,and supplementary

figures 1-3. The second is a zipped folder containing all files necessary for

replicating our Bayesian and parsimony analyses.

1151