Diversity of function and higher-order structure within HWE sensor histidine kinases

Igor Dikiy[1], Danielle Swingle[1,2], Kaitlyn Toy[1,3], Uthama R. Edupuganti[1,2], Giomar Rivera-Cancel[4], Kevin H. Gardner[1,3,5*]

[1]: Structural Biology Initiative, CUNY Advanced Science Research Center, New York, NY 10031

[2]: Ph.D. Program in Biochemistry, The Graduate Center – City University of New York, New York, NY 10016

[3]: Department of Chemistry and Biochemistry, City College of New York, New York, NY 10031

[4]: Department of Internal Medicine, UT Southwestern Medical Center, Dallas, TX 75390

[5]: Ph.D. Programs in Biochemistry, Biology, and Chemistry, The Graduate Center – City University of New York, New York, NY 10016

*: direct correspondence to kgardner@gc.cuny.edu

ORCIDs:

Igor Dikiy: 0000-0002-8155-7788

Danielle Swingle: 0000-0003-3691-4560

Kaitlyn Toy: 0000-0003-2595-8779

Uthama R. Edupuganti: 0000-0003-0571-4101

Giomar Rivera-Cancel: 0000-0002-4909-4103

Kevin H. Gardner: 0000-0002-8671-2556

Keywords: histidine kinase, bacterial signal transduction, conformational change, structure-function, biophysics, environmental sensing, two-component system, HWE family, HisKA2 family

Running title: Diversity of function and structure in HWE family sensor HKs

Abstract

Integral to the protein structure/function paradigm, oligomeric state is typically conserved

along with function across evolution. However, notable exceptions such as the hemoglobins

show how evolution can alter oligomerization to enable new regulatory mechanisms. Here we

examine this linkage in histidine kinases (HKs), a large class of widely distributed prokaryotic

environmental sensors. While the majority of HKs are transmembrane homodimers,

members of the HWE/HisKA2 family can deviate from this architecture as exemplified by our

finding of a monomeric soluble HWE/HisKA2 HK (EL346, a photosensing Light-Oxygen-Voltage

(LOV)-HK). To examine the possibility of a wider range of possible relationships between

oligomerization and regulation within this family, we biophysically and biochemically

characterized multiple EL346 homologs and found a diverse range of HK oligomeric states and

functions. Three LOV-HK homologs are dimers with differing structural and functional responses

to light, while two are Per-ARNT-Sim (PAS)-HKs that interconvert between differentially active

monomers and dimers, suggesting dimerization as a regulatory control. Finally, we examined

putative interfaces in a dimeric LOV-HK, finding that multiple domains contributed to

dimerization and function. Our findings suggest the potential for novel regulatory modes and

oligomeric states beyond those traditionally defined for this important family of environmental

sensors.

Introduction

Sensor histidine kinases (HKs) are signal transduction receptors that are widespread in

prokaryotes, enabling these organisms to sense and respond to diverse environmental stimuli

(4, 5). HKs are typically found in two-component systems (TCS), which are most simply

composed of a sensor HK and a downstream response regulator (RR) co-located in the same

operon. The RR responds to a signal from the HK by acting as a phosphorylation-dependent

transcription regulator, with activation often triggering RR DNA binding and altering the

expression of nearby genes (6-9).

The prototypical sensor HK contains N-terminal sensor and C-terminal kinase domains,

the latter of which can be subdivided into a two-helix dimerization histidine phosphotransfer

(DHp) domain and a catalytic ATP-binding (CA) domain (10). The detection of a stimulus by the

sensor domain (or in some cases, elsewhere (9)) modulates the CA domain

autophosphorylation of a conserved His residue in the DHp domain; the phosphate group in this

phospho-His adduct is subsequently transferred to a conserved Asp in a downstream RR (6).

Most characterized sensor HKs are obligate homodimeric transmembrane receptors,

facilitating the transmission of signals across the membrane from periplasmic sensor domains to

cytoplasmic kinase domains via symmetry-breaking conformational changes (10-13). The

regulation of typical sensor HKs is largely believed to require a dimeric complex, both for the

structural changes between the "off" and "on" state as well as determining whether

autophosphorylation occurs between protomers (*in trans*) or within a protomer (*in cis*) (10, 14).

Dimerization is mediated via the DHp domains of two monomers, forming a four-helix bundle

and facilitating pivot-, piston-, or scissoring-type movements as part of the activation process

(10, 15), echoing similar themes in integrins (16) and certain other transmembrane receptors. In

contrast, some sensors adopt different oligomeric states, such as the large multimeric

assemblies of chemotaxis HKs (17), hexameric KaiC circadian HKs (18) or the monomeric

photosensory EL346 HK (1, 19).

The traditional view of sensor HK structure has been built up from extensive structural

and biochemical studies (10) that have usually focused on HKs in only one of several

subfamilies of histidine kinases (the HisKA family within Pfam (20)). In contrast, the related

HWE (His-Trp-Glu)-HK and HisKA_2 families (referred to here as the HWE/HisKA2 family) of

HKs have been less studied from a structural perspective (15). Members of this group, which

contain specific sequence variations in the DHp and CA domains compared to members of the

canonical HisKA family, tend to signal to a more diverse set of output proteins which often lack

DNA-binding output domains (15). The HWE/HisKA2 family is also enriched in HKs lacking

transmembrane segments, likely localizing them to the cytosol. Notably, soluble sensor HKs are

relieved of the constraints of transmitting a signal across a lipid bilayer, which typically involves

signal transmission via motions of one protomer with respect to the other (10, 15). This raises

several questions: Are such soluble HKs able to adopt diverse oligomeric states, including non-

dimeric architectures? And could these varied architectures be equivalently varied in their

regulation? We view the relatively understudied HWE/HisKA2 family of HKs (15) as a suitable

testbed for these questions.

Our prior work on the HWE/HisKA2 family focused on three soluble HKs involved in the

general stress response of the marine alphaproteobacterium *Erythrobacter litoralis* HTCC2594

(EL346, EL362, and EL368) (1, 3, 19, 21). Each of these proteins senses blue light through a

light oxygen voltage (LOV) sensor domain (22, 23), a sub-class of the versatile Per-ARNT-Sim

(PAS) domains that are present in over 30% of HKs (5, 24). The response to blue light provides

a readily tractable way to explore the inactive and active states of histidine kinases, typically

corresponding to dark and light states of LOV-HKs (21, 25, 26). These states can be

conveniently distinguished spectrophotometrically: while the dark state spectra of LOV-HKs
exhibit a triplet peak at around 450 nm, characteristic of protein-bound flavin chromophores, the
light state spectrum has a broad, singlet absorption peak at 390 nm (27). This spectroscopic
change is due to the photochemically-triggered reduction of the flavin chromophore and
concomitant formation of a cysteine-C(4a) covalent adduct, a bond that is thermally reversible in
the dark on a timescale varying from seconds to hours among LOV domains (23).
While our prior studies indicate that EL368 is dimeric, EL346 is a soluble and functional
monomer (1, 3). The EL346 crystal structure reveals that it is held in a monomeric state by an
intramolecular association between its DHp-like domain and the LOV sensory domain, blocking
the face typically used for DHp domains to dimerize (1). Despite this deviation from current HK
signaling models, the monomeric EL346 undergoes light-induced increases in
autophosphorylation rate (1) as well as conformational changes involving the sensor, DHp, and
kinase domains (19). More broadly, the active monomer of EL346 suggests that soluble HKs
may not require dimerization as a key element in signal transduction, unlike conventional
transmembrane HKs.
To address this general issue in HK signaling, we set out to characterize the oligomeric
and functional states of several novel HWE/HisKA2 HKs. To do so, we used sequence similarity
searches to identify multiple soluble LOV- and PAS-HKs closely related to EL346, selecting five
proteins for in-depth analyses. We showed that three of these five are novel dimeric LOV-HKs,
each of which properly photocycles by UV-visible absorbance spectroscopy. Notably, these
LOV-HKs differed in the effect of light on their autophosphorylation activity, with a mix of dark-
and light-activated proteins among them. In addition, we also characterized two light-insensitive
PAS-HKs which equilibrate between monomeric and dimeric states with differing activities,
suggesting dimerization as a possible regulatory switch in this family of HKs. Finally, we
combined structural and sequence information to engineer a "monomerized dimer" by deleting
putative dimerization interfaces in a dimeric LOV-HK, finding that broadly distributed residues
contribute to both dimerization and autokinase activity. Taken together, our data reveal a wider
range of oligomerization states and functional activity in HKs than traditionally described.
124
Results
Bioinformatics analysis reveals numerous alphaproteobacterial EL346 homologs
To identify other potential monomeric histidine kinases, we searched for enzymes highly
similar in domain organization to the well-validated and functional EL346 monomer. In addition
to EL346, the top 100 BLASTp hits defined a set of proteins from *Alphaproteobacteria* which all

contained the same domain architecture of a single N-terminal PAS or LOV domain followed by

a C-terminal HWE/HisKA2 histidine kinase. None of these sequences contained any predicted

transmembrane segments, strongly suggesting all these candidates are soluble. A multiple

sequence alignment of these protein sequences and distance tree analysis revealed three

clusters (**Fig. 1a**), which split on increasingly diverse sensor types: EL346-like LOV-HKs, LOV-

HKs, and PAS-HKs (non-LOV). LOV domains were provisionally identified by the presence of

the characteristic "NCRFLQ" sequence and high homology (> 30% identity) to the EL346 LOV

domain (28), while the non-LOV PAS domains lacked this sequence motif (highlighted in **Fig.**

S1). Interestingly, a subgroup of the EL346-like LOV-HKs is missing the conserved

phosphoacceptor histidine (**Fig. S1**). We also included the previously-identified related EL368

(dimer) and EL362 (mixed oligomerization state) histidine kinases from *E. litoralis* HTCC2594

(3, 21), as well as the *Caulobacter crescentus* LovK (dimer) (29) in the alignment; all three of

these proteins grouped in the LOV-HKs cluster. Additionally, our search identified a third PAS-

HK cluster with proteins containing non-LOV PAS domains that shared 17-24% sequence

identity with the EL346 LOV domain. Notably, many of these domains contained a conserved

tryptophan in place of the cysteine utilized in LOV-type photochemistry (28) (**Fig. S1**). These

data show that the LOV/PAS-HK architecture is widespread throughout *Alphaproteobacteria* (as

are HWE/HisKA2 HKs in general (30, 31)) and contains members which could well respond to

stimuli other than blue light.

Figure 1: EL346 homologs form three distinct clusters. a) PAS-HKs (purple), EL346-like LOV HKs (pink), and LOV-HKs (blue). The names of the proteins from each cluster that were selected for this study are shown. All other names are omitted for simplicity and can be found in **Figure S2**. **b)** Schematic structures of monomeric (left) and dimeric (right) LOV-HKs in the dark. Monomeric schematic is based on the structure of EL346 (1) and dimeric is based on the engineered fusion protein YF1 (2).

LOV homologs are light sensing and appear mostly dimeric, but have diverse functional

responses to illumination

We posited that the EL346-like LOV-HK cluster might contain a mix of other monomeric

and dimeric HKs, respectively analogous to EL346 and most natural HKs (as well as the

engineered LOV-HK YF1 (2)), as illustrated in **Fig. 1b**. We selected several HKs from each

cluster and characterized their oligomeric state and function with a variety of biochemical

approaches. We started by determining the ability of the LOV-containing proteins to bind flavin

chromophores and undergo typical LOV photochemistry via UV-visible absorbance

spectroscopy. The absorbance spectra of dark and light state samples of four representative

	Organism	Protein Name	Uniprot ID	Identity to EL346	Oligomeric State
EL346-like LOV-HKs	*Erythrobacter litoralis HTCC2594*	EL346	Q2NB77	N/A	Monomer
	Rubellimicrobium thermophilum DSM 16684	RT349	S9SB69	41.6%	Dimer
	Roseivivax halodurans JCM 10272	RH376	X7EIB8	35.3%	Dimer
LOV-HKs	*Methylorubrum extorquens DM4*	ME354	C7CCX6	27.7%	Dimer
	Erythrobacter litoralis HTCC2594	EL368	Q4TL45	23.7%	Dimer
	Erythrobacter litoralis HTCC2594	EL362	Q2N9L9	22.3%	Mixed oligomers
PAS-HKs	*Rhizobium etli CFM 42*	RE356	Q2KD32	25.3%	Monomer/ Dimer
	Methylocella silvestris BL2	MS367	B8ELH4	26.0%	Monomer/ Dimer
Other	*Bacillus subtilis/ Bradyrhizobium japonicum*	YF1 (engineered YtvA/FixL fusion)	N/A (PDB: 4GCZ)	21.9%	Dimer

Table 1: Oligomeric states of known LOV-HKs and PAS-HKs include mix of monomers and dimers. Soluble HKs shown here are separated by cluster with the same coloring as Fig. 1 (EL346-like LOV HKs – pink, LOV-HKs – blue, PAS-HKs – purple, Other – grey) and include those investigated in this study (EL346, RT349, RH376, ME354, RE356, and MS367) as well as those identified in literature (EL368, EL362, and YF1) (2, 3). Oligomeric states were determined in this study by SEC-MALS.

proteins – EL346, ME354, RT349, and RH376 (see **Table 1** for protein details) – all showed the

characteristic flavin-LOV triplet (~450 nm) in the dark state which disappeared upon illumination

(**Fig. 2a**). Upon incubation in the dark, the 450 nm absorbance returned with first-order

exponential kinetics that varied by protein: the LOV-HKs measured here, such as ME354 (236

min), RT349 (240 min), and RH376 (62 min), had longer reversion constants τ than EL346 (33

min) (**Fig. 2b**). The particularly long reversion time constants of ME354 and RT349 can be

rationalized by the presence of several "slowing" mutations in their LOV domains relative to that

of EL346, specifically at positions 19, 21, 32, and 101 (EL346 numbering), as collated in a

review by Pudasaini and coworkers (**Fig S3**) (23).

We determined the oligomeric state of these LOV-HKs in the presence of ATP (1 mM) in

dark and light conditions using size exclusion chromatography coupled to multi-angle light

scattering (SEC-MALS). We began by confirming that EL346 is monomeric under both

conditions with minimal change in global shape upon illumination as revealed by SEC elution

profiles (**Fig. 2c**). In contrast, the three other LOV-HK proteins were stable dimers, with RT349

exhibiting a light-dependent shift in elution peak position and shape. Further investigation

showed that the RT349 elution profile was also impacted by ATP as well: in the absence of

ATP, we observed two new peaks at even higher elution volumes in dark and light conditions

(**Fig. S4**).

While the oligomeric states of three new LOV-HKs were all similar, analyses of

autophosphorylation activity indicated a diverse activity profile (**Fig. 3**). EL346 and RH376

exhibited the light-dependent increase in activity typically seen in LOV-HKs, albeit to different

extents. In contrast, RT349 displayed an inverted signaling logic, with higher

autophosphorylation rates in the dark state than in the light state. While this phenomenon has

been reported for engineered LOV-HKs such as YF1 variants (2) and the EL346 R175A mutant

(1), it does not seem to have been observed in a naturally-occurring LOV-HK. On the other end

of the spectrum, ME354 exhibited no detectable activity in both the dark and light), suggesting

that even though this protein properly photocycles, its kinase activity is not regulated directly by

light. Overall, our results illustrate that despite adopting similar oligomeric states, the LOV-HKs

studied here exhibit diverse autophosphorylation activities, suggesting varying forms of

regulation and different signaling paradigms.

Figure 2: UV-visible absorption spectra and SEC-MALS reveal diversity of LOV-HK dark state reversion kinetics and oligomerization states. a) UV-visible absorption spectra of EL346, ME354, RT349, and RH376, superimposing dark state scans (black) with scans post-illumination (red). All spectra display the disappearance of the 450 nm triplet upon exposure to light, consistent with LOV photochemistry. Spectra were recorded in the presence of 1 mM ATP. **b)** Dark state reversion kinetics, measured by the return of absorbance at 446 nm post illumination (black), fit to single exponentials (red) reveal an 8-fold range of time constants. **c)** Superdex 200 GL SEC-MALS chromatograms are represented by dRI in arbitrary units and MALS-derived masses are solid lines under the peaks. MWs and elution volumes of six standard proteins are shown in grey in the top panel. MALS traces show that EL346 is monomeric in solution, while ME354, RT349 and RH376 are all predominantly dimeric. A light-dependent shift in elution peak shape and position is seen in RT349.

Figure 3: LOV-HKs display differing activity levels. Autophosphorylation assays of EL346 and RH376 follow the typical trend for LOV-HKs found in nature, with light activation increasing activity levels, while RT349 produces opposite results: it is more active in the dark state than in the light. Each bar indicates the mean +/- one standard deviation (n=3).

Non-LOV PAS-HKs contain mixture of monomers and dimers, with each state exhibiting different activity levels and stabilities

The two PAS-HKs characterized here, from *Rhizobium etli* CFN 42 (RE356) and from *Methylocella silvestris* BL2 (MS367), exhibit 45% sequence identity overall to one another (32% over the PAS domain). In SEC-MALS experiments, both proteins appeared as mixtures of dimer and monomer, as well as an oligomeric or aggregated fraction (**Fig. 4a**). We found that the monomeric and dimeric states of MS367 and RE356 were in slow equilibrium, allowing them to be separately purified and re-injected onto SEC-MALS. We characterized the stability of the two different states by incubating mixed samples at 4°C for varying amounts of time and using size-exclusion chromatography to visualize elution peaks. For both PAS-HKs, we observed that the monomeric state was stable on the order of days, while the dimer slowly oligomerized or aggregated on this timescale (**Fig. 4b**).

Figure 4: The PAS-HKs MS367 and RE356 are monomer/dimer mixtures that display differential autophosphorylation activity and stabilities. a) SEC-MALS measurements of PAS-HKs MS367 and RE356 reveal the presence of both monomeric and dimeric states. Monomers (red traces) and dimers (blue traces) were separated and injected onto a Superdex200 GL SEC. The resulting MALS-derived masses (solid lines under elution peaks) show that both states remain stable upon separation. **b)** Superdex200 GL SEC traces of PAS-HK monomer/dimer mixtures after being stored at 4 °C for 0, 2, and 4 days at ~20 μM concentration, with both proteins exhibiting a similar trend of the monomer remaining stable over time, while the dimer tends to aggregate. **c)** Autophosphorylation assays (plots, above; phosphor images, below) of the monomer (blue traces) and dimer (red traces) of the PAS-HKs show that for MS367, the dimer has a higher rate of activity while for RE356, the monomer has a higher rate.

Though these proteins currently lack a known trigger analogous to light for the LOV-HKs,

we took advantage of the slow equilibrium between oligomeric states to determine whether they

exhibited differing levels of autophosphorylation. We observed that both MS367 and RE356

displayed autophosphorylation activity in their monomeric and dimeric states with the

characteristic plateau indicating an equilibrium between kinase and phosphatase activity (**Fig.**

4c). While the MS367 dimer appears to be more active than the monomer, RE356 produced

opposite results, with the monomer displaying more activity than the dimer. Despite similarity in

primary sequence and oligomeric assembly, the two PAS-HKs studied here have apparently

opposite regulation via dimerization, again illustrating the diversity of function across this family.

Engineering a "monomerized" LOV-HK

Next, we sought to identify the determinants of dimerization in this family of sensor HKs,

recognizing that sequence changes can influence the multimeric arrangement of proteins

belonging to the same family (32, 33). To guide our analysis, we compared monomeric EL346

with selected dimeric HKs from all three clusters using sequence alignment. These pairwise

alignments highlighted four regions of difference (named RH1-4) in the primary sequence

between EL346 and the other HKs; these regions were distributed differently throughout the

three clusters. One protein, the dimeric RH376 from the EL346-like cluster, contained all four

regions of insertions of amino acids between the two proteins (**Figs. 5a and S5**).

We set out to test the hypothesis that one or more of the RH1-4 regions might be

determinants of oligomeric state (**Figs. 5b and 5c**). These insertions range from 5 to 19

residues and are distributed throughout both proteins (**Fig. S5**). The one insertion unique to

EL346, RH4, corresponds almost entirely to the highly dynamic C-terminal residues we

previously deleted without effect on the monomeric state of that protein (19). Thus, we did not

pursue RH4 as a driver of monomer/dimer state and instead focused on the RH1-3 segments

unique to RH376.

Observing that the insert regions were either in previously-reported dimerization

interfaces for LOV domains (RH1, A'α helix) (2, 3, 34) and HKs (RH2, DHp domain) (35, 36), or

large insertions with predicted secondary structure elements (RH3), we hypothesized that these

segments could contribute to the dimerization of RH376 and perhaps other HKs. A homology

model of the RH376 sequence based on the engineered dimeric LOV-HK YF1 (2) illustrates the

potential involvement of these residues in dimerization interfaces (**Fig. 5c**). We thus deleted

each of the RH1, RH2, and RH3 regions from the RH376 sequence both individually and in

combination, expressed these variant proteins, and compared them to RH376 WT in terms of

LOV photocycle, oligomeric state, and autophosphorylation activity.

Figure 5: RH376 mutant deletion regions are designed in predicted dimer interfaces. RH376 deletion mutants RH1, RH2, and RH3 are highlighted in 3 schematics: **a)** Domain architecture with the predicted secondary structure at the location of each deletion. RH1 is located in the A'α helix of the LOV domain, RH2 in the linker between the LOV domain and α1 of the DHp domain, and RH3 is in the ATP lid. **b)** Monomeric homology model of RH376 based on the structure of EL346 (1). **c)** Dimeric homology model of RH376 based on the engineered YF1 sensor kinase (2). All LOV domains are colored blue, DHp domains green, and CA domains red. Mutations are highlighted and circled.

RH1, RH2, and RH3 proteins displayed the characteristic LOV absorbance triplet

centered at 450 nm and exhibited dark state reversion kinetics similar to the WT protein (**Fig.**

S6), suggesting that the deletions did not perturb the folding and function of the LOV domain. In

RH2, we detected a small change in the relative heights of the triplet peaks, indicating that flavin

binding is likely adversely affected (**Fig. S6**) despite the deletion occurring outside the LOV

domain. We determined the oligomeric state of each deletion mutant by SEC-MALS; while none

of the single deletion constructs eluted as a monomer regardless of dark or light conditions,

RH2 had altered measured mass and SEC elution volume which suggested it samples both

dimeric and monomeric conformations (**Fig. 6**).

We suspected that none of these deletions would have a major impact on

autophosphorylation activity of RH376, given their modest effects on photocycle and

oligomerization state. As a baseline, we found that the RH376 WT protein displays robust

autophosphorylation activity in the dark as well as a ~5x increase in initial velocity upon

illumination (**Table S1, Fig. 6**). The RH1 mutant showed autophosphorylation activity similar to

the WT in the light, but greater in the dark, resulting in a much lower degree of light-activation.

Initial velocity in the dark was about 3x the WT, and in the light was slightly lower than WT.

Similarly, RH2 retained some light activation, but showed very little activity in both dark and light

conditions (**Table S1, Fig. 6**). On the other hand, RH3 displayed activity in the dark that was

~4x the WT initial velocity, with a light-dependent decrease in activity (**Fig. 6**). In all, each

mutant had a markedly different effect on autophosphorylation activity despite having minimal

effects on dimerization.

Given that neither RH1 nor RH3 perturbed dimeric state and both displayed activity

levels comparable to WT, yet had divergent effects on light-activation (**Table S1, Fig. 6**), we

explored the effects of combining these deletions ("RH1+3"). While neither the LOV absorbance

spectrum nor photocycle of this protein were affected (**Fig. S6**), SEC-MALS showed a dramatic

change in oligomeric state, as RH1+3 existed mainly as a monomer, though neither deletion

alone showed monomeric character (**Fig. 6**). Additionally, we detected no autophosphorylation

activity in RH1+3 at conditions similar to the other proteins tested (**Table S1**). Combining these

observations with those of RH2 – which lost activity and adopted a monomer/dimer mix – we

suggest that the LOV-HK RH376 both a). links dimerization and activity and b). uses a

distributed set of residues to control oligomeric state.

Figure 6: RH376 mutants RH1, RH2 and RH3 are active dimers, while double mutant RH1+3 is an inactive monomer. Superdex 200 GL SEC-MALS results show that RH376 WT, RH1 and RH3 are clear dimers, while RH1+3 is primarily monomeric. The mass calculated for RH2 falls between a monomer and dimer, suggesting that the protein likely exists as a monomer/dimer mixture. Expected MWs (grey lines) were calculated based on protein sequence. Mutant activity, represented by initial velocity calculated from the linear part of the curve in at least three replicates for each protein/condition (bars represent average), is shown in the bottom left panel. No activity was detected for RH1+3. Light activation can be seen in RH376 WT, RH1 and RH2, but not RH3. Each bar indicates the mean +/- one standard deviation (n=3).

Discussion

Our work further expands the known behavior of sensor HKs to include two proteins in

slow monomer:dimer equilibria along with three obligate dimers, one of which exhibits a slight

conformational change upon activation, and all of which exhibit varying activities. Prior to this

study, the vast majority of characterized sensor HKs were dimers with the EL346 LOV-HK from

E. litoralis HTCC2594 (1, 3) as the only well-characterized, functional protein that we are aware

of as an exclusive monomer. Using sequence homology searches to find homologs of EL346,

we found a large family of related LOV- and PAS-HKs in the *Alphaproteobacteria*. Most of these

proteins were Pfam (20) HisKA_2 or HWE-HK-type histidine kinases, two groups of HKs that

have been less well-characterized structurally to date (1, 13, 15, 37), and we hypothesized that

a subset of these sequences with the highest homology to monomeric EL346 might contain

other monomers.

Contrary to this expectation, we found a diverse range of assemblies, conformational

changes, and autophosphorylation activity among five new sensor HKs characterized here. For

the LOV-HK proteins, we found several dimers, including two members of the EL346-like cluster

of LOV-HKs (RT349 and RH376) that were dimeric in SEC-MALS analyses in both dark and

light conditions. ME354, a newly characterized member of the other LOV-HK cluster is also

clearly dimeric, as are the established EL368 and *C. crescentus* LovK (3, 29), which both also

cluster within that second group.

While these new LOV-HKs are all dimeric, we observed differences in their structural

response to blue light activation as assessed by SEC-MALS: ME354 and RH376 showed

minimal light-dependent change, while RT349 eluted at later volumes in the light, suggesting

that it adopts a more compact structure in the light. Notably, this shift was dependent upon not

only light, but also the presence of ATP (**Fig. S4**). These light-dependent changes are more

substantial than we previously observed by SEC-MALS with the EL346 and EL368 LOV-HKs (1,

3, 19), and are perhaps similar to the kinds of changes seen by small angle X-ray scattering in

in the engineered YF1 protein (2, 38). Further studies are needed to more fully characterize

these motions and establish how they relate to the large conformational changes postulated to

accompany activation from crystal structures of inactive HKs (39, 40). RT349 is also very

distinctive in that it is more active in the dark than in the light **(Fig. 3).** While this behavior has

been observed before in variants of engineered (2) and mutants of natural LOV-HKs (1), we are

not aware of any naturally-occurring LOV-HKs that share this signaling polarity. We anticipate

that such a light-dependent change in net autophosphorylation has contributions from altered

kinase and phosphatase activities (41), and will be the focus of a subsequent investigation.

311   Turning to the two PAS-HKs, both RE356 and MS367 interconvert slowly enough

between dimeric and monomeric states that we could separate them by size exclusion

chromatography. We observed differing activity levels of the various oligomerization states of

these proteins – both monomers are active, with the MS367 dimer being more active than the

monomer, and RE356 displaying opposite results (**Fig. 4**) – implying that shifts in the

monomer:dimer equilibrium might regulate sensor HK enzymatic activity. Such regulation has

been demonstrated or postulated for a range of signaling proteins including certain sensor HKs

(42), receptor tyrosine kinases (43), and photoreceptors (44-47). Dimerization may be

modulated by potential ligand binding in the yet-uncharacterized PAS domains of these HKs, as

in other PAS-domain containing signaling proteins (48), or via other mechanisms. We also

propose that the relative instability of the dimer state, which forms higher-molecular-weight

aggregates over time, may serve as an intrinsic timer specifically on one of the two signaling

species to temporally modify a biological response post-activation.

324   Given the integral role of dimerization in this possible regulation, along with its obvious

necessity in *trans*-phosphorylation within HKs (14), we sought to understand the sequence

determinants of dimerization with an engineering approach. Using RH376 as a model dimer, we

systematically removed combinations of the three insertions it contains compared to the

monomeric EL346 (**Fig. 5, S5**), hypothesizing that one or more of these changes could create a

monomeric LOV-HK. While each deletion mutant was well-folded as evidenced by binding flavin

chromophores and undergoing canonical LOV photochemistry, we found that all three single

deletion mutants (RH1, RH2, RH3) remained largely dimeric despite the removal of 5-30 amino

acid residues apiece. Remarkably, the combination of two of these deletions to generate the

RH1+3 construct monomerized the protein and abolished autophosphorylation activity. From

these studies on RH376, we arrive at two general conclusions: 1) multiple regions determine

oligomeric states in this family of LOV-HKs, and 2) oligomeric state changes have a complex

relationship with autophosphorylation activity and signaling function.

337   The first of these conclusions, the use of distributed interfaces for dimerization, is

consistent with structural studies on the dimeric YF1 engineered sensor kinase (2) showing

intermonomer contacts via residues in both the LOV domain A'α helix and DHp domain, which

are typical dimerization interfaces for LOV photosensors (2, 49) and canonical HisKA histidine

kinases (14, 50) respectively. These correspond to our RH1 and RH2 deletions; neither of these

changes, nor the RH3 deletion in a large loop near the kinase ATP binding site, were sufficient

on its own to generate monomers, but each had markedly different effects on

autophosphorylation. These data suggest different types of dimers may be generated by these

deletions, some of which retain function better than others. Unexpectedly, the RH3 deletion –

which removes a large loop near the ATP binding site – affected dimerization when paired with

RH1. While this site has not been previously described as being involved in dimerization to the

best of our knowledge, we note that ATP itself bridges interactions between multiple domains in

two known HWE/HisKA2 HK crystal structures (1, 37), laying some precedent for interactions at

or near the nucleotide to be involved in dimerization. Clearly, further structural work is needed to

more fully assess how dimerization is encoded among multiple HKs.

352   The diversity of oligomeric states and function in a family of sensor HKs that we uncover

here opens the door to novel regulatory modes within this family, such as control of

dimerization, whether by ligand or inherent thermodynamic stability, or cooperativity. The closely

related bacterial chemoreceptor proteins, for instance, form large arrays in order to increase

signaling output through cooperativity (17). Similar use of monomer/dimer transitions and

evolution of variable quaternary structure is at the heart of acquiring cooperativity and novel

regulatory modes in the classic hemoglobin family of oxygen-binding proteins, which started as

ancestral monomers prior to dimerizing by gene duplication events and the acquisition of a

relatively small number of mutations (33). Our findings suggest that various oligomeric states

within the sensor HK family could allow for different potential for cooperativity, perhaps

depending on the input signal or output pathway, or integrating multiple points of control or

regulation to be added to the canonical signal transduction pathway of two-component systems.

Experimental Procedures

Bioinformatics

Sequences similar to EL346 were identified by a BLAST search with default parameters

against the non-redundant NCBI database, using EL346 full-length protein sequence (Uniprot

ID: Q2NB77) as a query. EL346 and the best 100 hits were aligned using Clustal Omega (51)

and a distance tree was calculated from this alignment using the same software. Graphic

display of this tree was accomplished using iTOL (Interactive Tree of Life, http://itol.embl.de/)

(52). This tree was used to select HKs for cloning from the strains available at DSMZ

(http://www.dsmz.de).

Cloning, protein expression and purification

Five HKs were selected for biochemical characterization, abbreviated RT349, RH376,

ME354, RE356, and MS367 from the combination of their species name and number of amino

acid residues (**Table 1**). DNA encoding these proteins were cloned into a pHis-Gβ1 vector (53),

with constructs verified by DNA sequencing. Proteins were overexpressed in *Escherichia coli*

BL21(DE3) (Stratagene) in LB as previously described (19). Cells were harvested, resuspended

in Buffer A (50 mM Tris pH 8.0, 100 mM NaCl, and 10 mM $MgCl_2$), and lysed by sonication.

Lysates were centrifuged at 48,000 x g and 4 °C for 45 min. Supernatants were loaded into a

Ni^{2+} Sepharose affinity column (GE Healthcare) and the His6-Gβ1 tagged protein was washed

with 4 column volumes of Buffer A supplemented with 15 mM imidazole and eluted with 4

column volumes of Buffer A supplemented with 250 mM imidazole. Eluted proteins were

exchanged into Buffer A by dialysis, then fusion tags were cleaved by incubation with His_6-TEV

protease overnight at 4 °C. The tag-less proteins were separated from the tags and His_6-TEV

protease by Ni^{2+} affinity chromatography and were further purified by size-exclusion

chromatography on a HiLoad 16/600 Superdex 200 column (GE Healthcare) equilibrated with

HSEC buffer containing 50 mM HEPES pH 7.5, 100 mM NaCl, 10 mM $MgCl_2$, and 0.5 mM DTT.

For light-sensitive proteins, all purification steps were performed under dim red light.

Concentrations were determined from the theoretical absorption coefficient, ε_{280} for PAS-HKs,

calculated from the sequence using the ExPASy ProtParam server (54), and $\varepsilon_{446} = 11,800$ M^{-1}

cm^{-1} for flavin-containing proteins.

RH376 deletion mutants were generated as follows: RH2 and RH3 were ordered as

synthetic constructs from Genewiz (South Plainfield, NJ) and inserted into the same pHis-Gβ1

vector as the WT protein. RH1 and RH1+3 mutants were generated by PCR using an N-

terminal primer lacking the sequence to be deleted using WT and RH3 plasmids as templates,

respectively.

Absorbance and dark state reversion kinetics

UV-visible absorbance spectroscopy measurements were acquired using a Varian Cary

50 spectrophotometer at 24 °C with a 1 cm path length quartz cuvette. First, a UV-visible

spectrum in the range 250-550 nm was acquired in light and dark conditions for each LOV-HK.

For dark measurements, all steps were performed under dim red light. For light measurements,

the samples were illuminated for 1 minute using a blue LED panel (225 LED bulbs in 30.5 ×

30.5-cm panel, 13.8 W, 465 nm maximum illumination wavelength; LEDwholesalers [Hayward,

CA]) prior to each scan. To measure dark state recovery, each LOV-HK sample was illuminated

under a blue LED panel for 1 minute and then transferred to the spectrophotometer. A_{446} of the

samples in HSEC buffer with 1 mM ATP was monitored at 5-min intervals for 400-800 min at

24 °C. Time constants τ were determined by fitting measurements to a monoexponential decay

using the following equation:

413             $Abs = A_{inf} - (A_{inf} - A_o)e^{-t/\tau}$, where $A_{inf} = A_{446}$ at infinity, $A_o = A_{446}$ initial, and t = time

414   Raw data were fit and plotted using in RStudio with the nls function; plots were edited in Adobe

415   Illustrator CS6.

417   ***SEC-MALS***

418   All samples and buffers were filtered with a 0.1-µm pore size filter before use. ATP was

added to each HK sample (~15 µM, 400 µL) to a final concentration of 1 mM. Samples were

then injected at a 0.4 mL/min flow rate onto a Superdex 200 GL 10/300 SEC column (GE

Healthcare) using HSEC buffer with 1 mM ATP. For lit experiments, samples were illuminated

with a blue LED panel for 1 min prior to injection and the blue LED panel was faced toward the

column during the run. For dark experiments, all steps were performed under dim red light.

Samples were detected post-elution by the inline miniDAWN TREOS light scattering and Optilab

rEX refractive index detectors (Wyatt Technology [Santa Barbara, CA]). FPLC runs were

performed at 4 °C, while MALS and refractive index measurements were at 25 °C. Data analysis

and molecular weight calculations were performed using the ASTRA V software (Wyatt

Technology). Raw data were plotted using RStudio.

Autophosphorylation assays

Autophosphorylation was measured as described previously (1, 19). A minimum of three

trials were conducted for each protein under each condition. Proteins were added to a reaction

buffer containing 50 mM Tris, 100 mM NaCl, 5 mM $MnCl_2$, and 1 mM DTT with pH 8.2. Protein

concentrations were confirmed by analysis of dilution series on Coomassie-stained SDS-PAGE.

Final reaction concentrations ranged from approximately 10 to 35 micromolar as listed in the

table below:

Protein	Concentration (µM)	Protein	Concentration (µM)
EL346	16.1	RE356 monomer	9.6
ME354	11.6	RE356 dimer	8.5
RT349	28.6	RH1	12.4
RH376 WT	14.3	RH2	30.4
MS367 monomer	10.7	RH3	34.7
MS367 dimer	11.1	RH1+3	9.1

A mixture of unlabeled ATP and 10 µCi [γ-^{32}P] ATP was added to each mixture to initiate

the reaction (final ATP concentration of 200 µM). Aliquots were removed at time points of 0.5, 1,

2, 4, 8, 6, and 32 min and placed into a 4x SDS-gel loading buffer to quench the reaction. For

dark measurements, all steps were performed under dim red light. For light measurements, the

samples were illuminated with a blue LED panel just prior to and throughout the course of the

experiment. Samples were subjected to SDS-PAGE analysis; dried gels were exposed for 1hr -

overnight and bands were visualized by phosphorimaging. Band intensities were measured

using Fiji (55, 56); initial velocity values were calculated in Microsoft Excel from the linear region

of the curve and were plotted in RStudio.

Data availability – All data for this work are contained in this manuscript.

Supporting information – This article contains supporting information, including Figures S1-7,

Table S1, and accompanying literature citations.

Acknowledgements – The authors would like to acknowledge Bright Shi for generating the

RH1+3 construct, as well as Matthew Cleere, Zaynab Jaber, Jaynee Hart, and other Gardner

lab members for fruitful discussions.

Funding and additional information – This work was supported by NIH grants R01 GM106239

(to K.H.G.) and F32 GM1119311 (to I.D.). The content is solely the responsibility of the authors

and does not necessarily represent the official views of the National Institutes of Health.

Conflict of interest – The authors declare that they have no conflicts of interest with the contents

of this article.

Abbreviations –HK, histidine kinase; LOV, Light-Oxygen-Voltage; PAS, Per-ARNT-Sim; TCS,

two-component system; RR, response regulator; DHp, dimerization histidine phosphotransfer;

CA, catalytic ATP-binding; SEC-MALS, size exclusion chromatography multi-angle light

scattering.

References

1.   Rivera-Cancel, G., Ko, W. H., Tomchick, D. R., Correa, F., and Gardner, K. H. (2014)

472   Full-length structure of a monomeric histidine kinase reveals basis for sensory regulation

473   Proc Natl Acad Sci U S A **111**, 17839-17844 10.1073/pnas.1413983111

2.   Diensthuber, R. P., Bommer, M., Gleichmann, T., and Moglich, A. (2013) Full-length

structure of a sensor histidine kinase pinpoints coaxial coiled coils as signal transducers

and modulators Structure **21**, 1127-1136 10.1016/j.str.2013.04.024

3.   Correa, F., Ko, W. H., Ocasio, V., Bogomolni, R. A., and Gardner, K. H. (2013) Blue light

regulated two-component systems: enzymatic and functional analyses of light-oxygen-

voltage (LOV)-histidine kinases and downstream response regulators Biochemistry **52**,

4656-4666 10.1021/bi400617y

4.   Galperin, M. Y. (2018) What bacteria want Environmental microbiology **20**, 4221-4229

10.1111/1462-2920.14398

5.   Mascher, T., Helmann, J. D., and Unden, G. (2006) Stimulus perception in bacterial

signal-transducing histidine kinases Microbiol Mol Biol Rev **70**, 910-938

10.1128/MMBR.00020-06

6.   West, A. H., and Stock, A. M. (2001) Histidine kinases and response regulator proteins

in two-component signaling systems Trends Biochem Sci **26**, 369-376,

https://www.ncbi.nlm.nih.gov/pubmed/11406410

7.   Wang, S. (2012) Bacterial Two-Component Systems: Structures and Signaling

Mechanisms In Protein Phosphorylation in Human Health, Huang C, ed. InTech, Chapter

15,

8.   Mechaly, A. E., Soto Diaz, S., Sassoon, N., Buschiazzo, A., Betton, J. M., and Alzari, P.

M. (2017) Structural Coupling between Autokinase and Phosphotransferase Reactions

in a Bacterial Histidine Kinase Structure **25**, 939-944 e933 10.1016/j.str.2017.04.011

9.   Kenney, L. J., and Anand, G. S. (2020) EnvZ/OmpR Two-Component Signaling: An

Archetype System That Can Function Noncanonically EcoSal Plus **9**,

10.1128/ecosalplus.ESP-0001-2019

10.   Bhate, M. P., Molnar, K. S., Goulian, M., and DeGrado, W. F. (2015) Signal transduction

499   in histidine kinases: insights from new structures Structure **23**, 981-994

500   10.1016/j.str.2015.04.002

501   11.   Szurmant, H., White, R. A., and Hoch, J. A. (2007) Sensor complexes regulating two-

502   component signal transduction Curr Opin Struct Biol **17**, 706-715

503   10.1016/j.sbi.2007.08.019

504   12.   Gushchin, I., and Gordeliy, V. (2018) Transmembrane Signal Transduction in Two-

505   Component Systems: Piston, Scissoring, or Helical Rotation? Bioessays **40**,

506   10.1002/bies.201700197

507   13.   Rinaldi, J., Fernandez, I., Shin, H., Sycz, G., Gunawardana, S., Kumarapperuma, I. *et al.*

508   (2021) Dimer Asymmetry and Light Activation Mechanism in Brucella Blue-Light Sensor

509   Histidine Kinase mBio **12**, 10.1128/mBio.00264-21

510   14.   Ashenberg, O., Keating, A. E., and Laub, M. T. (2013) Helix bundle loops determine

511   whether histidine kinases autophosphorylate in cis or in trans J Mol Biol **425**, 1198-1209

512   10.1016/j.jmb.2013.01.011

513   15.   Herrou, J., Crosson, S., and Fiebig, A. (2017) Structure and function of HWE/HisKA2-

514   family sensor histidine kinases Curr Opin Microbiol **36**, 47-54 10.1016/j.mib.2017.01.008

515   16.   Luo, B. H., Carman, C. V., and Springer, T. A. (2007) Structural basis of integrin

516   regulation and signaling Annual review of immunology **25**, 619-647

517   10.1146/annurev.immunol.25.022106.141618

518   17.   Muok, A. R., Briegel, A., and Crane, B. R. (2020) Regulation of the chemotaxis histidine

519   kinase CheA: A structural perspective Biochimica et biophysica acta Biomembranes

520   **1862**, 183030 10.1016/j.bbamem.2019.183030

521   18.   Kitayama, Y., Nishiwaki-Ohkawa, T., and Kondo, T. (2014) Intersubunit communications

522   within KaiC hexamers contribute the robust rhythmicity of the cyanobacterial circadian

523   clock Microb Cell **1**, 67-69 10.15698/mic2014.01.129

19. Dikiy, I., Edupuganti, U. R., Abzalimov, R. R., Borbat, P. P., Srivastava, M., Freed, J. H.

et al. (2019) Insights into histidine kinase activation mechanisms from the monomeric

blue light sensor EL346 Proc Natl Acad Sci U S A **116**, 4963-4972

10.1073/pnas.1813586116

20. El-Gebali, S., Mistry, J., Bateman, A., Eddy, S. R., Luciani, A., Potter, S. C. *et al.* (2019)

The Pfam protein families database in 2019 Nucleic Acids Res **47**, D427-D432

10.1093/nar/gky995

21. Swartz, T. E., Tseng, T. S., Frederickson, M. A., Paris, G., Comerci, D. J., Rajashekara,

G. *et al.* (2007) Blue-light-activated histidine kinases: two-component sensors in bacteria

Science **317**, 1090-1093 10.1126/science.1144306

22. Crosson, S., Rajagopal, S., and Moffat, K. (2003) The LOV domain family:

photoresponsive signaling modules coupled to diverse output domains Biochemistry **42**,

2-10 10.1021/bi026978l

23. Pudasaini, A., El-Arab, K. K., and Zoltowski, B. D. (2015) LOV-based optogenetic

devices: light-driven modules to impart photoregulated control of cellular signaling Front

Mol Biosci **2**, 18 10.3389/fmolb.2015.00018

24. Gao, R., and Stock, A. M. (2009) Biological insights from structures of two-component

proteins Annu Rev Microbiol **63**, 133-154 10.1146/annurev.micro.091208.073214

25. Tseng, T. S., Frederickson, M. A., Briggs, W. R., and Bogomolni, R. A. (2010) Light-

activated bacterial LOV-domain histidine kinases Methods Enzymol **471**, 125-134

10.1016/S0076-6879(10)71008-9

26. Moglich, A. (2019) Signal transduction in photoreceptor histidine kinases Protein Sci **28**,

1923-1946 10.1002/pro.3705

27. Zoltowski, B. D., and Gardner, K. H. (2011) Tripping the light fantastic: blue-light

photoreceptors as examples of environmentally modulated protein-protein interactions

Biochemistry **50**, 4-16 10.1021/bi101665s

28. Crosson, S., and Moffat, K. (2001) Structure of a flavin-binding plant photoreceptor

domain: insights into light-mediated signal transduction Proc Natl Acad Sci U S A **98**,

2995-3000 10.1073/pnas.051520298

29. Purcell, E. B., McDonald, C. A., Palfey, B. A., and Crosson, S. (2010) An analysis of the

solution structure and signaling mechanism of LovK, a sensor histidine kinase

integrating light and redox signals Biochemistry **49**, 6761-6770 10.1021/bi1006404

30. Karniol, B., and Vierstra, R. D. (2004) The HWE histidine kinases, a new family of

bacterial two-component sensor kinases with potentially diverse roles in environmental

signaling J Bacteriol **186**, 445-453, https://www.ncbi.nlm.nih.gov/pubmed/14702314

31. Staroń, A., and Mascher, T. (2010) General stress response in alpha-proteobacteria:

PhyR and beyond Mol Microbiol **78**, 271-277 10.1111/j.1365-2958.2010.07336.x

32. Perica, T., Chothia, C., and Teichmann, S. A. (2012) Evolution of oligomeric state

through geometric coupling of protein interfaces Proc Natl Acad Sci U S A **109**, 8127-

8132 10.1073/pnas.1120028109

33. Pillai, A. S., Chandler, S. A., Liu, Y., Signore, A. V., Cortez-Romero, C. R., Benesch, J.

L. P. *et al.* (2020) Origin of complexity in haemoglobin evolution Nature **581**, 480-485

10.1038/s41586-020-2292-y

34. Banerjee, A., Herman, E., Kottke, T., and Essen, L. O. (2016) Structure of a Native-like

Aureochrome 1a LOV Domain Dimer from Phaeodactylum tricornutum Structure **24**,

171-178 10.1016/j.str.2015.10.022

35. Qin, L., Dutta, R., Kurokawa, H., Ikura, M., and Inouye, M. (2000) A monomeric histidine

kinase derived from EnvZ, an Escherichia coli osmosensor Mol Microbiol **36**, 24-32,

https://www.ncbi.nlm.nih.gov/pubmed/10760160

36. Tomomori, C., Tanaka, T., Dutta, R., Park, H., Saha, S. K., Zhu, Y. *et al.* (1999) Solution

structure of the homodimeric core domain of Escherichia coli histidine kinase EnvZ

Nature structural biology **6**, 729-734 10.1038/11495

37. Rinaldi, J., Arrar, M., Sycz, G., Cerutti, M. L., Berguer, P. M., Paris, G. *et al.* (2016)

Structural Insights into the HWE Histidine Kinase Family: The Brucella Blue Light-

Activated Histidine Kinase Domain J Mol Biol **428**, 1165-1179

10.1016/j.jmb.2016.01.026

38. Berntsson, O., Diensthuber, R. P., Panman, M. R., Bjorling, A., Hughes, A. J., Henry, L.

et al. (2017) Time-Resolved X-Ray Solution Scattering Reveals the Structural

Photoactivation of a Light-Oxygen-Voltage Photoreceptor Structure **25**, 933-938 e933

10.1016/j.str.2017.04.006

39. Mechaly, A. E., Sassoon, N., Betton, J. M., and Alzari, P. M. (2014) Segmental helical

motions and dynamical asymmetry modulate histidine kinase autophosphorylation PLoS

Biol **12**, e1001776 10.1371/journal.pbio.1001776

40. Casino, P., Miguel-Romero, L., and Marina, A. (2014) Visualizing autophosphorylation in

histidine kinases Nat Commun **5**, 3258 10.1038/ncomms4258

41. Moglich, A., Ayers, R. A., and Moffat, K. (2009) Design and signaling mechanism of

light-regulated histidine kinases J Mol Biol **385**, 1433-1444 10.1016/j.jmb.2008.12.017

42. Lee, J., Tomchick, D. R., Brautigam, C. A., Machius, M., Kort, R., Hellingwerf, K. J. *et al.*

(2008) Changes at the KinA PAS-A dimerization interface influence histidine kinase

function Biochemistry **47**, 4051-4064 10.1021/bi7021156

43. Yarden, Y., and Schlessinger, J. (1987) Epidermal growth factor induces rapid,

reversible aggregation of the purified epidermal growth factor receptor Biochemistry **26**,

1443-1451 10.1021/bi00379a035

44. Liao, X., Liu, W., Yang, H. Q., and Jenkins, G. I. (2020) A dynamic model of UVR8

photoreceptor signalling in UV-B-acclimated Arabidopsis The New phytologist **227**, 857-

866 10.1111/nph.16581

45.   Sang, Y., Li, Q. H., Rubio, V., Zhang, Y. C., Mao, J., Deng, X. W. *et al.* (2005) N-terminal

601   domain-mediated homodimerization is required for photoreceptor activity of Arabidopsis

602   CRYPTOCHROME 1 Plant Cell **17**, 1569-1584 10.1105/tpc.104.029645

603   46.   Zoltowski, B. D., and Crane, B. R. (2008) Light activation of the LOV protein vivid

604   generates a rapidly exchanging dimer Biochemistry **47**, 7012-7019 10.1021/bi8007017

605   47.   Zoltowski, B. D., Motta-Mena, L. B., and Gardner, K. H. (2013) Blue light-induced

606   dimerization of a bacterial LOV-HTH DNA-binding protein Biochemistry **52**, 6653-6661

607   10.1021/bi401040m

608   48.   Scheuermann, T. H., Tomchick, D. R., Machius, M., Guo, Y., Bruick, R. K., and Gardner,

609   K. H. (2009) Artificial ligand binding within the HIF2alpha PAS-B domain of the HIF2

610   transcription factor Proc Natl Acad Sci U S A **106**, 450-455 10.1073/pnas.0808092106

611   49.   Key, J., Hefti, M., Purcell, E. B., and Moffat, K. (2007) Structure of the redox sensor

612   domain of Azotobacter vinelandii NifL at atomic resolution: signaling, dimerization, and

613   mechanism Biochemistry **46**, 3614-3623 10.1021/bi0620407

614   50.   Dutta, R., Qin, L., and Inouye, M. (1999) Histidine kinases: diversity of domain

615   organization Mol Microbiol **34**, 633-640 10.1046/j.1365-2958.1999.01646.x

616   51.   Sievers, F., Wilm, A., Dineen, D., Gibson, T. J., Karplus, K., Li, W. *et al.* (2011) Fast,

617   scalable generation of high-quality protein multiple sequence alignments using Clustal

618   Omega Molecular systems biology **7**, 539 10.1038/msb.2011.75

619   52.   Letunic, I., and Bork, P. (2019) Interactive Tree Of Life (iTOL) v4: recent updates and

620   new developments Nucleic Acids Res **47**, W256-W259 10.1093/nar/gkz239

621   53.   Sheffield, P., Garrard, S., and Derewenda, Z. (1999) Overcoming expression and

622   purification problems of RhoGDI using a family of "parallel" expression vectors Protein

623   Expr Purif **15**, 34-39 10.1006/prep.1998.1003

624   54.   Gasteiger, E., Hoogland, C., Gattiker, A., Duvaud, S. e., Wilkins, M. R., Appel, R. D. *et*

625   *al.* (2005) Protein Identification and Analysis Tools on the ExPASy Server In The

Proteomics Protocols Handbook, Walker JM, ed. Humana Press, Totowa, NJ Chapter

Chapter 52, 571-607

55.   Rueden, C. T., Schindelin, J., Hiner, M. C., DeZonia, B. E., Walter, A. E., Arena, E. T. *et*

al. (2017) ImageJ2: ImageJ for the next generation of scientific image data BMC

bioinformatics **18**, 529 10.1186/s12859-017-1934-z

56.   Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T. *et al.*

(2012) Fiji: an open-source platform for biological-image analysis Nature methods **9**,

676-682 10.1038/nmeth.2019

634