Diversity of function and higher-order structure within HWE sensor histidine kinases

Igor Dikiy[1], Danielle Swingle[1,2], Kaitlyn Toy[1,3], Uthama R. Edupuganti[1,2], Giomar Rivera-Cancel[4], Kevin H. Gardner[1,3,5*]

[1]: Structural Biology Initiative, CUNY Advanced Science Research Center, New York, NY 10031

[2]: Ph.D. Program in Biochemistry, The Graduate Center – City University of New York, New York, NY 10016

[3]: Department of Chemistry and Biochemistry, City College of New York, New York, NY 10031

[4]: Department of Internal Medicine, UT Southwestern Medical Center, Dallas, TX 75390

[5]: Ph.D. Programs in Biochemistry, Biology, and Chemistry, The Graduate Center – City University of New York, New York, NY 10016

*: direct correspondence to kgardner@gc.cuny.edu

ORCIDs:

Igor Dikiy: 0000-0002-8155-7788

Danielle Swingle: 0000-0003-3691-4560

Kaitlyn Toy: 0000-0003-2595-8779

Uthama R. Edupuganti: 0000-0003-0571-4101

Giomar Rivera-Cancel: 0000-0002-4909-4103

Kevin H. Gardner: 0000-0002-8671-2556

Keywords: histidine kinase, bacterial signal transduction, conformational change, structure-function, biophysics, environmental sensing, two-component system, HWE family, HisKA2 family

Running title: Diversity of function and structure in HWE family sensor HKs

Abstract

Integral to the protein structure/function paradigm, oligomeric state is typically conserved

along with function across evolution. However, notable exceptions such as the hemoglobins

show how evolution can alter oligomerization to enable new regulatory mechanisms. Here we

examine this linkage in histidine kinases (HKs), a large class of widely distributed prokaryotic

environmental sensors. While the majority of HKs are transmembrane homodimers,

members of the HWE/HisKA2 family can deviate from this architecture as exemplified by our

finding of a monomeric soluble HWE/HisKA2 HK (EL346, a photosensing Light-Oxygen-Voltage

(LOV)-HK). To further explore the diversity of oligomerization states and regulation within this

family, we biophysically and biochemically characterized multiple EL346 homologs and found a

range of HK oligomeric states and functions. Three LOV-HK homologs are primarily dimeric with

differing structural and functional responses to light, while two Per-ARNT-Sim (PAS)-HKs

interconvert between differentially active monomers and dimers, suggesting dimerization might

control enzymatic activity for these proteins. Finally, we examined putative interfaces in a

dimeric LOV-HK, finding that multiple regions contribute to dimerization. Our findings suggest

the potential for novel regulatory modes and oligomeric states beyond those traditionally defined

for this important family of environmental sensors.

Introduction

Sensor histidine kinases (HKs) are signal transduction receptors that are widespread in

prokaryotes, enabling these organisms to sense and respond to diverse environmental stimuli

(3, 4). HKs are typically found in two-component systems (TCS), which are most simply

composed of a sensor HK and a downstream response regulator (RR) co-located in the same

operon. The RR responds to a signal from the HK by acting as a phosphorylation-dependent

transcription regulator, with activation often triggering RR DNA binding and altering the

expression of nearby genes (5-8).

The prototypical sensor HK contains N-terminal sensor and C-terminal kinase domains,

the latter of which can be subdivided into a two-helix dimerization histidine phosphotransfer

(DHp) domain and a catalytic ATP-binding (CA) domain (9). The detection of a stimulus by the

sensor domain (or in some cases, elsewhere (8)) modulates the CA domain

autophosphorylation of a conserved His residue in the DHp domain; the phosphate group in this

phospho-His adduct is subsequently transferred to a conserved Asp in a downstream RR (5).

Most characterized sensor HKs are obligate homodimeric transmembrane receptors,

facilitating the transmission of signals across the membrane from periplasmic sensor domains to

cytoplasmic kinase domains via symmetry-breaking conformational changes (9-12). The

regulation of typical sensor HKs is largely believed to require a dimeric complex, both for the

structural changes between the "off" and "on" state as well as determining whether

autophosphorylation occurs between protomers (*in trans*) or within a protomer (*in cis*) (9, 13).

Dimerization is mediated via the DHp domains of two monomers, forming a four-helix bundle

and facilitating pivot-, piston-, or scissoring-type movements as part of the activation process (9,

14), echoing similar themes in integrins (15) and certain other transmembrane receptors. In

contrast, some sensors adopt different oligomeric states, such as the large multimeric

assemblies of chemotaxis HKs (16), hexameric KaiC circadian HKs (17) or the monomeric

photosensory EL346 HK (1, 18).

The traditional view of sensor HK structure has been built up from extensive structural

and biochemical studies (9) that have usually focused on HKs in only one of several subfamilies

of histidine kinases (the HisKA family within Pfam (19)). In contrast, the related HWE (His-Trp-

Glu)-HK and HisKA_2 families (referred to here as the HWE/HisKA2 family) of HKs have been

less studied from a structural perspective (14). Members of this group, which contain specific

sequence variations in the DHp and CA domains compared to members of the canonical HisKA

family, tend to signal to a more diverse set of output proteins which often lack DNA-binding

output domains (14). The HWE/HisKA2 family is also enriched in HKs lacking transmembrane

segments, likely localizing them to the cytosol. Notably, soluble sensor HKs are relieved of the

constraints of transmitting a signal across a lipid bilayer, which typically involves signal

transmission via motions of one protomer with respect to the other (9, 14). This raises several

questions: Are such soluble HKs able to adopt diverse oligomeric states, including non-dimeric

architectures? And could these varied architectures be equivalently varied in their regulation?

We view the relatively understudied HWE/HisKA2 family of HKs (14) as a suitable testbed for

these questions.

Our prior work on the HWE/HisKA2 family focused on three soluble HKs involved in the

general stress response of the marine alphaproteobacterium *Erythrobacter litoralis* HTCC2594

(EL346, EL362, and EL368) (1, 18, 20, 21). Each of these proteins senses blue light through a

light oxygen voltage (LOV) sensor domain (22, 23), a sub-class of the versatile Per-ARNT-Sim

(PAS) domains that are present in over 30% of HKs (4, 24). The response to blue light provides

a readily tractable way to explore the inactive and active states of histidine kinases, typically

corresponding to dark and light states of LOV-HKs (21, 25, 26). These states can be

conveniently distinguished spectrophotometrically: while the dark state spectra of LOV-HKs

exhibit a triplet peak at around 450 nm, characteristic of protein-bound flavin chromophores, the

light state spectrum has a broad, singlet absorption peak at 390 nm (27). This spectroscopic

change is due to the photochemically-triggered reduction of the flavin chromophore and

concomitant formation of a cysteine-C(4a) covalent adduct, a bond that is thermally reversible in

the dark on a timescale varying from seconds to hours among LOV domains (23).

While our prior studies indicate that EL368 is dimeric, EL346 is a soluble and functional

monomer (1, 20). The EL346 crystal structure reveals that it is held in a monomeric state by an

intramolecular association between its DHp-like domain and the LOV sensory domain, blocking

the face typically used for DHp domains to dimerize (1). Despite this deviation from current HK

signaling models, the monomeric EL346 undergoes light-induced increases in

autophosphorylation rate (1) as well as conformational changes involving the sensor, DHp, and

kinase domains (18). More broadly, the active monomer of EL346 suggests that soluble HKs

may not require dimerization as a key element in signal transduction, unlike conventional

transmembrane HKs.

To address this general issue in HK signaling, we set out to characterize the oligomeric

and functional states of several novel HWE/HisKA2 HKs. To do so, we used sequence similarity

searches to identify multiple soluble LOV- and PAS-HKs closely related to EL346, selecting five

proteins for in-depth analyses. We showed that three of these five are novel dimeric LOV-HKs,

each of which properly photocycles by UV-visible absorbance spectroscopy. Notably, these

LOV-HKs differed in the effect of light on their autophosphorylation activity, with a mix of dark-

and light-activated proteins among them. In addition, we also characterized two light-insensitive

PAS-HKs which equilibrate between monomeric and dimeric states with differing activities,

suggesting dimerization as a possible regulatory switch in this family of HKs. Finally, we

combined structural and sequence information to engineer a "monomerized dimer" by deleting

putative dimerization interfaces in a dimeric LOV-HK, finding that broadly distributed residues

contribute to dimerization. Taken together, our data reveal a wider range of oligomerization

states and functional activity in HKs than traditionally described.

Results

Bioinformatics analysis reveals numerous alphaproteobacterial EL346 homologs

To identify other potential monomeric histidine kinases, we searched for enzymes highly

similar in domain organization to the well-validated and functional EL346 monomer. In addition

to EL346, the top 100 BLASTp hits defined a set of proteins from *Alphaproteobacteria* which all

contained the same domain architecture of a single N-terminal PAS or LOV domain followed by

a C-terminal HWE/HisKA2 histidine kinase. None of these sequences contained any predicted

transmembrane segments, strongly suggesting all these candidates are soluble. A multiple

sequence alignment of these protein sequences and distance tree analysis revealed three

clusters (**Fig. 1a**), which split on increasingly diverse sensor types: EL346-like LOV-HKs, LOV-

HKs, and PAS-HKs (non-LOV). LOV domains were provisionally identified by the presence of

the characteristic "NCRFLQ" sequence and high homology (> 30% identity) to the EL346 LOV

domain (28), while the non-LOV PAS domains lacked this sequence motif (highlighted in **Fig.**

S1). Interestingly, a subgroup of the EL346-like LOV-HKs is missing the conserved

phosphoacceptor histidine (**Fig. S1**). We also included the previously-identified related EL368

(dimer) and EL362 (mixed oligomerization state) histidine kinases from *E. litoralis* HTCC2594

(20, 21), as well as the *Caulobacter crescentus* LovK (dimer) (29) in the alignment; all three of

these proteins grouped in the LOV-HKs cluster. Additionally, our search identified a third PAS-

HK cluster with proteins containing non-LOV PAS domains that shared 17-24% sequence

identity with the EL346 LOV domain. Notably, many of these domains contained a conserved

tryptophan in place of the cysteine utilized in LOV-type photochemistry (28) (**Fig. S1**). These

data show that the LOV/PAS-HK architecture is widespread throughout *Alphaproteobacteria* (as

are HWE/HisKA2 HKs in general (30, 31)) and contains members which could well respond to

stimuli other than blue light.

Figure 1: EL346 homologs form three distinct clusters. a) PAS-HKs (purple), EL346-like LOV
HKs (pink), and LOV-HKs (blue). The names of the proteins from each cluster that were selected for
this study are shown. All other names are omitted for simplicity and can be found in **Figure S2**.
b) Schematic structures of monomeric (left) and dimeric (right) LOV-HKs in the dark state. Monomeric
schematic is based on the structure of EL346 (1) and dimeric is based on the engineered fusion
protein YF1 (2).

LOV homologs are light sensing and appear mostly dimeric, but have diverse functional

responses to illumination

We posited that the EL346-like LOV-HK cluster might contain a mix of other monomeric

and dimeric HKs, respectively analogous to EL346 and most natural HKs (as well as the

engineered LOV-HK YF1 (2)), as illustrated in **Fig. 1b**. We selected several HKs from each

cluster and characterized their oligomeric state and function with a variety of biochemical

approaches. We started by determining the ability of the LOV-containing proteins to bind flavin

chromophores and undergo typical LOV photochemistry via UV-visible absorbance

spectroscopy. The absorbance spectra of dark and light state samples of four representative

proteins – EL346, ME354, RT349, and RH376 (see **Table 1** for protein details) – all showed the

	Organism	Protein Name	Uniprot ID	% ID EL346	Condition	MW avg ± sd [max, min] (kDa)	Expected MW (kDa)	Oligomeric State
EL346-like LOV-HKs	*Erythrobacter litoralis* HTCC 2594	EL346	Q2NB77	100%	D +ATP	36.7 ± 3.0 [29.7, 39.3]	38.6/77.2	Monomer
					L +ATP	37.0 ± 1.9 [33.5, 38.8]		Monomer
	Rubellimicrobium thermophilum DSM 16684	RT349	S9SB69	41.6%	D +ATP	68.5 ± 1.5 [64.8, 70.6]	37.6/75.1	Mostly Dimer
					L +ATP	67.3 ± 1.4 [64.2, 69.1]		Mostly Dimer
	Roseivivax halodurans JCM 10272	RH376	X7EIB8	35.3%	D +ATP	81.5 ± 1.1 [78.7, 82.6]	42.5/85.0	Dimer
					L +ATP	80.8 ± 1.7 [76.4, 82.8]		Dimer
		RH1	N/A	N/A	D +ATP	86.6 ± 1.5 [84.3, 90.2]	42.1/84.1	Dimer
					L +ATP	80.8 ± 0.4 [79.8, 81.7]		Dimer
		RH2			D +ATP	73.2 ± 0.3 [72.1, 73.6]	41.8/83.5	Mostly Dimer
					L +ATP	74.9 ± 0.8 [73.5, 76.3]		Mostly Dimer
		RH3			D +ATP	76.4 ± 1.1 [74.8, 81.3]	39.5/78.9	Dimer
					L +ATP	76.5 ± 1.2 [74.9, 81.7]		Dimer
		RH1+3			D +ATP	41.7 ± 1.8 [38.4, 46.8]	39.0/78.1	Mostly Monomer
					L +ATP	48.1 ± 2.3 [44.2, 51.9]		Mostly Monomer
LOV-HKs	*Methylorubrum extorquens* DM4	ME354	C7CCX6	27.7%	D +ATP	71.7 ± 3.1 [65.4, 74.8]	38.8/77.7	Mostly Dimer
					L +ATP	72.3 ± 4.0 [62.8, 75.6]		Mostly Dimer
	Erythrobacter litoralis HTCC2594	EL368	Q4TL45	23.7%	N/A	N/A	N/A	N/A
	Erythrobacter litoralis HTCC2594	EL362	Q2N9L9	22.3%	N/A	N/A	N/A	N/A
PAS-HKs	*Rhizobium etli* CFM 42	RE356	Q2KD32	25.3%	Monomer	37.8 ± 0.7 [36.6, 38.6]	39.9/79.7	Monomer
					Dimer	87.4 ± 1.6 [84.6, 89.8]		Dimer
	Methylocella silvestris BL2	MS367	B8ELH4	26.0%	Monomer	38.3 ± 3.6 [32.7, 43.0]	40.2/80.4	Monomer
					Dimer	82.7 ± 6.4 [71.9, 96.7]		Dimer
Other	*Bacillus subtilis/ Bradyrhizobium japonicum*	YF1 (YtvA/FixL fusion)	N/A (PDB:4GCZ)	21.9%	N/A	N/A	N/A	Dimer

159
Table 1: Oligomeric states of known LOV-HKs and PAS-HKs include mix of monomers and dimers.
Soluble HKs shown here are separated by cluster with the same coloring as Fig. 1 (EL346-like LOV HKs
– pink, LOV-HKs – blue, PAS-HKs – purple, Other – grey) and include those investigated in this study
(EL346, RT349, RH376, ME354, RE356, and MS367) as well as those identified in literature (EL368,
EL362, and YF1) (2, 20). Mass data were collected using Superdex200 GL SEC-MALS (n=1). Values
shown are MALS MW (average mass), minimum, maximum, and standard deviation among the points
that make up the mass line for each measurement, and oligomeric states were defined using a qualitative
five-bin scale.
168

characteristic flavin-LOV triplet (~450 nm) in the dark state which disappeared upon illumination

(**Fig. 2a**). Upon incubation in the dark, the 450 nm absorbance returned with first-order

exponential kinetics that varied by protein: the LOV-HKs measured here, such as ME354 (236

min), RT349 (240 min), and RH376 (62 min), had longer reversion constants τ than EL346 (33

min) (**Fig. 2b**). The particularly long reversion time constants of ME354 and RT349 can be

rationalized by the presence of several "slowing" mutations in their LOV domains relative to that

of EL346, specifically at positions 19, 21, 32, and 101 (EL346 numbering), as collated in a

review by Pudasaini and coworkers (**Fig S3**) (23).

We determined the oligomeric state of these LOV-HKs in the presence of ATP (1 mM) in

dark and light conditions using size exclusion chromatography coupled to multi-angle light

scattering (SEC-MALS). We began by confirming that EL346 is monomeric under both

conditions with minimal change in global shape upon illumination as revealed by SEC elution

profiles. In contrast, the three other LOV-HK proteins were dimeric or mostly dimeric, with

RT349 exhibiting a light-dependent shift in elution peak position and shape (**Fig. 2c, Table 1**).

Further investigation showed that the RT349 elution profile was also impacted by ATP as well:

in the absence of ATP, we observed two new peaks at even higher elution volumes in dark and

light conditions (**Fig. S4**).

While the oligomeric states of the three new LOV-HKs were all similar, analyses of

autophosphorylation activity indicated a diverse activity profile (**Fig. 3**). EL346 and RH376

exhibited the light-dependent increase in activity typically seen in LOV-HKs, albeit to different

extents. In contrast, RT349 displayed an inverted signaling logic, with higher

autophosphorylation rates in the dark state than in the light state. While this phenomenon has

been reported for engineered LOV-HKs such as YF1 variants (2) and the EL346 R175A mutant

(1), it does not seem to have been observed in a naturally-occurring LOV-HK. On the other end

of the spectrum, ME354 exhibited no detectable activity in both the dark and light, suggesting

that even though this protein properly photocycles, its kinase activity is not regulated directly by

light under these conditions. Overall, our results illustrate that despite adopting similar

oligomeric states, the LOV-HKs studied here exhibit diverse autophosphorylation activities,

suggesting varying forms of regulation and different signaling paradigms.

Figure 2: UV-visible absorption spectra and SEC-MALS reveal diversity of LOV-HK dark state reversion kinetics and oligomerization states. a) UV-visible absorption spectra of EL346, ME354, RT349, and RH376, superimposing dark state scans (black) with scans post-illumination (red). All spectra display the disappearance of the 450 nm triplet upon exposure to light, consistent with LOV photochemistry. Spectra were recorded in the presence of 1 mM ATP. **b)** Dark state reversion kinetics, measured by the return of absorbance at 446 nm post illumination (black), fit to single exponentials (red) reveal an 8-fold range of time constants. **c)** Superdex 200 GL SEC-MALS chromatograms are represented by dRI in arbitrary units and MALS-derived masses are solid lines under the peaks (n=1). MWs and elution volumes of six standard proteins are shown in grey in the top panel. MALS traces show that EL346 is monomeric in solution, while ME354, RT349 and RH376 are all predominantly dimeric (**Table 1**). A light-dependent shift in elution peak shape and position is seen in RT349.

Figure 3: LOV-HKs display differing activity levels. Autophosphorylation assays (plots, above; phosphor images, below) of EL346 and RH376 follow the typical trend for LOV-HKs found in nature, with light activation increasing activity levels, while RT349 produces opposite results: it is more active in the dark state than in the light. Each bar indicates the mean +/- one standard deviation (n=3).

Non-LOV PAS-HKs contain mixture of monomers and dimers, with each state exhibiting different activity levels and stabilities

The two PAS-HKs characterized here, from *Rhizobium etli* CFN 42 (RE356) and from *Methylocella silvestris* BL2 (MS367), exhibit 45% sequence identity overall to one another (32% over the PAS domain). In SEC-MALS experiments, both proteins appeared as mixtures of dimer and monomer, as well as an oligomeric or aggregated fraction. We found that the monomeric and dimeric states of MS367 and RE356 were in slow equilibrium, allowing them to be separately purified and re-injected onto SEC-MALS (**Fig. 4a**). We characterized the *in vitro* stability of the two different states by incubating mixed samples at 4°C for varying amounts of time and using size-exclusion chromatography to visualize elution peaks. For both PAS-HKs, we observed that the monomeric state was stable on the order of days, while the dimer slowly oligomerized or aggregated on this timescale (**Fig. 4b**).

Figure 4: The PAS-HKs MS367 and RE356 are monomer/dimer mixtures that display differential autophosphorylation activity and stabilities. a) SEC-MALS measurements of PAS-HKs MS367 and RE356 reveal the presence of both monomeric and dimeric states. Monomers (red traces) and dimers (blue traces) were separated and injected onto a Superdex200 GL SEC (n=1). The resulting MALS-derived masses (solid lines under elution peaks) show that, under these conditions, both states remain stable upon separation. **b)** Superdex200 GL SEC traces of PAS-HK monomer/dimer mixtures after being stored at 4 °C for 0, 2, and 4 days at ~20 µM concentration, with both proteins exhibiting a similar trend of the monomer remaining stable over time, while the dimer tends to aggregate. **c)** Autophosphorylation assays (n=3) (plots, above; phosphor images, below) of the monomer (blue traces) and dimer (red traces) of the PAS-HKs show that for MS367, the dimer has a higher rate of activity while for RE356, the monomer has a higher rate.

Though these proteins currently lack a known trigger analogous to light for the LOV-HKs, we took advantage of the slow equilibrium between oligomeric states to determine whether they exhibited differing levels of autophosphorylation. We observed that both MS367 and RE356 displayed autophosphorylation activity in their monomeric and dimeric states with the characteristic plateau indicating an equilibrium between kinase and phosphatase activity (**Fig. 4c**). While the MS367 dimer appears to be more active than the monomer, RE356 produced opposite results, with the monomer displaying more activity than the dimer. Despite similarity in primary sequence and oligomeric assembly, the two PAS-HKs studied here in an *in vitro* context have apparently opposite regulation via dimerization, again illustrating the diversity of function across this family.

Engineering a "monomerized" LOV-HK

Next, we sought to identify the determinants of dimerization in this family of sensor HKs,

recognizing that sequence changes can influence the multimeric arrangement of proteins

belonging to the same family (32, 33). To guide our analysis, we aligned the sequence of the

monomeric EL346 with those of selected dimeric HKs from all three clusters. These pairwise

alignments suggested four candidate regions that might influence dimerization; three regions

missing from EL346, but found in many of the other HKs (named RH1-3), as well a unique C-

terminal segment in EL346 (RH4) (**Figs. 5a and S5**).

To test the hypothesis that one or more of the RH1-4 regions might be determinants of

oligomeric state, we systematically deleted one or more of these regions from the dimeric

RH376 protein, which contained all four segments, in the hope of finding a constitutively

monomeric variant (**Figs. 5b and 5c**). As the RH4 segment corresponds almost entirely to the

highly dynamic C-terminal residues we previously deleted without effect on the monomeric state

of EL346 (18), we did not investigate RH4 as a driver of monomer/dimer state and instead

focused on the RH1-3 segments unique to RH376.

Observing that the insert regions were either in previously-reported dimerization

interfaces for LOV domains (RH1, A'α helix) (2, 20, 34) and HKs (RH2, DHp domain) (35, 36),

or large insertions with predicted secondary structure elements (RH3, ATP lid), we hypothesized

that these segments could contribute to the dimerization of RH376 and perhaps other HKs. A

RH376 homology model (37) based on the engineered dimeric LOV-HK YF1 protein (2)

illustrates the potential involvement of these residues in dimerization interfaces (**Fig. 5c**). We

thus deleted each of the RH1, RH2, and RH3 regions from the RH376 sequence both

individually and in combination, expressed these variant proteins, and compared them to RH376

WT in terms of LOV photocycle, oligomeric state, and autophosphorylation activity.

Figure 5: RH376 mutant deletion regions are designed in predicted dimer interfaces. RH376 deletion mutants RH1, RH2, and RH3 are highlighted in 3 schematics: **a)** Domain architecture with the predicted secondary structure at the location of each deletion. RH1 is located in the A'α helix of the LOV domain, RH2 in the linker between the LOV domain and α1 of the DHp domain, and RH3 is in the ATP lid. **b)** Monomeric homology model of RH376 based on the structure of EL346 (1). **c)** Dimeric homology model of RH376 based on the engineered YF1 sensor kinase (2). All LOV domains are colored blue, DHp domains green, and CA domains red. Mutations are highlighted and circled.

RH1, RH2, and RH3 proteins displayed the characteristic LOV absorbance triplet
centered at 450 nm and exhibited dark state reversion kinetics similar to the WT protein (**Fig.**
S6), suggesting that the deletions did not perturb the folding and function of the LOV domain. In
RH2, we detected a small change in the relative heights of the triplet peaks, indicating that flavin
binding is likely adversely affected (**Fig. S6**) despite the deletion occurring outside the LOV
domain. We determined the oligomeric state of each deletion mutant by SEC-MALS; while none
of the single deletion constructs eluted as a monomer regardless of dark or light conditions,
RH2 had an altered measured mass and SEC elution volume which suggest it samples both
dimeric and monomeric conformations (**Table 1, Fig. 6**).
We suspected that none of these deletions would have a major impact on
autophosphorylation activity of RH376, given their modest effects on photocycle and
oligomerization state. As a baseline, we found that the RH376 WT protein displays robust
autophosphorylation activity in the dark as well as a ~5x increase in initial velocity upon
illumination (**Table S1, Fig. 6**). The RH1 mutant showed autophosphorylation activity similar to
the WT in the light, but greater in the dark, resulting in a much lower degree of light-activation.
Similarly, RH2 retained some light activation, but showed very little activity in both dark and light
conditions (**Table S1, Fig. 6**). On the other hand, RH3 displayed activity in the dark that was
~4x the WT initial velocity, with a light-dependent decrease in activity (**Table S1, Fig. 6**). In all,
each mutant had a markedly different effect on autophosphorylation activity despite having
minimal effects on dimerization.
Given that neither RH1 nor RH3 perturbed dimeric state and both displayed activity
levels comparable to WT, yet had divergent effects on light-activation (**Table S1, Fig. 6**), we
explored the effects of combining these deletions ("RH1+3"). While neither the LOV absorbance
spectrum nor photocycle of this protein were affected (**Fig. S6**), SEC-MALS showed a dramatic
change in oligomeric state, as RH1+3 existed mainly as a monomer, though neither deletion
alone showed monomeric character (**Table S1, Fig. 6**). Additionally, we detected no
autophosphorylation activity in RH1+3 at conditions similar to the other proteins tested. Overall,
we conclude that RH376 uses a distributed set of residues to control oligomeric state.
279

Figure 6: RH376 mutants RH1, RH2 and RH3 are active and mostly dimeric, while double mutant RH1+3 is inactive and primarily monomeric. Superdex 200 GL SEC-MALS results (n=1) show that RH376 WT, RH1 and RH3 are clear dimers, while RH1+3 is primarily monomeric, and RH2 is primarily dimeric. Expected MWs (grey lines) were calculated based on protein sequence. Mutant activity, represented by initial velocity calculated from the linear part of the curve in at least three replicates for each protein/condition (bars represent average), is shown in the bottom left panel. No activity was detected for RH1+3. Light activation can be seen in RH376 WT, RH1 and RH2, but not RH3. Each bar indicates the mean +/- one standard deviation (n=3).

Discussion

Our work further expands the known behavior of sensor HKs to include two proteins in

slow monomer:dimer equilibria along with three chiefly dimeric proteins, one of which exhibits a

slight conformational change upon activation, and all of which exhibit varying activities. Prior to

this study, the vast majority of characterized sensor HKs were dimers with the EL346 LOV-HK

from *E. litoralis* HTCC2594 (1, 20) as the only well-characterized, functional protein that we are

aware of as an exclusive monomer. Using sequence homology searches to find homologs of

EL346, we found a large family of related LOV- and PAS-HKs in the *Alphaproteobacteria*. Most

of these proteins were Pfam (19) HisKA_2 or HWE-HK-type histidine kinases, two groups of

HKs that have been less well-characterized structurally to date (1, 12, 14, 38), and we

hypothesized that a subset of these sequences with the highest homology to monomeric EL346

might contain other monomers.

Contrary to this expectation, we found a diverse range of assemblies, conformational

changes, and autophosphorylation activity among the five new sensor HKs characterized here.

For the LOV-HK proteins, we found several dimers, including two members of the EL346-like

cluster of LOV-HKs (RT349 and RH376) which SEC-MALS analyses showed to be dimeric or

primarily dimeric in both dark and light conditions. ME354, a newly characterized member of the

other LOV-HK cluster is also mostly dimeric, similar to the established EL368 and *C. crescentus*

LovK (20, 29), which both also cluster within that second group.

While these new LOV-HKs are all mostly dimeric, we observed differences in their

structural response to blue light activation as assessed by SEC-MALS: ME354 and RH376

showed minimal light-dependent change, while RT349 eluted at later volumes in the light,

suggesting that it adopts a more compact structure under this condition. Notably, this shift was

dependent upon not only light, but also the presence of ATP (**Fig. S4**). These light-dependent

changes are more substantial than we previously observed by SEC-MALS with the EL346 and

EL368 LOV-HKs (1, 18, 20), and are perhaps similar to the kinds of changes seen by small

angle solution X-ray scattering in in the engineered YF1 protein (2, 39). Further studies are

needed to more fully characterize these motions and establish how they relate to the large

conformational changes postulated to accompany activation from crystal structures of inactive

HKs (40, 41). RT349 is also very distinctive in that it is more active in the dark than in the light

(Fig. 3). While this behavior has been observed before in variants of engineered (2) and

mutants of natural LOV-HKs (1), we are not aware of any naturally-occurring LOV-HKs that

share this signaling polarity. We anticipate that such a light-dependent change in net

autophosphorylation has contributions from altered kinase and phosphatase activities (42), and

this will be the focus of a subsequent investigation.

Turning to the two PAS-HKs, both RE356 and MS367 interconvert slowly enough

between dimeric and monomeric states that we could separate them by size exclusion

chromatography. We observed differing activity levels of the various oligomerization states of

these proteins – both monomers are active, with the MS367 dimer being more active than the

monomer, and RE356 displaying opposite results (**Fig. 4**). The *in vitro* data lead us to propose a

mechanism where shifts in the monomer:dimer equilibrium could regulate sensor HK enzymatic

activity. While demonstrating that monomer:dimer control of RE356 and MS367 activity *in vivo* is

outside the scope of this work, we note that comparable mechanisms are widely accepted or

suggested for a very broad range of signaling proteins including certain sensor HKs (43),

receptor tyrosine kinases (44), and photoreceptors (45-48). In this case, dimerization may be

modulated by potential ligand binding in the yet-uncharacterized PAS domains of these HKs, as

in other PAS-domain containing signaling proteins (49), or via other mechanisms. Perhaps the

relative instability of the dimer state, which forms higher-molecular-weight aggregates over time,

could serve as an intrinsic timer specifically on one of the two signaling species to temporally

modify a biological response post-activation.

Given the integral role of dimerization for many sensor HKs, we sought to understand

the sequence determinants of dimerization with an engineering approach. Using RH376 as a

model dimer, we systematically removed combinations of the three insertions it contains

compared to the monomeric EL346 (**Fig. 5, S5**), hypothesizing that one or more of these

changes could create a monomeric LOV-HK. While each deletion mutant was well-folded as

evidenced by binding flavin chromophores and undergoing canonical LOV photochemistry, we

found that all three single deletion mutants (RH1, RH2, RH3) remained largely dimeric despite

the removal of 5-30 amino acid residues apiece. Remarkably, the combination of two of these

deletions to generate the RH1+3 construct substantially monomerized the protein and abolished

autophosphorylation activity.

From these studies on RH376, we arrive at the general conclusion that multiple regions

determine oligomeric states in this family of LOV-HKs. The use of distributed interfaces for

dimerization is consistent with structural studies on the dimeric YF1 engineered sensor kinase

(2) showing intermonomer contacts via both the LOV and DHp domains, which are typical

dimerization interfaces for LOV photosensors (2, 50) and canonical HisKA histidine kinases (13,

51) respectively. These correspond to our RH1 and RH2 deletions; neither of these changes,

nor the RH3 deletion in a large loop near the kinase ATP binding site, were sufficient on its own

to generate monomers. Unexpectedly, the RH3 deletion – which removes a large loop near the

ATP binding site – affected dimerization when paired with RH1. While this site has not been

previously described as being involved in dimerization to the best of our knowledge, we note

that ATP itself bridges interactions between multiple domains in two known HWE/HisKA2 HK

crystal structures (1, 38), laying some precedent for interactions at or near the nucleotide to be

involved in dimerization. Clearly, further structural work is needed to more fully assess how

dimerization is encoded among multiple HKs.

The diversity of oligomeric states and function in a family of sensor HKs that we

uncovered here opens the door to novel regulatory modes within this family, such as control of

dimerization, whether by ligand or inherent thermodynamic stability, or cooperativity. The closely

related bacterial chemoreceptor proteins, for instance, form large arrays to increase signaling

output through cooperativity (16). Similar use of monomer/dimer transitions and evolution of

variable quaternary structure is at the heart of acquiring cooperativity and novel regulatory

modes in the classic hemoglobin family of oxygen-binding proteins, which started as ancestral

monomers prior to dimerizing by gene duplication events and the acquisition of a relatively small

number of mutations (33). Our findings suggest that various oligomeric states within the sensor

HK family could allow for different potential for cooperativity, perhaps depending on the input

signal or output pathway, or integrating multiple points of control or regulation to be added to the

canonical signal transduction pathway of two-component systems.

Experimental Procedures

Bioinformatics

Sequences similar to EL346 were identified by a BLAST search with default parameters

against the non-redundant NCBI database, using EL346 full-length protein sequence (Uniprot

ID: Q2NB77) as a query. EL346 and the best 100 hits were aligned using Clustal Omega (52)

and a distance tree was calculated from this alignment using the same software. Graphic

display of this tree was accomplished using iTOL (Interactive Tree of Life, http://itol.embl.de/)

(53). This tree was used to select HKs for cloning from the strains available at DSMZ

(http://www.dsmz.de).

Cloning, protein expression and purification

Five HKs were selected for biochemical characterization, abbreviated RT349, RH376,

ME354, RE356, and MS367 from the combination of their species name (e.g. *Rubellimicrobium*

thermophilum = RT) and number of amino acid residues (**Table 1**). DNA encoding these

proteins were cloned into a pHis-Gβ1 vector (54), with constructs verified by DNA sequencing.

Proteins were overexpressed in *Escherichia coli* BL21(DE3) (Stratagene) in LB as previously

described (18). Cells were harvested, resuspended in Buffer A (50 mM Tris pH 8.0, 100 mM

NaCl, and 10 mM MgCl$_2$), and lysed by sonication. Lysates were centrifuged at 48,000 x g and

4 °C for 45 min. Supernatants were loaded into a Ni^{2+} Sepharose affinity column (GE

Healthcare) and the His6-Gβ1 tagged protein was washed with 4 column volumes of Buffer A

supplemented with 15 mM imidazole and eluted with 4 column volumes of Buffer A

supplemented with 250 mM imidazole. Eluted proteins were exchanged into Buffer A by dialysis,

then fusion tags were cleaved by incubation with His$_6$-TEV protease overnight at 4 °C. The tag-

less proteins were separated from the tags and His$_6$-TEV protease by Ni^{2+} affinity

chromatography and were further purified by size-exclusion chromatography on a HiLoad

16/600 Superdex 200 column (GE Healthcare) equilibrated with HSEC buffer containing 50 mM

HEPES pH 7.5, 100 mM NaCl, 10 mM MgCl$_2$, and 0.5 mM DTT. For light-sensitive proteins, all

purification steps were performed under dim red light. Concentrations were determined from the

theoretical absorption coefficient, ε_{280} for PAS-HKs, calculated from the sequence using the

ExPASy ProtParam server (55), and $\varepsilon_{446} = 11,800$ M^{-1} cm^{-1} for flavin-containing proteins.

RH376 deletion mutants were generated as follows: RH2 and RH3 were ordered as

synthetic constructs from Genewiz (South Plainfield, NJ) and inserted into the same pHis-Gβ1

vector as the WT protein. RH1 and RH1+3 mutants were generated by PCR using an N-

terminal primer lacking the sequence to be deleted using WT and RH3 plasmids as templates,

respectively.

Absorbance and dark state reversion kinetics

UV-visible absorbance spectroscopy measurements were acquired using a Varian Cary

50 spectrophotometer at 24 °C with a 1 cm path length quartz cuvette. First, a UV-visible

spectrum in the range 250-550 nm was acquired in light and dark conditions for each LOV-HK.

For dark measurements, all steps were performed under dim red light. For light measurements,

the samples were illuminated for 1 minute using a blue LED panel (225 LED bulbs in 30.5 ×

30.5-cm panel, 13.8 W, 465 nm maximum illumination wavelength; LEDwholesalers [Hayward,

CA]) prior to each scan. To measure dark state recovery, each LOV-HK sample was illuminated

under a blue LED panel for 1 minute and then transferred to the spectrophotometer. A_{446} of the

samples in HSEC buffer with 1 mM ATP was monitored at 5-min intervals for 400-800 min at

24 °C. Time constants τ were determined by fitting measurements to a monoexponential decay

using the following equation:

$Abs = A_{inf} - (A_{inf} - A_{o})e^{-t/\tau}$, where $A_{inf} = A_{446}$ at infinity, $A_{o} = A_{446}$ initial, and $t = time$

Raw data were fit and plotted using RStudio with the nls function; plots were edited in Adobe

Illustrator CS6.

SEC-MALS

All samples and buffers were filtered with a 0.1-µm pore size filter before use. ATP was

added to each HK sample (~15 µM, 400 µL) to a final concentration of 1 mM. Samples were

then injected at a 0.4 mL/min flow rate onto a Superdex 200 GL 10/300 (or in the case of Figure

S4, Superdex 200 Increase GL 10/300) SEC column (GE Healthcare) using HSEC buffer with 1

mM ATP. For lit experiments, samples were illuminated with a blue LED panel for 1 min prior to

injection and the blue LED panel was faced toward the column during the run. For dark

experiments, all steps were performed under dim red light. Samples were detected post-elution

by the inline miniDAWN TREOS light scattering and Optilab rEX refractive index detectors

(Wyatt Technology [Santa Barbara, CA]). FPLC runs were performed at 4 °C, while MALS and

refractive index measurements were at 25 °C. Data analysis and molecular weight calculations

were performed using the ASTRA V software (Wyatt Technology). Raw data were plotted using

RStudio. Assignments of oligomeric state are based on a qualitative, five-bin scale, with proteins

considered to be "dimeric" or "monomeric" when their MALS-calculated mass falls within ±5% of

the sequence-derived molecular weight for either state. Between these ranges are three bins of

equal size: "mostly dimer", "monomer/dimer mixture", and "mostly monomer."

Autophosphorylation assays

Autophosphorylation was measured as described previously (1, 18). A minimum of three

trials were conducted for each protein under each condition. Proteins were added to a reaction

buffer containing 50 mM Tris, 100 mM NaCl, 5 mM $MnCl_2$, and 1 mM DTT with pH 8.2. Protein

concentrations were confirmed by analysis of dilution series on Coomassie-stained SDS-PAGE.

Final reaction concentrations ranged from approximately 10 to 35 micromolar as listed in the

table below:

Protein	Concentration (µM)	Protein	Concentration (µM)
EL346	16.1	RE356 monomer	9.6
ME354	11.6	RE356 dimer	8.5
RT349	28.6	RH1	12.4
RH376 WT	14.3	RH2	30.4
MS367 monomer	10.7	RH3	34.7
MS367 dimer	11.1	RH1+3	9.1

A mixture of unlabeled ATP and 10 µCi [γ-^{32}P] ATP was added to each mixture to initiate

the reaction (final ATP concentration of 200 µM). Aliquots were removed at time points of 0.5, 1,

2, 4, 8, 6, and 32 min and placed into a 4x SDS-gel loading buffer to quench the reaction. For

dark measurements, all steps were performed under dim red light. For light measurements, the

samples were illuminated with a blue LED panel just prior to and throughout the course of the

experiment. Samples were subjected to SDS-PAGE analysis; dried gels were exposed for 1hr -

overnight and bands were visualized by phosphorimaging. Band intensities were measured

using Fiji (56, 57); initial velocity values were calculated in Microsoft Excel from the linear region

of the curve and were plotted in RStudio.

Data availability – All data for this work are contained in this manuscript.

Supporting information – This article contains supporting information, including Figures S1-6,

Table S1, and accompanying literature citations.

Acknowledgements – The authors would like to acknowledge Bright Shi for generating the

RH1+3 construct, as well as Matthew Cleere, Zaynab Jaber, Jaynee Hart, and other Gardner

lab members for fruitful discussions.

Funding and additional information – This work was supported by NIH grants R01 GM106239

(to K.H.G.) and F32 GM1119311 (to I.D.). The content is solely the responsibility of the authors

and does not necessarily represent the official views of the National Institutes of Health.

Conflict of interest – The authors declare that they have no conflicts of interest with the contents

of this article.

Abbreviations –HK, histidine kinase; LOV, Light-Oxygen-Voltage; PAS, Per-ARNT-Sim; TCS,

two-component system; RR, response regulator; DHp, dimerization histidine phosphotransfer;

CA, catalytic ATP-binding; SEC-MALS, size exclusion chromatography multi-angle light

scattering.

References

1.   Rivera-Cancel, G., Ko, W. H., Tomchick, D. R., Correa, F., and Gardner, K. H. (2014)
      Full-length structure of a monomeric histidine kinase reveals basis for sensory regulation
      Proc Natl Acad Sci U S A **111**, 17839-17844 10.1073/pnas.1413983111

2.   Diensthuber, R. P., Bommer, M., Gleichmann, T., and Moglich, A. (2013) Full-length
      structure of a sensor histidine kinase pinpoints coaxial coiled coils as signal transducers
      and modulators Structure **21**, 1127-1136 10.1016/j.str.2013.04.024

3.   Galperin, M. Y. (2018) What bacteria want Environmental microbiology **20**, 4221-4229
      10.1111/1462-2920.14398

4.   Mascher, T., Helmann, J. D., and Unden, G. (2006) Stimulus perception in bacterial
      signal-transducing histidine kinases Microbiol Mol Biol Rev **70**, 910-938
      10.1128/MMBR.00020-06

5.   West, A. H., and Stock, A. M. (2001) Histidine kinases and response regulator proteins
      in two-component signaling systems Trends Biochem Sci **26**, 369-376,
      https://www.ncbi.nlm.nih.gov/pubmed/11406410

6.   Wang, S. (2012) Bacterial Two-Component Systems: Structures and Signaling
      Mechanisms In Protein Phosphorylation in Human Health, Huang C, ed. InTech, Chapter
      15,

7.   Mechaly, A. E., Soto Diaz, S., Sassoon, N., Buschiazzo, A., Betton, J. M., and Alzari, P.
      M. (2017) Structural Coupling between Autokinase and Phosphotransferase Reactions
      in a Bacterial Histidine Kinase Structure **25**, 939-944 e933 10.1016/j.str.2017.04.011

8.   Kenney, L. J., and Anand, G. S. (2020) EnvZ/OmpR Two-Component Signaling: An
      Archetype System That Can Function Noncanonically EcoSal Plus **9**,
      10.1128/ecosalplus.ESP-0001-2019

9.   Bhate, M. P., Molnar, K. S., Goulian, M., and DeGrado, W. F. (2015) Signal transduction

in histidine kinases: insights from new structures Structure **23**, 981-994

10.1016/j.str.2015.04.002

10.   Szurmant, H., White, R. A., and Hoch, J. A. (2007) Sensor complexes regulating two-

component signal transduction Curr Opin Struct Biol **17**, 706-715

10.1016/j.sbi.2007.08.019

11.   Gushchin, I., and Gordeliy, V. (2018) Transmembrane Signal Transduction in Two-

Component Systems: Piston, Scissoring, or Helical Rotation? Bioessays **40**,

10.1002/bies.201700197

12.   Rinaldi, J., Fernandez, I., Shin, H., Sycz, G., Gunawardana, S., Kumarapperuma, I. *et al.*

(2021) Dimer Asymmetry and Light Activation Mechanism in Brucella Blue-Light Sensor

Histidine Kinase mBio **12**, 10.1128/mBio.00264-21

13.   Ashenberg, O., Keating, A. E., and Laub, M. T. (2013) Helix bundle loops determine

whether histidine kinases autophosphorylate in cis or in trans J Mol Biol **425**, 1198-1209

10.1016/j.jmb.2013.01.011

14.   Herrou, J., Crosson, S., and Fiebig, A. (2017) Structure and function of HWE/HisKA2-

family sensor histidine kinases Curr Opin Microbiol **36**, 47-54 10.1016/j.mib.2017.01.008

15.   Luo, B. H., Carman, C. V., and Springer, T. A. (2007) Structural basis of integrin

regulation and signaling Annual review of immunology **25**, 619-647

10.1146/annurev.immunol.25.022106.141618

16.   Muok, A. R., Briegel, A., and Crane, B. R. (2020) Regulation of the chemotaxis histidine

kinase CheA: A structural perspective Biochimica et biophysica acta Biomembranes

1862, 183030 10.1016/j.bbamem.2019.183030

17.   Kitayama, Y., Nishiwaki-Ohkawa, T., and Kondo, T. (2014) Intersubunit communications

within KaiC hexamers contribute the robust rhythmicity of the cyanobacterial circadian

clock Microb Cell **1**, 67-69 10.15698/mic2014.01.129

18.   Dikiy, I., Edupuganti, U. R., Abzalimov, R. R., Borbat, P. P., Srivastava, M., Freed, J. H.

528   *et al.* (2019) Insights into histidine kinase activation mechanisms from the monomeric

529   blue light sensor EL346 Proc Natl Acad Sci U S A **116**, 4963-4972

530   10.1073/pnas.1813586116

531   19.   El-Gebali, S., Mistry, J., Bateman, A., Eddy, S. R., Luciani, A., Potter, S. C. *et al.* (2019)

532   The Pfam protein families database in 2019 Nucleic Acids Res **47**, D427-D432

533   10.1093/nar/gky995

534   20.   Correa, F., Ko, W. H., Ocasio, V., Bogomolni, R. A., and Gardner, K. H. (2013) Blue light

535   regulated two-component systems: enzymatic and functional analyses of light-oxygen-

536   voltage (LOV)-histidine kinases and downstream response regulators Biochemistry **52**,

537   4656-4666 10.1021/bi400617y

538   21.   Swartz, T. E., Tseng, T. S., Frederickson, M. A., Paris, G., Comerci, D. J., Rajashekara,

539   G. *et al.* (2007) Blue-light-activated histidine kinases: two-component sensors in bacteria

540   Science **317**, 1090-1093 10.1126/science.1144306

541   22.   Crosson, S., Rajagopal, S., and Moffat, K. (2003) The LOV domain family:

542   photoresponsive signaling modules coupled to diverse output domains Biochemistry **42**,

543   2-10 10.1021/bi026978l

544   23.   Pudasaini, A., El-Arab, K. K., and Zoltowski, B. D. (2015) LOV-based optogenetic

545   devices: light-driven modules to impart photoregulated control of cellular signaling Front

546   Mol Biosci **2**, 18 10.3389/fmolb.2015.00018

547   24.   Gao, R., and Stock, A. M. (2009) Biological insights from structures of two-component

548   proteins Annu Rev Microbiol **63**, 133-154 10.1146/annurev.micro.091208.073214

549   25.   Tseng, T. S., Frederickson, M. A., Briggs, W. R., and Bogomolni, R. A. (2010) Light-

550   activated bacterial LOV-domain histidine kinases Methods Enzymol **471**, 125-134

551   10.1016/S0076-6879(10)71008-9

26. Möglich, A. (2019) Signal transduction in photoreceptor histidine kinases Protein Sci **28**,

1923-1946 10.1002/pro.3705

27. Zoltowski, B. D., and Gardner, K. H. (2011) Tripping the light fantastic: blue-light

photoreceptors as examples of environmentally modulated protein-protein interactions

Biochemistry **50**, 4-16 10.1021/bi101665s

28. Crosson, S., and Moffat, K. (2001) Structure of a flavin-binding plant photoreceptor

domain: insights into light-mediated signal transduction Proc Natl Acad Sci U S A **98**,

2995-3000 10.1073/pnas.051520298

29. Purcell, E. B., McDonald, C. A., Palfey, B. A., and Crosson, S. (2010) An analysis of the

solution structure and signaling mechanism of LovK, a sensor histidine kinase

integrating light and redox signals Biochemistry **49**, 6761-6770 10.1021/bi1006404

30. Karniol, B., and Vierstra, R. D. (2004) The HWE histidine kinases, a new family of

bacterial two-component sensor kinases with potentially diverse roles in environmental

signaling J Bacteriol **186**, 445-453, https://www.ncbi.nlm.nih.gov/pubmed/14702314

31. Staroń, A., and Mascher, T. (2010) General stress response in alpha-proteobacteria:

PhyR and beyond Mol Microbiol **78**, 271-277 10.1111/j.1365-2958.2010.07336.x

32. Perica, T., Chothia, C., and Teichmann, S. A. (2012) Evolution of oligomeric state

through geometric coupling of protein interfaces Proc Natl Acad Sci U S A **109**, 8127-

8132 10.1073/pnas.1120028109

33. Pillai, A. S., Chandler, S. A., Liu, Y., Signore, A. V., Cortez-Romero, C. R., Benesch, J.

L. P. *et al.* (2020) Origin of complexity in haemoglobin evolution Nature **581**, 480-485

10.1038/s41586-020-2292-y

34. Banerjee, A., Herman, E., Kottke, T., and Essen, L. O. (2016) Structure of a Native-like

Aureochrome 1a LOV Domain Dimer from Phaeodactylum tricornutum Structure **24**,

171-178 10.1016/j.str.2015.10.022

35. Qin, L., Dutta, R., Kurokawa, H., Ikura, M., and Inouye, M. (2000) A monomeric histidine

kinase derived from EnvZ, an Escherichia coli osmosensor Mol Microbiol **36**, 24-32,

https://www.ncbi.nlm.nih.gov/pubmed/10760160

36. Tomomori, C., Tanaka, T., Dutta, R., Park, H., Saha, S. K., Zhu, Y. *et al.* (1999) Solution

structure of the homodimeric core domain of Escherichia coli histidine kinase EnvZ Nat

Struct Biol **6**, 729-734 10.1038/11495

37. Waterhouse, A., Bertoni, M., Bienert, S., Studer, G., Tauriello, G., Gumienny, R. *et al.*

(2018) SWISS-MODEL: homology modelling of protein structures and complexes

Nucleic Acids Research **46**, W296-W303 10.1093/nar/gky427

38. Rinaldi, J., Arrar, M., Sycz, G., Cerutti, M. L., Berguer, P. M., Paris, G. *et al.* (2016)

Structural Insights into the HWE Histidine Kinase Family: The Brucella Blue Light-

Activated Histidine Kinase Domain J Mol Biol **428**, 1165-1179

10.1016/j.jmb.2016.01.026

39. Berntsson, O., Diensthuber, R. P., Panman, M. R., Bjorling, A., Hughes, A. J., Henry, L.

et al. (2017) Time-Resolved X-Ray Solution Scattering Reveals the Structural

Photoactivation of a Light-Oxygen-Voltage Photoreceptor Structure **25**, 933-938 e933

10.1016/j.str.2017.04.006

40. Mechaly, A. E., Sassoon, N., Betton, J. M., and Alzari, P. M. (2014) Segmental helical

motions and dynamical asymmetry modulate histidine kinase autophosphorylation PLoS

Biol **12**, e1001776 10.1371/journal.pbio.1001776

41. Casino, P., Miguel-Romero, L., and Marina, A. (2014) Visualizing autophosphorylation in

histidine kinases Nat Commun **5**, 3258 10.1038/ncomms4258

42. Moglich, A., Ayers, R. A., and Moffat, K. (2009) Design and signaling mechanism of

light-regulated histidine kinases J Mol Biol **385**, 1433-1444 10.1016/j.jmb.2008.12.017

43. Lee, J., Tomchick, D. R., Brautigam, C. A., Machius, M., Kort, R., Hellingwerf, K. J. *et al.*

(2008) Changes at the KinA PAS-A dimerization interface influence histidine kinase

function Biochemistry **47**, 4051-4064 10.1021/bi7021156

44. Yarden, Y., and Schlessinger, J. (1987) Epidermal growth factor induces rapid,

reversible aggregation of the purified epidermal growth factor receptor Biochemistry **26**,

1443-1451 10.1021/bi00379a035

45. Liao, X., Liu, W., Yang, H. Q., and Jenkins, G. I. (2020) A dynamic model of UVR8

photoreceptor signalling in UV-B-acclimated Arabidopsis New Phytol **227**, 857-866

10.1111/nph.16581

46. Sang, Y., Li, Q. H., Rubio, V., Zhang, Y. C., Mao, J., Deng, X. W. *et al.* (2005) N-terminal

domain-mediated homodimerization is required for photoreceptor activity of Arabidopsis

CRYPTOCHROME 1 Plant Cell **17**, 1569-1584 10.1105/tpc.104.029645

47. Zoltowski, B. D., and Crane, B. R. (2008) Light activation of the LOV protein vivid

generates a rapidly exchanging dimer Biochemistry **47**, 7012-7019 10.1021/bi8007017

48. Zoltowski, B. D., Motta-Mena, L. B., and Gardner, K. H. (2013) Blue light-induced

dimerization of a bacterial LOV-HTH DNA-binding protein Biochemistry **52**, 6653-6661

10.1021/bi401040m

49. Scheuermann, T. H., Tomchick, D. R., Machius, M., Guo, Y., Bruick, R. K., and Gardner,

K. H. (2009) Artificial ligand binding within the HIF2alpha PAS-B domain of the HIF2

transcription factor Proc Natl Acad Sci U S A **106**, 450-455 10.1073/pnas.0808092106

50. Key, J., Hefti, M., Purcell, E. B., and Moffat, K. (2007) Structure of the redox sensor

domain of Azotobacter vinelandii NifL at atomic resolution: signaling, dimerization, and

mechanism Biochemistry **46**, 3614-3623 10.1021/bi0620407

51. Dutta, R., Qin, L., and Inouye, M. (1999) Histidine kinases: diversity of domain

organization Mol Microbiol **34**, 633-640 10.1046/j.1365-2958.1999.01646.x

52. Sievers, F., Wilm, A., Dineen, D., Gibson, T. J., Karplus, K., Li, W. *et al.* (2011) Fast,

scalable generation of high-quality protein multiple sequence alignments using Clustal

Omega Mol Syst Biol **7**, 539 10.1038/msb.2011.75

53. Letunic, I., and Bork, P. (2019) Interactive Tree Of Life (iTOL) v4: recent updates and

new developments Nucleic Acids Res **47**, W256-W259 10.1093/nar/gkz239

54. Sheffield, P., Garrard, S., and Derewenda, Z. (1999) Overcoming expression and

purification problems of RhoGDI using a family of "parallel" expression vectors Protein

Expr Purif **15**, 34-39 10.1006/prep.1998.1003

55. Gasteiger, E., Hoogland, C., Gattiker, A., Duvaud, S. e., Wilkins, M. R., Appel, R. D. *et*

al. (2005) Protein Identification and Analysis Tools on the ExPASy Server In The

Proteomics Protocols Handbook, Walker JM, ed. Humana Press, Totowa, NJ Chapter

Chapter 52, 571-607

56. Rueden, C. T., Schindelin, J., Hiner, M. C., DeZonia, B. E., Walter, A. E., Arena, E. T. *et*

al. (2017) ImageJ2: ImageJ for the next generation of scientific image data BMC

bioinformatics **18**, 529 10.1186/s12859-017-1934-z

57. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T. *et al.*

(2012) Fiji: an open-source platform for biological-image analysis Nat Methods **9**, 676-

682 10.1038/nmeth.2019

644