Are populations of economically important bonefish and queen conch 'open' or 'closed' in the northern Caribbean Basin?

Running title: Northern Caribbean gene flow

Marlis R. Douglas[1] | Tyler K. Chafin[1] | Julie E. Claussen[2] | David P. Philipp[2,3] | Michael E. Douglas[1]

[1]Department of Biological Sciences, University of Arkansas, Fayetteville, Arkansas, 72701

USA

[2]Fisheries Conservation Foundation, Champaign, IL 61820 USA

[3]Cape Eleuthera Institute, Rock Sound, Eleuthera, The Bahamas

Correspondence Michael E. Douglas med1@uark.edu/ https://orcid.org/0000-0001-9670-7825

KEY WORDS: ddRAD, dispersal, Eleuthera Island, Grand Bahama Island, outlier loci, SNPs

4   **Abstract**(243)(280)

Demographics of co-occurring species can often be diagnosed through population genomic analyses

6   of single nucleotide polymorphisms (SNPs). These data can define population structure, gene flow,

plus candidate regions in the genome that potentially reflect local adaptations. Importantly, they can

8   gauge if populations are demographically 'open' or 'closed' (i.e., with global or local recruitment). We

derived SNPs from double-digest restriction-site associated DNA (ddRAD) to test the demographics of

10   commercially important queen conch (*Lobatus gigas*) and bonefish (*Albula vulpes*) from two

northeast Caribbean Basin islands (Grand Bahama to the north and Eleuthera to the south).

12   Specifically, we tested the hypothesis that the strong west-to-east current in the Great Bahama

Canyon forms a vicariant barrier separating the two islands. We conducted Bayesian assignment

14   tests conducted on putatively neutral (*L. gigas*=64,863; *A. vulpes*=36,206 SNPs) and highly-

differentiated outlier SNP datasets (*L. gigas*=51–88; *A. vulpes*=79–123, respectively). For conch, both

16   analyses indicated weak structure, substantiating the vicariant hypothesis. However, results for

bonefish suggested adult mobility and strong easterly currents as potential drivers for source-sink

18   dynamics. These results, while preliminary, provide templates for future research endeavors with

these impacted species. Outlier loci, for example, can potentially place populations of each within a

20   demographic continuum, rather than within a dichotomous 'open'/'closed' framework, as well as

diagnose 'source' and 'sink' populations, as herein. These methodologies can then be applies to co-

22   distributed species with similar but less well understood ecologies so as to evaluate basin-wide

trends in connectivity and local adaptation.

1 | INTRODUCTION

24 Connectivity among populations is a fundamental consideration for marine conservation (Bell et al.

2014) and management (Miller et al. 2014), with aspects that resonate both theoretically (Kool et al.

26 2010, Ellingson & Krug 2015) and functionally (Hameed et al. 2016, Krueck et al. 2017). Benthic

marine invertebrates, for example, generally disperse as a component of the plankton so as to reach

28 areas beneficial for growth and development. This often extends across temporally variable periods

but with the end result being populations that are regarded as demographically 'open' (Domingues et

30 al. 2010). Many marine fishes follow a similar life-history strategy (Williamson et al. 2016), but with

an additional and innate capacity to 'self-recruit' back into their natal populations (Christie et al.

32 2010), a situation that serves to confound the traditional 'open' versus 'closed' dynamic. In a similar

vein, larger-bodied marine vertebrates often exhibit philopatry, defined as the return of reproducing

34 individuals to the locality from which they were spawned (Feldheim et al. 2014, Bonanomi et al.

2016, Salles et al. 2016). This, in turn, necessitates a capacity to home with great precision, in that

36 the natal areas were previously experienced only during the egg or larval stage. When this behavior

is manifested in both genders, a 'closed' population can result such that dynamics are driven not by

38 long-distance immigration but rather by local reproduction/ recruitment (Bentzen & Bradbury 2016).

A variety of mechanisms have been employed to evaluate dispersal and connectivity in

40 marine species, and consequently, to diagnose their open or closed status (Cowen et al. 2000, Cowen

& Sponaugle 2009). In this regard, much contemporary evidence has been provided via indirect

42 methods, with particular emphasis on genetic and genomic techniques that gauge 'effective

dispersal' (i.e., that due to survival and subsequent reproduction), rather than strictly by movements

44 identified among populations. The latter provide considerably greater resolution, such that results

establish larval relatedness (Ottmann et al. 2016), self-recruitment (Teske et al. 2016), panmixia

46   (Jasonowicz et al. 2017, Kensington et al. 2017, Tay et al. 2017), and adult dispersal (Veríssimo et al.

2017). Genomic assays have also been extended as well to estimate dispersal and connectivity

48   among deep-sea biodiversity, a group largely opaque to more traditional methods (Baco et al. 2016).

While methods of dispersal in marine biodiversity are many and varied, their variance can

50   often be a source of ambiguity, particularly when statistical tests of revelant hypotheses are

conducted (Weersing & Toonen 2009). For example, several studies have displayed varying results

52   when population connectivity was tested against the temporal duration of the planktonic larval

stage. In this sense, numeorus studies established a positive correlation between connectivity and

54   distance (Reisser et al. 2014, Ellingson & Krug 2015, Herrera et al. 2016, Morvezen et al. 2016, Tay et

al. 2016, Van Wyngaarden et al. 2016, Jasonowicz et al. 2017, Kensington et al. 2017, Martinez et al.

56   2017). Yet, numerous others have demonstrated just the reverse, i.e. an inverse relationship

between connectivity and distance (Kool et al. 2010, Jackson et al. 2014, Jorde et al. 2015, Cumming

58   et al. 2016, Gomes et al. 2016, Hameed et al. 2016, Teske et al. 2016). Clearly, many factors can

constrain or accelerate dispersal, with the end result being the promotion or depression of

60   connectivity.

In a similar vein, other studies have established global interconnectedness but surprisingly, as

62   a complex matrix of locally-adapted populations with larval transport varying both spatially and

temporally (Christie et al. 2010, Miller et al. 2014, Gould et al. 2017, McKeon et al. 2017, Segovia et

64   al. 2017). This, in turn, suggests a gradient in population structure, rather than a binary condition

with available choices established as either 'open' versus 'closed.'

66   Here, we investigated the spatial population structure of two economically important species

        in the northeastern Caribbean Basin, queen conch *Lobatus gigas* and bonefish *Albula vulpes*. We did

68   so by employing a double-digest restriction-site associated DNA (ddRAD) methodology that promotes

        the identification of single nucleotide polymorphisms (SNPs). Our study is the first to employ such

70   techniques with these economically important study species, and we hypothesized that our assay of

        neutral loci so derived would reflect the homogenizing effects of gene flow in each, thus emphasizing

72   their placement within the 'open' end of the demographic spectrum. We then sought to verify our

        general analyses of neutral loci within each species by deriving a more reduced and fine-grained

74   series of outlier SNPs that have the potential to demonstrate local adaptation. In doing so, we tested

        the hypothesis that sharp environmental gradients can counterbalance gene flow and instead

76   promote adaptive divergence.

        Finally, we recognized that our study species comprise exploited stocks, each of which

78   transects numerous national borders across the Caribbean Basin. As such, they experience varying

        levels of temporal and spatial exploitation. Thus, an understanding of coarse- and fine-grained

80   dispersal in these species, and indeed if they display connectivity, is a necessary baseline from which

        to derive an ecologically meaningful framework for sustainable conservation. In this sense, the

82   potential for local adaptation and source-sink dynamics between islands could seriously impact the

        management of conch and bonefish in the Caribbean.

84   **2 | MATERIALS AND METHODS**

        2.1 | Study site

86   The Caribbean Basin extends west from Florida along the Gulf Coast of North America, then south

      along the coast of México and Central America, continuing eastward across the northern coast of

88   South America, with end-points at the Greater and Lesser Antilles. Within this matrix, the

      Commonwealth of the Bahamas (Figure 1 top) represents a shallow series of carbonate banks

90   (~104,000 km^2) situated on a submerged continental border east of Florida (Andrews et al. 1970).

      Our Bahamian study sites are located along the coasts of two islands: Grand Bahama (153 km west-

92   east; 24 km north-south), and Eleuthera (~1.6 km west-east, 180 km north-south; Figure 1 top).

      Grand Bahama is located 86 km east of Palm Beach (FL) and represents the fourth largest island in

94   the Bahamian chain, whereas Eleuthera is some 370 km distant with the Atlantic Ocean on its east

      side and the shallow (<3m depth) Great Bahama Bank immediately west (Figure 2 top).

96   Approximately 1–3 km further west lie the abyssal waters of Exuma Sound (~1500m depth; "3" of

      Figure 2 top).

98   The Great Bahama Canyon (length=225 km, max width=37 km, max depth=6.9 km) is located

      northeast of the Great Bahama Bank ("1" of Figure 2 top), and effectively separates Grand Bahama

100   and Andros islands (to the north) from Eleuthera (to the south). It is divided into two main branches:

      'Northwest Providence Channel' extending north and west towards Grand Bahama Island, with

102   'Tongue of the Ocean' reaching south and east of Andros Island, towards Exuma Sound. The

      Northwest Providence Channel also absorbs a small but significant easterly contribution from the

104   Gulf Stream (Richardson & Finlen 1967), with currents extending west-to-east in the Canyon (green

      arrow, Figure 2 bottom). The occasionally turbid currents within the Great Bahama Canyon deposit a

106   large sediment fan as it merges with the Antilles Current of the Atlantic Ocean ~65km to the north-

northeast (red arrow, Figure 2 bottom) (Andrews 1970). The direction and extent of these currents

108   offer the potential for a vicariant separation between Grand Bahamas and Eleuthera islands.

2.2 | Bonefish

110   A complex of four bonefish species occur within the Atlantic Ocean/ Caribbean Basin: Bonefish

(*Albula vulpes*), Threadfin bonefish (*A. nemoptera*), Channel bonefish (*A. goreensis*), and an as yet

112   undescribed form *A. sp. cf. vulpes*. The three nominate species are partitioned non-geographically in

the Basin (Wallace & Tringali 2016). We restricted our evaluation to *A. vulpes*, a species that reflects

114   an elevated home range fidelity within the shallow flats proximate to Bahamian islands (Murchie et

al. 2013). However, deeper channels that separate the shallow flats can serve as conduits for travel,

116   or as refugia from rapid changes in water temperature (Humston et al. 2005). Inshore/nearshore

habitat are somewhat distant from home ranges, yet allow pre-spawning adults to congregate during

118   October-May. In this sense, pre-spawning bonefish are found on the south side of Grand Bahama

island where deeper waters are available for spawning (Murchie et al. 2015). Spawning movements

120   to offshore waters (>300 m) then occur on the full moon (Danylchuk et al. 2011). However, specific

spawning locations and actual spawning activities have yet to be identified/observed. Bonefish

122   produce leptocephali larvae that are planktonic, and subsequently drift for ~41–71 days before

eventually recruiting to inshore nurseries (Adams & Cook 2015).

124   A Basin-wide genetic analysis of *A. vulpes* (Wallace & Tringali 2016) revealed two somewhat

intermixed genetic populations. However, the extent of larval drift from local or distant sources

126   remains unknown for each. In this sense, populations of *A. vulpes* may reflect a situation found in

other high gene flow species, such as cardinalfish *Siphamia tubifer*, where genetic homogeneity

128   effectively conceals a high population turnover and a variable accumulation of larvae from divergent

sources (Gould et al. 2017). Brown crab *Cancer pagurus* displays a similar condition in the English

130   Channel and North Sea, where high levels of dispersal and gene flow effectively mask a genetic

patchiness due to recruitment that is both spatially and temporally variable (McKeon et al. 2017).

132   **2.3 | Queen conch**

Queen conch (*Lobatus gigas*) is not only the largest of six species in the Caribbean Basin, but also an

134   icon of considerable economic and cultural importance (Kough et al. 2017). It feeds on micro-and

macro-algae, but with habitat shifts that occur ontogenetically from shallower seagrass habitat into

136   deeper-water macroalgae plains (~40m) (Doerr & Hill 2013). Harvest (and over-harvest) are

promoted by a density-dependent reproductive strategy in which both genders not only reduce

138   movements but also aggregate along open sandy bottoms for broadcast spawning (Stoner 1997).

Spawning extends from April through October, with peak larval abundance in late June-

140   August when temperatures in the northeastern Caribbean Basin exceed 28°C (Stoner & Davis 1997a).

Planktonic larvae drift for ~60–75 days (Ballantine & Appeldoorn 1983), but with high natural

142   mortality and with distances controlled by oceanic conditions. For example, the surface environment

dictates the vertical distribution of larvae in the water column, with >80% of larvae occurring at 0.5-

144   5.0 m depth (Stoner & Davis 1997b).

In 1992, *L. gigas* was recognized by the Convention on International Trade of Endangered

146   Species (CITES: https://cites.org/eng/prog/queen_conch/introduction) and, in 1994, was added to

the Red List maintained by the International Union for the Conservation of Nature (IUCN;

148   http://www.fao.org/docrep/006/Y5261E/y5261e07.htm). A review of the *L. gigas* fishery is provided

by Brownell & Stevely (1981).

150   A broad-ranging spatial genetic study of *L. gigas* across the greater Caribbean (Truelove et al.

2017a,b) found that isolation-by-oceanic-distance dictated Basin-wide gene flow and pannmixia, but

152   surprisingly, not within the Bahamas. Similarly, a more regional study in the Mesoamerican

Caribbean again rejected panmixia (Machkour-M'Rabet et al. 2017), whereas a more homogeneous

154   gene flow was suggested for the Bahamas (Mitton et al. 1983). Also of note is the fact that overall

genetic homogeneity among populations (per the Bahamas) can be masked by the annual variation

156   in numbers of larvae that emerge from over-fished source populations.

2.4 | Collection, extraction, and amplification of samples

158   During 2014, we sampled fin clips from 57 adult bonefish collected by rod and reel from three

locations on the west and south sides of Eleuthera Island (EW, ES1, ES2; Table 1) and one from the

160   Atlantic Ocean (east) side (EEB; Table 1). Muscle samples were also gathered from 60 adults at Grand

Bahama Island, with 30 taken from the south (Flats) side (GSB) and 30 from the north (Atlantic) side

162   (GNB) (total N=117). Sampling was in compliance with the U.S. National Research Council Guide for

the Care and Use of Laboratory Animals (https://grants.nih.gov/grants/olaw/Guide-for-the-Care-and-

164   use-of-laboratory-animals.pdf). We also collected 30 individuals of queen conch from the south side

of Grand Bahama Island (GSC), and 30 from the Flats (or west) side of Eleuthera Island (EWC) (Table

166   1, Figure 1 bottom). Fin clips and muscle plugs were preserved on WHATMAN® FTA® Classic DNA cards,

with genomic DNA subsequently extracted using a GENTRA® Puregene DNA Purification Kit and stored

168   in hydrating solution (per manufacturer instructions). The quality of DNA was visualized on 2%

agarose gels, then quantified using a Qubit fluorometer (Thermo-Fisher Scientific, Inc.). Subsequent

170   library preparation followed previously published protocols (Peterson et al. 2012, as modfied in

Bangs et al. 2018).

172   Restriction digests were performed using 1µg of genomic DNA with PstI (5'-CTGCAG-3') and

MspI (5'-CCGG-3') in CutSmart buffer (New England Biosciences) for 20 hours at 37°C, after first

174   optimizing restriction enzyme and size selection choices using an *in silico* approach (FRAGMATIC; Chafin

et al. 2018). Digests were visualized on 2% agarose gels, purified using AMPure XP beads, and

176   quantified with a Qubit fluorometer. Digested samples were normalized to 0.1 µg of DNA and

subsequently ligated with barcoded Illumina adaptors, using custom oligos (Peterson et al. 2012).

178   Inline barcodes were designed to differ by at least 2 bases so as to minimize mis-assignments. Pooled

ligations were size-selected at 350-400 bps using the Pippin Prep automated size fractionator (Sage

180   Sciences). Adaptors were extended by PCR (10 cycles) with Phusion high-fideltiy DNA polymerase

(New England Biosciences). Size distribution and successful amplification of purified libraries was

182   then confirmed on the Agilent 2200 TapeStation. Indexed libraries were multiplexed with 100-bp

single-end sequencing on the Illumina HiSeq 2500 lane at the University of Oregon Genomics and Cell

184   Characterization Core Facility.

2.5 | Processing and assembly of ddRAD data

186   Raw reads were demultiplexed and processed in the PyRAD pipeline (Eaton 2014) using a 90%

identity threshold for clustering homologous reads. Read clusters representing putative loci within

188   individuals were filtered when alleles were > two (i.e., potentially overmerged or representing

paralogs), or when sequencing depth was <10. Following assembly, additional loci were discarded if

190   they exhibited: 1) <50% presence across individuals; 2) >60% heterozygosity per site among

individuals; 3) >10 heterozygous sites per consensus sequence; or 4) >10 consensus N bases. Further

192   data processing was performed using custom scripts (available at: github.com/tkchafin/scripts).

2.6 | Detection and spatial structure of local selection

194   We first sub-sampled all SNPs to a maximum of one per locus. We then utilized a Bayesian method

(BAYESCAN v.2.1, Foll & Gaggiotti 2008) with a burn-in period of 50,000 followed by 100,000

196   iterations to calculate per-locus F_{ST}, to determine a candidate list of potential loci under selection.

We used F_{ST}-etimates to derive three additional datasets per taxon: (1) All unlinked SNPs (=ALL); (2)

198   SNPs >2 standard deviations (sd) above mean F_{ST} (=SD2); and (3) SNPs >3 sd above the mean (=SD3).

We employed multiple methods to examine population structure for both species across all

200   three datasets. Hierarchical iterative clustering analysis using STRUCTURE v.2.2.4 (Pritchard et al. 2000)

was run in parallel (Besnier & Glover 2013) on the Jetstream cloud computing platform (Hancock et

202   al. 2019) using the GNU Parallel shell tool (Tange 2011). STRUCTURE was run for 250k repetitions with a

burn-in period of 100k, and with potential clusters for K-values ranging from 1-20. Optimal K for each

204   dataset was ascertained using the delta-K method (Evanno et al. 2005), as implemented and

visualized in the CLUMPAK pipeline (Kopelman et al. 2015). Differentiation among sampling localities

206   was also examined with Discriminant Analysis of Principal Components (DAPC) in the R package

ADEGENET (Jombart 2008, Jombart et al. 2008), with a minimal number of principal components

208   retained to explain 50% variance.

2.7 | Asymmetric migration among populations

210　Physical processes such as water current and tidal fluxes often create patterns of asymmetric gene

flow among populations, and consequently impact patterns of genetic diversity. Given our initial

212　hypothesis, we were interested in gauging the amount and direction of gene flow among the six

study populations of bonefish, and two of queen conch. In contrast to STRUCTURE, which employs a

214　Bayesian probabilistic model to assign individuals to ancestry clusters, we instead wished to estimate

the posterior probability of individual migratory histories, thereby allowing the direction and

216　magnitude of dispersal among populations to be dissected.

To do so, we employed a revised software optimized for large SNP datasets (i.e., BA3-SNPs,

218　Mussmann et al. 2019). We first tuned MCMC acceptance parameters using an automated iterative

procedure, then generated posterior estimates of migration rates using 250,000 iterations with a

220　burn-in period of 50,000. We further tested the hypothesis of source-sink dynamics by computing

net emigration rates as a sum of immigration rates subtracted from summed emigration rates

222　(Andreasen et al 2012), with positive numbers indicating a genetic source populations wheras

negative values indicated a sink population.

224　**3 | RESULTS**

After filtering, we obtained 56,046 loci for bonefish (*Albula vulpes*), of which 68.7% (N=36,206) were

226　parsimoniously informative (Table 2). Average number of loci per individual was 35,443, at a mean

depth of 77x. Subsampling only unlinked SNPs produced a matrix of 25,205 SNPs, from which SD2

228　and SD3 datasets were derived, and which contained 123 and 79 SNPs, respectively.

For queen conch (*Lobatus gigas*), 30,697 loci were available post-filtering, and represented

230　64,864 SNPs, of which 68.4% (N=44,334) were parsimony informative. Individuals were sequenced at

a depth of 58.6x, averging 33,409 loci per individual (Table 2). Subsampled unlinked SNPs totaled

232   19,716, from which 88 and 51 SNPs were derived as input to the SD2 and SD3 datasets.

3.1 | Spatial structure in Bonefish and Queen Conch

234   STRUCTURE runs for *A. vulpes* employing the ALL dataset (=neutral differentiation) revealed scant

differentiation among sample sites (Figure 3). The most highly-supported models for SD2 for SD3

236   were K=2 and K=3 groupings, both of which depicted GSB (Grand Bahama, south side) as clustereing

with EWB (Eleuthera, west side) and ES2 (south and west side of Eleuthera). The second cluster

238   placed GNB (Grand Bahama north side) with EEB (Eleuthera east or Atlantic side) and ES1 (south and

west side of Eleuthera) (Figure 3). The K=3 STRUCTURE run with SD2 and SD3 datasets was a bit mixed,

240   with GSB and EWB linked as one group, ES1, ES2, and EEB as a second group, with GNB as the third.

Subsequent K-groups were decidedly more mixed which rendered them indecipherable.

242   However, the DAPC clustering of the ALL dataset for bonefish (Figure 4 left) displayed

considerable overlap among sites (per STRUCTURE plot in Figure 3). Three populations with scant

244   separation were grouped within the center of the graph (i.e., ES1, ES2 and EEB). Three others were

somewhat more peripheral and reflected greater within-group variability (i.e., GNB, GSB, and EWB).

246   The same type of analysis, but employing the SD2 dataset (Figure 4 middle), separated GNB, EWB,

EEB, and ES1 without overlap, while grouping GSB with ES2. These relationships were maintained in

248   the DAPC clustering of the SD3 dataset, but with more overlap between the previously separated EEB

and ES1, whereas ES2 and GSB maintained their linkage. GNB and EWB remained distinct (Figure 4

250   right).

The STRUCTURE analysis of *L. gigas* populations using the ALL dataset revealed little genetic

252   differentiation, with mean F_{ST} (all SNPs) =0.0129. However both EWC and GSC showed significant

separation when both SD2 and SD3 datasets were evaluated (Figure 5). The DAPC clustering of the

254   ALL dataset (Figure 6 left) again showed considerable overlap between the two islands, whereas each

separated with considerably less overlap when SD2 was analysed, and became even more distinct

256   based upon the SD3 (Figure 6, middle, right).

3.2 | Asymmetric migration

258   The BAYESASS3 analysis of *L. gigas* found very high demographic independence among populations,

with 'self-migration' (e.g., recruitment) rates estimated at >98% for both Grand Bahama and Eleuthra

260   populations (Table 3). Results for bonefish revealed highly variable and asymmetric migration rates,

with EEB and EWB depicted as source populations, with outward migration rates to both Grand

262   Bahama and flats-side Eleuthra populations. This varied from 8.5-17.9% for EEB and 7.5-14.6% for

EWB (Table 3; Figure 7). Notably, the remaining populations lacked any strong signal of outward

264   migration, and all showed negative rates of net emigration (Figure 8), suggesting a sink dynamic.

4 | DISCUSSION

266   Populations are frequently classified as to the manner by which they are replenished. In this sense,

closed populations are defined as having recruitment via local reproduction, whereas open

268   populations are independent of source (Pinsky et al. 2012). This dichotomy underscores several

ecological, evolutionary, and conservation-based research agendas: Exploitation or protection (Yau

270   et al. 2014, Krueck et al. 2017); Life history and taxonomy (Mora & Sale 2002, Baco et al. 2016);

Recovery from disturbance (Bell et al. 2014); and Extent to which populations are locally adapted

272 (Jorde et al. 2015, Salles et al. 2016, Segovia et al. 2017). It also has obvious implications for how

populations are managed, yet with considerable resolution provided by the advent of genomic

274 techniques, and particularly with the estimation of local adaptation (Reisser et al. 2014).

4.1 | F_{ST} outliers for queen conch

276 For non-model organisms (as herein), two approaches are most often utilized: The first identifies loci

that display unusually high genetic differentiation among populations (i.e., F_{ST} outlier approach),

278 whereas the second (i.e., gene-environment association) seeks correlations between the allele

frequencies of populations and their local environments. We employed the first approach in this

280 study, and by so doing, found that outlier loci did indeed discriminate sessile populations of queen

conch *Lobatus gigas* separated by a potential vicariant barrier (the Great Bahama Canyon).

282 This is aptly demonstrated in Figure 5, where the south side of Grand Bahama Island and the

west side of Eleuthera Island are clearly discrete one from another. This condition is reinforced by

284 the DAPC analysis (Figure 6), where distributions for SD2, and SD3 datasets consistently diverge from

that depicted for the ALL dataset. F_{ST} values for these two populations are not significantly different

286 when computed from the ALL data, but are when compiled from the SD2 and SD3 datasets (Table 3).

Of interest is the fact that whereas neutral loci showed little differntiation across the study area, a

288 smaller proportion reflected strong patterns of differentiation consistent with local adaptation. This

scenario has also been manifested in other marine species that were similarly investigated using SNP

290 analyses (Cure et al. 2017, Thomas et al. 2017).

4.2 | F_{ST} outliers for bonefish

292   A similar analysis employing data for bonefish *Albula vulpes* failed to group samples on either side of

the Canyon into geographically congruent groups, suggesting instead a distribution more panmictic

294   in format (per Sᴛʀᴜᴄᴛᴜʀᴇ analysis and DAPC plot for the ALL data). Furthermore, Sᴛʀᴜᴄᴛᴜʀᴇ plots for

K=2 and K=3 groupings (Figure 3) deviated but little. For K=2, both SD2 and SD3 datasets linked the

296   south side of Grand Bahama Island (GSB) with the west side of Eleuthera (EWB) and with one (of

two) ES locations (i.e., ES2). This, in turn, suggests a level of connectivity across the Great Bahama

298   Canyon, a conjecture supported by Figure 4. However, the north side of Grand Bahama Island (GNB)

grouped with the eastern (Atlantic) side of Eleuthera (EEB) and with the second ES site in the K=2

300   grouping (i.e., ES1). This relationship separates at K=3, with GSB remaining linked with EWB, whereas

EEB and ES2 separate. These relationships are difficult to interpret biologically, particularly given

302   what we know of geomorphic and current patterns in the northeast Basin (Figure 2). A more

parsimonious explanation would point to the considerable mobility of adult bonefish superimposed

304   onto a lack of site specificity.

4.3 | Potential problems with F_{ST} outlier analyses

306   One potential issue with detecting selection via departure from neutrality is that each locus may

manifest an independent response that can hinge upon population structure and demography. This is

308   particularly problematic when the average level of differentiation among populations is relatively

high (Hoban et al. 2017). However, pairwise F_{ST} values for bonefish are neither elevated nor

310   significant (Table 3), suggesting this concern may not apply in our situation. An additional concern is

that some loci may demonstrate false signatures of selection, such as when small populations found

312   on the leading edge of an expansion are contributing disproportionately to observed frequency

differences. Again, our study populations are already established and thus the concept of a leading

314 edge of expansion is not a viable consideration. Similarly, cryptic hybridization and introgression can

invoke confounding effects as well, but again, these are not apparent in our data.

4.4 | Asymmetrical migration in bonefish and queen conch

The structure of the physical environment can promote differential movements in aquatic organisms,

318 and thus promote a metapopulation structure within a species (i.e., a source–sink dynamic). This in

turn can skew estimates of genetic differentiation among populations, and consequently promote

320 erroneous management decisions, particularly when the organisms are anthropogenically harvested

such that active management is based on an economic model. In this sense, we employed genetic

322 estimates of population migration so as to ascertain potential source-sink considerations in our study

organisms, and to determine if indeed movements between them were asymmetrical.

324 In this study, populations of conch were found to be highly demographically independent, as

seen in both estimates of contemporary migration (BA3-SNPs) and from estimates of population

326 structure using highly-differentiated SNPs. On the other hand, bonefish showed strong signals of

source-sink dynamics across islands. The two Eleuthra populations (EEB and EWB) seemingly act as

328 genetic sources, as evidenced by their high net emigration rates derived from BayesAss3 (Figure 8).

Their genetic differentiation may suggest the presence of discrete genetic stocks, reflecting either

330 spatial or temporal asynchrony with regards to spawning. Given a tendency for bonefish to move

into deeper waters for spawning (Danylchuk et al 2011), the differentiation between the two

332 Eleuthera populations may indicate spawning site-fidelity, with subsequent intermingling of

ancestries in flat-side feeding grounds. The latter may also explain the cryptic intra-site structure as

334   recorded in the assignment tests of Fst outlier SNP datasets (Figure 3).

4.5 | Dispersal and connectivity within the Caribbean Basin

336   We hypothesized that the population demographics of our study species would be impacted by the

currents passing through the Grand Bahama Canyon. Prior to test, we sought data from othere

338   species that might potentially shed light on this conjecture. Jackson et al. (2014) investigated genetic

differentiation among 19 spawning aggregations of Nassau grouper *Epinephelus striatus* in the Basin.

340   The life history of this species parallels that of bonefish (above) in that spawning aggregations are

cued by water temperature and phases of the moon over a 3-month period. As with bonefish,

342   breeding aggregations develop some distance from individual home ranges, with larvae drifting for a

similar duration (~40 days) before settling. Four Basin-wide genetic groups were identifed, to include

344   the Bahamas, but with little within-group differentiation, again in parallel with that found for

bonefish (per Wallace & Tringali 2016). A major barrier to larval dispersal was the separation of the

346   Basin into eastern and western regions (Jackson et al. 2014, Figure 3). These results implicate the

Great Bahama Canyon (Figure 1 top) as a vicariant barrier, similar to what was found herein with

348   conch.

In a second study, Kool et al. (2010) coupled results from a matrix-based projection model

350   with those based on a bio-physical approach to examine genetic dispersal among coral reef patches

in the Basin. Allele frequencies were found to shift along the length of the Bahamas, with the

352   sharpest demarcation occurring at the separation between Grand Bahama Island and the southern

Bahamas. This separation was not recognized in the original publication but indeed juxtaposes with

354 the presence of the Grand Bahama Canyon (Figure 1). Again, the results from Kool et al. (2010)

promote the Canyon as a major vicariant feature in the region.

356 **5 | CONCLUSIONS**

An understanding of genetic connectivity within exploited species is a crucial aspect, such that

358 appropriate geographic units can be subsequently designated for management purposes. This

includes the establishment of quotas for 'take,' as well as the delineation boundaries for marine

360 protected areas (MPAs). Furthermore, an understanding of Basin-wide linkages (or lack thereof)

among populations of commercially exploited species (as herein) promotes the realization among the

362 numerous stakeholders that fisheries management is a complex and continually evolving geopolitical

problem.

364   Understanding adaptive (i.e., ecologically relevant) variation potentially allows the

incorporation of such information into conservation and management practices (Allendorf 2017). For

366 instance, it could be used to define adaptive-based management units that, in high gene flow

species, may represent subsets within more traditional neutral-based units (Funk et al. 2012). Outlier

368 SNPs could also identify illegal conch fishing and help define stock boundaries and potential

management units (Milano et al. 2014).

370   These aspects cut to the definition of conservation genomics, which (in the narrow sense) is

defined as those approaches that differ conceptually and quantitatively from traditional genetics.

372 This, in turn, allows genome-wide signals of adaptation to be detected, and this in turn subsequently

provides answers to those questions impossible to address using genetic data alone (Garner et al.

374 2016). Importantly, such data can provide genomic evidence for potential heat tolerance (or the lack

thereeof), an issue that has serious reverberations for sessile marine invertebrates in particular (as

376 herein) (Thomas et al., 2017). These issues loom large as oceans acidify and their heat profiles

expand in lockstep with global climate change (Hurd et al., 2018).

378 **ACKNOWLEDGMENTS**

Funding was provided by the following generous endowments to the University of Arkansas: The

380 Bruker Professorship in Life Sciences (MRD), a Distinguished Doctoral Fellowship (TKC), and the 21st

Century Chair in Global Change Biology (MED). We gratefully acknowledge the Cape Eleuthera

382 Institute (CEI) and The Island School for logistical support and field assistance. Sampling was

supported under CEI collecting permits. Sampling was supported by grants from Patagonia's World

384 Trout Initiative, the Charles A. and Anne Morrow Lindbergh Foundation, the Baldwin Foundation, as

well as generous personal donations from B. Hallig, T. Rice, and J. Spring. Sequencing and analyses

386 were supported by the Arkansas Economic Development Commission (Arkansas Settlement Proceeds

Act of 2000) and the Arkansas High Performance Computing Center (AHPCC). Sequence data were

388 generated at the Genomics and Cell Characterization Core Facility, University of Oregon/ Eugene.

Raw datasets are available via NCBI SRA (accession to be added upon acceptance).

390 **AUTHOR'S CONTRIBUTIONS**

MRD/ JEC/ DPP/ MED designed the study; JEC/ DPP contributed to fieldwork; MRD/ TKC/ MED

392 contributed to lab work; TKC performed bioinformatic work and statistical analyses; MRD/ TKC/ MED

interpreted results and generated figures; MRD/ TKC/ MED drafted the manuscript; All authors gave

394 final approval for publication.

ORCID

396   *Michael E. Douglas* https://orcid.org/0000-0001-9670-7825

398   **DATA ACCESSIBILITY**

DNA SNP loci: Dryad Digital Repository (http://dx.doi.org/upon-acceptance).

REFERENCES

402   Adams, A. J., & Cooke, S. J. (2015). Advancing the science and management of flats fisheries for

bonefish, tarpon, and permit. *Environmental Biology of Fishes*, 98, 2123–2131

404   (doi:10.1007/s10641-015-0446-9)

Allendorf, F. W. (2017). Genetics and the conservation of natural populations: Allozymes to genomes.

406   *Molecular Ecology*, 26, 420–430 (doi:10.1111/mec.13948)

Andreasen, A. M., Stewart, K. M., Longland, W. S., Beckmann, J. P., & Forister, M. L. (2012).

408   Identification of source-sink dynamics in mountain lions of the Great Basin. *Molecular

Ecology*, 21, 5689–5701 (doi: 10.1111/j.1365-294X.2012.05740.x)

410   Andrews, J. E. (1970). Structure and sedimentary development of the outer channel of the Great

Bahama Canyon. *Geological Society of America Bulletin*, 81, 217–225 (10.1130/0016-

412   7606(1970)81[217:SASDOT]2.0.CO;2)

Andrews, J. E., Shepard, F. P., & Hurley, R. J. (1970). Great Bahama Canyon. *Geological Society of

414   America Bulletin*, 81. 1061–1078 (doi:10.1130/0016-7606(1970)81[1061:GBC]2.0)

Baco, A. R., Etter, R .J., Ribeiro, P. A., & Von Der Heyden, S., Beerli, P. & Kinlan, B. P. (2016). A

416   synthesis of genetic connectivity in deep-sea fauna and implications for marine reserve

design. *Molecular Ecology*, 25, 3276–3298 (doi:10.1111/mec.13689)

418   Ballantine, D. L., & Appeldoorn, R. S. (1983). Queen conch culture and future prospects in

Puerto Rico. *Proceedings of the Gulf and Caribbean Fisheries Institute,* 35, 57–63

420   (https://www.researchgate.net/publication/283924228)

Bangs, M. R., Douglas, M. R., Mussmann, S. M., & Douglas, M. E. (2018). Unraveling historic

422   introgression and resolving phylogenetic discord within *Catostomus* (Pisces:

   Catostomidae). *BMC Evolutionary Biology,* 18, 86 (doi:10.1186/s12862-018-1197-y)

424 Bell, J. J., Smith, D., Hannan, D., Haris, A., Jompa, J., & Thomas, L. (2014). Resilience to disturbance

   despite limited dispersal and self-recruitment in tropical barrel sponges: Implications for

426   conservation and management. *PLoS ONE*, 9, e91635

   (doi:10.1371/journal.pone.0091635)

428 Bentzen, P., & Bradbury, I. R. (2016). Don't bet against the natal homing abilities of marine

   fishes. *Molecular Ecology*, 25, 2691–2692 (doi:10.1111/mec.13591)

430 Besnier, F., & Glover, K. A. (2013). ParallelStructure: An R Package to distribute parallel runs of the

   population genetics program Structure on multi-core computers. PLoS ONE 8(7), e70651

432   (doi:10.1371/journal.pone.0070651)

Bonanomi, S., Therkildsen, N. O., Retzel, A., Hedeholm, R. B., Pedersen, M. W., Meldrup, D., …

434   Nielsen, B. C. (2016). Historical DNA documents long-distance natal homing in marine fish.

   Molecular Ecology, 25, 2727–2734 (doi:10.1111/mec.13580)

436 Brownell, W. N., & Stevely, J. M. (1981). The biology, fisheries and management of the queen

   conch, *Lobatus gigas*. *Marine Fish Review*, 43, 1-12 (spo.nmfs.noaa.gov/mfr437/mfr4371.pdf)

438 Chafin, T. K., Martin, B. T., Mussmann, S. M., Douglas, M. R., & Douglas, M. E. (2018). FRAGMATIC: in

   silico locus prediction and its utility in optimizing ddRADseq projects. *Conservation Genetics*

440   *Resources*, 10, 325-328 (doi: 10.1007/s12686-017-0814-1)

Christie, M. R., Johnson, D. W., Stallings, C. D., & Hixon, M. A. (2010). Self-recruitment and

442   sweepstakes reproduction amid extensive gene flow in a coral-reef fish. *Molecular Ecology*,

19, 1042–1057 (doi:10.1111/j.1365-294X.2010.04524.x)

444   Cowen, R. K., Lwiza, K. M. M., Sponaugle, S., Paris, C. B., & Olsen, D. B. (2000). Connectivity of

marine populations: Open or closed? *Science*, 287, 857–59

446   (doi:10.1126/science.287.5454.857)

Cowen R. K., & Sponaugle, S. (2009). Larval dispersal and marine population connectivity.

448   *Annual Review of Marine Science*, 1, 443–466

(doi:10.1146/annurev.marine.010908.163757)

450   Cumming, R. A., Nikula, R., Spencer, H. G., & Waters, J. M. (2016). Trans-Tasman genetic connectivity

in the intertidal air-breathing slug *Onchidella nigricans*. *Marine Ecology Progress Series*,   562,

452   93–100, 2016 (doi:10.3354/meps11926)

Cure, K., Thomas, L., Hobbs, J-P. A., Fairclough, D. B., & Kennington, W. J. (2017). Genomic signatures

454   of local adaptation reveal source-sink dynamics in a high gene flow fish species. *Scientific

Reports*, 7, 8618 (doi:10.1038/s41598-017-09224-y)

456   Danylchuk, A. J., Cooke, S. J,, Goldberg, T. L., Suski, C. D. , Murchie, K. J., Danylchuk, S. E., ...

Philipp, D. P. (2011). Aggregations and offshore movements as indicators of spawning

458   activity of bonefish (*Albula vulpes*) in The Bahamas. *Marine Biology*, 158, 1981–1999

(doi:10.1007/s00227-011-1707-6)

460   Domingues, C. P., Creer, S., Taylor, M. I., Queiroga, H., & Carvalho, G. R. (2010). Genetic structure of

Carcinus maenas within its native range: Larval dispersal and oceanographic variability.

462   *Marine Ecology Progress Series*, 410, 111–123 (doi:10.3354/meps08610)

Doerr, J. C., & Hill, R. L. (2013). Home range, movement rates, and habitat use of queen conch,

464   *Strombus gigas*, in St. John, U.S. Virgin Islands. *Caribbean Journal of Science*, 47, 251-259

(doi:10.18475/cjos.v47i3.a13)

466   Eaton, D. A. R. (2014). PʏRAD: Assembly of de novo RADseq loci for phylogenetic analyses.

Bioinformatics, 30, 1844–1849 (doi:10.1093/bioinformatics)

468   Ellingson, R. A., & Krug, P. J. (2015). Reduced genetic diversity and increased reproductive

isolation follow population-level loss of larval dispersal in a marine gastropod. *Evolution*,

470   70, 18–37 (doi:10.1111/evo.12830)

Evanno, G., Regnaut, S., & Goudet, J. (2005). Detecting the number of clusters of individuals using the

472   software Sᴛʀᴜᴄᴛᴜʀᴇ: A simulation study. *Molecular Ecology*, 14, 2611–2620 (doi:

0.1111/j.1365-294X.2005.02553.x)

474   Feldheim, K. A., Gruber, S. H., DuBattista, J. D., Babcock, E. A., Kessel, S. T., Hendry, A. P, …

Chapman, D. D. (2014). Two decades of genetic profiling yields first evidence of natal

476   philopatry and long-term fidelity to parturition sites in sharks. *Molecular Ecology*, 23, 110–

117 (doi:10.1111/mec.12583)

478   Foll, M., & Gaggiotti, O. (2008). A genome-scan method to identify selected loci appropriate for

both dominant and codominant markers: a Bayesian perspective. *Genetics,* 180, 977–993

480   (doi:10.1534/genetics.108.092221)

Funk, W. C., McKay, J. K., Hohenlohe, P. A., & Allendorf, F. W. (2012). Harnessing genomics for

482   delineating conservation units. *Trends in Ecology and Evolution*, 27, 489–496

(doi:10.1016/j.tree.2012.05.012)

484 Garner, B. A., Hand, B. K., Amish, S. J., Bernatchez, L., Foster, J. T., Miller, K. M., ... Luikart, G. (2016).

Genomics in conservation: Case studies and bridging the gap between data and application.

486   *Trends in Ecology and Evolution*, 31, 81–83 (doi:10.1016/j.tree.2015.10.009)

Gomes, I., Peteiro, L. G., Albuquerque, R., Nolasco, R., Dubert, J., Swearer, S. E., & Queiroga, H.

488   (2016). Wandering mussels: Using natural tags to identify connectivity patterns among

Marine Protected Areas. *Marine Ecology Progress Series*, 552, 159–176

490   (doi:10.3354/meps11753)

Gould, A. L., & Dunlap, P. V. (2017). Genomic analysis of a cardinalfish with larval homing potential

492   reveals genetic admixture in the Okinawa Islands. *Molecular Ecology*, 26, 3870–3882

(doi:10.1111/mec.14169)

494 Hameed, S. O., White, J. W., Miller, S. H., Nickols, K. J., & Morgan, S. G. (2016). Inverse approach

to estimating larval dispersal reveals limited population connectivity along 700 km of

496   wave-swept open coast. *Proceedings of the Royal Society of London Part B Biological*

Sciences, 283, 20160370 (doi:10.1098/rspb.2016.0370)

498   Hancock, D. Y., Stewart, C. A., Vaughn, M., Fischer, J., Lowe, J. M., Turner, G., ... Snapp-Childs, W.

(2019). Jetstream—Early operations performance, adoption, and impacts. *Concurrency and*

500   *Computation: Practice and Experience*, 31, e4683 (doi: 10.1002/cpe.4683)

Herrera, M., Nanninga, G. B., Planes, S., Jones, G. P., Thorrold, S. R., Saenz-Agudelo, P., ...

502   Berumen, M. L. (2016). Seascape and life-history traits do not predict self-recruitment in

a coral reef fish. *Biology Letters*, 12, 20160309 (doi:10.1098/rsbl.2016.0309)

504   Hoban, S., Kelley, J. L., Lotterhos, K. E., Antolin, M. F., Bradburd, G., Lowry, D. B., ... Whitlock, M. C.

(2017). Finding the genomic basis of local adaptation: Pitfalls, practical solutions, and future

506   directions. *American Naturalist*, 188, 379–397 (doi:10.1086/688018)

Humston, R., Ault, J. S., Larkin, M. F , & Luo, J. (2005). Movements and site fidelity of the

508   bonefish *Albula vulpes* in the northern Florida Keys determined by acoustic telemetry.

Marine Ecology Progress Series, 291, 237–248 (doi:10.3354/meps291237)

510   Hurd, C.L., Lenton, A., Tilbrook, B., & Boyd, P. W. (2018). Current understanding and challenges for

oceans in a higher-CO_2 world. *Nature Climate Change*, 8, 686–694 (doi:10.1038/s41558-018-

512   0211-0)

Jackson, A. M., Semmens, B. X., Sadovy de Mitcheson, Y., Nemeth, R. S., Heppell, S. A., Bush, P. G., ...

514   Bernardi, G. (2014). Population structure and phylogeography in Nassau Grouper

(*Epinephelus striatus*), a mass-aggregating marine fish. *PLoS ONE*, 9, e97508

516   (doi:10.1371/journal.pone.0097508)

Jasonowicz, A. J., Goetz, F. W., Goetz, G. W., & Nichols, K. M. (2017). Love the one you're with:

518   Genomic evidence of panmixia in the sablefish (*Anoplopoma fimbria*). *Canadian Journal of*

Fisheries and Aquatic Science, 74, 377–387 (doi:10.1139/cjfas-2016-0012)

520   Jombart, T. (2008). ADEGENET: An R package for the multivariate analysis of genetic markers.

Bioinformatics, 24, 1403–1405 (doi:10.1093/bioinformatics/btn129)

522   Jombart, T., Devillard, S., Dufour, A-B., & Pontier, D. (2008). Revealing cryptic spatial patterns in

genetic variability by a new multivariate method. *Heredity*, 101, 92–103

524   (doi:10.1038/hdy.2008.34)

Jorde, P. E., Søvik, G., Westgaard, J-I., Albretsen, J., André, C., Hvingel, C., ... Jørstad, K. B.

526   (2015). Genetically distinct populations of northern shrimp, *Pandalus borealis*, in the

North Atlantic: Adaptation to different temperatures as an isolation factor. *Molecular*

528   *Ecology*, 24, 1742–1757 (doi:10.1111/mec.13158)

Kensington, W. J., Keron, P. W., Harvey, E. S., Wakefield, C.B., Williams, C. B., Halafihi, A. J., ...

530   Newman, S. J. (2017). High intra-ocean, but limited inter-ocean genetic connectivity in

populations of the deep-water oblique-banded snapper *Pristipomoides zonatus* (Pisces:

532   Lutjanidae). *Fisheries Research*, 193, 242–249 (doi:10.1016/j.fishres.2017.04.015)

Kool, J. T., Paris, C. B., Andréfouët, S., & Cowen, R. K. (2010). Complex migration and the

534   development of genetic structure in subdivided populations: An example from   Caribbean

coral reef ecosystems. *Ecography*, 33, 597-606 (doi:10.1111/j.1600-0587.2009.06012.x)

536   Kopelman. N. M., Mayzel, J., Jackobsson, M. J., Rosenberg, N. A., & Mayrose, I. (2015). CLUMPAK: a

        program for identifying clustering modes and packaging population structure inferences

538   across K. *Molecular Ecology Resources*, 15, 1179–1191 (doi: 10.1111/1755-0998.12387)

      Kough, A. S., Cronin, H., Skubel, R., Belak, C. A., & Stoner, A. W. (2017). Efficacy of an established

540   marine protected area at sustaining a queen conch *Lobatus gigas* population during three

        decades of monitoring. *Marine Ecology Progress Series*, 573, 177–189

542   (doi:10.3354/meps12163)

      Krueck, N. C., Ahmadia, G. N., Green, A., Jones, G. P., Possingham, H. P., Riginos, C., ... Mumby,   P. J.

544   (2017). Incorporating larval dispersal into MPA design for both conservation and fisheries.

        Ecological Applications, 27, 925–941 (doi:10.1002/eap.1495)

546   Machkour-M'Rabet, S., Cruz-Medina, J., García-De León, F. J. , De Jesús-Navarrete. A., Hénaut,   Y.

        (2017). Connectivity and genetic structure of the queen conch on the Mesoamerican Reef.

548   *Coral Reefs*, 36, 535–548 (doi:10.1007/s00338-017-1551-3)

      Martinez, E., Buonaccorsi, V., Hyde, J. R., & Aguilar, A. (2017). Population genomics reveals high gene

550   flow in grass rockfish (*Sebastes rastrelliger*). *Marine Genomics*, 33, 57–63

        (doi:10.1016/j.margen.2017.01.004)

552   McKeon, N. J., Hauser, L., & Shaw, P. W. (2017). Microsatellite genotyping of brown crab *Cancer*

        pagurus reveals fine scale selection and 'non-chaotic' genetic patchiness within a high gene

554   flow system. *Marine Ecology Progress Series*, 566, 91–103   (doi:10.3354/meps12044)

Milano, I., Babbucci, M., Cariani, A., Atanassova, M., Bekkevold, D., Carvalho, G. R., … Bargelloni, L.

556   (2014). Outlier SNP markers reveal fine-scale genetic structuring across European hake

populations (*Merluccius merluccius*). *Molecular Ecology*, 23, 118-135

558   (doi:10.1111/mec.12568)

Miller, K. J., Mundy, C. N., & Mayfield, S. (2014). Molecular genetics to inform spatial

560   management in benthic invertebrate fisheries: A case study using the Australian

greenlip abalone. *Molecular Ecology*, 23, 4958–4975 (doi:10.1111/mec.12914)

562   Mitton, J. B., Berg, C. J. Jr., & Orr, K. S. (1989). Population structure, larval dispersal, and gene   flow

in the Queen Conch, *Lobatus gigas*, of the Caribbean. *Biological Bulletin*, 177, 356-362

564   (doi:10.2307/1541595)

Mora, C., & Sale, P. F. (2002). Are populations of coral reef fish open or closed? *Trends in*

566   *Ecology and Evolution*, 17, 422-428 (doi:10.1016/S0169-5347(02)02584-3)

Morvezen, R., Charrier, G., Boudry, P., Chauvaud, L., Breton, F., Strand, Ø., & Laroce J. (2016).

568   Genetic structure of a commercially exploited bivalve, the great scallop *Pecten maximus*,

along the European coasts. *Conservation Genetics*, 17, 57–67 (doi:10.1007/s10592-015-

570   0760-y)

Murchie, K. J., Cooke, S. J., Danylchuk, A. J., Danylchuk, S. E., Goldberg, T., Suski, C., & Philipp,   D.P.

572   (2013). Movement patterns of bonefish (*Albula vulpes*) in tidal creeks and coastal waters of

Eleuthera, The Bahamas. *Fisheries Research*, 147, 404-412

574   (doi:10.1016/j.fishres.2013.03.019)

Murchie, K. J., Schultz, A. D., Stein, J. A., Cooke, S. J., Lewis, J., Franklin, J., ... Philipp, D.P. (2015).

576   Defining adult bonefish (*Albula vulpes*) movement corridors around Grand Bahama in the

Bahamian Archipelago. *Environmental Biology of Fishes*, 98, 2203–2212 (doi:10.1007/s10641-

578   015-0422-4)

Mussmann, S. M., Douglas, M. R., Chafin, T. K., & Douglas, M. E. (2019). BA3-SNPs:

580   Contemporary migration reconfigured in BayesAss for next-generation sequence data.

Methods in Ecology and Evolution, 10, 1808-1813 (doi:10.1111/2041-210X.13252)

582 Ottmann, D., Grorud-Colvert, K., Sard, N. M., Huntington, B. E., Banks, M. A., & Sponaugle, S.

2016). Long-term aggregation of larval fish siblings during dispersal along an open coast.

584   *Proceedings of the National Academy of Sciences U.S.A.,* 113, 14067–14072

(doi:10.1073/pnas.1613440113)

586 Peterson, B. K., Weber, J. N., Kay, E. H., Fisher, H. S., & Hoekstra, H. E. (2012). Double digest

RADseq: An inexpensive method for de novo SNP discovery and genotyping in model and

588   non-model species. *PLoS ONE,* 7, e37135 (doi:10.1371/journal.pone.0037135)

Pinsky, M. L,, Palumbi, S. R., Andrefouet, S., & Purkis, S. J. (2012). Open and closed seascapes:

590   Where does habitat patchiness create populations with high fractions of self-

recruitment? *Ecological Applications*, 22, 1257–1267 (doi:10.1890/11-1240.1)

592 Pritchard, J. K., Stephens, M., & Donnelly, P. (2000). Inference of population structure using

multilocus genotype data. *Genetics*, 155, 945–959 (PMC1461096)

594   Reisser, C. M.O., Bell, J. J., & Gardner, J. P.A. (2014). Correlation between pelagic larval duration

        and realised dispersal: Long-distance genetic connectivity between northern New

596   Zealand and the Kermadec Islands archipelago. *Marine Biology*, 161, 297–312

        (doi:10.1007/s00227-013-2335-0)

598   Richardson, W. S., & Finlen, J. R. (1967). The transport of Northwest Providence Channel. *Deep-Sea*

        Research, 14, 361-367 (doi:10.1016/0011-7471(67)90078-2).

600   Salles, O. C., Pujol, B., Maynard. J. A., Almany, G. A., Berumen, M. L., Jones, G. P., ... Planes, S.

        (2016). First genealogy for a wild marine fish population reveals multigenerational

602   philopatry. *Proceedings of the National Academy of Sciences U.S.A.,* 113:13245–13250

        (doi:10.1073/pnas.1611797113)

604   Segovia, N. I., Gallardo-Escárate, C., Poulin, E., & Haye, P. A. (2017). Lineage divergence, local

        adaptation across a biogeographic break, and artificial transport, shape the genetic

606   structure in the ascidian *Pyura chilensis*. *Science Reports*, 7, 44559 (doi:10.1038/srep44559)

        Stoner, A. W. (1997). The status of queen conch, *Lobatus gigas*, research in the Caribbean.

608   *Marine Fisheries Review*, 59, 14-23   (https://www.researchgate.net/publication/242472717)

        Stoner, A. W., & Davis, M. (1997a). Abundance and distribution of queen conch veligers (*Lobatus*

610   *gigas*) in the central Bahamas. I. Horizontal patterns in relation to reproductive and nursery

        grounds. *Journal of Shellfish Research*, 16, 7-18

612   (http://www.biodiversitylibrary.org/bibliography/2179#/summary)

Stoner, A. W., & Davis, M. (1997b). Abundance and distribution of queen conch veligers (*Lobatus*

614   *gigas*) in the central Bahamas. II. Vertical patterns in nearshore and deep-water habitats.

             Journal of Shellfish Research, 16, 19-29

616   (http://www.biodiversitylibrary.org/bibliography/2179#/summary)

Tay, Y. C., Chng, M. W. P., Sew, W. W. G. , Rheindt, F. E., Tun,, K. P. P., & Meier, R. (2016).

618   Beyond the Coral Triangle: High genetic diversity and near panmixia in Singapore's

             populations of the broadcast spawning sea star *Protoreaster nodosus*. *Royal Society   Open*

620 *Science*, 3, 160253 (doi:10.1098/rsos.160253)

Tange, O. (2011), GNU PARALLEL: The command-line power tool

622   (https://www.usenix.org/system/files/login/articles/105438-Tange.pdf)

Teske, P. R., Sandoval-Castillo, J., van Sebille, E., Waters, J., & Beheregaray ,L. B. (2016).

624   Oceanography promotes self-recruitment in a planktonic larval disperser. *Science*

             Reports, 6, 34205 (doi:10.1038/srep34205)

626 Thomas, L., Kennington, W. J., Evans, R. D., Kendrick, G. A., & Stat, M. (2017). Restricted gene flow

             and local adaptation highlight the vulnerability of high-latitude reefs to rapid environmental

628   change. *Global Change Biology*, 23, 2197–2205 (doi:10.1111/gcb.13639)

Truelove, N. K., Kough, A. S., Behringer, D. C., Paris, C. B., Box, S. J., Preziosi, R. F., & Butler, M. J.

630   IV. (2017). Biophysical connectivity explains population genetic structure in a highly

             dispersive marine species. *Coral Reefs*, 36, 233–244 (doi:10.1007/s00338-016-1516-y)

632   Truelove, N. K., Box, S. J., Aiken, K. A., Blythe-Mallett, A., Boman, E. M., Booker, C. J., ... Stoner,   A. W.

(2017). Isolation by oceanic distance and spatial genetic structure in an overharvested

634   international fishery. *Diversity and Distributions*, 23, 1–9 (doi:10.1111/ddi.12626)

Van Wyngaarden, M., Snelgrove, P. V. R., DiBacco, C., Hamilton, L. C., Rodríguez-Ezpeleta, N.,

636   Jeffery, N. W., ... Bradbury, I. R. (2016). Identifying patterns of dispersal, connectivity, and

selection in the sea scallop, Placopecten magellanicus, using RADseq-derived SNPs.

638   *Evolutionary Appiications*, 10, 102–117 (doi:10.1111/eva.12432)

Veríssimo, A., Sampaio, I., McDowell, J. R., Alexandrino, P., Mucientes, G., Queiroz, N., ... Noble,

640   L. R. (2017). World without borders—genetic population structure of a highly migratory

marine predator, the blue shark (*Prionace glauca*). *Ecology and Evolution*, 7, 4768–4781

642   (doi:10.1002/ece3.2987)

Wallace, E. M., & Tringali, M. D. (2016). Fishery composition and evidence of population structure

644   and hybridization in the Atlantic bonefish species complex (*Albula* spp.). *Marine Biology*, 163,

142.(doi:10.1007/s00227-016-2915-x)

646   Weersing, K., & Toonen, R. J. (2009). Population genetics, larval dispersal, and connectivity in

marine systems. *Marine Ecology Progress Series*, 393, 1–12 (doi:10.3354/meps08287)

648   Williamson, D. H., Harrison, H. B., Almany, G. R., Berumen, M. L., Bode, M., Bonin, M. C., ...

Jones, G. P. (2016). Large-scale, multidirectional larval connectivity among coral reef fish

650   populations in the Great Barrier Reef Marine Park. *Molecular Ecology*, 25, 6039–6054

(doi:10.1111/mec.13908)

652   Yau, A. J., Lenihan, H. S., & Kendall, B. E. (2014) Fishery management priorities vary with self-

recruitment in sedentary marine populations. *Ecological Applications*, 24, 1490-1504

654   (doi:10.1890/13-1201.1)

TABLE 1 Collection data for bonefish (*Albula vulpes*) and queen conch (*Lobatus gigas*). Listed are: Species, Island, Side (of island with compass direction in parentheses), Locality, Date, N, and Sample label).

Species	Island	Side	Locality	Date	N	Label
Bonefish	Grand Bahama	Flats (N)	North	26-Jan-14	30	GNB
Bonefish	Grand Bahama	Ocean (S)	South	25-Jan-14	30	GSB
Bonefish	Eleuthra	Flats (W)	Broad Creek	2-Mar-14	30	EWB
Bonefish	Eleuthra	Ocean (E)	Half Sound	8-Mar-14	10	EEB
Bonefish	Eleuthra	Flats (SW)	Plum Creek	8-Mar-14	10	ES1
Bonefish	Eleuthra	Flats (SW)	Sound Side	10-Mar-14	7	ES2
Conch	Grand Bahama	South	Freeport	1-Jan-14	30	GSC
Conch	Eleuthra	Flats (W)	Schooner Cay 2	1-Mar-14	30	EWC

TABLE 2 Study species (= bonefish *Albules vulpes*; queen conch *Lobatus gigas*) evaluated in this study.
662 Also provided are number of genetic loci greater than 50% retention (= Loci > 50%); Average number
of loci per individual (= Av loci/ind); Average depth of coverage (= Av coverage); Number of SNPs
664 generated (= N SNPs); and Number of parsimonious-informative SNPs (= PI SNPs).

Species	Loci > 50%	Av loci/ind	Av coverage	N SNPs	PI SNPs
Bonefish	38,221	39,502	73X	36,797	25,039
Conch	30,697	33,409	59X	64,863	44,334

666 TABLE 3 BᴀʏᴇsAss3 results for bonefish (*Albula vulpes*) and queen conch (*Lobatus gigas*) sampled at Grand Bahama and Eleuthera islands. Values are expressed as proportion of individuals expected to
668 be of a given migrant or non-migrant origin. Migration rates along the diagonal are analogous to a self-recruitment rate, with high values representative of demographic independence. Sample site
670 abbreviations per Table 1.

	EWB	EEB	ES1	ES2	GNB	GSB	GNC	EWC
EWB	0.884	0.039	0.020	0.019	0.021	0.018	-	-
EEB	0.085	0.833	0.020	0.021	0.020	0.020	-	-
ES1	0.104	0.146	0.689	0.020	0.020	0.021	-	-
ES2	0.127	0.106	0.026	0.693	0.025	0.025	-	-
GNB	0.104	0.075	0.016	0.016	0.774	0.015	-	-
GSB	0.179	0.077	0.015	0.016	0.016	0.698	-	-
GNC	-	-	-	-	-	-	0.987	0.013
EWC	-	-	-	-	-	-	0.013	0.988

FIGURE LEGENDS

674   FIGURE 1 (Top) Map depicting The Commonwealth of Bahama (insert top right). The islands of Grand

Bahama, Abaco, Eleuthera and Andros (northeastern Basin) are depicted. Blue dots represent sample

676   sites for queen conch, whereas those for bonefish are designated with red dots. (Bottom) Sample

sites at Eleuthera Island for queen conch and bonefish. Colors as above. Labels as in Table 1.

FIGURE 2 (Top) Bathymetry map of the northwest Carribean Basin (modified from

680   http://oceanexplorer.noaa.gov/explorations/09deepseacorals/logs/summary/media/bahamas_regio

n_600.html). Red ellipse designates Eleuthera Island; 1 = Great Bahama Canyon; 2 = Great Bahama

682   Flat; 3 = Exuma Sound. (Bottom) Oceanic currents in the northern Caribbean Basin (modified from:

University of Miami Rosenthiel School of Marine and Atmospheric Science,

684   http://oceancurrents.rsmas.miami.edu/caribbean/caribbean-maps.html). Black lines depict currents;

Yellow arrow designates direction of Gulf Stream (east coast of Florida); Green arrow indicates

686   direction of current in Grand Bahama Canyon; 2 = Eleuthera Island; Red Arrow indicates direction of

the Antillean Current (Atlantic Ocean).

FIGURE 3. Structure plots for bonefish (*Albula vulpes*) sampled at Grand Bahama and Eleuthera

690   islands, The Bahamas. K=2 represent a 2-group analysis employing (1) the SD2 dataset containing

single nucleotide polymorphisms (SNPs) >2 standard deviations (sd) above mean Fst (N=99), and (2)

692   the SD3 dataset containing SNPs >3 sd above mean Fst (N=64). K = 3 represents a 3-groupl analysis

involving the same two datasets. GN = Grand Bahamas Flats North (Flats or North side); GS = Grand

694   Bahamas Ocean Side (South or Bahamian side); EW = Eleuthera Flats Broad Creek (West side); EE =

Eleuthera Ocean Half-Sound (Atlantic Ocean or East side); ES1 = Eleuthera Ocean Plum Creek (Flats or

696   Southwest side); ES2 = Eleuthera Ocean South Side (Flats or Southwest side). More details on sample

sites provided in Table 1.

FIGURE 4. Plots depicting discriminant analysis of principal components (DAPC) for six populations of

700   bonefish (*Albula vulpes*) along discriminant functions one (=DF1) and two (=DF2). ALL = plot utilising

All loci (N=38,221); SD2 = plot depicting single nucleotide polymorphisms (SNPs) > 2 standard

702   deviations (sd) above mean Fst (N=99); SD3 = plot employing SNPs > 3 sd above mean Fst (N=64). GN

= Grand Bahamas Flats North (Flats or North side); GS = Grand Bahamas Ocean Side (South or

704   Bahamian side); EW = Eleuthera Flats Broad Creek (West side); EE = Eleuthera Ocean Half-Sound

(Atlantic Ocean or East side); ES1 = Eleuthera Ocean Plum Creek (Flats or Southwest side); ES2 =

706   Eleuthera Ocean South Side (Flats or Southwest side). More details provided in Table 1.

708   FIGURE 5. Structure plots for queen conch (*Lobatus gigas*) sampled as Grand Bahama and Eleuthra

islands, the Bahamas, with individuals represented by vertical lines. Two groups are depicted [GBS =

710   Grand Bahama South Freeport (Flats or South side); EW = Eleuthera Flats Schooner Cay 2 (Flats or

West side)]; ALL = Plot employing all single nucleotide polymorphisms (SNPs) (N=30,697); SD2 = SNPs

712   > 2 standard deviations (sd) above mean Fst (N=88); and SD3 = plot depicting SNPs > 3 sd above

mean Fst (N=51).

FIGURE 6. Plots depicting discriminant analysis of principal components (DAPC) for two populations

716   of queen conch (*Lobatus gigas*), with GBS = Grand Bahama South Freeport (Flats or South side) and

EW = Eleuthera Flats Schooner Cay 2 (Flats or West side) plotted along discriminant function one

718   (=DF1). All = Plot employing all single nucleotide polymorphisms (SNPs) (N=30,697); SD2 = SNPs > 2

standard deviations (sd) above mean Fst (N=88); and SD3 = plot depicting SNPs > 3 sd above mean

720   Fst (N=51).

722   FIGURE 7. Graph depicting relative migration rates for six populations of bonefish (*Albula vulpes*),

where nodes represent populations, and edges are weighted according to the level of direction

724   relative migration rates (*m*), as derived from BA3-SNPs results. All edges below 0.02 were omitted.

Full results shown in Table 3. GN = Grand Bahamas Flats North (Flats or North side); GS = Grand

726   Bahamas Ocean Side (South or Bahamian side); EW = Eleuthera Flats Broad Creek (West side); EE =

Eleuthera Ocean Half-Sound (Atlantic Ocean or East side); ES1 = Eleuthera Ocean Plum Creek (Flats or

728   Southwest side); ES2 = Eleuthera Ocean South Side (Flats or Southwest side).

730   FIGURE 8. Asymmetric immigration recorded for bonefish (*Albula vulpes*) from Grand Bahama and

Eleuthra islands. Values are expressed as net emigration rates (summed emigration - summed

732   immigration) for each population, with positive values indicating 'source' populations and negative

values indicating 'sink' populations.