Are populations of economically important bonefish and queen conch 'open' or 'closed' in the northern Caribbean Basin?

Running title: Northern Caribbean gene flow

Marlis R. Douglas[1] | Tyler K. Chafin[1] | Julie E. Claussen[2] | David P. Philipp[2,3] | Michael E. Douglas[1]

[1]Department of Biological Sciences, University of Arkansas, Fayetteville, Arkansas, 72701 USA

[2]Fisheries Conservation Foundation, Champaign, IL 61820 USA

[3]Cape Eleuthera Institute, Rock Sound, Eleuthera, The Bahamas

Correspondence Michael E. Douglas med1@uark.edu/ https://orcid.org/0000-0001-9670-7825

Abstract

Demographics of co-occurring species can often be diagnosed through population genomic analyses of single nucleotide polymorphisms (SNPs). These data can define population structure, gene flow, and candidate regions in the genome that potentially reflect local adaptations. They can also gauge if populations are demographically 'open' or 'closed' (i.e., with global or local recruitment). We derived SNPs from double-digest restriction-site associated DNA (ddRAD) to test the demographics of commercially important bonefish *Albula vulpes* (N=117) and queen conch *Lobatus gigas* (N=60) from two northeast Caribbean Basin islands (Grand Bahama to the north and Eleuthera to the south). Specifically, we tested the hypothesis that the strong west-to-east current in the Great Bahama Canyon is a vicariant barrier separating the two islands. We conducted Bayesian assignment tests on putatively neutral (*A. vulpes*=36,206 SNPs; *L. gigas*=64,863) and highly-differentiated outlier datasets (*A. vulpes*=123 and 79 SNPs; *L. gigas*=88 and 51, respectively). For bonefish, results diagnosed asymmetrical gene flow and north-south differentiation, as potentially driven by adult mobility and easterly currents. However, both analyses indicated genetic structure in conch, substantiating the vicariant hypothesis. These results provide templates for future research endeavors with these impacted species. Outlier loci, for example, can potentially place populations of each within a demographic continuum, rather than within a dichotomous 'open/closed' framework, as well as diagnose 'source' and 'sink' populations, as herein. These methodologies can then be applied to co-distributed species with similar but less well understood ecologies so as to evaluate basin-wide trends in connectivity and local adaptation.

1 | INTRODUCTION

Connectivity among populations is a fundamental consideration for both marine conservation (Bell et al. 2014) and management (Miller et al. 2014), with aspects that resonate both theoretically (Kool et al. 2010, Ellingson & Krug 2015) and functionally (Hameed et al. 2016, Krueck et al. 2017). Benthic marine invertebrates and many sedentary fishes generally disperse as a component of the plankton so as to reach areas beneficial for growth and development. This often extends across temporally variable periods with the end result being populations diagnosed as demographically 'open' (Domingues et al. 2010). Many marine fishes follow a similar life-history strategy (Williamson et al. 2016), but with an additional and innate capacity to 'self-recruit' back into their natal populations (Christie et al. 2010), a situation that confounds the traditional 'open' versus 'closed' dynamic. In a similar vein, larger-bodied marine vertebrates often exhibit philopatry, defined as the return of reproducing individuals to the locality from which they were spawned (Feldheim et al. 2014, Bonanomi et al. 2016, Salles et al. 2016). This, in turn, necessitates a capacity to home with great precision, in that natal areas were previously experienced only during the egg or larval stage. When this behavior is manifested in both sexes, a 'closed' population can result such that dynamics are driven not by long-distance immigration but rather by local reproduction and recruitment (Bentzen & Bradbury 2016).

Numerous mechanisms have been employed to evaluate dispersal and connectivity in marine species, and consequently, to diagnose their open or closed status (Cowen et al. 2000, Cowen & Sponaugle 2009). In this regard, much contemporary evidence stems from indirect methods, with particular emphasis on genetic and genomic techniques that gauge

'effective dispersal' (i.e., that due to survival and subsequent reproduction), rather than direct observation of individual movements among populations. The former provide considerably greater resolution, and often establish larval relatedness (Ottmann et al. 2016), self-recruitment (Teske et al. 2016), panmixia (Jasonowicz et al. 2017, Kensington et al. 2017, Tay et al. 2017), and adult dispersal (Veríssimo et al. 2017). Genomic assays have also been extended as well to estimate dispersal and connectivity among deep-sea biodiversity, where studies are largely dominated by more conservative methods (Baco et al. 2016).

While methods of dispersal in marine biodiversity are many and varied, their variance can often be a source of ambiguity, particularly when statistical tests of revelant hypotheses are conducted (Weersing & Toonen 2009). For example, several studies have displayed varying results when population connectivity was tested against the temporal duration of the planktonic larval stage. Some recognized a positive correlation between connectivity and distance (Reisser et al. 2014, Ellingson & Krug 2015, Herrera et al. 2016, Morvezen et al. 2016, Tay et al. 2016, Van Wyngaarden et al. 2016, Jasonowicz et al. 2017, Kensington et al. 2017, Martinez et al. 2017). Yet, others have demonstrated an inverse relationship (Kool et al. 2010, Jackson et al. 2014, Jorde et al. 2015, Cumming et al. 2016, Gomes et al. 2016, Hameed et al. 2016, Teske et al. 2016). Clearly, dispersal can be constrained or accelerated by many factors , with the promotion or depression of connectivity being an end result.

In a similar vein, other studies have established global interconnectedness but, contrastingly, as a complex matrix of locally-adapted populations, with larval transport varying both spatially and temporally (Christie et al. 2010, Miller et al. 2014, Gould et al. 2017, McKeon et al. 2017, Segovia et al. 2017). This, in turn, suggests a gradient in population structure, rather than a binary condition where available choices are either 'open' *versus* 'closed.' Here, we investigated the existence of binary spatial population structure as potential endpoints of a continuum, and did so by examining populations of economically important marine biodiversity in the northeastern Caribbean Basin. Our focus on a distant corner of the Caribbean ecosystem was based on the presence of an extensive and deep marine trench (the Great Bahama Canyon) that may serve as a potential west-to-east vicariant barrier that prevents larval gene flow between populations of bonefish *Albula vulpes* and queen conch *Lobatus gigas* located at islands to the north and south, respectively.

1.1 | Bonefish

A complex of four bonefish species occur within the Atlantic Ocean/ Caribbean Basin: Bonefish (*Albula vulpes*), Threadfin bonefish (*A. nemoptera*), Channel bonefish (*A. goreensis*), and an as yet undescribed form *A. sp. cf. vulpes*. The three nominate species are partitioned non-geographically in the Basin (Wallace & Tringali 2016). We restricted our evaluation to *A. vulpes*, a species with an elevated home range fidelity within the shallow flats proximate to Bahamian islands (Murchie et al. 2013). However, deeper

channels separaing shallow flats can serve as conduits for travel or as refugia from rapid

changes in water temperature (Humston et al. 2005).

Inshore/nearshore habitats are somewhat distant from home ranges, yet allow pre-

spawning bonefish to congregate during October-May, such as on the south side of Grand

Bahama island (Figure 1) where deeper spawning waters are available (Murchie et al.

2015). Spawning movements into offshore waters then occur on the full moon (Danylchuk

et al. 2011), with adults gathering at depth (>130m) prior to a quick ascent (~67m in

<2min) to spawn within a distinct oceanographic layer of low turbulence and flow (i.e.,

optimal pycnocline/ thermocline) that subsequently retains the planktonic leptocephali

larvae (Lombardo et al. 2020). The latter subsequently drift for ~41–71 days before

eventually recruiting to unspecified inshore nurseries (Adams & Cook 2015).

A basin-wide mtDNA analysis of *A. vulpes* (Wallace & Tringali 2016) revealed two

somewhat intermixed genetic populations. However, the extent of their larval drift from

local or distant sources is unknown. Populations of *A. vulpes* may reflect a situation found

in other high gene flow species, such as the cardinalfish *Siphamia tubifer*, where genetic

homogeneity effectively conceals a high population turnover and a variable accumulation

of larvae from divergent sources (Gould et al. 2017). The brown crab, *Cancer pagurus,*

displays a similar condition in the English Channel and North Sea, where high levels of

dispersal and gene flow effectively mask a genetic patchiness that results from a

recruitment that is spatially and temporally variable (McKeon et al. 2017).

1.2 | Queen conch

Queen conch (*Lobatus gigas*) is not only an icon of considerable economic and cultural importance (Kough et al. 2017), but also represents the largest of six species in the Caribbean Basin. It feeds on micro-and macro-algae, but with habitat shifts that occur ontogenetically from shallower seagrass beds into deeper-water macroalgal plains (~40m in depth) (Doerr & Hill 2013). Harvest (and over-harvest) are promoted by a density-dependent reproductive strategy in which both sexes not only reduce movements but also aggregate along open sandy bottoms for broadcast spawning (Stoner 1997).

Spawning extends from April through October, with peak larval abundance in late June-August when temperatures in the northeastern Caribbean Basin exceed 28°C (Stoner & Davis 1997a). Planktonic larvae drift for ~60–75 days (Ballantine & Appeldoorn 1983), but with high natural mortality and with distances controlled by oceanic conditions. For example, the surface environment dictates the vertical distribution of larvae in the water column, with >80% of larvae occurring at 0.5-5.0m depth (Stoner & Davis 1997b).

In 1992, *L. gigas* was recognized by the Convention on International Trade of Endangered Species (CITES: https://cites.org/eng/prog/queen_conch/introduction) and was subsequently added in 1994 to the Red List maintained by the International Union for the Conservation of Nature (IUCN; http://www.fao.org/docrep/006/Y5261E/y5261e07.htm). A review of the *L. gigas* fishery is provided by Brownell & Stevely (1981).

A broad-ranging spatial genetic study of *L. gigas* across the greater Caribbean using nine microsatellite DNA markers (Truelove et al. 2017a,b) found that basin-wide gene flow

was constrained by oceanic distance that served to isolate local populations, but surprisingly, not within Bahamian populations where F_{ST} values dis not differ significantly. Similarly, a more regional study in the Mesoamerican Caribbean using five inter-microsatellite DNA (ISSR) primers again rejected panmixia (Machkour-M'Rabet et al. 2017), whereas an earlier protein electrophoretic study in the Bahamas indicated gene flow was more homogeneous (Mitton et al. 1983). Also of note is the fact that overall genetic homogeneity among populations (per the Bahamas) can be masked by the annual variation in numbers of larvae produced by over-fished source populations.

1.3 | Research objectives

In this study, we employed a double-digest restriction-site associated DNA (ddRAD) methodology to identify single nucleotide polymorphisms (SNPs) as diagnostic markers in bonefish and conch. Our study is the first to employ this techniques with these economically important study species, and we hypothesized that our derivation of neutral loci would reflect the homogenizing effects of gene flow in each, thus emphasizing their placement within the 'open' end of the demographic spectrum. We then sought to contrast our initial results employing neutral loci by deriving a more reduced and fine-grained series of outlier SNPs within each species that may potentially demonstrate local adaptation. By doing so, we tested the hypothesis that sharp environmental gradients within a vicariant marine topography can counterbalance gene flow and promote instead adaptive divergence.

Finally, we recognize that our study species comprise exploited stocks, each of which transects numerous national borders across the Caribbean Basin. As such, they experience varying levels of temporal and spatial exploitation. Thus, a more nuanced understanding of coarse- and fine-grained dispersal in these species, and indeed if they display connectivity, provides a necessary baseline from which to derive an ecologically meaningful framework for sustainable conservation. In this sense, the potential for local adaptation and source-sink dynamics between islands could provide a rationale from which to seriously recalibrate conch and bonefish management in the Caribbean.

2 | MATERIALS AND METHODS

2.1 | Study site

The Caribbean Basin extends westward from Florida along the Gulf Coast of North America, then south along the coast of México and Central America, continuing eastward across the northern coast of South America, with end-points at the Greater and Lesser Antilles. Within this matrix, the Commonwealth of the Bahamas (Figure 1 top) represents a shallow series of carbonate banks (~104,000km^2) situated on a submerged continental border east of Florida (Andrews et al. 1970). Our Bahamian study sites are located along the coasts of two islands: Grand Bahama (153km west-east; 24km north-south), and Eleuthera (~1.6km west-east, 180km north-south; Figure 1 top). Grand Bahama is located 86km east of Palm Beach (FL) and represents the fourth largest island in the Bahamian chain, whereas Eleuthera is some 370km distant, with the Atlantic Ocean on its east side and the shallow (<3m depth)

Great Bahama Bank immediately west (Figure 2 top). Approximately 1–3km southwest lie the abyssal waters of Exuma Sound (~1500m depth; "3" of Figure 2 top).

The Great Bahama Canyon (length=225km, max width=37km, max depth=6.9km;"1" of Figure 2 top) is located northeast of the Great Bahama Bank, and effectively separates Grand Bahama and Abaco islands (to the north) from Eleuthera (to the south; Figure 1). It is divided into two main branches: 'Northwest Providence Channel' extending north and west towards Grand Bahama Island, with 'Tongue of the Ocean' extending south along the east coast of Andros Island, facing Exuma Sound. The Northwest Providence Channel also absorbs a small but significant easterly contribution from the Gulf Stream (yellow arrow, Figure 2 bottom; Richardson & Finlen 1967), with currents extending west-to-east in the Canyon (green arrow, Figure 2 bottom). The occasionally turbid currents within the Great Bahama Canyon deposit a large sediment fan as they merge with the Antilles Current of the Atlantic Ocean ~65km to the north-northeast (red arrow, Figure 2 bottom) (Andrews 1970). Although the seasonal variability of these currents has not been established, their overall direction and extent offer the potential for a vicariant separation between Grand Bahamas and Eleuthera.

2.4 | Collection, extraction, and amplification of samples

During 2014, we sampled fin clips from 57 adult bonefish collected by rod and reel from three locations on the west and south sides of Eleuthera Island (EWB, ES1, ES2; Table 1) and one from the Atlantic Ocean (east) side (EEB; Table 1). Muscle samples were also gathered from 60 adults at Grand Bahama Island, with 30 taken from the south (Ocean)

side (GSB) and 30 from the north (Flats) side (GNB) (total N=117). Sampling was in compliance with the U.S. National Research Council Guide for the Care and Use of Laboratory Animals (https://grants.nih.gov/grants/olaw/Guide-for-the-Care-and-use-of-laboratory-animals.pdf).

We also collected 30 individuals of queen conch from the south side of Grand Bahama Island (GSC), and 30 from the Flats (or west) side of Eleuthera Island (EWC) (Table 1, Figure 1 bottom). Fin clips and muscle plugs were preserved on WHATMAN® FTA® Classic DNA cards, with genomic DNA subsequently extracted using a GENTRA® Puregene DNA Purification Kit and stored in hydrating solution (per manufacturer instructions). The quality of DNA was visualized on 2% agarose gels, then quantified using a Qubit fluorometer (Thermo-Fisher Scientific, Inc.). Subsequent library preparation followed previously published protocols (Peterson et al. 2012, as modifed in Bangs et al. 2018).

Restriction digests were performed using 1μg of genomic DNA with *Pst*I (5'-CTGCAG-3') and *Msp*I (5'-CCGG-3') in CutSmart buffer (New England Biosciences) for 20 hours at 37°C, after first optimizing restriction enzyme and size selection choices using an *in silico* approach (FRAGMATIC; Chafin et al. 2018). Digests were visualized on 2% agarose gels, purified using AMPure XP beads, and quantified with a Qubit fluorometer. Digested samples were normalized to 0.1μg of DNA and subsequently ligated with barcoded Illumina adaptors, using custom oligos (Peterson et al. 2012). Inline barcodes were designed to differ by at least 2 bases so as to minimize mis-assignments. Pooled ligations were size-selected at 350-400bps using the Pippin Prep automated size fractionator (Sage Sciences).

Adaptors were extended by PCR (10 cycles) with Phusion high-fideltiy DNA polymerase

(New England Biosciences). Size distribution and successful amplification of purified

libraries was then confirmed on the Agilent 2200 TapeStation. Indexed libraries were

multiplexed with 100-bp single-end sequencing on the Illumina HiSeq 2500 lane (Genomics

and Cell Characterization Core Facility, University of Oregon/Eugene).

2.5 | Processing and assembly of ddRAD data

Raw ~100bp reads were demultiplexed and processed in the PYRAD pipeline (Eaton 2014)

using a 90% identity threshold for clustering homologous reads. Read clusters representing

putative loci within individuals were excluded when sequencing depth was <10. Following

assembly, additional loci were discarded if they exhibited: 1) <50% presence across

individuals; or 2) >10 consensus N bases. We additionally removed potential paralogs or

over-merged loci by excluding those having: 1) >2 alleles in any individual; 2) >60%

heterozygosity per site among individuals; 3) >10 heterozygous sites per consensus

sequence. Further data processing was performed using custom scripts (available at:

https://github.com/tkchafin/scripts).

2.6 | Detection and spatial structure of local selection

We first sub-sampled all SNPs to a maximum of one per locus. We then utilized a Bayesian

method (BAYESCAN v.2.1, Foll & Gaggiotti 2008) with a burn-in period of 50,000 followed

by 100,000 iterations to generate a candidate list of potential loci under selection. We then

used F_{ST}-etimates to derive three additional datasets per taxon: (1) All unlinked SNPs

(=ALL); (2) SNPs >2 standard deviations (sd) above mean F_{ST} (=SD2); and (3) SNPs >3 sd

above the mean (=SD3).

We employed multiple methods to examine population structure for both species

across all three datasets. Hierarchical iterative clustering analysis (STRUCTURE v.2.2.4;

Pritchard et al. 2000) was run in parallel (Besnier & Glover 2013) on the Jetstream cloud

computing platform (Hancock et al. 2019) using the GNU Parallel shell tool (Tange 2011).

Structure was run for 250,000 repetitions with a burn-in period of 100,000, and with

potential clusters for K-values ranging from 1-20. Optimal K for each dataset was

ascertained using the delta-K method (Evanno et al. 2005), as implemented and visualized

in the CLUMPAK pipeline (Kopelman et al. 2015). Differentiation among sampling localities

was also examined with with a cross-validation approach (Discriminant Analysis of Principal

Components, DAPC) in the R package ADEGENET (Jombart 2008, Jombart et al. 2008). We

defined the number of retained principal component axes as that which minimized the

root-mean-square error (RMSE) of assignment for 10% of randomly selected individuals

(with the remaining 90% serving as a training set), and N=30 replicates per level of PC

retention. We specified the number of DF axes retained as equal to the number of

populations minus one.

2.7 | Asymmetric migration among populations

Physical processes such as water current and tidal fluxes often create patterns of

asymmetric gene flow among populations, and consequently impact patterns of genetic

diversity. Given our initial hypothesis, we were interested in gauging the amount and

direction of gene flow among the six study populations of bonefish, and two of queen

conch. In contrast to STRUCTURE, which employs a Bayesian probabilistic model to assign

individuals to ancestry clusters, we instead estimated the posterior probability of individual

migratory histories, thereby allowing the direction and magnitude of dispersal among

populations to be dissected.

To do so, we employed a revised software optimized for large SNP datasets (i.e.,

BA3-SNPs, Mussmann et al. 2019). We first tuned MCMC acceptance parameters using an

automated iterative procedure, then generated posterior estimates of migration rates

using 250,000 iterations with a burn-in period of 50,000. We further tested the hypothesis

of source-sink dynamics by computing net emigration rates as a sum of immigration rates

subtracted from summed emigration rates (Andreasen et al 2012), with positive numbers

indicating a genetic source population whereas negative values indicated a sink population.

3 | RESULTS

After filtering, we obtained 56,046 loci for bonefish *Albula vulpes*, of which 68.7%

(N=36,206) were parsimoniously informative (Table 2). Average number of loci per

individual was 35,443, at a mean coverage depth of 77x. Subsampling only unlinked SNPs

produced a matrix of 25,205 SNPs (=ALL), from which datasets SD2 (123 SNPs) and SD3 (79

SNPs) were derived, respectively.

For queen conch *Lobatus gigas*, 30,697 loci were available post-filtering, and

represented 64,864 SNPs, of which 68.4% (N=44,334) were parsimony informative.

Individuals were sequenced at a coverage depth of 58.6x, averaging 33,409 loci per

individual (Table 2). Subsampled unlinked SNPs totaled 19,716 (=ALL), from which datasets SD2 (88 SNPs) and SD3 (51 SNPs) were obtained.

3.1 | Spatial structure in Bonefish and Queen Conch

STRUCTURE runs for *A. vulpes* employing the ALL dataset (=neutral differentiation) revealed scant separation among sample sites (Figure 3). The most highly-supported models for SD2 and SD3 were K=2 and K=3 groupings, both of which depicted GSB (Grand Bahama, south side) as clustereing with EWB (Eleuthera, west side) and ES2 (south and west side of Eleuthera). The second cluster placed GNB (Grand Bahama north side) with EEB (Eleuthera east or Atlantic side) and ES1 (south and west side of Eleuthera) (Figure 3). The K=3 STRUCTURE run with SD2 and SD3 datasets was a bit mixed, with GSB and EWB as one group with ES1, ES2, and EEB as a second, and GNB as the third. Subsequent K-groups were decidedly more mixed which rendered them indecipherable.

In our DAPC analyses, we retained N=45, 25, and 15 PC axes for bonefish (ALL, SD2, and SD3 datasets, respectively), corresponding to 73%, 77%, and 66% of retained variance. In all cases, the number of retained DF axes (N=5) was equal to the number of populations minus one. Clustering of the ALL dataset for bonefish (Figure 4 left) displayed considerable overlap among sites (per STRUCTURE plot, Figure 3). Three populations with scant separation (i.e., ES1, ES2 and EEB) were grouped within the center of the graph. Three others were somewhat more peripheral and reflected greater within-group variability (i.e., GNB, GSB, and EWB). The same type of analysis, but employing the SD2 dataset (Figure 4 middle), separated GNB, EWB, EEB, and ES1 without overlap, while grouping GSB with ES2. These

relationships were maintained in the DAPC clustering of the SD3 dataset, but with more overlap between the previously separated EEB and ES1, whereas ES2 and GSB maintained their linkage. GNB and EWB also remained distinct (Figure 4 right).

The STRUCTURE analysis of *L. gigas* populations using the ALL dataset revealed little genetic differentiation, with mean F_{ST} =0.0129. However both EWC and GSC showed significant separation when both SD2 and SD3 datasets were evaluated (Figure 5). DAPC for conch with cross-validation retained N=15, 10, and 5 PC axes (ALL, SD2, and SD3, datasets, respectively). This yielded 16.7%, 44%, and 50.5% of PC variance. However, there is N=1 DF axis for conch, in that the number of DAPC discriminant functions is N_populations-1. The proportion variance explained by this single axis is irrelevant in that it is 100% by definition. The DAPC clustering of the ALL dataset (Figure 6 left) again showed considerable overlap between the two islands, whereas each separated with considerably less overlap when SD2 was analysed, and became even more distinct based upon SD3 (Figure 6, middle, right).

3.2 | Asymmetric migration

The BAYESASS3 analysis of bonefish revealed highly variable and asymmetric migration rates, with EEB and EWB depicted as source populations, with outward migration rates to both Grand Bahama and flats-side Eleuthera populations. This varied from 8.5-17.9% for EEB and 7.5-14.6% for EWB (Table 3; Figure 7). Notably, the remaining populations lacked any strong signal of outward migration, and all showed negative rates of net emigration (Figure 8), suggesting a sink dynamic. Results for *L. gigas* found very high demographic

independence among populations, with 'self-migration' (e.g., recruitment) rates estimated at >98% for both Grand Bahama and Eleuthera populations (Table 3).

4 | DISCUSSION

Populations are frequently classified as to the manner by which they are replenished. In this sense, closed populations are defined as having recruitment via local reproduction, whereas open populations are independent of source (Pinsky et al. 2012). This dichotomy underscores several ecological, evolutionary, and conservation-based research agendas: Exploitation or protection (Yau et al. 2014, Krueck et al. 2017); life history and taxonomy (Mora & Sale 2002, Baco et al. 2016); recovery from disturbance (Bell et al. 2014); and extent to which populations are locally adapted (Jorde et al. 2015, Salles et al. 2016, Segovia et al. 2017). It also has obvious implications for how populations are managed. The advent of genomic techniques has provided considerable resolution, particularly with regard to the estimation of local adaptation (Reisser et al. 2014).

4.1 | F_{ST} outliers for bonefish

For non-model organisms (as herein), two approaches are most often utilized: The first identifies loci that display unusually high genetic differentiation among populations (i.e., F_{ST} outlier approach), whereas the second (i.e., gene-environment association) seeks correlations between the allele frequencies of populations and their local environments. We employed the first approach in this study.

Our analysis failed to group samples of bonefish *Albula vulpes* into geographically congruent groups on either side of the Canyon, suggesting instead a distribution more panmictic in format (per STRUCTURE analysis and DAPC plot for the ALL data). Furthermore, STRUCTURE plots for K=2 and K=3 groupings (Figure 3) deviated but little. For K=2, both SD2 and SD3 datasets linked the south side of Grand Bahama Island (GSB) with the west side of Eleuthera (EWB) and with one (of two) ES locations (i.e., ES2). This, in turn, suggests a level of connectivity across the Great Bahama Canyon, a conjecture supported by Figure 4. However, the north side of Grand Bahama Island (GNB) grouped with the eastern (Atlantic) side of Eleuthera (EEB) and with the second ES site in the K=2 grouping (i.e., ES1). This relationship separates in K=3, with GSB remaining linked with EWB, whereas EEB and ES2 separate.

These relationships are difficult to interpret biogeographically, particularly given what we know of geomorphic and current patterns in the northeast Basin (Figure 2). A more parsimonious explanation would point to the considerable mobility of adult bonefish superimposed onto a lack of site specificity. Although dispersal pathways for bonefish are relatively unknown in the northeast Basin, their quantification would identify Bahamian populations that serve as sinks for bonefish spawning under investigation at South Abaco Island (Lombardo et al. 2020; Figure 1). In this sense, our results provide reasonable hypotheses.

4.2 | F_{ST} outliers for queen conch

A similar analysis did indeed discriminate sessile populations of queen conch *Lobatus gigas* separated by the Great Bahama Canyon (a potentially vicariant barrier). This is aptly demonstrated in Figure 5, where the south side of Grand Bahama Island and the west side of Eleuthera Island are clearly discrete one from another. This condition is reinforced by the DAPC analysis (Figure 6), where distributions for SD2, and SD3 datasets consistently diverge from that depicted for the ALL dataset. F_{ST} values for these two populations are not significantly different when computed from the ALL data, but are so when compiled from the SD2 and SD3 datasets (Table 3). Thus, while neutral loci demonstrated little differntiation across the study area, a smaller proportion reflected strong differentiation consistent with local adaptation. This scenario has also been manifested in other marine species similarly investigated using SNP analyses (Cure et al. 2017, Thomas et al. 2017).

4.3 | Potential problems with F_{ST} outlier analyses

One potential issue with detecting selection via departure from neutrality is that each locus may manifest an independent response that can hinge upon population structure and demography. This is particularly problematic when the average level of differentiation among populations is relatively high (Hoban et al. 2017). However, pairwise F_{ST} values for bonefish are neither elevated nor significant (Table 3), suggesting this may not apply in our situation. An additional concern is that some loci may demonstrate false signatures of selection, such as when small populations on the leading edge of an expansion contribute disproportionately to observed frequency differences. Again, our study populations are already established and thus the concept of a leading edge is not a viable consideration.

Similarly, cryptic hybridization and introgression can invoke confounding effects as well, but again, these are not apparent in our data.

4.4 | Asymmetrical migration in bonefish and queen conch

The structure of the physical environment can promote differential movements in aquatic organisms, and thus promote a metapopulation structure within a species (i.e., a source–sink dynamic). This in turn can skew estimates of genetic differentiation among populations, and consequently promote erroneous management decisions, particularly when active management is based on an economic model, such as when the organisms are harvested. We employed genetic estimates of population migration to ascertain potential source–sink considerations in our study organisms, and to determine if indeed movements between them were asymmetrical.

Bonefish in our study indeed demonstrated strong signals of source-sink dynamics across islands. The two Eleuthera populations (EEB and EWB) seemingly act as genetic sources, as evidenced by their high net emigration rates derived from BayesAss3 (Figure 8). Their genetic differentiation may suggest the presence of discrete genetic stocks on each side of the island, reflecting either spatial or temporal asynchrony with regards to spawning. Given a tendency for bonefish to move into deeper waters for spawning (Danylchuk et al 2011; Lombardo et al. 2020), the differentiation between the two Eleuthera populations may suggest spawning site-fidelity (potentially within deeper water off Abaco Island; per Lombardo et al. 2020), with increased intermingling subsequently due to the export of larvae to other Bahamian areas. The latter may also explain the cryptic

intra-site structure as recorded in the assignment tests of *Fst* outlier SNP datasets (Figure 3).

Populations of conch, on the other hand, were highly demographically independent. These results are apparent in both estimates of contemporary migration (BA3-SNPs), and from estimates of population structure using highly-differentiated SNPs.

4.5 | Dispersal and connectivity within the Caribbean Basin

We hypothesized that the population demographics of our study species would be impacted by the currents passing through the Grand Bahama Canyon. Prior to test, we sought data from other species that might potentially shed light on this conjecture.

Jackson et al. (2014) investigated genetic differentiation among 19 spawning aggregations of Nassau grouper *Epinephelus striatus* in the Basin. The life history of this species parallels that of bonefish, in that spawning aggregations are cued by water temperature and phases of the moon over a 3-month period. As with bonefish, breeding aggregations develop some distance from individual home ranges, with larvae drifting for a similar duration (~40 days) before settling. Four Basin-wide genetic groups were identifed, to include the Bahamas, but with little within-group differentiation, again in parallel with that found for bonefish (per Wallace & Tringali 2016). A major barrier to larval dispersal was the separation of the Basin into eastern and western regions (Jackson et al. 2014, Figure 3). These results implicate the Great Bahama Canyon (Figure 1 top) as a vicariant barrier, similar to what was found herein with conch.

In a second study, Kool et al. (2010) coupled results from a matrix-based projection model with those based upon a bio-physical approach to examine genetic dispersal among coral reef patches within the Basin. Allele frequencies were found to shift along the length of the Bahamas, with the sharpest demarcation occurring at the separation between Grand Bahama Island and the southern Bahamas. This separation was not recognized in the original publication, but indeed juxtaposes with the presence of the Grand Bahama Canyon (Figure 1). Again, the results from Kool et al. (2010) promote the Canyon as a major vicariant feature in the region.

5 | CONCLUSIONS

An understanding of genetic connectivity within exploited species is a crucial conservation aspect, in that once designated, appropriate geographic units can then be subsequently managed so as to avoid over-exploitation. This includes the establishment of quotas for 'take,' as well as the delineation boundaries for marine protected areas (MPAs). Furthermore, an understanding of Basin-wide linkages (or lack thereof) among populations of commercially exploited species (as herein) promotes the realization among numerous stakeholders that fisheries management is a complex and continually evolving geopolitical problem.

Conservation and management practices for targeted species can be greatly enhanced if indeed they are supplemented with adaptive (i.e., ecologically relevant) data (Allendorf 2017). For instance, adaptive-based management units can potentially be defined such that, in high gene flow species, may represent subsets within more traditional

(neutral-based) units (Funk et al. 2012). Outlier SNPs could also identify the results of illegal

fishing (such as for conch), and also serve to define stock boundaries and potential

management units (Milano et al. 2014). For bonefish, it may require the protection of

larger areas that can accommodate those fairly precise oceanographic conditions that

define where they spawn and which provide the predictable and stable dispersal

mechanism required by pelagic larvae.

These aspects cut to the definition of conservation genomics, which is defined (in

the narrow sense) as those approaches that differ conceptually and quantitatively from

traditional genetics. This, in turn, serves to amplify genome-wide signals of adaptation, and

promotes answers for those questions impossible to address using legacy approaches

alone (Garner et al. 2016). Importantly, such data can provide genomic evidence for

potential heat tolerance (or the lack thereeof), an issue that particularly impacts sessile

marine invertebrates (Thomas et al., 2017). Such issues loom large as oceans acidify and

their heat profiles expand in lockstep with global climate change (Hurd et al., 2018).

ACKNOWLEDGMENTS

Funding was provided by the following generous endowments to the University of

Arkansas: The Bruker Professorship in Life Sciences (MRD), a Distinguished Doctoral

Fellowship (TKC), and the 21[st] Century Chair in Global Change Biology (MED). We gratefully

acknowledge the Cape Eleuthera Institute (CEI) and The Island School for logistical support

and field assistance. Sampling was supported under CEI collecting permits. Sampling was

supported by grants from Patagonia's World Trout Initiative, the Charles A. and Anne

Morrow Lindbergh Foundation, the Baldwin Foundation, as well as generous personal donations from B. Hallig, T. Rice, and J. Spring. Computational analyses were supported by the Arkansas Economic Development Commission (Arkansas Settlement Proceeds Act of 2000), the Arkansas High Performance Computing Center (AHPCC), and an XSEDE Research Allocation (TG-BIO160065) to access the Jetstream cloud. Sequence data were generated at the Genomics and Cell Characterization Core Facility, University of Oregon/ Eugene. Raw datasets are available via NCBI SRA (accession added upon acceptance).

AUTHOR'S CONTRIBUTIONS

MRD/ JEC/ DPP/ MED designed the study; JEC/ DPP contributed to fieldwork; MRD/ TKC/ MED contributed to lab work; TKC performed bioinformatic work and statistical analyses; MRD/ TKC/ MED interpreted results and generated figures; MRD/ TKC/ MED drafted the manuscript; All authors gave final approval for publication.

ORCID

Marlis R. Douglas https://orcid.org/0000-0001-6234-3939

Tyler K. Chafin https://orcid.org/0000-0001-8687-5905

Michael E. Douglas https://orcid.org/0000-0001-9670-7825

DATA ACCESSIBILITY

ddRAD SNP loci and scripts can be found at: https://github.com/tkchafin/alb_conch_data

REFERENCES

Adams, A. J., & Cooke, S. J. (2015). Advancing the science and management of flats

    fisheries for bonefish, tarpon, and permit. *Environmental Biology of Fishes*, 98,

    2123–2131 (doi:10.1007/s10641-015-0446-9)

Allendorf, F. W. (2017). Genetics and the conservation of natural populations: Allozymes to

    genomes. *Molecular Ecology*, 26, 420–430 (doi:10.1111/mec.13948)

Andreasen, A. M., Stewart, K. M., Longland, W. S., Beckmann, J. P., & Forister, M. L. (2012).

    Identification of source-sink dynamics in mountain lions of the Great Basin.

    Molecular Ecology, 21, 5689–5701 (doi: 10.1111/j.1365-294X.2012.05740.x)

Andrews, J. E. (1970). Structure and sedimentary development of the outer channel of the

    Great Bahama Canyon. *Geological Society of America Bulletin*, 81, 217–225

    (10.1130/0016-7606(1970)81[217:SASDOT]2.0.CO;2)

Andrews, J. E., Shepard, F. P., & Hurley, R. J. (1970). Great Bahama Canyon. *Geological

    Society of America Bulletin*, 81. 1061–1078 (doi:10.1130/0016-

    7606(1970)81[1061:GBC]2.0)

Baco, A. R., Etter, R .J., Ribeiro, P. A., & Von Der Heyden, S., Beerli, P. & Kinlan, B. P. (2016).

    A synthesis of genetic connectivity in deep-sea fauna and implications for marine

    reserve design. *Molecular Ecology*, 25, 3276–3298 (doi:10.1111/mec.13689)

Ballantine, D. L., & Appeldoorn, R. S. (1983). Queen conch culture and future prospects in

Puerto Rico. *Proceedings of the Gulf and Caribbean Fisheries Institute,* 35, 57–63

(https://www.researchgate.net/publication/283924228)

Bangs, M. R., Douglas, M. R., Mussmann, S. M., & Douglas, M. E. (2018). Unraveling historic

introgression and resolving phylogenetic discord within *Catostomus* (Pisces:

Catostomidae). *BMC Evolutionary Biology,* 18, 86 (doi:10.1186/s12862-018-1197-y)

Bell, J. J., Smith, D., Hannan, D., Haris, A., Jompa, J., & Thomas, L. (2014). Resilience to

disturbance despite limited dispersal and self-recruitment in tropical barrel

sponges: Implications for conservation and management. *PLoS ONE*, 9, e91635

(doi:10.1371/journal.pone.0091635)

Bentzen, P., & Bradbury, I. R. (2016). Don't bet against the natal homing abilities of marine

fishes. *Molecular Ecology*, 25, 2691–2692 (doi:10.1111/mec.13591)

Besnier, F., & Glover, K. A. (2013). PARALLELSTRUCTURE: An R Package to distribute parallel

runs of the population genetics program STRUCTURE on multi-core computers. PLoS

ONE 8(7),   e70651 (doi:10.1371/journal.pone.0070651)

Bonanomi, S., Therkildsen, N. O., Retzel, A., Hedeholm, R. B., Pedersen, M. W., Meldrup, D.,

... Nielsen, B. C. (2016). Historical DNA documents long-distance natal homing in

marine fish. *Molecular Ecology*, 25, 2727–2734 (doi:10.1111/mec.13580)

Brownell, W. N., & Stevely, J. M. (1981). The biology, fisheries and management of the

   queen conch, *Lobatus gigas*. *Marine Fish Review*, 43, 1-12

   (spo.nmfs.noaa.gov/mfr437/mfr4371.pdf)

Chafin, T. K., Martin, B. T., Mussmann, S. M., Douglas, M. R., & Douglas, M. E. (2018).

   FRAGMATIC: in silico locus prediction and its utility in optimizing ddRADseq

   projects. *Conservation Genetics Resources*, 10, 325-328 (doi: 10.1007/s12686-017-

   0814-1)

Christie, M. R., Johnson, D. W., Stallings, C. D., & Hixon, M. A. (2010). Self-recruitment and

   sweepstakes reproduction amid extensive gene flow in a coral-reef fish. *Molecular

   Ecology*, 19, 1042–1057 (doi:10.1111/j.1365-294X.2010.04524.x)

Cowen, R. K., Lwiza, K. M. M., Sponaugle, S., Paris, C. B., & Olsen, D. B. (2000). Connectivity

   of marine populations: Open or closed? *Science*, 287, 857–59

   (doi:10.1126/science.287.5454.857)

Cowen R. K., & Sponaugle, S. (2009). Larval dispersal and marine population connectivity.

   Annual Review of Marine Science, 1, 443–466

   (doi:10.1146/annurev.marine.010908.163757)

Cumming, R. A., Nikula, R., Spencer, H. G., & Waters, J. M. (2016). Trans-Tasman genetic

   connectivity in the intertidal air-breathing slug *Onchidella nigricans*. *Marine

   Ecology Progress Series*, 562, 93–100, 2016 (doi:10.3354/meps11926)

Cure, K., Thomas, L., Hobbs, J-P. A., Fairclough, D. B., & Kennington, W. J. (2017). Genomic signatures of local adaptation reveal source-sink dynamics in a high gene flow fish species. *Scientific Reports*, 7, 8618 (doi:10.1038/s41598-017-09224-y)

Danylchuk, A. J., Cooke, S. J,, Goldberg, T. L., Suski, C. D. , Murchie, K. J., Danylchuk, S. E., ... Philipp, D. P. (2011). Aggregations and offshore movements as indicators of spawning activity of bonefish (*Albula vulpes*) in The Bahamas. *Marine Biology*, 158, 1981–1999 (doi:10.1007/s00227-011-1707-6)

Domingues, C. P., Creer, S., Taylor, M. I., Queiroga, H., & Carvalho, G. R. (2010). Genetic structure of *Carcinus maenas* within its native range: Larval dispersal and oceanographic variability. *Marine Ecology Progress Series*, 410, 111–123 (doi:10.3354/meps08610)

Doerr, J. C., & Hill, R. L. (2013). Home range, movement rates, and habitat use of queen conch, *Strombus gigas*, in St. John, U.S. Virgin Islands. *Caribbean Journal of Science*, 47, 251-259 (doi:10.18475/cjos.v47i3.a13)

Eaton, D. A. R. (2014). PʏRAD: Assembly of de novo RADseq loci for phylogenetic analyses. *Bioinformatics*, 30, 1844–1849 (doi:10.1093/bioinformatics)

Ellingson, R. A., & Krug, P. J. (2015). Reduced genetic diversity and increased reproductive isolation follow population-level loss of larval dispersal in a marine gastropod. *Evolution*, 70, 18–37 (doi:10.1111/evo.12830)

Evanno, G., Regnaut, S., & Goudet, J. (2005). Detecting the number of clusters of

individuals using the software Structure: A simulation study. *Molecular Ecology*, 14,

2611–2620 (doi: 0.1111/j.1365-294X.2005.02553.x)

Feldheim, K. A., Gruber, S. H., DuBattista, J. D., Babcock, E. A., Kessel, S. T., Hendry, A. P, …

Chapman, D. D. (2014). Two decades of genetic profiling yields first evidence of

natal philopatry and long-term fidelity to parturition sites in sharks. *Molecular*

Ecology, 23, 110–117 (doi:10.1111/mec.12583)

Foll, M., & Gaggiotti, O. (2008). A genome-scan method to identify selected loci

appropriate for both dominant and codominant markers: a Bayesian perspective.

Genetics, 180, 977–993 (doi:10.1534/genetics.108.092221)

Funk, W. C., McKay, J. K., Hohenlohe, P. A., & Allendorf, F. W. (2012). Harnessing genomics

for delineating conservation units. *Trends in Ecology and Evolution*, 27, 489–496

(doi:10.1016/j.tree.2012.05.012)

Garner, B. A., Hand, B. K., Amish, S. J., Bernatchez, L., Foster, J. T., Miller, K. M., … Luikart,

G. (2016). Genomics in conservation: Case studies and bridging the gap between

data and application. *Trends in Ecology and Evolution*, 31, 81–83

(doi:10.1016/j.tree.2015.10.009)

Gomes, I., Peteiro, L. G., Albuquerque, R., Nolasco, R., Dubert, J., Swearer, S. E., &

Queiroga, H. (2016). Wandering mussels: Using natural tags to identify connectivity

patterns among Marine Protected Areas. *Marine Ecology Progress Series*, 552, 159–

176 (doi:10.3354/meps11753)

Gould, A. L., & Dunlap, P. V. (2017). Genomic analysis of a cardinalfish with larval homing

potential reveals genetic admixture in the Okinawa Islands. *Molecular Ecology*, 26,

3870–3882 (doi:10.1111/mec.14169)

Hameed, S. O., White, J. W., Miller, S. H., Nickols, K. J., & Morgan, S. G. (2016). Inverse

approach to estimating larval dispersal reveals limited population connectivity

along 700 km of wave-swept open coast. *Proceedings of the Royal Society of London

Part B Biological Sciences*, 283, 20160370 (doi:10.1098/rspb.2016.0370)

Hancock, D. Y., Stewart, C. A., Vaughn, M., Fischer, J., Lowe, J. M., Turner, G., … Snapp-

Childs, W. (2019). Jetstream—Early operations performance, adoption, and

impacts. *Concurrency and Computation: Practice and Experience*, 31, e4683 (doi:

10.1002/cpe.4683)

Herrera, M., Nanninga, G. B., Planes, S., Jones, G. P., Thorrold, S. R., Saenz-Agudelo, P., …

Berumen, M. L. (2016). Seascape and life-history traits do not predict self-

recruitment in a coral reef fish. *Biology Letters*, 12, 20160309

(doi:10.1098/rsbl.2016.0309)

Hoban, S., Kelley, J. L., Lotterhos, K. E., Antolin, M. F., Bradburd, G., Lowry, D. B., …

Whitlock, M. C. (2017). Finding the genomic basis of local adaptation: Pitfalls,

practical solutions, and future directions. *American Naturalist*, 188, 379–397

(doi:10.1086/688018)

Humston, R., Ault, J. S., Larkin, M. F , & Luo, J. (2005). Movements and site fidelity of the

bonefish *Albula vulpes* in the northern Florida Keys determined by acoustic

telemetry. *Marine Ecology Progress Series*, 291, 237–248

(doi:10.3354/meps291237)

Hurd, C.L., Lenton, A., Tilbrook, B., & Boyd, P. W. (2018). Current understanding and

challenges for oceans in a higher-CO2 world. *Nature Climate Change*, 8, 686–694

(doi:10.1038/s41558-018-0211-0)

Jackson, A. M., Semmens, B. X., Sadovy de Mitcheson, Y., Nemeth, R. S., Heppell, S. A.,

Bush, P. G., … Bernardi, G. (2014). Population structure and phylogeography in

Nassau Grouper (*Epinephelus striatus*), a mass-aggregating marine fish. *PLoS ONE*,

9, e97508 (doi:10.1371/journal.pone.0097508)

Jasonowicz, A. J., Goetz, F. W., Goetz, G. W., & Nichols, K. M. (2017). Love the one you're

with: Genomic evidence of panmixia in the sablefish (*Anoplopoma fimbria*).

Canadian Journal of Fisheries and Aquatic Science, 74, 377–387 (doi:10.1139/cjfas-

2016-0012)

Jombart, T. (2008). ADEGENET: An R package for the multivariate analysis of genetic

markers. *Bioinformatics*, 24, 1403–1405 (doi:10.1093/bioinformatics/btn129)

Jombart, T., Devillard, S., Dufour, A-B., & Pontier, D. (2008). Revealing cryptic spatial

patterns in genetic variability by a new multivariate method. *Heredity*, 101, 92–103

(doi:10.1038/hdy.2008.34)

Jorde, P. E., Søvik, G., Westgaard, J-I., Albretsen, J., André, C., Hvingel, C., ... Jørstad, K. B.

(2015). Genetically distinct populations of northern shrimp, *Pandalus borealis*, in

the North Atlantic: Adaptation to different temperatures as an isolation factor.

Molecular Ecology, 24, 1742–1757 (doi:10.1111/mec.13158)

Kensington, W. J., Keron, P. W., Harvey, E. S., Wakefield, C.B., Williams, C. B., Halafihi, A. J.,

... Newman, S. J. (2017). High intra-ocean, but limited inter-ocean genetic

connectivity in populations of the deep-water oblique-banded snapper

Pristipomoides zonatus (Pisces: Lutjanidae). *Fisheries Research*, 193, 242–249

(doi:10.1016/j.fishres.2017.04.015)

Kool, J. T., Paris, C. B., Andréfouët, S., & Cowen, R. K. (2010). Complex migration and the

development of genetic structure in subdivided populations: An example from

Caribbean coral reef ecosystems. *Ecography*, 33, 597-606 (doi:10.1111/j.1600-

0587.2009.06012.x)

Kopelman. N. M., Mayzel, J., Jackobsson, M. J., Rosenberg, N. A., & Mayrose, I. (2015).

CLUMPAK: a program for identifying clustering modes and packaging population

structure inferences across K. *Molecular Ecology Resources*, 15, 1179–1191 (doi:

10.1111/1755-0998.12387)

Kough, A. S., Cronin, H., Skubel, R., Belak, C. A., & Stoner, A. W. (2017). Efficacy of an

   established marine protected area at sustaining a queen conch *Lobatus gigas*

   population during three decades of monitoring. *Marine Ecology Progress Series*,

   573, 177–189 (doi:10.3354/meps12163)

Krueck, N. C., Ahmadia, G. N., Green, A., Jones, G. P., Possingham, H. P., Riginos, C., ...

   Mumby, P. J. (2017). Incorporating larval dispersal into MPA design for both

   conservation and   fisheries. *Ecological Applications*, 27, 925–941

   (doi:10.1002/eap.1495)

Lombardo, S. M., Adams, A. J., Danylchuk, A. J., Luck, C. A., & Ajemian, M. J. (2020). Novel

   deep-water spawning patterns of bonefish (*Albula vulpes*), a shallow water fish.

   Marine Biology, 167, 187 (doi:10.1007/s00227-020-03799-3)

Machkour-M'Rabet, S., Cruz-Medina, J., García-De León, F. J. , De Jesús-Navarrete. A.,

   Hénaut, Y. (2017). Connectivity and genetic structure of the queen conch on the

   Mesoamerican Reef. *Coral Reefs*, 36, 535–548 (doi:10.1007/s00338-017-1551-3)

Martinez, E., Buonaccorsi, V., Hyde, J. R., & Aguilar, A. (2017). Population genomics reveals

   high   gene flow in grass rockfish (*Sebastes rastrelliger*). *Marine Genomics*, 33, 57–

   63 (doi:10.1016/j.margen.2017.01.004)

McKeon, N. J., Hauser, L., & Shaw, P. W. (2017). Microsatellite genotyping of brown crab

   Cancer pagurus reveals fine scale selection and 'non-chaotic' genetic patchiness

within a high gene flow system. *Marine Ecology Progress Series*, 566, 91–103

(doi:10.3354/meps12044)

Milano, I., Babbucci, M., Cariani, A., Atanassova, M., Bekkevold, D., Carvalho, G. R., ...

Bargelloni, L. (2014). Outlier SNP markers reveal fine-scale genetic structuring

across European hake populations (*Merluccius merluccius*). *Molecular Ecology*, 23,

118-135 (doi:10.1111/mec.12568)

Miller, K. J., Mundy, C. N., & Mayfield, S. (2014). Molecular genetics to inform spatial

management in benthic invertebrate fisheries: A case study using the Australian

greenlip abalone. *Molecular Ecology*, 23, 4958–4975 (doi:10.1111/mec.12914)

Mitton, J. B., Berg, C. J. Jr., & Orr, K. S. (1989). Population structure, larval dispersal, and

gene flowin the Queen Conch, *Lobatus gigas*, of the Caribbean. *Biological Bulletin*,

177, 356-362 (doi:10.2307/1541595)

Mora, C., & Sale, P. F. (2002). Are populations of coral reef fish open or closed? *Trends in

Ecology and Evolution*, 17, 422-428 (doi:10.1016/S0169-5347(02)02584-3)

Morvezen, R., Charrier, G., Boudry, P., Chauvaud, L., Breton, F., Strand, Ø., & Laroce J.

(2016). Genetic structure of a commercially exploited bivalve, the great scallop

Pecten maximus, along the European coasts. *Conservation Genetics*, 17, 57–67

(doi:10.1007/s10592-015-0760-y)

Murchie, K. J., Cooke, S. J., Danylchuk, A. J., Danylchuk, S. E., Goldberg, T., Suski, C., &

Philipp, D.P. (2013). Movement patterns of bonefish (*Albula vulpes*) in tidal creeks

and coastal waters of Eleuthera, The Bahamas. *Fisheries Research*, 147, 404-412

(doi:10.1016/j.fishres.2013.03.019)

Murchie, K. J., Schultz, A. D., Stein, J. A., Cooke, S. J., Lewis, J., Franklin, J., ... Philipp, D.P.

(2015). Defining adult bonefish (*Albula vulpes*) movement corridors around Grand

Bahama in the Bahamian Archipelago. *Environmental Biology of Fishes*, 98, 2203–

2212 (doi:10.1007/s10641-015-0422-4)

Mussmann, S. M., Douglas, M. R., Chafin, T. K., & Douglas, M. E. (2019). BA3-SNPs:

Contemporary migration reconfigured in BayesAss for next-generation sequence

data. *Methods in Ecology and Evolution*, 10, 1808-1813 (doi:10.1111/2041-

210X.13252)

Ottmann, D., Grorud-Colvert, K., Sard, N. M., Huntington, B. E., Banks, M. A., & Sponaugle,

S. 2016). Long-term aggregation of larval fish siblings during dispersal along an open

coast. *Proceedings of the National Academy of Sciences U.S.A.,* 113, 14067–14072

(doi:10.1073/pnas.1613440113)

Peterson, B. K., Weber, J. N., Kay, E. H., Fisher, H. S., & Hoekstra, H. E. (2012). Double

digest RADseq: An inexpensive method for de novo SNP discovery and genotyping

in model and non-model species. *PLoS ONE,* 7, e37135

(doi:10.1371/journal.pone.0037135)

Pinsky, M. L,, Palumbi, S. R., Andrefouet, S., & Purkis, S. J. (2012). Open and closed

seascapes: Where does habitat patchiness create populations with high fractions of

self-recruitment? *Ecological Applications*, 22, 1257–1267 (doi:10.1890/11-1240.1)

Pritchard, J. K., Stephens, M., & Donnelly, P. (2000). Inference of population structure using

multilocus genotype data. *Genetics*, 155, 945–959 (PMC1461096)

Reisser, C. M.O., Bell, J. J., & Gardner, J. P.A. (2014). Correlation between pelagic larval

duration and realised dispersal: Long-distance genetic connectivity between

northern New Zealand and the Kermadec Islands archipelago. *Marine Biology*, 161,

297–312 (doi:10.1007/s00227-013-2335-0)

Richardson, W. S., & Finlen, J. R. (1967). The transport of Northwest Providence Channel.

Deep-Sea Research, 14, 361-367 (doi:10.1016/0011-7471(67)90078-2).

Salles, O. C., Pujol, B., Maynard. J. A., Almany, G. A., Berumen, M. L., Jones, G. P., ... Planes,

S. (2016). First genealogy for a wild marine fish population reveals

multigenerational philopatry. *Proceedings of the National Academy of Sciences

U.S.A.,* 113:13245–13250 (doi:10.1073/pnas.1611797113)

Segovia, N. I., Gallardo-Escárate, C., Poulin, E., & Haye, P. A. (2017). Lineage divergence,

local adaptation across a biogeographic break, and artificial transport, shape the

genetic structure in the ascidian *Pyura chilensis*. *Science Reports*, 7, 44559

(doi:10.1038/srep44559)

Stoner, A. W. (1997). The status of queen conch, *Lobatus gigas*, research in the Caribbean.

    Marine Fisheries Review, 59, 14-23

    (https://www.researchgate.net/publication/242472717)

Stoner, A. W., & Davis, M. (1997a). Abundance and distribution of queen conch veligers

    (*Lobatus gigas*) in the central Bahamas. I. Horizontal patterns in relation to

    reproductive and nursery grounds. *Journal of Shellfish Research*, 16, 7-18

    (http://www.biodiversitylibrary.org/bibliography/2179#/summary)

Stoner, A. W., & Davis, M. (1997b). Abundance and distribution of queen conch veligers

    (*Lobatus gigas*) in the central Bahamas. II. Vertical patterns in nearshore and deep-

    water habitats. *Journal of Shellfish Research*, 16, 19-29

    (http://www.biodiversitylibrary.org/bibliography/2179#/summary)

Tay, Y. C., Chng, M. W. P., Sew, W. W. G. , Rheindt, F. E., Tun,, K. P. P., & Meier, R. (2016).

    Beyond the Coral Triangle: High genetic diversity and near panmixia in Singapore's

    populations of the broadcast spawning sea star *Protoreaster nodosus*. *Royal Society*

    Open Science, 3, 160253 (doi:10.1098/rsos.160253)

Tange, O. (2011), GNU PARALLEL: The command-line power tool.

    (https://www.usenix.org/system/files/login/articles/105438-Tange.pdf)

Teske, P. R., Sandoval-Castillo, J., van Sebille, E., Waters, J., & Beheregaray ,L. B. (2016).

    Oceanography promotes self-recruitment in a planktonic larval disperser. *Science*

    Reports, 6, 34205 (doi:10.1038/srep34205)

Thomas, L., Kennington, W. J., Evans, R. D., Kendrick, G. A., & Stat, M. (2017). Restricted

gene flow and local adaptation highlight the vulnerability of high-latitude reefs to

rapid environmental change. *Global Change Biology*, 23, 2197–2205

(doi:10.1111/gcb.13639)

Truelove, N. K., Kough, A. S., Behringer, D. C., Paris, C. B., Box, S. J., Preziosi, R. F., & Butler,

M. J. IV. (2017). Biophysical connectivity explains population genetic structure in a

highly dispersive marine species. *Coral Reefs*, 36, 233–244 (doi:10.1007/s00338-

016-1516-y)

Truelove, N. K., Box, S. J., Aiken, K. A., Blythe-Mallett, A., Boman, E. M., Booker, C. J., …

Stoner, A. W. (2017). Isolation by oceanic distance and spatial genetic structure in

an overharvested international fishery. *Diversity and Distributions*, 23, 1–9

(doi:10.1111/ddi.12626)

Van Wyngaarden, M., Snelgrove, P. V. R., DiBacco, C., Hamilton, L. C., Rodríguez-Ezpeleta,

N., Jeffery, N. W., … Bradbury, I. R. (2016). Identifying patterns of dispersal,

connectivity, and selection in the sea scallop, Placopecten magellanicus, using

RADseq-derived SNPs. *Evolutionary Appiications*, 10, 102–117

(doi:10.1111/eva.12432)

Veríssimo, A., Sampaio, I., McDowell, J. R., Alexandrino, P., Mucientes, G., Queiroz, N., …

Noble, L. R. (2017). World without borders—genetic population structure of a

highly migratory marine predator, the blue shark (*Prionace glauca*). *Ecology and

Evolution*, 7, 4768–4781 (doi:10.1002/ece3.2987)

Wallace, E. M., & Tringali, M. D. (2016). Fishery composition and evidence of population

structure and hybridization in the Atlantic bonefish species complex (*Albula*

spp.). *Marine Biology*, 163, 142.(doi:10.1007/s00227-016-2915-x)

Weersing, K., & Toonen, R. J. (2009). Population genetics, larval dispersal, and connectivity

in marine systems. *Marine Ecology Progress Series*, 393, 1–12

(doi:10.3354/meps08287)

Williamson, D. H., Harrison, H. B., Almany, G. R., Berumen, M. L., Bode, M., Bonin, M. C., …

Jones, G. P. (2016). Large-scale, multidirectional larval connectivity among coral

reef fish populations in the Great Barrier Reef Marine Park. *Molecular Ecology*, 25,

6039–6054 (doi:10.1111/mec.13908)

Yau, A. J., Lenihan, H. S., & Kendall, B. E. (2014) Fishery management priorities vary with

self-recruitment in sedentary marine populations. *Ecological Applications*, 24, 1490-

1504 (doi:10.1890/13-1201.1)

TABLE 1 Collection data for bonefish (*Albula vulpes*) and queen conch (*Lobatus gigas*) at Grand Bahama and Eleuthera islands, northern Caribbean Basin. Listed are: Species, Island, Side (of island with compass direction in parentheses), Locality, Date, N (=sample size), and Label (=sample abbreviation).

Species	Island	Side	Locality	Date	N	Label
Bonefish	Grand Bahama	Flats (N)	North	26-Jan-14	30	GNB
Bonefish	Grand Bahama	Ocean (S)	South	25-Jan-14	30	GSB
Bonefish	Eleuthera	Flats (W)	Broad Creek	2-Mar-14	30	EWB
Bonefish	Eleuthera	Ocean (E)	Half Sound	8-Mar-14	10	EEB
Bonefish	Eleuthera	Flats (SW)	Plum Creek	8-Mar-14	10	ES1
Bonefish	Eleuthera	Flats (SW)	Sound Side	10-Mar-14	7	ES2
Conch	Grand Bahama	South	Freeport	1-Jan-14	30	GSC
Conch	Eleuthera	Flats (W)	Schooner Cay 2	1-Mar-14	30	EWC

TABLE 2 Study species (= bonefish *Albules vulpes*; queen conch *Lobatus gigas*) evaluated in this study. Also provided are number of genetic loci greater than 50% retention (= Loci > 50%); Average number of loci per individual (= Av loci/ind); Average depth of coverage (= Av coverage); Number of parsimonious-informative SNPs (= PI SNPs); Number of unlinked SNPs (=ALL); and number of SNPs sampled at 2 (=SD2) and 3 (=SD3) standard deviations from the mean F_{ST}.

Species	Loci > 50%	Av loci/ind	Av coverage	PI SNPs	ALL	SD2	SD3
Bonefish	56,046	35,443	77X	36,206	25,205	123	79
Conch	30,697	33,409	59X	44,334	20,873	88	51

TABLE 3 BᴀʏᴇsAss3 results for bonefish (*Albula vulpes*) and queen conch (*Lobatus gigas*) sampled at Grand Bahama and Eleuthera islands. Values are expressed as proportion of individuals expected to be of a given migrant or non-migrant origin. Migration rates along the diagonal are analogous to a self-recruitment rate, with high values representative of demographic independence. Sample site abbreviations per Table 1.

	EWB	EEB	ES1	ES2	GNB	GSB	GNC	EWC
EWB	0.884	0.039	0.020	0.019	0.021	0.018	-	-
EEB	0.085	0.833	0.020	0.021	0.020	0.020	-	-
ES1	0.104	0.146	0.689	0.020	0.020	0.021	-	-
ES2	0.127	0.106	0.026	0.693	0.025	0.025	-	-
GNB	0.104	0.075	0.016	0.016	0.774	0.015	-	-
GSB	0.179	0.077	0.015	0.016	0.016	0.698	-	-
GNC	-	-	-	-	-	-	0.987	0.013
EWC	-	-	-	-	-	-	0.013	0.988

FIGURE LEGENDS

FIGURE 1 (Top) Map depicting The Commonwealth of Bahama (insert top right). The islands of Grand Bahama, Abaco, Eleuthera and Andros (northeastern Basin) are depicted. Blue dots represent sample sites for queen conch, whereas those for bonefish are designated with red dots. (Bottom) Sample sites at Eleuthera Island for queen conch and bonefish. Colors as above. Labels as in Table 1.

FIGURE 2 (Top) Bathymetry map of the northwest Carribean Basin (modified from http://oceanexplorer.noaa.gov/explorations/09deepseacorals/logs/summary/media/bahamas_region_600.html). Colors indicate depth: Blues = deepest (>3,000m); yellow-to-green = intermediate (500-2500m); beige = shallowest (<500m); Islands depicted in black. Red ellipse = Eleuthera Island; Great Bahama Canyon = 1; Great Bahama Flat = 2; Exuma Sound = 3. (Bottom) Oceanic currents in the northern Caribbean Basin (modified from: University of Miami Rosenthiel School of Marine and Atmospheric Science, http://oceancurrents.rsmas.miami.edu/caribbean/caribbean-maps.html). Black lines = currents; Yellow arrow = direction of Gulf Stream (east coast of Florida); Green arrow = current direction in Grand Bahama Canyon; Eleuthera Island = 2; Red Arrow = direction of Antillean Current (Atlantic Ocean).

FIGURE 3. STRUCTURE plots for bonefish (*Albula vulpes*) sampled at Grand Bahama and Eleuthera islands, The Bahamas. ALL = Plot employing all single nucleotide polymorphisms (SNPs) (N=25,205); K = 2 represent a 2-group analysis employing: (1) SD2 dataset containing single nucleotide polymorphisms (SNPs) >2 standard deviations (sd) above mean F_{ST} (N=123); (2) SD3 dataset containing SNPs >3 sd above mean F_{ST} (N=79). K = 3 represents 3-groupl analysis involving the same two datasets. GNB = Grand Bahamas Flats North (Flats or North side); GSB = Grand Bahamas Ocean Side (South or Bahamian side); EWB = Eleuthera Flats Broad Creek (West side); EEB = Eleuthera Ocean Half-Sound (Atlantic Ocean or East side); ES1 = Eleuthera Ocean Plum Creek (Flats or Southwest side); ES2 = Eleuthera Ocean South Side (Flats or Southwest side). Sample site details in Table 1.

FIGURE 4. Plots depicting discriminant analysis of principal components (DAPC) for six populations of bonefish (*Albula vulpes*) along discriminant functions one (=DF1) and two (=DF2). ALL = plot utilising all loci (N=25,205); SD2 = plot depicting single nucleotide polymorphisms (SNPs) >2 standard deviations (sd) above mean F_{ST} (N=123); SD3 = plot employing SNPs >3 sd above mean F_{ST} (N=79). GNB = Grand Bahamas Flats North (Flats or North side); GSB = Grand Bahamas Ocean Side (South or Bahamian side); EWB = Eleuthera Flats Broad Creek (West side); EEB = Eleuthera Ocean Half-Sound (Atlantic Ocean or East side); ES1 = Eleuthera Ocean Plum Creek (Flats or Southwest side); ES2 = Eleuthera Ocean South Side (Flats or Southwest side). Additional details in Table 1.

FIGURE 5. Structure plots for queen conch (*Lobatus gigas*) sampled as Grand Bahama and Eleuthera islands, the Bahamas, with individuals represented by vertical lines. Two groups are depicted [GSC = Grand Bahama South Freeport (Flats or South side); EWC = Eleuthera Flats Schooner Cay 2 (Flats or West side)]; ALL = Plot employing all single nucleotide polymorphisms (SNPs) (N=20,873); SD2 = SNPs >2 standard deviations (sd) above mean F_{ST} (N=88); and SD3 = plot depicting SNPs >3 sd above mean F_{ST} (N=51).

FIGURE 6. Plots depicting discriminant analysis of principal components (DAPC) for two populations of queen conch (*Lobatus gigas*), with GSC = Grand Bahama South Freeport (Flats or South side) and EWC = Eleuthera Flats Schooner Cay 2 (Flats or West side) plotted along discriminant function one (=DF1). ALL = Plot employing all single nucleotide polymorphisms (SNPs) (N=20,873); SD2 = SNPs >2 standard deviations (sd) above mean F_{ST} (N=88); and SD3 = plot depicting SNPs >3 sd above mean F_{ST} (N=51).

FIGURE 7. Graph depicting relative migration rates for six populations of bonefish (*Albula vulpes*), where nodes represent populations, and edges are weighted according to the level of direction relative migration rates (*m*), as derived from BA3-SNPs results. All edges below 0.02 were omitted. Full results shown in Table 3. GNB = Grand Bahamas Flats North (Flats or North side); GSB = Grand Bahamas Ocean Side (South or Bahamian side); EWB = Eleuthera Flats Broad Creek (West side); EEB = Eleuthera Ocean Half-Sound (Atlantic Ocean

or East side); ES1 = Eleuthera Ocean Plum Creek (Flats or Southwest side); ES2 = Eleuthera

Ocean South Side (Flats or Southwest side).

FIGURE 8. Asymmetric immigration recorded for bonefish (*Albula vulpes*) from Grand

Bahama and Eleuthera islands. Values are expressed as net emigration rates (summed

emigration - summed immigration) for each population, with positive values indicating

'source' populations and negative values indicating 'sink' populations. GNB = Grand

Bahamas Flats North (Flats or North side); GSB = Grand Bahamas Ocean Side (South or

Bahamian side); EWB = Eleuthera Flats Broad Creek (West side); EEB = Eleuthera Ocean

Half-Sound (Atlantic Ocean or East side); ES1 = Eleuthera Ocean Plum Creek (Flats or

Southwest side); ES2 = Eleuthera Ocean South Side (Flats or Southwest side).