1

Genetic variation in anesthesia success in a New Zealand freshwater snail

Qiudong Song[1*], Richard Magnuson[1*], Joseph Jalinsky[1], Marissa Roseman[1], and

Maurine Neiman[1,2]

[1]*Department of Biology, University of Iowa, Iowa City, IA, USA*

[2]*Department of Gender, Women's, and Sexuality Studies, University of Iowa, Iowa City,*

IA, USA

Co-first authors and contributed equally

Corresponding author: Maurine Neiman, maurine-neiman@uiowa.edu

Keywords: *Potamopyrgus antipodarum*, gastropod, mollusk, asexual, pain

Abstract

Anesthesia is used to immobilize organisms for experiments and surgical procedures

and to humanely mitigate pain. Despite its scientific importance, and though

invertebrates seem to experience pain, overarching regulations for the application of

anesthesia in invertebrate taxa do not exist. Indeed, the almost complete absence of

legal requirements for invertebrate anesthesia might explain at least in part why there

seems to have been little effort towards developing widely effective invertebrate

anesthesia techniques. Here, we demonstrate that genetic differences between snails

are a major driver of wide variation in menthol anesthesia success in an important

freshwater snail model system, *Potamopyrgus antipodarum*. This is the first example of

which we are aware of a role for genetic variation in anesthesia success in a mollusk.

These findings both illuminate a potential mechanism underlying previously documented

challenges in molluscan anesthesia and will set the stage for powerful and humane

manipulative experiments in *P. antipodarum*.

1. INTRODUCTION

Animal welfare regulations do not typically include invertebrates at least in part because how these organisms process pain is not well understood (AMVA, 2020; Gilbertson & Wyatt, 2016). Nevertheless, invertebrates exhibit clear responses to stimuli that could be perceived as perception of pain (AVMA, 2020). Indeed, Canada, the United Kingdom, and the European Union have animal research regulations that extend protections to cephalopods (Butler-Struben, Brophy, & Crook, 2018), potentially signifying a paradigm shift regarding which taxa deserve formal protections. The basis for animal welfare regulation of cephalopod research is the assumption, rooted in observational studies (e.g., Roper & Hochberg, 1988; Wells, 1978), that cephalopods perceive pain and exhibit emotional responses. This assumption has since been backed up empirically (e.g., Crook, Hanlon, & Walters 2013; Scientific Panel on Animal Health and Welfare, 2005). Researcher perception of the wellbeing of a research organism can also affect study outcomes. For example, as outlined by Poole (1997), conclusions reached from studies using organisms under distress may be unreliable and thus that pain experienced by research organisms can influence results and ultimately study conclusions (also see AVMA, 2020).

Other than cephalopods, there are no other general protections extended to invertebrates for use in research of which we are aware. In the absence of regulatory oversite, given the likelihood that at least some invertebrates perceive pain, and because some experimental procedures are facilitated by or require that the animal is immobilized, there is a clear demand for ability to reliably and effectively apply anesthesia in all invertebrates.

We here describe important steps forward in identifying an effective anesthesia

technique for *Potamopyrgus antipodarum* (GRAY, 1843), a New Zealand freshwater

snail model system that is a prominent model system for the evolution of sexual

reproduction (e.g., Neiman, Paczesniak, Soper, Baldwin, & Hehman, 2011),

ecotoxicology (e.g., Geiß et al., 2017), invasion biology (e.g., Donne, Neiman, Woodell,

Haase, & Verhaegen, 2020), and host-parasite interactions (e.g., Bankers et al., 2017).

Studies of *P. antipodarum* often incorporate tissue manipulation (e.g., tentacle severing

as a test of tissue regeneration; Krois, Cherukuri, Puttagunta, & Neiman, 2013) or

require determination of male vs. female status (e.g. Jalinsky, Logsdon, & Neiman,

2020). These studies either require or would be qualitatively improved by the availability

of an effective, safe, and easy-to-use anesthetic.

Menthol crystals are the preferred method for anesthetizing *P. antipodarum* (e.g.,

Krois et al., 2013), with the standard approach based on exposure of snails to crushed

menthol crystals for ~90 minutes (McCraw, 1958). While this method can be effective

(Krois et al., 2013), a considerable fraction (often >50%) of exposed *P. antipodarum* do

not become anesthetized (Magnuson, 2018). Similar intraspecific variation and

generally low anesthesia efficiency has been reported in other mollusks (e.g., Butler-

Struben, Brophy, & Crook, 2018; Fiorito et al., 2015).

Magnuson (2018) noted that some *P. antipodarum* lineages (defined as all snails

descended from a single, laboratory-isolated female) seemed to have higher anesthesia

success than others. This observation led us to hypothesize that this lineage effect

points to an important role of genetic background in anesthesia success in these snails.

This hypothesis finds indirect support from reports of genetically based differences in

anesthesia efficacy in rodents (e.g., Chesler et al., 2003; Moghil, Smith, O'Reilly, &

Plourde, 2005; Seltzer, 2014), insects (e.g., Guan, Scott, & Nash, 2000), and

roundworms (Hawasli, Saifee, Liu, Nonet, & Crowder, 2004; Morgan & Sedensky, 1994)

(reviewed in Nash, 2002; Steele, Morgan, & Sedensky, 2007).

Here, we performed the first test of which we are aware of a genetic basis for

anesthesia efficacy in mollusks by subjecting different genetic lineages of common

garden-born and raised *P. antipodarum* asexual lineages to the same menthol

anesthesia protocol. The presence of significance across-lineage variation in anesthesia

success would be consistent with this hypothesis, suggesting that some *P. antipodarum*

genetic backgrounds are more susceptible to menthol anesthesia than others. This

result will be useful to other *P. antipodarum* and mollusk researchers going forward,

highlighting a potential explanation for frequently reported challenges in anesthesia

success and indicating that at least some experiments requiring anesthesia can

succeed if susceptible lineages are used.

2. METHODS

2.1. Anesthesia experiment

We followed the definition of successful anesthesia provided by Lewbart & Mosley

(2011), who considered the snail to be anesthetized when the animal shows no tentacle

withdrawal or body movement following the gentle scrape of its foot with a needle

("needle test"). For the experiments presented herein, the anesthetization process

begins with the addition of 200 mL of carbon-filtered tap water to a ~1/2-L plastic cup.

We then added the *P. antipodarum* to be anesthetized to the cup, immediately followed

by the addition of 2 g of crushed menthol crystals. After 90 minutes of exposure, we

removed the snails from the menthol water. We defined anesthesia as successful if the

body of the snail was relaxed such that the tentacles, head, and foot had emerged from

the shell and there was no withdrawal or body movement during the needle test

(administered with a 25-gauge needle). We deemed anesthesia as unsuccessful if the

snail was fully contracted into the shell or if the snail was relaxed but failed the needle

test.

We then administered this anesthesia protocol to each of six groups of ten snails

(N = 60 snails per lineage) from each of 17 different *P. antipodarum* triploid asexual

lineages, for a total of 102 trials and 1020 snails (Table 1). While each of the 17

lineages was founded by a single unique asexual female and was cultured separately in

its own tank, three pairs of lineages (six of the 17 lineages) shared descent from the

same field-collected grandmother. These founding grandmothers were sampled from

natural populations two generations before the birth of the snails used here, providing

ample time to eliminate any effects of differences in the sampling environment. The

implications are that these three pairs of lineages (each pair defined by descent from

the same grandmother) will be genetically identical (barring *de novo* mutations) across

pair members (hereafter, "clonemates"). We were later able to back up these

assumptions by leveraging SNP data collected for another project (see Discussion). By

enabling comparisons between clonemates (N = three pairs) as well as across

genetically distinct lineages ("non-clonemates"; 14 lineages, including those three

lineage pairs), our set of lineages thus provides a powerful test of the extent to which

genetic factors influence anesthesia success.

All of the snails in our experiment were born and raised in the same constant-

temperature room, under identical feeding and culture conditions (following standard of

care for *P. antipodarum*; e.g., Zachar & Neiman, 2013). The snails used for the trials

were haphazardly selected from the adult snails in the 15 L tanks used to house each

lineage and were not reused across trials. Our measure of anesthesia success per trial

was the percentage of snails, out of the ten snails per lineage per trial, that were

successfully anesthetized.

Because the anesthesia success data were not distributed normally (Kolmogorov-

Smirnov test; test statistic = 0.111, df = 102, p = 0.003), we used a non-parametric

Kruskal-Wallis test to first address whether there was a significant effect of lineage

founder ("grandmother") on anesthesia success. A significant outcome of this Kruskal-

Wallis analysis will indicate that the descendants of different grandmothers respond

differently to anesthesia. Because the founding grandmothers constituted separately

sampled snails from highly diverse New Zealand populations (e.g., Paczesniak, Jokela,

Larkin, & Neiman, 2013; Verhaegen, McElroy, Bankers, Neiman, & Haase, 2018), this

outcome is consistent with a major role of genetic variation for anesthesia success in *P.*

antipodarum.

We next used the lineage-level anesthesia success data to address whether the

three pairs of clonemates were significantly more similar in anesthesia success rates

than were non-clonemate pairs. We began by calculating the absolute value of the

difference in anesthesia success rates ("difference scores") between the means of the

six success rates measured for each of the 17 lineages relative to each of the other

lineages. For example, this value for Alex 13 relative to Alex 27 is the absolute value of

0.23 (Alex 13 mean success rate) - 0.92 (Alex 27 mean success rate), equivalent to

0.69. There were a total of three unique pairwise difference scores for the clonemates

(reflecting the three pairs of clonemates) and a total of 136 unique pairwise difference

scores for the non-clonemate comparisons. We then used one-sample sign tests (as

implemented at

https://www.mathcelebrity.com/nonparam.php?pl=Generate+Practice+Problem) relative

to a test value of zero, to determine whether the clonemates and non-clonemate

difference scores, respectively, were significantly different than zero. In a scenario

where genetic background is an important determinant of anesthesia success, the

clonemate difference score would not be significantly different than zero, while the non-

clonemate score should be significantly higher than zero.

2.1. SNP Genotyping

As part of a large and wholly independent population genomics project, we used

KASP™ assays to genotype 14 of the 17 lineages at 49 of the 50 SNPs used in

Verhaegen et al. (2018) (Supporting Information Table S1). We began by using the

same chloroform-phenol DNA extraction method described in Sharbrough, Luse, Boore,

Logsdon, and Neiman (2018) on the dissected head tissue of one randomly selected

adult snail from each of the fourteen lineages. We then sent the DNA extractions to

LGC Genomics (Hoddesdon, UK) for SNP genotyping. Upon receiving the raw data, we

assigned SNP genotypes by using the infinite alleles model distance index, setting the

maximum distance threshold defining a genotype (Rogstad, Keane, & Beresh, 2002) to

zero differences (excluding the three sites of 686 total genotyped with missing data)

with GenoDive software v.2.0 (Meirmans & Van Tienderen, 2004).

3. RESULTS

Our analyses revealed wide across-lineage variation in anesthesia outcome between

lineages descended from different grandmothers (Kruskal-Wallis, $p < 0.001$; Figure 1),

ranging from a median of 95% of all snails anesthetized for several different lineages

from lake Alexandrina to ~8% for Grasmere 59 (Table 1). This result is as is expected if

there is a major genetic component to anesthesia success in *P. antipodarum*.

The difference score between clonemates was not significantly different than zero

(one-sample sign test; $p = 0.125$), with the caveat that we had only three comparisons

and thus low statistical power. By contrast, the difference score between non-

clonemates was significantly higher than zero ($p = 0.002$) and was nearly twice as high

(mean difference score 0.340 +/- 0231 SD) as the mean difference score for the

clonemates (mean difference score = 0.173 +/- 0.132 SD; Fig. 2). These results are also

consistent with a scenario where genetic factors influence anesthesia success in *P.*

antipodarum.

There were eight distinct SNP genotypes across the 14 lineages genotyped

(Table 1). These SNP data demonstrated that the two pairs of clonemates that were

genotyped (Alex 19_1 and Alex 19_2 and Grasmere 49_1 and Grasmere 49_2) indeed

shared SNP genotypes. These data also revealed that most other non-clonemate

lineages themselves had unique genotypes. Finally, we discovered that the two sets of

lineages with different grandmothers that nevertheless shared SNP genotypes 4 and

26, respectively, had strikingly similar anesthesia success rates relative to other

lineages with different SNP genotypes (Fig. 1).

4. DISCUSSION

Together, our data indicate that genetic background is a major contributor to menthol

anesthesia success in *P. antipodarum*, an important model system in evolutionary

biology, ecology, and ecotoxicology. This is the first demonstration of which we are

aware of a genetic component to variation in anesthesia efficacy in mollusks. One

obvious practical implication of these results for *P. antipodarum* researchers is the

opportunity to select lineages with higher anesthesia susceptibility for experiments that

require anesthetics.

We were able to perform a powerful independent validation of the likelihood that

genetic variation does underlie across-lineage variation in anesthesia success in *P.*

antipodarum by leveraging single-nucleotide polymorphism (SNP) data generated for a

different population genomic project. These data backed up our assumptions about the

genetic identity of clonemates relative to non-clonemates and also revealed that

lineages not previously known to be close relatives that shared a SNP genotype had

very similar anesthesia outcomes relative to lineages with different SNP genotypes.

To our knowledge, this evidence for genetic variation for anesthetic success in *P.*

antipodarum is the first report of such a phenomenon in mollusks. That genetic

background influences anesthesia success in a wide variety of common model systems

like flies, worms, and mice suggest that the same might be expected for other mollusk

taxa.

Two important questions worthy of future study emerge from our results. First,

what is the genetic basis of the across-lineage variation in anesthesia success?

Second, even though clonemates seem to share similar anesthesia success outcomes

relative to non-clonemates, we do not see identical responses (i.e., no examples of

100% success or 100% failure) within clonemates.

The first question can be addressed by using techniques such as RNA

sequencing and whole-genome sequencing to identify gene expression differences and

genetic variants differentiating lineages with very distinct anesthesia responses.

Because multiple studies indicate that asexual *P. antipodarum* lineages from the same

lake population are typically closely related relative to asexual lineages from other lakes

(e.g., Paczesniak et al., 2013), a powerful experiment might be one that compared

genomic sequence from lineages from the same lake with different anesthesia

responses. A good example of such lineages in the experiment we describe here are

provided by low-responding lineages from lake Alexandrina like Alex 13 (23.33%

anesthesia success) and Alex 21 (15% anesthesia success) as compared to Alex 27

(91.67% anesthesia success) and Alex 32 and 36 (95% anesthesia success).

The second question, regarding variation in anesthesia success even within

clones and clonemates, hints that non-genetic factors (e.., age, body condition) could

influence anesthesia outcomes (e.g., Gao et al., 2018; Guenette & Lair, 2006; Larsson

& Wahlström, 1998). While we made a distinct effort to minimize age effects by

choosing the largest snails possible from each lineage, we did not know the ages of

individual snails. Nor are we able to exclude the possibility that across-snail differences

in health or body condition could have come into play. Future experiments could

address these possibilities by comparing anesthesia success within a lineage as a

function of snail age or in the context of experimental manipulation of factors like food

availability.

Acknowledgements:

We gratefully acknowledge Molly Gallagher, Sydney Stork, Benjamin Ripperger, and

Alexander Kern for snail care. John Logsdon contributed to snail collections. David

("Davey") Neiman contributed his Excel expertise to data analysis. The University of

Iowa Department of Biology Honors program, the Iowa Center for Undergraduate

Research, and Rick and Linda Maxson logistically and financially supported several

undergraduate researchers involved in the project. The SNP genotyping described here

was also supported by NSF-MCB 1122176, NSF-DEB 1731657, and NSF-DEB

1601242, with contributions to lineage selection by collaborators Kristi Montooth and

Kyle McElroy.

References:

American Veterinary Medical Association. 2020. The AMVA Guidelines for the

Euthanasia of Animals: 2020 Edition.

Bankers, L., Fields, P., McElroy, K. E., Boore, J. L., Logsdon, J. M., Jr., & Neiman, M.

(2017). Genomic evidence for population-specific responses to coevolving

parasites in a New Zealand freshwater snail. *Molecular Ecology*, *26*, 3663-3675.

Butler-Struben, H. M., Brophy, S. M., & Crook, R. J. (2018). *In vivo* recording of neural

and behavioral correlates of anesthesia induction, reversal, and euthanasia in

cephalopod molluscs. *Frontiers in Physiology*, *9*, 109.

Chesler, E., Ritchie, J., Kokayeff, A., Lariviere, W. R., Wilson, S. G., & Moghil, J. S.

(2003). Genotype-dependence of gabapentin and pregabalin sensitivity: the

pharmacogenetic mediation of analgesia is specific to the type of pain being

inhibited. *Pain*, *106*, 325-335.

Crook, R. J., Hanlon, R. T., & Walters, E. T. (2013). Squid have nociceptors that display

widespread long-term sensitization and spontaneous activity after bodily injury.

Jour*nal of Neuroscience*, *33*, 10021-10026.

Donne, C., Neiman, M., Woodell, J. D., Haase, M., & Verhaegen, G. (2020). A layover

in Europe: reconstructing the invasion route of asexual lineages of a New

Zealand snail to North America. *Authorea* doi:

10.22541/au.158456440.01903363 (Accepted pending minor revisions in

Molecular Ecology).

d'Ovidio, D., Monticelli, P., Santoro, M., & Adami, C. (2019). Immersion anesthesia with

ethanol in African giant land snails (*Acathina fulica*). *Heliyon*, *5*, e01546.

Fiorito, G., Affuso, A., Basil, J., Cole, A., de Girolamo, P., D'Angelo, L., Dickel, L.,

Gestal, C., Grasso, F., Kuba, M., Mark, F., Melillo, D., Osorio, D., Perkins, K.,

Ponte, G., Shashar, N., Smith, D., Smith, J., & Andrews, P. L. R. (2015).

Guidelines for the care and welfare of cephalopods in research- a consensus

based on an initiative by CephRes, FELASA, and the Boyd group. *Laboratory*

Animals, *49*, 1-90.

Gao, J., Hu, Z., Shi, L., Li, N., Ouyang, Y., Shu, S., Yao, S., & Chen, X. (2018). HCN

channels contribute to the sensitivity of intravenous anesthetics in developmental

mice. *Oncotarget*, *9*, 12907–12917.

Geiß, C., Ruppert, K., Askem, C., Barroso, C., Faber, D., Ducrot, V., Holbech, H.,

Hutchinson, T. H., Kajankari, P., Kinnberg, K. L., Lagadic, L., Matthiessen, P.,

Morris, S., Neiman, M., Penttinen, O.-P., Sanchez-Marin, P., Tiegeler, M.,

Weltje, L., & Oehlmann, J. (2017). Validation of the OECD Reproduction Test

Guideline with the New Zealand mudsnail *Potamopyrgus antipodarum* using

trenbolone and prochloraz. *Ecotoxicology*, *26*, 370-382.

Gilbertson, C. R., & Wyatt, J. D. (2016). Evaluation of euthanasia techniques for an

invertebrate species, land snails (*Succinea putris*). *Journal of the American*

Association for Laboratory Animal Science, *55*, 577-581.

Guan, Z., Scott, R. L., & Nash, H. A. (2000). A new assay for the genetic study of

general anesthesia in *Drosophila melanogaster*: Use in analysis of mutations in

the X-chromosomal 12E region. *Journal of Neurogenetics*, *14*, 25-42.

Guenette, S. A., & Lair, S. (2006). Anesthesia of the leopard frog, *Rana pipiens*: A

comparative study between four different agents. *Journal of Herpetological*

Medicine and Surgery, *16*, 38-44.

Hawasli, A. H., Saifee, O., Liu, C., Nonet, M. L., & Crowder, C. M. (2004). Resistance of

volatile anesthetics by mutations enhancing excitatory neurotransmitter release

in *Caenorhabditis elegans*. *Genetics*, *168*, 831-843.

Krois, N. R., Cherukuri, A., Puttagunta, N., & Neiman, M. (2013). Higher rate of tissue

regeneration in polyploid asexual vs. diploid sexual freshwater snails. *Biology*

Letters, *9*, 20130422.

Larsson, J., & Wahlström, G. (1998). The influence of age and administration rate on

the brain sensitivity to propofol in rats. *Acta Anaesthesiologica Scandinavica, 42*,

987-994.

Lewbart, G. A., & Mosley, C. (2011). Clinical anesthesia and analgesia in invertebrates.

Journal of Exotic Pet Medicine, *21*, 59-70.

Lively, C. M. 1987 Evidence from a New Zealand snail for the maintenance of sex by

parasitism. *Nature*, *328*, 519-521.

Logsdon, J. M., Neiman, M., Boore, J., Sharbrough, J., Bankers, L., McElroy, K.,

Jalinsky, J., Fields, P., & Wilton, P. (2017). A very recent whole genome duplication in

Potamopyrgus   antipodarum predates multiple origins of asexuality & associated

polyploidy. *PeerJ   Preprints*, *5*, e3046v3041.

Magnuson, R. (2018). An exploration of alternative anesthesia in the difficult-to-

anesthetize New Zealand freshwater snail *Potamopyrgus antipodarum*. Iowa

Research Online.

Meirmans, P. G., & Van Tienderen, P. H. (2004). GENOTYPE and GENODIVE: two

programs for the   analysis of genetic diversity of asexual organisms. *Molecular*

Ecology Notes, *4*, 792–794.

McCraw, B. M. (1958). Relaxation of snails before fixation. *Nature*, *181*, 575.

Moghil, J. S., Smith, S. B., O'Reilly, M. K., Plourde, G. (2005). Influence of nociception

and stress-induced antinociception on genetic variation in isoflurane anesthetic

potency among mouse strains. *Anesthesiology*, *103*, 751-758.

Morgan, P. G., & Sedensky, M. M. (1994). Mutations conferring new patterns of

sensitivity to volatile anesthetics in *Caenorhabditis elegans*. *Anesthesiology*, *81*,

888-898.

Nash, H. A. (2002). *In vivo* genetics of anaesthetic action. *British Journal of*

Anaesthesia, *89*, 143-155.

Neiman, M., Larkin, K., Thompson, A. R., & Wilton, P. (2012). Male offspring production

by asexual *Potamopyrgus antipodarum*, a New Zealand snail. *Heredity, 109*, 57-

62.

Neiman, M., Jokela, J., & Lively, C. M. (2005). Variation in asexual lineage age in

Potamopyrgus antipodarum, a New Zealand snail. *Evolution*, *59*, 1945-1952.

Neiman, M., Paczesniak, D., Soper, D. M., Baldwin, A. T., & Hehman, G. (2011). Wide

variation in ploidy level and genome size in a New Zealand freshwater snail with

coexisting sexual and asexual lineages. *Evolution*, *65*, 3203–3216.

Paczesniak, D., Jokela, J., Larkin, K., & Neiman, M. (2013). Discordance between

nuclear and mitochondrial genomes in sexual and asexual lineages of the

freshwater snail *Potamopyrgus antipodarum*. *Molecular Ecology*, *22*, 4695-4710.

Poole T. (1997). Happy animals make good science. *Laboratory Animals*, *31*, 116-124.

Rogstad, S. H., Keane, B., & Beresh, J. (2002). Genetic variation across VNTR loci in

central North American *Taraxacum* surveyed at different spatial scales. *Plant*

Ecology, *161*, 111–121.

Roper C.F.E., & Hochberg F.G. (1988). Behavior and systematics of cephalopods from

Lizard Island, Australia, based on color and body patterns. *Malacologia*, *29*, 153-

193.

Scientific Panel on Animal Health and Welfare (AHAW). 2005. Opinion of the Scientific

Panel on Animal Health and Welfare (AHAW) on a request from the Commission

related to the 'Aspects of the biology and welfare of animals used for

experimental and other scientific purposes'. *EFSA Journal*, *292*, 1-136.

Seltzer, Z. (2014). Nothing in pain makes sense except in the light of genetics. *Pain*,

2014, 841-842.

Sharbrough, J., Luse, M., Boore, J. L., Logsdon, J. M., Jr., & Neiman, M. (2018).

Radical amino acid mutations persist longer in the absence of sex. *Evolution*, *72*,

808-824.

Steele, L. M., Morgan, P. G., & Sedensky, M. M. (2007). Genetics and the mechanisms

of action of inhaled anesthetics. *Current Pharmacogenomics*, *5*, 125-141.

Verhaegen, G., McElroy, K., Bankers, L., Neiman, M., & Haase, M. (2018). Adaptive

phenotypic plasticity in a clonal invader. *Ecology and Evolution*, *8*, 4465-4483.

Wells, M. J. (1978). Octopus: Physiology and behavior of an advanced invertebrate.

London, UK: Chapman and Hall.

Zachar, N., & Neiman, M. (2013). Profound effects of population density on fitness-

related traits in an invasive freshwater snail. *PLoS ONE, 8*, e80067.

Figure Legends and Table Caption:

Figure 1: Median percent of successfully anesthetized snails (6 trials of 10 snails each) subjected to the standard *P. antipodarum* menthol anesthesia protocol per each of 17 lineages founded by 14 different grandmothers. Triangle symbols (vs. circles) are used to denote the three lineage pairs that each share a grandmother. The numbers above the 14 of the 17 lineages that were genotyped indicate SNP genotype.

Figure 2: Median pairwise difference score in anesthesia success for clonemate vs. non-clonemate lineages. The non-clonemate pairwise difference scores were significantly higher than zero while the clonemate pairwise difference scores were not significantly higher than zero.

383   Table 1. Description of the *P. antipodarum* asexual lineages used in this study.

Lake of Origin	Lineage	Grandmother	Mean Anesthesia Success % (+/- SD)	Genotyped	SNP Genotype
Alexandrina	Alex 13	Alex 13	23.33 +/- 15.01	N	NA
Alexandrina	Alex 19_1	Alex 19	50.00 +/- 6.32	Y	3
Alexandrina	Alex 19_2	Alex 19	53.33 +/- 10.33	Y	3
Alexandrina	Alex 21_1	Alex 21	15.00 +/- 8.37	Y	2
Alexandrina	Alex 21_2	Alex 21	35.00 +/- 12.25	N	NA
Alexandrina	Alex 27	Alex 27	91.67 +/- 11.69	Y	4
Alexandrina	Alex 31	Alex 31	48.33 +/- 7.53	N	NA
Alexandrina	Alex 32	Alex 32	95.00 +/- 8.37	Y	4
Alexandrina	Alex 36	Alex 36	95.00 +/- 5.48	Y	5
Alexandrina	Alex 39	Alex 39	93.33 +/- 8.16	Y	4
Grasmere	Grasmere 49_1	Grasmere 49	76.67 +/- 15.01	Y	26
Grasmere	Grasmere 49_2	Grasmere 49	48.33 +/- 9.83	Y	26
Grasmere	Grasmere 54	Grasmere 54	83.33 +/- 13.66	Y	26
Grasmere	Grasmere 59	Grasmere 59	8.33 +/- 7.53	Y	24
Gunn	Gunn 6	Gunn 6	36.67 +/- 20.66	Y	29
Gunn	Gunn 7	Gunn 7	71.67 +/- 16.02	Y	26
Gunn	Gunn 9	Gunn 9	78.33 +/- 14.72	Y	31

384
385