1

Genetic variation in anesthesia success in a New Zealand freshwater snail

Qiudong Song[1*], Richard Magnuson[1*], Joseph Jalinsky[1], Marissa Roseman[1], and Maurine

Neiman[1,2]

1Department of Biology, University of Iowa, Iowa City, IA, USA

2Department of Gender, Women's, and Sexuality Studies, University of Iowa, Iowa City, IA, USA

Co-first authors and contributed equally

Corresponding author: Maurine Neiman, maurine-neiman@uiowa.edu

Keywords: *Potamopyrgus antipodarum*, gastropod, mollusk, asexual, pain

Abstract

Anesthesia is used to immobilize organisms for experiments and surgical procedures and to

humanely mitigate pain. Despite its scientific importance, and though invertebrates seem to

experience pain, overarching regulations for the application of anesthesia in invertebrate taxa do

not exist. Indeed, the almost complete absence of legal requirements for invertebrate anesthesia

might explain at least in part why there seems to have been little effort towards developing

widely effective invertebrate anesthesia techniques. Here, we demonstrate that genetic

differences between snails are a major driver of wide variation in menthol anesthesia success in

an important freshwater snail model system, *Potamopyrgus antipodarum*. This is the first

example of which we are aware of a role for genetic variation in anesthesia success in a mollusk.

These findings both illuminate a potential mechanism underlying previously documented

challenges in molluscan anesthesia and will set the stage for powerful and humane manipulative

experiments in *P. antipodarum*.

1. INTRODUCTION

Animal welfare regulations do not typically include invertebrates at least in part because how these organisms process pain is not well understood (AMVA, 2020; Gilbertson & Wyatt, 2016). Nevertheless, invertebrates exhibit clear responses to stimuli that could be perceived as perception of pain (AVMA, 2020). Indeed, Canada, the United Kingdom, and the European Union have animal research regulations that extend protections to cephalopods (Butler-Struben, Brophy, & Crook, 2018), potentially signifying a paradigm shift regarding which taxa deserve formal protections. The basis for animal welfare regulation of cephalopod research is the assumption, rooted in observational studies (e.g., Roper & Hochberg, 1988; Wells, 1978), that cephalopods perceive pain and exhibit emotional responses. This assumption has since been backed up empirically (e.g., Crook, Hanlon, & Walters 2013; Scientific Panel on Animal Health and Welfare, 2005). Researcher perception of the wellbeing of a research organism can also affect study outcomes. For example, as outlined by Poole (1997), conclusions reached from studies using organisms under distress may be unreliable and thus that pain experienced by research organisms can influence results and ultimately study conclusions (also see AVMA, 2020).

Other than cephalopods, there are no other general protections extended to invertebrates for use in research of which we are aware. In the absence of regulatory oversite, given the likelihood that at least some invertebrates perceive pain, and because some experimental procedures are facilitated by or require that the animal is immobilized, there is a clear demand for ability to reliably and effectively apply anesthesia in all invertebrates.

Here, we describe important steps forward in identifying an effective anesthesia

technique for *Potamopyrgus antipodarum* (GRAY, 1843), a New Zealand freshwater snail model

system that is a prominent model system for the evolution of sexual reproduction (e.g., Neiman,

Paczesniak, Soper, Baldwin, & Hehman, 2011), ecotoxicology (e.g., Geiß et al., 2017), invasion

biology (e.g., Donne, Neiman, Woodell, Haase, & Verhaegen, 2020), and host-parasite

interactions (e.g., Bankers et al., 2017). Studies of *P. antipodarum* often incorporate tissue

manipulation (e.g., tentacle severing as a test of tissue regeneration; Krois, Cherukuri,

Puttagunta, & Neiman, 2013) or require determination of male vs. female status (e.g. Jalinsky,

Logsdon, & Neiman, 2020). These studies either require or would be qualitatively improved by

the availability of an effective, safe, and easy-to-use anesthetic.

Menthol crystals are the preferred method for anesthetizing *P. antipodarum* (e.g., Krois

et al., 2013), with the standard approach based on exposure of snails to crushed menthol

crystals for ~90 minutes (McCraw, 1958). While this method can be effective (Krois et al., 2013),

a considerable fraction (often >50%) of exposed *P. antipodarum* do not become anesthetized

(Magnuson, 2018). Similar intraspecific variation and generally low anesthesia efficiency has

been reported in other mollusks (e.g., Butler-Struben, Brophy, & Crook, 2018; Fiorito et al.,

2015).

Magnuson (2018) noted that some *P. antipodarum* lineages (defined as all snails

descended from a single, laboratory-isolated female) seemed to have higher anesthesia success

than others. This observation led us to hypothesize that this lineage effect points to an important

role of genetic background in anesthesia success in these snails. This hypothesis finds indirect

support from reports of genetically based differences in anesthesia efficacy in rodents (e.g.,

Chesler et al., 2003; Moghil, Smith, O'Reilly, & Plourde, 2005; Seltzer, 2014), insects (e.g., Guan,

Scott, & Nash, 2000), and roundworms (Hawasli, Saifee, Liu, Nonet, & Crowder, 2004; Morgan &

Sedensky, 1994) (reviewed in Nash, 2002; Steele, Morgan, & Sedensky, 2007).

Here, we performed the first test of which we are aware of a genetic basis for anesthesia

efficacy in mollusks by subjecting different genetic lineages of common garden-born and raised

P. antipodarum asexual lineages to the same menthol anesthesia protocol. The presence of

significance across-lineage variation in anesthesia success would be consistent with this

hypothesis, suggesting that some *P. antipodarum* genetic backgrounds are more susceptible to

menthol anesthesia than others. This result will be useful to other *P. antipodarum* and mollusk

researchers going forward, highlighting a potential explanation for frequently reported

challenges in anesthesia success and indicating that at least some experiments requiring

anesthesia can succeed if susceptible lineages are used.

2. METHODS

We followed the definition of successful anesthesia provided by Lewbart & Mosley (2011), who

considered the snail to be anesthetized when the animal shows no tentacle withdrawal or body

movement following the gentle scrape of its foot with a needle ("needle test"). For the

experiments presented herein, the anesthetization process begins with the addition of 200 mL of

carbon-filtered tap water to a ~1/2-L plastic cup. We then added the *P. antipodarum* to be

anesthetized to the cup, immediately followed by the addition of 2 g of crushed menthol

crystals. After 90 minutes of exposure, we removed the snails from the menthol water. We

defined anesthesia as successful if the body of the snail was relaxed such that the tentacles,

head, and foot had emerged from the shell and there was no withdrawal or body movement

during the needle test (administered with a 25-gauge needle). We deemed anesthesia as

unsuccessful if the snail was fully contracted into the shell or if the snail was relaxed but failed

the needle test.

We then administered this anesthesia protocol to each of six groups of ten snails (N = 60

snails per lineage) from each of 17 different *P. antipodarum* triploid asexual lineages, for a total

of 102 trials and 1020 snails (Table 1). While each of the 17 lineages was founded by a single

unique asexual female and was cultured separately in its own tank, three pairs of lineages (six of

the 17 lineages) shared descent from the same field-collected grandmother. These founding

grandmothers were sampled from natural populations two generations before the birth of the

snails used here, providing ample time to eliminate any effects of differences in the sampling

environment. The implications are that these three pairs of lineages (each pair defined by

descent from the same grandmother) will be genetically identical (barring *de novo* mutations)

across pair members (hereafter, "clonemates"). We were later able to back up these

assumptions by leveraging SNP data collected for another project (see Discussion). By enabling

comparisons between clonemates (N = three pairs) as well as across genetically distinct lineages

("non-clonemates"; 14 lineages, including those three lineage pairs), our set of lineages thus

provides a powerful test of the extent to which genetic factors influence anesthesia success.

All of the snails in our experiment were born and raised in the same constant-

temperature room, under identical feeding and culture conditions (following standard of care for

P. antipodarum; e.g., Zachar & Neiman, 2013). The snails used for the trials were haphazardly

selected from the adult snails in the 15 L tanks used to house each lineage and were not reused

across trials. Our measure of anesthesia success per trial was the percentage of snails, out of the

ten snails per lineage per trial, that were successfully anesthetized.

2.1. Data analysis.

Because the anesthesia success data were not distributed normally (Kolmogorov-Smirnov test;

test statistic = 0.111, df = 102, *p* = 0.003), we used a non-parametric Kruskal-Wallis test to first

address whether there was a significant effect of lineage founder ("grandmother") on anesthesia

success. A significant outcome of this Kruskal-Wallis analysis will indicate that the descendants of

different grandmothers respond differently to anesthesia. Because the founding grandmothers

constituted separately sampled snails from highly diverse New Zealand populations (e.g.,

Paczesniak, Jokela, Larkin, & Neiman, 2013; Verhaegen, McElroy, Bankers, Neiman, & Haase,

2018), this outcome is consistent with a major role of genetic variation for anesthesia success in

P. antipodarum.

We next used the lineage-level anesthesia success data to address whether the three

pairs of clonemates were significantly more similar in anesthesia success rates than were non-

clonemate pairs. We began by calculating the absolute value of the difference in anesthesia

success rates ("difference scores") between the means of the six success rates measured for

each of the 17 lineages relative to each of the other lineages. For example, this value for Alex 13

relative to Alex 27 is the absolute value of 0.23 (Alex 13 mean success rate) - 0.92 (Alex 27 mean

success rate), equivalent to 0.69. There were a total of three unique pairwise difference scores

for the clonemates (reflecting the three pairs of clonemates) and a total of 136 unique pairwise

difference scores for the non-clonemate comparisons. We then used one-sample sign tests (as

implemented at

https://www.mathcelebrity.com/nonparam.php?pl=Generate+Practice+Problem) relative to a

test value of zero, to determine whether the clonemates and non-clonemate difference scores,

respectively, were significantly different than zero. In a scenario where genetic background is an

important determinant of anesthesia success, the clonemate difference score would not be

significantly different than zero, while the non-clonemate score should be significantly higher

than zero.

3. Results

Our analyses revealed wide across-lineage variation in anesthesia outcome between lineages

descended from different grandmothers (Kruskal-Wallis, $p < 0.001$; Figure 1), ranging from a

median of 95% of all snails anesthetized for several different lineages from lake Alexandrina to

~8% for Grasmere 59 (Table 1). This result is as is expected if there is a major genetic component

to anesthesia success in *P. antipodarum*.

The difference score between clonemates was not significantly different than zero (one-

sample sign test; $p = 0.125$), with the caveat that we had only three comparisons and thus low

statistical power. By contrast, the difference score between non-clonemates was significantly

higher than zero ($p = 0.002$) and was nearly twice as high (mean difference score 0.340 +/- 0231

SD) as the mean difference score for the clonemates (mean difference score = 0.173 +/- 0.132

SD; Fig. 2). These results are also consistent with a scenario where genetic factors influence

anesthesia success in *P. antipodarum*.

4. Discussion

Together, our data indicate that genetic background is a major contributor to menthol anesthesia success in *P. antipodarum*, an important model system in evolutionary biology, ecology, and ecotoxicology. This is the first demonstration of which we are aware of a genetic component to variation in anesthesia efficacy in mollusks. One obvious practical implication of these results for *P. antipodarum* researchers is the opportunity to select lineages with higher anesthesia susceptibility for experiments that require anesthetics.

After our experiment had concluded, we were fortuitously able to perform a powerful unplanned independent validation of the likelihood that genetic variation does underlie across-lineage variation in anesthesia success in *P. antipodarum* by leveraging single-nucleotide polymorphism (SNP) data generated for a different project (Neiman et al., unpublished). This other project provided SNP data for 14 of the 17 lineages used here, which were genotyped at the same 48-SNP set used by Verhaegen et al. (2018). We followed the DNA extraction procedure described in Sharbrough, Luse, Boore, Logsdon, and Neiman (2018), and applied the same genotyping approach and data analyses reported in Verhaegen et al. (2018) to assign multilocus SNP genotypes to these 14 lineages (Table 1).

There were seven distinct SNP genotypes across the 17 lineages. These SNP data demonstrated that the one pair of clonemates that were genotyped (Alex 19_1 and Alex 19_2) indeed shared an otherwise unique SNP genotype. These data also revealed that most other non-clonemate lineages themselves had unique genotypes. Finally, we discovered that the two sets of lineages with different grandmothers that nevertheless shared SNP genotypes 4 and 26,

respectively, had strikingly similar anesthesia success rates relative to other lineages with

different SNP genotypes (Fig. 1).

To our knowledge, this evidence for genetic variation for anesthetic success in *P.*

antipodarum is the first report of such a phenomenon in mollusks. That genetic background

influences anesthesia success in a wide variety of common model systems like flies, worms, and

mice suggest that the same might be expected for other mollusk taxa.

Two important questions worthy of future study emerge from our results. First, what is

the genetic basis of the across-lineage variation in anesthesia success? Second, even though

clonemates seem to share similar anesthesia success outcomes relative to non-clonemates, we

do not see identical responses (i.e., no examples of 100% success or 100% failure) within

clonemates.

The first question can be addressed by using techniques such as RNA sequencing and

whole-genome sequencing to identify gene expression differences and genetic variants

differentiating lineages with very distinct anesthesia responses. Because multiple studies

indicate that asexual *P. antipodarum* lineages from the same lake population are typically closely

related relative to asexual lineages from other lakes (e.g., Paczesniak et al., 2013), a powerful

experiment might be one that compared genomic sequence from lineages from the same lake

with different anesthesia responses. A good example of such lineages in the experiment we

describe here are provided by low-responding lineages from lake Alexandrina like Alex 13

(23.33% anesthesia success) and Alex 21 (15% anesthesia success) as compared to Alex 27

(91.67% anesthesia success) and Alex 32 and 36 (95% anesthesia success).

The second question, regarding variation in anesthesia success even within clones and

clonemates, hints that non-genetic factors (e.g., age, body condition) could influence anesthesia

outcomes (e.g., Gao et al., 2018; Guenette & Lair, 2006; Larsson & Wahlström, 1998). While we

made a distinct effort to minimize age effects by choosing the largest snails possible from each

lineage, we did not know the ages of individual snails. Nor are we able to exclude the possibility

that across-snail differences in health or body condition could have come into play. Future

experiments could address these possibilities by comparing anesthesia success within a lineage

as a function of snail age or in the context of experimental manipulation of factors like food

availability.

Acknowledgements:

We gratefully acknowledge Molly Gallagher, Sydney Stork, Benjamin Ripperger, and Alexander

Kern for snail care. John Logsdon contributed to snail collections. The University of Iowa

Department of Biology Honors program, the Iowa Center for Undergraduate Research, and Rick

and Linda Maxson logistically and financially supported several undergraduate researchers

involved in the project. The SNP genotyping described here was also supported by NSF-MCB

1122176 and NSF-DEB 1731657, with contributions to lineage selection by collaborator Kristi

Montooth.

References:

American Veterinary Medical Association. 2020. The AMVA Guidelines for the Euthanasia of

Animals: 2020 Edition.

Bankers, L., Fields, P., McElroy, K. E., Boore, J. L., Logsdon, J. M., Jr., & Neiman, M. (2017).

Genomic evidence for population-specific responses to coevolving parasites in a New

Zealand freshwater snail. *Molecular Ecology*, *26*, 3663-3675.

Butler-Struben, H. M., Brophy, S. M., & Crook, R. J. (2018). *In vivo* recording of neural and

behavioral correlates of anesthesia induction, reversal, and euthanasia in cephalopod

molluscs. *Frontiers in Physiology*, *9*, 109.

Chesler, E., Ritchie, J., Kokayeff, A., Lariviere, W. R., Wilson, S. G., & Moghil, J. S. (2003).

Genotype-dependence of gabapentin and pregabalin sensitivity: the pharmacogenetic

mediation of analgesia is specific to the type of pain being inhibited. *Pain*, *106*, 325-335.

Crook, R. J., Hanlon, R. T., & Walters, E. T. (2013). Squid have nociceptors that display widespread

long-term sensitization and spontaneous activity after bodily injury. Jour*nal of*

Neuroscience, *33*, 10021-10026.

Donne, C., Neiman, M., Woodell, J. D., Haase, M., & Verhaegen, G. (2020). A layover in Europe:

reconstructing the invasion route of asexual lineages of a New Zealand snail to North

America. *Authorea* doi: 10.22541/au.158456440.01903363 (Accepted pending minor

revisions in *Molecular Ecology*).

d'Ovidio, D., Monticelli, P., Santoro, M., & Adami, C. (2019). Immersion anesthesia with ethanol

in African giant land snails (*Acathina fulica*). *Heliyon*, *5*, e01546.

Fiorito, G., Affuso, A., Basil, J., Cole, A., de Girolamo, P., D'Angelo, L., Dickel, L., Gestal, C., Grasso,

F., Kuba, M., Mark, F., Melillo, D., Osorio, D., Perkins, K., Ponte, G., Shashar, N., Smith, D.,

Smith, J., & Andrews, P. L. R. (2015). Guidelines for the care and welfare of cephalopods

in research- a consensus based on an initiative by CephRes, FELASA, and the Boyd group.

Laboratory Animals, *49*, 1-90.

Gao, J., Hu, Z., Shi, L., Li, N., Ouyang, Y., Shu, S., Yao, S., & Chen, X. (2018). HCN channels

contribute to the sensitivity of intravenous anesthetics in developmental mice.

Oncotarget, *9*, 12907–12917.

Geiß, C., Ruppert, K., Askem, C., Barroso, C., Faber, D., Ducrot, V., Holbech, H., Hutchinson, T. H.,

Kajankari, P., Kinnberg, K. L., Lagadic, L., Matthiessen, P., Morris, S., Neiman, M.,

Penttinen, O.-P., Sanchez-Marin, P., Tiegeler, M., Weltje, L., & Oehlmann, J. (2017).

Validation of the OECD Reproduction Test Guideline with the New Zealand mudsnail

Potamopyrgus antipodarum using trenbolone and prochloraz. *Ecotoxicology*, *26*, 370-

382.

Gilbertson, C. R., & Wyatt, J. D. (2016). Evaluation of euthanasia techniques for an invertebrate

species, land snails (*Succinea putris*). *Journal of the American Association for Laboratory*

Animal Science, *55*, 577-581.

Guan, Z., Scott, R. L., & Nash, H. A. (2000). A new assay for the genetic study of general

anesthesia in *Drosophila melanogaster*: Use in analysis of mutations in the X-

chromosomal 12E region. *Journal of Neurogenetics*, *14*, 25-42.

Guenette, S. A., & Lair, S. (2006). Anesthesia of the leopard frog, *Rana pipiens*: A comparative

study between four different agents. *Journal of Herpetological Medicine and Surgery*, *16*,

38-44.

Hawasli, A. H., Saifee, O., Liu, C., Nonet, M. L., & Crowder, C. M. (2004). Resistance of volatile

anesthetics by mutations enhancing excitatory neurotransmitter release in

Caenorhabditis elegans. *Genetics*, *168*, 831-843.

Krois, N. R., Cherukuri, A., Puttagunta, N., & Neiman, M. (2013). Higher rate of tissue

regeneration in polyploid asexual vs. diploid sexual freshwater snails. *Biology Letters*, *9*,

20130422.

Larsson, J., & Wahlström, G. (1998). The influence of age and administration rate on the brain

sensitivity to propofol in rats. *Acta Anaesthesiologica Scandinavica, 42*, 987-994.

Lewbart, G. A., & Mosley, C. (2011). Clinical anesthesia and analgesia in invertebrates. *Journal of

Exotic Pet Medicine*, *21*, 59-70.

Lively, C. M. 1987 Evidence from a New Zealand snail for the maintenance of sex by parasitism.

Nature, *328*, 519-521.

Logsdon, J. M., Neiman, M., Boore, J., Sharbrough, J., Bankers, L., McElroy, K., Jalinsky, J., Fields,

P., & Wilton, P. (2017). A very recent whole genome duplication in *Potamopyrgus*

antipodarum predates multiple origins of asexuality & associated polyploidy. *PeerJ*

Preprints, *5*, e3046v3041.

Magnuson, R. (2018). An exploration of alternative anesthesia in the difficult-to-anesthetize New

Zealand freshwater snail *Potamopyrgus antipodarum*. Iowa Research Online.

McCraw, B. M. (1958). Relaxation of snails before fixation. *Nature*, *181*, 575.

Moghil, J. S., Smith, S. B., O'Reilly, M. K., Plourde, G. (2005). Influence of nociception and stress-

induced antinociception on genetic variation in isoflurane anesthetic potency among

mouse strains. *Anesthesiology*, *103*, 751-758.

Morgan, P. G., & Sedensky, M. M. (1994). Mutations conferring new patterns of sensitivity to

volatile anesthetics in *Caenorhabditis elegans*. *Anesthesiology*, *81*, 888-898.

Nash, H. A. (2002). *In vivo* genetics of anaesthetic action. *British Journal of Anaesthesia*, *89*, 143-

155.

Neiman, M., Larkin, K., Thompson, A. R., & Wilton, P. (2012). Male offspring production by

asexual *Potamopyrgus antipodarum*, a New Zealand snail. *Heredity, 109*, 57-62.

Neiman, M., Jokela, J., & Lively, C. M. (2005). Variation in asexual lineage age in *Potamopyrgus*

antipodarum, a New Zealand snail. *Evolution*, *59*, 1945-1952.

Neiman, M., Paczesniak, D., Soper, D. M., Baldwin, A. T., & Hehman, G. (2011). Wide variation in

ploidy level and genome size in a New Zealand freshwater snail with coexisting sexual and

asexual lineages. *Evolution*, *65*, 3203–3216.

Paczesniak, D., Jokela, J., Larkin, K., & Neiman, M. (2013). Discordance between nuclear and

mitochondrial genomes in sexual and asexual lineages of the freshwater snail

Potamopyrgus antipodarum. *Molecular Ecology*, *22*, 4695-4710.

Poole T. (1997). Happy animals make good science. *Laboratory Animals*, *31*, 116-124.

Roper C.F.E., & Hochberg F.G. (1988). Behavior and systematics of cephalopods from Lizard

Island, Australia, based on color and body patterns. *Malacologia*, *29*, 153-193.

Scientific Panel on Animal Health and Welfare (AHAW). 2005. Opinion of the Scientific Panel on

Animal Health and Welfare (AHAW) on a request from the Commission related to the

'Aspects of the biology and welfare of animals used for experimental and other scientific

purposes'. *EFSA Journal*, *292*, 1-136.

Seltzer, Z. (2014). Nothing in pain makes sense except in the light of genetics. *Pain*, *2014*, 841-

842.

Sharbrough, J., Luse, M., Boore, J. L., Logsdon, J. M., Jr., & Neiman, M. (2018). Radical amino acid

mutations persist longer in the absence of sex. *Evolution*, *72*, 808-824.

Steele, L. M., Morgan, P. G., & Sedensky, M. M. (2007). Genetics and the mechanisms of action

of inhaled anesthetics. *Current Pharmacogenomics*, *5*, 125-141.

Verhaegen, G., McElroy, K., Bankers, L., Neiman, M., & Haase, M. (2018). Adaptive phenotypic

plasticity in a clonal invader. *Ecology and Evolution*, *8*, 4465-4483.

Wells, M. J. (1978). Octopus: Physiology and behavior of an advanced invertebrate. London, UK:

Chapman and Hall.

Zachar, N., & Neiman, M. (2013). Profound effects of population density on fitness-related traits

in an invasive freshwater snail. *PLoS ONE*, *8*, e80067.

Figure Legends and Table Caption

Figure 1: Median percent of successfully anesthetized snails (6 trials of 10 snails each) subjected

to the standard *P. antipodarum* menthol anesthesia protocol per each of 17 lineages founded by

14 different grandmothers. Triangle symbols (vs. circles) are used to denote the three lineage

pairs that each share a grandmother. The numbers above the 14 of the 17 lineages that were

genotyped indicate SNP genotype.

Figure 2: Median pairwise difference score in anesthesia success for clonemate vs. non-

clonemate lineages. The non-clonemate pairwise difference scores were significantly higher than

zero while the clonemate pairwise difference scores were not significantly higher than zero.

Table 1. Description of the *P. antipodarum* asexual lineages used in this study.

Lake of Origin	Lineage	Grandmother	Mean Anesthesia Success % (+/- SD)	Genotyped	SNP Genotype
Alexandrina	Alex 13	Alex 13	23.33 +/- 15.01	N	NA
Alexandrina	Alex 19_1	Alex 19	50.00 +/- 6.32	Y	3
Alexandrina	Alex 19_2	Alex 19	53.33 +/- 10.33	Y	3
Alexandrina	Alex 21_1	Alex 21	15.00 +/- 8.37	Y	2
Alexandrina	Alex 21_2	Alex 21	35.00 +/- 12.25	N	NA
Alexandrina	Alex 27	Alex 27	91.67 +/- 11.69	Y	4
Alexandrina	Alex 31	Alex 31	48.33 +/- 7.53	N	NA
Alexandrina	Alex 32	Alex 32	95.00 +/- 8.37	Y	4
Alexandrina	Alex 36	Alex 36	95.00 +/- 5.48	Y	5
Alexandrina	Alex 39	Alex 39	93.33 +/- 8.16	Y	4
Grasmere	Grasmere 49_1	Grasmere 49	76.67 +/- 15.01	Y	26
Grasmere	Grasmere 49_2	Grasmere 49	48.33 +/- 9.83	Y	NA
Grasmere	Grasmere 54	Grasmere 54	83.33 +/- 13.66	Y	26
Grasmere	Grasmere 59	Grasmere 59	8.33 +/- 7.53	Y	24
Gunn	Gunn 6	Gunn 6	36.67 +/- 20.66	Y	29
Gunn	Gunn 7	Gunn 7	71.67 +/- 16.02	Y	26
Gunn	Gunn 9	Gunn 9	78.33 +/- 14.72	Y	31