1 Phylogenomics of 8,839 *Clostridioides difficile* genomes reveals recombination-driven

2 evolution and diversification of toxin A and B

Michael J. Mansfield[1*], Benjamin J-M Tremblay[1*], Ji Zeng[2,3], Xin Wei[1], Harold Hodgins[1], Jay

Worley[4,5], Lynn Bry[4,6], Min Dong[2,3,#], Andrew C. Doxey[1,#]

[1]Department of Biology, David R. Cheriton School of Computer Science, and Waterloo Centre

for Microbial Research, University of Waterloo, 200 University Ave. West, Waterloo, Ontario,

N2L 3G1, Canada.

[2] Department of Urology, Boston Children's Hospital, Boston, Massachusetts, USA

[3] Department of Microbiology, Harvard Medical School, Boston, Massachusetts, USA

[4] Massachusetts Host-Microbiome Center, Department of Pathology, Brigham and Women's

Hospital, Harvard Medical School, Boston, MA, USA

[5] National Center for Biotechnology Information, National Library of Medicine, National

Institutes of Health, Bethesda, MD, USA

[6] Division of Infectious Diseases, Department of Medicine, Brigham and Women's Hospital,

Harvard Medical School, Boston, Massachusetts, USA

[*]Co-first authors

[#]Correspondence should be addressed to A.C.D. (acdoxey@uwaterloo.ca) and M.D.

(min.dong@childrens.harvard.edu)

Abstract

Clostridioides difficile is the major worldwide cause of antibiotic-associated gastrointestinal infection. A pathogenicity locus (PaLoc) encoding one or two homologous toxins, toxin A (TcdA) and toxin B (TcdB) is essential for *C. difficile* pathogenicity. However, toxin sequence variation poses major challenges for the development of diagnostic assays, therapeutics, and vaccines. Here, we present a comprehensive phylogenomic analysis 8,839 *C. difficile* strains and their toxins including 6,492 genomes that we assembled from the NCBI short read archive. A total of 5,175 *tcdA* and 8,022 *tcdB* genes clustered into 7 (A1-A7) and 12 (B1-B12) distinct subtypes, which form the basis of a new method for toxin-based subtyping of *C. difficile*. We developed a haplotype coloring algorithm to visualize amino acid variation across all toxin sequences, which revealed that TcdB has diversified through extensive homologous recombination throughout its entire sequence, and formed new subtypes through distinct recombination events. In contrast, TcdA varies mainly in the number of repeats in its C-terminal repetitive region, suggesting that recombination-mediated diversification of TcdB provides a selective advantage in *C. difficile* evolution. The application of toxin subtyping is then validated by classifying 351 *C. difficile* clinical isolates from Brigham and Women's Hospital in Boston, demonstrating its clinical utility. Subtyping partitions TcdB into binary functional and antigenic groups generated by intragenic recombinations, including two distinct cell-rounding phenotypes, whether recognizing frizzled proteins as receptors, and whether can be efficiently neutralized by monoclonal antibody bezlotoxumab, the only FDA-approved therapeutic antibody. Our analysis also identifies eight universally conserved surface patches across the TcdB structure, representing ideal targets for developing broad-spectrum therapeutics. Finally, we established an open online database (DiffBase) as a central hub for collection and classification of *C. difficile* toxins, which will help clinicians decide on therapeutic strategies targeting specific toxin variants, and allow researchers to monitor the ongoing evolution and diversification of *C. difficile*.

Key words: C. difficile, toxin, TcdA, TcdB, toxin A, toxin B, recombination, subtype, bezlotoxumab, frizzled

Introduction

Clostridioides difficile (formerly *Clostridium difficile*) is a diverse group of Gram-positive spore-forming anaerobic bacteria[1]. Toxigenic strains have become important opportunistic pathogens to humans. Their spores are widespread and can colonize human and animal colons after disruption of the gut microflora, most notably due to antibiotic treatment. *C. difficile* infection (CDI) results in a range of symptoms from self-limiting diarrhea to severe pseudomembranous enterocolitis and death[2–7]. It is the most frequent cause of healthcare-associated gastrointestinal infections across developed countries worldwide[2–5,8].

Ribotyping (RT), which compares intergenic spacers between ribosomal RNA genes, is widely utilized to categorize *C. difficile* linages[5,9]. Various other methods including multilocus sequence typing based on allelic variation of housekeeping genes and whole genome sequencing analysis have also been adopted to further discriminate strains[5,9–13]. Phylogenetic analyses revealed a growing diverse population[1,14–16]. In recently years, there is an emergence and spreading of various epidemic hypervirulent strains such as the RT027 clonal lineage, which first caused outbreaks in 2000-2003 in North America and is associated with increased disease severity and mortality[17–20]. RT078 is an emerging hypervirulent lineage which is also the dominant type found in domesticated animals[21,22]. There are also geographic differences, for instance, RT017 has become a dominant lineage in Japan and Korea[23].

The major virulence factors in toxigenic *C. difficile* strains are two homologous large protein toxins, TcdA (~300 kDa) and TcdB (~270 kDa)[24–27]. Nontoxigenic *C. difficile* strains without these toxins exist and can colonize humans and animals, but do not cause diseases[28]. TcdA and TcdB share overall ~66% sequence similarity and belong to the large clostridial toxin (LCT) family, which include TcsH and TcsL in *Paeniclostridium sordellii*, Tcnα in *Clostridium novyi*, and TpeL in *Clostridium perfringens*[5,6,8,9,24,25,29–31]. TcsH and TcsL can be considered orthologs of TcdA and TcdB, respectively, with TcsH sharing ~77% sequence identity with TcdA and TcsL sharing ~76% identity with TcdB[32] (Fig. S1). TcdA and TcdB share a protein domain architecture consisting of an N-terminal glucosyltransferase domain (GTD), followed by a cysteine protease domain (CPD), an intermingled membrane translocation delivery domain and receptor-binding domain (DRBD),

and a large C-terminal combined repetitive oligopeptides domain (CROPs) (Fig. S1). After

binding, endocytosis, and translocation across endosomal membranes into the cytosol of host cells,

these toxins glucosylate and inactivate host Ras/Rho family of small GTPases, leading to

disruption of the actin cytoskeleton, cell rounding, and ultimately cell death[33].

TcdA and TcdB were first identified in the 1990s, and the toxin sequences from a reference strain

(VPI10463) have been widely used as the standard in diagnostic and therapeutic development.

However, sequence variations in the toxin genes exist across *C. difficile* strains and could affect

receptor-binding specificity, preferences toward distinct small GTPases, overall toxicity, and

antigenicity. For instance, strains such as R20291 (belonging to RT027) produces a TcdB variant

with ~8% of residue differences from the reference TcdB, which exhibited a significant impact on

its immunogenicity: mice immunized with the reference TcdB developed resistance to the same

TcdB, but all died when challenged with this variant TcdB[34], and several antibodies raised against

the reference TcdB, including the FDA approved therapeutic antibody bezlotoxumab, either do

not recognize or have lower efficacy against this TcdB variant[34–36]. Furthermore, this TcdB variant

also loses the ability to recognize frizzled (FZD) proteins, which are one of the major receptors for

the reference TcdB, due to residue changes at the FZD-binding interface[35,37–40].

These toxin variations pose a significant challenge for developing effective broad-spectrum

diagnostic assays, therapeutic antibodies, and vaccines. Understanding variations in toxins is a key

step to address this challenge and may also reveal their potential evolutionary paths and functional

differences. A toxinotyping method has been previously developed utilizing PCR-based

amplification of toxin gene fragments and analyzing polymorphism with restriction enzyme

digestions, which can distinguish over 34 toxinotypes[41,42]. Although toxinotyping highlights the

variation among toxin genes, it lacks the resolution to understand the molecular basis for

diversification of toxins and sequence-function relationships.

Rapid growth of genomic sequencing of *C. difficile* strains in recent years provides an opportunity

to analyze and categorize the diversification of TcdA and TcdB with single residue resolution.

Here we performed a comprehensive analysis of nearly all available *C. difficile* TcdA and TcdB

sequences, including assembly and analysis of 6,492 new genomes, with the goal to 1) build a

comprehensive central database of *C. difficile* toxin sequences; 2) better understand the
mechanisms underlying TcdA and TcdB diversification; and 3) develop a system to classify TcdA
and TcdB into subtypes that allow clinicians and researchers to categorize and predict functional-
immunological variations of any future sequenced *C. difficile* isolates.
121

Results

Collection of TcdA and TcdB sequences across 8,839 C. difficile genomes

To build a comprehensive database of TcdA and TcdB sequences, we combined data from NCBI

GenBank and the NCBI short-read archive (SRA). From 2,347 *C. difficile* genomes in GenBank,

we identified an initial set of 1,633 *tcdA* and 2,028 *tcdB* genes. We then developed a computational

pipeline for automated retrieval of *C. difficile* genomes from the SRA, *de novo* genome assembly,

genome annotation, and extraction of *tcdA* and *tcdB* genes (see Methods). Using this pipeline, we

assembled the genomes of 6,492 *C. difficile* isolates and identified an additional 3,542 *tcdA* and

5,994 *tcdB* genes (Table 1). Combining both sources, we identified 5,175 TcdA and 8,022 TcdB

encoding sequences.

We then carried out alignments of all toxin protein sequences. The TcdB alignment covered the

entire sequence (1-2366), with 712 (30%) of the positions showing variations across all domains.

The TcdA alignment possessed much lower variation than TcdB within the 1-1874 region as it had

only 168 (9%) variable sites, but its CROPs domain (1831-2710) contained an extremely high

degree of variation in the number length of repeats: from 3 repeats in the shortest variant to 45 in

the longest variant, and 32 in the reference TcdA variant from VPI 10463 (Fig. S2). This is likely

generated by homologous recombination due to the repetitive nature of this region. The CROPs

domain is composed of long repeats (LRs) of ~30 residues and short repeats (SRs) of ~19-24

residues[27]. The CROPs domain in TcdA is not only repetitive at a protein sequence level, but also

showed a high degree of repetitiveness at a DNA level, whereas the repetitiveness of the CROPs

domain in TcdB is largely limited to the protein level[43,44], which may account for frequent

recombination in TcdA-CROPs but not in TcdB-CROPs.

Classifying TcdA and TcdB into subtypes

In total, there were 116 unique TcdA protein sequences and 212 unique TcdB protein sequences.

We then clustered these sequences into distinct subfamilies ("subtypes") using average linkage

hierarchical clustering (see Methods). Analysis of TcdB is based on full-length sequences, but

TcdA is limited to the 1-1874 region to avoid the highly variable CROPs domain. In addition, we

also included TcsH and TcsL sequences in our analysis. Clustering produced 7 distinct TcdA

subtypes which we labeled A1-A7, and 12 distinct TcdB subtypes which we labeled B1-B12, with

the subtype number ranked based on their total frequency of occurrence in GenBank and NCBI-

SRA (Fig. 1). Each unique sequence was then further numbered following a period within its

subtype (e.g. B1.1, 1.2, 1.3, etc.). Sequences within the same TcdA and TcdB subtype demonstrate

strong pairwise similarities, and weak similarities between subtypes (Fig. 1a, 1d). Quantitative

analysis revealed that thresholds of 99.4% (TcdA) and 97% (TcdB) can be used to effectively

assign toxin sequences to these subtypes (Fig. S3). We then selected one representative sequence

for each subtype and carried out phylogenic analysis and pairwise comparison. TcdA subtypes A1

to A6 possess higher similarities (>97.9%) and clustered together, with A7 forming a divergent

lineage (Fig. 1b, 1c). A7 is a unique sequence with only 85.3% to 85.6% identity to others (Fig.

1c). The entire TcdA family was further outgrouped by TcsH as expected (Fig. 1b). TcdB also

formed a monophyletic family that was outgrouped by TcsL and a second lineage of TcsL-related

proteins (Fig. 1e). TcdB subtypes can be subdivided into three groups, one including B6, B7, B4,

B8, a second including B9, B2, B10, B11, and a third including B12, B1, B5, and B3 (Fig. 1e).

The lowest identity among TcdB subtypes is 85.3% (between B7 and B12, Fig. 1f). A7, B10, B11,

and B12 represent rare divergent subtypes recently reported: A7 is in strain RA09-70, which does

not express TcdB[45]; B10, B11, and B12 were identified recently from strains CD10-165, CD160,

and 173070, respectively[45], and all three strains do not express TcdA.

By mining unassembled *C. difficile* genomes from the SRA, we were able to discover 125 TcdA

and TcdB protein sequences that were not represented in GenBank. Most novel toxin variants

clustered into subtypes A1 (N = 25), B1 (N = 52), and subtypes A2 (N = 10) and B2 (N = 12)

(Table S1). However, three highly divergent TcdA variants identified from SRA datasets formed

new subtypes not represented in GenBank. These include subtypes A4 from strain ECDC-088

(SRS1486236), A5 from strain ECDC-009 (SRS1486256), and A6 from strain L;13.7548369.T

(SRS1486661), all of which are clinical isolates. All three of these strains contained

truncated/partial TcdB variants which represent putative pseudogenes.

To link our subtyping with known clinical *C. difficile* strains, we manually curated subtype

assignments for a set of 63 *C. difficile* strains selected from the literature, which covers known

toxinotypes, and compared subtypes with toxinotypes, ribotypes, and whether the strain produces

the third toxin known as *C. difficile* transferase toxin (CDT) (Table S2). The majority express an

A1/B1 subtype combination and include reference strains 630 and VPI 10463 that express the

widely used standard TcdA and TcdB sequence (defined as A1.1 and B1.1, Table S2). The group

that expresses a combination of A2/B2 is the second largest and includes hypervirulent RT027

strains R12087 and R20291. The group expressing A3/B3 include strains (e.g. M120 and NAP07)

classified as RT078. Subtype B4 is mainly expressed in strains (e.g. 1470) belonging to RT017,

which lacks TcdA. Other pairings in the table include A2/B9, A3/B5, A2/B6, and A1/B4. The

table includes many strains that do not express functional TcdA, which can express B1, B2, B3,

B4, B6, B7, B8, B10, B11, or B12; one strain that only expresses TcdA but not TcdB (A7 in RA09-

70); four strains that only express CDT; and one strain (SLO037) that expresses none of the three

toxins. This table represents a small portion of *C. difficile* strains and a full list of a total 1640 *C.*

difficile strains from the NCBI database with their toxin subtypes noted is included as Table S3.

In general, phylogenetic subtyping of *C. difficile* toxins correlated well with previously identified

toxinotypes, but at greater resolution by analyzing TcdA and TcdB separately (Table S2, see

Discussion). There was less congruence with ribotypes, however, as different subtypes were found

in the same ribotype strains, and the same subtype was found in different ribotype strains.

Therefore, neither toxinotype nor ribotype were able to accurately categorize toxins based on

phylogenetic relationships (Table S2). Subtyping was capable of capturing the full phylogenetic

diversity of TcdA and TcdB available in previously known and new strains.

Distribution of toxin subtypes across the C. difficile phylogeny

To evaluate the phylogenomic distribution of toxin subtypes across *C. difficile*, we constructed a

whole-genome based phylogeny of 1,934 complete *C. difficile* genomes based on 14,194 SNP

positions across 88 conserved marker genes (Fig. 2a, Table S3) (see Methods). The genome tree

is highly consistent with known phylogenetic relationships, as the previously identified clades 1-

5 are represented by distinct lineages[14] (Fig. 2a). Two of the three divergent environmental

lineages C-I and C-II are also present as divergent branches (Fig. 2a).

A total of 1,640 (84.8%) *C. difficile* strains were found to encode TcdA and/or TcdB, while the

remainder (294, 15.2%) lack toxin genes. The predicted toxin subtypes across the *C. difficile*

genome tree demonstrate strong clade associations, and therefore are highly congruent with strain

phylogenetic relationships. The congruency between subtype and phylogeny provides further

support for our toxin classification (Fig. 2a). For example, subtype A1/B1 which includes

reference strains 630 and VPI 10463 is most common among toxin-containing strains (979, 59.7%)

and associated with clade 1 (Fig. 2b). A2/B2 was second most common and associated with clade

2, A3/B5 with clade 3, -/B4 with clade 4, and A3/B3 with clade 5. Also prevalent were types -/B1,

A1/-, and A2/B6 (Fig. 2b). Deviations from the A1/B1 toxin type are often associated with the

emergence of numerous hypervirulent and epidemic outbreak strains such as A2/B2 (RT027),

A3/B3 (RT078), and -/B4 (RT017) (Fig. 2a). Interestingly, the highly divergent environmental

lineages encode the highly divergent TcdB subtypes B10 and B11 (C-I) and B12 (C-II) (Fig. 2a,

Table S2). This is consistent with an early divergence of B10-B12 in *C. difficile* evolution,

predating the emergence of TcdB subtypes found in the other clinical strains.

Interestingly, we also observed rare lateral transfer events involving only one of the two toxin

genes to create hybrid strains containing new subtype combinations. Examples include the

spontaneous emergence of an A1/<u>B4</u> strain within clade 1, and the emergence of an <u>A1</u>/B2 strain

in clade 2 (Fig. 2a). Thus, through lateral transfer and homologous recombination, subtype B4 has

likely replaced B1 in a clade 1 strain, and subtype A1 has likely replaced A2 in a clade 2 strain.

Furthermore, we observed many independent clades containing *tcdA-*/*tcdB-* *C. difficile* strains

(e.g., see six lineages marked by asterisks in Fig. 2a). This is consistent with previously reported

"defective" toxin clades[46], and indicates numerous independent losses of the pathogenicity locus

throughout *C. difficile* evolution.

Toxin subtyping of an independent dataset of clinical C. difficile isolates

As an independent test dataset for our toxin subtyping method, we examined 351 genomes of *C.*
difficile isolates derived from a clinical cohort from Brigham and Women's Hospital (BWH) in
Boston (Fig. 2c)[47]. As they were not included in our initial database, they are ideal for testing the
robustness and effectiveness of our subtype classification. All identified toxins could be accurately
assigned to our reference sequences, with most (97%) aligning with 100% identity to our database,
and the remainder aligning with >= 99.8% identity. Out of 351 total strains, 62 (17.7%) were toxin
deficient, while 289 (82.3%) contained TcdA and/or TcdB genes (Table S4). Of these, there were
12 distinct subtype combinations, with frequencies similar to those observed in the NCBI dataset.
A1/B1 strains were most common (N = 222), followed by A2/B2 (N = 24), -/B4 (N = 11), and
A3/B3 (N = 10) (Fig. 2c). Therefore, our method was able to rapidly and automatically classify a
large dataset of 351 clinically relevant *C. difficile* isolates, with all sequences represented in our
current classification.

Intragenic recombination drives TcdB diversification

We next focused on understanding the evolution of TcdA and TcdB variants and mechanisms for
their diversification. To visualize global patterns of variation within TcdA and TcdB, we
developed a haplotype coloring algorithm (https://github.com/doxeylab/haploColor) based on
previous methods for genome visualization[48] First, sequences are painted black where they
matched the reference sequence (i.e., B1.1). Then, remaining positions were painted different
colors where they matched selected other subtypes (Fig. 3a): blue when matching B3.1, gold when
matching B6.1, and green when matching TcsL. The result of this algorithm applied to the TcdB
alignment revealed a striking block-like and highly mosaic pattern of amino acid variation, which
strongly indicates recombination between subtypes (Fig. 3a). B1, B3, and B5 are composed of a
B1-like variation (black) pattern across their full-length sequences, while B6 and B7 are composed
of a B6-like pattern (gold) across their full-length sequences. B2, B4, B8, and B9, however, possess
a mosaic combination of B1-like and B6-like patterns. B4, B6, B7, and B8 share a distinct B6-like
pattern of amino acid variation across their N-terminal region including the GTD and CPD
domains, but when examining the DRBD, the B6-like pattern is shared by a different set of
subtypes (B2, B6, B7, B10, and B11). These patterns indicate ancestral within-gene ("intragenic")

recombination events involving distinct regions of TcdB. As a statistical test of recombination, we
further performed phylogenetic network analysis using SplitsTree[49]. Consistent with patterns of
amino acid variation and per-domain phylogenetic analysis, network analysis revealed significant
evidence of recombination within TcdB ($p = 0$; Phi test for recombination) (Fig. S4). In contrast
to TcdB, TcdA (1-1874) produced homogeneous patterns of variation across each subtype (Fig.
S5) and did not display evidence of recombination in network analysis ($p = 0.186$) (Fig. S4),
indicating that recombination occurs frequently only in TcdB, but not in TcdA.

We further performed separate phylogenetic analyses of each domain (GTD, CPD, DRBD, and
CROP) of TcdB (Fig. 3b). The phylogenetic tree of each domain produced two main groups
(labeled i and ii), which correspond with the B1-like and B6-like patterns revealed in the alignment
visualization (Fig. 3a). Each subtype can therefore be described as a chimeric combination of type
"i" (B1-like) or type "ii" (B6-like) domains (Fig. 3c). Based on the per-domain phylogenetic
relationships and recombination patterns, we formulated a potential evolutionary model for the
origin of TcdB subtypes (Fig. 3d). An early TcdB ancestor split into two main groups: (i) B1, B5,
and B3; and (ii) B6 and B7. Subtype B2 likely originated by a recombination event fusing an
ancestral type i and type ii toxin. B9 likely originated from a recombination event between B1 and
B2, B4 from a recombination event between B1 and a type ii toxin, and B8 from a recombination
event between B3 and a type ii toxin. Subtypes B10-B12, which are rare variants recently
identified, are early diverging lineages since they consistently outgrouped other subtypes in
phylogenetic analysis (Fig. 3b), consistent with their divergent lineages among other strains (Fig.
2a).

In addition to these major ancestral recombination events, we also identified a considerable degree
of "microrecombination" events involving exchange of small segments between subtypes. For
example, a single TcdB sequence (B1.59) from subtype B1 has acquired an N-terminal segment
that is clearly derived from subtype B2 or B9 (Fig. 3a, Figure S6). This unique TcdB gene, which
appears to be the result of a spontaneous recombination event between a B1 and B2-containing
strain, is derived from a newly assembled clinical isolate from a Fidaxomicin clinical trial
(SRS1378602). A second similar example is B1.58 from a clinical isolate (ECDC-040,
SRS1486176), which has acquired a DRBD and CROPS segment from a B2-containing strain

(Figure S6). Fourteen such cases of microrecombination including these are depicted in Figure S6.
TcdB in particular appears to have diversified through an extensive degree of intragenic
recombination involving both large and small segments.

Subtyping partitions TcdB into distinct functional and antigenic groups

The value of subtyping classification is to facilitate a molecular understanding of the impact of
sequence variations on function and antigenicity. For instance, our sequence alignment divides the
GTD into two groups: one contains B4, B6, B7, and B8; and the rest form another group (Fig. 3b,
Fig. S7). Previous studies have reported two types of cell-rounding effects: TcdB1 and B2 are
known to induce rounded cells with many protrusions remaining attached to cell culture plate,
whereas TcdB from the strain 1470 and 8864 have been reported to cause rounded cells without
protrusions, which is similar to TcsL[50]. It has been proposed that this is a result of the altered
specificity of their GTD in targeting different small GTPases[32,50]. TcdB in strain 1470 is classified
as B4, and the strain 8864 expresses B7, thus our classification predicts that the group containing
B4/6/7/8 induces TcsL-like cell rounding phenotype. This is indeed the case for two recently
reported clinical strains HSJD-312 and HMX152: both express toxins classified as B6 under our
subtyping system (Table S2) and have been reported to induce TcsL-like cell rounding[51].

Another well-characterized functional motif in TcdB is its FZD-binding interface, with key
residues clearly defined by the co-crystal structure[37,38]. It has been reported that B2 lost the ability
to bind FZDs due to residue variations at FZD-binding interfaces[35,39,40]. To survey whether these
variations may also exist in other subtypes, we aligned the key residues across all TcdB sequences
and visualized them in color. As shown in Fig. 4a, FZD-binding motif is highly conserved across
B1/3/4/5/8/9, while B2/6/7/10 share the same set of residue changes. Thus, B6/7/10 are predicted
to lose FZD-binding capability similar to B2. B11 contains a subset of residue changes found in
B2 within this region and likely also has reduced binding to FZDs. This pattern is consistent with
the phylogenic alignment of the DRBD domain, in which B1/3/4/5/8/9 form group i and
B2/6/7/10/11 form the group ii (Fig. 3b, 3c). Interestingly, although most B2 variants possess FZD-
binding site substitutions, there are a few exceptions that contain a largely in-tact FZD binding

site. In particular, B2.12 assembled from strain 2007223 (ERS001491) contains only a single

amino acid substitution (F1597S) in this region. Examination of the alignment reveals that this is

likely due to a microrecombination event that has replaced most of the FZD binding site with a

B1-like segment (Fig. 4a, Fig. S6). A similar scenario occurred in a member of subtype B6, in

which a B1-like segment has partially replaced this region (Fig. S6).

Sequence variations between subtypes could also have a drastic impact on efficacy of therapeutic

antibodies and vaccines. Bezlotoxumab from Merck is the only monoclonal antibody against TcdB

that was approved by the FDA and is currently used to reduce the recurrence of CDI[52]. This

antibody was generated using fragments of TcdB1 as antigens and its epitope sites (located at the

N-terminal of CROPs) have been established through crystallography[53]. We thus aligned all key

residues within its epitope across all TcdB sequences, which revealed extensive residue changes

largely conserved in B2/6/7/9/10/11 (Fig. 4a). This is consistent with our alignment of the CROPs

domain that group B2/6/7/9/10/11 together (Fig. 3b, 3c). It has been shown that bezlotoxumab

exhibited as low as over ~700-fold reduction in neutralization efficacy against TcdB from several

RT027 strains, which likely express B2, compared with its efficacy against B1 from VPI10463[36].

It also showed a similarly low efficacy against a strain 8864, which expresses B7. These results

indicate that bezlotoxumab does not have good efficacy against CDI caused by strains that express

B2/6/7/9/10/11. Furthermore, there are also a few amino acid changes within the epitope region in

B3/B8, and it has been shown that bezlotoxumab has ~60-fold reduction in efficacy against the

TcdB from a RT078 strain[36], which likely express B3 (Table S1). These results clearly indicate

that subtype classification of toxins will be able to guide the use of bezlotoxumab in clinic.

In addition to bezlotoxumab, we also examined another monoclonal antibody PA41, which is

under development[36], and a single-domain antibody (also known as VHH or nanobody) E3[54]. The

epitopes for both have been well established through co-crystal structures[54,55]. Both recognize the

GTD domain, with E3 recognizing the N-terminus of TcdB (Fig. 4a). The epitope site for PA41 is

highly conserved across most subtypes except a single residue change (Y323H) in B4. This is

consistent with the previous finding that PA41 can potently neutralize TcdB from many different

strains except RT017 strains, which express B4[36]. The epitope site for E3 is conserved in most

subtypes except a single residue change (I58T or A) in B4/6/7/8/10/11/12, and the impact of this
single residue change remains to be examined experimentally.
370
We finally mapped evolutionary conservation across all available TcdB sequences onto the
recently reported crystal structure of TcdB[54] (Fig. 4b). Relative to the reference TcdB1 sequence,
amino acid variants are common across the full-length TcdB sequence and occur throughout each
domain (Fig. 4a) but some regions (e.g., N-terminus of the GTD, C-terminus of CROPS domain,
segments of the pore-forming region of the DRBD and C-terminus of the CROPS domain) were
highly conserved. Based on structure, we identified eight conserved surface patches containing
universally conserved residues which represent potential key therapeutic targets for developing
broad-spectrum diagnostics, antibodies, and vaccines (Fig. 4b).
379
380

Diff-base: a central hub for storing and analyzing TcdA and TcdB sequences
To address the needs of the research and clinical community in understanding toxin subtyping and
variations, we developed an online open database freely accessible at *diffbase.uwaterloo.ca*.
DiffBase stores all unique TcdA and TcdB sequences identified to date from the NCBI and SRA
and organizes sequences into our subtype classification scheme. Different subtypes and individual
sequences can be explored and visualized in reference trees, with additional information such as
source strains, and links to other resources (Fig. S8). In addition, users can query their own TcdA
or TcdB sequences against the database using a built-in BLAST interface, which will report the
top matching sequences in the database and provide toxin classifications and other related
information. To keep up with new sequences and information concerning TcdA and TcdB,
DiffBase facilitates community feedback and allows users to submit new information to be added
to the next iteration of the database.
393

Discussion

Here we created the largest database to date capturing available TcdA and TcdB sequence diversity. This up-to-date collection includes genes from sequenced *C. difficile* isolates in GenBank, as well as thousands of genomes that were assembled, annotated and analyzed from the NCBI short-read archive. We clustered TcdA and TcdB variants into phylogenetic subtypes, which provided a robust classification that is both congruent with *C. difficile* genome phylogeny as well as variation in functional and therapeutically relevant amino acids including TcdB regions targeted by existing monoclonal antibodies. Our analysis revealed that TcdB undergoes extensive homologous recombination, and its potential evolutionary history is proposed based on recombination among various subtypes. Finally, our analysis revealed mapped eight conserved patches across the TcdB structure, which will facilitate future studies that aim to develop "universal" *C. difficile* therapeutics that broadly target all TcdB subtypes.

In general, there is some agreement between previously defined toxinotypes and our toxin subtypes, but subtyping provides additional information as it is able to describe TcdA and TcdB separately. For example, toxinotype 0 associates largely with the A1/B1 subtype, toxinotype III associates largely with A2/B2, toxinotype IV with A3/B5, toxinotype VIII associates with -/B4, toxinotype IX associates largely with A2/B6, toxinotype X associates with -/B7, and so on (Table S1). However, subtype A3/B3 associated with toxinotypes V, VI, VII, XVI, XXVIII, all of which are found in clade 5 strains. Moving forward, with improved abilities to perform genome sequencing of clinical isolates, it will be increasingly possible to classify strains based on their genome-wide phylogenetic relationships as well as their toxin subtypes.

In comparison to TcdA, our analysis identified a much greater degree of sequence variation within TcdB and a larger number of subtypes. Given that we see evidence for extreme recombination in TcdB but not TcdA, it is possible that there is a greater selective pressure for positive selection and diversification of TcdB. We speculate that intragenic recombination of TcdB may drive antigenic diversification, whereas in TcdA this process may be driven by truncation and variation of its C-terminal CROPS region. The CROPs domain showed similarity with carbohydrate-binding proteins and may contribute to toxin attachment to cells by binding to carbohydrate moieties (27, 42, 44). The CROPs domain may also act as a chaperone that protects other domains (45). Possibly

due to its repetitive nature, the CROPs domain is often the region that induces strong immune

responses. It remains to be determined whether frequent recombinations/changes in TcdA-CROPs

may alter its function and/or antigenicity.

These findings further suggest that TcdB may play a central role in *C. difficile* pathogenesis, which

is consistent with previous findings that TcdA-/TcdB+ mutant *C. difficile* strains are fully virulent,

whereas TcdA+/TcdB- strains are attenuated in multiple mouse models[26,56]. It has also been

suggested that TcdB is the primary factor for inducing the host immune and inflammatory

responses in mouse models[26]. The key role of TcdB in CDI is further confirmed by the findings

that an antibody that neutralizes TcdB (bezlotoxumab), but not another one that neutralizes TcdA

(actoxumab, Merck), conferred protection against CDI in gnotobiotic piglets [57] and reduced CDI

recurrence in humans[52,58] and it is also consistent with the fact that many clinical isolates only

express TcdB[59]. An exception to a dominant role for TcdB is the very rare TcdA+ TcdB- strain. It

is noteworthy that one such strain identified in GenBank contains the single most divergent TcdA

sequence (subtype A7)[45], which may have diverged to acquire a pathogenic functionality without

requiring TcdB.

For such recombination events to have occurred in TcdB, phylogenetically distinct *C. difficile*

strains containing different toxin subtypes must have coexisted within the same host individuals,

exchanged genetic material and recombined to produce new recombinant forms. Co-infection with

different *C. difficile* ribotypes has been recently reported in a clinical case study[60]. Theoretically,

co-infection does not need to occur frequently to promote recombination. A single individual

containing two or more *C. difficile* strains, may be sufficient to promote recombination, generating

hybrid toxins with different regions derived from different sequences or subtypes. The new

recombinant strain can then increase in frequency through transmission to other individuals. Our

analysis suggests that this process has not only occurred frequently in the past as a mechanism by

which different subtypes originated, but that it may be a frequent and ongoing process in new

clinical isolates (e.g., B1.59 from SRS1378602). Consideration of intragenic recombination and

how it may shape TcdB function and toxicity will be important in efforts to understand the

emergence of new *C. difficile* hypervirulent strains and develop targeted therapeutic interventions.

Recombination offers considerable adaptive benefits to proteins by facilitating rapid mutation of

a sequence by exchange of entire segments as opposed to the relatively slower process of single

point mutations. In this way, proteins can diversify by shuffling a few basic building blocks such

as protein domains. In pathogens, recombination plays a major role in pathoadaptive evolution by

facilitating rapid "switching" of virulence factors and antigenic proteins[61,62]. Antigenic

recombination can promote the sudden avoidance of immune recognition (antigenic escape),

which has been demonstrated for the *C. difficile* S-layer gene[63]. In the case of TcdB, intragenic

recombination may generate new hybrid toxins composed of different domains types and

functions. In theory, recombination could also generate resistance to therapeutics by replacing

entire binding interfaces with compatible regions from other toxins that possess drug-resistant

mutations. Recombination-mediated domain shuffling not only describes TcdB sequence patterns

and phylogenetic relationships, and also provides an explanation for important functional

differences between TcdB variants. For example, the exchange of a B6-like GTD between

subtypes B4, B8, B6, and B7, correlates with the TcsL-like clumping and rounding phenotype.

Also intriguing are the many microrecombination events that have occurred in the DRDB region

which overlap with FZD-binding site. For example, likely due to partial homologous

recombination with a B1-like toxin, one B2 variant (B2.12, strain 2007223 from ERS001491)

contains an in-tact FZD-binding interface with only a single amino acid substitution (F1597S).

This suggests that intragenic recombination in TcdB may promote rapid evolutionary switching

between receptor-binding activities or affinities.

Given the extent of TcdB diversification and its primary role in virulence, it is critically important

to identify conserved regions that can be targeted for therapeutic and diagnostic applications.

Sequence conservation mapped to protein structure also revealed at least 8 distinct surface patches

containing a high density of universally conserved residues across all TcdB subtypes, which

represent promising regions for the development of inhibitors. Importantly, the binding site for the

antibody therapeutic bezlotoxumab, which is commonly used to treat *C. difficile* infections, was

not among these and instead displayed considerable variation across TcdB subtypes with B2, B10,

B11, B9, B6, and B7 in particular displaying 7-8 likely destabilizing substitutions. Although the

common B1 subtype of TcdB is largely conserved across this region, based on analysis, it is

possible that intragenic recombination with other strains (e.g., a B2-containing strain) could

generate spontaneous resistance to bezlotoxumab by replacing this region with a B2-like segment.

Future efforts to target highly conserved clusters of surface-exposed residues on the TcdB structure

may yield promising candidates for therapeutic or vaccine development.

Finally, based on sequence-based classification of *tcdA* and *tcdB* genes, we propose a revised

scheme for naming these genes in future studies. In this scheme, a newly identified TcdA or TcdB

sequence may be aligned to our reference database and named based on the top hit according to

sequence identity, provided that the sequence exceeds thresholds used for our clustering (99.4%

for TcdA and 97.0% for TcdB). In order to enable automated subtyping of new *tcdA* and *tcdB*

genes and facilitate community collaboration and data sharing, we have developed a freely

available, online database (DiffBase) for use by *C. difficile* clinical and research community. In

the future, clinicians will be able to query toxin sequences from clinical isolates and immediately

determine the toxin subtype, which will help them decide on therapeutic strategies. For instance,

among the 351 clinical cases in the BWH dataset, there are 34 cases expressing B2/B6/B7/B9, for

which treatment of bezlotoxumab would not be effective. Therefore, toxin subtyping will guide

proper choices of clinical treatment in consideration of toxin variations and allow researchers to

monitor the ongoing evolution and diversification of *C. difficile*.

Methods

Dataset construction

Assembly of 6492 C. difficile genomes from the NCBI short read archive

A set of *Clostridiodes difficile* sequencing runs was retrieved from the NCBI short read archive (SRA) by text query for "*Clostridioides difficile*" on June 20th, 2019. Metagenomic samples were omitted, leaving only genomic samples to reduce the chance of contamination from other bacterial species. Sequencing runs were downloaded using the fasterq-dump module of the SRA toolkit. To account for multiple library preparation methods and adapters, the fastp tool[64] was used to perform adapter trimming and quality control of the sequencing reads. For each quality-controlled set of reads, SPAdes version 3.12[65] was used for genomes assembly, with *C. difficile* str. 630 as a conservative reference and the --untrusted-contigs and --careful options. Each assembly was automatically annotated using the Prokka pipeline[66] with a minimum contig length of 200. In order to verify the identity of the assembled genomes as strains of *C. difficile,* the predicted genes from Prokka were taxonomically annotated using Centrifuge[67] against their pre-compiled index of bacterial, archaeal, viral, and human genomes. Only samples that were statistically verified as belong to strains of *C. difficile* were kept.

To identify the *tcdA* and *tcdB* genes from all strains, the phmmer tool was used to search for matches to TcdA (uniprot accession # Q189K5_CLOD6) and TcdB (uniport accession # Q189K3_CLOD6) as queries. In order to distinguish true sequence variants from poorly assembled, low-quality, or chimeric variants, only hits that clearly represented well-assembled toxin sequences (that is, yielding a protein equal to or greater than 1,800 amino acids) were retained. Sequences with apparent N- or C-terminal truncations representing less than 1% of the total assembled data set were also removed. In total, the final re-assembled set of redundant TcdA and TcdB sequences consisted of 3,542 and 5,994 sequences, respectively. Redundancy in each of these sets was removed by clustering with CD-HIT version 4.6[68] at 100% identity. Non-redundant sets were aligned using the L-INS-i algorithm of the MAFFT package[69].

TcdA and TcdB sequences from the NCBI GenBank database and manually curated set

GenBank homologs of TcdA and TcdB were also identified via a BLAST search of the NCBI non-
redundant database on Feb 8, 2020. TcdB and TcdA sequences from *C. difficile* strain 630 were
used as queries. Homologs were filtered to those with *E*-value < 1e-10, 70% identity and query
alignment coverage, which removed partial sequences. In addition, we manually curated 63
reference *C. difficile* strains collected from previous studies[12,41]. For these 63 genomes, we
manually identified corresponding strains within the NCBI or SRA database and identified *tcdA*
and *tcdB* genes based on pre-computed genome annotations or through similarity searches.
Fourteen genomes could not be associated with *tcdA* and *tcdB* genes in the NCBI; for these cases,
raw genomic reads were retrieved from European Nucleotide Archive (ENA) and were assembled
using SPADES as described earlier.

Construction of TcdA and TcdB alignments
A combined dataset of TcdA and TcdB homologs was created by pooling SRA-derived, NCBI-nr
derived, and the manually curated set of sequences. The combined set of sequences were aligned
using MUSCLE[70] with default parameters as implemented in Seaview[71]. Due to significant length
variation at the C-terminus of TcdA alignment, only the CROP-less core region (1-1874) of the
alignment was kept for subsequent analysis; while the entire TcdB alignment (1-2366) was used.
Redundant sequences (100% identity) were removed as well as sequences annotated as partial that
contained truncations in the alignment.

Sequence clustering and analysis
TcdA and TcdB alignments were then processed separately using an analysis pipeline
implemented within R. For each case, the multiple sequence alignment was converted to a distance
matrix using the dist.alignment() function from the seqinr package[72]. Average linkage hierarchical
clustering was performed using the hclust() function. Pairwise sequence similarities were mapped
onto the clustering tree and visualized using the ComplexHeatmap package[73] and clustering
threshold were chosen to generate subtypes with strong internal (within-cluster) and lower external
(between-cluster) similarities based on visual analysis and quantitative analysis of percentage
identity distributions.

For analysis of amino acid variation, we converted the alignments into data matrices using the
alignment2matrix() function from the BALCONY R package[74]. We then identified all variant
residues across all alignment positions relative to the sequences of TcdA and TcdB from strain
630 as a reference. Residues implicated in frizzled binding[37], bezlotoxumab binding[53], PA41
binding[55], and E3 binding[54] were then analyzed in terms of their variation across subtypes. E3
binding residues were identified by analysis of PDB structure 6OQ5[54], by selecting atoms in chain
A (TcdB) within a 4 Å distance of chain E (E3) using PyMol's distance algebra functions. The
ComplexHeatmap R package was used for data visualization.

Analysis of repeats in TcdA was done using InterproScan as part of the InterPro 80.0 database[75].
The number of detected matches to ProSite's cell wall-binding repeat profile (PS51170) was
counted in the A1.1 reference sequence (UniProt P16154, 2710 aa), the longest (3070 aa) and
shortest (1889 aa) variants of TcdA in our database.

Structural analysis
To map sequence conservation on to the structure of TcdB, we used the ConSurf server[76] with the
TcdB alignment as input and the recently determined crystal structure of full-length TcdB (PDB
ID 6OQ5)[54] as the template. Default parameters (neighbor-joining with ML distance and Bayesian
calculation of conservation scores) were used. Structural visualization was done using PyMol
version 2.3.4, using the recommended script (https://consurf.tau.ac.il/pyMOL/consurf_new.py)
with insufficient data hidden from the image.

Construction and toxin subtyping of C. difficile genome phylogeny
We retrieved 2,118 assemblies for 1,934 representative *C. difficile* genomes
(https://www.ncbi.nlm.nih.gov/genome/tree/535) from the NCBI. The snippy pipeline
(https://github.com/tseemann/snippy) was used to map all genomes to the reference (strain 630,
GCA_000003215). For phylogenetic reconstruction, we analyzed 14,194 SNPs across 88
conserved marker genes (those present in *C. difficile*) derived from the PhyEco Firmicutes
dataset[77]. A phylogeny was reconstructed using RAXML with the GTRGAMMA model [78]. All
TcdA and TcdB homologs from the NCBI were then subtyped by BLAST against our database of
labeled toxin subtype sequences, using only the conserved portion (region 1-1874) of the TcdA

alignment, and the full 1-2366 regions from the TcdB alignments. An assignment script written in
Perl was used to parse BLAST output files and assign subtypes. The subtype "X" associated with
the best matching reference sequence (highest sequence identity) was assigned if the alignment
coverage exceeded 90% and labeled as complete; otherwise, it was labeled as a partial sequence.
601
602   *SplitsTree analysis of recombination*
603   TcdA and TcdB alignments were analyzed by SplitsTree version 4.0[49]. A NeighborNet tree
604   visualization was produced using protein maximum-likelihood distances according to the WAG
605   model of evolution. The Phi test for recombination was performed as implemented in SplitsTree
606   which selected a window size of 100 for TcdA with k = 3 and a window size of TcdB with k = 21.
607
608   *Haplotype visualization*
609   For visualization of recombinant blocks and haplotype structure within TcdA and TcdB protein
610   alignments, we developed a modified algorithm based on a previous method from Wang et al.[48]
611   for comparative genomic visualization. An implementation of this method in the R programming
612   language is available at https://github.com/doxeylab/haploColor. The algorithms works as follows:
613
614   (1) Assign first sequence as reference.
615   (2) Assign all residues of reference a new color C.
616   (3) Assign positions in other sequences that match the reference, the same color C.
617   (4) Identify sequence most dissimilar to the current reference across unassigned positions, and
618   assign it as the new reference.
619   (5) Repeat steps 2-3 for a defined number of iterations or until all sequences are completely
620   colored.
621
622   The algorithm was applied directly to the TcdA and TcdB alignments and run for both 4 and 16
623   iterations (TcdB) and 16 iterations (TcdA).
624
Development of the DiffBase web-server
The DiffBase web server was developed as an R shiny() application. Contained within DiffBase
is an implementation of BLAST+. Individual sequences can be submitted to the server, where the

blastp program is run to find matches from within the entirety of the server sequence repositories.
An *E*-value cutoff of 1e-10 is used to filter hits, and the results are sorted by percent identity
between query and target sequences. Toxin groups can also be viewed in a phylogenetic tree
visualized using ggtree R package[79]. Metadata about group members was obtained from the NCBI
Identical Protein Group (IPG) database. The source code is freely available at
https://github.com/doxeylab/diffBase.

634
635

636 **Data Availability**
637 Our open source online database is available at: https://diffbase.uwaterloo.ca and
638 https://github.com/doxeylab/diffbase.
639 All source code for analyses is available at: https://github.com/doxeylab/diffBaseAnalyses

640
641

Acknowledgments

A.C.D. acknowledges funding from the Natural Sciences and Engineering Research Council of
Canada (NSERC Discovery Grant, RGPIN-2019-04266; Discovery Accelerator Supplement,
RGPAS-2019-00004), and from the Government of Ontario (Early Researcher Award). A.C.D.
holds a University Research Chair at the University of Waterloo.

M.D. acknowledges support from National Institute of Health (NIH) (R01AI132387 and
R01AI139087 to M.D.), the NIH-funded Harvard Digestive Disease Center (P30DK034854), and
Boston Children's Hospital Intellectual and Developmental Disabilities Research Center
(P30HD18655). M.D. holds the Investigator in the Pathogenesis of Infectious Disease award from
the Burroughs Wellcome Fund.

J. W. acknowledges support from the Intramural Research Program of the National Library of
Medicine, NIH. L.B. acknowledges support from NIH (P30 DK034854), Hatch Family
Foundation, and Brigham and Women's Hospital Precision Medicine Institute.

657
658

659   **References**

661   1.   Knight, D. R., Elliott, B., Chang, B. J., Perkins, T. T. & Riley, T. V. Diversity and
662   evolution in the genome of *Clostridium difficile*. *Clin. Microbiol. Rev.* **28**, 721–741
663   (2015).
664   2.   Guh, A. Y. *et al.* Trends in U.S. burden of *Clostridioides difficile* infection and outcomes.
665   *N. Engl. J. Med.* **382**, 1320–1330 (2020).
666   3.   Heinlen, L. & Ballard, J. D. *Clostridium difficile* infection. *Am. J. Med. Sci.* **340**, 247–252
667   (2010).
668   4.   Rupnik, M., Wilcox, M. H. & Gerding, D. N. *Clostridium difficile* infection: New
669   developments in epidemiology and pathogenesis. *Nat. Rev. Microbiol.* **7**, 526–536 (2009).
670   5.   Martin, J. S. H., Monaghan, T. M. & Wilcox, M. H. *Clostridium difficile* infection:
671   Epidemiology, diagnosis and understanding transmission. *Nat. Rev. Gastroenterol.*
672   *Hepatol.* **13**, 206–216 (2016).
673   6.   Smits, W. K., Lyras, D., Lacy, D. B., Wilcox, M. H. & Kuijper, E. J. *Clostridium difficile*
674   infection. *Nat. Rev. Dis. Prim.* **2**, 16020 (2016).
675   7.   Abt, M. C., McKenney, P. T. & Pamer, E. G. *Clostridium difficile* colitis: Pathogenesis
676   and host defence. *Nat. Rev. Microbiol.* **14**, 609–620 (2016).
677   8.   Lessa, F. C. *et al.* Burden of *Clostridium difficile* infection in the United States. *N. Engl. J.*
678   *Med.* **372**, 825–834 (2015).
679   9.   Knetsch, C. W. *et al.* Current application and future perspectives of molecular typing
680   methods to study clostridium difficile infections. *Eurosurveillance* **18**, 20381 (2013).
681   10.   Griffiths, D. *et al.* Multilocus sequence typing of *Clostridium difficile*. *J. Clin. Microbiol.*
682   **48**, 770–8 (2010).
683   11.   Eyre, D. W. *et al.* Diverse sources of *C. difficile* infection identified on whole-genome
684   sequencing. *N. Engl. J. Med.* **369**, 1195–1205 (2013).
685   12.   Bletz, S., Janezic, S., Harmsen, D., Rupnik, M. & Mellmann, A. Defining and Evaluating
686   a Core Genome Multilocus Sequence Typing Scheme for Genome-Wide Typing of
687   *Clostridium difficile*. *J. Clin. Microbiol.* **56**, (2018).
688   13.   Janezic, S. & Rupnik, M. Development and Implementation of Whole Genome
689   Sequencing-Based Typing Schemes for *Clostridioides difficile*. *Front. Public Heal.* **7**, 309

(2019).

14.   Dingle, K. E. *et al.* Evolutionary history of the *Clostridium difficile* pathogenicity locus.

Genome Biol. Evol. **6**, 36–52 (2014).

15.   Janezic, S., Potocnik, M., Zidaric, V. & Rupnik, M. Highly divergent *Clostridium difficile*

strains isolated from the environment. *PLoS One* **11**, e0167101 (2016).

16.   He, M. *et al.* Evolutionary dynamics of *Clostridium difficile* over short and long time

scales. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 7527–7532 (2010).

17.   Collins, J. *et al.* Dietary trehalose enhances virulence of epidemic *Clostridium difficile*.

Nature **553**, 291–294 (2018).

18.   He, M. *et al.* Emergence and global spread of epidemic healthcare-associated *Clostridium*

difficile. *Nat. Genet.* **45**, 109–113 (2013).

19.   McDonald, L. C. *et al.* An epidemic, toxin gene-variant strain of *Clostridium difficile*. *N.*

Engl. J. Med. **353**, 2433–2441 (2005).

20.   Loo, V. G. *et al.* A predominantly clonal multi-institutional outbreak of *Clostridium*

difficile - Associated diarrhea with high morbidity and mortality. *N. Engl. J. Med.* **353**,

2442–2449 (2005).

21.   Goorhuis, A. *et al.* Emergence of *Clostridium difficile* infection due to a new

hypervirulent strain, polymerase chain reaction ribotype 078. *Clin. Infect. Dis.* **47**, 1162–

70 (2008).

22.   Jhung, M. A. *et al.* Toxinotype V *Clostridium difficile* in humans and food animals.

Emerg. Infect. Dis. **14**, 1039–1045 (2008).

23.   Collins, J., Danhof, H. & Britton, R. A. The role of trehalose in the global spread of

epidemic *Clostridium difficile*. *Gut Microbes* **10**, 204–209 (2019).

24.   Aktories, K., Schwan, C. & Jank, T. *Clostridium difficile* Toxin Biology. *Annu. Rev.*

Microbiol. **71**, 281–307 (2017).

25.   Voth, D. E. & Ballard, J. D. *Clostridium difficile* toxins: Mechanism of action and role in

disease. *Clin. Microbiol. Rev.* **18**, 247–263 (2005).

26.   Carter, G. P. *et al.* Defining the Roles of TcdA and TcdB in Localized Gastrointestinal

Disease, Systemic Organ Damage, and the Host Response during *Clostridium difficile*

Infections. *MBio* **6**, e00551 (2015).

27.   Pruitt, R. N. & Lacy, D. B. Toward a structural understanding of *Clostridium difficile*

toxins A and B. *Front. Cell. Infect. Microbiol.* **2**, 28 (2012).

28.   Gerding, D. N., Sambol, S. P. & Johnson, S. Non-toxigenic clostridioides (formerly

clostridium) difficile for prevention of *C. difficile* infection: From bench to bedside back

to bench and back to bedside. *Front. Microbiol.* **9**, 1700 (2018).

29.   Schirmer, J. & Aktories, K. Large clostridial cytotoxins: Cellular biology of Rho/Ras-

glucosylating toxins. *Biochim. Biophys. Acta - Gen. Subj.* **1673**, 66–74 (2004).

30.   Jank, T. & Aktories, K. Structure and mode of action of clostridial glucosylating toxins:

the ABCD model. *Trends Microbiol.* **16**, 222–229 (2008).

31.   Orrell, K. E., Mansfield, M. J., Doxey, A. C. & Melnyk, R. A. The *C. difficile* toxin B

membrane translocation machinery is an evolutionarily conserved protein delivery

apparatus. *Nat. Commun.* **11**, 432 (2020).

32.   Genth, H. *et al.* Haemorrhagic toxin and lethal toxin from *Clostridium sordellii* strain

vpi9048: Molecular characterization and comparative analysis of substrate specificity of

the large clostridial glucosylating toxins. *Cell. Microbiol.* **16**, 1706–1721 (2014).

33.   Davies, A. H., Roberts, A. K., Shone, C. C. & Acharya, K. R. Super toxins from a super

bug: structure and function of *Clostridium difficile* toxins. *Biochem. J.* **436**, 517–526

(2011).

34.   Lanis, J. M., Heinlen, L. D., James, J. A. & Ballard, J. D. *Clostridium difficile*

027/BI/NAP1 encodes a hypertoxic and antigenically variable form of TcdB. *PLoS*

Pathog. **9**, e1003523 (2013).

35.   Chung, S.-Y. *et al.* The Conserved Cys-2232 in *Clostridioides difficile* Toxin B Modulates

Receptor Binding. *Front. Microbiol.* **9**, 2314 (2018).

36.   Marozsan, A. J. *et al.* Protection against *Clostridium difficile* infection with broadly

neutralizing antitoxin monoclonal antibodies. *J. Infect. Dis.* **206**, 706–713 (2012).

37.   Chen, P. *et al.* Structural basis for recognition of frizzled proteins by *Clostridium difficile*

toxin B. *Science* **360**, 664–669 (2018).

38.   Tao, L. *et al.* Frizzled proteins are colonic epithelial receptors for *C. difficile* toxin B.

Nature **538**, 350–355 (2016).

39.   Peng, Z. *et al.* Designed Ankyrin Repeat Protein (DARPin) Neutralizers of TcdB from

Clostridium difficile Ribotype 027. *mSphere* **4**, e00596-19 (2019).

40.   López-Ureña, D. *et al.* Toxin B Variants from *Clostridium difficile* Strains VPI 10463 and

752    NAP1/027 Share Similar Substrate Profile and Cellular Intoxication Kinetics but Use

753    Different Host Cell Entry Factors. *Toxins (Basel).* **11**, (2019).

754  41. M, R. & S, J. An Update on *Clostridium Difficile* Toxinotyping. *J. Clin. Microbiol.* **54**,

755    (2016).

756  42. Rupnik, M. Clostridium difficile toxinotyping. *Methods Mol. Biol.* **646**, 67–76 (2010).

757  43. Von Eichel-Streiber, C., Sauerborn, M. & Kuramitsu, H. K. Evidence for a modular

758    structure of the homologous repetitive C-terminal carbohydrate-binding sites of

759    *Clostridium difficile* toxins and *Streptococcus mutans* glucosyltransferases. *J. Bacteriol.*

760    **174**, 6707–6710 (1992).

761  44. von Eichel-Streiber, C., Laufenberg-Feldmann, R., Sartingen, S., Schulze, J. & Sauerborn,

762    M. Comparative sequence analysis of the *Clostridium difficile* toxins A and B. *MGG Mol.*

763    *Gen. Genet.* **233**, 260–268 (1992).

764  45. M, M. *et al.* Clostridium Difficile: New Insights Into the Evolution of the Pathogenicity

765    Locus. *Sci. Rep.* **5**, 15023 (2015).

766  46. Stabler, R. A. *et al.* Comparative phylogenomics of *Clostridium difficile* reveals clade

767    specificity and microevolution of hypervirulent strains. *J. Bacteriol.* **188**, 7297–305

768    (2006).

769  47. Worley, J. N. *et al.* Genomic determination of relative risks for *Clostridioides difficile*

770    infection from asymptomatic carriage in ICU patients. *Clin. Infect. Dis.* In Press. (2020).

771    doi:doi:10.1093/cid/ciaa894

772  48. Wang, J. R., de Villena, F. P.-M. & McMillan, L. Comparative analysis and visualization

773    of multiple collinear genomes. *BMC Bioinformatics* **13 Suppl 3**, S13 (2012).

774  49. Huson, D. H. SplitsTree: Analyzing and visualizing evolutionary data. *Bioinformatics*

775    (1998). doi:10.1093/bioinformatics/14.1.68

776  50. Chaves-Olarte, E. *et al.* R-Ras glucosylation and transient RhoA activation determine the

777    cytopathic effect produced by toxin B variants from toxin A-negative strains of

778    *Clostridium difficile*. *J. Biol. Chem.* **278**, 7956–7963 (2003).

779  51. Ramírez-Vargas, G. *et al.* Novel Clade C-I *Clostridium difficile* strains escape diagnostic

780    tests, differ in pathogenicity potential and carry toxins on extrachromosomal elements.

781    *Sci. Rep.* **8**, 13951 (2018).

782  52. Wilcox, M. H. *et al.* Bezlotoxumab for Prevention of Recurrent *Clostridium difficile*

Infection. *N. Engl. J. Med.* **376**, 305–317 (2017).

53.   Orth, P. *et al.* Mechanism of action and epitopes of *Clostridium difficile* toxin B-
neutralizing antibody bezlotoxumab revealed by X-ray crystallography. *J. Biol. Chem.*
289, 18008–21 (2014).

54.   Chen, P. *et al.* Structure of the full-length *Clostridium difficile* toxin B. *Nat. Struct. Mol.*
Biol. **26**, 712–719 (2019).

55.   Kroh, H. K. *et al.* A neutralizing antibody that blocks delivery of the enzymatic cargo of
Clostridium difficile toxin TcdB into host cells. *J. Biol. Chem.* **293**, 941–952 (2018).

56.   Lyras, D. *et al.* Toxin B is essential for virulence of *Clostridium difficile*. *Nature* **458**,
1176–1179 (2009).

57.   Steele, J., Mukherjee, J., Parry, N. & Tzipori, S. Antibody against TcdB, but not TcdA,
prevents development of gastrointestinal and systemic *Clostridium difficile* disease. *J.*
Infect. Dis. **207**, 323–30 (2013).

58.   Gupta, S. B. *et al.* Antibodies to Toxin B Are Protective Against *Clostridium difficile*
Infection Recurrence. *Clin. Infect. Dis.* **63**, 730–734 (2016).

59.   Janezic, S., Marín, M., Martín, A. & Rupnika, M. A new type of toxin a-negative, toxin B-
positive *Clostridium difficile* strain lacking a complete tcdA gene. *J. Clin. Microbiol.* **53**,
692–695 (2015).

60.   Wang, L. *et al.* Coinfection with 2 *Clostridium difficile* ribotypes in China. *Medicine*
(Baltimore). **97**, e9946 (2018).

61.   Awadalla, P. The evolutionary genomics of pathogen recombination. *Nat. Rev. Genet.* **4**,
50–60 (2003).

62.   Wilson, D. J. *et al.* Rapid evolution and the importance of recombination to the
gastroenteric pathogen *Campylobacter jejuni*. *Mol. Biol. Evol.* **26**, 385–397 (2009).

63.   Dingle, K. E. *et al.* Recombinational switching of the *Clostridium difficile* S-layer and a
novel glycosylation gene cluster revealed by large-scale whole-genome sequencing. *J.*
Infect. Dis. **207**, 675–86 (2013).

64.   Chen, S., Zhou, Y., Chen, Y. & Gu, J. fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics **34**, i884–i890 (2018).

65.   Bankevich, A. *et al.* SPAdes: a new genome assembly algorithm and its applications to
single-cell sequencing. *J. Comput. Biol.* **19**, 455–77 (2012).

66. Seemann, T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* **30**, 2068–2069
(2014).
67. Kim, D., Song, L., Breitwieser, F. P. & Salzberg, S. L. Centrifuge: rapid and sensitive
classification of metagenomic sequences. *Genome Res.* **26**, 1721–1729 (2016).
68. Li, W. & Godzik, A. Cd-hit: a fast program for clustering and comparing large sets of
protein or nucleotide sequences. *Bioinformatics* **22**, 1658–1659 (2006).
69. Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. *Mol. Biol. Evol.* **30**, 772–80 (2013).
70. Edgar, R. C. MUSCLE: multiple sequence alignment with high accuracy and high
throughput. *Nucleic Acids Res.* **32**, 1792–7 (2004).
71. Gouy, M., Guindon, S. & Gascuel, O. SeaView version 4: A multiplatform graphical user
interface for sequence alignment and phylogenetic tree building. *Mol. Biol. Evol.* **27**, 221–
4 (2010).
72. Charif, D. & Lobry, J. R. SeqinR 1.0-2: A Contributed Package to the R Project for
Statistical Computing Devoted to Biological Sequences Retrieval and Analysis. in 207–
232 (Springer, Berlin, Heidelberg, 2007). doi:10.1007/978-3-540-35306-5_10
73. Gu, Z., Eils, R. & Schlesner, M. Complex heatmaps reveal patterns and correlations in
multidimensional genomic data. *Bioinformatics* **32**, 2847–9 (2016).
74. Płuciennik, A. *et al.* BALCONY: an R package for MSA and functional compartments of
protein variability analysis. *BMC Bioinformatics* **19**, 300 (2018).
75. Hunter, S. *et al.* InterPro: The integrative protein signature database. *Nucleic Acids Res.*
37, (2009).
76. Ashkenazy, H. *et al.* ConSurf 2016: an improved methodology to estimate and visualize
evolutionary conservation in macromolecules. *Nucleic Acids Res.* **44**, W344-50 (2016).
77. Wu, D., Jospin, G. & Eisen, J. A. Systematic Identification of Gene Families for Use as
'Markers' for Phylogenetic and Phylogeny-Driven Ecological Studies of Bacteria and
Archaea and Their Major Subgroups. *PLoS One* (2013).
doi:10.1371/journal.pone.0077033
78. Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. *Bioinformatics* **30**, 1312–1313 (2014).
79. Yu, G., Smith, D. K., Zhu, H., Guan, Y. & Lam, T. T. ggtree : an r package for

visualization and annotation of phylogenetic trees with their covariates and other

associated data. *Methods Ecol. Evol.* **8**, 28–36 (2017).

Figures and Tables

Table 1. Assembled *C. difficile* genomes from the NCBI SRA and associated statistics.

Property	Statistic (mean +/- SD)
Number of samples	6,492
Assembly length	4.2759 +/- 0.019 Mb
Number of contigs	403.6 +/- 544
GC content	28.33 +/- 4.14 % GC
Mean contig length	62.68 +/- 46.97 Kb
Contig N50	905,900.13 +/- 865,085
Contig N90	298,748 +/- 52,843.68

Figure 1. Clustering of TcdA and TcdB sequences derived from NCBI GenBank and SRA into subtypes. (**a**) Hierarchical clustering of TcdA sequences, split into 8 groups. (**b**) Neighbor-joining phylogenetic tree of representative sequences of each TcdA subtype. (**c**) Percentage identities between representative sequences. (**d**) Hierarchical clustering of TcdB sequences, split into 14 groups. (**e**) Neighbor-joining phylogenetic tree of representative sequences of each TcdB subtype. (**f**) Percentage identities between representative sequences. Hierarchical clustering was performed using the hclust() function in R, and cluster definitions were selected based on strong within-cluster sequence similarities and weak between-cluster similarities, as demonstrated visually and quantitatively. The reference strains (VPI 10463 and strain 630) are associated with TcdA group A1 and TcdB group B1. The hypervirulent ribotype 027 strains such as R12087 and R20291 are associated with TcdA group A2 and TcdB group B2. Also included are the homologs of TcdA and TcdB (TcdH and TcdL, respectively) from *P. sordellii*, which expectedly exhibit the highest divergence from other groups. The datasets include TcdA and TcdB sequences from the NCBI GenBank as well as an additional 125 sequences assembled from the SRA.

Figure 2. Toxin subtypes across the *C. difficile* phylogeny and occurrence of subtypes in a clinical CDI cohort. (**a**) TcdA (inner ring) and TcdB (outer ring) subtypes mapped onto a tree of 1934 *C. difficile* genomes. The genome tree is derived from the NCBI and is based on clustering of all-by-all genome BLAST scores. Lineages corresponding to previously identified *C. difficile* PaLoc clades (1 – 5) are labeled numerically. PaLoC clade 1 was subdivided into four sublineages labeled 1a-1d. Selected, clinically relevant strains are shown on the tree, with hypervirulent/epidemic outbreak strains indicated by stars. (**b**) Frequency of toxin subtypes detected in 1,934 representative, complete *C. difficile* genomes from NCBI/GenBank. A total of 1640 (84.8%) *C. difficile* strains contained TcdA and/or TcdB, while 294 (15.2%) were toxin deficient. (**c**) Frequency of toxin subtypes detected in a CDI clinical cohort from Brigham and Women's Hospital (BWH). The total dataset contained 351 *C. difficile* genomes derived from infected patients. Of these, 289 (82.3%) contained toxin genes, and 62 (17.7%) were toxin deficient.

Figure 3. **Evolutionary diversification of TcdB by intragenic recombination and domain shuffling.** (**a**) Visualization of amino acid variation patterns in TcdB using a newly developed haplotype coloring algorithm (HaploColor). The visualization shows patterns of amino acid variation across the TcdB alignment. In this algorithm, the first sequence (B1.1) is assigned a distinct color, and all other sequences are colored the same color where they match this first sequence. Then, the process is repeated using a second sequence (B6.1) as the new reference, and so on. This reveals multiple colored segments indicative of common ancestry (identity by descent). Mosaic patterns are indicative of intragenic recombination. (**b**) Phylogenetic trees of TcdB based on individual domains. Each domain tree can be subdivided into two types (labeled 1 and 2), which allows each subtype to be described based on its domain composition (**c**). This reveals that TcdB subtypes are composed of domains with variable evolutionary histories, indicative of domain shuffling and intragenic recombination. (**d**) Evolutionary model depicting relationships between subtypes and putative recombination events. Here, TcdB split early into two main groups (i and ii). Subtype B2 likely originated by a recombination event fusing an ancestral type i and type ii toxin. B9 likely originated from recombination between B1 and B2, B4 from recombination between B1 and a type ii toxin, and B8 from recombination between B3 and a type ii toxin.

Figure 4. Conservation and functional variation across TcdB subtypes. (**a**) Frequency of amino acid variants across all positions of TcdB. The height of the bar indicates the number of unique TcdB sequences that contain a substitution relative to the classical TcdB1 (B1.1) sequence from strain 630 and VPI10463. Below this is a plot of amino acid variation for key functional regions including the binding sites for the frizzled receptor (FZD) and the antibodies (E3, PA41, and bezlotoxumab). The alignment is colored gray for residues that match the common amino acid found in B1.1, and variants are colored blue (darkest blue = most common variant). E3 and PA41 binding sites

are highly conserved, whereas FZD and bezlotoxumab binding sites are highly variable. FZD and bezlotoxumab variants also co-occur with each other. (**b**) Evolutionary conservation mapped to the protein structure of full length TcdB based on PDB 6OQ5[54]. Eight highly conserved surface patches are indicated, and additional details are in Fig. S7. Center residues within each surface patch are indicated in bold font.

Supplementary Data

Figure S1. Phylogenetic and structural overview of the TcdA and TcdB protein family. (a) The TcdA family forms a monophyletic clade with TcsH from *Paeniclostridium sordellii* as a sister phylogenetic lineage. Similarly, the the TcdB family forms a monophyletic clade with TcsL from *Paeniclostridium sordellii* as a sister phylogenetic lineage. This implies a scenario whereby TcdA and TcdB evolved by an ancestral gene duplication that predates the speciation event leading to divergence of *C. difficile* and *P. sordellii*. (b) Representative crystal structures and domain architectures are shown for TcdA (above) and TcdB (below). The structure of TcdA lacks the CROPS domain and is derived from PDB ID 4F04. The full-length structure of TcdB is based on PDB ID (6OQ5), and was modified to remove bound antibodies. Domain definitions were derived from Aktories et al.

a

b

Figure S2. Alignment of TcdA sequences derived from GenBank and the NCBI short read archive, illustrating considerable variation in the length of the C-terminal CROPS region. (a) Complete alignment of 480 unique TcdA sequences. (b) Visualization of unaligned sequences to display C-terminal length variation following residue ~900.

Figure S3. Analysis of sequence similarities within and between subtypes of TcdA and TcdB. Pairwise sequence identities were calculated between all TcdA and TcdB sequences. The % identity distributions are plotted for sequences within ("w") the same subtype versus between ("b") subtypes for TcdA (left) and TcdB (right). As expected, the % identities are much higher within than between subtypes. For TcdA, a % identity threshold of 99.4 effectively distinguishes sequences within the same subtype, whereas for TcdB, a threshold of 97% effectively distinguishes sequences within the same subtype.

Figure S4. SplitsTree analysis of TcdA and TcdB and statistical detection of recombination. Split networks of TcdA and TcdB were generated using the SplitsTree software. Parallel edges suggest the existence of sites that are not compatible with a perfect monophyletic tree, which can result from recombination. An extremely long branch (A7) has been truncated in order to permit visualization.

Figure S5. Haplotype analysis of the CROP-less TcdA alignment. Visualization and analysis of amino acid variation patterns was performed using the HaploColor algorithm (https://github.com/doxeylab/haploColor), which was run for 16 iterations. Patterns of amino acid variation within each phylotype are highly homogeneous, and thus a lack of evidence for recombination. One potential exception is highlighted in yellow, involving two amino acid variants that occur within phylotype A1 that are lacking in most other A1 sequences but present in phylotypes A2 and A3. However, this pattern may also be due to ancestral variation rather than recombination. Overall, compared to TcdB, the TcdA displays considerably less sequence variation and lacks the mosaic patterns that would result from recombination.

Figure S6. Visualization of amino acid variation patterns in TcdB highlighting putative microrecombination events. The TcdB multiple sequence alignment was colored using the HaploColor algorithm (https://github.com/doxeylab/haploColor), which was run for 16 iterations. Fourteen example segments containing amino acid variants that are unexpected for their subtype are shown by white ovals. These represent putative between-subtype microrecombination events. These fourteen are not a complete list as many more can be seen visually.

Figure S7. a) Phylogenetic tree of the GTD domains. B) alignment of region 280-490. According to the tree and the alignment, and the functions, GTD domains could be classed into to groups, one group is TcsL-like which gives vero cells rounding and clumping phenotypes (strain 1470 and 8846); the other group is the classical TcdB-like group which only give the rounding phenotype.

Figure S8. A screenshot of the DiffBase online database.

Table S1. Subtypes of novel TcdA and TcdB sequences identified in the NCBI short read archive. Novel sequences contain at least one substitution not observed in existing sequences derived from NCBI GenBank.

Subtype	#
A1	25
A2	10
A3	3
A4	1
A5	1
A6	1
B1	52
B2	12
B3	2
B4	1
B5	2
B6	7
B7	1
B8	4
B9	3

Table S2. List of clinically relevant and previously studied *C. difficile* strains, associated toxin phylotypes, and toxinotypes. Different groups are assigned unique colors. The table is based on information compiled from Rupnik and Janezic (2016), Bletz et al. (2018), and NCBI genome metadata.

Toxinotype	Strain	Ribotype	Clade	TcdA subtype	TcdB subtype	Combined subtype	Toxin Production	Notes
0	VPI 10463	003/087		A1.1	B1.1	A1/B1	A+B+CDT−	TcdB1
0	630	012	1	A1.4	B1.1	A1/B1		TcdB1
0	E28	012		A1.38	B1.1	A1/B1		
0	T3	012		A1.4	B1.1	A1/B1		
0	E14	014/020		A1.1	B1.2	A1/B1		
0	CD166	014/020		A1.1	B1.5	A1/B1		
	CD111	014/020		A1.1	B1.2	A1/B1		
	CD109	014/020		A1.1	B1.2	A1/B1		
	CD90	014/020		A1.1	B1.2	A1/B1		
0	CD43	027			B1.5	-/B1		
0	E12	106		A1.1	B1.4	A1/B1		
0	5555-DH/ST42	002		A1.1	B1.46	A1/B1		
0	CD002	002			B3.1	-/B3		
0/V	597B	131		A1.14	B1.56	A1/B1	A+B+CDT+	
I	EX 623	102	1	A1.1	B1.109	A1/B1	A+B+CDT+	
II	AC 008	103	1	A1.1	B1.3	A1/B1	A+B+CDT+	
III	R20291	027	2	A2.1	B2.1	A2/B2		TcdB2
IIIb	R 12087 (=CD196)	027	2	A2.1	B2.1	A2/B2	A+B+CDT+	
IIIa	SE 844	080	2	A2.5	B9.1	A2/B9	A+B+CDT+	
IIIc	CH6230	251	2	A2.7	B2.3	A2/B2	A+B+CDT+	
IIIe	AI 541	251	2	A2	B2.7	A2/B2	A-B+CDT+	
IIId	3073	SLO 042	2	A2.11	B2.24	A2/B2	A+B+CDT+	
IV	55767	023	3	A3.2	B5.1	A3/B5	A+B+CDT+	
V	SE 881	045	5	A3.4	B3	A3/B3	A+B+CDT+	
V	M120	078	5	A3.1	B3.1	A3/B3		
	NAP07	078		A3.1	B3.1	A3/B3		
VI	51377	127	5	A3.1	B3.1	A3/B3	A+B+CDT+	
VII	57267	063	5	A3.1	B3.7	A3/B3	A+B+CDT+	
VIII	1470	017	4		B4.1	-/B4	A−B+CDT−	
VIII	M68	017			B4.1	-/B4		
VIII	E13	017			B4.1	-/B4		
IXa	51680	019	2	A2.2	B6.1	A2/B6	A+B+CDT+	
Ixb	TFA/V20-1	244	2	A2.6	B6.2	A2/B6		
IXc	8785	109	5	A2	B6.5	A2/B6	A+B+CDT+	

IXd	1732874	SLO 228 036 /	2	A2.8	B6.13	A2/B6	A+B+CDT+
Xa	8864	591(CE)	2		B7.1	-/B7	A−B+CDT+
Xb	J9965	SLO 032	2		B7.4	-/B7	A−B+CDT+
XIa	IS 58	033	5			-/-	A−B−CDT+
XId	OCD 5/2	033	5			-/-	A−B−CDT+
XIb	R 11402	288 (CE)	5			-/-	A−B−CDT+
XII	TFA/V14-10	153(CE)	2			-/-	A−B−CDT+
XII	IS 25	258	1	A1.1	B1.10	A1/B1	A+B+CDT-
XIII	R 9367	070	1		B1.2	-/B1	
XIVa	R 10870	111	2	A2	B6.3	A2/B6	A+B+CDT+
XIVb	R 9385	122	2	A2.12	B6.6	A2/B6	A+B+CDT+
XVI	SUC36	078	5	A3.8	B3.6	A3/B3	A+B+CDT+
XVIII	K095	014	1	A1.1	B1.105	A1/B1	A+B+CDT−
XIX	TR13	018	1	A1.2	B1.2	A1/B1	A+B+CDT−
XX	TR14	SLO 005	1	A1.1	B1.16	A1/B1	A+B+CDT−
XXI	CH6223	SLO 035	4	A1.35	B4.11	A1/B4	A+B+CDT−
XXII	CD07-468	027	2	A1.36	B2.1	A1/B2	A+B+CDT+
XXV	7325	027	2	A2.1	B2.1	A2/B2	A+B+CDT+
XXVI	7459	050 (CE)	1		B1.6	-/B1	A-B+CDT−
XXVII	KK2443/200 6	SLO 037	1			-/-	A-B-CDT−
XXVIII	CD08-070	126	5	A3.1	B3.1	A3/B3	A+B+CDT+
XXIX	CD07-140	001	1	A1.1	B1.2	A1/B1	A+B+CDT−
	CD92	001		A1.1	B1.2	A1/B1	
XXX	ES 130	SLO 101	5		B8.1	-/B8	A-B+CDT+
XXXI	WA 151	SLO 098			B8.2	-/B8	A-B+CDT+
XXXII	173070	151(CE)	C-II		B12.1	-/B12	A-B+CDT-
XXXIII	2402	SLO 086	1	A1.37	B4.10	A1/B4	A+B+CDT-
XXXIV	CD10-055	SLO 201			B2.7	-/B2	A-B+CDT-
XXXIII	2402	SLO 086		A1.37	B4.10	A1/B4	A+B+CDT-
	RA09-70			A7.1	N	A7/-	A+B-CDT-
	CD160				B11.1		
	HMX-149			N	B11.2	-/B11	A-B+CDT-
	CD10-165			C-I	N	B10.1	-/B10	A-B+CDT-
	SA10-050			C-I	N	B10.2	-/B11	A-B+CDT-
	HSJD-312			N	B6.9	-/B6	A-B+CDT+
	HMX152			N	B6.9	-/B6	A-B+CDT+