DNA methylation site loss for plasticity-led novel trait genetic fixation

Takafumi Katsumura[1,2,*], Suguru Sato[3], Kana Yamashita[3], Shoji Oda[4], Takashi Gakuhari[5], Shodai Tanaka[3], Kazuko Fujitani[6], Toshiyuki Nishimaki[1], Tadashi Imai[7], Yasutoshi Yoshiura[7], Hirohiko Takeshima[8], Yasuyuki Hashiguchi[9], Yoichi Sekita[10], Hiroshi Mitani[4], Motoyuki Ogawa[1], Hideaki Takeuchi[2,11] and Hiroki Oota[1,12]

[1]Department of Anatomy, Kitasato University School of Medicine, Japan

[2]The Graduate School of Natural Science and Technology, Okayama University, Japan

[3]Department of Biological Sciences, Kitasato University School of Science, Japan

[4]Department of Integrated Biosciences, The University of Tokyo, Japan

[5]Institute of Human and Social Sciences, Kanazawa University, Japan

[6]Gene Analysis Center, Kitasato University School of Medicine, Japan

[7]National Research Institute of Fisheries and Environment of Inland Sea, Japan Fisheries Research and Education Agency, Japan

[8]Department of Marine Biology, Tokai University School of Biological Sciences, Japan

[9]Department of Biology, Osaka Medical College, Japan

[10]Department of Biosciences, Kitasato University School of Science, Japan

[11]Graduate School of Life Science, Tohoku University, Japan

[12]Department of Biological Sciences, Graduate School of Science, The University of Tokyo, Japan

*Correspondence to:

Takafumi Katsumura, Department of Anatomy, Kitasato University School of Medicine, Japan, +81 42 778 9022, tk@med.kitasato-u.ac.jp

Abstract

Many organisms exhibit phenotypic plasticity that changes their traits in response to their environment.

Whether or not this plasticity contributes to adaptive evolution is a fundamental question in

evolutionary biology because various studies report that natural populations adapt to rapid

environmental changes via plasticity, which often "triggers" novel adaptive traits. While such

alternative phenotypes could be induced by nongenetic perturbations that include gene expression

noise or epigenetic modification caused by environmental change, it remains unknown what molecular

mechanism genetically fix the alternative phenotypes as adaptive traits passed into the next generation.

Here we show that a decrease in methylated CpG sites leads to loss of plasticity, which triggers genetic

fixation of novel traits, in medaka fish (*Oryzias latipes*). We found that the gut length was correlated

with the number of methylated CpG sites upstream of the *Plxnb3* gene. The medaka, in which the

methylated DNA region is deleted by CRISPR/Cas9, showed a loss of plasticity in gut length.

Moreover, standing variation in the promoter region of another gene, *Ppp3r1*, made a longer gut stably

in wild medaka groups that lost the gut-length plasticity. Furthermore, our phylogenetic analysis

revealed the timing of these evolutionary events, indicating that the loss of phenotypic plasticity by

nucleotide substitutions initiates the process of genetic fixation of the novel trait. That is, while

phenotypic plasticity plays a role as a buffer of evolution and contributes to environmental adaptation,

as previously thought, our molecular data suggest that mutation on CpG site causing the loss of

phenotypic plasticity, is the trigger for a generation of novel traits.

Introduction

Genetic and environmental factors that alter a phenotype are driving forces leading to novel traits. In

current biology, a novel mutation that is considered to cause a novel trait spreads into a population

through generations if it is advantageous. Meanwhile, traits often show variances induced by

environmental changes (phenotypic plasticity[1]). The altered trait, being exposed under selective

pressure over generations, can be expressed regardless of environmental factors by subsequent genetic mutations through generations, even in nature[2]. This process is called, the "plasticity-first" evolution (PFE), which is a candidate for an alternative pathway of adaptive evolution[3]. However, it has been discussed over the years whether the existing evolutionary framework can explain the PFE process[4] because the molecular mechanism of how novel traits that originated from plasticity lead to genetic fixation remains unknown.

The PFE process involves the expression of phenotypic plasticity in ancestral lineages and genetic fixation in derived lineages[3]. In other words, PFE is a process in which environmentally induced traits lose their environmental responsiveness and is fixed genetically during their evolution. Therefore, in order to investigate molecular mechanism involved in the PFE process, it is reasonable to divide this process into two steps: (1) loss of phenotypic plasticity and (2) genetic fixation of an induced trait. The expression of phenotypic plasticity is considered to occur as a result of changes in gene expression patterns caused by epigenetic changes (such as DNA methylation[5–7]). Moreover, the PFE hypothesis states that following random mutations fix plastic traits induced by environmental change and are transmitted to offspring. If this hypothesis is true, the genetic mutation inducing the altered trait as a novel trait may exist in the epigenetic modification region related to the expression of phenotypic plasticity. Therefore, we sought the molecular basis of these two steps, phenotypic plasticity and the genetic fixation of novel traits, and examined whether they involved in such a PFE process using molecular biological experiments. To this end, we focus on gut length, which shows typical phenotypic plasticity known to be an adaptive trait that enhances nutrient absorption efficiency responding in the feeding environment[8] and genetic variation in various animals[9,10].

The medaka (*Oryzias latipes*), also known as Japanese rice fish, is a small, omnivorous freshwater fish that inhabits low to high latitudes in East Asia. Their local populations adapt genetically to the various environments of each region, and show geographic phenotypic plasticity and latitudinal variations for the traits associated with gut length[11,12]. Moreover, their evolutionary history

is well-studied at the genomic level[13,14], and various molecular biological techniques can be adapted to

them[15], which makes medaka an excellent model to test whether the focal trait meets the critical

criteria (phenotypic plasticity in the ancestral lineage and genetic fixation in the derived lineage) of

PFE[2] and allows us to investigate the molecular mechanism of the PFE process. Thus, we aimed to

show the PFE process underlying their gut length differences and reveal the molecular basis and

evolutionary process from the loss of plasticity to genetic fixation.

These evolutionary-ecological features, coupled with two genome-wide approaches and

subsequent genome-editing, allowed us to describe the molecular mechanism of PFE underlying gut-

length variations of local medaka populations. First, we found that the evolution of the medaka gut

fulfills the criteria of PFE by measuring the gut-length, which revealed seasonal and local geographical

variations. Second, we found candidate genes involved in the two steps of the PFE process by genome-

wide methylation and population genomic analyses. Finally, by analyzing the functions and molecular

evolution of the two genes, we revealed that a longer gut became genetically expressed through

fixation of mutations after the loss of gut-length plasticity due to a decrease in CpG sites. The new

finding is that PFE is not a process of genetically fixing phenotypic plasticity by the following

mutation, but a process in which plasticity, a buffer that reduces environmental pressure, is lost by

mutation and often replaced by an existing favorable mutation.

Results

We first verified whether or not the medaka gut shows the phenotypic plasticity on ancestral lineages

which are one of the requirements of PFE. We investigated the gut length, collecting wild medaka in

three rivers, the Kabe, Ejiri, and Shihodo, in Kagawa Prefecture, Japan (Fig. 1A and table S1). We

measured the gut length of 18–23 medakas in each river in two seasons, winter and summer, over a 2-

year period, and found statistically significant changes in the gut length relative to standard length in

the Kabe and Ejiri rivers between seasons (fig. S1). Especially in the Kabe river, the medaka in the

winter had significantly shorter guts than did those in the summer, which was observed repeatedly over

two years (Fig. 1B, Bayesian generalized linear mixed model [GLMM], estimate = -0.44±0.05, 95%

credible interval [CI] did not include zero, table S2 and fig. S1). By plotting body and gut lengths in

each season, we found no correlation between them in the summer when the gut is longer (fig. S2). To

investigate whether or not this change in gut length caused by mixing genetically distinct populations,

we analyzed genetic diversity change between winter and summer based on the single nucleotide

polymorphisms (SNPs) in the mtDNA and chromosomes and did not detect any noticeable changes of

genetic diversity including gene flow from other populations (fig. S3). Taken together, with the

lifespan of medaka in the wild (1–1.5 years)[16], no overwintering twice[17] and no significant difference

in the body length between seasons (fig. S1), these results indicated that the gut length changes in the

Kabe river medaka must be phenotypic plasticity within individuals in response to the environment.

The gut-length change was often related to the food source[10,18]. To examined whether food

source changes between the seasons, we conducted stable isotope analysis of the caudal fins in three

rivers. The stable isotope analysis based on $\delta^{13}C$ and $\delta^{15}N$ showed no correlation with gut length (fig.

S4), indicating food diversity or carnivorous habits did not strongly governed this gut-length plasticity.

To test whether or not the food source affects the gut length, we transported the wild captured medaka

from the Kabe river into our outdoor breeding facility and bred them in an artificial condition. The

outdoor breeding experiment showed the same plasticity even under the condition of feeding with

artificial food (fig. S5, table S3), indicating no relationship between the plasticity and the food source.

The gut-length plasticity could be caused by other environmental factors that we could not measure,

such as the amount of food consumed.

Figure 1 | Phenotypic plasticity and genetic variation of the gut length in medaka. (A) A map

of sampling locations (gray circles) in Kagawa Prefecture, Japan. (B) Seasonal gut length

changes in the three rivers. The blue and orange dots, their error bars indicate the mean gut

length and 95% CI (credible interval) after adjusting for the effect of body length in winter (W)

and summer (S), respectively, as estimated by Bayesian GLMM. The red dash line joins the

mean in each season representing the seasonal variation. (C) Geographic variations of gut

length in 40 local populations with example photos (white bars: 1 cm). The phylogenetic tree

and ADMIXTURE barplot were constructed using genome-wide single nucleotide

polymorphisms (SNPs) published in our prior study[13]. The right plot shows the geographical

variation of the gut length corrected by body length among medaka populations. Their points

and error bars were fitted values with 95% CIs from the Bayesian GLMM. Abbreviations on

the far right: S1–S4, N1, 2 and K are: SJPN (Southern Japanese group) 1–4, NJPN (Northern

Japanese group)1, 2 and KOR (Korean/Chinese group), respectively. The asterisk indicates

that the NJPN1 medaka (the red branches in the phylogenetic tree) have a significantly

longer gut than do the other subgroups.

We subsequently verified whether or not the medaka gut shows genetic fixation on derived lineages, which is also one of the requirements of PFE, using wild-derived laboratory stocks consisting of 81 geographical local populations. Our previous genome-wide SNP study showed that these stocks cover a genetic diversity of wild medaka and were genetically divided into seven subgroups (SJPN1-4, NJPN1, NJPN2, and KOR) derived from three lineages from the northern Japanese (NJPN), the southern Japanese (SJPN), and the Korean/Chinese (KOR) medaka groups[13]. Based on the population genetic background of these stocks, we sampled 10 individuals from 40 out of 81 populations to avoid bias from population structure (table S4) and measured the gut length dissected. All local populations were reared in the same environment in our outdoor facility. Nevertheless, the NJPN1 subgroup showed a significantly longer gut compared to the average body length than did those of the other subgroups (Fig. 1C, Bayesian GLMM, estimate = -0.11 – -0.26, 95% CIs did not include zero, table S4 and S5), indicating that a longer gut length of the NJPN1 subgroup is fixed by genetic factors common within this subgroup. Meanwhile, the gut lengths within the SJPN and KOR groups vary, indicating that the gut length diversity in the SJPN and KOR groups is defined by genetic factor(s) or environmental factors. As in the wild medaka in Kagawa (Fig. 1B), to examine whether or not the gut of these stocks also showed seasonal plasticity, we compared gut length between seasons using the available stocks from each group (fig. S6A). In NJPN1, the gut length was longer than the body length regardless of the season, although an exception was observed. In NJPN2, there was no significant variation in gut length by season, except in the Toyooka population. In contrast, SJPN, which shares ancestry with the Kagawa medaka, showed large seasonal fluctuations in gut length. Our GLMM analysis showed that the genetic background of SJPN interacted with the seasons and influenced the gut length (fig. S6B, table S6). Together with the seasonal plasticity of the Kagawa medaka which belong to the SJPN group (Fig. 1B), these results indicate the phenotypic plasticity of gut length exists at least in the SJPN lineage. Therefore, based on the evolutionary history of medaka[13], the ancestors of

medaka originally had phenotypic plasticity in gut length, and the derived subgroup (i.e., NJPN1) of

medaka probably acquired a longer gut length to adapt. Thus, we concluded that the evolution of the

medaka gut meets the requirements for PFE.

Under the PFE hypothesis, DNA methylation, which occurs as environmental responses to

seasons and is stable over a long time if the seasonal environment remains unchanged[19], is a potential

candidate as a molecular basis of seasonal phenotypic plasticity. To test whether the DNA methylation

is related to gut length plasticity, methyl-CpG binding domain sequencing (MBD-seq) was used to

detect the methylation of DNA extracted from the gut of Kabe river medakas which we observed as

exhibiting the strongest seasonal plasticity (Fig. 1B, fig. S1). The MBD-seq analysis revealed hypo- or

hypermethylated regions associated with winter or summer, respectively, for 2 years (Fig. 2A and fig.

S7). We detected a total of 206 seasonally varying methylated regions: 14 and 192 regions

hypermethylated in winter and summer, respectively. Annotated these regions and explored the genes

in their vicinity (fig. S8), a total of 71 genes were present in the regions within 2 kb (kilobase)

upstream and downstream (two and 69 genes in winter and summer methylated regions, respectively)

(Fig. 2A, table S7). We conducted the enrichment analysis about the gene lists using Reactome

(https://reactome.org/) and Gene Ontology (GO) (http://geneontology.org/) analyses, and found that the

semaphorin-plexin signaling related genes (*Plexin a3* [*Plxna3*], *Plexin a4* [*Plxna4*], and *Plexin b3*

[*Plxnb3*]) concerning the neuronal axon guidance were significantly enriched by both analyses (Fig.

2A, table S8). Notably, the *Plxnb3* gene, which is involved in the inhibition of neurite outgrowth, had a

CpG island (CGI) in the upstream region showing seasonal methylation (Fig. 2B). We then performed

both the whole-genome bisulfite sequence (WGBS) and the targeted bisulfite sequence (tBS) to

determine which CpG sites in the upstream regions were seasonally methylated. The WGBS analysis

reconfirmed methylation changes in 63 regions detected by the MBD-seq (fig. S9), and the Reactome

analysis again showed enrichment of the semaphorin-plexin signaling-related pathway using the list of

adjacent genes in these regions (table S7). In the tBS analysis, a statistically significant seasonal

difference in methylation was observed at the second CpG site (fig. S10), although a G/A SNP was

detected at the 5th CpG site which led to the difference in the methylation level. Furthermore, the

methylations of the second and the adjacent CpG sites were significantly correlated and seasonally

varied (fig. S11). These correlations were not observed in the populations collected from the Ejiri and

Shihodo rivers where seasonal changes in gut length were unclear (fig. S11). A real-time quantitative

polymerase chain reaction (PCR) showed that the gene expression of *Plxnb3* in summer was lower

than that in winter ($P < 0.01$), indicating the gene expression was correlated with the seasonal

methylation on the upstream region of *Plxnb3* (Fig. 2C). Considering the *Plxnb3* gene function and

correlation, the upstream region showing the seasonal methylation might be the molecular basis for

regulating the gut length plasticity.

Figure 2 I DNA methylation map and *Plxnb3* gene expression pattern correlated with winter

and summer. (A) Two hundred six differentially methylated regions (DMRs) were filtered

using Reactome, GO analyses, and the position of the methylated region for the gene. The

color difference of the heat map shows the difference in the degree of methylation, i.e.,

orange and blue show the hypermethylated and hypomethylated region, respectively. The

four different colors of bar on the top of heatmap represent the combinations of season and

year, which are also assigned to Fig. 2B and 2C. WsWs and wSwS represent

hypermethylation in winter and summer, respectively. (B) The upstream region of *Plxnb3*

showed seasonal methylation, which included the CpG island (denoted by the gray CGI bar).

(C) *Plxnb3* gene expression was suppressed in the summer. The inset graph shows the

difference in the gene expression between seasons for two years (2014 + 2015) at the time.

A significant difference in gene expression between seasons was detected using the

Wilcoxon rank-sum test (**$P < 0.01$).

      The plasticity observed in the wild population was that the change of the gut length responded

to the seasonal environment. Thus, if the environment is constant, the gut length will be constant

regardless of body length in individuals that show plasticity. In contrast, in individuals that have lost

plasticity, the gut length will increase with body length without responding to the constant

environment. To examine whether the upstream sequence of *Plxnb3* is associated with seasonal

plasticity of the gut, we generated a genome-edited medaka, *ΔupPlxnb3*, in which the region showing

seasonal DNA methylation was deleted using CRISPR/Cas9. We then reared a population of these

three genotypes obtained by heterozygous mating in a laboratory constant environment (14/10

light/dark cycle at 28°C) and examined the relationship between body and gut length among the

genotypes. At first, testing the segregation ratio of genotypes after a heterozygous mating showed the

homozygous and heterozygous rates of medaka with the deletion were about half that compared with

the expected segregation rates (Fig. 3A, $P = 0.011$ using the chi-square test). Next, comparing

genotypes, no significant difference in gut length was detected, but a significant difference in body

length was observed between the wild type and mutant (Fig. 3B, fig. S12). Finally, by plotting body

and gut lengths (Fig. 3C, fig. S12) for each genotype, we found dependency between those lengths in

the mutants but not in the wild types (Bayesian generalized linear model [GLM], the effect size of the

deletion and interaction between standard length and the deletion; estimates = -1.67±0.67 and

0.08±0.03; each 95% CI did not include zero, table S9). These results show that the mutants could have

a gut that lengthens in proportion to its body length without responding to the laboratory environment,

indicating that the mutants with deletion of the *Plxnb3* upstream sequence lost gut plasticity.

The lower survival rate and shorter body length of the mutants led us to hypothesize that the

gut of each individual lacking the upstream sequence of *Plxnb3* undergoes some histological changes.

Therefore, we investigated the distribution of neuronal axons involved in the *Plxnb3* by

immunostaining, using neurofilament H protein as a marker (Fig. 3D). Compared with the wild-type

intestinal villi, we found that the immunohistological signal disappeared at the tip of the intestinal

villus in the mutants (Fig. 3D and E). Moreover, the *Plxnb3* expression level of the mutants showed

significantly greater variance than that of the wild type, indicating the gene expression regulation

became unstable (Fig. 3F). These results suggest that the seasonal methylated region includes the

regulatory region of *Plxnb3*, which that deletion may lead to the disruption of neural mechanisms that

senses the seasonal changes of feeding environment (such as sensory-cell mediated food detection[20]).

Then, it may have caused inadequate nutrient absorption adapted to the environmental change,

resulting in a short body length and a lower survival rate (Fig. 3A and B). Overall, our data indicate

that the upstream region of *Plxnb3* contributes to the plasticity of gut length, and it is plausible that the

seasonal methylation that occurs in the upstream region is linked to the plasticity of gut length.

Figure 3 | Effects of the deletion of the *Plxnb3* upstream region on gut plasticity. +/+, +/-, and

-/- represent the wild type, the heterozygote, and the homozygote of the *Plxnb3* upstream

deletant (i.e., the wild type/*ΔupPlxnb3* and *ΔupPlxnb3*/*ΔupPlxnb3*), respectively. (A) The

deviation of the segregation ratio among genotypes was evaluated using the chi-squared

test. (B) Comparisons of standard and gut lengths among genotypes were performed in the

Bayesian generalized linear model (GLM) framework. Points and error bars were fitted values

and 95% Bayesian CIs (full data in fig. S12). The asterisk indicates that -/- exhibited

significantly shorter body length than did the other genotypes. (C) The gut length is

associated with the standard length in the mutants with deletion, not the wild type. Lines and

shaded areas are fitted values and 95% CIs from the Bayesian GLM. The yellow and blue in

(B) and (C) indicate the wild type and mutants (both of the heterozygote and homozygote),

respectively. Full data of (B) and (C), and their estimates, are in fig. S12 and table S2,

respectively. (D) Immunochemical staining revealed the decrease of neurofilament-positive

cells at the tip of the intestinal villus. (E) The difference in the degree of immunochemical

staining (D) was statistically quantified and detected using the Wilcoxon rank-sum test (*$P <$

0.05). (F) *Plxnb3* expression showed the uneven variance between the wild types (+/+) and

the mutants (-/-), which was evaluated using the F-test (*$P < 0.05$), indicating the unstable

Plxnb3 expression in the mutant gut. The orange and blue in (E) and (F), indicate the wild

type and the homozygote of the *Plxnb3* upstream deletant, respectively.

Under the PFE hypothesis, a genetic mutation should occur and fix a plasticity trait. Because

the NJPN1 subgroup exhibited a long genetically fixed gut (Fig. 1C), the causative genetic mutations

should be presumed to be highly frequent in the subgroup. In order to examine which gene is

responsible for the longer gut of the NJPN1, 62 K SNPs obtained by RAD-seq[21] were used to search

genomic regions that were differentiated in the NJPN1 subgroup and associated with the gut length

(Fig. 4A). We found that a candidate signal strongly associated with gut length, resulting in a highly

differentiated population, was detected in the *Protein phosphatase 3 regulatory subunit B, alpha*

(*Ppp3r1*) gene on chromosome 1, which is also involved in neuronal axon outgrowth[22]. Bayesian

GLMM analysis confirmed that the T/T genotype had a significant effect on gut length compared to

the C/C genotype (Fig. 4B, table S10). This mutation was a synonymous SNP, with no linked non-

synonymous SNPs in the coding region within the NJPN1 subgroup (fig. S13), which predicted that

the changes in the *Ppp3r1* expression contributed to the gut length. Given that a previous study of

conditional *Ppp3r1* KO mice showed hypoplasia of the small intestine[23], we could hypothesize that the

difference of this gene expression in medaka would lead to geographical differences in gut length. To

test this hypothesis, we examined the relationship between *Ppp3r1* expression levels and gut length

using outdoor-reared wild-derived laboratory stocks in the winter because a remarkable difference in

gut length was expected. We found that the gut length was significantly correlated with the *Ppp3r1*

expression level unless genetic background was controlled (Fig. 4C, fig. S14, table S11), and the

NJPN1 medakas with a homozygous T allele showed higher *Ppp3r1* expression than other subgroups

with a homozygous C allele (Fig. 4C, $P < 0.01$). In addition, in the same genetic background, a

homozygous T allele showed higher *Ppp3r1* expression than a homozygous C allele (fig. S15A, B),

and the gut length was significantly correlated with the *Ppp3r1* expression level (fig. S15C). These

results indicate that the mutation associated with the up-regulation of the *Ppp3r1* expression is

involved in the genetic fixation for the longest gut among the medaka subgroups.

Figure 4 | Differentiation and association SNPs with gut length across the medaka

populations. (A) F_{ST} was calculated between NJPN1 and the others. The red line indicates

0.59592 Weir and Cockerham's weighted F_{ST}. Association P-values with gut length were

calculated using 22,235 SNPs filtered by a minor allele frequency of >0.05. The blue dashed

line shows the P-value in which the top 10 SNPs distinguished from the others (table S12).

The red points show that significant SNPs fulfilled the above criteria. (B) The effect size of

Ppp3r1 genotype of gut-length associated SNPs (Cytosine [C] or Thymine [T]) was estimated

from the posterior mean of a Bayesian GLMM and is given relative to the model intercept at

the C/C genotype (dashed line). Vertical bars indicate 95% CI. (C) *Ppp3r1* expression was

significantly correlated with gut length estimated by Bayesian GLMM (effect size of the

Ppp3r1 expression: estimate =0.42±0.12; 95% CIs did not include zero, full data in fig. S14).

Line and shaded area are fitted value and 95% CIs from the Bayesian GLMM. The inset

graph shows the difference in the *Ppp3r1* expression between the C/C or T/T genotypes,

which were examined using winter-sampled individuals. The statistical significance was

evaluated using the Wilcoxon rank-sum test (**$P < 0.01$).

Finally, we analyzed the molecular evolution of two identified regions based on the medaka

population phylogeny. First, we determined the *Plxnb3* upstream sequences to estimate the evolution

of methylated regions associated with plasticity in the medaka populations (Fig. 5A, fig. S16). A

comparison of the upstream sequences revealed that the number of CpG sites varied between the

populations, indicating nucleotide substitutions resulted in the gain and/or loss of CpG sites (fig. S17).

Ancestral sequence estimates indicated that the common ancestor of SJPN and NJPN had eight CpG

sites in this upstream region, and the number of CpG sites diversified from seven to 10 upon branching

to each SJPN lineage. On the other hand, the number of CpG sites in the *Plxnb3* upstream was reduced

to six in the common ancestor of NJPN1 and NJPN2, which did not meet the criteria of CpG islands

(GC >50%, Obs/Exp >0.6, and length >100 bp). We subsequently examined our RAD-seq data and

observed the *Ppp3r1* mutation in the NJPN1 subgroup and the SJPN1 Iwata population, which occur

both on the Sea of Japan and Pacific Ocean sides of the eastern Japanese archipelago, respectively.

These results suggest that the mutation appeared at least in the common ancestor of SJPN and NJPN

that spread throughout the Japanese archipelago and subsequently existed as a standing variation.

Considering the phylogenetic relationship among groups (Fig. 1C, fig. S18), the mutation in *Ppp3r1*

must have spread in NJPN1 after the number of CpG sites in the *Plxnb3* upstream decreased in the

common ancestor of NJPN1 and NJPN2.

To investigate the relationship between *Plxnb3* and *Ppp3r1*, we estimated the effects of the

number of CpG sites in the *Plxnb3* upstream and the presence or absence of the *Ppp3r1* mutation on

gut length using GLMM. We found a positive correlation between the number of CpG sites in the

Plxnb3 upstream and gut length among the populations with or without the *Ppp3r1* mutation,

respectively (Bayesian GLMM, estimate =0.06, 95% CIs did not include zero, table S13) and that the

long guts tended to appear depending on the number of CpG sites (Fig. 5B). In contrast, the *Ppp3r1*

mutation had the effect of inducing a long gut regardless of the number of CpG sites in the *Plxnb3*

upstream (Fig. 5B, table S13). Furthermore, the plot (Fig. 5B) showed that the variability in gut length

was small for each *Ppp3r1* genotype when the number of CpG sites was lowest. These results suggest

that the number of CpG sites in the *Plxnb3* upstream is involved in the expression of plasticity.

Supporting this possibility, we found that NJPN1 and NJPN2 medaka with the lowest CpG sites lost

seasonal plasticity in gut length (fig. S6). Furthermore, the *Ppp3r1* mutation was not observed in the

Kabe river populations showing seasonal plasticity but in the NJPN1 wild population (fig. S19). These

additional results more strongly suggest the gut-length plasticity and genetic fixation of a long gut are

associated with the number of CpG sites in the *Plxnb3* upstream and *Ppp3r1* mutation, respectively.

Taken together with the molecular evolution of these two regions, our data suggest that the mutation

spread to the NJPN1 subgroup after loss of plasticity due to reduction of the number of CpG sites in

the *Plxnb3* upstream, after which a long gut became expressed in northern populations due to the

standing variation of *Ppp3r1* (Fig. 5A).

Figure 5 | Evolutionary relationship between the *Plxnb3* upstream region and the *Ppp3r1*

mutation. (A) The number of CpG sites increased or decreased at each node on the

population tree. The red arrowheads indicate evolutionary events, which were inferred from

the current proportion of the number of CpG sites on the *Plxnb3* upstream region and the

allele frequency of gut-length associated SNPs on the *Ppp3r1* (gray: C-allele; yellow: T-

allele). The gradient from gray to dark green at the top of the horizontal bar graph indicates

the number of CpG site on the *Plxnb3* upstream region. The right plot shows the effect sizes

of each genetic background estimated as a fixed effect in the Bayesian GLMM framework.

The effect size is given relative to the model intercept at NJPN1 (dashed line), and each 95%

CI of effect size did not include zero, indicating the long gut are fixed genetically in the

NJPN1 subgroup. (B) A correlation between the number of CpG sites in the *Plxnb3* upstream

and gut length in each local population. Gray and yellow points indicate C- and T-alleles of

Ppp3r1 fixed in each population, respectively. The white line and shaded area are fitted

values and 95% CIs from the Bayesian GLMM. (C) A relationship between gut length and

annual temperature among subgroups ($r = 0.869$, $P = 0.025$). X-axis and Y-axis indicate the

standard deviation (of annual temp.) of their original habitats and the average of coefficient of

variation (of relative gut length) within genetic group, respectively. The abbreviations and

colored points in (A) and (C) are the same as those in Fig. 1C.

Discussion

In this study, we hypothesized that DNA methylation regulates phenotypic plasticity and that loss of

the CpG site to be methylated leads to loss of phenotypic plasticity, and found the correlation between

seasonal variation in gut length and methylation of the upstream region of the *Plxnb3* gene in medaka.

We also found that deletion of the upstream region of *Plxnb3* leads to a loss of gut plasticity (Fig. 5B).

Furthermore, a seasonal variation in gut length was not observed in populations with the lowest

number of CpG sites (fig. S6). These results indicate that the seasonal variation of CpG site

methylation in the upstream region of *Plxnb3* contributes to the gut-length plasticity in medaka. In

addition, it is not because substitutions of the CpG sites, that were seasonally methylated, resulted in

the loss of gut length plasticity, since those CpG sites are conserved among medaka populations (fig.

S17). Thus, the loss of plasticity may not be due to the loss of CpG sites with seasonal methylation but

rather to the loss of nearby CpG sites that are stably methylated through seasons. Considering that

methylation of promoter regions affects transcription factor binding[24], the decrease of CpG sites in the

upstream region of *Plxnb3* would weaken the influence of seasonal CpG methylation on transcription

factor binding. In other words, constantly methylated CpG sites in the upstream region may have

limited the binding of transcription factors only to seasonally methylated CpG sites, consequently

making the effects of seasonal methylation of CpG sites more obvious. Under this hypothesis, it can be

predicted that the difference in the number of stably methylated CpG sites in the upstream region

results in the variability in the *Plxnb3* gene expression. In practice, the variability of the *Plxnb3* gene

expression in the population with more CpG sites was smaller than that in the population with fewer

CpG sites (fig. S20). Although it remains to be addressed whether or not the stably methylated CpG

sites enhance the influence of the seasonal methylation of CpG sites in the vicinity, our results provide

strong evidence that DNA methylation is involved in regulating plasticity and that loss of CpG sites

results in the loss of plasticity.

Genome-wide SNP analysis indicated that a *Ppp3r1* mutation, which is present as a standing

variation, contributed to genetically fixing of the long gut observed in NJPN1 (Fig. 4). This gene is the

most crucial candidate because it is involved in neuronal axon outgrowths like *Plxnb3* and small

intestine formation[23]. Because increase in the *Ppp3r1* gene expression strongly correlates with gut

elongation (Fig. 4 and fig. S15), a mutation responsible for gene expression might exist in the

regulatory region, which is linked to the synonymous mutation in *Ppp3r1* found in this study.

Moreover, NJPN1 showed the *Ppp3r1* expression higher than the other subgroups (fig. S15A),

suggesting that lineage-specific mutation(s) may be present. On the other hand, this associated

mutation (or a linked unknown mutation) explains only the gut length variation in NJPN1 and a part of

the SJPN1 subgroup (Fig. 5). The long guts of the other subgroups may be attributable to genetic

variations originated by other mutations that occurred and/or were selected independently in each

population or a plastic variation induced by environmental pressures under artificial rearing conditions.

To address this matter, crossbred experiments will be necessary in future studies. Nevertheless, it is an

important finding to understand the process of PFE in which a pre-existing genetic mutation in the

genomic region completely different from those involved in gut length plasticity, is involved in the

genetic fixation of the long gut in medaka.

This study shows that two genes involved in neuronal axons development contribute to

phenotypic plasticity and genetic polymorphism of gut length. It was clarified at the molecular level

that the plasticity was associated with seasonal methylation of CpG in the *Plxnb3* upstream and that the

genetic mutation in *Ppp3r1* spread to populations that the methylation site had reduced by the mutation

(i.e., nucleotide substitution). In short, these molecular mechanisms and evolutionary processes

indicate that the mutation-driven evolution occurred in gut length. This differs from "plasticity-driven

evolution," which is based on phenotypic plasticity, in which the adaptive trait is led by natural

selection and the following mutation fixes it (summarized as, "selection-driven evolution" in Nei

[2013][25]). Furthermore, our results indicate that genetic fixation of a plasticity-led trait would occur

under the following process. The phenotypic plasticity regulated by CpG methylation plays the role of

a buffer for environmental pressure, which contributes to the accumulation and maintenance of *de novo*

mutations as standing variations. When plasticity is lost, however, the adaptive mutation is selected

from standing variations, and then the adaptive phenotype is genetically fixed. Our data indicate that

this process is likely to be the PFE observed in nature, and the hereditary phenomenon of the

environmentally induced trait could be explained by the genetic framework of loss of CpG sites and

fixation by standing variations.

17   Why has a genetic fixation of the long gut occurred in the NJPN1 medaka? SJPN and NJPN

medaka originate in northern Kyushu and the Tajima-Tango region, respectively[13]. In SJPN medaka,

populations with similar genetic backgrounds inhabit the different climatic environments. Because the

foraging environment is susceptible to annual temperature changes and the gut length has become

varied in response to the average temperature variation (Fig. 5C), the gut length would have been

plastically adjusted for efficient digestion and absorption in various foraging environments. On the

other hand, NJPN1 is a group that has spread rapidly to the north from small habitats where the

environment is relatively uniform[13]. In the habitat of NJPN2, both temperature and gut length show

minimal variability (Fig. 5C), while the NJPN1 habitats show a large temperature variation and short

growing period because of the high latitudes of their habitats[11]. Therefore, the NJPN2 medaka did not

need to change their gut length, where as the NJPN1 medaka must feed during the short summer and

prepare for the winter. Under these circumstances for the NJPN1 medaka, maintaining a long gut

throughout the season may have served as food storage, rather than regulating gut length to increase

absorption efficiency as in SJPN medaka. Moreover, the foraging behavior of NJPN1 medaka that has

expanded into the higher latitude regions is more frequent than that of medaka in the lower latitude

regions[11]. This indicates that genetic mutations which can change gut length may be driving

geographic differences in behavior of foraging. Indeed, the neurofilament protein involved in sensory

neural regulation in the gut[26] was not stably expressed in medaka that had lost plasticity (Fig. 3D). This

result suggests that the gut could not be detecting the appropriate amount of feeding, which may lead

to excessive feeding behavior. I.e., the gut-brain interaction may have enhanced the genetic fixation of

advantageous mutation.

The identified molecular mechanisms and the above evolutionary inference leads to the

conclusion that plasticity is easily lost under a stable environment, and that after loss of plasticity, a

favorable mutation can easily be fixed on foraying into harsh habitats. In a macroscopic-type view, this

phenomenon may appear as though the evolution of acquired traits has occurred because it occurs

continuously.

Materials and Methods

Ethics statement

The Institutional Animal Care and Use Committee of Kitasato University and The University of Tokyo

approved all the experimental procedures (No. 4111 and C-09-01, respectively).

Sampling and measurement of gut length

To detect the seasonal plasticity of gut length, we collected 228 medaka individuals from three rivers

in Kagawa prefecture from January 2014 to August 2015 (Fig. 1A). The details are in supplementary

table S1. To explore the genetically-fixed gut length in medaka populations, we sampled 400

individuals from 40 wild-derived laboratory stocks that originated from wild populations and consisted

of seven genetic backgrounds (SJPN1-4, NJPN1, NJPN2, and KOR)[13]. Each stock is a descendant of

populations collected from each wild habitat in 1983[27] and has been maintained in independent tanks in

common environments (i.e., the same food and feeding times) at our outdoor breeding facility at The

University of Tokyo[13,28] (see fig. S9 in Katsumura et al. [2014])[28]. The sampled stocks were: SJPN1

(PO.NEH): Ichinoseki, Toyota, Mishima, Iwata, Sakura, Shingu; SJPN2 (Sanyo/Shikoku/Kinki):

Ayabe, Tanabe, Okayama, Takamatsu, Kudamatsu, Misho; SJPN3 (San-in): Kasumi, Tsuma, Tottori,

Hagi; SJPN4 (Northern and Southern Kyushu): Kusu, Arita, Hisayama, Fukue, Kazusa, Izumi, Hiwaki,

Nago; NJPN1 (derived NJPN): Kamikita, Yokote, Niigata, Kaga, Maizuru, Miyazu; NJPN2 (ancestral

NJPN): Amino, Kumihama, Toyooka, Kinosaki, Hamasaka; KOR (KOR/CHN): Yongcheon, Maegok,

Bugang, Sacheon, Shanghai. The subgroup names in parentheses are defined in our prior study[13]. We

isolated the gut (from the esophagus to the anus), fixed each one in 4% paraformaldehyde for 1 hour on

ice, and took photos of them laid out on glass slides to measure their lengths. Unfortunately, 32

medaka individuals died before this analysis, and 27 individuals were removed because of presenting

with an abnormal form due to aging (e.g., spondylosis). Finally, we obtained the standard and gut

lengths from photos of 341 individuals (table S4) using Image J software[29]. Statistical analyses were

performed using R software[30].

Detecting gut length changes using a Bayesian GLMM framework

The relationships between gut length and seasons, genetic background, *Plxnb3* upstream, *Ppp3r1*

genotypes, and gene expressions were modeled using a GLMM framework with the MCMC method in

the R package brms[31] in R v3.5.2. The gut length was modeled with a gamma error structure and log

link function. Fixed effects included the standard length and sex in each analysis, and the random

effect was subject identity and genetic background. The Markov chain Monte Carlo (MCMC)

conditions are described in each supplementary table.

Detecting seasonal methylated regions and filtering using Reactome and GO databases

We extracted genomic DNA from two male and two female guts per season (total 16 guts) from the

Kabe river medaka using the phenol/chloroform method described previously[13]. The quality and

quantity of genomic DNA were assessed using 0.5% agarose electrophoresis, a nanophotometer

(IMPLEN), and the Qubit BR dsDNA Assay Kit (Thermo Fisher Scientific). After adjusting the DNA

concentration to 40 ng/µl, 1.2 µg of DNA was sheared to target 150–200 bp by using two protocols of

400 bp and 200 bp implemented in the S220 Focused-ultrasonicator (Covaris). Sheared DNAs were

purified using the MinElute PCR Purification Kit (Qiagen) according to the manufacture's protocol,

and their quality and quantity were measured by the 4200 TapeStation system (Agilent Technologies)

and Qubit dsDNA BR Assay Kit (Thermo Fisher Scientific). A total of 500 ng of sheared DNAs was

input for methyl-CpG binding domain (MBD) enrichment using the EpiXplore Methylated DNA

Enrichment Kit (Clontech) according to the manufacturer's instruction. Subsequently, 30 ng of

methylated DNA was used to generate the NGS (Next-Generation Sequencing) library by using the

NEBNext ULTRA DNA Library Prep Kit for Illumina (New England Biolabs) and the NEBNext

Multiplex Oligos for Illumina (New England Biolabs) according to the manufacturers' instructions.

The quality and quantity of genomic DNA were assessed using the TapeStation D1000HS screen tape

(Agilent Technologies) and the Qubit dsDNA HS Assay Kit (Thermo Fisher Scientific). The

sequencing process was outsourced to Macrogen Japan in Kyoto and was performed by Illumina Hiseq

2500 with 51 bp single-end reads to obtain an average of 40 million reads per sample. The data have

been submitted to the DDBJ Sequence Read Archive (DRA) database under accession number:

DRA010581.

Our single-end reads were filtered using the FASTQ Quality Trimmer and Filter in FASTX-

Toolkit ver.0.0.13 (http://hannonlab.cshl.edu/fastx_toolkit/download.html) using the following options:

"-t 20 -l 30 -Q 33" and "-Q 33 -v -z -q 20 -p 80", respectively. The draft genome of the medaka

sequenced by the PacBio sequencer (Medaka-Hd-rR-pacbio_version2.2.4.fasta;

http://utgenome.org/medaka_v2/#!Assembly.md) was used to align the reads using BWA (Burrows-

Wheeler Aligner) backtrack 0.7.15-r1140[32] using the "-n 0.06 -k 3" option. After the mapping process,

the multi-mapped reads were removed using SAMtools v1.9[33] and the "-q 1 -F 4 -F 256 -F 2048"

option.

The detection of differentially methylated regions (DMRs) was performed using

MethylAction[34] with "fragsize=200, winsize=50" on R ver.3.4.4 in RStudio ver.1.0.136. First,

significant DMRs in one of the groups at least (Adjusted analysis of deviation [ANODEV] P-values

[Benjamini-Hochberg] <0.05) were explored by comparing between the four groups (14W and 14S,

15W and 15S). Next, the regions that were TRUE in the "frequent column", which the "frequent" state

corresponds to those where the methylation status of the samples is two-thirds or more agreement in

each group, were filtered. And then, we extracted regions with patterns that hypermethylation and

hypomethylation coincided with seasonal changes repeatedly, as seasonal varying methylated regions.

To determine if type I error rates is acceptable, we calculated a false discovery rate (FDR) for each

pattern of DMR among the groups by the MethylAction-implemented bootstrapping approach (2500

reps). We then annotated the DMRs (length distribution, overlap with open chromatin location

information from ATAC-seq data[35], and TE search using RepeatMasker[36]) using BEDTools[37] and

SeqMonk ver.1.48.0 (https://www.bioinformatics.babraham.ac.uk/projects/seqmonk/), and finally

listed genes within 2,000 bp upstream and downstream of the DMRs.

To identify the biological processes of genes uniquely associated with each seasonally-

correlated region, we performed enrichment analyses based on Reactome pathway, GO terms, and

human genome annotations using Enrichr program[38,39], without correcting for gene size variation.

Before this Enrichr analysis, we estimated medaka orthologous genes to human genes using biomart in Ensembl database and ORTHOSCOPE[40] because we used an information of gene function in human. Significant Reactome pathways (Adjusted *P*-value <0.05) and the top 10 GO terms for "biological processes" were used as the filter to list the candidate genes. Adding the gene list with the seasonally methylated region on their upstream, we visualized as a Venn diagram and identified the functional gene, *Plxnb3*, associated with seasonal plasticity (Fig. 2A). Finally, we identified a CpG island on upstream of this gene using MethPrimer[41] with the following default setting: GC >50%, O/E >0.6, and length >100 bp (Fig. 2B).

Analyses of whole genome bisulfite sequencing and targeted bisulfite sequencing

We used the same DNA as that used in MBD-seq. The whole genome bisulfite sequencing (WGBS) was outsourced to Macrogen Japan Inc. In brief, genomic DNA was bisulfite converted using the EZ DNA Methylation-Gold Kit (ZYMO RESEARCH). Sequencing libraries were prepared using the Accel-NGS™ Methyl-Seq DNA Library Kit (Swift BIOSCIENCE). Each library was sequenced using NovaSeq6000 (Illumina) to obtain raw data of more than 45 Gb paired-end (about 56 times more than medaka genome).

WGBS analysis was performed using programs: Bismark ver.0.23.0[42] and SeqMonk ver.1.48.0 (https://www.bioinformatics.babraham.ac.uk/projects/seqmonk/). The raw fastq reads (average 51.5 Gb/sample) were filtered using fastp[43] with the following option: "-q 20 -n 10 -f 15 -F 15 -t 15 -T 15 -l 20". The clean fastq reads (average 39.7 Gb/sample) were aligned to medaka reference genome (Medaka-Hd-rR-pacbio_version2.2.4.fasta) with Bismark using the following parameters: "--score_min L,0,-0.6". The duplicate reads were removed using "deduplicate_bismark" in Bismark. The methylation calls were performed using bismark_methylation_extractor and bismark2bedGraph command in Bismark. The methylation levels were determined for each CpG site within the regions detected by MBD-seq with SeqMonk using the "Difference quantification" function (as a percent of

methylated CpGs calls over total calls for each site) and retrieved using the "Annotated probe report"

function. Logistic regression filter on probes in MBD-seq detected peaks where summer vs winter had

a significance below 0.1 after Benjamimi and Hochberg correction with a minimum number of

observations of 10 in all samples. Also, the differences in methylation between season was visualized

by SeqMonk.

      For targeted bisulfite sequencing (tBS) analysis, we extracted genomic DNA from two male

and two female guts per season (total 16 guts) from the Ejiri and Shihodo rivers as well as above. The

genomic DNA was bisulfite converted and sequenced using MiSeq (Illumina) as described

previously[44,45]. The primer pairs used to amplify the target were: oPlexnb3_MR_Nxt_1F:

TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGTATATGTGATGTTTAGAGGATTGG;

oPlexnb3_MR_Nxt_1R:

GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAGTTTTTAAACTAACTCAAAATTCTAAC

C. The methylation level for each CpG site was calculated by QUMA[46] using the automated shell

scripts in Sekita et al. (2021) and the following condition to exclude low quality sequences: "Upper

limit of unconverted CpHs, 5; Lower limit of percent converted CpHs, 0.95; Upper limit of alignment

mismatches, 20; Lower limit of percent identity, 0.95." Statistical difference in methylation between

seasons on each CpG site was evaluated by GLMM using "lmerTest" R package. The methylation (the

number of methylated and unmethylated CpG) was modeled with a binomial error structure and logit

link function. Fixed effect included the season (winter or summer), and the random effect was

sampling year. The short read data of tBS and WGBS have been submitted to the DRA database under

the accession numbers: DRA015161 and DRA015162, respectively.

Finding the high-differentiated and gut-length-associated SNPs

For 341 gut-length-measured medaka, genomic DNAs were extracted from muscle using NucleoSpin

Tissue (Macherey-Nagel) according to the manufacturer's protocol. After the quality check using a

Nanophotometer (IMPLEN) and 0.5% agarose gel electrophoresis, six individuals were removed

because of low DNA concentrations. For 335 individuals, we arranged the original RAD-seq protocol[21]

adding an indexing PCR step to adjust the sample size and generated 14 RAD-seq libraries (24

individuals per library, 23 individuals in the last library) by the following method. The designed P1-

adaptor included 24 in-line barcodes which had two nucleotide differences in each, and the P1-adaptors

and Sbf I-HF (New England Biolabs)-digested DNAs for 90 min at 37˚C were ligated by T4 ligase

(New England Biolabs). After DNA pooling, DNAs were sonicated using an S220 Focused-

ultrasonicator (Covaris) to target 300 bp and purified using the GeneRead Size Selection Kit (Qiagen)

according to the manufacturers' protocols. The End-repair, A-tailing, and P2 adaptor ligation steps

were performed using the NEBNext Ultra DNA Library Prep Kit for Illumina (New England Biolabs).

After size selection using AMPure XP beads (Beckman Coulter), indexing PCR was performed using

Q5 (New England Biolabs) under the following conditions: initial denaturing step at 90˚C for 30 sec,

14 cycles of denaturation at 98˚C for 10 sec, annealing at 68˚C for 30 sec, extension at 72˚C for 20 sec,

and a final extension step at 72˚C for 5 min. The PCR products were purified with AMPure XP beads

and then were validated using the Qubit dsDNA HS Assay Kit (Thermo Fisher Scientific) for DNA

concentration, 4200 TapeStation (Agilent Technologies) for fragment length, and Miseq (Illumina) for

library quality. Finally, RAD-seq data were generated using three lanes of Hiseq 2500 (Illumina) with

51 bp single-end reads settings conducted by Macrogen Japan. The data have been submitted to the

DRA database under accession number: DRA010605.

Our single-end reads were filtered by Cutadapt ver1.12[47] using the following options: "-m 50 -

e 0.2," and were demultiplexed by process_radtags (v1.44) implemented in Stacks[48] using the

following options: "-c -r -t 44 -q -s 0 --barcode_dist_1 2." After quality filtering and check, one

individual (from the Kazusa population) was removed because of low reads. The draft genome of the

medaka sequenced by the PacBio sequencer (Medaka-Hd-rR-pacbio_version2.2.4.fasta;

http://utgenome.org/medaka_v2/#!Assembly.md) was used to align the reads using BWA backtrack

0.7.15-r1140[32] using the "-n 0.06 -k 3" option. After the mapping process, the multi-mapped reads

were removed using SAMtools v1.9[33] and the "-q 1 -F 4 -F 256 -F 2048" option. SNP call was

performed by Stacks pipeline: pstacks -m 2 --model_type snp --alpha 0.05; cstacks -g -n 1; sstacks -g;

rxstacks --lnl_filter --lnl_lim -10 --conf_filter --conf_lim 0.75 --prune_haplo --model_type bounded --

bound_low 0 --bound_high 0.1; cstacks -g -n 1; sstacks -g; populations -r 0.5 -p 7 -m 6 -f p_value -a

0.0 --p_value_cutoff 0.1 --lnl_lim -10. Assigned medaka populations into seven genetic groups based

on our prior study[13], genotype imputation was performed by Beagle (v4.1) using GT format[49]. Finally,

data set was generated using VCFtools[50] to merge the seven genotype-imputed data, and included

63,265 SNPs.

To detect high-differentiated and gut-length-associated genes, we calculated Weir and

Cockerham's F_{ST} and the degree of association with the gut length using VCFtools[50] and PLINK

v1.90b3.46[51] (https://www.cog-genomics.org/plink/1.9/) using the SNP data. Although the gut length

was highly correlated with the population structure of medaka (Fig. 1C), we did not control the

structure information because this reduced the power of the analysis and could lead to false

negatives[52]. To avoid the false positives without controlling the population structure as much as

possible, we designed the sampling strategy to choose multiple subgroups from each group and to use

balancing samples across subgroup populations to homogenize allele frequencies[52]. Focusing on SNPs

with a minor allele frequency of more than 0.05, we extracted the top 10 SNPs above the global F_{ST}

value (weighted F_{ST}) and with the highest $-\log_{10}$ P-value from 22,235 SNPs. Association of gut length

and SNPs were assessed by permutation test with 10,000,000 times resampling, implemented in

PLINK v1.90b3.46[51].

Phylogenetic tree reconstruction using RAD-seq data

We used 22,235 SNPs data, which is the same data above association analysis, for reconstruction of the

phylogenetic tree. The data was analyzed via the IQ-TREE program[53] to reconstruct a maximum

likelihood tree with general time-reversible (GTR) + ascertainment bias correction model and 1,000

ultrafast-bootstrappings[54]. Then, we used the FigTree ver.1.4.4 program

(http://tree.bio.ed.ac.uk/software/figtree/) to visualize the phylogenetic tree.

Generating a genome-editing medaka by the CRISPR/Cas9 system

The d-rR medaka line was used to generate mutants deleted from the seasonal methylation region. Fish

were maintained at 28°C under a 14-h light/10-h dark cycle. We collected the d-rR eggs every

morning. Microinjection was carried out in medaka embryos at the one-cell stage using FemtoJet

(Eppendorf) and InjectMan NI 2 (Eppendorf). Cas9 protein (500 ng/μL), tracrRNA (200 ng/μL), two

crRNAs (each 200 ng/μL), and saturated phenol red solution were mixed in a 4:2:2:1 ratio, and

injected into the embryos. The crRNAs were designed to delete the seasonal methylation region of 334

bp using CRISPRdirect[55], which identified the specific regions on the medaka genome, and their

crRNA sequences were: oPlxnB3upCGI_F1,

GAUGCUGUUGCCUGGCCAUAguuuuagagcuaugcuguuuug; oPlxnB3upCGI_R1,

GAUGACACAGUGGAGCAUCGguuuuagagcuaugcuguuuug. The crRNAs, tracrRNA, and

recombinant Cas9 protein were obtained from FASMAC. After injection, the embryos (F0) were

maintained at 28°C using an air incubator. After hatching and maturing, the injected medaka were

intercrossed for the identification of the germline founders with mutations in each target locus by

genotyping their fertile eggs. The genotyping was performed using PCR and 1.5% agarose gel

electrophoresis using EX Taq Hot Start Version (TaKaRa Bio) according to the manufacturer's

protocol and two designed primers: oPlxnb3CpG1F2, 5'-TTGCATCTGGCTTTGCATGAATC-3';

oPlxnb3CpG1R2, 5'-ATAACATCACCATGGCAACAACG-3'. The germline founders (F1) were

outcrossed and generated their off-springs (F2). Heterozygous mutation carriers in the F2 generation

were identified by above PCR-based genotyping using their fin-extracted DNAs. Finally, sixty F3

individuals generated by heterozygous mating of the heterozygous mutation carriers in the F2 were

used for subsequent analyses.

Analysis of *Plxnb3* upstream-deleted medaka

To examine a phenotype of *Plxnb3* upstream-deleted (Δ*upPlxnb3*) medaka, we conducted three tests:

segregation of genotypes, a comparison of gut and body length, and an immunohistological

comparison. For these tests, we performed heterozygous mating using the F2 generation and obtained

their fertile eggs. We bred the medaka larvae in a tank (W52×D21×H23cm) and raised them until 2

weeks after hatching, divided the juveniles into groups of four to six individuals, and transferred them

to 2 L tanks to raise them for an average of 103 days after hatching. Each genotype was determined

using the PCR-based method described above, and the segregation ratio was evaluated using the chi-

squared test. The significant differences in gut lengths between genotypes were detected using a

Bayesian GLM, performed with the brms R package with the default MCMC setting on RStudio

ver.1.3. The gut length was modeled with a gamma error structure and log link function. Fixed effects

included the standard length and sex in each analysis. For immunohistological analysis, 5 μm serial

tissue sections from wild types and mutants (n = 5 each) were prepared using our previously described

method[56]. The sections were incubated with a mouse monoclonal anti-phosphorylated neurofilament H

(BioLegend, #smi-31, 1:1000) of which cross-reactivity for medaka had been confirmed in a previous

study[57]. Quantitative analysis of an immunohistological-staining signal was performed using ImageJ

software[29]. We obtained an average of six images per individual at a 100× objective using light

microscopy (BX63, Olympus) equipped with a digital camera (DP74, Olympus). We then split each

image into red, green, and blue using the color channels option and subtracted green from the red

image with the image calculator. Setting the same threshold, we measured the proportion of the stained

area for one intestinal villus of each individual. After standardizing the percentage of the stained area

in each experiment, we evaluated the difference of the immunohistological-staining signals between

the wild types and the mutants using the Wilcoxon rank-sum test ($P < 0.05$).

RT-qPCR

We performed reverse transcription-quantitative PCR (RT-qPCR) analyses to evaluate the gene

expressions of the *Plxnb3* and *Ppp3r1* genes. We sampled three individuals from each season in the

Kabe river (12 individuals totally) for the *Plxnb3* gene expression, and four individuals from four

geographical populations (Kamikita and Yokote from NJPN1, Amino from NJPN2, Ichinoseki and

Iwata from SJPN1, and Izumi from SJPN4) for the *Plxnb3* and/or the *Ppp3r1* gene expression. The

guts were isolated from ethanol-fixed Kabe river samples stored at -20˚C and wild-derived laboratory

stocks in winter (29 Jan. 2018). The gut was homogenized using a BioMasher (Nippi), and total RNA

was isolated from each sample using a combination of the TRIzol (Invitrogen) and NucleoSpin RNA

(Macherey-Nagel).

cDNA, which was used as a template for qPCR, was synthesized using a PrimeScript RT

reagent kit with gDNA Eraser (TaKaRa Bio) according to the manufacturer's protocol. *Ribosomal*

protein L7 (*Rpl7*) was used as an internal reference gene because medaka *Rpl7* was expressed with the

less variance among the same tissue samples, in different tissues and stages of development[58], and

between winter and summer[59]. The negative control was treated with RNase free water and RT minus

products. The reaction mixture consisted of 2 μl of cDNA template, 10 μl of TB Green® Premix Ex

Taq™ (Tli RNaseH Plus) (TaKaRa Bio), 0.4 μl of forward and reverse primers (10 μM), and 7.2 μl of

RNase-free water. The reaction procedure in a LightCycler® 480 Real-Time PCR System (Roche

Diagnostic) was: 95˚C, 30 sec; 95˚C, 5 sec + 60˚C, 30 sec (for amplification, 40 cycles); 95˚C, 5 sec +

60˚C, 60 sec + 97˚C, 0.11˚C/sec (for the melting curve); 40˚C, 30 sec (for cooling). The primer pairs

were: oPlxnb3ex32F1:TGGTGAAGAGCAGTGAAGATCC;

oPlxnb3ex33R1:TAGTGTTCCCTTCATGGACAGC for *Plxnb3* (PCR efficiency: 105%),

oCANB1ex4F1:TAAACATGAAGGGAAGGCTGGAC;

oCANB1ex3R1:ATTCGCCTTCAGGATCTACGAC, for *Ppp3r1* (PCR efficiency: 95%), and

oRPL7ex3-4F1:ATCCGAGGTATCAACGGAGTC; oRPL7ex5R1:TGCCGTAGCCACGTTTGTAG

for *Rpl7* (PCR efficiency: 95%). The specificity of all PCR primer pairs was confirmed as a single

peak in the melting curve analysis. Two technical replicates were run for the RT-qPCR analyses, and

the relative expressions of each sample were calculated using the ΔΔCt method with the LightCycler®

(Roche Diagnostic). The statistical tests were performed using R ver.4.1.1 and RStudio ver.1.4.1717.

Outdoor breeding experiment in artificial environment

To test whether the food source affects the gut length, we transported the wild medaka from the Kabe

river into the outdoor breeding facility on the Kashiwa campus, The University of Tokyo, in August

2015. Then in September 2016 and February 2017, we sampled and compared the gut lengths of bred

and wild medakas in the outdoor breeding facility and the Kabe river, respectively. Their gut lengths

were measured according to the method described above.

Common garden experiment

To examine whether NJPN1 medaka show a shorter gut in winter than in summer, we sampled and

measured the medaka gut lengths on 3 February and 30 August 2021 using available wild-derived

laboratory stocks. The sampled stocks were: NJPN1: Yokote (n = 10, 10), Miyazu (n = 10, 10), Kaga

(n = 7, 10); NJPN2: Kumihama (n = 9, 10), Amino (n = 10, 10), Toyooka (n = 11, 8); SJPNs as

control: Ichonoseki (n = 8, 10), Okayama (n = 10, 5), Mishima (n = 10, 9). The significant differences

between seasons were detected by the Steel-Dwass test implemented in an NSM3 R package. We used

the Bayesian GLMM to detect the gut length variations using the brms R package, as described above.

Fixed effects included the standard length, sex, season (winter and summer), genetic background, and

the interaction between season and genetic background. Random effects were subject identity and the

sampling year. Four MCMC chains were run for 2,000 samples, with a 1,000 sample burn-in, by using

every sample. This gave a total of 4,000 samples used. R-hat is the Germany-Rubin convergence

statistic for estimating the degree of convergence of a random Markov chain and indicates insufficient

convergence at values greater than 1.1.

Principal component analysis

To test whether or not the population structure had changed in the Kabe river between winter and

summer, we performed a principal component analysis (PCA) using the SNPRelate program[60] in R

ver.3.2.2. A total of 2,167,650 bi-allelic SNPs were generated from the bam files used to detect the

methylation regions by MBD-seq. The procedure was such that the SNPs were extracted using

SAMtools v1.9[33] mpileup and filtered vcfutils.pl with varFilter -d 6 option, and then only bi-allelic

sites were extracted using VCFtools[50]. The PCA graph was drawn using ggplot2[61].

Calculation of d_A distances between seasons in each river based on mtDNA D-loop sequences

The genetic distance matrix was obtained by the calculation made from the number of base differences

per site and from the estimation of the net average between groups of mtDNA D-loop sequences using

the MEGA7[62]. The D-loop sequences were generated by the method previously described[63]. These

nucleotide sequences were deposited into the international DNA database DDBJ/EMBL/GenBank

(accession nos. LC719344 – LC719462).

Relationships between the number of CpG sites on the upstream of *Plxnb3* and the mutation in

the *Ppp3r1* region

To examine the genetic variation on the upstream of *Plxnb3* among geographic populations, we

performed PCR direct sequencing as described previously[28]. The two primers were

o*Plxnb3*CpG1F1:GTTCATTTAAGGGAGGAACCAAAGG and

o*Plxnb3*CpG1R1:CTCCACTGTGTCATCAAAAGAAGC. Sequences were determined and aligned across 40 populations, then the CpG sites on these sequences were counted using MEGA X software[64]. Note that the sequences obtained from the Kaga, Shingu, and Shanghai local populations were excluded from the analysis because their CpG sites of upstream sequences or *Ppp3r1* mutations existed as polymorphisms. Thus, the regression line between the number of CpG sites and gut lengths was calculated using the populations with homozygotes of the C allele on *Ppp3r1*. A phylogenetic tree was reconstructed to estimate the number of CpG sites on ancestral upstream sequences of *Plxnb3* using the maximum likelihood method under the GTR model implemented in MEGA X[64]. Those numbers were plotted on the medaka genetic group tree obtained from the previous study[13], and the allele frequencies of the *Plxnb3* upstream sequence and *Ppp3r1* mutations in each group were calculated as the number of alleles divided by the number of chromosomes. Note that data obtained from the Yongcheon and Shanghai populations were excluded from these analyses because those populations might have experienced artificial gene-flow via the strain maintenance process mentioned in our prior study[13].

To test whether the dispersion of the gut length was large, according to the dispersion of the air temperature of each region, we calculated the coefficient of variation (CV) of the gut length and standard deviation of the average annual temperature for each group using the Global Solar Atlas database (https://globalsolaratlas.info). We plotted the standard deviation of the average annual temperature on the X-axis and the mean CV of the gut length on the Y-axis and performed a Pearson's product-moment correlation test in R and RStudio.

Acknowledgments

We thank Mrs. Shizuko Chiba, Mrs. Sumiko Tomizuka, Dr. Atsuko Shimada, and Prof. Emeritus Akihiro Shima (The University of Tokyo) for maintaining the medaka stocks from wild populations. We also thank Robert E. Brandt, Founder, CEO, and CME, of MedEd Japan, for editing and formatting the manuscript.

This study was supported in part by grants-in-aid for Young Scientists (B) no. JP16K21352,

Early-Career Scientists no. JP19K16201, Scientific Research on Innovative Areas no. JP19H05737,

Scientific Research (A) no. JP17H01453, and Scientific Research (B) no. 17H03738. T.K. was also

supported by a grant-in-aid from JSPS Research Fellowships no. JP16J07227.

Authors' contributions

T.K., H.Takeshima, Y.H., H.Takeuchi, and H.O. designed the experiments; T.K., S.S., K.Y., S.O.,

T.G., S.T., K.F., T.N., T.I., Y.H., and H.Takeshima performed the sampling and experiments; T.K.,

S.S., and K.Y. analyzed the data; T.K., T.G., T.N., Y.S., H.M., M.O., H.Takeuchi, and H.O. interpreted

the results of the experiments; T.K. and H.O. wrote the manuscript. All authors have read and

approved the final manuscript.

References

1.   Gilbert, S. F., Bosch, T. C. G. & Ledón-Rettig, C. Eco-Evo-Devo: developmental symbiosis and

developmental plasticity as evolutionary agents. *Nat. Rev. Genet.* **16**, 611–622 (2015).

2.   Levis, N. A., Isdaner, A. J. & Pfennig, D. W. Morphological novelty emerges from pre-existing

phenotypic plasticity. *Nat. Ecol. Evol.* **2**, 1289–1297 (2018).

3.   Levis, N. A. & Pfennig, D. W. Evaluating "Plasticity-First" Evolution in Nature: Key Criteria and

Empirical Approaches. *Trends Ecol. Evol.* **31**, 563–574 (2016).

4.   Laland, K. *et al.* Does evolutionary theory need a rethink? *Nature* **514**, 161–164 (2014).

5.   Baerwald, M. R. *et al.* Migration-related phenotypic divergence is associated with epigenetic

modifications in rainbow trout. *Mol. Ecol.* **25**, 1785–1800 (2016).

6.   Kronholm, I. & Collins, S. Epigenetic mutations can both help and hinder adaptive evolution.

Mol. Ecol. **25**, 1856–1868 (2016).

7.   Wang, X. *et al.* DNA methylation mediates differentiation in thermal responses of Pacific oyster (Crassostrea gigas) derived from different tidal levels. *Heredity (Edinb)* **126**, 10–22 (2021).

8.   Zaldúa, N. & Naya, D. E. Digestive flexibility during fasting in fish: a review. *Comp. Biochem. Physiol. A Mol. Integr. Physiol.* **169**, 7–14 (2014).

9.   Zandonà, E., Auer, S. K., Kilham, S. S. & Reznick, D. N. Contrasting Population and Diet Influences on Gut Length of an Omnivorous Tropical Fish, the Trinidadian Guppy (Poecilia reticulata). *PLoS One* **10**, e0136079 (2015).

10. Starck, J. M. Shaping up: how vertebrates adjust their digestive system to changing environmental conditions. *Anim. Biol. Leiden Neth.* **53**, 245–257 (2003).

11. Suzuki, Y., Miyake, T. & Yamahira, K. An acquisition trade-off with fast growth in a fish, the medaka Oryzias latipes: why do low-latitude ectotherms grow more slowly? *Evol. Ecol.* **24**, 749–759 (2010).

12. Yamahira, K. & Nishida, T. Latitudinal variation in axial patterning of the medaka (Actinopterygii: Adrianichthyidae): Jordan's rule is substantiated by genetic variation in abdominal vertebral number. *Biol. J. Linn. Soc. Lond.* **96**, 856−866 (2009).

13. Katsumura, T., Oda, S., Mitani, H. & Oota, H. Medaka Population Genome Structure and Demographic History Described via Genotyping-by-Sequencing. *G3 (Bethesda)* **9**, 217–228 (2019).

14. Hilgers, L. & Schwarzer, J. The untapped potential of medaka and its wild relatives. *Elife* **8**, e46994 (2019).

15. Kirchmaier, S., Naruse, K., Wittbrodt, J. & Loosli, F. The genomic and genetic toolbox of the teleost medaka (Oryzias latipes). *Genetics* **199**, 905–918 (2015).

16. Leaf, R. T., Jiao, Y., Murphy, B. R. & Kramer, J. Life-History Characteristics of Japanese Medaka Oryzias latipes. *Copeia* **2011**, 559–565 (2011).

17. Edeline, E., Terao, O. & Naruse, K. Empirical evidence for competition-driven semelparity in

wild medaka. *Popul. Ecol.* **58**, 371–383 (2016).

18. Wagner, C. E., McIntyre, P. B., Buels, K. S., Gilbert, D. M. & Michel, E. Diet predicts intestine

length in Lake Tanganyika's cichlid fishes. *Funct. Ecol.* **23**, 1122–1131 (2009).

19. Artemov, A. V. *et al.* Genome-Wide DNA Methylation Profiling Reveals Epigenetic Adaptation

of Stickleback to Marine and Freshwater Conditions. *Mol. Biol. Evol.* **34**, 2203–2213 (2017).

20. Schneider, C., O'Leary, C. E. & Locksley, R. M. Regulation of immune responses by tuft cells.

Nat. Rev. Immunol. **19**, 584–593 (2019).

21. Baird, N. A. *et al.* Rapid SNP discovery and genetic mapping using sequenced RAD markers.

PLoS One **3**, e3376 (2008).

22. Graef, I. A. *et al.* Neurotrophins and netrins require calcineurin/NFAT signaling to stimulate

outgrowth of embryonic axons. *Cell* **113**, 657–670 (2003).

23. Fujita, M. *et al.* Calcineurin B1 Deficiency in Glial Cells Induces Mucosal Degeneration and

Inflammation in Mouse Small Intestine. *Biol. Pharm. Bull.* **41**, 786–796 (2018).

24. Héberlé, É. & Bardet, A. F. Sensitivity of transcription factors to DNA methylation. *Essays*

Biochem. **63**, 727–741 (2019).

25. Nei, M. *Mutation-Driven Evolution. OUP Oxford, New York*, (2013).

26. Latorre, R., Sternini, C., De Giorgio, R. & Greenwood-Van Meerveld, B. Enteroendocrine cells: a

review of their role in brain-gut communication. *Neurogastroenterol. Motil.* **28**, 620–630 (2016).

27. Shima A. First listing of wild stocks of the Medaka Oryzias latipes currently kept by Zoological

Institute, Faculty of Science, University of Tokyo. *J. Fac. Sci. Univ. Tokyo. Sec. IV* **16**, 27–35

(1985).

28. Katsumura, T. *et al.* Natural allelic variations of xenobiotic-metabolizing enzymes affect sexual

dimorphism in Oryzias latipes. *Proc. Biol. Sci.* **281**, 20142259 (2014).

29. Schneider, C. A., Rasband, W. S. & Eliceiri, K. W. NIH Image to ImageJ: 25 years of image analysis. *Nat. Methods* **9**, 671–675 (2012).

30. R Core Team. R: A Language and Environment for Statistical Computing. Preprint at https://www.R-project.org/ (2018).

31. Bürkner, P.-C. Advanced Bayesian Multilevel Modeling with the R Package brms. *R J.* **10**, 395–411 (2018).

32. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* **25**, 1754–1760 (2009).

33. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079 (2009).

34. Bhasin, J. M., Hu, B. & Ting, A. H. MethylAction: detecting differentially methylated regions that distinguish biological subtypes. *Nucleic Acids Res.* **44**, 106–116 (2016).

35. Li, Y. *et al.* Dynamic transcriptional and chromatin accessibility landscape of medaka embryogenesis. *Genome Res.* **30**, 924–937 (2020).

36. *Smit, A. F. A., Hubley, R. & Green, P. RepeatMasker Open-4.0.* Preprint at http://repeatmasker.org (2013–2015).

37. Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics* **26**, 841–842 (2010).

38. Kuleshov, M. V. *et al.* Enrichr: a comprehensive gene set enrichment analysis web server 2016 update. *Nucleic Acids Res.* **44**, W90-W97 (2016).

39. Chen, E. Y. *et al.* Enrichr: interactive and collaborative HTML5 gene list enrichment analysis tool. *BMC Bioinformatics* **14**, 128 (2013).

40. Inoue, J. & Satoh, N. ORTHOSCOPE: An Automatic Web Tool for Phylogenetically Inferring Bilaterian Orthogroups with User-Selected Taxa. *Mol. Biol. Evol.* **36**, 621–631 (2019).

41. Li, L.-C. & Dahiya, R. MethPrimer: designing primers for methylation PCRs. *Bioinformatics* **18**, 1427–1431 (2002).

42. Krueger, F. & Andrews, S. R. Bismark: a flexible aligner and methylation caller for Bisulfite-Seq applications. *Bioinformatics* **27**, 1571–1572 (2011).

43. Chen, S., Zhou, Y., Chen, Y. & Gu, J. fastp: an ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics* **34**, i884–i890 (2018).

44. Kawasaki, Y. *et al.* A Novel method for the simultaneous identification of methylcytosine and hydroxymethylcytosine at a single base resolution. *Nucleic Acids Res.* **45**, e24 (2017).

45. Sekita, Y. *et al.* AKT signaling is associated with epigenetic reprogramming via the upregulation of TET and its cofactor, alpha-ketoglutarate during iPSC generation. *Stem Cell Res. Ther.* **12**, 510 (2021).

46. Kumaki, Y., Oda, M. & Okano, M. QUMA: quantification tool for methylation analysis. *Nucleic Acids Res.* **36**, W170-W175 (2008).

47. Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet J.* **17**, 10–12 (2011).

48. Catchen, J., Hohenlohe, P. A., Bassham, S., Amores, A. & Cresko, W. A. Stacks: an analysis tool set for population genomics. *Mol. Ecol.* **22**, 3124–3140 (2013).

49. Browning, S. R. & Browning, B. L. Rapid and accurate haplotype phasing and missing-data inference for whole-genome association studies by use of localized haplotype clustering. *Am. J. Hum. Genet.* **81**, 1084–1097 (2007).

50. Danecek, P. *et al.* The variant call format and VCFtools. *Bioinformatics* **27**, 2156–2158 (2011).

51. Chang, C. C. *et al.* Second-generation PLINK: rising to the challenge of larger and richer datasets. *Gigascience* **4**, 7 (2015).

52. Brachi, B., Morris, G. P. & Borevitz, J. O. Genome-wide association studies in plants: the missing heritability is in the field. *Genome Biol.* **12**, 232 (2011).

53. Nguyen, L.-T., Schmidt, H. A., von Haeseler, A. & Minh, B. Q. IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol. Biol. Evol.* **32**, 268–274 (2015).

54. Hoang, D. T., Chernomor, O., von Haeseler, A., Minh, B. Q. & Vinh, L. S. UFBoot2: Improving the Ultrafast Bootstrap Approximation. *Mol. Biol. Evol.* **35**, 518–522 (2018).

55. Naito, Y., Hino, K., Bono, H. & Ui-Tei, K. CRISPRdirect: software for designing CRISPR/Cas guide RNA with reduced off-target sites. *Bioinformatics* **31**, 1120–1123 (2015).

56. Nagata, K. *et al.* In vivo 3D analysis of systemic effects after local heavy-ion beam irradiation in an animal model. *Sci. Rep.* **6**, 28691 (2016).

57. Uemura, N. *et al.* Viable neuronopathic Gaucher disease model in Medaka (Oryzias latipes) displays axonal accumulation of alpha-synuclein. *PLoS Genet.* **11**, e1005065 (2015).

58. Zhang, Z. & Hu, J. Development and validation of endogenous reference genes for expression profiling of medaka (Oryzias latipes) exposed to endocrine disrupting chemicals by quantitative real-time RT-PCR. *Toxicol. Sci.* **95**, 356–368 (2007).

59. Shimmura, T. *et al.* Dynamic plasticity in phototransduction regulates seasonal changes in color perception. *Nat. Commun.* **8**, 412 (2017).

60. Zheng, X. *et al.* A high-performance computing toolset for relatedness and principal component analysis of SNP data. *Bioinformatics* **28**, 3326–3328 (2012).

61. Wickham, H. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag, New York https://ggplot2.tidyverse.org (2016).

62. Kumar, S., Stecher, G. & Tamura, K. MEGA7: Molecular Evolutionary Genetics Analysis Version 7.0 for Bigger Datasets. *Mol. Biol. Evol.* **33**, 1870–1874 (2016).

63. Katsumura, T. *et al.* Genetic differentiation among local populations of medaka fish (Oryzias latipes) evaluated through grid- and deme-based sampling. *Gene* **443**, 170–177 (2009).

1   64. Kumar, S., Stecher, G., Li, M., Knyaz, C. & Tamura, K. MEGA X: Molecular Evolutionary

2   Genetics Analysis across Computing Platforms. *Mol. Biol. Evol.* **35**, 1547–1549 (2018).