Examining the boundary sharpness coefficient as an index of cortical microstructure and its relationship with age and sex in autism spectrum disorder

Authors: Emily Olafson, BSc[1,2*], Saashi Bedford, MSc[1,3], Gabriel A. Devenyi, PhD[1,4], Raihaan Patel, BSc[1,5], Stephanie Tullo, PhD[1,3], Min Tae M. Park, PhD[6], Evdokia Anagnostou, PhD[7] , Simon Baron-Cohen, PhD[8] , Edward T. Bullmore, PhD[9] , Lindsay R. Chura, PhD[8], Michael C. Craig, PhD[10,12], Christine Ecker, PhD[,10,13] , Dorothea L. Floris, PhD[24,25] , Rosemary J. Holt, PhD[8] , Rhoshel Lenroot, PhD[15] , Jason P. Lerch, PhD[17,18], Michael V. Lombardo, PhD[8,19], Declan G. M. Murphy, PhD[10], Armin Raznahan, MD/PhD[11], Amber N. V. Ruigrok, PhD[8] , Michael D. Spencer, MD/PhD[8] , John Suckling, PhD[9] , Margot J. Taylor, PhD[20] , MRC AIMS Consortium[#], Meng-Chuan Lai, PhD[8,16,21,22,23], M. Mallar Chakravarty, PhD[1,3,4,5*]

1. Cerebral Imaging Centre, Douglas Mental Health University Institute, Montreal, Canada.

2. Department of Neuroscience, Weill Cornell Graduate School of Medical Sciences, New York City, USA.

3. Integrated Program in Neuroscience, McGill University, Montreal, Canada.

4. Department of Psychiatry, McGill University, Montreal, Canada.

5. Department of Biological and Biomedical Engineering, McGill University, Montreal, Canada

6. Department of Psychiatry, Schulich School of Medicine and Dentistry, Western University, London, ON, Canada

7. Holland Bloorview Kids Rehabilitation Hospital, Toronto, Canada, and Department of pediatrics, University of Toronto, ON, Canada

8. Autism Research Centre, Department of Psychiatry, University of Cambridge, Cambridge, UK.

9. Brain Mapping Unit, Department of Psychiatry, University of Cambridge, Cambridge, UK.

10. Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK.

11. Developmental Neurogenomics Unit, Human Genetics Branch, National Institute of Mental Health, Bethesda, MD, USA.

12. National Autism Unit, Bethlem Royal Hospital, London, UK.

13. Department of Child and Adolescent Psychiatry, Psychosomatics, and Psychotherapy, Goethe University, Frankfurt am Main, Germany.

14. Hassenfeld Children's Hospital at NYU Langone Department of Child and Adolescent Psychiatry, Child Study Center, New York City, New York, USA.

15. Department of Psychiatry, University of New South Wales, Sydney, NSW, Australia.

16. Program in Neurosciences and Mental Health, The Hospital for Sick Children, Toronto, Canada.

17. Department of Medical Biophysics, The University of Toronto, Toronto, Canada.

18. Wellcome Centre for Integrative Neuroimaging, University of Oxford, Oxford, UK.

19. Laboratory for Autism and Neurodevelopmental Disorders, Center for Neuroscience and Cognitive Systems @UniTn, Istituto Italiano di Tecnologia, Corso Bettini, 31, 38068 Rovereto, Italy

20. Diagnostic Imaging, The Hospital for Sick Children, Toronto, Canada.

21. The Margaret and Wallace McCain Centre for Child, Youth & Family Mental Health and Campbell Family Mental Health Research Institute, Centre for Addiction and Mental Health, Toronto, Canada.

22. Department of Psychiatry, University of Toronto, Toronto, Canada.

23. Department of Psychiatry, National Taiwan University Hospital and College of Medicine, Taipei, Taiwan

24. Donders Center for Brain, Cognition and Behavior, Radboud University Nijmegen, Nijmegen, The Netherlands

25. Department for Cognitive Neuroscience, Radboud University Medical Center Nijmegen, Nijmegen, The Netherlands

*Corresponding Authors:

Emily Olafson; Cerebral Imaging Centre,

Douglas Mental Health University Institute,

Verdun, Quebec, Canada. E-mail: olafsonemily@gmail.com

M. Mallar Chakravarty; Cerebral Imaging Centre,

 Douglas Mental Health University Institute,

Verdun, Quebec, Canada. E-mail: mallar@cobralab.ca

Acknowledgements:

*This study was performed by EO while at the Cerebral Imaging Centre, Douglas Mental Health University Institute, Montreal, Canada. E.O. is supported by the Natural Science and Engineering Research Council of Canada (NSERC) and the Fonds de la recherches en santé du Québec (FRQS).

The Cambridge Family Study of Autism was funded by a Clinician Scientist Fellowship from the MRC (G0701919) to MDS.

This study was funded by the MRC UK as the Autism Imaging Multicentre Study (AIMS), grant number GO 400061 to PIs Murphy, Bullmore, Baron-Cohen.

The Toronto cohorts were funded by the Canadian Institutes of Health Research (CIHR) MOP-106582, MOP119541 and MOP-142379 to Taylor and Anagnostou

GAD is supported in part by funding provided by Brain Canada, in partnership with Health Canada, for the Canadian Open Neuroscience Platform initiative.

M.V.L. is supported by a European Research Council (ERC) Starting Grant (ERC-2017-STG; AUTISMS; no. 755816)

A.R. is supported by the intramural program of the National Institutes of Mental Health (Clinical trial NCT00001246, clinicaltrials.gov; NIH Annual Report Number, 1ZIAMH002949-03).

M.M.C is supported by Canadian Institutes of Health Research (CIHR), Natural Science and Engineering Research Council of Canada (NSERC), Fonds de la recherches en santé du Québec

Abstract

Autism spectrum disorder (ASD) is associated with atypical brain development. However, the phenotype of regionally specific increased cortical thickness observed in ASD may be driven by several independent biological processes that influence the gray/white matter boundary, such as synaptic pruning, myelination, or atypical migration. Here, we propose to use the boundary sharpness coefficient (BSC), a proxy for alterations in microstructure at the cortical gray/white matter boundary, to investigate brain differences in individuals with ASD, including factors that may influence ASD-related heterogeneity (age, sex, and intelligence quotient). Using a vertex-based meta-analysis and a large multi-center magnetic resonance structural imaging (MRI) dataset, with a total of 1136 individuals, 415 with ASD (112 female; 303 male) and 721 controls (283 female; 438 male), we observed that individuals with ASD had significantly greater BSC in the bilateral superior temporal gyrus and left inferior frontal gyrus indicating an abrupt transition (high contrast) between white matter and cortical intensities. Increases were observed in different brain regions in males and females, with larger effect sizes in females. Individuals with ASD under 18 had significantly greater BSC in the bilateral superior temporal gyrus and right postcentral gyrus; individuals with ASD over 18 had significantly increased BSC in the bilateral precuneus and superior temporal gyrus. BSC correlated with ADOS-2 CSS in individuals with ASD in the right medial temporal pole. Importantly, there was a significant spatial overlap between maps of the effect of diagnosis on BSC when compared to cortical thickness. These results invite studies to use BSC as a possible new measure of cortical development in ASD and to further examine the microstructural underpinnings of BSC-related differences and their impact on measures of cortical morphology.

Introduction

Neuroimaging studies of autism spectrum disorder (ASD) have repeatedly observed early increases in cortical thickness (1–5) and altered structural and functional connectivity in individuals with ASD, particularly early in life (6–8). The early developmental period, which is coincident with ASD-onset, is a particularly sensitive time with respect to neuronal migration and cortical myelination. Indeed, neuronal migration disruptions from the ventricular and subventricular zones during early neocortex development have been observed in ASD (9–11), potentially leading to the presence of supernumerary neurons proximal to the gray/white matter boundary. Additionally, maturational abnormalities of intracortical myelin, a key cortical maturational feature (12,13), have been associated with ASD (14–16). This developmental variation may underlie measures seeking to capture so-called 'blurring' at the interface of the cortex and the underlying superficial white matter, as measured using structural magnetic resonance imaging (MRI) (17–20). Despite advances in ASD research generated by studying the ratio of the tissue intensities between these two compartments, there are limitations that may complicate the interpretation of this measure as an index of cortical microstructure. Firstly, the ratio of intensities between the tissue classes is dependent on the placement of the boundary that separates them. However, the boundary placement may be influenced by signal ambiguity that arises from a less defined gray/white matter boundary, thereby potentially confounding the measurement of interest. Further, histological studies have shown that more convex regions such as gyral crowns are more myelinated in deeper cortical layers compared to concave regions, such as sulcal folds (21,22); thus measures of cortical microstructure are negatively correlated with cortical curvature, as T1-relaxation times are observed to be higher in gyral crowns and lower in sulcal folds (Sereno et al., 2013; Waenert et al., 2016). In ASD,

microstructural measurements obtained using the standard ratio of tissue intensities may be confounded by the well-characterized group differences in gyrification and curvature (23,24).

Here, we propose to partly overcome these limitations using our newly-developed metric, boundary sharpness coefficient (BSC), defined as the growth rate parameter of a sigmoid curve fit to the cortical intensity profile running perpendicular to the boundary surface (inspired by (25)). Using this new measure that potentially reflects perturbations to neuronal migration and/or intracortical myelination, we performed a large-scale analysis using images collected from multiple acquisition sites and analyzed them with a meta-analytic technique (1). In light of previous findings from our group (1) and others (2–4,26) of increased cortical thickness in ASD, we sought to further examine if previous studies of cortical thickness increases in ASD may have been influenced by cortical boundary abnormalities. Given results from previous neuroimaging and histological studies (18–20,25), we expected to find a greater degree of cortical blurring in individuals with ASD relative to typically developing individuals, or a decrease in BSC. Another goal of this study was to examine the influence of clinical heterogeneity on BSC. Based on previous studies of cortical anatomy, we expected age-related differences in BSC to be greatest in younger ASD individuals (1,27), to vary by sex (28–32) and full-scale IQ (FIQ) (1,33) .

Methods

Study participants: Data included here were acquired from previous studies by the Hospital for Sick Children (Canada), the Cambridge Family Study of Autism (UK) and the UK Medical Research Council Autism Imaging Multicentre Study (UK MRC AIMS). We also included publicly available data from the Autism Brain Imaging Data Exchange (ABIDE) I and II datasets (34,35)

(Table 1 and 2). All data used in this study was preprocessed by SB for analysis in (1), with an original sample size of 3145 participants (1415 individuals with ASD (1165 male/250 female) and 1730 controls (1172 male/558 female), aged 2-65 years. Sample characteristics (including image processing, and which subjects were excluded due to poor motion or segmentation quality) are equivalent between studies, with the exception of the NIMH site, which was not included in this study because the partial volume effects from its lower resolution would compromise the accuracy of BSC measurement. Quality control methodology is outlined in detail in a recent manuscript from our group (1). Supplementary materials of that work (Supplementary Methods: 2. Quality control (QC) and site elimination; 3. Image processing; 7. Quality control analysis) provide detailed accounts of our quality control choices for both the raw and processed data. In total, after segmentation quality control and motion quality control, there remained 1136 subjects: 415 with ASD (303 male/112 female) and 721 controls (438 male/283 female).

Image processing: Data was preprocessed using the minc-bpipe-library pipeline, including N4 bias field inhomogeneity correction (36). Data was then processed through the CIVET 1.1.12 cortical segmentation and cortical thickness pipeline which generated gray/white and pial surfaces and transformed subject brain volumes into standard MNI space (37). All intensity sampling and surface generation was performed in standard space.

BSC calculation: For each of the 77,212 vertices (which exclude all vertices on the midline wall) on the cortical surface of each subject, signal intensity was measured at 10 surfaces that span the region around the gray/white boundary. Together, the 10 intensity samples derived from these new

surfaces cover the bottom quarter of the cortex (the quarter that interfaces with the gray/white

boundary) and a small portion of white matter below the gray/white boundary (Fig. 1).

Gray matter surfaces were created at increasing percentile fractions of the cortical thickness

(0%, 6.25%, 12.5%, 18.75%, 25%, 50%), from the boundary to the pial surface (Fig. 1a,b). New

white matter surfaces were generated at the same distance from the gray/white boundary surface as

the gray matter surfaces, but in the direction of the white matter, with the exception of the white

matter equivalent of the 50% gray matter surface, which was not included because it crossed into

neighboring cortex in thin gyral crowns at certain vertices. See Supplementary Methods section S2

"Surface generation" for more details.

A sigmoid curve (Equation 1) was then fit to the 10 sample points using a non-linear least

squares estimator (Fig. 1c); see Supplementary Methods section S3 "Model fit" for more details.

$$profile_i = a + e^k + \frac{-e^k}{1 + e^{-BSC_i(x-d)}}$$

Equation 1. Sampled intensities at vertex i (*profile$_i$*) and their distance along the axis perpendicular
to the gray matter boundary surface were input to a nonlinear least-squares estimator to fit a
sigmoid function. The d parameter reflects the translation of the sigmoid along the x-axis, where x
is a vector of the displacement along the axis that runs orthogonal to the boundary, increasing
towards the pial surface, parameter a translates the sigmoid along the T1 intensity, parameter k
reflects the height of the sigmoid, and parameter BSC_i reflects the growth rate (steepness) of the
sigmoid.

BSC at each vertex (BSC parameter of the sigmoid curve) was extracted from the estimated

sigmoid curve (Fig. 1d), where a higher BSC reflects greater boundary "sharpness", or a faster

transition in intensity moving from gray matter to white matter (effectively a measure of contrast at

the gray/white matter surface, with higher BSC reflecting high contrast which we take to mean a

bigger difference between the microstructural composition of tissue compartments). See Supplementary Fig. S3 for representative sigmoid fits and associated BSC parameters, and Supplementary Fig. S4 for a spatial distribution of model convergence failures prior to spatial smoothing. These values were then log-transformed and smoothed with a 20 mm full-width half-max (FWHM) smoothing kernel to reduce the effect of noise and simulate a Gaussian distribution, and then values were resampled to a common surface mesh to enable cross-subject comparisons (38).

By measuring T1 signal at multiple distances on either side of the gray/white boundary and fitting an adequately flexible sigmoid curve to the measurements, BSC is theoretically able to capture cortical blurring wherever it occurs relative to the segmented boundary and thus avoids the circular boundary placement problem mentioned in the Introduction. Finally, given that BSC maps correlated moderately with mean curvature (Supplementary Fig. S2b) in certain parts of the cortex, BSC values were further residualized for mean curvature at the vertex-wise level across all subjects within each site in standard space (21).

Figure 1. Intensity sampling and sigmoid fit method used to calculate BSC for each vertex. At each vertex, the T1w image intensity was measured at 10 cortical surfaces surrounding the gray/white boundary, including the mid-surface, gray/white boundary surface, and a total of 8 newly-generated gray and white surfaces equally spaced apart (A, B). A sigmoid curve (Equation 1) was fit to the 10 sample points (C) and parameter *BSC*, reflecting the sigmoid growth rate (D), was log-transformed to create the measure that is the BSC at that vertex. Higher BSC values reflect a quicker transition between gray and white matter and a less blurred cortical boundary, whereas lower BSC values reflect a slower transition between gray and white matter and a more blurred cortical boundary (D). The tissue contrast ratio was calculated by dividing the intensity sampled at the gray 25% surface by the intensity sampled at the white 25% surface.

Statistical Analysis: Data was amassed from multiple sites in order to be sufficiently powered to analyse sources of heterogeneity in ASD. However, analysing multiple sites presents the challenge of site-specific confounds, such as scanner model, scan acquisition protocols, and sample demographics. These limitations were addressed by the prospective meta-analysis technique, where each site was treated as a separate study, and the results were pooled across sites to determine significance at each vertex (1,39,40). Meta-analyses were performed at a vertex-wise level (i.e., performing regressions at each vertex across the brain). Specifically, within each site, for each vertex across the brain where BSC was estimated, a linear regression model was performed to derive per-site Cohen's d effect sizes for the main effect of each variable of interest. Then, the final effect size representing the contribution of all sites was calculated in a random-effects meta-analysis (41) using the *metafor* package in R 3.4.0 (https://www.r-project.org/). See Supplementary Methods section S5 "Statistical Models" for more details. How models were chosen to study the impact of age, sex, and IQ are further detailed below. Final p-values were adjusted for multiple comparisons across all tests performed (all vertices of all analyses performed; as previously in (1)) using the False Discovery Rate (FDR) correction, which controls the proportion of null hypotheses that are falsely rejected (42).

Akaike Information Criterion (AIC) analysis of variable importance: The importance of age (linear term), age (quadratic term), sex, and FIQ were examined by using a vertex-wise Akaike Information Criteria (AIC) analysis. The AIC is a measure that assesses the relative quality of statistical models, where a model with the lowest AIC is considered the best fit for the data (43). AIC takes into account both accuracy and parsimony, because it carries a penalty for increasing the

number of free parameters in the model. Within each site, the AIC was calculated at each vertex for the linear regression model with and without the variable of interest and its interaction with diagnosis. The percentage of sites for which the model including age, age-squared, sex or FIQ was a better fit was calculated for each vertex as a weighted average based on site size (number of subjects scanned at each site). Based on the comparison of the models using AIC, it was determined that age and sex (but not FIQ or age-squared) were important explanatory variables at a substantial proportion of vertices across the brain for BSC, thus motivating our investigation into how these factors influence BSC in ASD (Supplementary Fig. S5). See Supplementary Methods section S5 "Statistical Models" for more details.

Sex-focused analysis: Sex-specific patterns of BSC were assessed by stratifying subjects by sex and performing the same case-control meta-analysis in males and females separately, including age and FIQ as covarites.

Age-focused analyses: The impact of age was examined by stratifying subjects by age, performing separate meta-analyses in subjects who were 18 years old and below, and in subjects who were above 18 years old, including sex and FIQ as covariates. Additionally, an age-centered analysis was performed to assess the trajectory of group differences by shifting the age at which group differences are assessed by 4-year intervals. This type of analysis allows for the interpretation of group differences at various 4-year cross-sections without splitting the dataset into age ranges, maximizing statistical power. The Cohen's d effect size for the main effect of diagnosis was

calculated for each site in each model (one for each age interval) and pooled in a random-effects meta-analysis in the same manner as the case-control comparisons.

Associations between BSC and ASD symptoms/characteristics: Since consistent measures of ASD symptoms or characteristics were not available across all sites, analyses were performed on a subset of individuals who had the same clinical measures. We chose the measure which had the largest number of individuals available, which was the ADOS-2 Calibrated Severity Scores (CSS) to examine overall symptom severity (N =172; also conducted separately in males [N =139] and females [N =33]). The analysis of the relationship between BSC and severity measures (as measured by ADOS-2 CSS) was performed by conducting a multiple regression analysis, calculating the semi-partial correlation of the ADOS-2 scores with BSC at each vertex, per site, with age, sex, and FIQ included in the model, in individuals with ASD only. The semi-partial correlation was then pooled across sites in a random effects meta-analysis, in the same fashion as the Cohen's d effect size in a typical meta-analysis (1).

Assessing spatial overlap between BSC and cortical thickness maps: The spatial correspondence between maps of the effect size of diagnosis on BSC and the effect size of diagnosis on cortical thickness was determined using a 'spin test' (44), which generates null estimates of overlap by applying random rotations to spherical projections of a cortical surface. The Pearson's correlation coefficient between the two original spatial maps is compared to the Pearson's correlation coefficient measured between one original spatial map and the other map's rotated permutations.

Maps of cortical thickness were developed in (1) from a near-identical dataset (excluding the NIMH dataset here).

Results

The tissue intensity ratio correlates with mean curvature: To determine the extent to which tissue intensity ratio may be influenced by cortical curvature, we assessed the correlation between mean curvature and the tissue intensity ratio (see Figure 1 and Supplementary Methods section S4 "Tissue intensity ratio" for methods). In a vertex-wise single-subject analysis, we found that the tissue intensity ratio was lower in gyri compared to sulci (Supplementary Fig. S1). Furthermore, in a cross-subject analysis to determine in which areas of the brain this relationship is present, we found a negative correlation between the tissue intensity ratio and mean curvature for most vertices in the brain (Supplementary Fig. S2a, c, d), including vast areas of the frontal, parietal, and occipital lobes. These results are consistent with the hypothesis that the impact of curvature on cortical T1w intensity extends to the tissue intensity ratio, and calls for a measure of cortical contrast between the gray and white matter that is less susceptible to bias by other cortical features. When assessed for its relationship to curvature, we found that BSC correlated minimally with mean curvature in isolated regions of the anterior frontal, parietal, and temporal lobes (Supplementary Fig. S2b), and did not correlate in any regions after regressing the values against mean curvature at the vertex level.

Greater BSC in individuals with ASD: We observed regions of significantly greater BSC in individuals with ASD compared to controls in the bilateral superior temporal gyrus, inferior temporal gyrus, and left inferior frontal gyrus ($<5\%$ FDR, peak Cohen's d = 0.36) (Fig. 2a), corresponding to a faster transition from cortical gray matter to white matter relative to controls.

Effect sizes varied moderately between sites but were mostly positive (Fig. 2b, Supplementary Fig. S6).

Figure 2. Diagnostic group comparisons of BSC. (A) Individuals with ASD had significantly higher BSC measures (<5% FDR, peak Cohen's d = 0.38) in the bilateral superior temporal gyrus, inferior temporal gyrus, and left inferior frontal gyrus. (B) Forest plot displaying site-specific effect sizes at a peak vertex in the left frontal gyrus represented by an asterix in (A). Sex-specific patterns of group differences in BSC were observed, with females showing significantly increased BSC in the bilateral superior parietal gyrus and superior temporal gyrus (C), and mxtales displaying significantly greater BSC in the bilateral inferior

temporal gyrus and left inferior frontal lobe (D). Further, the magnitude of the effect size was greater in females, where the peak Cohen's d value observed in females was 0.63, and in males, it was 0.32.

Sex-specific boundary alterations: In both the male and female subgroups, individuals with ASD had significant increases in BSC, but these increases were observed in different brain regions in males and females and with larger effect sizes in the female group (Fig. 2c, d). Females with ASD had significantly greater BSC in the bilateral superior parietal gyrus and superior temporal gyrus, with a peak Cohen's d of 0.63 (Fig. 2c). Males with ASD displayed significantly greater BSC in the bilateral inferior temporal gyrus and left inferior frontal lobe, with a peak Cohen's d of 0.32 (Fig. 2d).

Age-specific patterns of boundary alterations: In the age-stratified analysis, individuals with ASD above and below 18 both showed significantly greater BSC than their typically developing counterparts, though these group differences showed age-specific patterning and effect sizes, and the main effect of diagnosis was stronger in the age group over 18. Individuals with ASD under 18 had significantly greater BSC in the bilateral superior temporal gyrus and right postcentral gyrus, with a peak Cohen's d of 0.41 (Fig. 3a). Individuals with ASD over 18 had significantly increased BSC in the bilateral precuneus and superior temporal gyrus, with a peak Cohen's d of 0.62 (Fig. 3b). In the age-centered analysis, which examined group differences at specific ages, group differences were greatest between the ages of 12 and 20 in the right superior temporal gyrus and left inferior temporal gyrus (Fig 3c, d).

Figure 3. Age-stratified and age-centered analyses. Individuals under 18 with ASD had significantly higher BSC measures in the bilateral superior temporal gyrus as well as the left precentral gyrus (<5% FDR, peak Cohen's d = 0. 41) (A). Individuals over 18 with ASD showed significantly increased BSC in the bilateral precuneus and superior temporal gyrus (<5% FDR, peak Cohen's d = 0.62). (B) For the age-centered analysis, group differences in BSC were greatest between the ages of 12 and 20 in the right superior temporal gyrus and left inferior temporal gyrus. (D) Zoom of the left superior temporal lobe (top) and right inferior temporal lobe (bottom). (E) Plot of BSC across age in a single site (Toronto) at vertex highlighted with a white asterix in A.

Minimal correlation between BSC and ASD severity measures: We observed a significant

positive correlation between BSC and ADOS-2 CSS in individuals with ASD in the right medial

temporal pole (Fig. 4a) Given our findings of sex-specific regions of BSC differences in subjects

with ASD, we explored the relationship between BSC and ASD severity separately in males and

females. In the female group, we found a significant positive association between BSC and

ADOS-2 CSS in the left parietal lobe. No correlations between severity score and BSC were

observed in the male-only subset.

Figure 4. Relationship between BSC and ADOS-calibrated severity scores (CSS). Across all subjects with
ASD with severity scores, ADOS-CSS was positively correlated with BSC, shown for a peak vertex in the

right medial temporal gyrus (A). Correlations between ADOS-CSS and BSC were also observed in the female-only group in the left parietal lobe (B).

Overlap between maps of diagnostic effect on cortical thickness and BSC: We observed a

significant overlap between the maps of Cohen's d effect size of diagnosis on BSC and the map of

Cohen's d effect size of diagnosis on CT (p < 0.01) (Fig. 5a,c), as well as between the maps of

FDR-corrected q-values (p < 0.01) (Fig. 5b, Supplementary Fig. S7).

Figure 5. Cohen's d effect size maps of BSC (A) and cortical thickness (B - used with permission, from Bedford et al., 2019). Spatial correspondence assessed by the Pearson correlation coefficient in a permutation-based 'spin-test' analysis between BSC and cortical thickness is demarcated in red (C) was significant (p= 0.00; p<<0.001 as per the software output) with 1000 null spatial permutations. FDR-thresholded q-value maps of significant increases in BSC and increases in cortical thickness in individuals with ASD (D).

Discussion

In this study we employed a vertex-wise meta-analysis on a large multi-site dataset to investigate BSC in ASD, finding significant group-level increases in BSC in lateral frontal and temporal regions as well as sex-specific and age-specific patterns of BSC increases in individuals with ASD. As BSC is parameterized to capture cortical blurring which may be a product of differences in neural migration and intracortical myelination, there are several considerations for interpreting our observation of increased BSC in ASD. Neuronal migration defects are predicted to result in supernumerary neurons in the white matter compartment directly below the cortex (19,25,45). In previous neuroimaging studies examining cortical blurring at the gray/white boundary using the tissue intensity ratio, the presence of ectopic neurons is thought to be captured by a lower intensity difference between cortex and white matter (19). Thus, our finding of an increased intensity difference (as indexed by greater BSC) between cortex and white matter is unlikely to reflect differences in neuronal migration.

Since T1w intensity is thought to reflect underlying myeloarchitecture more so than cytoarchitecture (46), BSC may also reflect intracortical myelination: a reduction of myelin content (and presumably, of T1w signal) in the lower layers of the cortex would appear in T1w MRI as a sharper transition in intensity moving from gray to white matter. This assumption is supported by the general agreement between the group-average BSC map (Supplementary Fig. S8) and the intracortical myelin map derived using the T1w/T2w ratio by Glasser and Van Essen (47) and is further supported by evidence from genetic (48) and molecular (16,49) studies, preclinical mouse models (14,50), and postmortem histology (15) suggesting altered intracortical myelination in ASD.

Mounting evidence from diffusion imaging and resting state fMRI studies support the characterization of ASD as a connectopathy, or a brain network disorder. Since our proposed measure is parameterized to capture myelination in the lower layers of the cortex (a region through which fibers that connect distant brain regions pass), the higher BSC observed in ASD is supported by studies that find reduced long-range cortical connectivity (51,52) and thalamo-cortical connectivity (53,54) in ASD. As such, the contribution of myelination is increasingly relevant to understanding the neurodevelopmental mechanisms of ASD.

Two neuroimaging studies to date investigated boundary microstructure in ASD using a different metric called the tissue intensity ratio, reporting lower tissue intensity ratio in ASD which may indicate greater intracortical myelination or the presence of supernumerary neurons in the superficial white matter compartment directly beneath the gray/white boundary (17–19). This discrepancy to the present findings could be due, in part, to differences in preprocessing and analysis methods (particularly the use of the tissue intensity ratio). The tissue intensity ratio suffers from the ambiguity arising from the blurring around the cortical boundary which, in turn, alters placement of the gray/white boundary that serves as a reference point for the component gray and white measurements. Additionally, the present study uses a sample size that is considerably larger than previous studies and is more suitably powered to detect group differences in high-variability populations. Examining the directionality in each of the forest plots reveals how different studies may, indeed, yield different findings and how they may be dependent on age range and demographic composition. The data in the present study were rigorously filtered for motion and image processing artefacts (1) whereas quality control procedures were not described in detail in previous studies, making it difficult to assess the effect of motion or inaccurate segmentation on

reported results. It is possible that measures of cortical blurring using the tissue intensity ratio in previous ASD studies have been obscured by motion-induced blurring of the gray-white matter boundary (55,56), especially considering individuals with ASD are more likely to move during the scan than controls (1,57). It is also possible that factors such as age, sex, FIQ, and symptom severity are influencing case-control differences in previous studies of cortical blurring in autism (58). Therefore, we attempted to determine the degree to which diagnostic differences in BSC are modulated by these factors.

The male bias in ASD prevalence (3:1 males:females diagnosed (59) as well as sex differences in behaviour and key ASD-related phenotypes such as restrictive and repetitive behaviours (60,61) have spurred the investigation of neuroanatomical sex differences in ASD (32). Given their scarcity, females with ASD are a difficult population to recruit, and, as such, our understanding of the modulating effect of sex on neuroanatomy in individuals with ASD is relatively rudimentary. However, this knowledge gap is beginning to narrow as a result of efforts to curate large-scale publicly-available neuroimaging datasets (35) and an increased awareness of the importance of female representation in ASD studies (31). By combining publicly-available datasets with several multicentre consortium datasets, we were able to incorporate data from 117 females with ASD in this study. We observed sex-specific patterns of BSC, where females with ASD displayed higher BSC in the superior temporal and parietal lobes, whereas males with ASD displayed a greater BSC in the inferior temporal and frontal lobes. Sex-specific patterns of cortical neuroanatomy in autism have recently been reported, in both gross volumetric measures (1,62) and altered connectivity (28,29). The maximum effect size of diagnosis in the female sample was almost twice as high as the effect size of diagnosis observed in the male sample. BSC also correlated with

ADOS-CSS in the female-only sample. Taken together, these findings support a differential

neuroanatomical presentation of autism in males and females.

Age is another factor that is known to modulate diagnostic group differences. Though most

brain differences in ASD have been investigated in a child or adolescent population, there is

evidence that differences in brain anatomy related to ASD are present even in adulthood (63–65).

Intracortical myelination has shown to be ongoing even past adolescence, with accelerated

myelination until ~30 years of age, followed by a period of stability, and then a decrease in

myelination from the late 50s (12,66). Though our use of cross-sectional data limits our

interpretation of the age-centered results as being reflective of developmental processes, the peak

diagnostic group differences observed in adolescence and adulthood may be the result of reduced or

protracted myelination in ASD relative to the rate of myelination in typical development, as

observed in postmortem histology (15) and mouse models of ASD (14,67).

Intracortical myelination has been shown to drive MRI-based measures of cortical thickness

in the visual cortex across development (68), potentially by pushing the gray-white matter boundary

deeper into the cortex. In our analysis, we found a significant spatial overlap between maps of the

effect size of diagnosis on BSC and the effect size of diagnosis on cortical thickness (1). These

results suggest that the neuro-phenotype of increased cortical thickness in ASD, as observed in (1)

and many other studies, may be partially driven by lower levels of intracortical myelination in ASD

relative to controls. Moving forward, a longitudinal design would allow us to determine in the same

individual, the evolution of the pattern of BSC across development, and would allow for relating

patterns of BSC over time with respect to behaviours and autism symptomatology. Additionally,

since age may substantially moderate the pattern of sex differences, a longitudinal framework would allow for a more precise investigation into this time-sensitive relationship.

However, while cortical thickness differences observed in our previous work (1) correlate with autism severity scores in many regions of the brain, we found minimal correlations between BSC and ASD severity. This disparity suggests that intracortical myelination may inflate MRI-based measures of cortical thickness in ASD for several areas of the brain, but the cortical overgrowth phenotype itself is driven by separate biological processes (such as reduced synaptic pruning) that are more clinically relevant to ASD severity. On the other hand, deficits in intracortical myelination may be a neurobiological hallmark of the disorder that does not alter ASD behavioural severity beyond a certain point, and thus does not correlate with the ADOS-CSS score.

There are several limitations to this study. First, although our BSC maps correspond well with established patterns of intracortical myelination, the T1w intensity also reflects other biological properties including water content, iron, and dendrite density, and therefore may reflect more than the degree of myelination in the cortex (69). Second, abnormalities in the superficial white matter may also drive some of the changes to BSC. Though a vertex-wise analysis of the intracortical or superficial white matter intensities independently is precluded by between-scanner intensity profile differences and by the distorting effect of inhomogeneity gradients across images, it would be possible to evaluate differences in normalized intracortical intensities that have been scaled to the intensity of a predetermined sample of white matter within each subjects' brain. Finally, since the distance sampled in the white matter is based on a percentage of the cortical thickness, the depth at which white matter is measured varies between regions with different cortical thicknesses.

Given these findings of increased BSC in individuals with ASD, which may reflect the degree of myelination in the lower layers of the cortex, it will be pertinent to investigate how subcortical volume changes in ASD relate to BSC in cortical sensory regions with high thalamic input (26,70). Considering the importance of intracortical myelination on the fidelity of neural connections and the maintenance of networks, these results may help us better understand the cognitive and behavioural atypicalities seen in ASD. Finally, the significant spatial correspondence observed between maps of cortical thickness and of BSC calls for a reconsideration of what biological phenomena may underlie the MRI-derived measures of cortical thickness increases in autism.

Supplementary Information

Supplementary Methods

S1. Sample details

S1.1 Data processing: All data used in this study was preprocessed by SB for analysis in Bedford et al., (2019), and sample characteristics (including image processing, and which subjects were excluded due to poor motion or CIVET quality) are equivalent between studies, with the exception of the NIMH site, which was not included here. See Bedford et al. (2019) for imaging parameters and preprocessing considerations for each site. The data used in this study was obtained from scans whose CIVET/1.1.12 segmentations quality had already been assessed. As such, a newer version CIVET was not re-run on this data.

S1.2 Quality Control: In-scanner movements cause motion artefacts in MR images, which can influence certain morphological measurements like cortical thickness and cortical volume (55,56). As such, performing quality control (QC) assessments for motion artefacts is essential to eliminate statistical bias in disorders associated with movement abnormalities, such as ASD. All data in this report was rated for motion by three independent raters (SB, MMC, and ST). Data from subjects with excessive motion scores were excluded from analyses (1). In addition to motion QC, data from subjects whose CIVET QC scores were below a certain threshold were excluded (1). In total, after CIVET and motion QC, there remained 1056 subjects: 427 with ASD (310 male/117 female) and 629 controls (323 male/306 female).

S2. Surface Generation

S2.1 New gray matter surfaces A gray matter surface at 25% of the distance from the gray/white boundary surface to the pial surface was generated by averaging the vertices of the gray/white boundary surfaces (0%) and mid-surfaces (50%), which were created in standard space by CIVET/1.1.12, using the function *average_surfaces*. For each subject, the 12.5% gray matter surfaces were generated by combining the 0% surface with the 25% surface, 6.25% surfaces by combining the 0% surface with the 12.5% surface, and 18.75% surfaces by combining the 12.5% surfaces with the 25% surfaces.

S2.2 New white matter surfaces Each new white matter surface was generated by first creating vectors between each vertex on the gray/white boundary surface and the corresponding vertex on the gray matter surfaces (repeated separately for each gray matter surface generated (S2.1), with the exception of the 50% or mid-surface). The vectors were then inverted in the x, y, and z dimensions and added to the coordinates of the graywhite boundary surface vertices, such that white matter surfaces were created at the same distance from the gray/white boundary as the gray matter surfaces, but in the opposite direction (toward the superficial white matter instead of toward the pial surface).

S3. Model fit A sigmoid curve (Equation 1) was fit to the 10 sampled intensities by finding the optimal least squares solution over parameters (a, k, c, d) between a[min(yvalues), 2000], k[0, 100], c[0,1], and d[-50,50], using the nls() function and "port" algorithm in R (Figure S3). A base sigmoid function (71) was reparameterized to constrain the shape of the curve to be increasing from the pial

surface to the superficial white matter. In a single-subject analysis, convergence was achieved for ~98% of vertices that were regularly distributed across the brain (Figure S4).

S4. Tissue Intensity Ratio The tissue intensity ratio was calculated by dividing the gray matter intensity measured at 25% of the cortical thickness by the white matter intensity measured at an equal distance (25% of the cortical thickness) in the direction of the white matter. Values were resampled to a common mesh and smoothed, except for the single-subject analysis (Supplementary Fig 1), in which values were not resampled before smoothing so as to not introduce correlations with the curvature that are used to guide surface-based registration.

S5. Statistical models For the case-control comparisons, the following model was used, with i representing each vertex across the brain:

Case-control analysis: $BSC_i \sim \beta_0 Intercept + \beta_1 Age + \beta_2 Sex + \beta_3 Diagnosis + \beta_4 FIQ + \varepsilon$

Sex-stratified analysis: $BSC_i \sim \beta_0 Intercept + \beta_1 Age + \beta_2 Diagnosis + \beta_3 FIQ + \varepsilon$

Age-centered analysis: $BSC_i \sim \beta_0 Intercept + \beta_1 (Age-a)*Diagnosis + \beta_2 Sex + \beta_3 FIQ + \varepsilon$

      Where *a* is the age at which group differences are assessed.

Age-stratified analysis: $BSC_i \sim \beta_0 Intercept + \beta_1 Age + \beta_2 Sex + \beta_3 Diagnosis + \beta_4 FIQ + \varepsilon$

Model assessment: To test the importance of sex, age and FIQ, an Aikaike Information Criteria (AIC) was used. The AIC is a measure that assesses the relative quality of statistical models, where a model with the lowest AIC is considered the best fit for the data.

Sex:

$$BSC_i = \beta_0 + \beta_1 Diagnosis + \beta_2 Sex + \beta_3 (Diagnosis*Sex) + \beta_4 Age + \beta_5 FIQ + \varepsilon_i$$

$$BSC_i = \beta_0 + \beta_1 Diagnosis + \beta_2 Age + \beta_3 FIQ + \varepsilon_i$$

Age:

$$BSC_i = \beta_0 + \beta_1 Diagnosis + \beta_2 Sex + \beta_3 Age + \beta_4 (Diagnosis*Age) + \beta_5 FIQ + \varepsilon_i$$

$$BSC_i = \beta_0 + \beta_1 Diagnosis + \beta_2 Sex + \beta_5 FIQ + \varepsilon_i$$

Age-squared:

$$BSC_i = \beta_0 + \beta_1 Diagnosis + \beta_2 Sex + \beta_3 Age + \beta_4 Age^2 + \beta_5 (Diagnosis*Age^2) + \beta_6 FIQ + \varepsilon_i$$

$$BSC_i = \beta_0 + \beta_1 Diagnosis + \beta_2 Sex + \beta_5 Age + \beta_4 FIQ + \varepsilon_i$$

FIQ:

$$BSC_i = \beta_0 + \beta_1 Diagnosis + \beta_2 Sex + \beta_3 Age + \beta_4 FIQ + \beta_4 (Diagnosis*FIQ) + \varepsilon_i$$

$$BSC_i = \beta_0 + \beta_1 Diagnosis + \beta_2 Sex + \beta_3 Age + \varepsilon_i$$

For the AIC comparisons of polynomial terms of age, the following models were compared

$$BSC_i = \beta_0 + \beta_1 Diagnosis + \beta_2 Age + \beta_3 (Diagnosis*Age) + \beta_4 Sex + \varepsilon_i$$

$$BSC_i = \beta_0 + \beta_1 Diagnosis + \beta_2 Age + \beta_3 (Diagnosis*Age) + \beta_4 Age^2 + \beta_5 (Diagnosis*Age^2) + \beta_6 Sex + \varepsilon_i$$

Tables

Table 1. Subject demographics of the cohort under study. Group differences in age, sex distribution, and FIQ were also examined using t-tests (for continuous variables) and Chi-squared tests (for categorical variables).

Subjects by site Before QC (and after QC)	Age (years) [Median]	Total (after QC)	Female ASD (after QC)	Male ASD (after QC)	Female Control (after QC)	Male Control (after QC)	Full-scale IQ (FIQ) [Median]
ABIDE I - 17 sites (4 sites)	6-64 [14.7] (6-58 [16.9])	1101 (263)	64 (14)	466 (89)	99 (38)	472 (122)	41-148 [109] (69-148 [110])
ABIDE II - 16 sites (4 sites)	5-64 [11.7] (5-34 [11.1])	1044 (292)	73 (25)	414 (82)	175 (69)	382 (116)	49-149 [112] (49-144 [110])
SickKids Hospital, Toronto	4-65 [14.0] (4-49[14.8])	521 (321)	25 (18)	106 (62)	194 (117)	196 (124)	69-149 [111] (69-149 [112])
Cambridge Family Autism Study	12-18 [14.7] (12-18 [15.0])	96 (52)	17 (12)	39 (15)	20 (14)	20 (11)	73-146 [108] (82-139 [107])

UK MRC Autism Imaging Multicentre Study (University of Cambridge, King's College London)	18-52 [25.7] (18-52 [27.7])	253 (208)	54 (43)	72 (55)	54 (45)	73 (65)	73-137 [115] (73-137 [117])
Total Before QC	**2-65 [13.8]**	**3015**	233	1097	542	1143	41-149 [111]
Difference between sites (ANOVA/X²; total sample)	F(29, 2913) = 101.9, p < 0.001		$X^2(87, N = 3015) = 544,35, p < 0.001$				F(29, 2893) = 7.985, p < 0.001
Total After QC	**2-65 [14.0]**	**1136**	112	303	283	438	49-149 [112]
Difference (ANOVA/X²; QC'd sample)	F(9, 1126) = 118.1, p < 0.001		$X^2(27, N = 1136) = 148.2, p < 0.001$				F(9, 1126) = 0.94, p =0.49

Table 2. Subject demographics broken down by ABIDE I and II site.

Subjects by ABIDE	Age	Total	F- ASD	M- ASD	F- Ctl	M- Ctl	Full-scale

site Before QC (and after QC)	(years) [Median]	(after QC)	(after QC)	(after QC)	(after QC)	(after QC)	IQ (FIQ) [Median]
Kennedy Krieger Institute (KKI; ABIDE I&II)	8-13 [10.3] (8-13 [10.4])	256 (148)	19 (11)	58 (21)	65 (45)	123 (71)	63-143 [113] (69-143 [114])
Ludwig Maximilians University Munich (MAX MUN; ABIDE I)	7-58 [26.0] (7-58 [29.5])	57 (38)	3 (3)	21 (14)	4 (4)	29 (17)	79-133 [112] (93-133 [112])
New York University (NYU; ABIDE I & II)	5-39 [10.9] (5-39 [12.1])	289 (179)	19 (10)	135 (74)	28 (19)	107 (76)	67-148 [109] (67-148 [109])
Oregon Health and Science University (OHSU; ABIDE I & II)	7-15 [11.0] (8-15 [11.0])	121 (97)	7 (5)	43 (33)	29 (25)	42 (34)	69-140 [115] (69-136 [114])
San Diego State University (SDSU; ABIDE I &	7-18 [13.8] (8-18	94 (45)	8 (6)	39 (16)	8 (3)	39 (20)	66-141 [105] (66-139

II)	[14.4])						[104])
University of Michigan (UM; ABIDE I)	8-28 [14.0] (9-28 [15.8])	145 (55)	10 (4)	58 (17)	18 (12)	59 (22)	76-147 [108] (78-147 [110])

Supplementary Figures

Supplementary Figure S1. Left: tissue intensity ratio values displayed on a single subject's mid-cortical surface (i.e. the surface at 50% of the distance between the pial and white matter surface). Right: vertex-wise correlation between the tissue intensity ratio calculated as the ratio of white matter to gray matter intensities and mean curvature. The tissue intensity ratio is significantly negatively correlated with mean curvature for all subjects (<5% FDR, data not shown) in a single exploratory site (Sick Kids; 88 ASD/314 CTL), meaning that the tissue intensity ratio is higher in convex regions, such as sulci, and lower in concave regions, like gyri. Vertex-wise correlations were assessed in the native space before resampling to a common surface mesh.

Supplementary Figure S2. Regions of significant (<5% FDR) cross-subject correlation of the tissue intensity ratio with mean curvature (A) and of BSC with mean curvature (B) at each vertex in a single exploratory site (Sick Kids; 88 ASD/314 CTL). Relationship between mean curvature and tissue intensity ratio across subjects at a vertex in the right inferior frontal sulcus, Pearson's correlation coefficient = -0.56 (C). Map of the correlation coefficient between the tissue intensity ratio and mean curvature (D); white asterisk on the left lateral surface represents the vertex shown in (C).

Supplementary Figure S3. Image intensity samples (black dots) and estimated sigmoid curve (black line) from 12 randomly chosen vertices in a single subject. The x-axis displays distance along the cortical profile, starting at the outermost surface in the superficial white matter (denoted as -25%) and ending at the gray matter midsurface (50%). BSC values prior to log-transformation are displayed in the title of each plot.

Supplementary Figure S4. Distribution of convergence failures (i.e. where a sigmoid curve could not be adequately fit) for a single subject, shown in red. The subcortical surface is masked in black.

Supplementary Figure S5. Akaike information criterion (AIC) analysis reflecting the importance of sex, age, and FIQ on BSC. Color maps represent the percentage of sites (calculated as a weighted average, based on site size after QC) for which BSC is better fit by a linear model including sex, age, age squared, or FIQ and their interaction with diagnosis, than one without. The model including the sex term and its interaction with diagnosis was a better fit in 0-82% of sites, the model including age and its interaction with diagnosis was a better fit at 0-100% of sites, and the model including FIQ and its interaction with diagnosis was a better fit at 0-67% of sites. The left hemisphere on the left, right hemisphere on the right, for each parameter tested.

A Site   Cohen's d [95% CI]

Site		%	Cohen's d [95% CI]
KKI		12.77%	0.55 [0.22, 0.88]
MAX_MUN		3.07%	-0.20 [-0.88, 0.47]
NYU		15.75%	0.41 [0.11, 0.70]
OHSU		8.48%	0.23 [-0.18, 0.63]
SDSU		3.84%	0.36 [-0.24, 0.96]
UM		4.52%	0.17 [-0.39, 0.73]
CAM		4.50%	0.27 [-0.29, 0.82]
TORONTO		28.71%	0.27 [0.05, 0.49]
IoP		8.96%	0.32 [-0.07, 0.71]
Cambridge		9.42%	0.15 [-0.24, 0.53]
RE Model		100.00%	0.30 [0.18, 0.42]

-1   0 0.5 1
Observed Outcome

q-values
0.01

0.05

B Site   Cohen's d [95% CI]

Site		%	Cohen's d [95% CI]
KKI		13.10%	0.28 [-0.05, 0.61]
MAX_MUN		3.07%	0.11 [-0.57, 0.78]
NYU		15.94%	0.23 [-0.07, 0.52]
OHSU		8.40%	0.33 [-0.08, 0.74]
SDSU		3.89%	0.12 [-0.47, 0.72]
UM		4.52%	0.15 [-0.41, 0.70]
CAM		4.50%	0.24 [-0.32, 0.80]
TORONTO		28.54%	0.32 [0.10, 0.54]
IoP		8.65%	0.62 [0.22, 1.02]
Cambridge		9.40%	0.14 [-0.24, 0.53]
RE Model		100.00%	0.28 [0.17, 0.40]

-1 -0.5   0 0.5   1 1.5
Observed Outcome

Supplementary Figure S6. Significant vertex-wise group differences (ASD vs. control) in BSC across all individuals at 5% FDR. Forest plots show effect sizes per imaging site at two regions of interest highlighted with a red arrow: the left posterior postcentral gyrus (A) and right inferior temporal gyrus (B). Top row displays lateral views (left hemisphere on the left, right hemisphere on the right), and the bottom row displays medial views (left hemisphere on the left, right hemisphere on the right).

Supplementary Figure S7. Null correlation density from overlap between maps of cortical thickness and BSC FDR-corrected q-values of main effect of ASD diagnosis (p = 0.02) generated by rotating a cortical surface randomly and comparing the Pearson's correlation coefficient measured in the true overlap with those measured between the null models of overlap.

Supplementary Figure S8. Average BSC map versus average T1/T2 ratio myelin map from Glasser et al., (2011). Olafson et al. BSC map depicts the average BSC values across control subjects aged 22-35 for a single site (Sick Kids) for correspondence with the HCP dataset used by Glasser et al. (2011) to derive T1/T2 ratio maps. Regions of greatest BSC include the motor-somatosensory strip in the central sulcus, visual cortex in the occipital lobe, and early auditory areas in the Sylvian fissure, matching areas that were reported to contain the heaviest intracortical myelination as measured by the T1/T2 ratio (47). The top row displays lateral views (left hemisphere on the left, right hemisphere on the right), and the bottom row displays medial views, for each map.

References:

1.   Bedford SA, Park MTM, Devenyi GA, Tullo S, Germann J, Patel R, et al. Large-scale analyses of the relationship between sex, age and intelligence quotient heterogeneity and cortical morphometry in autism spectrum disorder. Mol Psychiatry [Internet]. 2019 Apr 26; Available from: http://dx.doi.org/10.1038/s41380-019-0420-6

2.   Anagnostou E, Taylor MJ. Review of neuroimaging in autism spectrum disorders: What have we learned and where we go from here [Internet]. Molecular Autism. 2011. Available from: http://dx.doi.org/10.1186/2040-2392-2-4

3.   Hazlett HC, Gu H, Munsell BC, Kim SH, Styner M, Wolff JJ, et al. Early brain development in infants at high risk for autism spectrum disorder. Nature. 2017 Feb 15;542(7641):348–51.

4.   Courchesne E, Mouton PR, Calhoun ME, Semendeferi K, Ahrens-Barbeau C, Hallet MJ, et al. Neuron number and size in prefrontal cortex of children with autism. JAMA. 2011 Nov 9;306(18):2001–10.

5.   Park MTM, Raznahan A, Shaw P, Gogtay N, Lerch JP, Chakravarty MM. Neuroanatomical phenotypes in mental illness: identifying convergent and divergent cortical phenotypes across autism, ADHD and schizophrenia. J Psychiatry Neurosci. 2018 May;43(3):201–12.

6.   Hahamy A, Behrmann M, Malach R. The idiosyncratic brain: distortion of spontaneous connectivity patterns in autism spectrum disorder. Nat Neurosci. 2015 Feb;18(2):302–9.

7.   Rudie JD, Brown JA, Beck-Pancer D, Hernandez LM, Dennis EL, Thompson PM, et al. Altered functional and structural brain network organization in autism. Neuroimage Clin. 2012 Nov 16;2:79–94.

8.   Just MA, Keller TA, Malave VL, Kana RK, Varma S. Autism as a neural systems disorder: a theory of frontal-posterior underconnectivity. Neurosci Biobehav Rev. 2012 Apr;36(4):1292–313.

9.   Pinto D, Delaby E, Merico D, Barbosa M, Merikangas A, Klei L, et al. Convergence of genes and cellular pathways dysregulated in autism spectrum disorders. Am J Hum Genet [Internet]. 2014; Available from: http://dx.doi.org/10.1016/j.ajhg.2014.03.018

10. Huguet G, Ey E, Bourgeron T. GG14CH26-Bourgeron The Genetic Landscapes of Autism Spectrum Disorders. Annu Rev Genomics Hum Genet [Internet]. 2013; Available from: http://dx.doi.org/10.1146/annurev-genom-091212-153431

11. Reiner O, Karzbrun E, Kshirsagar A, Kaibuchi K. Regulation of neuronal migration, an emerging topic in autism spectrum disorders. J Neurochem. 2016 Feb;136(3):440–56.

12. Grydeland H, Walhovd KB, Tamnes CK, Westlye LT, Fjell AM. Intracortical myelin links

with performance variability across the human lifespan: results from T1- and T2-weighted MRI myelin mapping and diffusion tensor imaging. J Neurosci. 2013 Nov 20;33(47):18618–30.

13. Deoni SCL, Dean DC 3rd, Remer J, Dirks H, O'Muircheartaigh J. Cortical maturation and myelination in healthy toddlers and young children. Neuroimage. 2015 Jul 15;115:147–61.

14. Graciarena M, Seiffe A, Nait-Oumesmar B, Depino AM. Hypomyelination and Oligodendroglial Alterations in a Mouse Model of Autism Spectrum Disorder. Front Cell Neurosci. 2018;12:517.

15. Zikopoulos B, Barbas H. Changes in Prefrontal Axons May Disrupt the Network in Autism. Journal of Neuroscience [Internet]. 2010; Available from: http://dx.doi.org/10.1523/jneurosci.2257-10.2010

16. Canali G, Garcia M, Hivert B, Pinatel D, Goullancourt A, Oguievetskaia K, et al. Genetic variants in autism-related CNTNAP2 impair axonal growth of cortical neurons. Hum Mol Genet. 2018 Jun 1;27(11):1941–54.

17. Norbom LB, Doan NT, Alnæs D, Kaufmann T, Moberget T, Rokicki J, et al. Probing Brain Developmental Patterns of Myelination and Associations With Psychopathology in Youths Using Gray/White Matter Contrast. Biol Psychiatry. 2019 Mar 1;85(5):389–98.

18. Mann C, Bletsch A, Andrews D, Daly E, Murphy C, MRC AIMS Consortium, et al. The effect of age on vertex-based measures of the grey-white matter tissue contrast in autism spectrum disorder. Mol Autism. 2018 Oct;9(1):49.

19. Andrews DS, Avino TA, Gudbrandsen M, Daly E, Marquand A, Murphy CM, et al. In Vivo Evidence of Reduced Integrity of the Gray-White Matter Boundary in Autism Spectrum Disorder. Cereb Cortex [Internet]. 2017; Available from: http://dx.doi.org/10.1093/cercor/bhw404

20. Bezgin G, Lewis JD, Evans AC. Developmental changes of cortical white–gray contrast as predictors of autism diagnosis and severity. Transl Psychiatry. 2018 Nov 16;8(1):249.

21. Sereno MI, Lutti A, Weiskopf N, Dick F. Mapping the human cortical surface by combining quantitative T1 with retinotopy. Cereb Cortex [Internet]. 2013; Available from: http://dx.doi.org/10.1093/cercor/bhs213

22. Bok ST. Histonomy of the cerebral cortex. Amsterdam: Elsevier; 1959.

23. Libero LE, Schaer M, Li DD, Amaral DG, Nordahl CW. A Longitudinal Study of Local Gyrification Index in Young Boys With Autism Spectrum Disorder. Cereb Cortex. 2019 Jun 1;29(6):2575–87.

24. Kohli JS, Kinnear MK, Fong CH, Fishman I, Carper RA, Müller R-A. Local Cortical Gyrification is Increased in Children With Autism Spectrum Disorders, but Decreases Rapidly

in Adolescents. Cereb Cortex. 2018;1–12.

25. Avino TA, Hutsler JJ. Abnormal cell patterning at the cortical gray-white matter boundary in autism spectrum disorders. Brain Res [Internet]. 2010; Available from: http://dx.doi.org/10.1016/j.brainres.2010.08.091

26. Schuetze M, Park MTM, Cho IY, MacMaster FP, Chakravarty MM, Bray SL. Morphological Alterations in the Thalamus, Striatum, and Pallidum in Autism Spectrum Disorder. Neuropsychopharmacology. 2016 Oct;41(11):2627–37.

27. Khundrakpam BS, Lewis JD, Kostopoulos P, Carbonell F, Evans AC. Cortical Thickness Abnormalities in Autism Spectrum Disorders Through Late Childhood, Adolescence, and Adulthood: A Large-Scale MRI Study. Cereb Cortex [Internet]. 2017; Available from: http://dx.doi.org/10.1093/cercor/bhx038

28. Irimia A, Torgerson CM, Jacokes ZJ, Van Horn JD. The connectomes of males and females with autism spectrum disorder have significantly different white matter connectivity densities. Sci Rep. 2017 Apr 11;7:46401.

29. Zeestraten EA, Gudbrandsen MC, Daly E, de Schotten MT, Catani M, Dell'Acqua F, et al. Sex differences in frontal lobe connectivity in adults with autism spectrum conditions. Transl Psychiatry. 2017 Apr 11;7(4):e1090.

30. Greenberg DM, Warrier V, Allison C, Baron-Cohen S. Testing the Empathizing-Systemizing theory of sex differences and the Extreme Male Brain theory of autism in half a million people. Proc Natl Acad Sci U S A. 2018 Nov 27;115(48):12152–7.

31. Lai M-C, Lombardo MV, Auyeung B, Chakrabarti B, Baron-Cohen S. Sex/gender differences and autism: setting the scene for future research. J Am Acad Child Adolesc Psychiatry. 2015 Jan;54(1):11–24.

32. Lai MC, Lerch JP, Floris DL, Ruigrok ANV, Pohl A, Lombardo MV, et al. Imaging sex/gender and autism in the brain: Etiological implications [Internet]. Journal of Neuroscience Research. 2017. Available from: http://dx.doi.org/10.1002/jnr.23948

33. Lotspeich LJ, Kwon H, Schumann CM, Fryer SL, Goodlin-Jones BL, Buonocore MH, et al. Investigation of Neuroanatomical Differences between Autism and Asperger Syndrome. Arch Gen Psychiatry [Internet]. 2004; Available from: http://dx.doi.org/10.1001/archpsyc.61.3.291

34. Di Martino A, O'Connor D, Chen B, Alaerts K, Anderson JS, Assaf M, et al. Enhancing studies of the connectome in autism using the autism brain imaging data exchange II. Sci Data. 2017 Mar 14;4:170010.

35. Di Martino A, Yan C-G, Li Q, Denio E, Castellanos FX, Alaerts K, et al. The autism brain imaging data exchange: towards a large-scale evaluation of the intrinsic brain architecture in autism. Mol Psychiatry. 2014 Jun;19(6):659–67.

36. Tustison NJ, Avants BB, Cook PA, Zheng Y, Egan A, Yushkevich PA, et al. N4ITK: improved N3 bias correction. IEEE Trans Med Imaging. 2010 Jun;29(6):1310–20.

37. Ad-Dab'bagh Y. E al. The CIVET Image-Processing Environment: A Fully Automated Comprehensive Pipeline for Anatomical Neuroimaging Research. In: Corbetta M, editor. In Proceedings of the 12th Annual Meeting of the Organization for Human Brain Mapping, [Internet]. Neuroimage; 2006. Available from: http://www.bic.mni.mcgill.ca/users/yaddab/Yasser-HBM2006-Poster.pdf

38. Lerch JP, Evans AC. Cortical thickness analysis examined through power analysis and a population simulation. Neuroimage. 2005 Jan 1;24(1):163–73.

39. Thompson PM, Stein JL, Medland SE, Hibar DP, Vasquez AA, Renteria ME, et al. The ENIGMA Consortium: Large-scale collaborative analyses of neuroimaging and genetic data. Brain Imaging Behav [Internet]. 2014; Available from: http://dx.doi.org/10.1007/s11682-013-9269-5

40. van Erp TGM, Hibar DP, Rasmussen JM, Glahn DC, Pearlson GD, Andreassen OA, et al. Subcortical brain volume abnormalities in 2028 individuals with schizophrenia and 2540 healthy controls via the ENIGMA consortium. Mol Psychiatry. 2016 Apr;21(4):547–53.

41. Borenstein M, Hedges LV, Higgins JPT, Rothstein HR. A basic introduction to fixed-effect and random-effects models for meta-analysis. Res Synth Methods. 2010 Apr;1(2):97–111.

42. Genovese CR, Lazar NA, Nichols T. Thresholding of statistical maps in functional neuroimaging using the false discovery rate. Neuroimage [Internet]. 2002; Available from: http://dx.doi.org/10.1006/nimg.2001.1037

43. Mazerolle M. Improving data analysis in herpetology: using Akaike's Information Criterion (AIC) to assess the strength of biological hypotheses. Amphib-reptil. 2006 Jan 1;27(2):169–80.

44. Alexander-Bloch AF, Shou H, Liu S, Satterthwaite TD, Glahn DC, Shinohara RT, et al. On testing for spatial correspondence between maps of human brain structure and function. Neuroimage. 2018 Sep;178:540–51.

45. Chun JJ, Shatz CJ. Interstitial cells of the adult neocortical white matter are the remnant of the early generated subplate neuron population. J Comp Neurol. 1989 Apr 22;282(4):555–69.

46. Eickhoff S, Walters NB, Schleicher A, Kril J, Egan GF, Zilles K, et al. High-resolution MRI reflects myeloarchitecture and cytoarchitecture of human cerebral cortex. Hum Brain Mapp. 2005 Mar;24(3):206–15.

47. Glasser MF, Van Essen DC. Mapping human cortical areas in vivo based on myelin content as revealed by T1- and T2-weighted MRI. J Neurosci. 2011 Aug 10;31(32):11597–616.

48. Richetto J, Chesters R, Cattaneo A, Labouesse MA, Gutierrez AMC, Wood TC, et al. Genome-Wide Transcriptional Profiling and Structural Magnetic Resonance Imaging in the

Maternal Immune Activation Model of Neurodevelopmental Disorders. Cereb Cortex. 2017 Jun 1;27(6):3397–413.

49. Lee H, Thacker S, Sarn N, Dutta R, Eng C. Constitutional mislocalization of Pten drives precocious maturation in oligodendrocytes and aberrant myelination in model of autism spectrum disorder. Transl Psychiatry. 2019 Jan 17;9(1):13.

50. Shen H-Y, Huang N, Reemmer J, Xiao L. Adenosine Actions on Oligodendroglia and Myelination in Autism Spectrum Disorder. Front Cell Neurosci. 2018 Dec 7;12:482.

51. Kikuchi M, Yoshimura Y, Hiraishi H, Munesue T, Hashimoto T, Tsubokawa T, et al. Reduced long-range functional connectivity in young children with autism spectrum disorder. Soc Cogn Affect Neurosci. 2015 Feb;10(2):248–54.

52. O'Reilly C, Lewis JD, Elsabbagh M. Is functional brain connectivity atypical in autism? A systematic review of EEG and MEG studies. PLoS One. 2017 May 3;12(5):e0175870.

53. Tomasi D, Volkow ND. Reduced Local and Increased Long-Range Functional Connectivity of the Thalamus in Autism Spectrum Disorder. Cereb Cortex. 2019 Feb 1;29(2):573–85.

54. Nair A, Treiber JM, Shukla DK, Shih P, Müller R-A. Impaired thalamocortical connectivity in autism spectrum disorder: a study of functional and anatomical connectivity. Brain. 2013 Jun;136(Pt 6):1942–55.

55. Alexander-Bloch A, Clasen L, Stockman M, Ronan L, Lalonde F, Giedd J, et al. Subtle in-scanner motion biases automated measurement of brain anatomy from in vivo MRI. Hum Brain Mapp [Internet]. 2016; Available from: http://dx.doi.org/10.1002/hbm.23180

56. Reuter M, Tisdall MD, Qureshi A, Buckner RL, van der Kouwe AJW, Fischl B. Head motion during MRI acquisition reduces gray matter volume and thickness estimates. Neuroimage [Internet]. 2015; Available from: http://dx.doi.org/10.1016/j.neuroimage.2014.12.006

57. Pardoe HR, Kucharsky Hiess R, Kuzniecky R. Motion and morphometry in clinical and nonclinical populations. Neuroimage. 2016 Jul 15;135:177–85.

58. Lombardo MV, Lai M-C, Baron-Cohen S. Big data approaches to decomposing heterogeneity across the autism spectrum. Mol Psychiatry [Internet]. 2019 Jan 7; Available from: http://dx.doi.org/10.1038/s41380-018-0321-0

59. Baxter AJ, Brugha TS, Erskine HE, Scheurer RW, Vos T, Scott JG. The epidemiology and global burden of autism spectrum disorders. Psychol Med. 2015 Feb;45(3):601–13.

60. Knutsen J, Crossman M, Perrin J, Shui A, Kuhlthau K. Sex differences in restricted repetitive behaviors and interests in children with autism spectrum disorder: An Autism Treatment Network study. Autism. 2019 May;23(4):858–68.

61. Mandy W, Chilvers R, Chowdhury U, Salter G, Seigal A, Skuse D. Sex differences in autism

spectrum disorder: evidence from a large sample of children and adolescents. J Autism Dev Disord. 2012 Jul;42(7):1304–13.

62. Retico A, Giuliano A, Tancredi R, Cosenza A, Apicella F, Narzisi A, et al. The effect of gender on the neuroanatomy of children with autism spectrum disorders: A support vector machine case-control study. Mol Autism [Internet]. 2016; Available from: http://dx.doi.org/10.1186/s13229-015-0067-3

63. Lazar M, Miles LM, Babb JS, Donaldson JB. Axonal deficits in young adults with High Functioning Autism and their impact on processing speed. Neuroimage Clin. 2014 Feb 7;4:417–25.

64. Ecker C, Ronan L, Feng Y, Daly E, Murphy C, Ginestet CE, et al. Intrinsic gray-matter connectivity of the brain in adults with autism spectrum disorder. Proc Natl Acad Sci U S A. 2013 Aug 6;110(32):13222–7.

65. Ecker C, Suckling J, Deoni SC, Lombardo MV, Bullmore ET, Baron-Cohen S, et al. Brain anatomy and its relationship to behavior in adults with autism spectrum disorder: a multicenter magnetic resonance imaging study. Arch Gen Psychiatry. 2012 Feb;69(2):195–209.

66. Tullo S, Patel R, Devenyi GA, Salaciak A, Bedford SA, Farzin S, et al. MR-based age-related effects on the striatum, globus pallidus, and thalamus in healthy individuals across the adult lifespan. Hum Brain Mapp. 2019 Aug;

67. Ellegood J, Anagnostou E, Babineau BA, Crawley JN, Lin L, Genestine M, et al. Clustering autism: using neuroanatomical differences in 26 mouse models to gain insight into the heterogeneity. Mol Psychiatry. 2015 Feb;20(1):118–25.

68. Natu VS, Gomez J, Barnett M, Jeska B, Kirilina E, Jaeger C, et al. Apparent thinning of visual cortex during childhood is associated with myelination, not pruning. bioRxiv [Internet]. 2018; Available from: http://dx.doi.org/10.1101/368274

69. Stüber C, Morawski M, Schäfer A, Labadie C, Wähnert M, Leuze C, et al. Myelin and iron concentration in the human brain: a quantitative study of MRI contrast. Neuroimage. 2014 Jun;93 Pt 1:95–106.

70. Uddin LQ. Idiosyncratic connectivity in autism: developmental and anatomical considerations. Trends Neurosci. 2015 May;38(5):261–3.

71. Weisstein EW. Sigmoid Function. http://mathworld.wolfram.com/ [Internet]. [cited 2019 Jul 22]; Available from: http://mathworld.wolfram.com/SigmoidFunction.html