GRIDSS2: harnessing the power of phasing and single breakends in somatic structural variant detection

Daniel L. Cameron[1,2,5], Jonathan Baber[2,3], Charles Shale[2,3], Jose Espejo Valle-Inclan[4], Nicolle Besselink[4], Edwin Cuppen[3,4], Peter Priestley[2,3], Anthony T. Papenfuss[1,5,6,7]

[1] Bioinformatics Division, Walter and Eliza Hall Institute of Medical Research, Parkville, Australia
[2] Hartwig Medical Foundation Australia, Sydney, Australia
[3] Hartwig Medical Foundation, Science Park 408, Amsterdam, The Netherlands
[4] Center for Molecular Medicine and Oncode Institute, University Medical Center Utrecht, Heidelberglaan 100, Utrecht, The Netherlands
[5] Department of Medical Biology, University of Melbourne, Australia
[6] Peter MacCallum Cancer Centre, Melbourne, Australia
[7] Sir Peter MacCallum Department of Oncology, University of Melbourne, Australia

Correspondence: Anthony T. Papenfuss (papenfuss@wehi.edu.au) and Daniel Cameron (cameron.d@wehi.edu.au)

Abstract

Here we present GRIDSS2, a general purpose structural variant caller optimised for tumour/normal somatic calling. Using cell line, patient sample validation and cohort-level comparisons, we show GRIDSS2 outperforms recent state-of-the-art tools. We demonstrate GRIDSS2 retains high sensitivity and precision even for small events by identifying a small (32-100bp) duplication signature strongly associated with colorectal cancer using 3,782 metastatic cancers that have been deeply sequenced by the Hartwig Medical Foundation. Essential to the high precision achieved by GRIDSS2 is the novel reporting of single breakend variants: structural variants in which only one side can be unambiguously determined. We show that the inclusion of single breakends reduces the false negative rate from 10.4% to 3.4%. Demonstrating the power single breakend calling has in genomic regions traditionally considered inaccessible to short read callers, we find that 47% of somatic centromeric breaks are repaired to non-centromeric sequence, with chromosome 1 exhibiting a unique centromeric rearrangement signature. Finally, we show that somatic structural variants are highly clustered with GRIDSS2 able to phase 16% of somatic structural variants in the Hartwig cohort from short read sequencing alone.

Introduction

The reliable detection of structural variants (SVs) is critical to the understanding of the role genome architecture plays in health and disease. This is especially important in cancer and precision medicine where structural variation can be a key driver mutation [1,2]. Over the past decade, many tools have been developed for the detection of genomic rearrangements, which have been the subject of extensive recent benchmarks [3,4]. These tools fall broadly into two camps: those that detect changes in DNA abundance, known as copy number variant or aberration (CNV/CNA) callers, and those that detect non-reference DNA adjacencies, known as structural variant (SV) or breakpoint callers. Whilst CNAs and SVs are merely two different viewpoints of the underlying genomic rearrangements, the methods of detection are fundamentally different. Here, we address the problem of SV detection using short read sequencing data, specifically in the detection of somatic SVs in paired tumour/normal samples.

Short read-based SV detection algorithms identify breakpoints by finding clusters of reads that do not support the reference allele. Typically these use discordant read pairs [5], or split reads[6], with some callers also considering reads with unmapped mates [7] and soft-clipped reads [8]. More sophisticated callers incorporate assembly either through de novo assembly [9], targeted breakpoint assembly [10], or breakend assembly [11]. These callers report breakpoints, that is, novel adjacencies. When reads cannot be unambiguously mapped on either side, a breakpoint call cannot be made. Some callers have attempted to address this by considering multiple alignment locations for each read [12] but this only works for regions with a small number of potential alignment locations. Such callers have not gained widespread adoption due to the computational cost of multi-mapping alignment and variant calling and the complexity of downstream interpretation.

The Variant Call Format (VCF)[13] defines a single breakend as a breakpoint in which only one side is observed. Single breakends occur when one side of the breakpoint cannot be placed, either due to the presence of multiple potential reference locations, or an adjacency with non-reference sequence (such as retroviral integration). By reporting single breakends in addition to breakpoints, structural variants involving repetitive sequence can be identified, which can contribute to improving short read caller sensitivity above the 50% reported in recent benchmarking[3,4,13]. Additional potential uses of this capability include viral integration detection [14,15] and improved single nucleotide accurate copy number segmentation.

As we move closer to a world in which CNAs and SVs primitives can be reliably detected, accurate interpretation of the causative biological events becomes increasingly possible. Whilst progress has been made on derivative chromosome reconstruction using long reads [16], reconstruction of complex events such as chromothripsis has been problematic for short reads [17,18]. To date, SV phasing has been used to reduce the complexity of reconstruction for long read based approaches [19] but has not been done by short read callers. The ability of phase somatic structural variants is limited by the read length and, for short read data, by the library fragment size - typically less than 500bp. Furthermore, the paired-end nature of short read data presents a unique challenge: transitive calls.

Another factor confounding high-level interpretation of short read based SV calls is the existence of transitive calls. DNA fragments shorter than the library fragment size can have read

pairs fully spanning the fragment. These will result in the reporting of breakpoints between the sequencing flanking either side of the short fragment. These transitive calls are particularly problematic for event interpretation and karyotype reconstruction as, had they been real, these SVs must have occurred on different derivative chromosomes than the true SVs involving the short DNA fragment. This problem can be resolved by transitive call reduction once the underlying breakpoints are identified. Although transitive call reduction has been utilised downstream of calling [20], it has not previously been incorporated into a SV caller, nor utilised to phase SVs.

Here, we present GRIDSS2, a multi-sample general-purpose structural variant caller specifically tuned for tumour/normal somatic calling. GRIDSS2 has the unique abilities to phase structural variants and report single breakends. We show favourable benchmarking results on cell line, patient sample and cohort data. Running GRIDSS2 on 3,782 metastatic solid tumours with matched normal samples from the Hartwig Medical Foundation cohort, report a somatic small (32-100bp) duplication signature; demonstrate the utility of phasing of somatic structural variants using short read data; and use single breakend variant calls to provide insight into centromeric rearrangements typically considered inaccessible even to multi-mapping aware short read callers. GRIDSS2 lays the groundwork for the development of downstream tools that enable a deeper understanding of the nature of genomic rearrangements.

Results

GRIDSS2 utilises the same high-level approach as the first version of GRIDSS, assembling all reads that potentially support a structural variant using a positional de Bruijn graph algorithm[11]. Breakend contigs are then realigned back to the reference to identify breakpoints and probabilistic structural variant calling is performed based on both the aligned reads and assembled contigs. Single breakend variant calling uses the same probabilistic variant calling approach as breakpoint calling, but instead of split reads, discordant read pairs, and assembly contigs with chimeric alignments support, single breakends are called based on soft-clipped reads, reads with unmapped or ambiguously mapping mates, and assemblies with unmapped or ambiguously mapping breakend sequence (Figure 1a). SV phasing is performed based on assembly contigs, and the presence of transitive calls (Figure 1b). SVs are phased cis if an assembly spans both breaks, or a transitive call is found, and phased trans if an assembly involves one SV but supports the reference at the other. Since assembly contig length is limited by the library fragment size, only nearby SVs can be phased. GRIDSS2 includes a 16-step somatic filter specifically tuned for deeply sequenced tumour/normal samples. Full details can be found in the Online Methods.

a)

Assemble

- Discordant read pairs
- Soft-clipped reads
- Split reads
- Reads with unmapped mates
- Indel-containing reads

Breakend assembly of fragments
sharing anchoring alignment

Align to reference

Single alignment

Zero | alignments
Multiple | alignments

Split alignment

Breakpoint call

Single breakend call

Phased breakpoints

Annotate unplaced sequence with:
- candidate reference genome locations
- RepeatMasker repeat type and class
- viral sequence matches

b)

Spanning assemblies
phase adjacent calls cis

Imprecise transitive calls replaced by cis phasing of precise calls

Figure 1: GRIDSS overview. a) contigs are assembled from a single locus of reads mutually supporting the same putative break junction. If the other side cannot be uniquely determined, the contig supports a single breakend call at the break junction position. If different portions of the contig sequence uniquely align to different genomic loci, the assembly supports multiple cis phased breakpoints. b) Nearby structural variants will have discordant read pairs spanning across multiple breakpoints. These generate spurious transitive calls that are collapsed into the underlying breakpoints, phasing them cis.

Benchmarking performance

To estimate precision and sensitivity of GRIDSS2, we used a recently generated somatic SV truth set for the COLO829 melanoma cell line and the COLO829BL cell line, which was derived from a normal cell from the same individual, using a combination of Illumina, PacBio, Oxford Nanopore, and 10X Genomics sequencing, followed by targeted capture and PCR-based validations and manual curation (J.E.V., E.C., manuscript in preparation). To test sensitivity and reproducibility, we ran GRIDSS2, manta [10], svaba[10,21], and novobreak[10,21,22] on 3 independent 40x normal/100x tumour replicates of the COLO829BL/COLO829T normal/tumour cell lines. GRIDSS2 achieved an average sensitivity/precision of 94%/83% compared to 88%/52% for manta, 75%/11% for svaba and 70%/7% for novobreak (Figure 2a).

To evaluate performance at lower sequencing depths and sample purity, we use in-silico downsampling and mixing to simulate a matched normal at 40x and a 60x tumour sample at 8%-100% purity corresponding to 5x, 10x, 15x, 20x, 25x, 30x, 45x, 50x, and 60x effective tumour coverage. Above 10x effective tumour coverage GRIDSS2 achieved higher sensitivity and specificity. At 10x and below, GRIDSS2 retained higher precision, but at lower sensitivity (Figure 2a).

Figure 2: Somatic benchmarks. a) COLO829T/BL tumour and blood cell lines were sequenced in triplicate to 100x/40x. In-silico purity downsampling was performed at 40x normal, and 60x tumour coverage. Results are compared against a PCR validated somatic truth set generated from multiple sequencing technologies. b) GRIDSS2/manta validation results on 13 patient samples for 50bp+ events. c) GRIDSS2/Strelka validation results for 32-50bp events. d) False negative rate (FNR) inferred from the presence of SVs copy number transitions broken down by magnitude of copy number change for 60 PCAWG samples. Comparison is between GRIDSS2/PURPLE and the PCAWG consensus call set. e) Inferred FNR for 3,782 100x tumour samples from the Hartwig Medical Foundation cohort. Single breakend variant calling is crucial to the low FNR in this cohort. f) Per sample counts of 32-100bp somatic tandem duplications in the Hartwig cohort. These mutations are enriched in colorectal cancer and associated with ATM driver mutations. g) Size distribution of small (32-100bp) tandem duplications across the Hartwig cohort. This is a distinct signature not associated with microsatellite expansion.

Validation on patient samples

To further validate somatic performance, we performed independent validation of GRIDSS2 and manta breakpoint calls from 13 patient tumor samples from the Hartwig Medical Foundation cohort [2,10,21,22] with a high burden of structural variants. Since the default minimum reported event sizes of GRIDSS2 and Manta are 32 and 50bp respectively, we compared 32-50bp events to the short indel caller, Strelka [23]. We used a hybrid capture approach with target probes flanking and overlapping break-junctions to independently validate over 5,000 calls identified by any tool. 3,403 of 3,666 (93%) GRIDSS2 calls were validated compared to 2,685 of 4,299 (65%) for Manta (Figure 2b). Of the private Manta calls not found by GRIDSS2, just 230 of 1777 (13%) were validated compared to 836 of 1031 (81%) GRIDSS private calls. No imprecise (that is, not base-pair accurate) GRIDSS2 calls passed somatic filtering, whereas imprecise manta calls validated at a rate (40/288, 14%) similar to manta private calls. All validated imprecise manta calls were called by GRIDSS2 precisely. In the 32-50bp range, 329 of 343 (96%) of GRIDSS2 calls validated against 142 of 182 (78%) for Strelka (Figure 2c). 95% (219 of 232) of 32-50bp calls private to GRIDSS2 were validated, compared to 47% (35 of 74) for Strelka. Notably, GRIDSS finds many short duplications of 32-100 bases which are largely missed by both Strelka and Manta.

Somatic short duplication signature

In addition to reidentifying known kilobase and megabase length duplication signatures, we find a signature consisting of short non-microsatellite tandem duplications of less than 100 bases (Figure 2f). There is a median of just 4 such events per tumor across the cohort but they are enriched in a small number of samples, with the majority of enriched samples in colorectal cancers (Figure 2g). Enrichment of samples with 20 or more short duplications is positively

associated with colorectal cancer (q=1.7e-09) and driver mutations in PARK2 (q=0.0003) and ATM (q=0.008).

These short tandem duplications are too large to be reliably called by most somatic indel callers, but too short to be reliably called by many SV callers. In part this is due to the weak read pair signal due to the short variant length, but also since most callers do not report variants shorter than 50bp threshold used for variant databases such as dbVar. Popular callers such as lumpy [24] and delly [25] do not call duplications shorter than 100 and 300bp respectively [4], and no duplications shorter than 300bp were included in the PCAWG consensus call set[1].

Structural variant/copy number consistency

Structural variant and copy number calls are intrinsically related. Any breakpoint must have either a compensating breakpoint (for example, as with inversions), or a copy number change at that SV position. Using this principle, we can estimate a false negative rate (FNR) from the number of unexplained copy number transitions. To generate matching SV and copy number calls, we ran GRIDSS2 and PURPLE [2] on 60 samples from the PCAWG WGS cohort and compared results with the state-of-the-art PCAWG consensus call set[20]. Copy number transitions in or within 100kb of centromeres or a gap in the reference genome were excluded.

Across the 60 samples, GRIDSS2 identified breakpoints for 90.8% of copy number transitions and single breakends for a further 2.6% with an estimated 6.6% FNR. The PCAWG consensus call set identified breakpoints for 88.9% of copy number transitions (11% FNR). This reduction in FNR was present for both clonal and subclonal variants (Figure 2d).

To evaluate GRIDSS2 on high quality, deeply sequenced samples, GRIDSS2 and PURPLE were run on 3,782 40x normal/100x tumour samples from the Hartwig Medical Foundation cohort. Excluding those occurring within 1kb of a gap in the reference genome, 60,139 of the 2,168,386 copy number transitions in the Hartwig cohort (3.4%) lacked a corresponding GRIDSS SV (Figure 2e). Explaining 7.0% of copy number transitions, single breakend variant calls were essential for achieving such a low FNR.

Novel centromeric break signature

Although only one side of single breakend variant calls can be uniquely placed, the assembled sequencing flanking the break can be used to classify retroviral integration, mobile element transposition, rearrangements involving centromeric and telomeric sequence, and other events. In the Hartwig cohort, RepeatMasker annotation reveals that the majority of somatic single breakend calls are caused by LINE insertions, or rearrangements involving centromeric sequence (Figure 3a). In the case of LINE insertions, one or both break junctions of a LINE insertion event may be called as a single breakend due to both the presence of multiple identical L1HS and L1P1 repeats and the characteristic polyA tail sequence of mobile element insertions[26] that causes de Bruijn graph assembly loops and truncation.

91% of the Hartwig cohort samples contain at least one copy number transition occurring in centromeric sequence. Being able to resolve the partners of the centromeric breaks explaining these copy number changes is critical to the accurate reconstruction of the derivative chromosomes. Single breakends into ALR/Alpha and HSATII centromeric repeats are able to give significant insight into the nature of these centromeric breaks. As each human centromere has a slightly different dominant repeat sequence, a mapping between each centromeric single breakend and their most likely centromeric breakpoint partner is possible. To do this, we aligned the single breakend sequences containing a centromeric or peri-centromeric repeat against the hg38 reference genome using BLAT, annotating each with the most likely centromeric partner. Using this approach, we were able to explain 5,614 of 11,996 (47%) centromeric copy number changes, implying that approximately half of centromeric rearrangements are centromere to centromere, and the remainder centromere to non-centromeric sequence. Of the 21,587 centromeric single breakends detected 3,148 (15%) had no copy centromeric copy number change, 6,850 (32%) had no copy number change but had multiple single breakends linked to the same chromosome, 3,358 (16%) had a single breakend associated with a centromere with copy number change, and the remaining 8,231 (38%) associated with a centromere with copy number change with multiple breakends mapping to that centromere in that sample.

Figure 3 classification of single breakends. a) RepeatMasker annotations indicate the majority of somatic single breakends are due to LINE translocations, or centromeric breaks. b) Most likely centromere for single breakends containing centromeric or peri-centromeric repeats based on realignment of breakend sequence to hg38. Shading indicates whether prediction is consistent with the copy number change across the centromere. Chromosome 1 has an excess of inter-chromosomal breaks to centromeric sequence. Chromosomes 13, 14, 15, 21, 22 have insufficient non-gap p-arm sequence for a centomeric copy number change to be called. c) Location of single breakends to centromeric sequence and corresponding centromere. Chromosome 1 has an excess of inter-chromosomal breaks to centromeric sequence, particularly to 5 and 19. d) Location of single breakends connected to centromeric sequence on the same chromosome. Red events left of the chromosomes are intra-chromosomal, and yellow events to the right are inter-chromosomal.

The centromeric single breakend rate can be further broken down by chromosome (Figure 3b), and based on the location of the single breakend (Figure 3c). Chromosome 1 is a clear outlier with an overabundance of centromeric inter-chromosomal rearrangements, particularly to chromosomes 5 and 19. Although the high level of sequence similarity between the centromeres of 1, 5, and 19 [27] could be a cause of false positive predictions, this relationship holds even when restricting the analysis to single breakends with an associated centromeric copy number change (Supplementary Figure 1), implying that the centromeric similarity between 1, 5 and 19, results in an increased rate of centromeric rearrangements between these chromosomes. In contrast, the lack of copy number supported single breakends to chromosome 13,14, 15, 21, and 22 centromeres is an artifact caused by missing p arm copy number due to gaps in the reference genome. Similarly, the centromeric sequence homologies between 13, 14, 21, and 22 combined with the lack of confirmatory copy number support, make it difficult to determine how much of the high inter-chromosomal centromeric rearrangement of chromosome 13 is due to misattribution of rearrangements to other chromosomes.

In general, intra-chromosomal single breakends to centromeric sequences occur close to the centromere (Figure 3d), with this effect less pronounced for inter-chromosomal breaks. Chromosome 1 is enriched for inter-chromosomal breaks, particularly to chromosomes 5 and 19, with inter-chromosomal breaks from these chromosomes to the centromere on 1 (Supplementary Figure 2) occurring in a pattern similar to the intra-chromosomal breaks of other chromosomes.

Somatic phasing

The breakend assembly approach taken by GRIDSS2 also enables the assembly-based phasing of nearby variants. When two structural variants occur in close proximity, they can be phased as cis if the contig aligns across both, and trans if the contig aligning across one aligns to the reference sequence at the other (Figure 4a). Segments shorter than 30bp are not typically uniquely alignable by BWA and unaligned short DNA segments are treated as insert sequences of an SV connecting the longer flanking segments. Since breakend assembly contig lengths are

limited by the fragment size distribution of the DNA library sequenced, only nearby variants can be phased.

For the Hartwig Medical Foundation cohort, variants could be phased up to around 500 base pairs. We found that multiple nearby somatic structural variants are frequent, with 22% of all structural variants having an adjacent variant within 1,000bp. This is far in excess of the 0.02% expected if the breakpoints were uniformly randomly distributed (Figure 4c). Of these, GRIDSS2 was able to phase 70% (Figure 4b) with 72% cis and 28% trans. This phasing information greatly assists downstream derivative chromosome reconstruction, as it exponentially reduces the number of possible paths through the breakpoint graph.

Figure 4: Structural Variant Phasing. a) Phasing of structural variants can be determined when breakend assembly contigs span multiple breakpoints. b) The majority of variants within 600bp can be phased using breakend assembly. c) Somatic SVs are highly clustered with 22% of all SVs in the Hartwig cohort potentially involving a DNA fragment of 1kbp or less.

Discussion

Through cell line, patient validation, and cohort-level comparisons, we have shown GRIDSS2 has excellent somatic performance above 10x effective tumour coverage. Our identification of a small (32-100bp) duplication signature highlights the importance of robust software tested across a wide range of variant types and sizes. The presence of a signature overlapping the widely accepted but arbitrary 50bp threshold separating indels from structural variants brings into question the wisdom of continuing to use this threshold as the minimum reported event size for structural variants.

Single breakend variant calling represents a significant conceptual advancement in the treatment of genomic rearrangements. Even though only one side can be unambiguously mapped to the reference genome, sequence classification of the unmapped side produces useful results. The identification of frequent centromeric rearrangements demonstrates the utility single breakend variant calling has in regions of the genome traditionally considered inaccessible to short read sequencing. It is not just in repetitive sequence that single breakend variant calling has value. The Hartwig cohort results show that merely reporting the presence of a break junction on the mappable side of a breakpoint enables improved copy number segmentation. Single breakend variant calling also potentially simplifies viral integration detection without the need for special tools since these integration sites can be identified by aligning single breakend sequences to a viral database.

One biologically significant finding coming from GRIDSS2's ability to phase structural variants is the degree to which somatic structural variants are clustered. In the Hartwig cohort of metastatic solid tumours, 22% of somatic structural variants potentially involve DNA fragments of less than 1000bp with GRIDSS2 able to phase 70% of these. Long read sequencing is considered the gold standard for structural variant phasing and phasability is indeed better: 10kb long reads increase this theoretical phasability to 31%. The plethora of very short fragments in cancer genomes, combined with the high indel error rate of PacBio and ONT sequencing presents a current drawback of long read sequence: simple long read based SV detection approaches are likely to misidentity complex rearrangements involving short DNA fragments as inserted sequence, since alignment of short fragments of long reads is only possible if error correction is performed prior to alignment. On COLO829, we found 5 instances (of 67 true positives) in which GRIDSS was able to correctly place a short DNA fragment the long read caller reported breakpoint inserted sequence. Care must be taken when comparing or combining short and long read variant calls to ensure the different representations of the same event are reconciled and cis phased. GRIDSS2's ability to phase breakpoints involving short DNA fragments is of great utility to downstream rearrangement event classification and karyotype reconstruction as it exponentially reduces the number of possible paths through the breakpoint graph.

GRIDSS2 is a new state-of-the-art somatic structural variant caller. Single breakend variant calling enables a sensitivity and specificity unprecedented amongst Illumina-based somatic structural variant calling. The highly clustered nature of somatic SVs mean that short read sequencing is surprisingly competitive when it comes to phasing somatic variants, and that sophisticated analysis and interpretation of somatic genomic rearrangements does not necessarily require long read sequencing. Going forward, we expect more sophisticated variant calling and assembly methods will be developed that are better able to handle the noise present in low complexity sequences. As single breakends are a genomic rearrangement primitive similar to but distinct from breakpoints, the vast majority of existing software is not able to correctly handle these variants. Effective utilisation of single breakend variant calls will require the development of an ecosystem of tools that fully support the VCF single breakend specifications.

Methods

GRIDSS2 extends the GRIDSS [11]software suite with additional features, tools and capabilities. GRIDSS2 is composed of the following 5 phases: (i) preprocessing, (ii) assembly, (iii) variant calling, (iv) annotation, and (v) somatic filtering.

Preprocessing

GRIDSS2 takes one or more aligned samples in the SAM/BAM[28] file format. These files are pre-processed on a per-file basis and all reads supporting a structural variant are extracted, and all fields or tags referring to another record are corrected. Reads with chimeric alignments (i.e. split reads), reads with a soft or hard clipped alignment CIGAR of at least 5bp, read pairs in which only one read is mapped, and discordant read pairs are extracted. The library insert size distribution is estimated from the first 10,000,000 reads using picard tools (http://broadinstitute.github.io/picard) and read pairs considered discordant if they fall outside the 99.5% distribution of fragment size lengths. The clipped bases of soft clipped non-chimeric reads are realigned to the reference genome using bwa mem [29] and converted to a chimeric split read alignments if an alignment is found. Inconsistencies in the mate chromosome and position are corrected (since tools such as GATK indel realignment adjust read alignment positions without updating the mate record), hard clips converted to soft clips, the NM, SA, MC, MQ tags recalculated, and the R2 tag is populated. Improving performance over GRIDSS, GRIDSS2 performs this in a two-pass manner with samtools [28] used for name/coordinate sorting the output of the first/second pass respectively.
As with GRIDSS, reads with low alignment sequence entropy and reads with a mapping quality (mapq) less than 20 (c.f. mapq<10 GRIDSS) are treated as unmapped, soft-clipped reads with clipped sequence having high homology with standard adapter sequences are ignored, reads marked as duplicates, and regions above 50,000x (c.f. 10,000x) coverage are ignored. Read alignments containing an insertion or deletion under 5bp are considered consistent with the reference.

Assembly

GRIDSS2 uses the same genome-wide positional de Bruijn graph break-end assembler used by GRIDSS. Reads are split into kmers and associated positions based on the anchoring alignment: kmers from split reads must be assembled only with kmers at the positions inferred by the anchoring alignment, and kmers of unmapped mate reads are assembled at any position consistent with the library fragment size distribution and the anchoring read alignment position. For assembly purpose, split reads and indel alignments are considered multiple soft clipped alignments, and discordant read pairs are treated as multiple read pairs with one read aligned.

The output of the assembly is a set of 'soft-clipped' contigs with anchoring bases supporting the reference, and non-reference bases supporting a putative breakpoint at a given position. This contig is iteratively realigned to the reference using bwa mem and converted to a split read alignment. Assemblies longer than the 1.5x maximum fragment size distribution, as well as assemblies supporting the reference sequence are ignored. Assembly alignments with a mapq

of less than 20 are treated as unmapped. GRIDSS2 introduces a number of refinements to the assembly calling process.

Assembly support is tracked per base pair. Fragments are considered to support a breakpoint only if the fragment support spans at least one base pair beyond any breakpoint homology on both sides. This ensures that when a single contig spans both a germline indel and a somatic SV, the fragments originating from the matched normal sample will not be considered as supporting the somatic breakpoint. This also improves variant allele fraction calculations in regions of complex rearrangement.

GRIDSS2 performs compound realignment of the entire assembly contig. BWA is used to align the entire assembly contig. Assembly contig bases which are soft clipped in the primary alignment reported by BWA are fed back to BWA for realignment. This process is repeated until either all bases are aligned, or no alignment can be found for the remaining bases. Assembly contigs that do not overlap with the locus of origin of the assembly are filtered out. To ensure that valid assemblies are not unnecessarily filtered, GRIDSS 2 includes both reads of each fragment in the assembly, and up to 300bp of anchoring reference-supporting sequence is included in the assembled contig. The remaining contigs are treated as split read alignments. To rectify over-alignment of the primary alignment location in the presence of imperfect breakpoint microhomology, the bounds of each split are adjusted to minimise the edit distance to the reference. The originating alignment is tracked using OA SAM tag and contigs that do not partially align to the originating assembly location and strand are filtered.

gridss.SoftClippedToSplitReads invokes bwa, -L 0,0 is added to the command line to remove the soft-clipping alignment penalty. This prevents 1bp non-template inserted sequences being over-aligned and reported as clean breakpoints with a flanking SNV.

Worse-case assembly performance has been improved by adding an assembly graph path count threshold. Generating 3 assembly contigs with more than 50,000 alternative paths through the assembly graph without advancing the assembly window will flush the assembly window. The maximum assembly window size has been reduced by 2.5x and more aggressive assembly read downsampling in high coverage regions is performed.

The presence of a contig with at least three non-overlapping alignments results in the breakpoints supported by that assembly being phased cis. If the initially soft-clipped portion of an assembly realigns across one breakpoint but not another, these breakpoints are phased trans.

Variant Calling

Breakpoints are called using a probabilistic model based on the empirical distribution of CIGAR operators, the library fragment size distribution, and mapping rate. Each read/read pair is given a phred-scaled quality score based on the mapping quality and the probability of encountering that read/read pair given the library empirical distribution. Split reads and soft clipped reads use the distribution of soft clipping CIGAR operators. Discordant read pairs use the discordant

mapping mate if distal or the library fragment size distribution if falling within the range reported by Picard tools CollectInsertSizeMetrics. Reads with unmapped mates use the unmapped mate fragment mapping rate, and indels based on rate of alignments with insertion/deletion CIGAR elements of matching lengths. As with GRIDSS, split reads and breakpoint-supporting assemblies incorporate the mapping quality scores on both sides of the supported break.

The key novel feature of the GRIDSS2 variant calling processing is the reporting of single breakend variants. Single breakends variants are called based on supporting soft clipped reads, assembly contigs, and reads with unmapped mates. Single breakend calling uses the same two-pass approach as breakpoint calling with all maximum cliques first calculated, then evidence uniquely assigned to the highest scoring clique.

In addition to single breakend variant calling, the variant caller has been improved by: reducing the default minimum called event to 10bp; preferentially allocating reads to variants supported by an assembly containing the read; requiring two supporting fragments to call a variant; and excluding inversion-like breakpoints from the minimum variant size filter to prevent filtering of foldback inversions.

Annotation

GRIDSS2 provides a full per-sample breakdown of all supporting evidence for each variant through the following VCF INFO and FORMAT fields:

- AS, RAS, CAS: counts of assembly contigs supporting a breakpoint originating locally, from the other side of the breakpoint, and from another location respectively. CAS assemblies support multiple variants and provide linking information about those variants.

- ASSR, ASRP: total number of split/soft clipped/indel-containing reads, and discordant read pairs/reads with unmapped mate contributing to any breakpoint-supporting assembly contig at the breakpoint location. Note that read/read pairs that are assembled into a contig but whose interval of support does not span the breakpoint are not counted. The interval of support for a read/read pair is defined as the interval between the first and the last contig base for which that read/read pair contributed to the assembly.

- SR, RP, IC: counts of split reads and discordantly mapped read pairs, and indel-containing reads that directly support the breakpoint.

- BA: counts of assembly contigs support a single breakend at this position. Such contigs are aligned only to the local breakend with the breakend sequence either aligning ambiguously, or unable to be aligned to the reference genome by bwa.

- BASSR, BASRP: total number of split reads or soft clipped reads, and discordant read pairs or reads contributing to any breakend-supporting assembly contig at the variant location.

- BSC, BUM: counts of soft-clipped reads, and reads with unmapped mates at the variant location

- ASQ, RASQ, CASQ, SRQ, RPQ, IQ, BAQ, BSCQ, BUMQ: corresponding quality score contribution for the supporting evidence.

- QUAL, BQ: total contribution to the breakpoint/breakend quality score.

- BANRP, BANSR, BANRPQ, BANSRQ: counts of read pairs/split reads not supporting this breakpoint but assembled into a contig that supports this breakpoint and their corresponding assembly quality score contribution.

- REF/REFPAIR: count of reads/read pairs spanning the local variant position that support the reference allele. Only reads/read pairs that span across the breakpoint microhomology interval (if present) are counted.

- VF/BVF: count of unique fragments supporting the breakpoint/breakend. By tracking unique fragments supporting the variant, a more accurate variant allele fraction can be calculated. This approach prevents double-counting of discordantly mapped fragments for which one of the reads contains a split read alignment. A fragment can support a variant either directly through split read, soft clipped read or discordant alignment of a read pair, indirectly through incorporation of one or both of the constituent reads in an assembly supporting the variant, or both directly and indirectly.

- RF: count of unique fragments supporting the reference allele.

- CQ: variant quality score prior to evidence reallocation.

- BEALN: Potential alignment locations of breakend sequence as determined by *gridss.AnnotateInsertedSequence.*

- BEID, BEIDL, BEIDH: identifiers of assembly contigs and the corresponding local and remote alignment base offsets. Single breakend variants do not have a remote breakend, and only breakpoint variants include breakpoint-supporting assemblies. Variants containing the same BEID are phased cis.

- CIPOS: For IMPRECISE variants, CIPOS encodes the interval in which the breakpoint could occur and for precise variants, CIPOS encodes the homology interval.

- CIRPOS: corresponding CIPOS of the remote breakend.

- IHOMPOS: interval of inexact homology. A Smith-Waterman alignment of the breakpoint sequence against the reference sequence is performed at both breakends. The reference and breakpoint sequence are extended 300bp from the break on either side with the reference extended an additional 10bp to account for potential indels in the alignment. The homology length is the length that the sequence alignment could be extended from the common sequence into the breakpoint/reference sequence. Alignments containing a soft clip on the common sequence side are classified as alignment errors and ignored. The SSW library[30] is used for which we implemented a JNI wrapper. Alignment scored 1, -4, 6, 1 for match, mismatch, gap open, and gap extend respectively which correspond to bwa mem alignment scores.

- SC: CIGAR encoding of the anchoring bases that at least one read/read pair/assembly is aligned to and supports the variant. This is encoded as a CIGAR string with a match for each anchoring base that provides support for the variant call, XNX for the interval over which the breakpoint could occur (due to microhomology or an imprecise call), and a deletion CIGAR element for any intervals over which there is no support (such as a small flanking deletion). Variants with an anchoring SC 10bp further from the break than a nearby variant are considered to be phased trans.

- SB: Strand bias of the reads supporting the variant. 1 indicates that reads would be aligned to the positive strand if the reference was changed to the variant allele. 0 indicates that reads bases would be aligned to the negative strand if the reference was changed to the variant allele. Strand bias is calculated purely from supporting reads and exclude read pair support since these are intrinsically 100% strand bias. Note that reads both directly supporting the variant and supporting via assembly will be double-counted. Both breakpoint and breakend supporting reads are included.

- IMPRECISE, HOMLEN, HOMSEQ, PARID, EVENT, CIEND, END, and SVTYPE fields carry their usual meaning as per the VCF file format specifications.

- MQ, MQN, MQX, BMQ, BMQN, BMQX mean, min, and max MAPQ score of reads/assembly contigs providing breakpoint/breakend support.

After initial annotation, *gridss.AnnotateInsertedSequence* aligns any single breakend sequences or non-template inserted breakpoint sequences to an arbitrary reference genome and adds an annotation reporting the alignment location. Integrated viral sequence is identified by aligning to a reference of viral sequences. By default, the same reference as the input files were aligned to is used. If a RepeatMasker bed file generated by BedOps[31] rmsk2bed is supplied, inserted sequences will be annotated with the RepeatMasker class and type corresponding to the BEALN alignments.

Somatic filtering

By default, GRIDSS2 is a sensitive caller and reports all putative variants supported by at least two well-mapped reads. To generate a set of high and low confidence somatic call sets, a somatic filter was developed. Variants with 3% of the supporting reads originating from the normal, or deletion or duplication breakpoints under 1000bp that have any direct split read support in the normal, are hard filtered. Variants are classified as low confidence if any of the following conditions are met: breakend coverage of less than 8 fragments in the normal; allelic fraction of less than 0.5% in the tumour; imprecise variant call; breakend variants without an assembly containing at least one discordant read pair; single breakends with a poly-C or poly-G run of at least 16bp in the breakend sequence; deletion or duplication breakpoints under 1000bp with a split read strand bias of 0.95 or greater; breakpoints with a microhomology of over 50bp; breakpoints with an inexact microhomology of over 50bp which are not deletion or duplications under 1000bp; deletion or duplication breakpoints under 1000bp with no split read support either directly, or through assembly; breakpoints with no discordant read pair support (either directly, or via assembly) which are not deletions or duplications under 1000bp; deletion or duplication breakpoints under 1000bp that have any direct split read support in the normal; 100-800bp deletion breakpoints with an inexact microhomology length of 6bp or greater; inversion-like breakpoints 40bp or less that have at least 6bp of microhomology; deletion-like breakpoints under 1000bp whose length of sequence inserted at the breakpoint is within 5bp of the deletion length, except those whose edit distance to the deleted bases is at least 0.5 per base, and less than 0.2 per base to the reverse complement. Breakpoint variants are filtered if either breakend is filtered.

Somatic variants are panel-of-normal (PON) filtered if a match within 2bp was found in a panel of normals. The default hg19 was constructed from the 40x coverage WGS matched normals for 3,972 patients from the Hartwig cohort using the *gridss.GeneratePonBedpe* utility. If multiple samples for a patient existed, only the normal for the first sample was included in the PON. Variants were aggregated across samples using the default setting of ignoring the FILTER field, and excluding imprecise calls and breakpoints/single breakends with a QUAL score of less than 75/428.

Viral insertions are annotated using *gridss.AnnotateInsertedSequence*. Single breakend sequences and non-template inserted sequences that do not have an alignment to the reference genome were aligned to a set of human viral reference sequences. Viral reference sequences were obtained from the virus host database [32] and filtered to include only viruses associated with the homo sapiens taxid of 9606. The viral sequences were then masked using RepeatMasker with "-no_is -s -noint -norna -species human" parameters. Generation scripts can be found at https://github.com/hartwigmedical/scripts/tree/master/virus.

Assembly linking: pairs of breakpoints mutually supported by a common assembly contig were annotated as linked by assembly. For assembly contigs spanning more than 2 breakpoints, each adjacent pair was linked with a unique identifier to enable unambiguous traversal of the breakpoint graph.

Transitive linking: chains of precise breakpoint variants were phased trans if an imprecise spanning transitive breakpoint call could be found. To identify transitive calls, a breadth-first search over the breakpoint graph was performed. Variants were considered transitive if the start and end breakends overlapped the start and end breakpoints in a path of precise breakpoint calls. Paths were limited to 1,000bp and 4 segments, with each segment required to be at least 20bp in length. Paths could not self-intersect. To prevent exponential runtime in highly rearranged genomes, at most 100,000 paths and at most 1,000 paths per starting breakpoint were considered.

Simple inversion annotation: pairs of breakpoints with orientations consistent with a simple inversion were annotated as simple inversions if the matching breakends were within 35bp on both sides, no other simple event annotation could be applied, and fragments supporting the constituent variants differed by at most threefold.

Templated insertion annotation: breakend/breakpoint and breakend/breakend pairs were annotated as simple templated insertions if the breakends had opposite orientations, were within 35bp, no other simple event annotation could be applied, and fragments supporting the constituent variants differed by at most threefold.

Reciprocal translocation: breakpoint/breakpoint pairs were annotated as reciprocal translocations if the breakends on both sides had opposite orientations, were within 35bp, no other simple event annotation could be applied, and fragments supporting the constituent variants differed by at most threefold.

Equivalent: variants were annotated as equivalent if variants had a breakend within 5bp of each other and they shared a common breakend sequence. Breakend sequences were truncated to the length of the shorter sequence and were considered matching when the per-base edit distance between breakend sequences was 0.1 or less. For the purposes of this comparison, the nominal breakend sequence was used for single breakends, and the reference sequence of the partner breakend was used for breakpoint variants. For breakpoint variants, the length of the breakend sequence was the maximum of 20 bases, and the width of the interval over which the fragments supporting the partner breakend had anchoring alignments.

Finally, a quality filter was applied to breakpoint variants with a QUAL score of less than 350 and single breakend variants with a QUAL score of less than 1000. Variants linked to a variant passing the qual filter other than through equivalence were rescued from the quality filtering and were considered to have passed regardless of the actual variant quality score. For each input file, two output files were generated: a high confidence call set containing calls passing all filters and a low confidence call set containing all calls except those failing the normal support filter or short events with split read support in the normal.

Independent validation of SV calls

13 samples from the Hartwig Medical Foundation metastatic cancer cohort were selected for capture panel validation of the structural variant calls. Each variant called in GRIDSS2 was

compared with variants called from Manta (for variants longer than 50 bases) and/or Strelka (for variants from 32-50 bases in length) to determine if the variant is shared or private. Variants were marked as matching GRIDSS2 if the start and end chromosomes and orientation both matched and start end positions (including confidence intervals) were within 20 bases of each other.Hybrid capture probes were created for each of the shared and private variants. For each breakpoint variant 3 probes of 120 bases each were created: Two reference probes leading up to the breakends from either side, as well as another SV probe going through the structural variant with the break junction close to the middle of the probe. The reference probes were designed to end 20 bases before the start of each structural variant breakend. The SV probe consists of any insert sequence flanked by equal number of bases from each side of the structural variant. For each single breakend variant 2 probes were created: One reference probe as described above and one SV probe which includes no more than 60 bases of the insert sequence with the remainder coming from the reference leading to the break point.

Together, this created a total of 17,125 capture probes of 120 nt in length, targeting 5,821 break-junctions (see supplementary table 1) which were ordered as custom target capture probes from Twist Biosciences (catalog ID 100533). For each of the 13 samples, 50 ng input DNA was used for indexed library construction with enzymatic fragmentation (Twist kit catalog IDs 100253, 100255 and 100401) according to the manufacturer's protocol. A bead-based size selection was performed after PCR to remove the remaining larger fragments (>700bp). Multiplexed hybridization was performed using the Twist Hybridization (ID100254), Blockers (PN100856) and Wash Kits (PN100214, 100215, 100216)) using Dynabeads™ MyOne™ Streptavidin T1 (Invitrogen PN65604D) following standard Twist protocol. Enriched library molecules were amplified by PCR for 11 cycles and sequenced on the Illumina NextSeq500 2x 150bp High Output run according to manufacturer's standard protocol.

We created a set of predicted alternate contigs from the shared and private structural variant calls using the same technique from above for generating the (non-reference) SV probes and added these to the reference genome. We then mapped each of the reads from the capture panel output with BWA to a hybrid genome including both the GRCH37 reference genome and the novel alternate contig.

We assessed the viability of the probes by mapping each of the 120 base SV probes to the 2,000 base alternate contigs to determine its mapping quality. Of the 5,821 SV probes, 80 had 0 mapping quality and 5,377 had a perfect mapping quality of 60. Probes with a mapping quality of less than 20 were ignored as well as 77 micro-satellite probes that were unable to be unambiguously validated. Resultant BAM files were examined for evidence of the SV alternate contigs in the SV source sample BAM as well as the BAM files of each of the other samples as controls for systemic effects for each of the predicted variants. Specifically, the read depth on the alternate contig at the variant location was used to assess the validation status of the variant. SV calls were marked as validated if all the following criteria were met (and not validated otherwise):

- At least 2 reads were mapped to the alternate contig in the predicted sample

- The support rate for the alternate contig was significantly higher (Poisson model, p=0.001) in the predicted sample than the maximum of the other 13 samples

- <40 reads in total across all 13 control samples were mapped to the predicted alternate contig

Comparison to PCAWG

Copy number data was obtained as for the Hartwig cohort running PURPLE [2] with default settings. PCAWG consensus SV and CN calls were obtained from https://dcc.icgc.org/releases/PCAWG/consensus_sv, and https://dcc.icgc.org/releases/PCAWG/consensus_cnv. Copy number transitions were matched with structural variants with a 100kb margin for PCAWG calls, and a 0bp margin for GRIDSS2/PURPLE. Copy number transitions in or within 100kb of centromeres or a gap in the reference genome were excluded from analysis. Copy number transitions matched by both a single breakend and a breakpoint, were considered breakpoint matches.

Hartwig metastatic tumour cohort

GRIDSS2 was run on 3,782 paired tumour/normal samples from the Hartwig Medical Foundation cohort of metastatic solid cancers with a 32bp minimum event size. Samples were aligned with bwa against a GRCH37 reference genome containing only primary contigs. Single breakend RepeatMasker annotations were obtained by running *gridss.AnnotateInsertedSequence* against the UCSC GRCH37 (hg19) RepeatMasker track downloaded from http://hgdownload.cse.ucsc.edu/goldenpath/hg19/bigZips/hg19.fa.out.gz after converting to BED formart using bedops *rmsk2bed*.

Single breakends with a RepeatMasker annotation associated with centromeric or pericentromeric repeats were considered centromeric single breakends. The matching chromosome was considered to be the chromosome with a BLAT [33] score of at least 900 and at least 25 higher than the best alignment on a different chromosome when aligning against hg38.

Phasability of the Hartwig cohort was calculated by determining, for each break junction, the length of the DNA segment if it was phased with the first break junction encountered in the appropriate orientation. Known phasing information was ignored for this analysis. Expected phasability was calculated by simulating 3,782 randomly fragmented paired genomes with the same number of break junctions as the corresponding Hartwig sample.

COLO829 somatic benchmark

The COLO829T/COLO829BL cell lines (ATCC® CRL-1974™ and 1980™ respectively) were each sequenced three times to 100x/40x using the HMF workflow [2,4] and aligned against GRCH37 without alt contigs using BWA 0.7.15. GRIDSS 2.9.3, manta 1.5.0, svaba 1.1.0, and smufin 0.9.3 [34] were run with default parameters. Programs were allocated 8,16,16,20 cores

and 32, 32, 50, 500Gb of memory respectively. No smufin results were obtained in any replicate as smufin failed to complete in the 100,000 CPU hours/3 months wall time allocated.

Call matching was performed using the StructuralVariantAnnotation BioConductor package (DOI 10.18129/B9.bioc.StructuralVariantAnnotation). A 100bp matching margin was allowed around the break junction position. Tandem duplication calls matched with insertion calls if the size difference between the duplication and insertion was within 25bp. False positive calls under 50bp were filtered after matching so as not to penalise a caller reporting an event slightly larger than 50bp in the truth set, but slightly smaller than 50bp in the call set. If multiple calls in a call set matched a single truth set call, all except the highest QUAL call were ignored.

COLO829 truth set generation

The COLO829 somatic SV truthset was generated using an orthogonal sequencing strategy. We sequenced the COLO829BL and COLO829T cell lines using Illumina HiSeqX (ENA run accessions ERR2752449 and ERR2752450 for COLO829BL and COLO829T, respectively), Oxford Nanopore (ERR2752451 and ERR2752452), Pacific Biosciences (ERR2752447 and ERR2752448) and 10X genomics (ERR2820166 and ERR2820167). All data is grouped under ENA study accession PRJEB27698.

Raw data was analysed for structural variants using the following tools:

- Illumina data was mapped using BWA 0.7.5, SV calling was performed with GRIDSS 2.0.1 and somatic SVs were filtered using gridss_somatic_filter.R.

- Nanopore data was mapped using NGMLR 0.2.6 and SV calling was performed with both NanoSV 1.2.0 and Sniffles 1.0.8 separately for COLO829T and COLO829BL. All SVs were merged with an overlap window of 200 base pairs using SURVIVOR 1.0.6 and SVs not present in COLO829BL were kept.

- Pacbio data was mapped using minimap2 2.11-r797 and SVs were called using pbsv 2.1.0 in joint calling mode for COLO829T and COLO829BL. Only SVs with no evidence in COLO829BL were kept.

- 10X genomics data was processed using Longranger 2.2.2 with default settings for COLO829BL and somatic mode for COLO829T. SV calls for both cell lines were merged with an overlap window of 200 base pairs using SURVIVOR 1.0.6 and SVs not present in COLO829BL were kept.

Somatic SV calls for each technology were merged with an overlap window of 200 base pairs using SURVIVOR 1.0.6. and all candidate breakpoints were subjected to independent validation by targeted capture and/or PCR-based approaches. SVs detected with two or more techniques that failed in these validation experiments were curated by manual inspection of the mapped reads using IGV [35]. A total of 69 SVs were finally considered as true somatic SVs for COLO829T.

References

1. Li, Y. *et al.* Patterns of somatic structural variation in human cancer genomes. *Nature* vol. 578 112–121 (2020).

2. Priestley, P. *et al.* Pan-cancer whole-genome analyses of metastatic solid tumours. *Nature* **575**, 210–216 (2019).

3. Kosugi, S. *et al.* Comprehensive evaluation of structural variation detection algorithms for whole genome sequencing. *Genome Biol.* **20**, 117 (2019).

4. Cameron, D. L., Di Stefano, L. & Papenfuss, A. T. Comprehensive evaluation and characterisation of short read general-purpose structural variant calling software. *Nat. Commun.* **10**, 3240 (2019).

5. Fan, X., Abbott, T. E., Larson, D. & Chen, K. BreakDancer: Identification of Genomic Structural Variation from Paired-End Read Mapping. *Current Protocols in Bioinformatics* 15.6.1–15.6.11 (2014) doi:10.1002/0471250953.bi1506s45.

6. Schröder, J. *et al.* Socrates: identification of genomic rearrangements in tumour genomes by re-aligning soft clipped reads. *Bioinformatics* vol. 30 1064–1072 (2014).

7. Ye, K., Schulz, M. H., Long, Q., Apweiler, R. & Ning, Z. Pindel: a pattern growth approach to detect break points of large deletions and medium sized insertions from paired-end short reads. *Bioinformatics* vol. 25 2865–2871 (2009).

8. Wang, J. *et al.* CREST maps somatic structural variation in cancer genomes with base-pair resolution. *Nat. Methods* **8**, 652–654 (2011).

9. Liu, S. *et al.* Discovery, genotyping and characterization of structural variation and novel sequence at single nucleotide resolution from de novo genome assemblies on a population scale. *Gigascience* **4**, 64 (2015).

10. Chen, X. *et al.* Manta: rapid detection of structural variants and indels for germline and cancer sequencing applications. *Bioinformatics* **32**, 1220–1222 (2016).

11. Cameron, D. L. *et al.* GRIDSS: sensitive and specific genomic rearrangement detection using positional de Bruijn graph assembly. *Genome Res.* **27**, 2050–2060 (2017).

12. Sindi, S. S., Onal, S., Peng, L. C., Wu, H.-T. & Raphael, B. J. An integrative probabilistic model for identification of structural variation in sequencing data. *Genome Biol.* **13**, R22 (2012).

13. Danecek, P. *et al.* The variant call format and VCFtools. *Bioinformatics* **27**, 2156–2158 (2011).

14. Wang, Q., Jia, P. & Zhao, Z. VirusFinder: software for efficient and accurate detection of viruses and their integration sites in host genomes through next generation sequencing data. *PLoS One* **8**, e64465 (2013).

15. Hajirasouliha, I. *et al.* Detection and characterization of novel sequence insertions using paired-end next-generation sequencing. *Bioinformatics* **26**, 1277–1283 (2010).

16. Aganezov, S., Zban, I., Aksenov, V., Alexeev, N. & Schatz, M. C. Recovering rearranged cancer chromosomes from karyotype graphs. *BMC Bioinformatics* **20**, 641 (2019).

17. Baca, S. C. *et al.* Punctuated evolution of prostate cancer genomes. *Cell* **153**, 666–677 (2013).

18. Cortés-Ciriano, I. *et al.* Comprehensive analysis of chromothripsis in 2,658 human cancers using whole-genome sequencing. *Nat. Genet.* **52**, 331–341 (2020).

19. Cretu Stancu, M. *et al.* Mapping and phasing of structural variation in patient genomes using nanopore sequencing. *Nat. Commun.* **8**, 1326 (2017).

20. ICGC/TCGA Pan-Cancer Analysis of Whole Genomes Consortium. Pan-cancer analysis of whole genomes. *Nature* **578**, 82–93 (2020).

21. Wala, J. A. *et al.* SvABA: genome-wide detection of structural variants and indels by local assembly. *Genome Res.* **28**, 581–591 (2018).

22. Chong, Z. & Chen, K. Structural Variant Breakpoint Detection with novoBreak. *Methods Mol. Biol.* **1833**, 129–141 (2018).

23. Saunders, C. T. *et al.* Strelka: accurate somatic small-variant calling from sequenced tumor-normal sample pairs. *Bioinformatics* **28**, 1811–1817 (2012).

24. Layer, R. M., Chiang, C., Quinlan, A. R. & Hall, I. M. LUMPY: a probabilistic framework for structural variant discovery. *Genome Biol.* **15**, R84 (2014).

25. Rausch, T. *et al.* DELLY: structural variant discovery by integrated paired-end and split-read analysis. *Bioinformatics* **28**, i333–i339 (2012).

26. Tubio, J. M. C. *et al.* Mobile DNA in cancer. Extensive transduction of nonrepetitive DNA mediated by L1 retrotransposition in cancer genomes. *Science* **345**, 1251343 (2014).

27. Hayden, K. E. Human centromere genomics: now it's personal. *Chromosome Research* vol. 20 621–633 (2012).

28. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079 (2009).

29. Li, H. & Durbin, R. Fast and accurate long-read alignment with Burrows-Wheeler transform. *Bioinformatics* **26**, 589–595 (2010).

30. Zhao, M., Lee, W.-P., Garrison, E. P. & Marth, G. T. SSW library: an SIMD Smith-Waterman C/C++ library for use in genomic applications. *PLoS One* **8**, e82138 (2013).

31. Neph, S. *et al.* BEDOPS: high-performance genomic feature operations. *Bioinformatics* **28**, 1919–1920 (2012).

32. Mihara, T. *et al.* Linking Virus Genomes with Host Taxonomy. *Viruses* **8**, 66 (2016).

33. Kent, W. J. BLAT--the BLAST-like alignment tool. *Genome Res.* **12**, 656–664 (2002).

34. Moncunill, V. *et al.* Comprehensive characterization of complex structural variations in cancer by directly comparing genome sequence reads. *Nat. Biotechnol.* **32**, 1106–1112 (2014).

35. Robinson, J. T. *et al.* Integrative genomics viewer. *Nature Biotechnology* vol. 29 24–26 (2011).

36. Zook, J. M. *et al.* A robust benchmark for detection of germline large deletions and

Acknowledgements

S. E. de Garis for manuscript feedback and copyediting.

This publication and the underlying study have been made possible partly on the basis of the data that Hartwig Medical Foundation and the Center of Personalised Cancer Treatment (CPCT) have made available to the study.

Author Contributions

DLC designed and implemented GRIDSS2. DC, PP, JB, CS designed and performed dry lab experiments and analysis. JEV provided COLO829 golden reference data. NB performed independent break junction validation experiments. ATP, PP, EC designed and supervised experiments. DLC, ATP, PP, EC contributed to writing of the manuscript.

Competing Interests

The authors declare no competing financial interests.

Data access

GRIDSS2 is freely available as open source software from https://github.com/PapenfussLab/gridss under a GPLv3 license. Analysis scripts used to generate results are available from https://github.com/PapenfussLab/gridss/tree/master/scripts/gridss2_manuscript.

Hartwig cohort data was obtained from the Hartwig Medical Foundation (Data request DR-005). Standardized procedures and request forms for access to this data can be found at https://www.hartwigmedicalfoundation.nl/en.

Raw and analyzed data for the creation of the COLO829T/COLO829BL tumor/normal cell line pair structural variant truth set are available grouped under ENA study accession PRJEB27698. The COLO829 truth VCF is available from https://github.com/UMCUGenetics/COLO829_somaticSV.

Capture panel validations of 13 patient tumor samples are available under the controlled access dataset accession EGAD00001005525.

HG002 BAMs were downloaded from https://ftp-trace.ncbi.nlm.nih.gov/ReferenceSamples/giab/data/

Figures

Figure 1: GRIDSS overview. a) contigs are assembled from a single locus of reads mutually supporting the same putative break junction. If the other side cannot be uniquely determined, the contig supports a single breakend call at the break junction position. If different portions of the contig sequence uniquely align to different genomic loci, the assembly supports multiple cis phased breakpoints. b) Nearby structural variants will have discordant read pairs spanning across multiple breakpoints. These generate spurious transitive calls that are collapsed into the underlying breakpoints, phasing them cis.

Figure 2: Somatic benchmarks. a) COLO829T/BL tumour and blood cell lines were sequenced in triplicate to 100x/40x. In-silico purity downsampling was performed at 40x normal, and 60x tumour coverage. Results are compared against a PCR validated somatic truth set generated from multiple sequencing technologies. b) GRIDSS2/manta validation results on 13 patient samples for 50bp+ events. c) GRIDSS2/Strelka validation results for 32-50bp events. d) False negative rate (FNR) inferred from the presence of SVs copy number transitions broken down by magnitude of copy number change for 60 PCAWG samples. Comparison is between GRIDSS2/PURPLE and the PCAWG consensus call set. e) Inferred FNR for 3,782 100x tumour samples from the Hartwig Medical Foundation cohort. Single breakend variant calling is crucial to the low FNR in this cohort. f) Per sample counts of 32-100bp somatic tandem duplications in the Hartwig cohort. These mutations are enriched in colorectal cancer and associated with ATM driver mutations. g) Size distribution of small (32-100bp) tandem duplications across the Hartwig cohort. This is a distinct signature not associated with microsatellite expansion.

Figure 3 classification of single breakends. a) RepeatMasker annotations indicate the majority of somatic single breakends are due to LINE translocations, or centromeric breaks. b) Most likely centromere for single breakends containing centromeric or peri-centromeric repeats based on realignment of breakend sequence to hg38. Shading indicates whether prediction is consistent with the copy number change across the centromere. Chromosome 1 has an excess of inter-chromosomal breaks to centromeric sequence. Chromosomes 13, 14, 15, 21, 22 have insufficient non-gap p-arm sequence for a centomeric copy number change to be called. c) Location of single breakends to centromeric sequence and corresponding centromere. Chromosome 1 has an excess of inter-chromosomal breaks to centromeric sequence, particularly to 5 and 19. d) Location of single breakends connected to centromeric sequence on the same chromosome. Red events left of the chromosomes are intra-chromosomal, and yellow events to the right are inter-chromosomal.

Figure 4: Structural Variant Phasing. a) Phasing of structural variants can be determined when breakend assembly contigs span multiple breakpoints. b) The majority of variants within 600bp can be phased using breakend assembly. c) Somatic SVs are highly clustered with 22% of all SVs in the Hartwig cohort potentially involving a DNA fragment of 1kbp or less.

Supplementary Figures

Supplementary Figure 1: Heatmap of single breakends to centromeric sequence by chromosomal arm. The x axis indicates the arm of the single breakend with the y axis indicating the most likely chromosomal arm the single breakend is connected to based on the breakend sequence, and the copy number profile across the centromere. ? indicates the arm is unknown due to the lack of copy number change across the centromere, and p and q indicate a centromeric copy number gain to that arm.

Supplementary Figure 2: location of single breakends associated with the chromosome 1 centromere. Red indicates a single breakend associated with centromeric sequence on the same chromosome, yellow indicates a single breakend associated with centromeric sequence on chromosome 1. Single breakends to chromosome 1 occurring on chromosomes 5 and 19 follow a similar distribution to intra-chromosomal single breakends.