DNMT3A haploinsufficiency results in behavioral deficits and global epigenomic dysregulation shared across neurodevelopment disorders

Diana L. Christian[1,6], Dennis Y. Wu[1,6], Jenna R. Martin[1], J. Russell Moore[1], Yiran R. Liu[1], Adam W. Clemens[1], Sabin A. Nettles[1], Nicole M. Kirkland[2], Cheryl A. Hill[2], David F. Wozniak[3,5], Joseph D. Dougherty[3,4,5], Harrison W. Gabel[1,*]

[1]Department of Neuroscience, Washington University School of Medicine, St. Louis, MO 63110-1093, USA

[2]Department of Pathology and Anatomical Science, University of Missouri School of Medicine, Columbia, MO 65212, USA

[3]Department of Psychiatry, Washington University School of Medicine, St. Louis, MO 63110-1093, USA

[4]Department of Genetics, Washington University School of Medicine, St. Louis, MO 63110-1093, USA

[5.]Intellectual and Developmental Disabilities Research Center, Washington University School of Medicine, St. Louis, MO, USA.

[6]These authors contributed equally: Diana L. Christian, Dennis Y. Wu.

[*]Email: gabelh@wustl.edu

Summary

Mutations in DNA methyltransferase 3A (DNMT3A) have been detected in autism and related disorders, but how these mutations disrupt nervous system function is unknown. Here we define the effects of neurodevelopmental disease-associated DNMT3A mutations. We show that diverse mutations affect different aspects of protein activity yet lead to shared deficiencies in neuronal DNA methylation. Heterozygous DNMT3A knockout mice mimicking DNMT3A disruption in disease display growth and behavioral alterations consistent with human phenotypes. Strikingly, in these mice we detect global disruption of neuron-enriched non-CG DNA methylation, a binding site for the Rett syndrome protein MeCP2. Loss of this methylation leads to enhancer and gene dysregulation that overlaps with models of Rett syndrome and autism. These findings define effects of DNMT3A haploinsufficiency in the brain and uncover

30 disruption of the non-CG methylation pathway as a convergence point across

31 neurodevelopmental disorders.

Introduction

Precise regulation of transcription through epigenetic mechanisms is critical for nervous

system development (Cholewa-Waclaw et al., 2016). Exome sequencing studies have revealed

mutations of genes encoding epigenetic modifiers of chromatin structure as a major underlying

cause of neurodevelopmental diseases (NDD), including autism spectrum disorder (ASD) (McRae

et al., 2017; Sanders et al., 2015; Satterstrom et al., 2019). A challenge emerging from these

discoveries is to define the cellular functions of the disrupted proteins during normal

development and search for shared pathways between these proteins that can potentially be

targeted for therapeutic development.

Gene regulation mediated by DNA methylation has emerged as an epigenetic mechanism

that plays a critical role in nervous system function (Kinde et al., 2015). In addition to the classical

methylation of cytosines found at CG dinucleotides (mCG), neurons contain uniquely high levels

of methyl-cytosine (mC) in a non-CG context, with this mark occurring primarily at CA

dinucleotides (mCA) (Guo et al., 2014; Lister et al., 2013a; Xie et al., 2012). mCA is deposited *de*

novo through the activation of the DNA methyltransferase 3A (DNMT3A) enzyme during the early

postnatal period (1-6 weeks of age in mice). Levels of mCA increase specifically in neurons until

the number of methylation sites in the non-CG context are nearly equivalent to mCG sites (Guo

et al., 2014; Lister et al., 2013a; Xie et al., 2012). A critical function for mCA is to serve as a binding

site for a neuron-enriched chromatin protein, Methyl-CpG binding Protein 2 (MeCP2) (Chen et

al., 2015; Gabel et al., 2015; Guo et al., 2014). MeCP2 was initially defined by its high affinity for

mCG, but biochemical and genomic studies indicate that it preferentially interacts with mCA to

down-regulate transcription of genes with essential functions in the brain (Boxer et al., 2019;

Gabel et al., 2015; Kinde et al., 2016; Lagger et al., 2017; Lyst and Bird, 2015). Loss of MeCP2

leads to the severe neurological disorder Rett syndrome, while duplication causes MeCP2-

duplication syndrome, an ASD, suggesting that read-out of mCA is critical to nervous system

function (Amir et al., 1999; Van Esch et al., 2005).

Notably, human exome sequencing studies have recently identified *de novo* mutations in

DNMT3A in individuals with ASD (Feliciano et al., 2019; Sanders et al., 2015; Satterstrom et al.,

2019). Separate studies have also defined heterozygous disruption of DNMT3A as the underlying

cause of Tatton-Brown Rahman syndrome (TBRS), a heterogeneous NDD characterized by

intellectual disability, overgrowth, craniofacial abnormalities, anxiety, and high penetrance of

ASD (Tatton-Brown et al., 2014, 2018). While a portion of the mutations identified in affected

individuals are truncations that are predicted to cause complete inactivation of the enzyme, a

majority of disease-associated alleles are missense mutations, raising questions about whether

loss-of-function effects are a primary mechanism of disruption in DNMT3A disorders (Tatton-

Brown et al., 2014, 2018). In addition, while heterozygous loss of DNMT3A has been studied in

the context of oncogenesis in the hematopoietic system (Cole et al., 2017), the effects of partial

loss of DNMT3A on nervous system function *in vivo* have not been examined; therefore, the

consequences of possible methylation changes on neuronal gene regulation and behavior are

unknown.

Here we examine the molecular effects of neurodevelopmental disease-associated

DNMT3A mutations and explore the consequences of heterozygous DNMT3A mutation on the

neuronal epigenome. Our results indicate that missense mutations across canonical domains of

DNMT3A disrupt different aspects of protein function, yet mutations in all domains reduce the

capacity of the enzyme to deposit neuronal mCA. We detect altered growth and behavior in

DNMT3A heterozygous deletion mice, supporting haploinsufficiency as a driver of pathology in

DNMT3A disorders. Through integrated epigenomic analysis, we reveal disruption of mCA

throughout the brain of DNMT3A mutant mice. Strikingly, we show that this loss of mCA leads to

disruption of distal regulatory enhancer activity and changes in gene expression that overlap with

models of MeCP2 disorders and other ASDs. These findings define the effects of NDD-associated

DNMT3A mutations for the first time and reveal disruption of mCA-mediated epigenomic

regulation as a convergence site across clinically distinct NDDs.

Results

Functional analysis of disease-associated DNMT3A mutations

Multiple DNMT3A mutations have been identified in individuals with ASD and TBRS.
However, the large number of missense mutations identified and the phenotypic heterogeneity
of individuals with these mutations raise the possibility that alterations of amino acids within
different protein domains may have distinct consequences and may dictate the nature and
severity of disease. We therefore sought to assess the effects of disease-associated DNMT3A
mutations on protein expression, cellular localization, and catalytic activity, looking for common
effects that may be core to the development of NDD.

We engineered amino-acid alterations homologous to human disease mutations into a
FLAG-tagged DNMT3A protein expression vector and assessed multiple mutations found within
each functional domain of the protein (Figure 1A). These analyses included mutations in the
chromatin interacting proline-tryptophan-tryptophan-proline (PWWP) domain, the auto-
inhibitory Histone H3 lysine 4 interacting ATRX-DNMT3-DNMT3L (ADD) domain, and the well-
defined methyltransferase catalytic domain (Gowher and Jeltsch, 2018). Transfection into
heterologous cells facilitated rapid assessment of protein expression by western blot, cellular
localization by immunocytochemistry, and catalytic activity using an *in vitro* methyltransferase
assay (Figure 1). Mutations in the PWWP domain resulted in a reduction in DNMT3A protein
expression and loss of nuclear localization compared to wild-type controls (Figure 1B-D, Figure
S1A-C). When expressed at equal levels to that of wild-type protein however, these mutations
exhibited substantial catalytic activity (Figure 1E,F). In contrast, mutations found in the catalytic
methyltransferase domain of DNMT3A showed wild-type expression and localization but
displayed deficits in catalytic activity in the *in vitro* methyltransferase analysis (Figure 1B-F, Figure
S1A-C). Mutations in the ADD domain of DNMT3A displayed normal protein localization and
expression levels and exhibited equal or higher methylation activity compared to wild-type
protein *in vitro* (Figure 1B-F, Figure S1A-C).

To further evaluate the functional effects of disease-associated DNMT3A mutations in the
context of endogenous chromatin, we tested the capacity of DNMT3A mutants to establish DNA
methylation across the genome in mouse cortical neurons. For this analysis, we focused on the
global build-up of mCA in postmitotic neurons that requires DNMT3A (Gabel et al., 2015; Lister
et al., 2013a). Cultured neurons isolated from the cerebral cortex at embryonic day 14.5

accumulate mCA *in vitro* and this build-up can be blocked by lentiviral-mediated delivery of Cre
recombinase to DNMT3A^{flx/flx} cells at 3 days *in vitro* (DIV) (Figure 2A, Figure S1D,E). We co-
transduced wild-type or mutant DNMT3A lentivirus at equal levels (Figure S1F) to test the
capacity of each protein to rescue deposition of DNA methylation. Analysis of mutations across
the major domains of DNMT3A detected deficits in mCA accumulation for all disease-associated
mutations tested (Figure 2B). Notably, mutations in the ADD domain that exhibited robust
catalytic activity *in vitro* displayed moderate-to-severe deficits in mCA deposition in neurons. The
ADD domain has been implicated in both histone binding and auto-inhibition of the protein (Guo
et al., 2015), thus effects in this neuronal assay may indicate that loss of ADD function blocks the
capacity of the enzyme to engage with chromatin and promote DNMT3A methylation activity in
cells. Together our results indicate that although NDD-associated mutations in DNMT3A affect
different protein domains and alter distinct aspects of protein function (e.g. localization,
chromatin interaction, catalysis), these mutations share a common outcome of reduced
enzymatic activity on neuronal DNA, with many mutations resulting in functionally null proteins.
131

***In vivo* effects of heterozygous DNMT3A disruption**
In light of our findings that multiple NDD-associated missense mutations in DNMT3A
result in complete or near-complete loss of function, we next sought to understand the effects
of heterozygous inactivation of DNMT3A *in vivo*. Previous studies have demonstrated severe
developmental deficits and perinatal lethality associated with complete loss of DNMT3A
(homozygous null mutation) in mice (Okano et al., 1999). However, the relevance of
heterozygous mutation of DNMT3A to neurodevelopmental disease has only recently been
uncovered and growth and behavioral effects of partial DNMT3A inactivation have not been
systematically assessed. We therefore carried out growth, behavioral, and molecular analyses of
mice carrying a constitutive heterozygous deletion of exon 19 of *Dnmt3a* (see *methods*) (Kaneda
et al., 2004). We find that this mutation leads to 50% reduction of RNA and protein expression,
allowing us to study the *in vivo* effects of heterozygous null mutation of DNMT3A (referred to as
DNMT3A^{KO/+}) (Figure S2A-C).

We first examined phenotypes with relevance to the overgrowth in individuals with
heterozygous DNMT3A mutations (Tatton-Brown et al., 2018), including enlarged body size and
obesity (body weight), tall stature (long-bone length), and macrocephaly (skull dimensions).
DNMT3A$^{KO/+}$ mice showed similar body weight to controls in the early postnatal period but were
significantly heavier than controls as mature adults (Figure 3A). This phenotype mimics a
maturity-associated trend toward increasing body weight observed in TBRS patients (Tatton-
Brown et al., 2018). Measurements of bone length indicated a small but significant increase in
tibia length in DNMT3A$^{KO/+}$ mice, with a trend towards longer femur length (Figure 3B, Figure
S3A-C). Morphometric analyses of the cranium and mandible indicated very subtle differences
between DNMT3A$^{KO/+}$ mice and their wild-type littermates (Figure S3D-F). One linear distance
spanning the rostrocaudal length of the interparietal bone is larger in DNMT3A$^{KO/+}$ mice relative
to wild-type littermates. Two linear distances in the facial region were significantly larger in wild-
type mice, while all other comparisons were not significantly different (Figure S3D). This suggests
very slight disruptions in growth of the facial region in DNMT3A$^{KO/+}$ mice. Together these findings
uncover effects on long bone length that mirror aspects of the human disorder, while skull
development in DNMT3A$^{KO/+}$ mice shows more limited effects. Additionally, enlarged body mass
in these mice appears to mimic overgrowth and obesity detected in individuals with TBRS
(Tatton-Brown et al., 2014).

To examine neurological and behavioral phenotypes in DNMT3A$^{KO/+}$ mice, we assessed
basic measures of sensation and motor performance such as balance (ledge test, platform test),
grip strength (inverted screen test), motor coordination (walking initiation, rotarod), and
sensorimotor gating (pre-pulse inhibition). DNMT3A$^{KO/+}$ mice were not significantly different in
these assays (Figure S4A-G), indicating that heterozygous loss of DNMT3A does not grossly
disrupt sensorimotor function. This allowed us to accurately assess more complex aspects of
behavior and cognition.

We carried out a panel of assays with relevance to neuropathology observed in humans
with DNMT3A mutations, including anxiety, autism, and intellectual disability. DNMT3A$^{KO/+}$ mice
displayed reduced exploratory behavior during open field testing, including reduced distance
traveled and rearing (Figure 3C-D). DNMT3A$^{KO/+}$ mice also displayed anxiety-like behaviors in this

assay, as they spent less time in the center of the open field arena (Figure 3E). In tests of climbing

behavior, DNMT3A$^{KO/+}$ mice showed longer latency to climb to the bottom of a pole and to the

top of mesh screens (Figure S4H-J), suggesting DNMT3A$^{KO/+}$ mice display differences in volitional

movement. To further assess anxiety-like behavior, we tested mice in the elevated plus maze and

observed that DNMT3A$^{KO/+}$ mice spent less time exploring the open arms of the maze with no

change in percent entries made into all arms (Figure 3F, Figure S4K). Overall, these results

demonstrate that the DNMT3A$^{KO/+}$ mice display changes in exploratory behavior, suggesting

altered emotionality and increased anxiety-like behaviors.

Analysis of common phenotypes examined in mouse models of autism (marble burying,

three chamber social approach, ultrasonic vocalizations) revealed additional changes in behavior.

We detected a significant reduction in marble burying activity for DNMT3A$^{KO/+}$ mice, indicating

alterations in repetitive digging behavior (Figure 3G). Testing of social interaction in the three

chamber social approach test for adult mice (Yang et al., 2011) detected trends toward similar

sociability and preference for social novelty in DNMT3A$^{KO/+}$ compared to controls (Figure S4L).

However, mutant mice showed reduced time investigating both mice and objects, as well as

reduced overall activity (Figure 3H,I, Figure S4M). These results may further reflect the overall

trend towards reduced exploration and anxiety-like phenotypes in these animals, instead of

changes in sociability (Nygaard et al., 2019). Reduced maternal-isolation induced ultrasonic

vocalizations (Barnes et al., 2017) were detected in DNMT3A$^{KO/+}$ mice at postnatal day five,

suggesting deficits in early pro-social behaviors or slight developmental delay in the normal

acquisition of this behavior (Figure 3J). Together these results indicate alterations in behaviors

commonly assessed in mouse models of autism (Chang et al., 2017; Simola and Granon, 2019;

Takumi et al., 2019), with our findings suggesting a reduction in activity and exploration, as well

as some changes in communication behaviors.

Intellectual disability is observed in patients with DNMT3A mutations, so we assessed

learning and memory in the DNMT3A$^{KO/+}$ mice using fear conditioning and Morris water maze

tests. The mutant mice displayed largely similar recall performance to that of control mice in

foot-shock induced fear conditioning, with similar levels of shock sensitivity (Figure 3K-M, Figure

S4N). However, DNMT3A$^{KO/+}$ mutants showed heightened freezing response during training, as

well as contextual and auditory recall phases of conditioned fear testing (Figure 3K-M). Mutant
mice also showed delayed extinction of freezing behavior in response to the auditory cue alone,
which may indicate altered emotionality or cognition (see *methods* for discussion). Assessment
of spatial and contextual memory by Morris water maze testing demonstrated that DNMT3A$^{KO/+}$
mice were slower to learn to find a visible platform and did not learn the location of the hidden
platform over time to the level of wild-type controls (Figure S4O-P). DNMT3A$^{KO/+}$ mice also
showed no differences in swimming speed compared to wild types (Figure S4Q-R). There were
no significant effects on distance traveled in target zone or platform crossings in the probe trial,
though DNMT3A$^{KO/+}$ mice trended towards fewer platform crossings (Figure S4S,T). These
findings suggest that DNMT3A$^{KO/+}$ mutants do not show frank deficits in learning and memory
but do display differences in task performance that further suggest altered emotionality or
cognition in these mice. Our analyses demonstrate that heterozygous deletion of DNMT3A
results in altered behavior in mice with relevance to anxiety and memory associated behaviors
observed in patients with DNMT3A mutations. These data support a model in which DNMT3A
haploinsufficiency can alter behavioral circuits to drive phenotypes in NDD.
218
Global disruption of DNA methylation in the DNMT3A$^{KO/+}$ brain
We next investigated the epigenomic defects that may underlie the altered behaviors
observed in DNMT3A$^{KO/+}$ mice. We first used sparse whole-genome bisulfite sequencing to
efficiently survey effects on global DNA methylation levels for multiple brain regions and liver
tissue isolated from wild type and DNMT3A$^{KO/+}$ mice. This analysis detected limited reductions in
genome-wide mCG levels in the DNMT3A$^{KO/+}$ brain that were not apparent in the liver, a non-
neural tissue (Figure 4A). In contrast, mCA levels were reduced by 30-50% across all brain regions
examined in DNMT3A$^{KO/+}$ mice (Figure 4B). DNA methylation across postnatal development in
the cerebral cortex, the brain region with the highest levels of mCA at 8 weeks, suggests that
deficits in mCA appear during initial accumulation of this methyl mark at 1-6 weeks (Figure 4B).
Thus, global mCA levels in the brain appear to be highly sensitive to heterozygous DNMT3A
disruption, while overall global mCG levels are largely maintained.

DNA methylation at specific genomic elements, including promoters, enhancers, and gene bodies is thought to play an important role in regulating transcription. Alterations in methylation at these regions can impact gene expression to affect the development and function of the brain (Clemens et al., 2019; Nord and West, 2019; Stroud et al., 2017a). We therefore assessed changes in methylation at base-pair resolution by high-depth whole-genome bisulfite sequencing to identify potential changes in mCA and mCG at these important regulatory sites. For this analysis we focused on the cerebral cortex, as this region is enriched for mCA (Figure 4B) and is disrupted in ASD and MeCP2 disorders (Clemens et al., 2019; Satterstrom et al., 2019; Sceniak et al., 2016; de la Torre-Ubieta et al., 2016).

High-resolution analysis of mCG confirmed the subtle reduction in mCG across all classes of genomic elements (Figure 5A,D). We considered that CG dinucleotides in specific sites in the neuronal genome may be more sensitive to a partial reduction in DNMT3A activity. For example, in the hematopoietic system, heterozygous disruption of DNMT3A leads to reductions in DNA methylation in genomic regions that can be identified as sensitive to complete loss of DNMT3A (Cole et al., 2017). We therefore evaluated developmentally-regulated adult-specific CG-differentially methylated regions (CG-DMRs) previously identified in the cortex (Figure S5A) (Lister et al., 2013a). Because DNMT3A is the only *de novo* methyltransferase expressed in the postnatal brain, we hypothesized that adult-specific CG-DMRs might be sensitive to a reduction in enzyme activity. Indeed, we found that these sites build up mCG during postnatal development and do not become methylated in a brain-specific DNMT3A conditional knockout mouse (DNMT3A cKO) (Stroud et al., 2017a) (Figure S5B). Analysis of adult-specific CG-DMRs in the DNMT3A$^{KO/+}$ model indicated that these sites are particularly sensitive to partial inactivation of DNMT3A compared to other regions genome wide (Figure 5A,C,D).

To further search for local sites of altered mCG in the DNMT3A$^{KO/+}$, we performed *de novo* calling of mCG differentially methylated regions using the BSmooth algorithm (Hansen et al., 2012). We identified 843 hypo- and 71 hyper-CG-DMRs across the genome that met high stringency filters for size and reproducibility (Figure 5A-C,E, Figure S5C, see *methods*). These hypo-DMRs significantly overlap with the previously identified adult-specific CG-DMRs (Lister et al., 2013a) (Figure 5F), further supporting the idea that DNMT3A is haploinsufficient for postnatal

mCG deposition at these sites. Examination of the genomic distribution of all DNMT3A[KO/+] CG-DMRs revealed significant overlap of hypo-DMRs with putative enhancer regions, gene bodies, and promoters (Figure 5A-C,F). DMRs were also highly enriched for overlap with CpG island shores, regions disrupted in studies of DNMT3A mutation outside of the nervous system (Cole et al., 2017; Spencer et al., 2017) (Figure 5F). Because a substantial percentage of mCG in neurons can occur in an oxidized, hydroxymethyl form (hmCG), we further performed oxidative bisulfite sequencing analysis of DNA from the cortex. This analysis revealed no clear evidence of differential effects on the oxidized or unoxidized forms of mCG across the genome in the DNMT3A[KO/+] (Figure S5D). Together these findings indicate that a small subset of mCG sites are particularly sensitive to heterozygous loss of DNMT3A. The localization of these CG-DMRs to regulatory elements suggests that these methylation changes could impact gene expression.

We next examined the profile of mCA at higher resolution, assessing genomic elements of different scales that have relevance to gene regulation. In contrast to the limited mCG changes in the DNMT3A[KO/+], analysis of mCA levels detected consistent 30-50% reductions at nearly all genomic regions examined (Figure 5G). This was true of gene bodies, promoters, and CpG island shores. CpG island sites, which show very low mCA levels in wild-type cortex, displayed less reduction of mCA, possibly due to floor effects in bisulfite-sequencing (see *methods*). Comparison of mCA changes in each class of genomic elements as a function of wild-type mCA levels suggested that consistent reductions occurred across the genome independent of the normal levels of mCA (Figure S5E). This suggests that changes in mCA levels in the DNMT3A[KO/+] do not preferentially impact specific classes of genomic elements or become substantially more severe in some regions based on the level of mCA that normally is deposited.

Recent analysis has demonstrated that topologically-associating domains (TADs) of chromatin folding are regions of organization for mCA that can impact gene regulation (Clemens et al., 2019). The "set-point" level of mCA within TADs is associated with the level of mCA at enhancers within TADs and high-mCA enhancers found in high-mCA TADs are particularly robust targets of repression by MeCP2 (Clemens et al., 2019). We therefore specifically assessed mCA levels at TADs and enhancers genome-wide. This analysis detected reductions in TAD mCA levels that were similar to global reductions in mCA at other genomic elements (Figure 5G). Enhancers

also showed this pervasive depletion of mCA (Figure 5G). Like other genomic elements, these

effects were consistent for TADs and enhancers with differing wild-type levels of mCA (Figure

S5E). Thus, widespread loss of mCA for TADs and enhancer elements occurs in DNMT3A[KO/+] mice

and has the potential to impact epigenetic control of regulatory elements by MeCP2.

Enhancer dysregulation results from methylation deficits in DNMT3A[KO/+] mice

We next examined how disruption of DNA methylation can affect epigenetic regulation in

DNMT3A[KO/+] neurons to alter gene expression and disrupt nervous system function. Recent

analysis indicates that mCA serves as a binding site for MeCP2 to mediate neuron-specific gene

regulation, in part by controlling the activity of distal regulatory enhancer elements (Clemens et

al., 2019). Loss of MeCP2 in mice leads to genome-wide upregulation of the activating mark

Histone H3 lysine 27 acetylation (H3K27ac) at enhancers that contain high levels of mCA and mCG

sites, while overexpression of MeCP2 leads to reciprocal downregulation of highly methylated

sites. Alterations in enhancer activity in MeCP2 mutants are linked to dysregulation of genes that

can then drive nervous system dysfunction. These findings suggest that reduced CA methylation

in the DNMT3A[KO/+] would remove binding sites for MeCP2 within enhancers. This mCA reduction

could then result in dysregulation of enhancer activity that partially phenocopies the effects we

have observed in MeCP2 mutant mice.

To investigate this possibility directly, we quantified the change in mCA binding sites in

the DNMT3A[KO/+] for enhancers significantly repressed by MeCP2 (Clemens et al., 2019). These

enhancers contain a large number of mCA sites due to high mCA/CA levels and an enrichment of

CA dinucleotides within these sequences. As a result, we found that the global 30-50% reduction

of mCA in the DNMT3A[KO/+] leads to a larger loss in the total number of mCA sites at MeCP2-

repressed enhancers than at other enhancers genome-wide (Figure 6A,B). Thus MeCP2-

repressed enhancers are particularly susceptible to mCA binding site loss from heterozygous

mutation of DNMT3A.

To determine if the reduction of mCA sites at MeCP2-repressed enhancers affects their

activity, we assessed changes in enhancer activation level by H3K27ac ChIP-seq analysis of the

DNMT3A[KO/+] and wild-type cerebral cortex. This analysis revealed significant changes in

acetylation at MeCP2-repressed enhancers (Figure 6A,C). Consistent with these effects arising

from 30-50% loss of the mCA that normally builds up post-mitotically at enhancers, we detect

changes that are concordant with, but smaller than, those caused by complete loss of post-

mitotic mCA in the DNMT3A cKO (Clemens et al., 2019) (Figure 6C).

Although significantly dysregulated enhancers can be detected in MeCP2 mutants, broad

sub-significance-threshold effects also occur genome-wide upon MeCP2 mutation, with nearly

all enhancers across the genome undergoing dysregulation that is proportional to the number of

mC binding sites at these regions (Figure 6D) (Clemens et al., 2019). Analysis of H3K27ac changes

at enhancers based on the normal density of mCA sites in these sequences genome-wide

revealed broad mCA-associated derepression of enhancers in DNMT3A$^{KO/+}$ cortex that is similar

to, but smaller in magnitude than, the effects observed in DNMT3A cKO and MeCP2 knockout

mice (MeCP2 KO). These effects are also reciprocal to effects observed in MeCP2 overexpression

mice (MeCP2 OE). Consistent with the limited disruption of mCG genome wide in the DNMT3A$^{KO/+}$

mice, there was more limited association between changes in enhancer activity and the level of

mCG at these sequences. This contrasts with MeCP2 mutants in which loss of protein binding at

both mCG and mCA sites leads to enhancer dysregulation that is associated with both mCA and

mCG (Clemens et al., 2019) (Figure 6D). Together, these findings demonstrate that loss of half of

the normal mCA sites in the DNMT3A$^{KO/+}$ cortex results in enhancer dysregulation that overlaps

with MeCP2 mutant mice, uncovering a role for shared neuronal chromatin pathology between

DNMT3A and MeCP2 disorders.

Overlapping transcriptional pathology between DNMT3A$^{KO/+}$, MeCP2 disorders, and ASD

The epigenetic alterations we observe in DNMT3A$^{KO/+}$ cerebral cortex can have direct

consequences on gene expression to drive neurological dysfunction in mice. Furthermore, the

overlapping effects on enhancers that we observe between DNMT3A$^{KO/+}$ and MeCP2 mutant

mice suggests that there may be shared transcriptional pathology occurring upon loss of mCA in

DNMT3A disorders and through disruption of MeCP2 in Rett syndrome and MeCP2-duplication

syndrome. We therefore assessed changes in gene expression in DNMT3A$^{KO/+}$ mice, interrogating

the extent to which these effects overlap with those observed in MeCP2 mutants and upon

complete disruption of mCA in the DNMT3A cKO. RNA-seq of DNMT3A[KO/+] cerebral cortex

identified subtle changes in gene expression that are consistent in magnitude with effects

observed in other heterozygous NDD models (Fazel Darbandi et al., 2018; Gompers et al., 2017;

Katayama et al., 2016) (Figure S6A). Gene set enrichment analysis on differential expression data

revealed dysregulation of Gene Ontology terms relating to neuronal development and function

(Figure S6B), suggesting that alterations in gene expression resulting from DNMT3A heterozygous

disruption could drive the behavioral alterations that we observed.

While a limited gene set is detected as significantly dysregulated in the DNMT3A[KO/+], we

considered if genome-wide alterations in enhancer activity could lead to wide-spread, subtle

dysregulation of gene expression that is below the threshold of detection for individual genes. In

this way, the transcriptional pathology in the DNMT3A[KO/+] brain could overlap with the

subthreshold genome-wide effects observed upon loss of neuronal mCA (DNMT3A cKO) and in

models of Rett syndrome (MeCP2 KO) and ASD (MeCP2 OE) (Clemens et al., 2019; Gabel et al.,

2015). Indeed, the significantly dysregulated genes in the DNMT3A[KO/+] overlapped extensively

with genes identified as significantly dysregulated in DNMT3A cKO and MeCP2 mutant mice

(Clemens et al., 2019), supporting the notion of shared gene expression effects between these

mouse models (Figure 6E). To more comprehensively assess the degree to which transcriptome-

wide changes in the DNMT3A[KO/+] phenocopy these MeCP2 mutant and DNMT3A cKO models, we

performed Generally Applicable Gene-set Enrichment (GAGE) analysis (Luo et al., 2009) of all

genes detected as dysregulated in these mutant models. This revealed highly significant,

concordant changes in gene expression in the DNMT3A[KO/+] for dysregulated gene sets detected

upon loss of mCA in the DNMT3A cKO and in MeCP2 mutant models (Figure 6F).

Having detected overlap in transcriptomic pathology between models of DNMT3A and

MeCP2 disorders, we sought to explore if shared gene expression signatures in the DNMT3A[KO/+]

mice extend to models of disorders that do not have as clear mechanistic links to DNMT3A

disorders. We therefore tested if DNMT3A[KO/+] mice show significant alterations in gene sets

identified as dysregulated in other mouse models of NDD and human gene sets implicated as

altered in the autistic brain. GAGE analysis across multiple datasets detected highly significant

dysregulation of gene sets identified in CHD8 and PTEN mouse models of overgrowth and ASD

(Gompers et al., 2017; Katayama et al., 2016; Tilot et al., 2016) as well as the SetD5 model of NDD

(Sessa et al., 2019) (Figure 7A). These findings support a role for overlapping gene dysregulation

underlying common symptomology found in affected individuals carrying mutations in distinct

genes.

Analysis of human gene sets detected as dysregulated in ASD postmortem brains (Gandal

et al., 2018; Voineagu et al., 2011) showed significant changes in the DNMT3A$^{KO/+}$ cortex (Figure

7B). This analysis also indicated upregulation of candidate genes linked to ASD from human

genetics studies (Abrahams et al., 2013; Banerjee-Basu and Packer, 2010) (Figure 7B). In addition,

analysis of co-expression modules of human brain development (Parikshak et al., 2013) showed

overlap with several neurodevelopmental modules including those that increase during early

cortical development and are enriched for ASD risk genes (M13, M16, and M17) (Figure S6C).

Modules involved in regulation of nucleic acids and gene regulation that are expressed early in

development and decrease over time are also increased upon heterozygous loss of DNMT3A (M2

and M3) (Figure S6C). These results indicate that important sets of genes with opposing

developmental trajectories and function are altered upon loss of DNMT3A regulation. Notably,

control resampling analysis indicated that significant dysregulation of these mouse and human

gene sets was not driven by enriched expression of these genes in the cortex (Figure S6D).

Together these findings suggest that the DNMT3A$^{KO/+}$ mouse shares overlapping transcriptional

pathology with gene expression changes underlying ASD.

Discussion

Our functional analysis of NDD-associated DNMT3A mutations together with our *in vivo*

studies provide an initial working model of molecular etiology in DNMT3A disorders. Diverse *de*

novo missense mutations that arise in affected individuals disrupt enzyme function by disabling

the capacity of the enzyme to localize to chromatin in the nucleus, altering the ADD-regulatory

domain, or disrupting the activity of the methyltransferase domain. Loss-of-function effects

resulting from these missense mutations, as well as early truncations or gene deletions, lead to

insufficient DNMT3A activity. This causes deficits in deposition of mCG at specific sites during

development and a massive deficit in postnatal mCA accumulation throughout the brain. These

changes in DNA methylation lead to alterations in epigenomic regulation, including subtle but

wide-spread disruption of mCA-MeCP2-mediated enhancer regulation in adult neurons, resulting

in gene expression changes that can drive deficits in nervous system function.

Our studies of DNMT3A mutations not only provide insight into the molecular etiology of

DNMT3A disorders, but also serve as a model for understanding the functional effects of diverse

de novo mutations underlying neurodevelopmental disorders. Exome sequencing studies have

identified a large and growing list of mutations in genes encoding epigenetic regulators in

individuals with NDD. Many of these are missense mutations and occur as heterozygous

disruptions (McRae et al., 2017; Satterstrom et al., 2019), leaving it unclear if simple loss-of-

function effects are sufficient to drive pathology through haploinsufficiency, or if more complex

effects play a role when individual amino acids are altered. In addition, while identification of

multiple mutations in a gene can implicate disruption of the gene as causative for NDD, it remains

possible that a subset of the mutations identified in affected individuals, particularly missense

mutations, are not in fact deleterious or causative. Functional testing of these variants is

therefore necessary to determine if they may underlie disease. Here, our analysis of DNMT3A

mutations in multiple functional assays has uncovered diverse mechanisms by which the protein

can be disrupted while pointing to a shared loss of function in the deposition of neuronal DNA

methylation. Notably, it is only by assessing multiple aspects of protein function (i.e. expression,

localization, activity, and cellular mCA levels) that we can detect deficits for each mutation tested.

For example, mutation of the ADD domain disrupts deposition of mCA, possibly due to loss of

regulation that can only be assessed in the endogenous chromatin context. Together our findings

establish the deleterious effects of diverse DNMT3A mutations and underscore the importance

of multidimensional analysis of *de novo* mutations to fully assess their potential role in NDD.

Our *in vivo* analyses show that heterozygous deletion of DNMT3A mirrors multiple key

features of DNMT3A disorders, including tall stature (increased long bone length), increased body

weight, and behavioral alterations. Detection of robust anxiety-like phenotypes in multiple

assays, deficits in pro-social communication, and alterations in repetitive behaviors align with

observed human phenotypes. In contrast, lack of strong deficits in learning and memory assays

in our mouse model may indicate that some regions and systems in humans are more susceptible

to DNMT3A disruption than in mice. However, we do detect alterations in behavior in these

assays (Figure 3K-M, Figure S4O-T) and the lack of strong deficits may also reflect insensitivity of

the methods used to measure specific aspects of disrupted cognition. In all, our *in vivo* analysis

indicates that heterozygous deletion of DNMT3A results in effects which can guide future studies

of molecular, cellular, and organismal dysfunction caused by mutation of DNMT3A.

We employed the DNMT3A$^{KO/+}$ mouse experimental system to assess how heterozygous

DNMT3A disruption impacts epigenetic regulation in the brain. Our analysis of DNA methylation

in tissues from DNMT3A$^{KO/+}$ mice detected very subtle changes in genome-wide mCG levels

across brain regions, with no global mCG effects in non-neural tissue (Figure 4A). Analysis of local

changes in mCG in the brain detected evidence of disrupted CG methylation at sites methylated

during postnatal development (i.e. adult hyper CG-DMRs). In addition, multiple hypo-CG-DMRs

can be detected at regulatory elements including enhancers. While limited, these effects have

the potential to alter gene expression and contribute to neurological alterations in these mice.

The limited nature of mCG effects is likely due to the redundant function of the other DNA

methyltransferases. The maintenance methyltransferase DNMT1 has the capacity to preserve

existing mCG patterns during cell divisions (Jeltsch et al., 2018). In addition, the *de novo*

methyltransferase DNMT3B is expressed with DNMT3A in many tissues during early development

and could provide critical redundancy for mCG patterning (Okano et al., 1999). Nonetheless, the

site-specific changes in mCG are also likely to occur in early development and in non-neural

tissues. For example, constitutive heterozygous deletion of DNMT3A has been shown to disrupt

mCG patterns in the blood and alter hematopoietic lineages (Cole et al., 2017). These changes in

mCG may contribute to changes in growth and other phenotypes observed in mice and humans.

In contrast to mCG, we detect a global reduction in mCA to approximately 30-50% of wild-

type levels in DNMT3A$^{KO/+}$ cortex, striatum, cerebellum, and hippocampus (Figure 4B). These

results generalize and extend findings in the hypothalamus (Sendžikaitė et al., 2019),

demonstrating the susceptibility of broad neuronal types and circuits to heterozygous loss of

DNMT3A. The susceptibility of mCA to heterozygous loss of DNMT3A is likely due to several

related factors. For example, DNMT3B is not expressed in postnatal neurons (Lister et al., 2013a),

and DNMT1 is not capable of depositing mCA (Jeltsch et al., 2018), making all mCA build-up in

neurons dependent on DNMT3A. In addition, the enzyme has slow kinetics for activity on CA sites
(Zhang et al., 2018) and deposition of mCA genome-wide by DNMT3A must take place in a
restricted time window (1-6 weeks) when the enzyme is highly expressed and active in neurons
(Clemens et al., 2019; Lister et al., 2013a; Stroud et al., 2017a). These constraints may make
enzyme levels limiting for mCA accumulation in neurons, providing an explanation for why global
mCA in the brain is sensitive to DNMT3A gene dosage. Notably, our findings suggest that
manipulations that activate the remaining DNMT3A, or prolong its high early postnatal
expression, might rescue deficits in mCA deposition. Conversely, duplication of the DNMT3A gene
could result in too much deposition of mCA and possibly cause significant neural dysfunction akin
to those effects seen in MeCP2 duplication disorder. Future studies can assess the feasibility of
rescue approaches and explore if DNMT3A duplication alters brain function.

Our analysis of chromatin changes downstream of altered DNA methylation has
uncovered a striking point of shared molecular disruption across models of DNMT3A disorders,
Rett syndrome, and MeCP2 duplication syndrome. While the clinical profile and pathophysiology
of DNMT3A disorders is clearly distinct from MeCP2 disorders, we have shown here that loss of
approximately a quarter of MeCP2 binding sites across the neuronal genome in the DNMT3A$^{KO/+}$
cortex results in subtle but wide-spread disruption of mCA-associated enhancer regulation that
partially phenocopies loss of MeCP2. This enhancer dysregulation can be linked to shared
alterations in gene expression across these models (Clemens et al., 2019) (Figure 6). Given the
critical roles of MeCP2-regulated genes for nervous system function (Gabel et al., 2015; Lagger
et al., 2017; Lyst and Bird, 2015), these epigenomic and transcriptomic effects likely contribute
to aspects of neurologic dysfunction observed in DNMT3A disorders. The persistence of many
mCA and mCG binding sites for MeCP2 in the DNMT3A$^{KO/+}$ may partially explain how DNMT3A
mutations manifest with less severe symptomology than in Rett Syndrome. In addition, absence
of DNMT3A early in prenatal development can contribute to overgrowth and other non-
overlapping aspects of DNMT3A and MeCP2 disorders. Together, our findings show that
disruption of mCA-MeCP2 mediated enhancer regulation likely contributes to three disorders
with distinct symptomology, defining a site of convergent molecular etiology underlying
heterogeneous clinical syndromes.

Our transcriptomic analysis of changes of ASD/NDD gene sets in DNMT3A mice has
further detected overlap with NDD beyond MeCP2 disorders, including both mouse models of
NDD/ASD (CHD8) and gene sets identified in human idiopathic ASD. As additional transcriptomic
studies of mouse models and human NDD brain emerge, systematic analyses of gene expression
effects can identify shared aspects of transcriptional pathology that can contribute to cognitive
and social deficits across diverse causes of NDD. Notably, the large number of chromatin
modifying enzymes mutated in these disorders raises the possibility that shared transcriptomic
effects emerge from common chromatin pathology. Our study has identified alterations in mCA
and enhancer regulation as a potential site of convergent dysfunction in MeCP2 and DNMT3A
disorders. Future studies may identify additional gene disruptions in which alterations in mCA
and enhancer dysregulation contribute to molecular pathology, expanding the role of
"methylopathies" in neurodevelopmental disease.
504

Acknowledgements
We thank the Division of Comparative Medicine at Washington University in Saint Louis for their
assistance with mouse husbandry and veterinary support. We are grateful for sequencing
support from the Genome Technology Access Center and the Center for Genome Sciences and
Systems Biology Spike in Cooperative at Washington University in Saint Louis. We thank members
of the Dougherty lab including S. Maloney, K. McCullough, and M. Rieger for assistance in USV
experiments. We thank J. Edwards, J. Goodman, and J. Yi for critical feedback on the experimental
design and manuscript. We thank A. Smith and T. Ley for helpful discussions. This work was
supported by NIH 5T32GM00815133 to D.L.C, by NIH 5T32GM007067 and F31NS108574 to
A.W.C., and by the Klingenstein-Simons Fellowship Fund, the G. Harold and Leila Y. Mathers
Foundation, the Brain and Behavior Research Foundation, the Simons Foundation for Autism
Research Initiative, and NIMH R01MH117405 to H.W.G.

Author Contributions
D.L.C and D.Y.W. are joint first authors, as each led critical components of the project and
analysis. D.L.C., J.R.Ma., and Y.R.L. generated and analyzed *in vitro* biochemical data. Y.R.L. and
S.A.N. generated primary neuronal culture samples. D.L.C. and J.R.Ma. generated skeletal
samples. N.M.K. and C.A.H. carried out craniofacial analysis and D.L.C. carried out long bone
analysis. D.L.C., J.R.Ma., D.F.W., and J.D.D. carried out behavioral tests and analysis. D.L.C.,
J.R.Ma., J.R.Mo., Y.R.L., and A.W.C. generated genomic data. D.Y.W. developed analysis
algorithms and pipelines. D.L.C., D.Y.W., J.R.Mo, and A.W.C. completing genomic analyses.
H.W.G. conceived the project and H.W.G., D.L.C., and D.Y.W. designed the experiments. H.W.G.,
D.L.C., and D.Y.W. wrote the manuscript and all authors contributed to manuscript editing and
revisions.

Declaration of interests

The authors declare no competing interests.

Figure Legends

Figure 1

Disease-associated DNMT3A mutations disrupt distinct aspects of protein function.

(**A**) Schematic of human DNMT3A protein showing canonical domains and disease-associated mutations identified in previous studies (Sanders et al., 2015; Tatton-Brown et al., 2018). (**B**) Example immunoblot of DNMT3A mutant protein expression. (**C**) Example images of DNMT3A protein immunocytochemistry from wild type and PWWP domain mutant. Scale bar = 20µm. (**D**) Quantification of DNMT3A mutant protein localization (#, $P<0.0001$; *, $P<0.05$; n=6-16 images; Generalized Linear Model test of percent nuclear expression per image for mutants compared to WT with Bonferroni correction). (**E**) Schematic of *in vitro* methylation assay for DNMT3A mutant proteins. (**F**) Activity of DNMT3A mutant proteins in the *in vitro* methylation assay. (***, $P<0.001$; **, $P<0.01$; *, $P<0.05$; n=4-19; one-sample Student's T-Test from normalized WT mean of 1 with Bonferroni correction). Bar graphs indicate mean with SEM error bars.

Figure 2

Disease-associated DNMT3A mutations prevent buildup of neuronal CA methylation.

(**A**) Schematic of DNMT3A functional analysis in primary culture neurons. Cortical neurons are collected from DNMT3A$^{flx/flx}$ mice at E14.5 and cultured. After 3 days *in vitro* (DIV), neurons are virally transduced with Cre recombinase and WT or mutant FLAG-tagged DNMT3A. On DIV 12.5, DNA and RNA are collected. Equal DNMT3A mRNA expression is verified by qRT-PCR (Figure S1D) and DNA is used for whole genome bisulfite sequencing analysis. (**B**) Relative mCA amount compared to Cre only and Cre+WT DNMT3A controls (#, $P<0.0001$; ***, $P<0.001$; **, $P<0.01$; *, $P<0.05$; n=4-11; one-sample Student's T-Test from normalized WT mean of 1 with Bonferroni correction). Bar graphs indicate mean with SEM error bars.

Figure 3

Heterozygous disruption of DNMT3A *in vivo* leads to growth and behavioral alterations.

(**A**) Body weight of DNMT3A$^{KO/+}$ and WT mice at three developmental timepoints (Male $P=0.038$ genotype by age interaction effect, $F_{(2,50)}=3.494$, n=6-18; Female $P=0.0032$ genotype by age interaction effect, $F_{(2,48)}=6.498$; Female $P=0.0016$ genotype effect, $F_{(1,48)}=11.18$, n=5-17; two-way ANOVA). (**B**) Lengths of femur and tibia bones measured by dual X-ray imaging shown as standard deviations from the WT mean for the DNMT3A$^{KO/+}$ mice (**, $P<0.01$; n=12; paired Student's T-Test). (**C**) Total ambulations of mice during 1-hour open-field testing, split into 10-minute bins ($P=0.0008$ effect by genotype, $F_{(1,46)}=13.02$, n=21,27; two-way repeated-measures ANOVA with Sidak's multiple comparison test; *, $P<0.05$; ****, $P<0.0001$). (**D**) Number of rearing events of mice during 1-hour open-field testing, split into 10-minute bins ($P=0.0103$ effect by genotype, $F_{(1,46)}=7.161$, n=21,27; two-way repeated-measures ANOVA with Sidak's multiple comparison test; *, $P<0.05$). (**E**) Schematic of open-field testing center and edge zones (left). Total time spent in the center zone of field during open-field testing (right) ($P=0.0075$, n=21,27; unpaired

Student's T-Test). (**F**) Schematic of the elevated plus maze indicating closed and open arms (left),
and percent of time mice spent in the open arms compared to all arms during first day of testing
(right) (*P*=0.0069; n=33,39; unpaired Student's T-Test). (**G**) Example images of marble burying
assay (left) and quantification of marbles buried during 30 minutes of testing split into 5-minute
bins (right)(*P*=0.0374 effect by genotype, $F_{(1,25)}$=4.834, n=14,13; two-way repeated-measures
ANOVA with Sidak's multiple comparison test; *, *P*<0.05). (**H**) Schematic of 3-chamber task in
which subject mouse can freely explore apparatus containing a novel mouse or an empty cup
(left). Quantification of time spent in zones closest to each cup (right) (Empty, *P*=0.0026; Novel,
P=0.0095; n=33,39; unpaired Student's T-Test). (**I**) Schematic of 3-chamber task in which subject
mouse can freely explore apparatus and interact with novel mouse or familiar mouse (left). Time
spent in zones closest to novel mouse and familiar mouse (right) (Familiar, *P*=0.29; Novel, *P*=0.24;
n=33,39; unpaired Student's T-Test). (**J**) Number of ultrasonic calls from pup isolated from the
nest for 3-minute testing over developmental time points (Analysis run on days 5-9 as these were
timepoints in which all animals tested had data; *P*=0.0378 effect by genotype, $F_{(1,285)}$=4.355, n=9-
46; two-way ANOVA with Sidak's multiple comparisons test; *, *P*<0.05). (**K-M**) Percent time spent
freezing in (**K**) Conditioned fear training (Baseline: *P*=0.0071 effect by genotype, $F_{(1,50)}$=7.897;
Cue: *P*=0.0013 effect by genotype, $F_{(1,50)}$=11.7; n=26; two-way repeated-measures ANOVA with
Sidak's multiple comparisons test; **, *P*<0.01; ***, *P*<0.001), (**L**) contextual fear trials (*P*=0.0215
effect by genotype, $F_{(1,50)}$=5.633, n=26; two-way repeated-measures ANOVA), and (**M**) cued fear
trials (Baseline: *P*=0.0606 effect by genotype, $F_{(1,50)}$=3.685; Cue: *P*<0.0001 effect by genotype,
$F_{(1,50)}$=17.03; n=26; two-way repeated-measures ANOVA with Sidak's multiple comparisons test;
*, *P*<0.05; ****, *P*<0.0001). Line graphs and bar graphs indicate mean with SEM error bars. Box
plots contain 10th-90th percentiles of data, with remaining data represented as individual points.
595
Figure 4
Global DNA methylation levels upon heterozygous loss of DNMT3A.
(**A**) Global mCG levels in DNA isolated from tissues of 8-week old mice (left) (*, *P*<0.05; unpaired
Student's T-Test with Bonferroni correction), and developmental time course of global mCG
(right), as measured by sparse whole genome bisulfite sequencing (WGBS). (**B**) Global mCA levels
in DNA isolated from tissues of 8-week old mice (left) (***, *P*<0.001; *, *P*<0.05; unpaired
Student's T-Test with Bonferroni correction), and developmental time course of global mCA
(right), as measured by sparse WGBS (*P*<0.0001 effect by genotype, $F_{(1,27)}$=1024; *P*<0.0001 effect
by age $F_{(5,27)}$=884.6; n=3-4; two-way ANOVA). Line graphs indicate mean with SEM error bars.
605
606
Figure 5
High-resolution analysis of DNA methylation changes in the DNMT3A$^{KO/+}$ cerebral cortex.
(**A**) Genome browser views of mCA and mCG in WT and DNMT3A$^{KO/+}$ cerebral cortex as measured
by high-depth WGBS. Broad view showing global reduction in mCA (left). Grey dashed line in mCA
plots at 0.03 to facilitate visual comparison of global mCA levels between genotypes. Zoomed-in
view of a DNMT3A$^{KO/+}$ CG-hypo-DMR that overlaps an enhancer (center) and a DNMT3A$^{KO/+}$ CG-
hypo-DMR at a CpG-island shore that overlaps with an adult-specific DMR (right). WT H3K27ac
ChIP-seq signal (Clemens et al., 2019), peaks of enhancer-associated H3K4me1
(Stamatoyannopoulos et al., 2012), peaks of promoter-associated H3K4me3

 (Stamatoyannopoulos et al., 2012), CpG islands, and gene annotations (Haeussler et al., 2019)
are shown below to illustrate overlap between DMRs and functional elements in the genome.
(**B,C**) Overlay of mCG signal for DMR regions shown in **A**. (**D**) Mean mCG/CG level per replicate
(top) and percent reduction, (DNMT3A$^{KO/+}$-WT)/WT, (bottom) in WT and DNMT3A$^{KO/+}$ cerebral
cortex across indicated classes of genomic regions. Adult-specific DMRs were identified in the
cortex (Lister et al., 2013a) (*, $P<0.05$; **, $P<0.01$; n=4 per genotype; paired Student's T-Test with
Bonferroni correction). (**E**) Heat map of CG DMRs called in the DNMT3A$^{KO/+}$ cortex. Biological
replicates (B1,B2) and technical replicates (R1,R2) are indicated. (**F**) Observed and expected (see
methods) overlap between DNMT3A$^{KO/+}$ cortex CG-DMRs and various genomic regions (****,
$P<0.0001$; ***, $P<0.001$; Fisher's Exact Test with Bonferroni correction). (**G**) Mean mCA/CA levels
per replicate (top) and percent reduction, (DNMT3A$^{KO/+}$-WT)/WT, (bottom) in WT and
DNMT3A$^{KO/+}$ cortex across indicated classes of genomic regions (*, $P<0.05$; **, $P<0.01$; ***,
$P<0.001$; n=4; paired Student's T-Test with Bonferroni correction). Box plots indicate median and
quartiles.
630
Figure 6
DNMT3A$^{KO/+}$ enhancer dysregulation and transcriptomic pathology overlaps with MeCP2
mutants.
(**A**) Genome browser view of DNA methylation and H3K27ac ChIP-seq data from WT and
DNMT3A$^{KO/+}$ cerebral cortex (top). Overlaid H3K27ac signal and mCA/CA levels at enhancer
regions highlighted in blue that were identified as dysregulated enhancers upon disruption of
mCA or MeCP2 (Clemens et al., 2019) (bottom). (**B**) Mean mCA sites/kb in WT and DNMT3A$^{KO/+}$
cortex (top) and number of mCA sites/kb lost in the DNMT3A$^{KO/+}$ cortex (bottom) for enhancers
significantly dysregulated in MeCP2 mutants (*, $P<0.05$; **, $P<0.01$; ***, $P<0.001$; ****,
$P<0.0001$; n=4; paired Student's T-Test with Bonferroni correction). (**C**) Boxplot of fold-change
in H3K27ac signal in DNMT3A cKO and the DNMT3A$^{KO/+}$ cortex for enhancers defined as
significantly dysregulated in MeCP2 mutants. (***, $P<10^{-8}$; **** $P<10^{-12}$; n=5 biological replicates
of DNMT3A$^{KO/+}$ and WT; Wilcoxon test) (**D**) Heatmap of changes in H3K27ac signal indicated
mutants across deciles of enhancers sorted by wild-type mCA or mCG sites ($P<2.2e-16$,
DNMT3A$^{KO/+}$ mCA/kb; $P<2.2e-16$, DNMT3A$^{KO/+}$ mCG/kb; Spearman Rho correlation). (**E**)
Observed versus expected overlap of significantly dysregulated genes (padj. < 0.1) in the
DNMT3A$^{KO/+}$ and genes dysregulated in DNMT3A cKO or MeCP2 mutant mice (***, $P<1e-5$; ****,
$P<1e-10$; hypergeometric test). (**F**) Significance of gene set expression changes in the indicated
direction in the DNMT3A$^{KO/+}$ cortex for GAGE analysis of gene sets identified as dysregulated in
DNMT3A cKO or MeCP2 mutant mice (Clemens et al., 2019). Note: legend is shared in **B** and **C**.
Box plots indicate median and quartiles.
652
Figure 7
Gene dysregulation in the DNMT3A$^{KO/+}$ overlaps with other ASD/NDD disorders.
(**A**) GAGE analysis of expression changes in DNMT3A$^{KO/+}$ for dysregulated gene sets identified in
studies of NDD mouse models (Gompers et al., 2017; Katayama et al., 2016; Sessa et al., 2019;
Tilot et al., 2016) (n=7 biological replicates of DNMT3A$^{KO/+}$ and WT). (**B**) GAGE analysis of
expression changes in DNMT3A$^{KO/+}$ for gene sets identified in studies of human ASD. Gene sets:
ASD module 12 (synaptic) and 16 (immune) identified in weighted-gene coexpression analysis of

human ASD brain (Voineagu et al., 2011), and ASD-dysregulated genes previously identified
(Abrahams et al., 2013; Gandal et al., 2018).

Figure S1 Related to Figures 1 and 2.
(**A**) Example images of DNMT3A mutant protein immunocytochemistry. Scale bar = 20μm. (**B**) Full
example immunoblot from Figure 1b with the truncated S312fs11x DNMT3A mutant protein
indicated with an asterisk. (**C**) Quantification of immunoblot signal of DNMT3A (α-FLAG) from
mutant proteins (***, $P<0.001$; n=6-30; unpaired Student's T-Test with Bonferroni correction).
(**D**) Genome wide mCA levels over time in neuronal cortical cultures, as measured by sparse
WGBS ($P=0.0035$ effect by time, $F_{(3,4)}=29.45$, n=2; one-way ANOVA). (**E**) Genome-wide mCA levels
in DNMT3A mutant add-back cortical cultures (****, $P<0.0001$; *, $P<0.05$; n=7-11; planned
unpaired Student's T-Tests with Bonferroni correction). (**F**) qRT-PCR of *Dnmt3a* normalized to
Actb for samples chosen for WGBS. (****, $P<0.0001$; n=4-11; one-sample Student's T-Test with
Bonferroni correction). Bar graphs indicate mean with SEM error bars.

Figure S2 Related to Figures 3,4,5,6 and 7.
(**A**) Normalized *Dnmt3a* mRNA and protein expression from 2-week cortices of DNMT3A$^{KO/+}$ and
wild-type littermates (****, $P<0.0001$; *, $P<0.05$; mRNA n=5, protein n=4, unpaired Student's T-
Test). (**B**) Protein expression of DNMT3A normalized to α-Tubulin measured by western blotting
for cerebral cortex of DNMT3A$^{KO/+}$ and wild-type littermates over developmental time ($P=0.0157$
effect by genotype, $F_{(1,16)}=7.303$, n=2-5; two-way ANOVA). (**C**) Protein expression of DNMT3A
normalized to α-Tubulin measured by western blotting for hippocampus, striatum, and
cerebellum ($P=0.0010$ effect by genotype, $F_{(1,18)}=15.48$, n=4; two-way ANOVA). Line plots indicate
mean with SEM error bars.

Figure S3 Related to Figure 3.
(**A,B**) Measurements of (**A**) femur and (**B**) tibia by dual x-ray imaging in WT and DNMT3A$^{KO/+}$ mice
at 8, 20, and 25 weeks of age. Lines indicate mean with SEM error bars. (**C**) Example dual x-ray
image of mouse body with femur and tibia indicated. (**D**) Example of reconstructed skull from
μCT imaging with landmarks used for craniofacial analysis shown. Red line indicates distance that
is significantly larger in the WT compared to the DNMT3A$^{KO/+}$, while blue line indicates distance
that is significantly smaller in the WT compared to the DNMT3A$^{KO/+}$ ($P<0.05$). (**E,F**) Principal
component analysis of (**E**) cranial and (**F**) mandibular shape shows no clear separation between
groups along PC1 or PC2.

Figure S4 Related to Figure 3.
(**A-J**) Comparison of DNMT3A$^{KO/+}$ and WT mice across a battery of sensorimotor assays.
DNMT3A$^{KO/+}$ mice show no significant difference in (**A**) walking initiation, or latency to fall off (**B**)
ledge or (**C**) platform. (**D-E**) DNMT3A$^{KO/+}$ mice show no difference compared to WT littermates in
motor coordination as evidenced by time on a continuous (**D**) and accelerating (**E**) rotarod. (**F**)
DNMT3A$^{KO/+}$ mice show no difference in grip strength compared to WT littermates evidenced by
no change in time on an inverted screen. (**G**) Mean % pre-pulse inhibition shows no significant
difference between genotypes. DNMT3A$^{KO/+}$ mice show a significant increase in time to (**H**) climb
down a pole ($P=0.016$, n=21,27; unpaired Student's T-Test), and to the top of a (**I**) 60° inclined

screen (P=0.039, n=21,27; unpaired Student's T-Test) or a (**J**) 90° inclined screen (P=0.045,
n=21,27; unpaired Student's T-Test). (**K**) DNMT3A$^{KO/+}$ mice show no deficit in elevated plus maze
exploration as measured by percent entries into open arms. (**L**) Percent preference for novel
conspecific in the 3-chambered social approach task for WT and DNMT3A$^{KO/+}$ mice using time
spent in zones as calculated: Mouse/(Mouse+Object)x100 or Novel/(Novel+Familiar)x100 within
each animal. (**M**) Total distance traveled during the 3-chambered social approach task for WT
and DNMT3A$^{KO/+}$ mice shows a broad reduction of distance traveled across all trials by
DNMT3A$^{KO/+}$ mice (P=0.018 effect by genotype, $F_{(1,70)}$=5.862, n=33,39; two-way ANOVA). (**N**)
Shock sensitivity during conditioned fear test as indicated by the minimum shock needed to
exhibit a behavioral response in mice shows no significant difference between genotypes. (**O-R**)
Path distance to escape platform and swim speeds in the Morris water maze task. DNMT3A$^{KO/+}$
mice show increased path distance to escape platform in both (**O**) cued trials (P=0.0012 effect by
genotype, $F_{(1,46)}$=11.93; P=0.0433 interaction effect of genotype and trial block, $F_{(3,138)}$=2.784;
n=21,27; two-way repeated-measures ANOVA) and (**P**) place trials (P=0.0408 interaction effect
of genotype and trial block, $F_{(4,184)}$=2.55 n=21,27; two-way repeated-measures ANOVA). No
significant difference is seen in swimming speed during (**Q**) cued trials (P=0.0634 effect by
genotype, $F_{(1,46)}$=3.619 n=21,27; two-way repeated-measures ANOVA) and (**R**) place trials
(P=0.098 effect by genotype, $F_{(1,46)}$=2.845, n=21,27; two-way repeated-measures ANOVA). (**S**)
DNMT3A$^{KO/+}$ mice show no significant difference in time spent in the target quadrant of a Morris
water maze compared to WT littermates. (**T**) DNMT3A$^{KO/+}$ mice show a trend towards a reduction
in platform crossings in the probe trial (P=0.0609; unpaired Student's T-Test). Bar graphs and line
plots indicate mean with SEM error bars. Box plots contain 10th-90th percentiles of data, with
remaining data represented as individual points.
727
Figure S5 Related to Figures 4,5, and 6.
(**A**) Boxplots of CG methylation over postnatal development at regions called as having higher
methylation in the frontal cortex of fetal versus adult tissue (Fetal-Specific DMR), or called as
having higher methylation in the frontal cortex of the adult versus fetal tissue (Adult-specific
DMR). DMRs from Lister et al. 2013, methylation data from Stroud et al. 2017. (#, P<2.2e-16;
Wilcoxon rank sum test with Bonferroni correction) (**B**) Boxplots of cortical methylation of WT
and DNMT3A cKO at 8 weeks postnatal within developmental DMRs shown in **A** (#, P<2.2e-16;
Wilcoxon rank sum test with Bonferroni correction). (**C**) Boxplots of cortical methylation of WT
and DNMT3A$^{KO/+}$ at 8 weeks postnatally within DMRs defined in the DNMT3A$^{KO/+}$ model. DMRs
are called on this data set, so no additional statistics were run on genotype differences. (**D**)
mCG/CG (left) and hmCG/CG (right) from WT and DNMT3A$^{KO/+}$ cortices in various genomic
contexts as measured by oxidative bisulfite sequencing. CpG islands were obtained from the
UCSC table browser (Haeussler et al., 2019), and CpG Shores were defined as the 8kb surrounding
them (*, P<0.05; paired Student's T-Test with Bonferroni correction). (**E**) Smooth scatter plots of
WT and DNMT3A$^{KO/+}$ mCA/CA for classes of genomic regions. Box plots indicate median and
quartiles.
744
Figure S6 Related to Figures 6 and 7.
(**A**) Volcano plot of DESeq log$_2$ fold changes of the DNMT3A$^{KO/+}$ versus WT. Genes reaching a
significance of $p_{adj.}$<0.1 are colored in red. (**B**) Top ten up- and down-regulated Gene Ontology

terms from Broad GSEA Molecular Signatures Database version 7.0 (Subramanian et al., 2005).
All terms are significant at an FDR<0.1. (**C**) GAGE analysis of developmental expression modules
(Parikshak et al., 2013). Significant modules (q-value<0.1) are colored in red (left). Expression
matched resampling of each gene set was performed 1,000 times and analyzed using GAGE for
enrichment in DNMT3A$^{KO/+}$ fold-change data (gray violin). This was compared with the true gene
set p-value (red point) to test for significance (right). Only the direction of dysregulation in which
the gene sets showed significance (i.e. DNMT3A$^{KO/+}$ greater or less) is shown. (**D**) Expression
matched resampling of GAGE analysis for gene sets displayed in Figures 6 and 7. Only the
direction of dysregulation in which the gene set showed significance (i.e. DNMT3A$^{KO/+}$ greater or
less) is shown.

758
759
760 **STAR Methods**

761 **Key Resources Table**

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Rabbit polyclonal anti-Histone H3 (acetyl K27)	Abcam	Cat# ab4729; RRID:AB_2118291
Rabbit Anti-DDDDK tag Polyclonal Antibody	Abcam	Cat#: ab1162; RRID:AB_298215
Mouse Anti-Dnmt3a Monoclonal Antibody, Clone 64B1446	Abcam	Cat#: ab13888 RRID: AB_300714
alpha Tubulin antibody [EP1332Y]	Abcam	Cat#: ab52866 RRID: AB_869989
Goat Anti-Rabbit IgG (H+L) Antibody, Alexa Fluor 568	ThermoFisher	Cat#: A-11011 RRID: AB_143157
IRDye 800CW Donkey anti-Rabbit IgG antibody	LI-COR Biosciences	Cat#: 926-32213 RRID: AB_621848
IRDye 800CW Goat anti-Rabbit IgG antibody	LI-COR Biosciences	Cat#: 926-32211 RRID: AB_621843
IRDye 800CW Goat anti-Mouse IgG antibody	LI-COR Biosciences	Cat#: 926-32210 RRID: AB_621842
Bacterial and Virus Strains		
Biological Samples		
Chemicals, Peptides, and Recombinant Proteins		
Critical Commercial Assays		
RNeasy Mini Kit	Qiagen	Cat#: 74104
DNeasy Kit	Qiagen	Cat#: 69504
Epitect Bisulfite Kit	Qiagen	Cat#: 59824
AllPrep DNA/RNA Kit	Qiagen	Cat#: 80204
NEBNext Ultra Directional RNA Library Prep Kit for Illumina	NEB	Cat#: E7420S
NEBNext Mulitplex Oligos for Illumina (Index Primers Set 1)	NEB	Cat#: E7335S
NEBNext rRNA Depletion Kit (Human/Mouse/Rat)	NEB	Cat#: E6310L
Ovation Ultralow Library System V2	NuGEN	Cat#: 0344-32
Ovation Ultralow Methyl-Seq Kit	NuGEN	Cat#: 0335-32

TrueMethyl oxBS plugin	NuGEN	Cat#: 0414-32
High-Capacity cDNA Reverse Transcription Kit	ThermoFisher	Cat#: 4368814
Power SYBR™ Green PCR Master Mix	ThermoFisher	Cat#: 4368577
Deposited Data		
RNA-sequencing data	This paper	GEO:
ChIP-sequencing data (H3K27ac)	This paper	GEO:
Bisulfite-sequencing data	This paper	GEO:
RNA, ChIP, Bisulfite-sequencing data	(Clemens et al., 2019)	GEO: GSE123373
DNMT3A ChIP-seq, Bisulfite-sequencing data	(Stroud et al., 2017b)	GEO: GSE104298
Dixon Hi-C contact matrices	(Dixon et al., 2012)	http://chromosome.sdsc.edu/mouse/hi-c/cortex.norm.tar.gz
Bisulfite-sequencing data	(Lister et al., 2013b)	GEO: GSE47966
Mus musculus mm9 genome assembly	UCSC	http://hgdownload.soe.ucsc.edu/goldenPath/mm9/
Ensembl gene models	UCSC	https://genome.ucsc.edu/cgi-bin/hgTables
Experimental Models: Cell Lines		
HEK293T	ATCC	Cat#: HEK293T
Neuro-2a	ATCC	Cat#: CCL-131
Experimental Models: Organisms/Strains		
Mouse: DNMT3A$^{KO/+}$	Derived from DNMT3A fl/fl strain (described in (Kaneda et al., 2004) provided by M. Goodell) and CMV-Cre (IMSR Cat# JAX:006054, RRID:IMSR_JAX:006054) mice obtained from the Jackson Laboratory. After DNMT3A heterozygous deletion, Cre negative mice were backcrossed to pure C57BL6 mice (IMSR Cat# JAX:000664, RRID:IMSR_JAX:000664) obtained from the Jackson Laboratory.	JAX: 006054; RRID:IMSR_JAX:006054 JAX: 000664; RRID:IMSR_JAX:000664
Mouse: *Dnmt3a*$^{fl/fl}$	(Kaneda et al., 2004)	Provided by M. Goodell
Oligonucleotides		
Actb Forward	IDT	AAGGCCAACCGTGAAAAGAT
Actb Reverse	IDT	GTGGTACGACCAGAGGCATAC
Dnmt3a Forward	IDT	GGCCTTCTCGACTCCAGATG
Dnmt3a Reverse	IDT	TTCCTCTTCTCAGCTGGCAC

Recombinant DNA		
Software and Algorithms		
DESeq2 (v1.14.1)	(Love et al., 2014)	http://www.bioconductor.org/packages/release/bioc/html/DESeq2.html
edgeR (v3.16.5)	(Robinson et al., 2010)	https://bioconductor.org/packages/release/bioc/html/edgeR.html
SAMtools (v1.3)	(Li and Durbin, 2009)	https://sourceforge.net/projects/samtools/files/
BEDtools2 (v2.25.0)	(Quinlan and Hall, 2010a)	https://github.com/arq5x/bedtools2
Bowtie2 (v2.2.5)	(Langmead and Salzberg, 2012)	http://bowtie-bio.sourceforge.net/bowtie2/index.shtml
STAR	(Dobin et al., 2013)	https://github.com/alexdobin/STAR
fastQC		https://www.bioinformatics.babraham.ac.uk/projects/fastqc/
MACS2 (v2.1.0)	(Zhang et al., 2008)	https://github.com/taoliu/MACS
Trim galore		https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/
Tadtree	(Weinreb and Raphael, 2016)	http://compbio.cs.brown.edu/projects/tadtree/
BS-seeker2	(Guo et al., 2013)	https://github.com/BSSeeker/BSseeker2
GREAT	(McLean et al., 2010)	http://great.stanford.edu
BSmooth	(McLean et al., 2010)	https://www.bioconductor.org/packages/release/bioc/html/bsseq.html
Gene Set Enrichment Analysis	Broad Institute	http://software.broadinstitute.org/gsea/downloads.jsp
Molecular Signatures Database	Broad Institute	http://software.broadinstitute.org/gsea/msigdb
ImageJ		https://imagej.nih.gov/ij/
Avizo		http://www.vsg3d.com/

Stratovan Checkpoint		https://www.stratovan.com/products/checkpoint
MorphoJ	(Dobin et al., 2013)	https://morphometrics.uk/MorphoJ_page.html
GraphPad Prism 7.03a		https://www.graphpad.com/

Contact for Reagent and Resource Sharing

Requests for reagents and resources should be directed towards the Lead Contact, Harrison Gabel (gabelh@wustl.edu). This study did not generate new unique reagents.

Experimental Model and Subject Details

Animal Husbandry
All animal protocols were approved by the Institutional Animal Care and Use Committee and the Animal Studies Committee of Washington University in St. Louis, and in accordance with guidelines from the National Institutes of Health (NIH). Mice were housed in a room on a 12:12 hour light/dark cycle, with controlled room temperature (20-22°C) and relative humidity (50%). Home cages measured 28.5 cm x 17.5 cm x 12 cm and were supplied with corncob bedding and standard laboratory chow and water. All mice were group-housed and adequate measures were taken to minimize animal pain or discomfort.

Transgenic animals
Male and female homozygous $Dnmt3a^{flx/flx}$ mice (Kaneda et al., 2004) were bred together for viral-mediated DNMT3A replacement assay culture experiments. To generate the DNMT3A heterozygous mouse model, $Dnmt3a^{flx/flx}$ mice were crossed to CMV:Cre (B6.C-Tg(CMV-cre)1Cgn/J) to generate $Dnmt3a^{KO/+}$Cre:CMV$^{+/-}$ offspring. $Dnmt3a^{KO/+}$Cre:CMV$^{+/-}$ progeny were bred to C57BL/6J to outcross the cre recombinase and generate experimental genotype (DNMT3A$^{KO/+}$). Subsequent experimental animals were generated from $Dnmt3a^{KO/+}$ males mated to C57BL/6J females to generate $Dnmt3a^{KO/+}$ and $Dnmt3a^{+/+}$ experimental and control animals for experiments. $Dnmt3a^{KO/+}$ females were not used for breeding to avoid social differences in mothering from mutant dams. Mice were genotyped with ear-DNA by PCR for $Dnmt3a$ and Cre, and recombination was tested. Mice were weighed at a variety of timepoints to assess growth.

Method Details
Immunocytochemistry
Staining. Neuro-2a cells (ATCC, CCL-131) were grown on coverslips and transfected with FLAG-tagged WT or mutant mouse DNMT3A plasmids and GFP plasmid. Coverslips were fixed with 4% paraformaldehyde in PBS for 20 minutes at room temperature, permeabilized with 0.1% Triton X-100 in PBS for 10 minutes at room temperature, and blocked with 1% BSA in PBS for 1 hour at room temperature. Coverslips were incubated overnight at 4°C in anti-DDDDK tag (FLAG-tag) primary antibody (Abcam, 1:5000, ab1162). Coverslips were then washed in PBS and incubated

for 1 hour at room temperature with fluorescent secondary antibody (ThermoFisher, 1:500, A-
11011) and counterstained with DAPI. *Imaging.* Images were captured using a Nikon A1Rsi
confocal microscope with a 20x air objective. Laser settings were kept constant for each image.
Analysis/Quantification. Cells were counted using an automatic threshold in FIJI and manually
classified as displaying nuclear or non-nuclear signal by a blinded observer. This was determined
by evaluating the overlap of FLAG signal (DNMT3A) with DAPI signal (nucleus). For mutants that
did not reach expression levels comparable to the WT or for images that had too few positive
cells, cell number was counted manually. 8 separate transfections were run, with each mutant
being counted over 3 or more independent experiments. Sample sizes are as follows: WT, 15
images, 880 cells; W297del, 8 images, 435 cells; I310N, 8 images, 492 cells; S312fs11x, 12 images,
321 cells; G532S, 9 images, 695 cells; M548K, 7 images, 333 cells; V665L, 6 images, 635 cells;
Y735C, 16 images, 613 cells; R749C, 8 images, 667 cells. P904L, 7 images, 692 cells. Percent
nuclear was assessed per image and a generalized linear model was run comparing each mutant
to WT. *P* values for each mutant were then Bonferroni corrected. We chose to use a generalized
linear model with Bonferroni correction to allow for us to compare ratios of percent nuclear
signal while taking into account experimental and biological replicates.
815

Modeling of DNMT3A disease mutations
HEK293T (ATCC, ACS-4500) or Neuro-2a cells (ATCC, CCL-131) were transfected with GFP and
FLAG-tagged WT or mutant mouse DNMT3A plasmids. Collected cell lysates were ruptured by 3
freeze/thaw cycles using liquid nitrogen, or sonication ~42 hours after transfection. Samples
were then either used for western blotting, the *in vitro* radioactive methyltransferase assay, or
RNA isolation for qRT-PCR.
822

qRT-PCR
RNA was reverse transcribed using the High-Capacity cDNA Reverse Transcription Kit
(Applied Biosystems). *Dnmt3a* and *Actb* were measured by qPCR using the Power SYBR™ Green
PCR Master Mix and primers *Actb* (F:AAGGCCAACCGTGAAAAGAT,
R:GTGGTACGACCAGAGGCATAC) and *Dnmt3a* (F:GGCCTTCTCGACTCCAGATG,
R:TTCCTCTTCTCAGCTGGCAC). Relative quantity of *Actb* and *Dnmt3a* cDNA was determined by
comparing the Ct of each primer set in each sample to a standard curve and then normalizing the
DNMT3A signal by the ACTB signal. We chose to compare experimental conditions to WT samples
using Student's T-Tests with Bonferroni correction, as these are two normally distributed groups
(visually checked) with similar variability.
833

In vitro **radioactive methyltransferase assay**
30 µl of cell lysate was used in the methyltransferase reaction previously described (Russler-
Germain et al., 2014). Lysates were incubated at 37˚C for 20 hours in 5 µl reaction buffer of 20
mM HEPES, 30 mM NaCl, 0.5 mM DTT, 1 mM EDTA, 0.2 mg/ml BSA, 5 mM 3 H-labeled SAM
(PerkinElmer, NET155050UC) and 500 ng/µl Poly(dI-dC) substrate (Sigma P4929). Substrate was
purified (Macherey-Nagel NucleoSpin Gel and PCR Clean-up) and radioactivity measured using a
scintillation counter. In instances where DNMT3A mutant showed altered protein expression, cell
lysate was re-balanced to match protein expression of WT DNMT3A. Only experimental replicates
where WT DNMT3A showed a 1.5-fold increase compared to GFP alone were used for

subsequent analysis. Outliers beyond 2 standard deviations above or below the mean were
removed. Number of independent replicates are as follows: W297del, 18; I310N, 19; S312fs11x,
4; G532S, 10; M548K, 15; V665L, 11; Y375C, 13; R749C, 7; P904L, 14. Significance was assessed
using a one-sample student's t-test, as we are comparing groups normalized to WT and GFP back
to the normalized value of 1.
848
Viral-mediated DNMT3A replacement assay
Functional activity of DNMT3A mutants in cortical neurons was determined by measuring
methylation build-up *in vitro*. Timed mating of DNMT3A$^{flx/flx}$ females and DNMT3A$^{flx/flx}$ males was
performed to collect embryonic cortical DNMT3A$^{flx/flx}$ neurons at embryonic day 14.5. At E14.5,
DNMT3A$^{flx/flx}$ cortical neurons were isolated and plated (DIV 0). On DIV 3, neurons were either
not perturbed or virally transduced with one of three conditions: 1) Cre only, 2) Cre and WT
DNMT3A, or 3) Cre and mutant DNMT3A. DNA and RNA were isolated on DIV 12.5 using the
AllPrep DNA/RNA Kit (Qiagen, 80204). DNA was used for whole genome bisulfite sequencing, and
RNA was used for qRT-PCR for DNMT3A. Number of independent replicates are as follows:
W297del, 5; S312fs11x, 4; G532S, 7; M548K, 6; V665L, 10; Y375C, 11; R749C, 6; P904L, 8. Buildup
of methylation over development was done using timed mating of C57BL/6J mice and collected
and prepared as described above. Significance was assessed using a one-sample student's t-test,
as we are comparing groups normalized to WT and GFP back to the normalized value of 1.
862
Ultrasonic vocalization and analysis
A total of 76 DNMT3A$^{KO/+}$ (n=30, 16 male and 14 female) and litter-matched WT (n=46, 25 male
and 21 female) mice were used for ultrasonic pup vocalization (USV) recording and analysis as
previously described (Barnes et al., 2017; Dougherty et al., 2013; Holy and Guo, 2005). Dams
were removed from the nest for a 10-minute acclimation, and individual pups had their body
temperature measured using an infrared laser thermometer. Pups were then removed from their
nest and placed in a dark, enclosed chamber. Ultrasonic vocalizations were recorded for 3
minutes with an Avisoft UltraSoundGate CM16 microphone and 416H amplifier using Avisoft
Recorder software (gain = 6 dB, 16 bits, sampling rate = 250 kHz). Following this, pups were
weighed and returned to their nest and littermates. All mice were recorded at postnatal days 5,
7, and 9, and on either day 11 or 15. Because not all animals were recorded from on day 11 or
15, only days 5, 7, and 9 were used for repeated measures ANOVA analysis. Frequency sonograms
were prepared and analyzed in MATLAB (frequency range = 40 kHz to 120 kHz, FFT size = 256,
overlap = 50%) with individual syllables identified and counted according to previously published
methods (Dougherty et al., 2013; Holy and Guo, 2005). Significance was assessed using a within-
subjects repeated measures ANOVA over timepoints 5-9, as these were when there was data
from all experimental subjects, and these are optimal testing times where number of calls was
highest.
881
Marble burying
A total of 27 DNMT3A$^{KO/+}$ (n=13, 8 male and 5 female) and litter-matched WT (n=14, 7 male and
7 female) mice were used for marble burying. Marble burying is a natural murine behavior and
has been used to indicate repetitive digging as well as anxiety-related behaviors. Protocol was
adapted from previously published methods (Lazic, 2015; Maloney et al., 2019a). In brief, 8-week

old mice were placed in a transparent enclosure (28.5 cm x 17.5 cm x 12 cm) with clean aspen
bedding and 20 dark blue marbles evenly spaced in a 4 x 5 grid on top of the bedding. Animals
were allowed to explore freely for 30 minutes. The number of buried marbles were counted every
5 minutes by two independent blinded observers. Marbles were considered "buried" if they were
at least two-thirds covered by bedding. Enclosure and marbles were cleaned thoroughly
between animals. Significance was assessed using a within-subjects repeated measures ANOVA
to determine if rate of burying marbles is different between genotypes. These statistical methods
are more appropriate than a simple t-test at 30 minutes, as mice may have buried all marbles
before this timepoint, and significant changes in marble burying behavior may have occurred at
an earlier timepoint in the assay.
897
Adult behavioral battery
A total of 72 DNMT3A$^{KO/+}$ (n=39, 18 male and 21 female) and litter-matched WT (n=33, 15 male
and 18 female) mice were used for adult behavioral testing. Mice were housed in mixed genotype
home cages with 2-5 animals per cage, and all tests were performed during the light cycle. All
experimenters were blinded to genotype during testing. For increased experimental rigor and
reproducibility, we used three separate cohorts of mice to ensure quality and consistency in any
observed phenotypes.
905
Testing started when mice were 3-4 months of age. The sequence of behavioral testing was
designed to minimize carry-over effects across behavioral tests. Cohorts, ages, and testing order
are in Supplementary Table 4. The majority of the assays were performed on cohorts 1 and 2
with cohort 3 being performed to test for reproducibility in some assays (Supplementary Table
4). Because of differences in testing sequences and exposure of mice to prior tests between
cohorts, we examined separate cohorts individually and looked at combined cohorts
(Supplementary Table 4). Testing was performed by the Washington University in St. Louis Animal
Behavior Core.
914
915 **One-hour locomotor activity**
916 Locomotor activity was evaluated by computerized photobeam instrumentation in transparent
917 polystyrene enclosures (47.6 cm x 25.4 cm x 20.6 cm) as previously described (Wozniak et al.,
918 2004). Activity variables such as ambulations and vertical rearings were measured in addition to
919 time spent in a 33 cm x 11 cm central zone.
920
Sensorimotor battery
Mice were assayed in walking initiation, balance (ledge and platform tests), volitional movement
(pole and inclined screens), and strength (inverted screen) as previously described (Grady et al.,
2006; Wozniak et al., 2004). For the walking initiation test, mice were placed on the surface in
the center of a 21 cm x 21 cm square marked with tape and the time for the mouse to leave the
square was recorded. During the balance tests, the time the mouse remained on an elevated
plexiglass ledge (0.75 cm wide) or small circular wooden platform (3.0 cm in diameter) was
recorded. During the Pole test, mice were placed at the top of a vertical pole with nose pointing
upwards. The time for the mouse to turn and climb down the pole was recorded. For the inclined
screen tests, a mouse was placed (oriented head-down) in the middle of an elevated mesh grid

measuring 16 squares per 10 cm angled at 60° or 90°. Time for the mouse to turn 180° and climb
to the top was recorded. For the inverted screen test, a mouse was placed on a similar screen
and when the mouse appeared to have a secure grasp of the screen, the screen was inverted
180° and the latency for the mouse to fall was recorded. All tests had a duration of 60 seconds,
except for the pole test which was 120 seconds. Two separate trials were done on subsequent
days and averaged time of both trials was used for analysis. Data from the walking initiation,
ledge, and platform tests were not normally distributed and therefore analyzed using Mann-
Whitney U tests.
939
Continuous and accelerating rotarod
Motor coordination and balance were assessed using the rotarod test (Rotamex-5, Columbus
Instruments, Columbus, OH) with three conditions: a stationary rod (60-second maximum), a
rotating rod at constant 5 rpm (60-second maximum), and a rod with accelerating rotational
speed (5 – 20 rpm, 180-second maximum) as previously described (Grady et al., 2006). This
protocol is designed to minimize learning and instead measure motor coordination, so testing
sessions were separated by 4 days to allow for extinction. Testing included one trial on stationary
rod, and two trials on both the constant-speed rotarod and accelerating rotarod. Later timepoints
in the constant speed rotarod test failed tests of normality, as the majority of mice stayed on the
rotating rod for all 60 seconds. However, data were analyzed with two-way repeated-measures
ANOVA.
951
Morris water maze
Spatial learning was assessed as previously described (Wozniak et al., 2004). Cued trials (visible
platform, variable location) and place trials (submerged, hidden platform, consistent location)
were conducted in which escape path latency, length, and swimming speeds were recorded.
Animal tracking was done using a computerized system (ANY-maze, Stoelting). During cued trials,
animals underwent 4 trials per day over 2 consecutive days with the platform being moved to a
different location for each trial with few distal spatial cues available. Each trial lasted no longer
than 60 seconds, with a 30-minute interval between each trial. Performance was analyzed across
four blocks of trials (2 trials/block). After a three-day rest period, animals were tested on place
trials, in which mice were required to learn the single location of a submerged platform with
several salient distal spatial cues. Place trials occurred over 5 consecutive days of training, with 2
blocks of 2 consecutive trials (60-second trial maximum, 30-second inter-trial-interval after the
mouse has reached the platform) with each block separated by 2 hours. Mice were released into
different quadrants over different trials. Place trials were averaged over each of the five
consecutive days (4 trials/block). One hour after the final block, a probe trial occurred (60-second
trial maximum) in which the platform is removed, and the mouse is released from the quadrant
opposite where the platform had been located. The time spent in pool quadrants, and the
number of crossings over the exact platform location were recorded. DNMT3A$^{KO/+}$ mice showed
a small, but significant reduction in target zone time in cohort 2, though there was no difference
in cohort 1 (Supplementary Table 4). Additionally, female mice had significantly faster swimming
speeds than male mice across both genotypes (Supplementary Table 4). These findings and the
observation that the DNMT3A$^{KO/+}$ were slower moving on some of the sensorimotor tests, make

path length a more appropriate variable than escape latency for evaluating performance, as
escape latency can be confounded by the differences in swimming speeds.
976
3-Chamber social approach
Sociability was assayed as previously described (Moy et al., 2004; Silverman et al., 2011). Mice
were tested in a rectangular all-Plexiglas apparatus (each chamber measuring 19.5 cm x 39 cm x
22cm) divided into three chambers with walls containing rectangular openings (5 cm x 8 cm) and
sliding doors. The apparatus was in a room with indirect light and was cleaned between tests
with Nolvasan solution. Stimulus mice were contained within a small stainless-steel withholding
cage (10 cm height x 10 cm diameter; Galaxy Pencil/Utility Cup, Spectrum Diversified Designs),
allowing minimal contact between mice without allowing fighting. Between tests, withholding
cages were cleaned with 75% ethanol solution. A digital video camera recorded movement of the
mouse within the apparatus and allowed for tracking with ANY-maze (Stoelting). Distance and
time spent in each chamber and investigation zones surrounding the withholding cages were
recorded. Zones were defined as 12 cm in diameter from the center of withholding cages.
989
The test sequence consisted of 4 consecutive 10-minute trials in which the test mouse is placed
in the middle chamber and allowed to freely explore the environment. In the first trial, the mouse
is placed in the middle chamber with the doors to other chambers shut. In the second trial, the
mouse is placed in the middle chamber and can explore all three chambers of the task, allowing
it to acclimate to the environment. Neither genotype tested showed a preference towards a side
of the chamber during this habituation. For the third trial, a sex-matched novel conspecific was
placed within a withholding cage with the other cage remaining empty. For the fourth trial, the
same sex-matched conspecific was in one withholding cage, while a new unfamiliar sex-matched
stimulus mouse was placed in the other withholding cage. The locations of stimuli mice were
counterbalanced across groups for the third trial and randomized novel or familiar for the fourth
trial.
1001
Elevated plus maze
Anxiety-like behaviors were examined using the elevated plus maze as previously described
(Boyle, 2006). The apparatus contains a central platform (5.5 cm x 5.5 cm) with two opposing
open arms and two opposing closed arms (each 36 cm x 6.1 cm x 15 cm) constructed of black
Plexiglas. Mouse position is measured using beam-breaks from pairs of photocells configured in
a 16 x 16 matrix and outputs are recorded using an interface assembly (Kinder Scientific) and
analyzed using software (MotoMonitor, Kinder Scientific) to determine time spent, distance
traveled, and entries made into open arms, closed arms, and the center area. Test sessions were
conducted in a dimly lit room with each session lasting 5 minutes and each mouse tested over 3
consecutive days. Data shown are from day 1. All mice showed a decrease in time, distance, and
entries into open arms on days 2 and 3. There was no significant difference between genotypes
in percent entries into open arms (Figure S4K; P=0.137; unpaired Student's T-Test) or total entries
into arms (data not shown), indicating that both genotypes explored the maze. Percent distance
traveled in open arms showed similar effects to percent time in open arms (Percent distance
traveled: P=0.027; unpaired Student's T-Test). Analysis of these data in individual cohorts
detected DNMT3A$^{KO/+}$ significant effects for the percent of open arm time on the first day in

cohorts 1 and 3, with no evidence of an effect in cohort 2 (Supplementary Table 4). Individual cohorts also showed no significant difference between genotypes in percent open arm entries (Supplementary Table 4) suggesting that mice explored the elevated plus maze sufficiently to detect anxiety-like behaviors.

Acoustic startle/prepulse inhibition

Sensorimotor gating was evaluated as previously described (Dougherty et al., 2013; Gallitano-Mendel et al., 2008; Hartman et al., 2001). In short, mice were presented with an acoustic startle response (120 dB auditory stimulus pulse, 40 ms broadband burst) and a pre-pulse (response to pre-pulse plus startle pulse). Stimulus onset began at 65 seconds, and 1ms force readings were obtained and averaged to produce an animal's startle amplitude. 20 startle trials were presented in 20 minutes. The first 5 minutes were an acclimation period where no stimuli above the 65 dB background were presented. The session started and finished with 5 consecutive startle (120 dB pulse) trials. The middle 10 trials were interspersed with pre-pulse trials, consisting of an additional 30 presentations of 120 dB startle stimuli preceded by pre-pulse stimuli of 4, 12, or 20 dB above background (10 trials for each PPI trial type). To calculate percent pre-pulse inhibition, we used $\%PPI = 100 \times (ASR_{\text{startle pulse alone}} - ASR_{\text{prepulse + startle pulse}})/ASR_{\text{startle pulse alone}}$.

Conditioned fear

Fear conditioning was done as previously described (Maloney et al., 2019a, 2019b). Mice were habituated to an acrylic chamber (26 cm x 18 cm x 18 cm) containing a metal grid floor and an odorant and was illuminated by LED light which remained on for the duration of the trial. Day 1 testing lasted 5 minutes in which an 80 dB tone sounded for 20 seconds at trial timepoints 100, 160, and 220 seconds. A 1.0 mA shock (unconditioned stimulus) occurred within the last 2 seconds of the tone (conditioned stimulus). Baseline freezing behavior during the first 2 minutes and the freezing behavior during the last 3 minutes was quantified using image analysis (Actimetrics, Evanston, Illinois). On Day 2, testing lasted for 8 minutes in which the light was illuminated but no tones or shocks were presented. On Day 3, testing lasted for 10 minutes in which the mouse was placed in an opaque chamber with a different odorant than the original test chamber. The 80 dB tone began at 120 seconds and lasted for the remainder of the trial and freezing behavior to the conditioned auditory stimulus was quantified for the remaining 8 minutes. Small elevated freezing levels of the DNMT3A$^{KO/+}$ mice for the contextual fear and auditory cue data could be interpreted as evidence for an increased baseline propensity to freeze or stronger fear conditioning in the mutant mice. However, an alternative hypothesis is that the exaggerated freezing levels displayed by the DNMT3A$^{KO/+}$ mice may reflect an emotional hypersensitivity to the footshock as was documented by their freezing levels during tone-shock training. In support of the latter hypothesis, evaluation of baseline freezing levels in individual cohorts showed that they were only significantly different in one of the two cohorts tested.

Statistical analysis for behavioral tests

Behavioral data were analyzed with R v3.3.2 (including the ANOVA function from the Car package in R (Fox and Weisberg, 2011)) and plots were made using GraphPad Prism 7.03a. Normality was assessed using the Shapiro-Wilkes test and visually confirmed. Data not normally distributed were analyzed using non-parametric tests, with the exception of continuous rotarod data. Sexes

were considered separately with genotype and times/block as fixed factors with no consistent
sex effects observed, therefore data were collapsed across sex. Statistical testing was performed
using planned assay-specific methods, such as using Student's T-Tests for single parameter
comparisons between genotypes, and within-subjects two-way repeated-measures ANOVA for
comparisons across timepoints. Individual timepoints within repeated measures tests were
evaluated using Sidak's multiple comparisons test. Individual cohorts were analyzed separately
and in aggregate with similar trends seen across cohorts (Supplementary Table 4), therefore data
from all cohorts were included together.
1070
1071 **Tissue**
1072 Brain tissue was dissected from DNMT3A$^{KO/+}$ and WT littermate mice in ice-cold PBS, flash-frozen
1073 in liquid nitrogen, and stored at -80°C.
1074
Western blotting
Western blotting from cell culture. Neuro-2a or HEK293T cells were collected and combined with
2x laemmli buffer with 5% β-mercaptoethanol. Samples were passed through a Wizard Column
(Fisher, Wizard Minipreps Mini Columns, PRA7211), boiled for 5 minutes, and run on a BioRad 4-
12% acrylamide gel at 125 V for 60 minutes. Samples were then transferred to a nitrocellulose
membrane, which was bisected between 37kDa and 50kDa bands. Membranes were blocked
with 3% bovine serum albumin in TBS-T for 1 hour at room temperature and then the lower
membrane was immunostained with anti-GFP (ThermoFisher, 1:2000, A-11122) while the upper
membrane was immunostained with anti-DDDDK (Abcam, 1:1000, ab1162) for 12-16 hours at
4°C. All primary and secondary antibodies were diluted in 3% Bovine Serum Albumin in TBS-T.
Membranes were then washed with TBS-T then incubated for 1 hour at room temperature with
IR-dye secondary antibody (IRDye 800CW Donkey anti-Rabbit, LI-COR Biosciences, 1:15,000,
product number: 926-32213). Blots were then washed in PBS, and imaged using the LiCOR
Odyssey XCL system, and quantified using Image Studio Lite software (LI-COR Biosciences). FLAG
(DDDDK) and GFP levels were normalized to a standard curve, and protein levels are expressed
as normalized DDDDK values divided by normalized GFP values to enable comparison of FLAG
(DDDDK) levels between blots. Each blot included a standard curve and WT samples. Outliers
beyond 2 standard deviations above or below the mean were removed. Number of independent
replicates are as follows: WT, 29; W297del, 7; I310N, 7; S312fs11x, 12; G532S, 7; M548K, 9;
V665L, 7; Y375C, 8; R749C, 6; P904L, 7. Significance was assessed using a one sample T-Test, as
protein expression levels were normalized to GFP and WT, and mutant protein expression was
compared to the normalized WT value of 1.
1097
Western blotting from tissue. Brain tissue samples were homogenized with a dounce
homogenizer in buffer with protease inhibitors (10mM HEPES pH 7.9, 10mM KCl, 1.5mM MgCl$_2$,
1mM DTT, 10mM EDTA). A portion of the lysate was removed and 1% SDS was added. Samples
were boiled for 10 minutes, followed by a 10-minute spin at 15,000g. Supernatant was collected
and run through a Wizard Column (Fisher, Wizard Minipreps Mini Columns, PRA7211), then
diluted in LDS sample buffer with 5% β-mercaptoethanol. Samples were boiled for 5 minutes, run
on an 8% acrylamide gel for 60 minutes at 125 V, and transferred to a nitrocellulose membrane.
Membrane was bisected between 75kDa and 100kDa. Membranes were blocked with 3% bovine

serum albumin in TBS-T for 1 hour at room temperature, and the upper membrane was
immunostained with anti-DNMT3A (Abcam, 1:1000, ab13888) while the lower membrane was
immunostained with anti-α-Tubulin (Abcam, 1:1000, ab52866) for 12-16 hours at 4°C. All primary
and secondary antibodies were diluted in 3% Bovine Serum Albumin in TBS-T. Membranes were
then washed with TBS-T then incubated for 1 hour at room temperature with IR-dye secondary
antibody (IRDye 800CW Goat anti-Rabbit, or IRDye 800CW Goat anti-Mouse, LI-COR Biosciences,
1:15,000, product numbers: 926-32211 and 926-32210 respectively). Blots were then washed in
PBS, imaged using the LiCOR Odyssey XCL system, and quantified using Image Studio Lite
software (LI-COR Biosciences). DNMT3A and α-Tubulin levels were normalized to a standard
curve, and protein levels are expressed as normalized DNMT3A values divided by normalized α-
Tubulin values to enable comparison of DNMT3A levels between blots. For brain region analysis,
sample sizes of n=4 per genotype (2 male and 2 female pairs) were used. For time course analysis,
sample sizes of n=2 per genotype (1 male and 1 female pairs) were used for all time points except
the 2-week timepoint in which n=6 (3 male and 3 female pairs) was used. Significance was
assessed using a two way ANOVA considering genotype and time to determine if there was a
detectable difference in protein expression over time.
1122
Bisulfite sequencing
Whole genome bisulfite sequencing from cortical cultures. Samples were chosen for whole
genome bisulfite sequencing if mutant and WT samples expressed equal amounts of *Dnmt3a*
mRNA as measured by qRT-PCR (Figure S2D). DNA from cortical cultures was bisulfite converted
and prepared for sequencing using the Tecan Ovation Ultralow Methyl-Seq Kit (Tecan, 0335-32)
and the Epitect Bisulfite Kit (Qiagen, 59824) was used for bisulfite conversion. We used alternate
bisulfite conversion cycling conditions ([95°C, 5 min; 60°C, 20 min] x 4 cycles, 20°C hold) to ensure
lowest possible bisulfite non-conversion rate. Libraries were PCR-amplified for 10-11 cycles.
Libraries were then pooled and sequenced at a depth of 0.01-0.03x genomic coverage using an
Illumina MiSeq 2x150 through the Spike-In Cooperative at Washington University in St. Louis.
Significance was assessed using a one-sample student's t-test, as we are comparing groups
normalized to WT and GFP back to the normalized value of 1.
1135
Whole genome bisulfite sequencing from tissue. DNA was isolated from tissue using the DNEasy
Kit (Qiagen). 300 ng of DNA was prepared for sequencing using the Ovation Ultralow Methyl-Seq
Kit (Tecan, 0335-32) with and the Epitect Bisulfite Kit (Qiagen, 59824) was used for bisulfite
conversion. For these samples, 300 ng of DNA was fragmented for 45 seconds with the Covaris
E220 sonicator (10% Duty Factory, 175 Peak Incidence Power, 200 cycles per burst, milliTUBE
200µL AFA Fiber). DNA was then purified using 0.7 volumes of Agencourt Beads to select for long
DNA inserts for sequencing. We used alternate bisulfite conversion cycling conditions ([95°C, 5
min; 60°C, 20 min] x 4 cycles, 20°C hold) to ensure lowest possible bisulfite non-conversion rate.
Libraries were PCR-amplified for 12 cycles. Libraries were then pooled and sequenced using an
Illumina MiSeq 2x150 through the Spike-In Cooperative at Washington University in St. Louis.
Samples for shallow-depth sequencing (Figure 4A,B) were sequenced at 0.01-0.03x genomic
coverage. For brain region and liver methylation, n=2 per genotype per region (one male pair,
one female pair). For developmental time course methylation, n=3-4 per genotype per timepoint,
with at least one male and one female pair. 8-week cortex samples for deep sequencing (Figure

5, Figure 6A,B,D, Figure S5C-E) were sequenced at 6.4-7.8x coverage per biological rep (two
technical reps per biological rep, two biological reps per genotype) using only male genotype
pairs. For shallow sequencing experiments, significance was assessed using either a two-sample
Student's T-Test to compare global methylation values of the cortex between two genotypes, or
using a two way ANOVA to compare broad methylation changes across a variety of brain regions
between genotypes. Genomic element comparisons were done using two-sample Student's T-
Tests with Bonferroni correction.

1157

Oxidative bisulfite sequencing from tissue. DNA was isolated from tissue using the DNEasy Kit
(Qiagen, 69504). 450 ng of DNA was prepared for sequencing using the Ovation Ultralow Methyl-
Seq Kit (Tecan, 0335-32) with TrueMethyl oxBS plugin (Tecan, 0414-32). For these samples, 450
ng of DNA was fragmented for 45 seconds with the Covaris E220 sonicator (10% Duty Factory,
175 Peak Incidence Power, 200 cycles per burst, milliTUBE 200µL AFA Fiber). DNA was then
purified using 0.7 volumes of Agencourt Beads to select for long DNA inserts for sequencing. 300
ng of DNA was used for OxBS libraries, whereas the remaining 150 ng of DNA was used for
bisulfite libraries. We used alternate bisulfite conversion cycling conditions ([95°C, 5 min; 60°C,
20 min] x 2 cycles; 95°C, 5 min; 60°C, 40 min; 95°C, 5 min; 60°C, 45 min; 20°C hold) to ensure
lowest possible bisulfite non-conversion rate. Bisulfite and oxidative bisulfite libraries were PCR-
amplified for 11 and 13 cycles respectively. Libraries were then pooled and sequenced using an
Illumina MiSeq 2x150 through the Spike-In Cooperative at Washington University in St. Louis.
Samples were sequenced at 0.8-2.2x genomic coverage per replicate (two replicates per
genotype). Genomic element comparisons were done using two-sample Student's T-Tests with
Bonferroni correction.

1173

Whole-genome bisulfite analysis
Bisulfite sequencing analysis was performed as previously described (Clemens et al., 2019).
Briefly, data were adapter-trimmed, mapped to mm9, then deduplicated and called for
methylation using BS-seeker2. Methylation levels across regions were assessed using bedtools
map -o sum, summing the number of reads mapping to Cs (supporting mC) and the amount of
coverage in the region, then dividing those two numbers(Quinlan and Hall, 2010b).
Hydroxymethylation was calculated as the percent methylation found in the BS-seq minus the
percent methylation found in the matching oxBS-seq. Due to count noise, this occasionally
resulted in apparent negative hydroxymethylation. During bisulfite sequencing not all DNA can
be efficiently bisulfite converted. Though our methods should maximize the amount of converted
unmethylated C, there is still a small percentage of unmethylated cytosines that are called as
methylated due to non-conversion (0.2-0.3%). Due to this non-conversion, very lowly methylated
regions (e.g. mCA at CpG islands) may not show the same percent reduction in mCA as highly
methylated regions. Data were visualized using the UCSC genome browser
(http://genome.ucsc.edu) (Kent et al., 2002). Average methylation per-sample is normally
distributed in all regions examined, and variance between genotypes is similar, fitting the
assumptions of a 2-sample t-test. Methylation levels for individual elements are not necessarily
normally distributed, so non-parametric tests were used instead.

1192

1193

Differentially methylated region detection

BSmooth (Hansen et al., 2012) was used to call differentially CpG methylated regions between DNMT3A$^{KO/+}$ and WT mice, using two technical replicates each of two biological replicates. CG sites were filtered for requiring at least 2x coverage in all replicates and differentially methylated regions were called with a statistical threshold of t-stat >2.0. These regions were further filtered for a length >100 bp and a requirement that the smoothed per-rep methylation values were consistent. For hypomethylated regions, all WT mCG/CG values needed to be greater than any KO mCG/CG value and all KO methylation values needed to be higher than all WT methylation values for hypermethylated regions. Data fit the assumptions and requirements of BSmooth. Data were distributed evenly between chromosomes, and the overlap between DMRs and regions of interest fit a hypergeometric distribution, making a fisher's exact test appropriate.

RNA sequencing

Total RNA isolation was carried out as previously described (Clemens et al., 2019). In brief, cerebral cortex was dissected in ice-cold PBS from DNMT3A$^{KO/+}$ and WT littermates at 8 weeks of age (n=7 pairs, 3 male, 4 female). Cortex was lysed in RLT buffer following the RNeasy Mini Kit (Qiagen, 74104). RNA libraries were generated from 250 ng of RNA with NEBNext Ultra Directional RNA Library Prep Kit for Illumina (NEB) using a modified amplification protocol (37°C, 15 minutes; 98°C, 30 seconds; [98°C, 10 seconds; 65°C, 30 seconds; 72°C, 30 seconds]x13; 72°C, 5 minutes; 4°C hold). RNA libraries were pooled at a final concentration of 10nM and sequenced using Illumina HiSeq3000 1x50bp with the Genome Technology Access Center at Washington University in St. Louis, typically yielding 15-30 million single-end reads per sample.

RNA sequencing analysis

RNA sequencing analysis was performed as previously described (Clemens et al., 2019). Briefly, raw FASTQ files were trimmed with Trim Galore and rRNA sequences were filtered out with Bowtie. Remaining reads were aligned to mm9 using STAR (Dobin et al., 2013) with the default parameters. Reads mapping to multiple regions in the genome were then filtered out, and uniquely mapping reads were converted to BED files and separated into intronic and exonic reads. Finally, reads were assigned to genes using bedtools coverage -counts (Quinlan and Hall, 2010b).

For gene annotation we defined a "flattened" list of longest transcript forms for each gene, generated on Ensgene annotations and obtained from the UCSC table browser. For each gene, Ensembl IDs were matched up to MGI gene names. Then, for each unique MGI gene name, the most upstream Ensgene TSS and the most downstream TES were taken as that gene's start and stop. Based on these Ensembl gene models, we defined TSS regions and gene bodies. Differentially expressed genes were identified using a Wald test through DESeq2, running using default parameters on exonic reads from the DNMT3A$^{KO/+}$ and WT.

Chromatin immunoprecipitation protocol

Chromatin immunoprecipitation was performed as previously described (Clemens et al., 2019; Cohen et al., 2011). Cerebral cortex was dissected on ice in PBS from DNMT3A$^{KO/+}$ and WT

littermates at 8-weeks old (n=5 pairs, 3 male, 2 female). The tissue was flash-frozen in liquid
nitrogen and stored at -80°C. Chromatin were fragmented with the Covaris E220 sonicator (5%
Duty Factory, 140 Peak Incidence Power, 200 cycles per burst, milliTUBE 1mL AFA Fiber). ChIP
was performed with H3K27ac antibody (0.025-0.1µg; Abcam, ab4729) and libraries were
generated using Ovation Ultralow Library System V2 (Tecan, 0344NB-32). Libraries were pooled
to a final concentration of 8-10nM and sequenced using Illumina HiSeq 3000 with the Genome
Technology Access Center at Washington University in St. Louis, typically yielding 15-40 million
single-end reads per sample.
1246
Chromatin immunoprecipitation analysis
ChIP sequencing analysis was performed as previously described (Clemens et al., 2019). Briefly,
reads were mapped to mm9 using bowtie2 and reads were extended based on library sizes and
deduplicated. Bedtools coverage –counts was used to quantify ChIP signal at the transcriptional
start site (TSS), gene body (GB), and transcriptional end site (TES) (Quinlan and Hall, 2010b).
edgeR was then used to determine differential ChIP-signal across genotypes. Data were
visualized using the UCSC genome browser (http://genome.ucsc.edu) (Kent et al., 2002).
1254
1255 **Controlled resampling**
1256 A similar resampling approach was used as previously described (Clemens et al., 2019). Briefly,
1257 for every entry in a sample set (e.g. DNMT3A-dysregulated genes), an entry in the control set
1258 (e.g. all other genes) with a similar desired characteristic (e.g. expression) was selected,
1259 generating a control set of the same size and variable distribution as the sample set.
1260
Identification of dysregulated enhancers
Enhancer regions from Clemens et al. 2019 were used, and enhancers dysregulated in the
DNMT3A$^{KO/+}$ were called using the same method. Briefly, H3K27ac ChIP-seq reads were
quantified in all acetyl peak regions, and edgeR was used to identify peaks with significantly
different amounts of H3K27ac signal. Peak regions were then divided into promoters, enhancers,
and non-identified peaks. Data fits the assumptions of BSmooth. Overlap between misregulated
enhancers in different genotypes fit a hypergeometric distribution.
1268
1269
GAGE
Gene set enrichment analysis for the gene sets described was performed using the Generally
Applicable Gene-set Enrichment (GAGE) program (Luo et al., 2009). Analysis was performed
directionally on input of shrunken, log-normalized exonic fold changes output from DESeq2
analysis of DNMT3A$^{KO/+}$ versus WT RNA-seq data. Gene sets with an FDR q-value below 0.1 and
an adjusted p-value below 0.5 following expression matched resampling repeated 1,000 times
were considered statistically significant. Gene sets were selected for analysis from both human
and mouse studies of autism associated genes. SFARI genes (Abrahams et al., 2013) with scores
of equal to or less than 3 were considered. Date accessed: 6/20/2019.
1279
GSEA

Gene Set Enrichment Analysis (GSEA) (version 7.0, the Broad Institute of MIT and
Harvard, http://software.broadinstitute.org/gsea/downloads.jsp) was performed on shrunken,
log-normalized exonic fold changes from DESeq2 between DNMT3A[KO+] and WT RNA-seq data.
GSEA calculated a gene set Enrichment Score (ES) that analyzed genes were enriched in the
biological signal conduction on the MsigDB (Molecular Signatures
Database, http://software.broadinstitute.org/gsea/msigdb). Background was set to all expressed
genes in this study and 1,000 permutations were set to generate a null distribution for
enrichment score in the hallmark gene sets and functional annotation gene sets. The gene sets
database used for enrichment analysis were 'c5.all.v7.0.symbols.gmt', 'c5.bp.v7.0.symbols.gmt',
'c5.cc.v7.0.symbols.gmt'and 'c5.mf.v7.0.symbols.gmt' and FDR <0.1 was defined as the cut-off
criteria for significance.
1292
Craniofacial morphological analyses
A total of 24 sex-matched littermate paired mice (DNMT3A[KO/+] n=12, 7 male, 5 female; WT n=12,
7 male, 5 female) across 3 time-points (8 weeks DNMT3A[KO/+] n=4, WT n=4; 20 weeks DNMT3A[KO/+]
n=4, WT n=4; 25 weeks DNMT3A[KO/+] n=4, WT n=4) were fixed in 4% paraformaldehyde through
intracardiac perfusions. Whole mouse heads were scanned at the Musculoskeletal Research
Center at Washington University in St. Louis using a Scanco μCT40 machine. CT images had voxel
dimensions of 0.018 millimeters and were reconstructed on a 2048x2048 pixel grid. The CT
images were converted to 8bit images using ImageJ (https://imagej.nih.gov/ij/) and surface
reconstructions were acquired in Avizo (http://www.vsg3d.com/). Thirty-five three-dimensional
landmarks were collected from surface reconstructions of the cranium and mandible using
Stratovan Checkpoint (https://www.stratovan.com/products/checkpoint).
1304
Generalized Procrustes Analysis in MorphoJ software was used to explore the differences and
similarities of shape between the DNMT3A[KO/+] mice and their WT littermates as previously
described (Hill et al., 2013). To control for possible differences in size, the landmark coordinate
data were natural log-transformed and analyzed with a linear regression model. Additionally, to
localize differences in form to specific linear distances, landmark data were analyzed using
Euclidean Distance Matrix Analysis (EDMA).
1311
Bone length measurements
A total of 24 sex-matched littermate paired mice (DNMT3A[KO/+] n=12, 7 male, 5 female; WT n=12,
7 male, 5 female) across 3 time-points (8 weeks DNMT3A[KO/+] n=4, WT n=4; 20 weeks DNMT3A[KO/+]
n=4, WT n=4; 25 weeks DNMT3A[KO/+] n=4, WT n=4) were fixed in 4% paraformaldehyde through
intracardiac perfusions. Decapitated mouse bodies were scanned at the Musculoskeletal
Research Center at Washington University in St. Louis using a Faxitron Model UltraFocus100 Dual
X-Ray machine. Bone lengths were measured using ImageJ. Data were taken over three age time-
points: 8 weeks, 20 weeks, and 25 weeks of age for male and female mice. There was no
significant difference in bone lengths based upon sex, but there was a difference based by age.
To normalize for this age effect, data were expressed as DNMT3A[KO/+] bone lengths normalized to
the WT lengths within groups. Left and right bones were measured and the larger was used for
analysis.
1324

Experimental design

Authenticated cell lines from ATCC (HEK293T, NEURO2A) were used, and no mycoplasma contamination testing was needed. Sample sizes were chosen based upon previously published studies using similar techniques. Statistical tests and exclusion criteria (values beyond 2 standard deviations of the group mean) were similar to that of previously published studies and indicated in the appropriate methods. For all animal experiments, experimenters were blinded to genotype during data collection. No treatment conditions were used, so no samples or animals were allocated to experimental groups and no randomization was needed. Tests that assume equal variance were only run if group variances were similar, otherwise alternative tests were used.

Data availability statement

The data that support the findings of this study are available from the corresponding author upon request. DOIs for all published gene sets used in comparison and enrichment analysis:

Lister et al. 2013: https://doi.org/10.1126/science.1237905;

Clemens et al. 2019: https://doi.org/10.1016/j.molcel.2019.10.033;

Stroud et al. 2017: https://doi.org/10.1016/j.cell.2017.09.047;

Gompers et al. 2017: https://doi.org/10.1038/nn.4592;

Katayama et al. 2016: https://dx.doi.org/10.1038/nature19357;

Tilot et al. 2016: https://doi.org/10.1038/mp.2015.17;

Sessa et al. 2019: https://doi.org/10.1016/j.neuron.2019.07.013;

Gandal et al. 2018: https://doi.org/10.1126/science.aat8127;

Voineagu et al. 2011: https://doi.org/10.1038/nature10110;

Abrahams et al. 2013: https://doi.org/10.1186/2040-2392-4-36;

Parikshak et al. 2013: https://dx.doi.org/10.1038/nature20612;

Raw and aggregate bisulfite-seq, raw and gene-count data for RNA-seq, and raw and peak call data for ChIP seq will be available on GEO.

Supplemental Information

Table S1. DNMT3A Mutation Table, Related to Figure 1, Figure 2, Figure S1

Table S2. BSsmooth-defined Differentially Methylated Regions, Related to Figure 4, Figure 5, Figure S5

Table S3. Table of differentially expressed genes, Related to Figure 6, Figure 7, Figure S6

Table S4. Behavioral Test Table, Related to Figure 3, Figure S3, Figure S4

References

Abrahams, B.S., Arking, D.E., Campbell, D.B., Mefford, H.C., Morrow, E.M., Weiss, L.A., Menashe, I., Wadkins, T., Banerjee-Basu, S., and Packer, A. (2013). SFARI Gene 2.0: a community-driven knowledgebase for the autism spectrum disorders (ASDs). Mol. Autism *4*, 36.

Amir, R.E., Van den Veyver, I.B., Wan, M., Tran, C.Q., Francke, U., and Zoghbi, H.Y. (1999). Rett syndrome is caused by mutations in X-linked MECP2 , encoding methyl-CpG-binding protein 2. Nat. Genet. *23*, 185–188.

Banerjee-Basu, S., and Packer, A. (2010). SFARI Gene: an evolving database for the autism
research community. Dis. Model. Mech. *3*, 133–135.

Barnes, T.D., Rieger, M.A., Dougherty, J.D., and Holy, T.E. (2017). Group and Individual
Variability in Mouse Pup Isolation Calls Recorded on the Same Day Show Stability. Front. Behav.
Neurosci. *11*.

Boxer, L.D., Renthal, W., Greben, A.W., Whitwam, T., Silberfeld, A., Stroud, H., Li, E., Yang, M.G.,
Kinde, B., Griffith, E.C., et al. (2019). MeCP2 Represses the Rate of Transcriptional Initiation of
Highly Methylated Long Genes. Mol. Cell.

Boyle, M.P. (2006). Forebrain Glucocorticoid Receptors Modulate Anxiety-Associated
Locomotor Activation and Adrenal Responsiveness. J. Neurosci. *26*, 1971–1978.

Chang, Y., Cole, T.B., and Costa, L.G. (2017). Behavioral Phenotyping for Autism Spectrum
Disorders in Mice. Curr. Protoc. Toxicol. *72*.

Chen, L., Chen, K., Lavery, L.A., Baker, S.A., Shaw, C.A., Li, W., and Zoghbi, H.Y. (2015). MeCP2
binds to non-CG methylated DNA as neurons mature, influencing transcription and the timing
of onset for Rett syndrome. Proc. Natl. Acad. Sci. *112*, 5509–5514.

Cholewa-Waclaw, J., Bird, A., von Schimmelmann, M., Schaefer, A., Yu, H., Song, H.,
Madabhushi, R., and Tsai, L.-H. (2016). The Role of Epigenetic Mechanisms in the Regulation of
Gene Expression in the Nervous System. J. Neurosci. *36*, 11427–11434.

Clemens, A.W., Wu, D.Y., Moore, J.R., Christian, D.L., Zhao, G., and Gabel, H.W. (2019). MeCP2
Represses Enhancers through Chromosome Topology-Associated DNA Methylation. Mol. Cell.

Cohen, S., Gabel, H.W., Hemberg, M., Hutchinson, A.N., Sadacca, L.A., Ebert, D.H., Harmin, D.A.,
Greenberg, R.S., Verdine, V.K., Zhou, Z., et al. (2011). Genome-Wide Activity-Dependent MeCP2
Phosphorylation Regulates Nervous System Development and Function. Neuron *72*, 72–85.

Cole, C.B., Russler-Germain, D.A., Ketkar, S., Verdoni, A.M., Smith, A.M., Bangert, C.V., Helton,
N.M., Guo, M., Klco, J.M., O'Laughlin, S., et al. (2017). Haploinsufficiency for DNA
methyltransferase 3A predisposes hematopoietic cells to myeloid malignancies. J. Clin. Invest.
127, 3657–3674.

Dixon, J.R., Selvaraj, S., Yue, F., Kim, A., Li, Y., Shen, Y., Hu, M., Liu, J.S., and Ren, B. (2012).
Topological domains in mammalian genomes identified by analysis of chromatin interactions.
Nature *485*, 376–380.

Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M.,
and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner. Bioinformatics *29*, 15–21.

Dougherty, J.D., Maloney, S.E., Wozniak, D.F., Rieger, M.A., Sonnenblick, L., Coppola, G.,
Mahieu, N.G., Zhang, J., Cai, J., Patti, G.J., et al. (2013). The Disruption of *Celf6*, a Gene

Identified by Translational Profiling of Serotonergic Neurons, Results in Autism-Related
Behaviors. J. Neurosci. *33*, 2732 LP – 2753.

Fazel Darbandi, S., Robinson Schwartz, S.E., Qi, Q., Catta-Preta, R., Pai, E.L.-L., Mandell, J.D.,
Everitt, A., Rubin, A., Krasnoff, R.A., Katzman, S., et al. (2018). Neonatal Tbr1 Dosage Controls
Cortical Layer 6 Connectivity. Neuron *100*, 831-845.e7.

Feliciano, P., Zhou, X., Astrovskaya, I., Turner, T.N., Wang, T., Brueggeman, L., Barnard, R.,
Hsieh, A., Snyder, L.G., Muzny, D.M., et al. (2019). Exome sequencing of 457 autism families
recruited online provides evidence for autism risk genes. Npj Genomic Med. *4*, 1–14.

Fox, J., and Weisberg, S. (2011). An R Companion to Applied Regression (SAGE Publications).

Gabel, H.W., Kinde, B., Stroud, H., Gilbert, C.S., Harmin, D.A., Kastan, N.R., Hemberg, M., Ebert,
D.H., and Greenberg, M.E. (2015). Disruption of DNA-methylation-dependent long gene
repression in Rett syndrome. Nature *522*, 89–93.

Gallitano-Mendel, A., Wozniak, D.F., Pehek, E.A., and Milbrandt, J. (2008). Mice Lacking the
Immediate Early Gene Egr3 Respond to the Anti-Aggressive Effects of Clozapine Yet are
Relatively Resistant to its Sedating Effects. Neuropsychopharmacol. Off. Publ. Am. Coll.
Neuropsychopharmacol. *33*, 1266–1275.

Gandal, M.J., Zhang, P., Hadjimichael, E., Walker, R.L., Chen, C., Liu, S., Won, H., van Bakel, H.,
Varghese, M., Wang, Y., et al. (2018). Transcriptome-wide isoform-level dysregulation in ASD,
schizophrenia, and bipolar disorder. Science *362*, eaat8127.

Gompers, A.L., Su-Feher, L., Ellegood, J., Copping, N.A., Riyadh, M.A., Stradleigh, T.W., Pride,
M.C., Schaffler, M.D., Wade, A.A., Catta-Preta, R., et al. (2017). Germline Chd8
haploinsufficiency alters brain development in mouse. Nat. Neurosci. *20*, 1062–1073.

Gowher, H., and Jeltsch, A. (2018). Mammalian DNA methyltransferases: new discoveries and
open questions. Biochem. Soc. Trans. *46*, 1191–1202.

Grady, R.M., Wozniak, D.F., Ohlemiller, K.K., and Sanes, J.R. (2006). Cerebellar Synaptic Defects
and Abnormal Motor Behavior in Mice Lacking α- and β-Dystrobrevin. J. Neurosci. *26*, 2841–
2851.

Guo, J.U., Su, Y., Shin, J.H., Shin, J., Li, H., Xie, B., Zhong, C., Hu, S., Le, T., Fan, G., et al. (2014).
Distribution, recognition and regulation of non-CpG methylation in the adult mammalian brain.
Nat. Neurosci. *17*, 215–222.

Guo, W., Fiziev, P., Yan, W., Cokus, S., Sun, X., Zhang, M.Q., Chen, P.-Y., and Pellegrini, M.
(2013). BS-Seeker2: a versatile aligning pipeline for bisulfite sequencing data. BMC Genomics
14, 774.

Guo, X., Wang, L., Li, J., Ding, Z., Xiao, J., Yin, X., He, S., Shi, P., Dong, L., Li, G., et al. (2015).
Structural insight into autoinhibition and histone H3-induced activation of DNMT3A. Nature
517, 640–644.

Haeussler, M., Zweig, A.S., Tyner, C., Speir, M.L., Rosenbloom, K.R., Raney, B.J., Lee, C.M., Lee,
B.T., Hinrichs, A.S., Gonzalez, J.N., et al. (2019). The UCSC Genome Browser database: 2019
update. Nucleic Acids Res. *47*, D853–D858.

Hansen, K.D., Langmead, B., and Irizarry, R.A. (2012). BSmooth: from whole genome bisulfite
sequencing reads to differentially methylated regions. Genome Biol. *13*, R83.

Hartman, R.E., Wozniak, D.F., Nardi, A., Olney, J.W., Sartorius, L., and Holtzman, D.M. (2001).
Behavioral Phenotyping of GFAP-ApoE3 and -ApoE4 Transgenic Mice: ApoE4 Mice Show
Profound Working Memory Impairments in the Absence of Alzheimer's-like Neuropathology.
Exp. Neurol. *170*, 326–344.

Hill, C.A., Martínez-Abadías, N., Motch, S.M., Austin, J.R., Wang, Y., Jabs, E.W., Richtsmeier, J.T.,
and Aldridge, K. (2013). Postnatal brain and skull growth in an Apert syndrome mouse model.
Am. J. Med. Genet. A. *161*, 745–757.

Holy, T.E., and Guo, Z. (2005). Ultrasonic Songs of Male Mice. PLoS Biol. *3*, e386.

Jeltsch, A., Broche, J., and Bashtrykov, P. (2018). Molecular Processes Connecting DNA
Methylation Patterns with DNA Methyltransferases and Histone Modifications in Mammalian
Genomes. Genes *9*, 566.

Kaneda, M., Okano, M., Hata, K., Sado, T., Tsujimoto, N., Li, E., and Sasaki, H. (2004). Essential
role for de novo DNA methyltransferase Dnmt3a in paternal and maternal imprinting. Nature
429, 900–903.

Katayama, Y., Nishiyama, M., Shoji, H., Ohkawa, Y., Kawamura, A., Sato, T., Suyama, M., Takumi,
T., Miyakawa, T., and Nakayama, K.I. (2016). CHD8 haploinsufficiency results in autistic-like
phenotypes in mice. Nature *537*, 675–679.

Kent, W.J., Sugnet, C.W., Furey, T.S., Roskin, K.M., Pringle, T.H., Zahler, A.M., and Haussler, and
D. (2002). The Human Genome Browser at UCSC. Genome Res. *12*, 996–1006.

Kinde, B., Gabel, H.W., Gilbert, C.S., Griffith, E.C., and Greenberg, M.E. (2015). Reading the
unique DNA methylation landscape of the brain: Non-CpG methylation, hydroxymethylation,
and MeCP2. Proc. Natl. Acad. Sci. *112*, 6800–6806.

Kinde, B., Wu, D.Y., Greenberg, M.E., and Gabel, H.W. (2016). DNA methylation in the gene
body influences MeCP2-mediated gene repression. Proc. Natl. Acad. Sci. *113*, 15114–15119.

Lagger, S., Connelly, J.C., Schweikert, G., Webb, S., Selfridge, J., Ramsahoye, B.H., Yu, M., He, C.,
Sanguinetti, G., Sowers, L.C., et al. (2017). MeCP2 recognizes cytosine methylated tri-nucleotide

and di-nucleotide sequences to tune transcription in the mammalian brain. PLOS Genet. *13*,
e1006793.

Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2. Nat.
Methods *9*, 357–359.

Lazic, S.E. (2015). Analytical strategies for the marble burying test: avoiding impossible
predictions and invalid p-values. BMC Res. Notes *8*, 141.

Li, H., and Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler
transform. Bioinformatics *25*, 1754–1760.

Lister, R., Mukamel, E.A., Nery, J.R., Urich, M., Puddifoot, C.A., Johnson, N.D., Lucero, J., Huang,
Y., Dwork, A.J., Schultz, M.D., et al. (2013a). Global epigenomic reconfiguration during
mammalian brain development. Science *341*.

Lister, R., Mukamel, E.A., Nery, J.R., Urich, M., Puddifoot, C.A., Johnson, N.D., Lucero, J., Huang,
Y., Dwork, A.J., Schultz, M.D., et al. (2013b). Global Epigenomic Reconfiguration During
Mammalian Brain Development. Science *341*, 1237905.

Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol *15*, 550.

Luo, W., Friedman, M.S., Shedden, K., Hankenson, K.D., and Woolf, P.J. (2009). GAGE: generally
applicable gene set enrichment for pathway analysis. BMC Bioinformatics *10*, 161.

Lyst, M.J., and Bird, A. (2015). Rett syndrome: a complex disorder with simple roots. Nat. Rev.
Genet. *16*, 261–275.

Maloney, S.E., Rieger, M.A., Al-Hasani, R., Bruchas, M.R., Wozniak, D.F., and Dougherty, J.D.
(2019a). Loss of CELF6 RNA binding protein impairs cocaine conditioned place preference and
contextual fear conditioning. Genes Brain Behav. *18*, e12593.

Maloney, S.E., Yuede, C.M., Creeley, C.E., Williams, S.L., Huffman, J.N., Taylor, G.T., Noguchi,
K.N., and Wozniak, D.F. (2019b). Repeated neonatal isoflurane exposures in the mouse induce
apoptotic degenerative changes in the brain and relatively mild long-term behavioral deficits.
Sci. Rep. *9*, 1–14.

McLean, C.Y., Bristor, D., Hiller, M., Clarke, S.L., Schaar, B.T., Lowe, C.B., Wenger, A.M., and
Bejerano, G. (2010). GREAT improves functional interpretation of cis-regulatory regions. Nat.
Biotechnol. *28*, 495–501.

McRae, J.F., Clayton, S., Fitzgerald, T.W., Kaplanis, J., Prigmore, E., Rajan, D., Sifrim, A., Aitken,
S., Akawi, N., Alvi, M., et al. (2017). Prevalence and architecture of de novo mutations in
developmental disorders. Nature *542*, 433–438.

Moy, S.S., Nadler, J.J., Perez, A., Barbaro, R.P., Johns, J.M., Magnuson, T.R., Piven, J., and
Crawley, J.N. (2004). Sociability and preference for social novelty in five inbred strains: an
approach to assess autistic-like behavior in mice. Genes Brain Behav. *3*, 287–302.

Nord, A.S., and West, A.E. (2019). Neurobiological functions of transcriptional enhancers. Nat.
Neurosci.

Nygaard, K.R., Maloney, S.E., and Dougherty, J.D. (2019). Erroneous inference based on a lack of
preference within one group: Autism, mice, and the social approach task. Autism Res. *12*, 1171–
1183.

Okano, M., Bell, D.W., Haber, D.A., and Li, E. (1999). DNA methyltransferases Dnmt3a and
Dnmt3b are essential for de novo methylation and mammalian development. Cell *99*, 247–257.

Parikshak, N.N., Luo, R., Zhang, A., Won, H., Lowe, J.K., Chandran, V., Horvath, S., and
Geschwind, D.H. (2013). Integrative Functional Genomic Analyses Implicate Specific Molecular
Pathways and Circuits in Autism. Cell *155*, 1008–1021.

Quinlan, A.R., and Hall, I.M. (2010a). BEDTools: a flexible suite of utilities for comparing
genomic features. Bioinformatics *26*, 841–842.

Quinlan, A.R., and Hall, I.M. (2010b). BEDTools: a flexible suite of utilities for comparing
genomic features. Bioinforma. Oxf. Engl. *26*, 841–842.

Robinson, M.D., McCarthy, D.J., and Smyth, G.K. (2010). edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. Bioinformatics *26*, 139–140.

Russler-Germain, D.A., Spencer, D.H., Young, M.A., Lamprecht, T.L., Miller, C.A., Fulton, R.,
Meyer, M.R., Erdmann-Gilmore, P., Townsend, R.R., Wilson, R.K., et al. (2014). The R882H
DNMT3A Mutation Associated with AML Dominantly Inhibits Wild-Type DNMT3A by Blocking Its
Ability to Form Active Tetramers. Cancer Cell *25*, 442–454.

Sanders, S.J., He, X., Willsey, A.J., Ercan-Sencicek, A.G., Samocha, K.E., Cicek, A.E., Murtha, M.T.,
Bal, V.H., Bishop, S.L., Dong, S., et al. (2015). Insights into Autism Spectrum Disorder Genomic
Architecture and Biology from 71 Risk Loci. Neuron *87*, 1215–1233.

Satterstrom, F.K., Kosmicki, J.A., Wang, J., Breen, M.S., De Rubeis, S., An, J.-Y., Peng, M., Collins,
R., Grove, J., Klei, L., et al. (2019). Large-scale exome sequencing study implicates both
developmental and functional changes in the neurobiology of autism. BioRxiv 484113.

Sceniak, M.P., Lang, M., Enomoto, A.C., James Howell, C., Hermes, D.J., and Katz, D.M. (2016).
Mechanisms of Functional Hypoconnectivity in the Medial Prefrontal Cortex of Mecp2 Null
Mice. Cereb. Cortex N. Y. NY *26*, 1938–1956.

Sendžikaitė, G., Hanna, C.W., Stewart-Morgan, K.R., Ivanova, E., and Kelsey, G. (2019). A
DNMT3A PWWP mutation leads to methylation of bivalent chromatin and growth retardation
in mice. Nat. Commun. *10*, 1–16.

Sessa, A., Fagnocchi, L., Mastrototaro, G., Massimino, L., Zaghi, M., Indrigo, M., Cattaneo, S.,
Martini, D., Gabellini, C., Pucci, C., et al. (2019). SETD5 Regulates Chromatin Methylation State
and Preserves Global Transcriptional Fidelity during Brain Development and Neuronal Wiring.
Neuron *104*, 271-289.e13.

Silverman, J.L., Turner, S.M., Barkan, C.L., Tolu, S.S., Saxena, R., Hung, A.Y., Sheng, M., and
Crawley, J.N. (2011). Sociability and motor functions in Shank1 mutant mice. Brain Res. *1380*,
120–137.

Simola, N., and Granon, S. (2019). Ultrasonic vocalizations as a tool in studying emotional states
in rodent models of social behavior and brain disease. Neuropharmacology *159*, 107420.

Spencer, D.H., Russler-Germain, D.A., Ketkar, S., Helton, N.M., Lamprecht, T.L., Fulton, R.S.,
Fronick, C.C., O'Laughlin, M., Heath, S.E., Shinawi, M., et al. (2017). CpG Island
Hypermethylation Mediated by DNMT3A Is a Consequence of AML Progression. Cell *168*, 801-
816.e13.

Stamatoyannopoulos, J.A., Snyder, M., Hardison, R., Ren, B., Gingeras, T., Gilbert, D.M.,
Groudine, M., Bender, M., Kaul, R., Canfield, T., et al. (2012). An encyclopedia of mouse DNA
elements (Mouse ENCODE). Genome Biol. *13*, 418.

Stroud, H., Su, S.C., Hrvatin, S., Greben, A.W., Renthal, W., Boxer, L.D., Nagy, M.A., Hochbaum,
D.R., Kinde, B., Gabel, H.W., et al. (2017a). Early-Life Gene Expression in Neurons Modulates
Lasting Epigenetic States. Cell *171*, 1151-1164.e16.

Stroud, H., Su, S.C., Hrvatin, S., Greben, A.W., Renthal, W., Boxer, L.D., Nagy, M.A., Hochbaum,
D.R., Kinde, B., Gabel, H.W., et al. (2017b). Early-Life Gene Expression in Neurons Modulates
Lasting Epigenetic States. Cell *171*, 1151-1164.e16.

Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A., Paulovich,
A., Pomeroy, S.L., Golub, T.R., Lander, E.S., et al. (2005). Gene set enrichment analysis: A
knowledge-based approach for interpreting genome-wide expression profiles. Proc. Natl. Acad.
Sci. *102*, 15545–15550.

Takumi, T., Tamada, K., Hatanaka, F., Nakai, N., and Bolton, P.F. (2019). Behavioral neuroscience
of autism. Neurosci. Biobehav. Rev. S0149763418303725.

Tatton-Brown, K., Seal, S., Ruark, E., Harmer, J., Ramsay, E., Del Vecchio Duarte, S., Zachariou,
A., Hanks, S., O'Brien, E., Aksglaede, L., et al. (2014). Mutations in the DNA methyltransferase
gene DNMT3A cause an overgrowth syndrome with intellectual disability. Nat. Genet. *46*, 385–
388.

Tatton-Brown, K., Zachariou, A., Loveday, C., Renwick, A., Mahamdallie, S., Aksglaede, L.,
Baralle, D., Barge-Schaapveld, D., Blyth, M., Bouma, M., et al. (2018). The Tatton-Brown-
Rahman Syndrome: A clinical study of 55 individuals with de novo constitutive DNMT3A
variants. Wellcome Open Res. *3*, 1–16.

Tilot, A.K., Bebek, G., Niazi, F., Altemus, J.B., Romigh, T., Frazier, T.W., and Eng, C. (2016). Neural
transcriptome of constitutional Pten dysfunction in mice and its relevance to human idiopathic
autism spectrum disorder. Mol. Psychiatry *21*, 118–125.

de la Torre-Ubieta, L., Won, H., Stein, J.L., and Geschwind, D.H. (2016). Advancing the
understanding of autism disease mechanisms through genetics. Nat. Med. *22*, 345–361.

Van Esch, H., Bauters, M., Ignatius, J., Jansen, M., Raynaud, M., Hollanders, K., Lugtenberg, D.,
Bienvenu, T., Jensen, L.R., Gécz, J., et al. (2005). Duplication of the MECP2 Region Is a Frequent
Cause of Severe Mental Retardation and Progressive Neurological Symptoms in Males. Am. J.
Hum. Genet. *77*, 442–453.

Voineagu, I., Wang, X., Johnston, P., Lowe, J.K., Tian, Y., Horvath, S., Mill, J., Cantor, R.M.,
Blencowe, B.J., and Geschwind, D.H. (2011). Transcriptomic analysis of autistic brain reveals
convergent molecular pathology. Nature *474*, 380–384.

Weinreb, C., and Raphael, B.J. (2016). Identification of hierarchical chromatin domains.
Bioinformatics *32*, 1601–1609.

Wozniak, D.F., Hartman, R.E., Boyle, M.P., Vogt, S.K., Brooks, A.R., Tenkova, T., Young, C., Olney,
J.W., and Muglia, L.J. (2004). Apoptotic neurodegeneration induced by ethanol in neonatal mice
is associated with profound learning/memory deficits in juveniles followed by progressive
functional recovery in adults. Neurobiol. Dis. *17*, 403–414.

Xie, W., Barr, C.L., Kim, A., Yue, F., Lee, A.Y., Eubanks, J., Dempster, E.L., and Ren, B. (2012).
Base-Resolution Analyses of Sequence and Parent-of-Origin Dependent DNA Methylation in the
Mouse Genome. Cell *148*, 816–831.

Yang, M., Silverman, J.L., and Crawley, J.N. (2011). Automated Three-Chambered Social
Approach Task for Mice. Curr. Protoc. Neurosci. *56*, 8.26.1-8.26.16.

Zhang, Y., Liu, T., Meyer, C.A., Eeckhoute, J., Johnson, D.S., Bernstein, B.E., Nusbaum, C., Myers,
R.M., Brown, M., Li, W., et al. (2008). Model-based Analysis of ChIP-Seq (MACS). Genome Biol.
9, R137.

Zhang, Z.M., Lu, R., Wang, P., Yu, Y., Chen, D., Gao, L., Liu, S., Ji, D., Rothbart, S.B., Wang, Y., et
al. (2018). Structural basis for DNMT3A-mediated de novo DNA methylation. Nature *554*, 387–
391.

1601

Figure 1

Figure 2

A

B

Figure 3

Figure 4

A

B

Figure 5

Figure 6

Figure 7

Supplementary Figure 2

A

B

C

Supplementary Figure 3

A
WT — DNMT3A^{KO/+}
Femur length (mm)
18, 17, 16, 15, 14, 13, 12
8 weeks   20 weeks   25 weeks

B
WT — DNMT3A^{KO/+}
Tibia length (mm)
22, 20, 18, 16, 14
8 weeks   20 weeks   25 weeks

C
Femur
Tibia

D
Key: WT significantly larger than DNMT3A mice
WT significantly smaller than DNMT3A mice

E
Cranium
Principal component 2
0.02, 0.01, 0.00, -0.01, -0.02, -0.03
-0.02 -0.01 0.00 0.01 0.02 0.03
Principal component 1
WT
DNMT3A^{KO/+}

F
Mandible
Principal component 2
0.02, 0.01, 0.00, -0.01, -0.02, -0.03
-0.02 -0.01 0.00 0.01 0.02 0.03
Principal component 1
WT
DNMT3A^{KO/+}

Supplementary Figure 4

Supplementary Figure 5

Supplementary Figure 6

A

B

GO Biological Process

C

D