1   A formin-mediated cell wall- plasma membrane- cytoskeleton continuum is required for

2   symbiotic infections in *Medicago truncatula*

<u>*Short title:*</u> The CW-PM-Cytoskeleton continuum during symbiotic infections

Pengbo Liang [a], Clara Schmitz [a], Beatrice Lace [a], Franck Anicet Ditengou [a], Jean Keller [b], Cyril

Libourel [b], Pierre-Marc Delaux [b], Thomas Ott [a,c,*]

[a] University of Freiburg, Faculty of Biology, Cell Biology, Schänzlestr. 1, 79104 Freiburg,

Germany

[b] Laboratoire de Recherche en Sciences Végétales (LRSV), Université de Toulouse, CNRS,

UPS, Castanet Tolosan, France

[c] CIBSS – Centre of Integrative Biological Signalling Studies, University of Freiburg,

Germany

* Correspondence author: Thomas.Ott@biologie.uni-freiburg.de

<u>*Keywords*</u>: Nodulation; symbiosis; actin; rhizobial infection; legume

ABSTRACT

Plant cell infections are tightly orchestrated by cell wall (CW) alterations, plasma membrane

(PM) resident signalling processes and dynamic remodelling of the cytoskeleton. During root

nodule symbiosis these processes result in morpho-dynamic responses including root hair

swelling and curling, PM invagination and polar growth of a tubular infection structure, the

infection thread (IT). However, the molecular details driving and guiding these PM

remodelling events remain to be unravelled. Here, we studied a formin protein (SYFO1) in *M.*

truncatula that is specifically induced during rhizobial infection. Phenotypical analysis of *syfo1*

mutants clearly indicates that the encoded protein is required for efficient rhizobial colonization

of root hairs. SYFO1 itself creates a proteinaceous bridge between the CW and the polarized

cytoskeleton. It binds to CW components via a proline-rich N-terminal segment, which is

indispensable for its function. On the cytoplasmic side of the PM SYFO1 is associated with

actin accumulations supporting the hypothesis that it contributes to cell polarization *in vivo*.

This is further sustained by the fact that cell shape changes can be induced in a stimulus-

dependent manner in root protoplasts expressing SYFO1. Taken together we provide evidence

for the evolutionary re-wiring of a generic cytoskeleton modulator into a symbiosis-specific

response.

38
39

INTRODUCTION

Legumes have the unique ability to symbiotically associate with rhizobia to maintain a nitrogen-fixing mutualism. Intracellular colonization of *Medicago truncatula* roots by the compatible rhizobium *Sinorhizobium meliloti* initiates from young, growing root hairs. In the course of the interaction an organogenetic program is executed in the root cortex and the pericycle that results in the development of nodules, in which symbiotic nitrogen fixation takes place [1, 2]. The first morphological step to establish this symbiosis comprises a rhizobial trap, where a growing root hair engulfs the symbiont by physically curling around it [3, 4]. Phenomena such as root hair deformation and root hair branching, steps that precede bacterial trapping in legumes [5, 6], have been generally observed in plants in response to the microtubule-stabilizing agent taxol [7], in mutants like the kinesin *mrh2* [8] or upon over-expression of the formin protein AtFH8 [9] and the ROP GTPase RHO-OF PLANTS 2 [10]. In contrast, incomplete curls were only observed at low frequency in *Arabidopsis* mutants affected in the loci *CEN1*, *CEN2*, *CEN3* [11] and in the *scn1-1* mutant, where the corresponding locus encodes the RhoGTPase GDP dissociation inhibitor SUPERCENTIPEDE1 [12]. This indicates full root hair curling to represent a rather specific invention. The entrapment of the symbiont completes with its full enclosure between root hair cell walls in a structure called the 'infection chamber' (IC) [3-5, 13]. This is followed by exocytotic secretion of host-derived cell wall loosening enzymes such as NODULE PECTATE LYASE [14] which, most likely, subsequently allows the formation of a negatively curved plasma membrane (PM) structure that further elongates into a tube-like channel, the 'infection thread' (IT) [3, 4, 15]. Parallel to these morphological changes a set of transcription factors genetically re-programs host root cells to allow transcellular IT progression and nodule organogenesis [16-22].

Molecularly, initial root hair responses are triggered upon the recognition of bacterial signalling molecules, called Nod Factors, by host LysM-type receptor-like kinases [23-28].

These include root hair swelling, deformation and branching that precede root hair curling.

This is aided by a tip-localized cytosolic calcium gradient [29, 30], global actin re-

arrangements and dense subapical fine actin bundles that are required for the delivery of Golgi-

derived vesicles to the root hair tip [5, 31-33]. However, the molecular machinery steering actin

reorganisation and polarity during root hair curling remains elusive. Altered actin dynamics

during early responses to NFs or rhizobia have been shown in mutants such *Lotus japonicus*

pir1/nap1 and *scarn* that are affected in the actin-related SCAR/WAVE complex [34, 35]. In

addition, the presence of highly dynamic F-actin plus ends in swelling root hairs and during

root hair deformation implies that newly accumulated actin physically pushes towards the PM

to re-initiate the growth [36]. But, different to metazoan cells, where membrane protrusions

such as filopodia are driven, among others, by formin-mediated polar growth of actin filaments

[37], any membrane protrusions in plant cells ultimately require local cell wall modifications

and thus this membrane-cell wall continuum to be coordinatively regulated.

Here, we identified SYMBIOTIC FORMIN 1 (SYFO1) as an essential gene controlling

symbiotic responses in root hairs. SYFO1 transiently re-localizes to the root hair tip upon

inoculation of *M. truncatula* root hairs with *S. meliloti* and mediates the formation of polar

actin accumulations. Its function strictly depends on the protein extracellular domain that

mediates cell wall attachment. Taken together we demonstrate SYFO1 as the first player

regulating the continuum between the plasma membrane and the cell wall during the onset of

rhizobial infections.

RESULTS

Evolutionary and transcriptional patterns identify symbiotically regulated formins

Since formins are well known proteins conferring polar growth of actin filaments, we searched

for candidates within this family that could control symbiotically induced root hair responses.

Based on the presence of a conserved Formin Homology 2 (FH2) domain, we identified 19

candidates in the *Medicago truncatula* genome (Table S1). Using publicly available

transcriptome data, two of them (*Medtr5g036540.1* and *Medtr8g062830.1*) were found to be

transcriptionally up-regulated during early stages of symbiotic interactions [38]. We

independently verified these data using quantitative RT-PCR (qRT-PCR) with

Medtr5g036540.1 being induced by about 60-fold at one day post inoculation (dpi) of roots

with *S. meliloti* while only a weak induction could be confirmed for *Medtr8g062830.1* at 5 dpi

(Fig. S1A). Therefore, we named the genes *SYMBIOTIC FORMIN 1* (*SYFO1*

(*Medtr5g036540.1*, *MtrunA17_Chr5g0414941* in the v5r1.6 *M. truncatula* genome version)

and SYFO1-like (*SYFO1L, Medtr8g062830.1, MtrunA17_Chr8g0364331*). Both encoded

proteins contain a predicted signal peptide (SP) in the extracellular domain (ECD) followed by

a single-span transmembrane domain (TMD), and FH2 domain in the cytoplasmic side.

To spatially resolve the observed transcriptional patterns for *SYFO1*, we generated a

fluorescent reporter where a nuclear localized tandem GFP was driven by the endogenous

SYFO1 promoter (*ProSYFO1-NLS-2xGFP*). Consistent with the qRT-PCR results (Fig. S1A),

the *SYFO1* promoter was activated at 1 dpi in root hairs and cortical cells (Fig.S1 B-C) while

no activity was observed in the absence of the symbiont.

To supplement the identification strategy and to search for putative genetic redundancy, we

studied evolutionary patterns within the formin family. The *SYFO1/1L* clade contains genes

from all Eudicots species included in the analysis, and *SYFO1* and *SYFO1L* derived from the

Papilionoideae duplication. In *A. thaliana*, three co-orthologs of *SYFO1* and *SYFO1L* are found:

AtFH4, *AtFH7*, *AtFH8* likely deriving from the Brassicaceae triplication (Fig. S2). While

AtFH4 and AtFH8 resemble a similar protein domain structure compared to SYFO1 and

SYFO1L, AtFH7 has lost the signal peptide and displays an altered TMD domain. Mining the

gene expression atlas of other Papillionoideae, we found *SYFO1* (in common bean,

Phvul.002G077100.2) or *SYFO1L* (in *Lotus japonicus Lj0g3v0115049.1*) upregulated during

nodulation, pinpointing for a potential shared function in nodulation within this clade.

Based on the phylogeny, we tested for relaxed or intensified selective pressure acting on

different branches of interest in eudicot sequences (represented by circles) (Fig. S2;

Supplemental Table S3). We identified relaxed selective pressure indicated by a higher overall

ratio of non-synonymous (dN) vs. synonymous (dS) mutations (dN/dS) on the branch

supporting the clade that includes all species that develop the root nodule symbiosis (NFN;

K=0.54, LRT=85.3 and p-val < 1e-16). This relaxation was individually confirmed for Fabales

(which includes legumes; K=0.66, LRT=18.3 and p-val< 14.1e-09), Cucurbitales (K=0.78,

LRT=25.0 and p-val=5.6e-07) and Fagales (K=0.65, LRT=18.3 and p-val< 1.9e-05) but not

for Rosales (K=1.17, LRT=7.4 and p-val< 6.7e-03) for which a slight intensification of

selective pressure (lower overall dN/dS ratio) was detected. In contrast, Brassicaceae species

are under strong intensification of the selective pressure (K=4.6, LRT=84.5 and p-val < 1e-16)

that is not even relaxed by the triplication of the branch in this family. The relaxed selection

pressure detected at the base of the NFN clade and at the base of the Fabales may both reflect

neofunctionalization that would have occurred for the recruitment of SYFO1/1L in root nodule

symbiosis. Interestingly, the corresponding orthologs of *Parasponia andersonii*, a non-legume

tree that forms a Frankia-type symbiosis with rhizobia, are not induced during the symbiotic

interaction. This further supports a possible functional specialisation of these formins in

legumes /Papilionoidae [39].

SYFO1 controls rhizobial infection and root hair responses

In order to genetically assess the function of SYFO1 and SYFO1L, we identified two independent *Tnt1* transposon insertion lines for *SYFO1* [*syfo1-1* (NF9730) at 485bp and *syfo1-2* (NF9495) at 1834bp downstream of the start codon] and *SYFO1L* [*syfo1L-1* (NF20350) at 1279bp and *syfo1L-2* (NF15608) at 1370bp downstream of the start codon] in Medicago wild type R108 (Fig. 1A). Endogenous *SYFO1* and *SYFO1L* transcripts were significantly reduced in both lines, respectively (Fig. S3), while only the *syfo1-1* and *syfo1-2* alleles showed a significant nodulation phenotype developing fewer nodules per root at 3 weeks post inoculation (wpi) with about half of them being aborted and/or white, indicative of non-functional nodules (Fig. 1B-D). In contrast, bacterial infection patterns of Medicago wild-type and *syfo1L-1* and *syfo1L-2* mutant nodules were identical (Fig. 1C, D and Fig. S4A, B). The infection zones were found to be reduced in white but elongated (Fig. S4C) or spherical (Fig. S4D) *syfo1* mutant nodules. The latter failed to maintain a persistent meristem (Fig. S4D). Therefore, we selected SYFO1 as the prime candidate of interest.

Although cytoskeleton-related mutants, like Lotus *pir1, nap1* and *scarn [34, 35]*, exhibit an impaired root hair growth, this was not observed for *syfo1-1* and *syfo1-2* where root hair length was indistinguishable from wild-type (Fig. S5). In agreement with this, we did not observe any differences in actin arrangement in growing root hairs within the symbiotically susceptible infection zone [4] under non-inoculated conditions (Fig. S6). These data demonstrate that SYFO1 is not required for normal root hair development under non-symbiotic conditions. As *SYFO1* transcripts were upregulated at 1-3 dpi, a stage where we observe most root hairs responding to the presence of the symbiont by root hair deformation (1-2 dpi) (Fig. 2A) and curling (2-3 dpi) (Fig. 2B), we phenotypically assessed *syfo1* mutants at these stages. Both mutant alleles showed significantly fewer deformations (Fig. 2C) as well as infection chambers and infection threads (Fig. 2E) at the respective time points. Both phenotypes were fully

rescued when generating independent transgenic roots expressing a *ProSYFO1:SYFO1-GFP*

(e.g. *syfo1-1c*) construct in these mutant backgrounds (Fig. 2D, F), demonstrating that the

insertion in the *SYFO1* locus is causative for these phenotypes.

As it was previously demonstrated that the inoculation of Lotus root hairs with its symbiont *M.*

loti resulted in a strong polarization and bundling of actin filaments with a strong accumulation

of F-actin in the root hair tip [34, 35], we tested whether this pattern is affected in *syfo1* mutants.

In the absence of *S. meliloti*, longitudinal actin filaments were observed in young growing root

hairs within the infection zone of Medicago wild-type plants (Fig. S7A). In agreement with the

observations in Lotus, actin strongly bundled and polarized with an accumulation of F-actin at

the apex (Fig. S7B-C) or occasionally at the apical shank of responding root hairs (Fig. S7D)

in almost 90% of all root systems of wild-type plants (Fig. S7E). However, this pattern was

strongly reduced in both *syfo1* mutant alleles where only about 30% of all tested roots contained

root hairs responding with the above-mentioned pattern (Fig. S7E).

SYFO1 associates with polar actin assemblies under symbiotic conditions

Since our data indicated a symbiosis-specific role of SYFO1 in root hair polarization, we

investigated spatial and temporal dynamics of SYFO1 at subcellular resolution. In roots

expressing the *ProSYFO1:SYFO1-GFP* construct, we observed a weak homogenous signal of

the SYFO1 protein at the PM of root hairs in the absence of rhizobia (Fig. 3A). The underlying

low, basal expression was also detected by qRT-PCR (Fig. S3A) while it was most likely too

weak when using the nuclear-localized GFP reporter to test promoter activity (Fig. S1).

Interestingly, SYFO1 strongly accumulated in subapical and apical foci at root hair tips prior

to deformation at 2 dpi with *S. meliloti* (Fig. 3B, C), which strongly resembled actin patterns

observed upon Nod Factor application as reported earlier [34]. In root hairs that

morphologically responded by deformation (Fig. 3D) and curling (Fig. 3E), SYFO1 distributed

again along the PM with only mild accumulations at the apical region (Fig. 3D). SYFO1 also

resided along the infection thread membrane even though to a much weaker extent (Fig. 3F-

F'). When co-localizing actin and SYFO1 (here: *ProUbi-SYFO1-GFP*) we frequently observed

transient SYFO1 accumulations in close proximity to enlarged nuclei, a hallmark for

symbiotically activated root hairs [40], with actin bundles orienting towards a nucleation centre

at the apical shank of the root hair (Fig. 3G-G').

Ligand-dependent morphological change mediated by SYFO1

To test whether SYFO1 acts in membrane-associated actin nucleation and polar filament

elongation, we made use of the fact that extension of actin filaments can drive membrane

protrusions as shown in cells lacking a rigid cell wall such as *Drosophila* Schneider 2 and

mouse P19 cells [41, 42]. For this, we generated a transgenic Medicago root organ culture

(ROC) constitutively expressing SYFO1-GFP for 8 weeks. When isolating protoplasts from

this culture, SYFO1 resided in the PM where it co-localized with the membrane stain FM4-64

with some accumulations in the sub-membrane space (Fig. 3H-H''). While no membrane

protrusions were formed in the absence of bacteria, we observed these membrane outgrowths

upon inoculation of protoplasts with *S. meliloti* for 5 hours. They coincided with SYFO1

accumulations inside these structures and often formed adjacent to bacterial aggregates (Fig.

3I-I''). To unambiguously verify that SYFO1 is the key driver of these protrusions we isolated

protoplasts from our *syfo1-1* and *syfo1L-1* mutants. No protrusions were found in the absence

of rhizobia in any of the used genotypes. Upon inoculation of protoplasts with *S. meliloti*, those

expressing wild-type *SYFO1* (wild-type, *syfo1L-1*) or over-expressing SYFO1-GFP (OE)

developed protrusions while they were entirely absent on protoplasts isolated from the *syfo1-1*

mutant (Fig. 3J). This demonstrates that focal accumulation of SYFO1 can drive cell

polarisation and membrane deformations in a stimulus-dependent manner.

A SYFO1-mediated cell wall-plasma membrane-cytoskeleton continuum is required for symbiotic responses in root hairs.

As actin binding of formins is generally mediated by the FH2 domain that is also present in the cytosolic domain of SYFO1 (Fig. 1A'), we examined the extracellular region, which is less prominently found among formin proteins. Further sequence analysis of the SYFO1^{ECD} (extracellular domain) revealed the presence of a proline-rich repeat (PRR) (e.g. Ser-Pro-Pro-Pro-Ser-Pro-Ser-Ser [SPPPSPSS]) between SP and TMD, which resembles a canonical motif of extensin proteins that have been proposed to contribute to cell wall architecture and tensile strength [43, 44]. Thus we investigated a possible role in cell wall association of the ECD. For this, we fused the 82 amino acids of the SYFO1 ECD to GFP (SYFO1^{ECD}-GFP) and expressed it in *Nicotiana benthamiana* leaf epidermal cells. The fluorescent label was found in the cell periphery in control cells (Fig. 4A-A'') while it remained predominantly associated with the cell wall upon plasmolysis (Fig. 4B-B''). In order to test whether this extensin-like motif contributes to the lateral immobilization of SYFO1 via cell wall anchoring we performed Fluorescence Recovery After Photobleaching (FRAP) experiments on root hairs constitutively expressing a full-length SYFO1 (SYFO1-GFP) or a mutant variant where we deleted the proline-rich repeat (SYFO1$^{\Delta PRR}$-GFP). These experiments revealed a slow recovery of the bleached region and a mobile fraction of about 24% for wild-type SYFO1 whereas the mobile fraction for SYFO1$^{\Delta PRR}$ was significantly higher (57%) (Fig. 4C-E). This clearly indicates that the PRR segment within the SYFO1 ECD anchors this formin to the cell wall. To address whether the cell wall association is required for SYFO1 function we conducted genetic complementation experiments, we generated transgenic roots expressing SYFO1$^{\Delta PRR}$ under the control of the native *SYFO1* promoter in our *syfo1-1* and *syfo1-2* alleles. In contrast to the full-length SYFO1 (Fig. 2), the deletion of the PRR fully abolished the ability to complement the root hair deformation phenotype of *syfo1* mutants (Fig. 4F), demonstrating that a SYFO1-

mediated cell wall-plasma membrane-actin continuum is required for symbiotic responsiveness

of root hairs in *M. truncatula*.

DISCUSSION

The engulfment of a single rhizobium by a tightly curled root hair represents a fascinating

process that allows most legumes to keep full control over the onset of root hair infection and

might minimize the risk of infection by non-symbiotic bacteria. Considering that the formin

protein SYFO1 serves symbiotic but not general functions in root hair responsiveness (Fig. S5

and Fig. S6) demonstrates that Medicago genetically re-wired few components of the actin

machinery to drive root hair curling. However, the generic set of actin and polarity factors such

as NAP1/PIR1, members of the SCAR/WAVE complex [34], and RHO OF PLANTS 10 [45]

are additionally required for root hair growth *per se*. Consequently, these mutants show

impaired growth also under non-symbiotic conditions. In contrast, legumes specifically

evolved SCARN, another member of the SCAR/WAVE complex, which is required for

rhizobial infection but not for growth under fully fertilized conditions [35]. The fact that the

number of infection chambers was higher in Lotus *scarn* mutants compared to wild-type plants

[35], places SCARN downstream of SYFO1. Since the bacterial colonization phenotypes in

nodules are comparable between both mutants, both may contribute to actin function at this

stage.

SYFO1 with its transmembrane helix and the extracellular proline-rich repeat carries all

features of a group 1 formin [46]. As previously described for AtFH1 [47], the PRR mediates

cell wall association and consequently restricts the lateral mobility also in SYFO1 (Fig. 4E).

Its ability to initiate membrane protrusions in cell wall-depleted protoplasts further suggests

that SYFO1 is involved in actin nucleation and filament elongation (Fig. 3G-G'; Fig. S8). In

line with non-symbiotic formins, such as AtFH4 from Arabidopsis that re-localizes to infection

sites of the powdery mildew fungus *Blumeria graminis* [48], we hypothesize that SYFO1

evolved to specifically mediate targeted secretion of cell wall constituents and other cargo

material to sustain symbiotic root hair responses including root hair curling and later stages of

infection. This entirely depends on the ability of SYFO1 to associate with the cell wall (Fig. 4)

where it maintains a cell wall-plasma membrane-cytoskeleton continuum that cannot be

functionally complemented by other, non-symbiotic formins that remain being expressed in

syfo1 mutants.

MATERIALS AND METHODS

Plant growths and phenotypical analysis

For phenotypical analysis *Medicago truncatula* wild-type R108, *syfo1-1*, *syfo1-2*, *syfo1L-1* and *syfo1L-2* seeds were scarified and sterilized before being sown on 1% agar plates for germination and kept in darkness at 4°C for 3 days for vernalization. Germination was allowed for up to 24 hours at 24°C before transferring the seedlings to plates containing Fahraeus medium [49] for 4 days in the presence of 1 mM nitrate before being transferred to a plate culture system without nitrogen for phenotyping studies. Plants were inoculated with 1ml *Sinorhizobium meliloti* 2011 (mCherry) at an OD_{600} of 0.05 (on plates or open pots with 1:1 ratio of vermiculite and sand mixture). Symbiotic responses including root hair deformations, infection chamber formation and IT development were scored 5 dpi of plants with *S. meliloti* on plates. Soil-based nodulation phenotyping samples were harvested and quantified at 3 wpi. Nodules were embedded in 7% Agar and sectioned with a thickness of 60 μm using a vibratome.

Genotyping of *Tnt1* insertion lines and quantitative Real-Time PCR

R0 or R1 seeds of *M. truncatula* R108 *Tnt1* transposon insertion lines were obtained from the Noble Research Institute (OK, USA) and insertions were verified using primers listed in Table S4. Total RNA of control and insertion lines was extracted using a commercial kit (Spectrum™ Plant Total RNA Kit, Sigma life science) following the supplier's instructions. An additional DNaseI treatment was performed. Synthesis of cDNA and qRT-PCR were conducted as described earlier [50] using the SuperScriptIII reverse Transcriptase (Invitrogen). All data were normalized to Ct values of the housekeeping gene ubiquitin [51] using primers listed in Table S4.

Hairy root transformation and inoculation of rhizobia

M. truncatula hairy root transformation was performed as previously described [52] using the *Agrobacterium rhizogenes* strain ARqua1. Plants were transferred weekly to fresh plates containing Fahraeus medium (pH 6.0) supplemented with 0.5 mM NH_4NO_3 and followed by 2 days of growth on nitrogen-free Fahraeus medium containing 0.1 μM AVG prior to inoculation. Images for localisation studies and root hair phenotyping analyses were taken on plants inoculated for 2 days and 5 days, respectively.

Phylogenetic and selective pressure analysis

SYFO1 (Medtr5g036540.1) and SYFO1L (Medtr8g062830.1) protein sequences were used as queries for a tBLASTn v2.8.1+ (10.1186/1471-2105-10-421) search against a database of 101 Angiosperms genomes (Table S2, sequences can be downloaded from the SymDB database (10.1038/s41477-020-0613-7): http://www.polebio.lrsv.ups-tlse.fr/symdb) with an e-value threshold of 1e-10. Sequences were then aligned using MAFFT v7.407 (10.1093/molbev/mst010) with default parameters. The resulting alignment was trimmed using trimAl v1.4 rev22 (10.1093/bioinformatics/btp348) to remove positions containing more than 20% of gaps. The cleaned alignment was then subjected to a Maximum Likelihood (ML) analysis using IQ-TREE v1.6.7 (10.1093/molbev/msu300) as described here after. First, the best-fitting evolutionary model was tested using ModelFinder (10.1038/nmeth.4285). Then a ML search was performed using 10,000 replicates of SH-aLRT (10.1093/sysbio/syq010) for testing branches support. The tree was finally visualized and annotated with iTOL v4.4 (10.1093/nar/gkw290).

Signal peptide and transmembrane domains were predicted from proteins using signal v5.0 (10.1038/s41587-019-0036-z) and TMHMM v2.0c (PMID: 9783223G) respectively using default parameters.

To look for relaxation (K<1) or intensification (K>1) of selection acting on different lineages

of interest in Eudicots (Table S3), we used the RELAX program (10.1093/molbev/msu400).

This method calculates different synonymous and non-synonymous substitution rates (ω =

dN/dS) using the phylogenetic tree topology for both foreground and background branches.

Because (i) the programs used does not accept gaps in codon sequences and (ii) there is a

negative correlation between the number of sequences and the number of ungapped positions,

we used different numbers of input sequences for RELAX analysis. Protein sequences from

SYFO1 and SYFO1L orthologs were aligned using MUSCLE v3.8.382. Short sequences were

excluded to maximize sequence number while limiting gapped positions compared to SYFO1

and SYFO1L sequences of Medicago using a custom R script. We opted for 151 CDS

sequences corresponding to 1008 positions (Table S3).

Construct design

The constructs used in this study were designed using Golden Gate cloning [53]. 2.5 kb

upstream of the SYFO1 start codon were chosen as putative promoter region. A Golden Gate

compatible full-length genomic DNA version (Medtr5g036540.1) was synthesized

(GENEWIZ, Germany) by removing the *BpiI* and *BsaI* restriction sites via silent mutations.

All cloning primers are listed in Table S4. To select transgenic roots a *pUbi:NLS-mCherry* or

pUbi:NLS-2xCerulean cassette was additionally inserted into the different T-DNAs containing

the transgenes of choice as previously described [50]. Level II and level III constructs were

assembled based on the principle described earlier [53]. An overview about all designed

constructs is provided in Table S5.

Confocal Laser-Scanning Microscopy and FRAP

For imaging the NLS-GFP reporter module, sectioned nodules, protein localisation and plasmolysis we used a Leica TCS SP8 confocal microscope equipped with a 20x HCX PL APO water immersion lenses (Leica Microsystems, Mannheim, Germany). GFP was excited with a White Light Laser (WLL) at 488 nm and the emission was detected at 500-550 nm. mCherry fluorescence was excited using a WLL at 561nm and emission was detected between 575-630 nm. Samples, co-expressing two fluorophores were imaged in sequential mode between frames. FRAP analysis was conducted using a Zeiss LSM 880 Airyscan confocal microscope. For this, the VP (Virtual Pinhole) mode was adapted based on the fluorescence intensity of probe and Airyscan processing was performed using the ZEN (black edition) software package. The bleaching region, reference region and background region were selected at identical size. A pre-bleaching time of 5 seconds was chosen. Bleaching was set to stop upon the intensity dropping to 50% of the initial intensity before fluorescence recovery was recorded for 10 minutes. Using the FRAP data process package in ZEN (black version), the mobile fraction was calculated by the following equation (mobile fraction= $I1/ IE$), where $I1$ represents the dropped intensity and the IE represents the recovered intensity normalized to the reference intensity.

Root organ culture, protoplast extraction, and inoculation

Transgenic Root Organ Culture (ROC) of *M. truncatula* expressing SYFO1-GFP were obtained via hairy root transformation according to [52]. Fully transformed root segments were cut and initially transferred to M-Media plates (Becard & Fortin, 1988) containing Augmentin (Sigma). 1 g of amoxicillin–200 mg of clavulanic acid) (400 mg/L) for two weeks and then subcultured on plates supplemented with 200 mg/L Augmentin for additional two weeks to remove *A. rhizogenes* contamination. Plates were sealed with micropore tape and incubated at

24°C in dark. Afterwards, the ROC was transferred to M-media plates without Augmentin to

support faster tissue growth. Continuous expression of the transformation marker was

monitored throughout the entire experiment.

Protoplasts were isolated from ROC by cutting the roots in small pieces of 2-5 mm length and

processed as described previously [54]. For inoculation, an *S. meliloti* culture was diluted in

W5 solution [54] to an OD_{600} of 0.05 before being added to the protoplasts.

Actin phalloidin staining and plasma membrane FM4-64 staining

Phalloidin-based actin staining was performed according to a published protocol [31]. In brief,

Medicago roots were transferred into Fahraeus medium containing 300 μM MBS (m-

maleimidobenzoyl- N-hydroxysuccinimide ester) for 30 minutes to stabilize the actin filaments.

The material was then fixed in 2 % formaldehyde in actin-stabilizing buffer (ASB) solution

[34]. Phalloidin was added to a final concentration of 16 μM and staining was performed in the

dark for 30 minutes. Root-derived protoplasts were submerged in a FM4-64 solution with a

final concentration of 20 μM and incubated on ice for 5-10 mins before imaging.

ACKNOWLEDGEMENTS

This study was supported by the project Engineering Nitrogen Symbiosis for Africa (ENSA) currently supported through a grant to the University of Cambridge by the Bill & Melinda Gates Foundation (OPP1172165) and UK government's Department for International Development (DFID) as well as by the China Scholarship Council (CSC) (grant no. 201506350004 to PL). JK, CL and PMD belong to the LRSV, which is part of the TULIP LABEX (ANR-10-LABX-41). We thank the staff of the Life Imaging Center (LIC) in the Center for Biological Systems Analysis (ZBSA) of the Albert-Ludwigs-University of Freiburg for help with their confocal microscopy resources, and the excellent support in image recording. The *Medicago truncatula* plants utilized in this research project, which are jointly owned by the Centre National De La Recherche Scientifique, were obtained from Noble Research Institute, LLC and were created through research funded, in part, by a grant from the National Science Foundation, NSF-0703285. We are grateful to the genotoul bioinformatics platform Toulouse Occitanie (Bioinfo Genotoul, doi: 10.15454/1.5572369328961167E12) for providing computing resources.

We would explicitly thank all the members of our team for the fruitful discussions and for providing their individual expertise throughout the course of the project.

The authors declare no competing interests.

Author Contributions

Conceptualization, P.L. and T.O.; Investigation, P.L., C.S., B.L., F.A.D., J.K., C.L., P.M.D. and T.O.; Writing –Original Draft, P.L. and T.O.; Writing –Review & Editing, P.L., C.S., B.L., F.A.D., J.K., C.L., P.M.D. and T.O.; Funding Acquisition, P.L., P.M.D. and T.O.; Supervision, T.O.

REFERENCES

1. Mylona, P., Pawlowski, K., and Bisseling, T. (1995). Symbiotic Nitrogen-Fixation. Plant Cell *7*, 869-885.

2. Oldroyd, G.E.D., and Downie, J.M. (2008). Coordinating nodule morphogenesis with rhizobial infection in legumes. Annu Rev Plant Biol *59*, 519-546.

3. Brewin, N.J. (2004). Plant cell wall remodelling in the rhizobium-legume symbiosis. Crit Rev Plant Sci *23*, 293-316.

4. Gage, D.J. (2004). Infection and invasion of roots by symbiotic, nitrogen-fixing rhizobia during nodulation of temperate legumes. Microbiol Mol Biol R *68*, 280-+.

5. Esseling, J.J., Lhuissier, F.G.P., and Emons, A.M.C. (2003). Nod factor-induced root hair curling: Continuous polar growth towards the point of nod factor application. Plant Physiology *132*, 1982-1988.

6. Oldroyd, G.E.D., and Downie, J.A. (2004). Calcium, kinases and nodulation signalling in legumes. Nat Rev Mol Cell Bio *5*, 566-576.

7. Bibikova, T.N., Blancaflor, E.B., and Gilroy, S. (1999). Microtubules regulate tip growth and orientation in root hairs of Arabidopsis thaliana. Plant Journal *17*, 657-665.

8. Yang, G.H., Gao, P., Zhang, H., Huang, S.J., and Zheng, Z.L. (2007). A Mutation in MRH2 Kinesin Enhances the Root Hair Tip Growth Defect Caused by Constitutively Activated ROP2 Small GTPase in Arabidopsis. Plos One *2*.

9. Yi, K., Guo, C., Chen, D., Zhao, B., Yang, B., and Ren, H. (2005). Cloning and functional characterization of a formin-like protein (AtFH8) from Arabidopsis. Plant Physiol *138*, 1071-1082.

433   10.   Jones, M.A., Shen, J.J., Fu, Y., Li, H., Yang, Z.B., and Grierson, C.S. (2002). The

434   Arabidopsis Rop2 GTPase is a positive regulator of both root hair initiation and tip

435   growth. Plant Cell *14*, 763-776.

436   11.   Parker, J.S., Cavell, A.C., Dolan, L., Roberts, K., and Grierson, C.S. (2000). Genetic

437   interactions during root hair morphogenesis in Arabidopsis. Plant Cell *12*, 1961-1974.

438   12.   Carol, R.J., Takeda, S., Linstead, P., Durrant, M.C., Kakesova, H., Derbyshire, P., Drea,

439   S., Zarsky, V., and Dolan, L. (2005). A RhoGDP dissociation inhibitor spatially

440   regulates growth in root hair cells. Nature *438*, 1013-1016.

441   13.   Fournier, J., Teillet, A., Chabaud, M., Ivanov, S., Genre, A., Limpens, E., de Carvalho-

442   Niebel, F., and Barker, D.G. (2015). Remodeling of the infection chamber before

443   infection thread formation reveals a two-step mechanism for rhizobial entry into the

444   host legume root hair. Plant Physiology *167*, 1233-1242.

445   14.   Xie, F., Murray, J.D., Kim, J., Heckmann, A.B., Edwards, A., Oldroyd, G.E.D., and

446   Downie, A. (2012). Legume pectate lyase required for root infection by rhizobia. P Natl

447   Acad Sci USA *109*, 633-638.

448   15.   Gage, D.J., and Margolin, W. (2000). Hanging by a thread: invasion of legume plants

449   by rhizobia. Curr Opin Microbiol *3*, 613-617.

450   16.   Heckmann, A.B., Lombardo, F., Miwa, H., Perry, J.A., Bunnewell, S., Parniske, M.,

451   Wang, T.L., and Downie, J.A. (2006). Lotus japonicus nodulation requires two GRAS

452   domain regulators, one of which is functionally conserved in a non-legume. Plant

453   Physiology *142*, 1739-1750.

454   17.   Bek, A.S., Sauer, J., Thygesen, M.B., Duus, J.O., Petersen, B.O., Thirup, S., James, E.,

455   Jensen, K.J., Stougaard, J., and Radutoiu, S. (2010). Improved Characterization of Nod

456   Factors and Genetically Based Variation in LysM Receptor Domains Identify Amino

Acids Expendable for Nod Factor Recognition in Lotus spp. Molecular Plant-Microbe

Interactions *23*, 58-66.

18.   Madsen, L.H., Tirichine, L., Jurkiewicz, A., Sullivan, J.T., Heckmann, A.B., Bek, A.S.,

Ronson, C.W., James, E.K., and Stougaard, J. (2010). The molecular network

governing nodule organogenesis and infection in the model legume Lotus japonicus.

Nat Commun *1*, 10.

19.   Cerri, M.R., Wang, Q.H., Stolz, P., Folgmann, J., Frances, L., Katzer, K., Li, X.L.,

Heckmann, A.B., Wang, T.L., Downie, J.A., et al. (2017). The ERN1 transcription

factor gene is a target of the CCaMK/CYCLOPS complex and controls rhizobial

infection in Lotus japonicus. New Phytologist *215*, 323-337.

20.   Kawaharada, Y., James, E.K., Kelly, S., Sandal, N., and Stougaard, J. (2017). The

ethylene responsive factor required for nodulation 1 (ERN1) transcription factor is

required for infection-thread formation in Lotus japonicus. Molecular plant-microbe

interactions *30*, 194-204.

21.   Schauser, L., Roussis, A., Stiller, J., and Stougaard, J. (1999). A plant regulator

controlling development of symbiotic root nodules. Nature *402*, 191-195.

22.   Combier, J.P., Frugier, F., de Billy, F., Boualem, A., El-Yahyaoui, F., Moreau, S.,

Vernie, T., Ott, T., Gamas, P., Crespi, M., et al. (2006). MtHAP2-1 is a key

transcriptional regulator of symbiotic nodule development regulated by microRNA169

in Medicago truncatula. Gene Dev *20*, 3084-3088.

23.   Ben Amor, B. (2003). The NFP locus of Medicago truncatula controls an early step of

Nod factor signal transduction upstream of a rapid calcium flux and root hair

deformation (vol 34, pg 495, 2003). Plant Journal *35*, 140-140.

24. Limpens, E., Franken, C., Smit, P., Willemse, J., Bisseling, T., and Geurts, R. (2003).

LysM domain receptor kinases regulating rhizobial Nod factor-induced infection.

Science *302*, 630-633.

25. Madsen, E.B., Madsen, L.H., Radutoiu, S., Olbryt, M., Rakwalska, M., Szczyglowski,

K., Sato, S., Kaneko, T., Tabata, S., Sandal, N., et al. (2003). A receptor kinase gene of

the LysM type is involved in legume perception of rhizobial signals. Nature *425*, 637-

640.

26. Arrighi, J.F., Barre, A., Ben Amor, B., Bersoult, A., Soriano, L.C., Mirabella, R., de

Carvalho-Niebel, F., Journet, E.P., Gherardi, M., Huguet, T., et al. (2007). The

Medicago truncatula lysine motif-receptor-like kinase gene family includes NFP and

new nodule-expressed genes (vol 142, pg 265, 2006). Plant Physiology *143*, 1078-1078.

27. Smit, P., Limpens, E., Geurts, R., Fedorova, E., Dolgikh, E., Gough, C., and Bisseling,

T. (2007). Medicago LYK3, an entry receptor in rhizobial nodulation factor signaling.

Plant Physiology *145*, 183-191.

28. Radutoiu, S., Madsen, L.H., Madsen, E.B., Felle, H.H., Umehara, Y., Gronlund, M.,

Sato, S., Nakamura, Y., Tabata, S., Sandal, N., et al. (2003). Plant recognition of

symbiotic bacteria requires two LysM receptor-like kinases. Nature *425*, 585-592.

29. de Ruijter, N.C.A., Rook, M.B., Bisseling, T., and Emons, A.M.C. (1998). Lipochito-

oligosaccharides re-initiate root hair tip growth in Vicia sativa with high calcium and

spectrin-like antigen at the tip. Plant Journal *13*, 341-350.

30. Cardenas, L., Feijo, J.A., Kunkel, J.G., Sanchez, F., Holdaway-Clarke, T., Hepler, P.K.,

and Quinto, C. (1999). Rhizobium Nod factors induce increases in intracellular free

calcium and extracellular calcium influxes in bean root hairs. Plant Journal *19*, 347-

352.

31.   Miller, D.D., de Ruijter, N.C.A., Bisseling, T., and Emons, A.M.C. (1999). The role of
505   actin in root hair morphogenesis: studies with lipochito-oligosaccharide as a growth
506   stimulator and cytochalasin as an actin perturbing drug. Plant Journal *17*, 141-154.

507   32.   Miller, D.D., Klooster, H.B.L.-t., and Emons, A.M.C. (2000). Lipochito-
508   oligosaccharide nodulation factors stimulate cytoplasmic polarity with longitudinal
509   endoplasmic reticulum and vesicles at the tip in vetch root hairs. Molecular plant-
510   microbe interactions *13*, 1385-1390.

511   33.   Zepeda, I., Sanchez-Lopez, R., Kunkel, J.G., Banuelos, L.A., Hernandez-Barrera, A.,
512   Sanchez, F., Quinto, C., and Cardenas, L. (2014). Visualization of Highly Dynamic F-
513   Actin Plus Ends in Growing Phaseolus vulgaris Root Hair Cells and Their Responses
514   to Rhizobium etli Nod Factors. Plant Cell Physiol *55*, 580-592.

515   34.   Yokota, K., Fukai, E., Madsen, L.H., Jurkiewicz, A., Rueda, P., Radutoiu, S., Held, M.,
516   Hossain, M.S., Szczyglowski, K., Morieri, G., et al. (2009). Rearrangement of Actin
517   Cytoskeleton Mediates Invasion of Lotus japonicus Roots by Mesorhizobium loti. Plant
518   Cell *21*, 267-284.

519   35.   Qiu, L.P., Lin, J.S., Xu, J., Sato, S., Parniske, M., Wang, T.L., Downie, J.A., and Xie,
520   F. (2015). SCARN a Novel Class of SCAR Protein That Is Required for Root-Hair
521   Infection during Legume Nodulation. Plos Genet *11*.

522   36.   Zepeda, I., Sanchez-Lopez, R., Kunkel, J.G., Banuelos, L.A., Hernandez-Barrera, A.,
523   Sanchez, F., Quinto, C., and Cardenas, L. (2014). Visualization of highly dynamic F-
524   actin plus ends in growing phaseolus vulgaris root hair cells and their responses to
525   Rhizobium etli nod factors. Plant Cell Physiol *55*, 580-592.

526   37.   Yang, C., Czech, L., Gerboth, S., Kojima, S.I., Scita, G., and Svitkina, T. (2007). Novel
527   roles of formin mDia2 in lamellipodia and filopodia formation in motile cells. Plos
528   Biology *5*, 2624-2645.

38.   Larrainzar, E., Riely, B.K., Kim, S.C., Carrasquilla-Garcia, N., Yu, H.J., Hwang, H.J.,

Oh, M., Kim, G.B., Surendrarao, A.K., Chasman, D., et al. (2015). Deep Sequencing

of the Medicago truncatula Root Transcriptome Reveals a Massive and Early

Interaction between Nodulation Factor and Ethylene Signals. Plant Physiology *169*,

233-+.

39.   van Velzen, R., Holmer, R., Bu, F.J., Rutten, L., van Zeijl, A., Liu, W., Santuari, L.,

Cao, Q.Q., Sharma, T., Shen, D.F., et al. (2018). Comparative genomics of the

nonlegume Parasponia reveals insights into evolution of nitrogen-fixing rhizobium

symbioses. P Natl Acad Sci USA *115*, E4700-E4709.

40.   Oldroyd, G.E.D. (2013). Speak, friend, and enter: signalling systems that promote

beneficial symbiotic associations in plants. Nature Reviews Microbiology *11*, 252-263.

41.   Matusek, T., Gombos, R., Szecsenyi, A., Sanchez-Soriano, N., Czibula, A., Pataki, C.,

Gedai, A., Prokop, A., Rasko, I., and Mihaly, J. (2008). Formin proteins of the DAAM

subfamily play a role during axon growth. J Neurosci *28*, 13310-13319.

42.   Gombos, R., Migh, E., Antal, O., Mukherjee, A., Jenny, A., and Mihaly, J. (2015). The

Formin DAAM Functions as Molecular Effector of the Planar Cell Polarity Pathway

during Axonal Development in Drosophila. J Neurosci *35*, 10154-10167.

43.   Estevez, J.M., Kieliszewski, M.J., Khitrov, N., and Somerville, C. (2006).

Characterization of synthetic hydroxyproline-rich proteoglycans with arabinogalactan

protein and extensin motifs in Arabidopsis. Plant Physiol *142*, 458-470.

44.   Lamport, D.T., Kieliszewski, M.J., Chen, Y., and Cannon, M.C. (2011). Role of the

extensin superfamily in primary cell wall architecture. Plant Physiol *156*, 11-19.

45.   Lei, M.J., Wang, Q., Li, X.L., Chen, A.M., Luo, L., Xie, Y.J., Li, G., Luo, D., Mysore,

K.S., Wen, J.Q., et al. (2015). The Small GTPase ROP10 of Medicago truncatula Is

Required for Both Tip Growth of Root Hairs and Nod Factor-Induced Root Hair

Deformation. Plant Cell *27*, 806-822.

46.   Deeks, M.J., Hussey, P.J., and Davies, B. (2002). Formins: intermediates in signal-

transduction cascades that affect cytoskeletal reorganization. Trends in Plant Science *7*,

492-498.

47.   Martiniere, A., Gayral, P., Hawes, C., and Runions, J. (2011). Building bridges:

formin1 of Arabidopsis forms a connection between the cell wall and the actin

cytoskeleton. Plant Journal *66*, 354-365.

48.   Sassmann, S., Rodrigues, C., Milne, S.W., Nenninger, A., Allwood, E., Littlejohn, G.R.,

Talbot, N.J., Soeller, C., Davies, B., Hussey, P.J., et al. (2018). An Immune-Responsive

Cytoskeletal-Plasma Membrane Feedback Loop in Plants. Curr Biol *28*, 2136-2144

e2137.

49.   Fahraeus, G. (1957). The Infection of Clover Root Hairs by Nodule Bacteria Studied

by a Simple Glass Slide Technique. J Gen Microbiol *16*, 374-&.

50.   Liang, P., Stratil, T.F., Popp, C., Marin, M., Folgmann, J., Mysore, K.S., Wen, J., and

Ott, T. (2018). Symbiotic root infections in Medicago truncatula require remorin-

mediated receptor stabilization in membrane nanodomains. Proc Natl Acad Sci U S A

115, 5289-5294.

51.   Satge, C., Moreau, S., Sallet, E., Lefort, G., Auriac, M.C., Rembliere, C., Cottret, L.,

Gallardo, K., Noirot, C., Jardinaud, M.F., et al. (2016). Reprogramming of DNA

methylation is critical for nodule development in Medicago truncatula. Nature Plants *2*.

52.   Boisson-Dernier, A., Chabaud, M., Garcia, F., Becard, G., Rosenberg, C., and Barker,

D.G. (2001). Agrobacterium rhizogenes-transformed roots of Medicago truncatula for

the study of nitrogen-fixing and endomycorrhizal symbiotic associations. Mol Plant

Microbe Interact *14*, 695-700.

53.   Binder, A., Lambert, J., Morbitzer, R., Popp, C., Ott, T., Lahaye, T., and Parniske, M.

(2014). A Modular Plasmid Assembly Kit for Multigene Expression, Gene Silencing

and Silencing Rescue in Plants. Plos One *9*.

54.   Jia, N., Zhu, Y.L., and Xie, F. (2018). An Efficient Protocol for Model Legume Root

Protoplast Isolation and Transformation. Frontiers in Plant Science *9*.

Figures and Legends

Figure 1. SYFO1 but not SYFO1L is required for nodulation. (*A-A'*) Isolation of independent mutant alleles
with *Tnt1* transposon insertions mapping to different regions of the *SYFO1* and *SYFO1L* gene (*A*) and protein (*A'*)
models, the extracellular domain is indicated as ECD and the orange box indicated the location of proline-rich
repeat (PRR). (*B-B''*) Nodule phenotypes observed on the different genotypes at 3 wpi with *S. meliloti* in open
pots using WT R108 plants as a control. Scale bars indicate 200 μm. (*C-D*) Quantification of nodule numbers (*C*)
and the ratio of aborted/wild-type like nodules (*D*) (n=10). Asterisks indicate a significant statistical difference
based on a Tukey–Kramer multiple-comparison test with p-values <0.01 (**), < 0.001 (***). Data are shown as
mean ± SE.

Figure 2. *syfo1* mutants are impaired in symbiotic root hair responses. Images show deformed root hairs (*A*), infection chambers (ICs) and infection threads (ITs) (*B*) on wild-type plants to illustrate the scored structures. Scale bars indicate 10 μm. *syfo1* mutants show significantly reduced responsiveness to the presence of compatible rhizobia when assessing root hair deformations (*C*) and infection-related structures such as ICs + ITs (*E*). These phenotypes were complemented by introducing a genomic version of a full-length *SYFO1* gene driven by the endogenous *SYFO1* promoter (*syfo1-1c, syfo1-2c*) with an empty vector (ev) transformation control aside (*D and F*). Asterisks indicate a significant statistical difference based on a Tukey–Kramer multiple-comparison test with p-values <0.01 (**), < 0.001 (***). Data are shown as mean ± SE with independent 9-14 plants for phenotypical analysis, and 10 plants for complementation analysis.

Figure 3. SYFO1 functions as a symbiotic polarity factor in root hairs. SYFO1-GFP localizes homogeneously
to the PM of root hairs under mock conditions (*A*). At 2 dpi with *S. meliloti*, SYFO1 transiently polarizes at
subapical (*B*) and apical regions (*C*) of root hairs, before distributing equally along the PM during root hair
deformation (*D*) and curling (*E*). The arrowhead marks cell wall autofluorescence around the IC (*E and F*).
SYFO1 remains on the IT membrane (*F* and inset *F'*). Co-localisation between SYFO1-GFP and actin marker
ABD2:mCherry (*G-G'*). The nucleus encircled with a dash line in *G-G'* is based on corresponding transmitted
light image (not shown). The arrowheads point towards actin bundles orienting towards a nucleation centre at the
apical shank of the root hair. Scale bars indicate 10 μm. Protoplasts of a ROC expressing SYFO1 and
counterstained with the styryl dye FM4-64 remain uniformly round at 0 dpi (*H-H''*) while focal membrane
protrusions with accumulated SYFO1 were found at 5 hpi with Ds-Red expressing *S. meliloti* (*I-I''*). Scale bars
indicate 5 μm. (*J*) Quantification of membrane protrusions in protoplasts using different genetic backgrounds.
Asterisks indicate a significant statistical difference based on an ANOVA followed by a Fisher LSD test, with p-
values <0.05 (*), < 0.01 (**). Data are shown as mean ± SE of 3 independent biological replicates with *n* indicating
the total number of protoplasts being scored.

626
627
Figure 4. Cell wall association of SYFO1 is essential for its function. The constitutively expressed ECD of
SYFO1 labelled the cell periphery in non-plasmolysed cells (*A-A''*) and remained at the cell wall upon plasmolysis
(*B-B''*). Arrowheads and arrows mark the cell wall and the retracted plasma membrane, respectively. FRAP
experiments on roots hairs revealed a low mobility of full-length SYFO1 (*C-C'''*) while deletion of the PRR
resulted in an increased mobility of the protein (*D-D'''*). Scale bars indicate 10 μm. (*E*) Quantification of the
mobile fractions of SYFO1 (n=17) and SYFO1$^{\Delta PRR}$ (n=12), asterisks indicate a significant statistical difference
based on a student t test. (*F*) The SYFO1$^{\Delta PRR}$ variant failed to genetically complement both *syfo1* mutant alleles
in comparison to roots transformed with the empty vector (ev) scoring n=10 independent root systems per
genotype. Asterisks indicate a significant statistical difference based on a Tukey–Kramer multiple-comparison
test with p-values <0.001 (***), p<0.01 (**). Data are shown as mean ± SE.
638
639