Suppression of chromosome instability limits acquired drug resistance

Elizabeth A Crowley[1], Nicole M Hermance[1], Conor P Herlihy[1], Amity L Manning[1*]

[1] Department of Biology and Biotechnology, Worcester Polytechnic Institute, Worcester, MA, 01609 USA

[*]Corresponding author:
Amity L Manning
Email: almanning@wpi.edu
tel: 508-831-4961

The authors declare no potential conflicts of interest.

Key words:
Mitosis, genomic instability, Aurora B, Gefitinib, NSCLC

Running Title: Suppression of CIN limits drug resistance

Statement of Significance:

Modulation of microtubule dynamics in cells that exhibit chromosome instability (CIN) is sufficient to promote mitotic fidelity, reduce genomic heterogeneity, and limit acquisition of drug resistance.

Abstract:

Numerical chromosome instability, or nCIN, defined as the high frequency of whole chromosome gains and losses, is prevalent in many solid tumors. nCIN has been shown to promote intra-tumor heterogeneity and corresponds with tumor aggressiveness, drug resistance and tumor relapse. While increased nCIN has been shown to promote the acquisition of genomic changes responsible for drug resistance, the potential to modulate nCIN in a therapeutic manner has not been well explored. Here we assess the role of nCIN in the acquisition of drug resistance in non small cell lung cancer. We show that generation of whole chromosome segregation errors in non small cell lung cancer cells is sensitive to manipulation of microtubule dynamics and that enhancement of chromosome cohesion strongly suppresses nCIN and reduces intra-tumor heterogeneity. We demonstrate that suppression of nCIN has no impact on non small cell lung cancer cell proliferation *in vitro* nor in tumor initiation in mouse xenograft models. However, suppression of nCIN alters the timing and molecular mechanisms that drive acquired drug resistance. These findings suggest mechanisms to suppress nCIN may serve as effective co-therapies to limit tumor evolution and sustain drug response.

Introduction

Whole chromosome instability, or numerical CIN (nCIN), is generated by underlying defects in

mitosis [1, 2]. The aneuploidy that results from mitotic segregation errors promotes intra-tumor

heterogeneity and is a driving force in cancer that contributes to tumor evolution and drug

resistance [1-7]. Chromosome segregation is exquisitely sensitive to the regulation of dynamic

microtubule attachments and defects that either increase or decrease the stability of microtubule

attachments can corrupt mitotic fidelity and contribute to nCIN [8, 9]. Conversely, perturbations

that reduce nCIN are proposed to limit acquired drug resistance and may hold therapeutic

potential.

Aurora B kinase is a master regulator of kinetochore-microtubule dynamics during mitosis. Its

overexpression is common in many cancer contexts and recent analyses of over 10,000 cancer

genomes from the Cancer Genome Atlas show that, across cancer subtypes, Aurora B

expression corresponds with degree of aneuploidy [10]. Increased expression of Aurora B also

correlates with poor patient prognosis in a variety of cancer contexts [11]. Aurora B is a

component of the Chromosome Passenger Complex (CPC). The CPC localizes to the

centromere where Aurora B kinase activity regulates localization and activity of numerous

kinetochore components responsible for binding and stabilizing kinetochore microtubule

attachments, thereby promoting satisfaction of the spindle assembly checkpoint and regulating

chromosome segregation [12]. Consistent with this function, both decreased and increased

Aurora B activity at the centromere results in chromosome segregation errors [13-18].

Here we demonstrate that in a panel of non small cell lung cancer (NSCLC) cell lines, where

Aurora B is highly expressed and rates of chromosome segregation are high, the experimental

reduction of centromere-localized Aurora B levels, or the independent stabilization of

kinetochore microtubule attachments, is sufficient to minimize genomic heterogeneity and limit

whole chromosome amplifications. We additionally find that suppression of nCIN limits acquired

drug resistance both *in vitro* and *in vivo*, and that sustained nCIN is rate limiting for tumor

relapse.

Materials and Methods

Cell culture, protein expression and depletion

PC9, H1299, A549, and HBEC3-KT (ATCC) cells were grown in RPMI1640 (Gibco, #61870036)

medium supplemented with 10% fetal bovine serum (Sigma, #16140071) and 1%

penicillin/streptomycin (Gibco, #15070063). All cells were cultured at 37°C and 5% CO_2. High

resolution immunofluorescence imaging with DNA stain (DAPI) is used to monitor and confirm

cell lines are free of *Mycoplasma* contamination. For short term knock down of targets, 50nM

pool siRNAs (non-targeted pool or a pool of 4 siRNAs to Wapl; Dharmacon-ON-TARGETplus

Human Wapl siRNA-SMARTpool) were transfected using RNAiMax (ThermoFisher, #13778100)

transfection reagent according to the manufacturer's instruction, or alternatively infected with a

lentiviral construct containing a shRNA hairpin for constitutive (pLK0.1-Puro) or doxycycline

inducible (tet-pLK0-Puro) depletion. Stable hairpin-expressing clones were selected with

Puromycin for 7-10 days. Induced depletion was achieved by the addition of 2μg/ml doxycycline

for a minimum of 48 hours.

Immunofluorescence and Fluorescence *in situ* hybridization

Cultured cells or single cell suspensions derived from tumors were grown on coverslips, fixed,

and stained for Aurora B (BD Biosciences, #611082), ACA (Antibodies Inc, #15-234), tubulin

(Santa Cruz, #5286), CENPA S7p (Cell Signaling, #2187), and/or CENPA (Enzo Life Sciences,

ADI-KAM-CC006-E) as previously described [19, 20]. To assess kinetochore microtubule

stability cells were incubated in ice cold RPMI for 5 minutes (PC9 and A549) or 10 minutes

(H1299) prior to fixation and staining. Fluorescence *in situ* hybridization with centromeric probes

and quantification of numerical heterogeneity was performed as previously described in [21]

using alpha satellite-specific probes for chromosomes 2,6,7,8 and 10 (Cytocell, #LPE002R,

006G, 007G, 008R, 010R). For FISH from tumors, single cell suspensions were made by gently

grinding tumors between two glass slides and culturing with continued drug treatment for 3-5

days. Cells were then collected and processed as above. Clonal Numerical Heterogeneity (NH)

of a population was determined by scoring >300 cells per clone/tumor for copy number for

chromosome probe (assessed in pairs) to determine the modal copy number, and the fraction of

cells deviating from that number (i.e., the numerical heterogeneity). A given copy number with

greater than 20% prevalence in a population was considered to be a stable subclone and not

included in the numerical heterogeneity score. Data is represented for individual chromosomes,

or alternatively, as average NH across all chromosomes measured for a given sample. A

students' 2-tailed t-test was used to determine statistical significance.

***In vivo* tumor growth and relapse assays**

Eight-week old male and female Crl:NU-*Foxn1^{nu}* mice (stock #088) were purchased from

Charles River Laboratory and maintained in a pathogen-free facility. All animal experiments

were performed in accordance with institutional regulations after protocol review and approval

by Worcester Polytechnic Institute's Institutional Animal Care and Use Committee.

For *in vivo* growth assays, three cohorts of 5 mice each were injected subcutaneously in the

flank with 5×10^6 PC9 cells. 2 cohorts received PC9 cells expressing a tetracycline-inducible

shWapl expression construct. One of these two cohorts was administered 2 μg/ml doxycycline

in the drinking water. All three cohorts were monitored daily and tumor size measured 3x/week.

Mice were humanly sacrificed, and tumor tissue collected when tumor volume reached 300mm^3.

For tumor relapse studies, tumor xenografts were generated as described above in 40 mice.

Xenograft tumor formation was observed in 38 mice. Once tumors reached a size of 300mm^3

mice were randomized equally into a minus or plus doxycycline group. All mice were given

50mg/kg Gefitinib (Selleckchem, #ZD1839) by oral gavage following a 5 day on and 2 day off

cycle. Mice were monitored daily and tumor size measured 3x/week to monitor tumor regression

and relapse. Once relapsed tumors reached a size of 300mm^3 mice were humanly sacrificed,

tumors harvested and frozen or cultured for subsequent experimental analysis.

Please see supplemental material for extended methods.

Data Availability

The data generated in this study are available within the article and its supplementary data files.

Results

Chromosome segregation errors in NSCLC cells correspond with high expression and

enhanced mitotic centromere localization of Aurora B

Non small lung cancer (NSCLC) cells frequently exhibit high levels of whole chromosome

instability (nCIN), a feature that has been experimentally linked to acquired drug resistance [1,

3-6, 22]. To understand the molecular changes that correspond with segregation errors and

nCIN in NSCLC cells we first identified a panel of NSCLC cell lines that exhibit mitotic defects

and chromosome copy number heterogeneity consistent with nCIN (A549, H1299, and PC9;

[21]). Using immunofluorescence microscopy to assess chromosome segregation we confirm

that NSCLC lines A549, H1299, and PC9 all exhibit high rates of lagging chromosomes during

anaphase that are indicative of mitotic chromosome segregation errors (Figure 1A). Frequent

high rates of whole chromosome segregation errors during mitosis contribute to genomic

heterogeneity within a cell population that can be assessed using FISH-based approaches to

measure population-level numerical heterogeneity (NH) for individual chromosomes. Using

centromeric probes for chromosomes 6, 7, 8, and 10 we find that, consistent with frequent

Figure 1: Enrichment of cohesion in NSCLC cells decreases Aurora B localization and activity at the centromere A) Quantification of anaphase lagging chromosomes in NSCLC cell lines B) NSCLC cell populations labelled with centromere enumeration FISH probes for chromosomes 6, 7, 8, and 10 and quantified for numerical heterogeneity. A minimum of 300 cells for each chromosome probe were scored and averaged per cell line. C-E) Representation and quantification of centromere localized Aurora B, CENPA (total) and pCENPA (pSer7) in PC9 cells treated with non-targeting (siControl) or Wapl specific (siWapl) siRNA. A minimum of 90 kinetochore pairs were measured (3/cell for 30 cells), for each of 3 biological replicates. Insets are of individual kinetochore pairs at 4x magnification. Scale bars are 5μm. Here and throughout *: p< 0.05, **: p<0.01; ***: p<0.001.

mitotic errors, populations of A549, H1299 and PC9 cells exhibit NH scores ranging from ~7-

25% (Figure 1B).

Aurora B kinase, an important regulator of mitotic chromosome segregation and the mitotic

spindle assembly checkpoint, is commonly overexpressed in non small cell lung cancer, where

nCIN is also prevalent [23-25]. Consistent with these earlier studies, we find that Aurora B

kinase is highly expressed in our panel of NSCLC cells, when compared to the normal lung

epithelial cell line HBEC3-KT (Supplemental Figure 1A). Aurora B regulation of mitotic

chromosome segregation is largely dependent on its localization at centromeres where it

phosphorylates key substrates that regulate the stability of kinetochore-microtubule attachments

[26-30] Importantly, we find that centromere localization of Aurora B in our panel of NSCLC cell

lines mirrors overall expression levels of this kinase such that Aurora B localization at

centromeres is increased, on average, two to five-fold over that seen in HBEC3-KT cells

(Supplemental Figure 1B).

Centromere localization and activity of Aurora B in NSCLC is sensitive to cohesion

Aurora B localization is sensitive to changes in cohesin regulation such that redistribution of the

cohesin complex from pericentromere enrichment to distribution along chromosome arms leads

to a concurrent redistribution of Aurora B along chromosome arms and a reduction in its

centromere localization [19, 31-33] Therefore, to test the impact of high centromere Aurora B

levels in the mitotic defects observed in these NSCLC lines without altering overall Aurora B

protein levels, we first experimentally manipulated the distribution of the cohesin complex.

Enhancement of chromosome cohesion along chromosome arms was achieved using si- and

sh-RNA approaches to deplete Wapl, a well-characterized negative regulator of the cohesin

complex [34] (Supplemental Figure 2A & B). Wapl associates with the cohesin complex to

regulate its dynamic association with chromatin throughout the cell cycle [35] such that Wapl

depletion blocks cohesin complex dissociation from chromosomes during early mitosis. We

used immunofluorescence imaging of metaphase cells to identify and measure inter-kinetochore

distances as a readout of functional centromere cohesion. This demonstrated that Wapl

knockdown enhanced cohesion, as evidenced by a reduction in interkinetochore distance

(Supplemental Figure 2C).

Using quantitative immunofluorescence, we find that depletion of Wapl perturbs Aurora B

localization such that centromere-localized Aurora B levels in all three NSCLC lines is reduced

by ~50% when Wapl is depleted (Figure 1C and Supplemental Figure 3A). Western blot

analysis of mitotic cells indicate that this change in Aurora B localization does not arise due to

Wapl-dependent changes in Aurora B expression (Supplemental Figure 2B). Consistent with

reduced centromere localization and activity of Aurora B, we see comparable reduction in the

phosphorylation, but not localization, of key Aurora B substrate CENPA (Total and CENPA pS7:

Figure 1D & E, Supplemental Figure 3B & C).

Modulation of Aurora B activity suppresses kinetochore-microtubule dynamics and

chromosome segregation errors

Aurora B kinase is a master regulator of microtubule dynamics such that Aurora B localization

and kinase activity promotes the dynamic turnover of kinetochore microtubule attachments. The

fidelity of chromosome segregation has been shown to be sensitive to both increased and

decreased Aurora B activity, and hence corresponding decreased and increased stability of

kinetochore-microtubule attachments [8, 16, 17, 36, 37]. Consistent with this activity, we find

that kinetochore fibers in Wapl-depleted cells, where Aurora B localization at centromeres had

been reduced, exhibit decreased sensitivity to cold-induced microtubule depolymerization

compared to control cells (Figure 2A). This suggests kinetochore microtubule attachments are

stabilized by the redistribution of Aurora B away from the centromere that occurs following Wapl

depletion. We next assessed mitotic fidelity in these cells and find that enhancement of

chromosome arm cohesion, via Wapl depletion, is sufficient to reduce the incidence of lagging

chromosomes during anaphase in both H1299 and PC9 cells (Figure 2C). Similarly, partial

Figure 2: Anaphase segregation errors in NSCLC cells are reduced by microtubule stabilizing perturbations A & B) Representation and quantification of cold stable microtubules in PC9 cells depleted of Wapl (A) or treated with 5nM Taxol (B). C) Representation and quantification of anaphase lagging chromosomes in NSCLC cells following Wapl depletion, or treatment with 5nM Taxol, or 10nM Hesperdain. White arrow heads indicate individual lagging chromosomes. A minimum of 30 metaphase or anaphase cells were scored per population for each of 3 biological replicates. Scale bars are 5µm.

inhibition of Aurora B activity with the small molecule inhibitor Hesperdain, is sufficient to

suppress mitotic defects in all three NSCLC lines (Figure 2C).

To directly test the role of microtubule dynamics in the high frequency of mitotic errors observed

in this panel of NSCLC cells, we treated cells with a microtubule-stabilizing concentration of

Taxol prior to assessing mitotic fidelity. We find that following acute treatment (1h) with 5nM

Taxol, or long-term treatment (6 weeks) with 1nM Taxol, cells exhibit both resistance to cold-

induced microtubule depolymerization and reduction in anaphase lagging chromosomes,

comparable to that seen following Wapl depletion (Figure 2B & C, Supplemental Figure 4A & B).

To next assess if enhanced cohesion and reduction of anaphase defects is sufficient to

suppress nCIN, PC9 cells were engineered to constitutively express one of two different shRNA

hairpin constructs designed to target Wapl mRNA for depletion. Single cell clones were derived

from parental and Wapl-deficient PC9 cells and analyzed for numerical heterogeneity (NH).

Individual clones derived from PC9 cells expressing an empty PLK0.1 vector exhibit NH values

of 18-31% for chromosome 6, and up to 20-40% for chromosome 2. NH for both chromosomes

was reduced ~2-3-fold in Wapl-deficient PC9 cell clones (chromosome 6: 3-12%, chromosome

2: 5-24%) (Supplemental Figure 2D), indicating that enhanced cohesion is sufficient to suppress

nCIN. Together, these data support a model whereby increased Aurora B activity and/or highly

dynamic kinetochore microtubule attachments underlie nCIN in NSCLC cells.

Suppression of CIN limits drug tolerance in NSCLC cells.

Activating mutations within the EGFR gene that drive tumor cell proliferation are present in

nearly a quarter of all NSCLCs [38]. Patients with such mutations are commonly treated with

EGFR tyrosine kinase inhibitors (TKIs) [39]. However, the majority of patients treated with

EGFR TKIs ultimately develop resistance with nearly 60% of resistant or relapsed tumors

exhibiting resistance-conferring mutations in the EGFR gene, making this the most frequent

mechanism of EGFR TKI resistance [40]. PC9 cells exhibit an activating deletion in exon 19 of

EGFR that drives cell proliferation and renders them sensitive to EGFR TKIs [41]. Like tumors in

patients, these cells commonly acquire resistance to EGFR TKIs via acquisition of a secondary

Threonine to Methionine mutation in EGFR (T790M) [40]. While nCIN in NSCLC, and other

cancer contexts, has been correlated with acquired drug resistance [5, 23, 42], the impact of

nCIN on mutation-based mechanisms of acquired drug resistance remain unclear.

Drug response is sensitive to cell proliferation rates and the impact of aneuploidies that result

from nCIN have alternatively been demonstrated to promote tumor cell growth or to reduce

proliferative capabilities [43, 44]. Therefore, we first assessed proliferative capacity of PC9 cells

with and without constitutive nCIN/Wapl depletion (Mock and Wapl KD, respectively) or

following acclimation to microtubule-stabilizing concentrations of Taxol. Importantly, in all clones

tested, proliferation rates with or without Wapl depletion or Taxol treatment, when grown in the

absence of TKI treatment, were comparable (Supplemental Figure 4C-E). Similarly, anchorage

independent colony formation assays of growth in soft agar revealed similar colony number and

size, irrespective of Wapl/nCIN status (Supplemental Figure 4F), indicating that suppression of

nCIN alone does not alter drug naïve PC9 cell growth.

Next we assessed the response of cells, with and without Wapl depletion or Taxol treatment, to

the EGFR TKI Gefitinib. EGFR activity results in phosphorylation of EGFR and downstream

targets (such as ERK) and promotes cell proliferation. Western blot analyses indicate that PC9

cells with either mock or induced Wapl depletion or treated with 1nM Taxol are initially similarly

responsive to Gefitinib: all populations show dramatic reduction of EGFR-dependent

phosphorylation (phospho EGFR: Tyr1068 & phospho ERK: Tyr 202/Tyr 204) (Figure 3A).

Nevertheless, following 4 weeks of continuous treatment with a sub-lethal dose of Gefitinib

drug-tolerant cells slowly form colonies that can be detected with crystal violet stain. Following

long-term exposure to Gefitinib, Wapl-depleted and Taxol-treated PC9 cells exhibit a dramatic

reduction in the number and size of drug tolerant colonies that arise while under Gefitinib

treatment (Figure 3B & C).

Figure 3: Suppression of nCIN limits drug tolerance A) Total and phosphorylated levels of EGFR and ERK in PC9 cells with induced depletion of Wapl or 1nM Taxol treatment following Gefitinib treatment. B & C) Crystal Violet staining and quantification of drug tolerant colonies of control and Wapl depleted PC9 cells following extended culture of Wapl depleted (B) or Taxol treated (C) cells in presence of 1mM Gefitinib D) Crystal Violet staining and quantification of drug tolerant colonies of control and Wapl depleted NSCLC cells following extended culture in the presence of Vincristine or Taxol. Representative images and quantification reflect data from three independent biological replicates.

Acquired resistance to other clinically relevant therapeutics such as Vincristine and Paclitaxel

have been reported in NSCLC [45-50]. Therefore, to next test whether the impact of

suppression of CIN on drug tolerance is specific to Gefitinib or instead may similarly limit the

outgrowth of cells tolerant to other therapeutic agents, we performed *in vitro* drug tolerance

assays in A549, H1299, and PC9 cells with and without constitutive depletion of Wapl in the

presence of either Vincristine or Paclitaxel. Over the course of 14 weeks the concentration of

these drugs was doubled every 4 days until drug tolerant cells emerged. Comparative measures

of cell numbers in each condition were approximated using crystal violet staining. Following

long-term treatment with either Vincristine or Paclitaxel, all three NSCLC lines constitutively

depleted of Wapl show reduction in the number of drug-tolerant cells (Figure 3D). Importantly,

cells constitutively depleted of Wapl for the duration of the drug sensitivity assay show no

change in proliferation rates, suggesting decreased tolerance to Vincristine or Paclitaxel cannot

be accounted for by changes in proliferation rate following Wapl depletion (Supplemental Figure

4E). These data suggest that nCIN permits acquisition of drug tolerance to promote continued

proliferation.

Chromosome Instability informs mechanism of TKI drug resistance.

To better understand the relationship between nCIN-dependent genomic changes that may

promote or permit resistance to drug therapy we characterized four drug-tolerant clones from

each of the parental and Wapl-depleted populations that were exposed to long-term Gefitinib

treatment. Clones were selected and expanded under culture conditions that maintained 1μM

Gefitinib and mock or induced Wapl depletion, as appropriate. Importantly, in both Wapl-

depleted and Wapl-proficient contexts, drug tolerant PC9 cells that persist following long term

exposure to Gefitinib continue to exhibit features consistent with Wapl status: single colonies

derived from Mock-depleted cells have frequent anaphase defects and a high measure of NH

and whole chromosome amplifications while Wapl-depleted cells have reduced anaphase

defects and less numerical heterogeneity (Figure 4A-C).

Figure 4: Suppression of nCIN influences mechanisms of drug tolerance in PC9 cells. A) Quantification of anaphase cells exhibiting lagging chromosomes in clonal populations of Gefitinib-tolerant cells with (Wapl KD) or without (Mock) induced Wapl depletion. B & C) Gefitinib-tolerant PC9 clonal populations derived with or without Wapl depletion labelled with centromere enumeration FISH probes for chromosomes 6, 7, 8, and 10 and quantified for chromosome copy number. Represented as average numerical copy number across all chromosomes assessed (B) and fraction of cells with modal or amplified copies of the indicated chromosomes (C). A minimum of 300 cells per condition per centromere probe assessed were scored. D) Western blot showing Wapl levels and EGFR pathway status in four each of mock and Wapl-depleted Gefitinib-tolerant clones. E) qPCR analysis of mRNA levels indicative of EMT pathway activation (Slug, Vimentin) in Mock and Wapl-depleted Gefitinib-tolerant clones.

Cells that acquire tolerance or resistance to TKI activity do so by restoring or bypassing EGFR

kinase function to activate downstream pathway components and promote proliferation [40].

Although the emergence of drug tolerant clones is decreased significantly by Wapl depletion,

molecular characterization of EGFR pathway function indicates that drug tolerant clones that do

arise from both Mock- and Wapl-depleted populations similarly exhibit phosphorylation of ERK,

a downstream target in the EGFR pathway (Figure 4D). In contrast, Wapl-depleted PC9 clones,

but not Mock-depleted PC9 clones, exhibit EGFR auto phosphorylation (Figure 4D). This

phosphorylation is dependent on EGFR kinase activity and suggests drug-tolerant clones that

lack nCIN (Wapl KD) primarily arise through the acquisition of resistance-conferring mutations

that impair TKI binding, while clones that sustain nCIN (Mock) likely exploit other mechanisms of

drug tolerance [38-40, 47].

Alternative pathways reported to promote TKI resistance in NSCLC include amplification of

EGFR, amplification of the receptor tyrosine kinase MET [47], and/or activation of epithelial to

mesenchymal transition (EMT) [51-53]. To understand if these alternative pathways could

explain tolerance of Mock- depleted PC9 clones to Gefitinib treatment, we first assessed copy

number of chromosome 7, on which both EGFR and MET genes are located. Drug naïve PC9

cells have a modal copy number of four for chromosome 7, as do drug tolerant Wapl-depleted

clones. In contrast, nearly 60% of drug tolerant Mock-depleted cells have >8 copies of

chromosome 7 (Figure 4C, Supplemental Figure 5). We next looked at expression of key EMT

transcription factor SLUG and found it to be upregulated 2-fold in 3 out of 4 Mock-depleted

clones, but not in any of the Wapl-depleted clones (Figure 4E). The gene that encodes for

SLUG, SNAI2, is located on chromosome 8p, adjacent to the centromere. In support of a model

whereby CIN influences the mechanism of TKI resistance, single cell analysis of chromosome 8

copy number indicates that the same 3 Mock-depleted clones with increased SLUG expression

also exhibit amplification of chromosome 8 (Figure 4C, Supplemental Figure 5). Consistent with

SNAI2 amplification and increased Slug expression, Vimentin, a widely used marker of EMT

and a Slug target gene is also increased 2 to 12-fold in Mock-depleted clones (Figure 4E).

Together these data suggest that nCIN influences the mechanism of acquired drug resistance.

Without nCIN, mutation-driven mechanisms of drug resistance dominate, but the acquisition of

such mutations are slow, particularly in drug treated populations with limited proliferation, and as

a result drug resistant colonies are slow to emerge. Through whole chromosome gains and

losses hundreds to thousands of genes may become mis-regulated in a single cell division. In

this way nCIN has the capacity to promote cellular changes that contribute to drug tolerance.

Wapl depletion is sufficient to suppress nCIN *in vivo*

To assess how cohesion-dependent suppression of nCIN impacts tumor initiation and growth *in*

vivo, PC9 cells with or without a tetracycline-inducible shWapl construct were injected

subcutaneously into the flanks of nude mice to generate 3 cohorts: one with parental PC9 cells,

and two with PC9 cells that harbor the inducible shWapl construct (Supplemental Figure 6A).

One cohort of PC9 shWapl mice received doxycycline in the drinking water to induce expression

of the Wapl-targeting hairpin throughout the duration of tumor initiation and growth. The cohort

injected with the PC9 parental cell line similarly received doxycycline in the water as a negative

control. Tumor initiation was comparable in all three cohorts of mice, as was rate of tumor

growth (Supplemental Figure 6B, Supplemental Figure 7A). To confirm the efficiency of Wapl

depletion and the suppression of nCIN in response to doxycycline treatment *in vivo*, tumors from

mice in each cohort were excised when they reached $300cm^3$ and subjected to analysis.

Quantitative PCR analyses confirm that depletion of Wapl mRNA and protein levels were

achieved and sustained during tumor initiation in response to doxycycline (Supplemental Figure

6C). Consistent with *in vitro* assays, cells derived from Wapl-deficient tumors exhibit fewer

anaphase defects (Supplemental Figure 6D) and less intra-tumor chromosome numerical

heterogeneity (Supplemental Figure 6E). Together these data indicate that suppression of nCIN

in NSCLC cells is not in itself sufficient to limit tumor initiation and growth *in vivo*.

nCIN is a driver of drug resistance *in vivo*

To test the role of nCIN in tumor relapse, the xenograft model described above was used to

establish mice that harbored inducible PC9 shWapl tumors. Once tumors reached a volume of

$300mm^3$, mice were put on a 5 days on, 2 days off (5+2) regimen of 50 mg/kg Gefitinib

treatment and randomly assigned to two cohorts. One cohort received doxycycline in the

drinking water to induce Wapl depletion concurrent with Gefitinib treatment, the other did not

(Figure 5A). Both cohorts of mice exhibited similar initial response to Gefitinib, with all tumors

exhibiting >50% recession within 2 weeks. In the 100 days following initial tumor recession mice

were sustained on a 5+2 drug regimen. During this time 37% (7/19) of Mock-depleted tumors

and 21% (4/19) of Wapl depleted tumors relapsed. Relapse of Wapl depleted tumors was

delayed by nearly 3 weeks compared to those without induced Wapl depletion (66 days vs 46

days for Mock depleted tumors) (Figure 5B, Supplemental Figure 7B). A similar frequency of the

Figure 5: nCIN is a driving force for acquired drug resistance. A) Diagram of experimental set up: PC9 cells carrying an inducible shWapl construct were injected into the flank of the mice. Once tumors reached 300mm³ mice were split into two cohorts: one was administered doxycycline in drinking water to induce Wapl depletion concurrent with Gefitinib via oral gavage, the other received Gefitinib alone. All mice were monitored for initial tumor recession and relapse. B) Time to relapse of tumors with uninduced (n=19) and induced Wapl depletion (n= 19). Red lines indicate tumors that relapsed in this time frame, gray lines indicate those that remained responsive to treatment. Shaded box indicates the minimum relapse free response to Gefitinib in each cohort. C) Visualization and quantification of C → T mutation that results in the T790M mutation in EGFR that confers resistance to TKI in tumors from each cohort. D) qPCR analysis of Wapl levels in control (Mock) and induced Wapl depletion (Wapl KD) that relapsed or not following Gefitinib treatment.

TKI resistance-conferring EGFR T790M mutation was detected in relapsed tumors from both

cohorts (Figure 5C). Interestingly, anaphase defects were similarly prevalent in all relapsed

tumors, regardless of cohort, indicating a selective pressure to maintain CIN during acquisition

of drug resistance (Supplemental Figure 7C). Consistent with this, all residual/non-relapsed

tumors analyzed from the Wapl depleted cohort sustained Wapl depletion while tumors that

relapsed expressed Wapl at levels comparable to the Mock-depleted cohort (Figure 5D). These

data support a model whereby re-establishment of Wapl expression/nCIN, or expansion of

clones that fail to silence Wapl/sustain nCIN, is limiting for acquired drug resistance and tumor

relapse (Figure 6).

Figure 6: Mitotic defects that promote CIN enable generation of drug tolerant aneuploidies that permit continued growth and increased incidence of acquired drug resistance.

Discussion

Clonal mutations that enable resistance to targeted or chemotherapeutic approaches pose a

clinical challenge and remain a major cause of death in many cancer types [54]. Clinically and

experimentally, the degree of intra-tumor genomic heterogeneity, and underlying defects in

mitotic cell division have been functionally linked to tumor evolution, drug resistance, and

metastasis [3, 4, 6, 7, 55, 56]. In the context of NSCLC, identification of driver mutations in

EGFR and the initial clinical success of TKI treatment has been hampered by rapid and

prevalent acquisition of drug resistance [38]. Here, we present mechanistic evidence that nCIN

may arise from mis-regulation of cohesin-sensitive Aurora B kinase activity at centromeres and

deregulation of spindle microtubule dynamics. We propose that subsequent chromosome

amplifications contribute to a high incidence of acquired resistance to therapy. The reduction in

emergence of drug tolerant clones following experimental suppression of nCIN demonstrate that

whole chromosome copy number changes create a favorable environment for continued

proliferation while resistance-conferring mutations are attained. Consistent with this, we find that

a selective pressure exists *in vivo* to maintain or re-establish nCIN to facilitate robust acquisition

of drug resistance and tumor relapse (Figures 5 &6).

Whole chromosome segregation errors enable tumor evolution and drug resistance

Our analysis examined isogenic PC9 cells that primarily differ, at least initially, in their nCIN

status. Our results indicate that nCIN contributes to drug resistance by allowing for the

generation of whole chromosome amplifications that harbor key drug tolerance genes (like

EGFR, MET, and SNAI2/SLUG) and promote continued proliferation in the presence of TKI. In

the absence of an increased mutation rate, this continued proliferation is key to enable the

acquisition of replication-dependent mutations that confer robust drug resistance and tumor

relapse. Such adaptive mutability may be particularly relevant to the mis-segregation and

subsequent selection for amplification of chromosome 7, which contains both MET and EGFR

genes. Increased EGFR gene copy number both promotes proliferation [40] and, by virtue of

having more EGFR gene templates for replication-acquired mutation, increases the apparent

mutability of individual cells [57]. These findings are consistent with previous studies showing

selective pressure for cancer cells to sustain nCIN and demonstrating that high levels of nCIN

can drive the generation and selection of drug-resistant clones [6, 7, 58, 59].

High mutation rates may preclude the need for nCIN in acquired drug resistance.

In the absence of nCIN, resistance is limited to the clonal amplification of cells with pre-existing

amplifications or mutations, and those that acquire chromosome amplifications through rare

segregation errors. An increase in mutation rate may negate the need for nCIN by increasing

the frequency at which resistance-conferring mutations are generated in each cell cycle.

Consistent with this view, high mutation rates and nCIN have been found to be mutually

exclusive in various cancer contexts [60].

nCIN as a therapeutic target

Similar to what we demonstrate here, work from other groups has shown that presence of nCIN

can promote acquisition of drug resistance and is a mechanism to evade oncogene addiction [5,

58]. Our data additionally show that suppression of segregation errors in a cancer context is

achievable and that reduction of nCIN can limit mechanisms of acquired drug resistance.

Indeed, due to its role in regulation of chromosome segregation, Aurora B is a provocative drug

target, and its inhibition has recently been shown to be efficacious in limiting proliferation of TKI-

resistant NSCLC cells [61]. Together these data propose that pathways that promote nCIN may

serve as valuable drug targets, alone, or as co-therapies to enhance or prolong response to

targeted therapeutic approaches.

Acknowledgements

This work was supported by a Smith Family Award for Excellence in Biomedical Research and

R00CA182731 to ALM.

Author Contributions

ALM conceived and devised the study. ALM, EAC, and NMH designed the experiments. ALM,

NMH, EAC, and CPH performed the experiments. ALM wrote the manuscript with input and

approval from all authors.

Declaration of Interests

Authors declare no competing interests

References

Baker DJ, Jin F, Jeganathan KB, van Deursen JM. Whole chromosome instability caused by Bub1 insufficiency drives tumorigenesis through tumor suppressor gene loss of heterozygosity. *Cancer Cell* 2009; 16: 475-486.

Weaver BA, Silk AD, Montagna C, Verdier-Pinard P, Cleveland DW. Aneuploidy acts both oncogenically and as a tumor suppressor. *Cancer Cell* 2007; 11: 25-36.

Choi CM, Seo KW, Jang SJ, Oh YM, Shim TS, Kim WS *et al*. Chromosomal instability is a risk factor for poor prognosis of adenocarcinoma of the lung: Fluorescence in situ hybridization analysis of paraffin-embedded tissue from Korean patients. *Lung Cancer* 2009; 64: 66-70.

McClelland SE, Burrell RA, Swanton C. Chromosomal instability: a composite phenotype that influences sensitivity to chemotherapy. *Cell Cycle* 2009; 8: 3262-3266.

Sotillo R, Schvartzman JM, Socci ND, Benezra R. Mad2-induced chromosome instability leads to lung tumour relapse after oncogene withdrawal. *Nature* 2010; 464: 436-440.

Lukow DA, Sausville EL, Suri P, Chunduri NK, Wieland A, Leu J *et al*. Chromosomal instability accelerates the evolution of resistance to anti-cancer therapies. *Dev Cell* 2021; 56: 2427-2439 e2424.

Ippolito MR, Martis V, Martin S, Tijhuis AE, Hong C, Wardenaar R *et al*. Gene copy-number changes and chromosomal instability induced by aneuploidy confer resistance to chemotherapy. *Dev Cell* 2021; 56: 2440-2454 e2446.

Bakhoum SF, Genovese G, Compton DA. Deviant kinetochore microtubule dynamics underlie chromosomal instability. *Curr Biol* 2009; 19: 1937-1942.

Godek KM, Kabeche L, Compton DA. Regulation of kinetochore-microtubule attachments through homeostatic control during mitosis. *Nat Rev Mol Cell Biol* 2015; 16: 57-64.

Taylor AM, Shih J, Ha G, Gao GF, Zhang X, Berger AC *et al*. Genomic and Functional Approaches to Understanding Cancer Aneuploidy. *Cancer Cell* 2018; 33: 676-689.e673.

Nagy Á, Lánczky A, Menyhárt O, Győrffy B. Validation of miRNA prognostic power in hepatocellular carcinoma using expression data of independent datasets. *Sci Rep* 2018; 8: 9227.

Hindriksen S, Lens SMA, Hadders MA. The Ins and Outs of Aurora B Inner Centromere Localization. *Front Cell Dev Biol* 2017; 5: 112.

González-Loyola A, Fernández-Miranda G, Trakala M, Partida D, Samejima K, Ogawa H *et al*. Aurora B Overexpression Causes Aneuploidy and p21Cip1 Repression during Tumor Development. *Mol Cell Biol* 2015; 35: 3566-3578.

Abe Y, Sako K, Takagaki K, Hirayama Y, Uchida KS, Herman JA *et al*. HP1-Assisted Aurora B Kinase Activity Prevents Chromosome Segregation Errors. *Dev Cell* 2016; 36: 487-497.

Broad AJ, DeLuca KF, DeLuca JG. Aurora B kinase is recruited to multiple discrete kinetochore and centromere regions in human cells. *J Cell Biol* 2020; 219.

Hauf S, Cole RW, LaTerra S, Zimmer C, Schnapp G, Walter R *et al*. The small molecule Hesperadin reveals a role for Aurora B in correcting kinetochore-microtubule attachment and in maintaining the spindle assembly checkpoint. *J Cell Biol* 2003; 161: 281-294.

Muñoz-Barrera M, Monje-Casas F. Increased Aurora B activity causes continuous disruption of kinetochore-microtubule attachments and spindle instability. *Proc Natl Acad Sci U S A* 2014; 111: E3996-4005.

Huang H, Lampson M, Efimov A, Yen TJ. Chromosome instability in tumor cells due to defects in Aurora B mediated error correction at kinetochores. *Cell Cycle* 2018; 17: 2622-2636.

Kleyman M, Kabeche L, Compton DA. STAG2 promotes error correction in mitosis by regulating kinetochore-microtubule attachments. *J Cell Sci* 2014; 127: 4225-4233.

Herlihy CP, Hahn S, Hermance NM, Crowley EA, Manning AL. Suv420 enrichment at the centromere limits Aurora B localization and function. *J Cell Sci* 2021; 134.

Manning AL, Benes C, Dyson NJ. Whole chromosome instability resulting from the synergistic effects of pRB and p53 inactivation. *Oncogene* 2014; 33: 2487-2494.

Swanton C, Nicke B, Schuett M, Eklund AC, Ng C, Li Q *et al*. Chromosomal instability determines taxane response. *Proc Natl Acad Sci U S A* 2009; 106: 8671-8676.

Yu J, Zhou J, Xu F, Bai W, Zhang W. High expression of Aurora-B is correlated with poor prognosis and drug resistance in non-small cell lung cancer. *Int J Biol Markers* 2018; 33: 215-221.

Vischioni B, Oudejans JJ, Vos W, Rodriguez JA, Giaccone G. Frequent overexpression of aurora B kinase, a novel drug target, in non-small cell lung carcinoma patients. *Mol Cancer Ther* 2006; 5: 2905-2913.

Takeshita M, Koga T, Takayama K, Ijichi K, Yano T, Maehara Y *et al*. Aurora-B overexpression is correlated with aneuploidy and poor prognosis in non-small cell lung cancer. *Lung Cancer* 2013; 80: 85-90.

Lampson MA, Renduchitala K, Khodjakov A, Kapoor TM. Correcting improper chromosome-spindle attachments during cell division. *Nat Cell Biol* 2004; 6: 232-237.

Liang C, Zhang Z, Chen Q, Yan H, Zhang M, Zhou L *et al*. Centromere-localized Aurora B kinase is required for the fidelity of chromosome segregation. *J Cell Biol* 2020; 219.

Gregan J, Polakova S, Zhang L, Tolic-Norrelykke IM, Cimini D. Merotelic kinetochore attachment: causes and effects. *Trends Cell Biol* 2011; 21: 374-381.

Welburn JP, Vleugel M, Liu D, Yates JR, 3rd, Lampson MA, Fukagawa T *et al*. Aurora B phosphorylates spatially distinct targets to differentially regulate the kinetochore-microtubule interface. *Mol Cell* 2010; 38: 383-392.

Cimini D, Wan X, Hirel CB, Salmon ED. Aurora kinase promotes turnover of kinetochore microtubules to reduce chromosome segregation errors. *Curr Biol* 2006; 16: 1711-1718.

Meppelink A, Kabeche L, Vromans MJ, Compton DA, Lens SM. Shugoshin-1 balances Aurora B kinase activity via PP2A to promote chromosome bi-orientation. *Cell Rep* 2015; 11: 508-515.

Zhou L, Liang C, Chen Q, Zhang Z, Zhang B, Yan H *et al*. The N-Terminal Non-Kinase-Domain-Mediated Binding of Haspin to Pds5B Protects Centromeric Cohesion in Mitosis. *Curr Biol* 2017; 27: 992-1004.

Haarhuis JH, Elbatsh AM, van den Broek B, Camps D, Erkan H, Jalink K *et al*. WAPL-mediated removal of cohesin protects against segregation errors and aneuploidy. *Curr Biol* 2013; 23: 2071-2077.

Sherwood R, Takahashi TS, Jallepalli PV. Sister acts: coordinating DNA replication and cohesion establishment. *Genes Dev* 2010; 24: 2723-2731.

Kueng S, Hegemann B, Peters BH, Lipp JJ, Schleiffer A, Mechtler K *et al*. Wapl controls the dynamic association of cohesin with chromatin. *Cell* 2006; 127: 955-967.

Liu D, Vader G, Vromans MJ, Lampson MA, Lens SM. Sensing chromosome bi-orientation by spatial separation of aurora B kinase from kinetochore substrates. *Science* 2009; 323: 1350-1353.

Cimini D. Detection and correction of merotelic kinetochore orientation by Aurora B and its partners. *Cell Cycle* 2007; 6: 1558-1564.

Gazdar AF. Activating and resistance mutations of EGFR in non-small-cell lung cancer: role in clinical response to EGFR tyrosine kinase inhibitors. *Oncogene* 2009; 28 Suppl 1: S24-31.

Gerber DE. EGFR Inhibition in the Treatment of Non-Small Cell Lung Cancer. *Drug Dev Res* 2008; 69: 359-372.

Stewart EL, Tan SZ, Liu G, Tsao MS. Known and putative mechanisms of resistance to EGFR targeted therapies in NSCLC patients with EGFR mutations-a review. *Transl Lung Cancer Res* 2015; 4: 67-81.

Arao T, Fukumoto H, Takeda M, Tamura T, Saijo N, Nishio K. Small in-frame deletion in the epidermal growth factor receptor as a target for ZD6474. *Cancer Res* 2004; 64: 9101-9104.

Vargas-Rondón N, Villegas VE, Rondón-Lagos M. The Role of Chromosomal Instability in Cancer and Therapeutic Responses. *Cancers (Basel)* 2017; 10.

Bakhoum SF, Compton DA. Chromosomal instability and cancer: a complex relationship with therapeutic potential. *J Clin Invest* 2012; 122: 1138-1143.

Chunduri NK, Storchová Z. The diverse consequences of aneuploidy. *Nat Cell Biol* 2019; 21: 54-62.

Zhou C, Zhu Y, Lu B, Zhao W, Zhao X. Survivin expression modulates the sensitivity of A549 lung cancer cells resistance to vincristine. *Oncol Lett* 2018; 16: 5466-5472.

Kavallaris M, Kuo DY, Burkhart CA, Regl DL, Norris MD, Haber M *et al*. Taxol-resistant epithelial ovarian tumors are associated with altered expression of specific beta-tubulin isotypes. *J Clin Invest* 1997; 100: 1282-1293.

Engelman JA, Zejnullahu K, Mitsudomi T, Song Y, Hyland C, Park JO *et al*. MET amplification leads to gefitinib resistance in lung cancer by activating ERBB3 signaling. *Science* 2007; 316: 1039-1043.

McDermott U, Pusapati RV, Christensen JG, Gray NS, Settleman J. Acquired resistance of non-small cell lung cancer cells to MET kinase inhibition is mediated by a switch to epidermal growth factor receptor dependency. *Cancer Res* 2010; 70: 1625-1634.

Nurwidya F, Takahashi F, Murakami A, Kobayashi I, Kato M, Shukuya T *et al*. Acquired resistance of non-small cell lung cancer to epidermal growth factor receptor tyrosine kinase inhibitors. *Respir Investig* 2014; 52: 82-91.

Lee AF, Chen MC, Chen CJ, Yang CJ, Huang MS, Liu YP. Reverse epithelial-mesenchymal transition contributes to the regain of drug sensitivity in tyrosine kinase inhibitor-resistant non-small cell lung cancer cells. *PLoS One* 2017; 12: e0180383.

Tulchinsky E, Demidov O, Kriajevska M, Barlev NA, Imyanitov E. EMT: A mechanism for escape from EGFR-targeted therapy in lung cancer. *Biochim Biophys Acta Rev Cancer* 2019; 1871: 29-39.

Yochum ZA, Cades J, Wang H, Chatterjee S, Simons BW, O'Brien JP *et al*. Targeting the EMT transcription factor TWIST1 overcomes resistance to EGFR inhibitors in EGFR-mutant non-small-cell lung cancer. *Oncogene* 2019; 38: 656-670.

Zhu X, Chen L, Liu L, Niu X. EMT-Mediated Acquired EGFR-TKI Resistance in NSCLC: Mechanisms and Strategies. *Front Oncol* 2019; 9: 1044.

Redmond KL, Papafili A, Lawler M, Van Schaeybroeck S. Overcoming Resistance to Targeted Therapies in Cancer. *Semin Oncol* 2015; 42: 896-908.

Rajagopalan H, Lengauer C. Aneuploidy and cancer. *Nature* 2004; 432: 338-341.

Gao C, Furge K, Koeman J, Dykema K, Su Y, Cutler ML *et al*. Chromosome instability, chromosome transcriptome, and clonal evolution of tumor cell populations. *Proc Natl Acad Sci U S A* 2007; 104: 8995-9000.

Andersson DI, Slechta ES, Roth JR. Evidence that gene amplification underlies adaptive mutability of the bacterial lac operon. *Science* 1998; 282: 1133-1135.

Salgueiro L, Buccitelli C, Rowald K, Somogyi K, Kandala S, Korbel JO *et al*. Acquisition of chromosome instability is a mechanism to evade oncogene addiction. *EMBO Mol Med* 2020; 12: e10941.

Orr B, Talje L, Liu Z, Kwok BH, Compton DA. Adaptive Resistance to an Inhibitor of Chromosomal Instability in Human Cancer Cells. *Cell Rep* 2016; 17: 1755-1763.

Lengauer C, Kinzler KW, Vogelstein B. Genetic instabilities in human cancers. *Nature* 1998; 396: 643-649.

Bertran-Alamillo J, Cattan V, Schoumacher M, Codony-Servat J, Giménez-Capitán A, Cantero F *et al*. AURKB as a target in non-small cell lung cancer with acquired resistance to anti-EGFR therapy. *Nat Commun* 2019; 10: 1812.