Abnormal craniofacial and spinal bone development with *col2a1*a depletion in a zebrafish
model of CHARGE syndrome

Maximilian Breuer[1#; *]; Maximilian Rummler[2#]; Charlotte Zaouter[1]; Bettina M. Willie[2##];
Shunmoogum A. Patten[1,3##; *]

[#], [##] authors contributed equally

Affiliations:

1. INRS – Centre Armand Frappier Santé Biotechnologie, 531 Boulevard des Prairies, Laval,
QC, Canada, H7V 1B7
2. Research Centre, Shriners Hospital for Children-Canada, Department of Pediatric Surgery,
McGill University, 1003 Decarie Blvd, Montreal, Canada H4A 0A9
3. Centre d'Excellence en Recherche sur les Maladies Orphelines - Fondation Courtois
(CERMO-FC), Université du Québec à Montréal (UQAM), Montréal, QC, Canada

*Correspondence: Correspondence should be addressed to
Dr. Maximilian Breuer
INRS- Centre Armand-Frappier Santé et Biotechnologie
531 Boulevard des Prairies
Laval, Quebec
H7V 1B7
Canada
Maximilian.Breuer@iaf.inrs.ca
+1 (438) 341-0445

23
24
25
26

32 **Conflict of Interest**

33 The authors declare no conflicts of interest

Abstract

CHARGE syndrome patients commonly display craniofacial abnormalities. Furthermore, most patients show features of idiopathic scoliosis, reduced bone mineral density and in a few cases osteopenia. While several clinical cases and studies have documented the skeletal deformities in CHARGE syndrome bearing *CHD7* mutations, the underlying mechanisms of the disorder remain elusive. Here, we detect and quantitatively analyze skeletal abnormalities in larval and adult *chd7*$^{-/-}$ zebrafish.

We show that young *chd7*$^{-/-}$ larvae present with abnormal craniofacial development, especially related to cartilage. We also observe scoliosis-like spinal deformations at 9 dpf. Gene expression analysis confirmed the reduction of osteoblast markers and Pparγ targets. MicroCT analyses identified abnormal craniofacial structures, Weberian apparatus and vertebral body morphology in *chd7*$^{-/-}$ mutants, with highly mineralized inclusions, along with significant variances in bone mineral density and bone volume. Notably, we detect a specific depletion of Col2a1a in the cartilage of craniofacial regions and vertebrae, in line with a significantly reduced number of chondrocytes.

Our study is the first to elucidate the mechanisms underlying morphological changes in craniofacial structure and vertebrae of adult *chd7*$^{-/-}$ zebrafish. The *chd7*$^{-/-}$ mutant zebrafish will be beneficial in future investigations of the underlying pathways of both craniofacial and spinal deformities commonly seen in CHARGE syndrome.

Keywords: CHARGE Syndrome, Chd7, idiopathic scoliosis, zebrafish, Col2a1

Introduction

Impaired bone development, craniofacial dysmorphism and spinal abnormalities are a major health concern in various genetic disorders. Among these, idiopathic scoliosis is the most common skeletal abnormality observed in children (1). Idiopathic scoliosis is a curvature of the spine with unidentified genetic cause. However, many risk genes have been connected to the underlying mechanisms (2). One of these related genetic disorders is the congenital multisystemic CHARGE syndrome (CS) named after the major characteristics of **C**oloboma, **H**eart defects, **A**tresea chonae, **R**etarded growth, **G**enital and **E**ar abnormalities (3-5). Though, considering the wide variety of observed phenotypes, the list of characteristics has been continuously augmented to include craniofacial abnormalities and spinal deformations (6-8). Along these lines, CS is closely associated with skeletal deformities such as craniofacial dysmorphisms, idiopathic scoliosis, kyphosis and hemivertebrae (9). In fact, idiopathic scoliosis is observed in a majority of CS cases, with studies showing over 60% of patients having diagnosed scoliosis at an average age of just over 6 years (10-12). In some cases, the areal bone mineral density (aBMD) is reduced (13).

CS is most commonly caused by a mutation in the chromodomain ATP-dependant helicase 7 (CHD7). Mutations are distributed evenly throughout the gene and the vast majority of cases are sporadic, with only very few cases of familial CS (7, 14-16). Analysis of pathways related to CHD7 show involvement in neural crest differentiation/proliferation/migration and stem cell quiescence (17-19). Furthermore, studies focusing on CS have identified regulatory mechanisms of CHD7 in the immune response and more strikingly in brain development (20, 21). Most recently, CHD7 has been connected to abnormal GABAergic development resulting in an autistic-like behavior (Jamadagni et al., unpublished data). While the function of CHD7 has been extensively studied, little attention has been given to the underlying mechanisms in skeletal development (22). Some studies have directly linked CHD7 to osteogenesis (23, 24).

Investigation in cell cultures have shown a dependency on CHD7 for successful differentiation,

more specifically to a differentiation complex with SETD1B, NLK and SMAD1 resulting in

depleted osteogenesis upon depletion of CHD7 in favour of adipogenesis by regulating PPARγ

target genes (23, 24). Various models to investigate this role in vivo have been proposed. A

mouse model for Chd7 deficiency, termed "looper" presents with ear ossicle malformations,

but no spinal deformities (25). However, mouse models for Chd7 deficiency have limitations,

as null mutants are embryonically lethal by 10 days (26). Yet, a recent zebrafish model has

proven valuable in the modelling of *chd7* dependent CS revealing a reduction in vertebrae

mineralization of young larvae (27).

Zebrafish have become increasingly relevant in the study of fundamental bone development

and bone related disorders (28-30), including scoliosis, osteoporosis, age related osteoarthritis

(31, 32). Notably, the spinal structure in zebrafish has been characterized in detail and

comprises three major regions: the Weberian apparatus, a specialized structure consisting of the

first four vertebrae connecting the auditory system and the swim bladder to amplify sound

vibrations, the precaudal vertebrae which are connected to neural arch and spines and caudal

vertebrae of the tail region with neural and hemal arch (33). Simplicity of analysis in zebrafish

to investigate spinal structures has been used to understand the effects of mechanical loading

on bone mineralization (30, 34). Zebrafish have specific advantages in the analysis of skeletal

development such as the closely related structure to humans. They also have rapid bone

development with first mineralization of vertebrate occurring after only 5 days post fertilization

(dpf). Additionally, the *ex utero* development and transparency of young larvae simplifies the

use of *in vivo* staining and transgenic lines to allow for the analysis of early calcification in

zebrafish bony structures (35-38). Furthermore, the simplicity of high-throughput drug

screening in this teleost model makes the model highly useful for investigating new drug targets

and effect on skeletal development (39, 40). These advantages along with their high

remodelling efficiency of skeletal structures including their possibility to regenerate fins, as well as the effectiveness to reproduce spinal deformity phenotypes have raised interest in the zebrafish model (41, 42).Studies in zebrafish have shown a close link to collagen related genes (43, 44). These highly conserved pathways between zebrafish and humans allow us to investigate the underlying pathomechanisms linking CHD7 to idiopathic scoliosis and other skeletal abnormalities.

In the present study we describe for the first time a detailed analysis of craniofacial and skeletal anomalies in a zebrafish $chd7^{-/-}$ mutant model for CS. Early larvae screens reveal a striking dysmorphism of craniofacial structures, as well as delay of mineralization of vertebrae bodies linked to deficient osteoblast differentiation. Gene expression and IHC analysis reveals striking reduction in osteoblast, chondrocyte, and collagen matrix markers, particularly displaying abolished levels of Col2a1. Finally, extensive microCT analysis in adults show abnormal mineralization and morphology in all major structures of the zebrafish skeleton.

Results

Zebrafish $chd7^{-/-}$ larvae show craniofacial and spinal deformities

We recently generate a chd7 knockout zebrafish line using CRISPR/Cas9 with a single nucleotide insertion causing a frame-shifting mutation and a premature stop codon 8 amino acids after the mutation site (Jamadagni et al., unpublished data). Consistent with our previous findings using a transient chd7 morpholino knockdown model (27), this stable $chd7^{-/-}$ zebrafish mutant using CRISPR/Cas9 that replicates hallmarks of CS including craniofacial and skeletal defects (Fig. 1). To further investigate the bone deformities in $chd7^{-/-}$ zebrafish in detail, we first screened young $chd7^{-/-}$ larvae at 6 dpf and 9 dpf for morphological abnormalities at early developmental stages. At 6 dpf we observed craniofacial changes, such as reduced length of the palatoquadrate and increased angle of the ceratohyal (Fig 1A). Calcein staining also reveals reduced mineralization of the facial structures, noticeably towards the quadrate and opercle (Fig

1B). At this stage $chd7^{-/-}$ larvae were minimally, yet significantly, smaller than control (Cntrl)

larvae, however this is in line with the expected CS phenotype of growth retardation and neural

crest abnormalities in patients (Supplemental Fig. 1A).

39% of $chd7^{-/-}$ larvae exhibited highly varying, scoliosis-like phenotypes at both precaudal and

caudal regions of the spine (Fig. 1C). Additionally, some of these also presented with a

kyphosis-like phenotype (Supplemental Fig. 1B). We found that $chd7^{-/-}$ larvae were still

reactive to a touch response, even with the severe morphological phenotype.

$chd7^{-/-}$ larvae have a spinal mineralization deficit that is nutrient dependent

Given the spinal deformities in 9 dpf $chd7^{-/-}$ larvae, we next sought to investigate mineralization

during larval development. Calcein staining for fluorochrome labeling of bones revealed no

delay in the onset of mineralization of the first precaudal vertebrae at 6 dpf. Upon further

development, at 9 dpf, $chd7^{-/-}$ larvae showed a significant reduction in the number of calcified

vertebrae (Fig. 1D, E). Notably, $chd7^{-/-}$ larvae were less efficient in complete calcification of

vertebrae towards the posterior region of the spine (Fig. 1D).

Since calcification of the bone matrix is dependent on mineral uptake from food, we tested how

the onset and development of vertebral bone structure calcification in 9 dpf $chd7^{-/-}$ zebrafish

was dependent on nutrition accessibility. Notably, wildtype and $chd7^{-/-}$ larvae showed no

difference in the number of mineralized vertebrae in a starvation situation between 5 dpf and 9

dpf. Expectedly, we observed significant differences when food was available from 5 dpf until

9 dpf, with $chd7^{-/-}$ zebrafish having a reduced number of mineralized vertebrae compared to

controls. The $chd7^{-/-}$ larvae had the same number of mineralized vertebrae when food was

available or absent (Fig. 1 D, E). Starvation in zebrafish larvae between 5 dpf and 9 dpf also

resulted in smaller yolks, which is to be expected, considering the high dependency on nutrition

supply by the yolk. Surprisingly, we found that Weberian vertebrae, which are the first 4

vertebrae of the spinal column, were more rapidly calcified in *chd7*$^{-/-}$ larvae than in controls

(Fig. 1D arrows, Fig 1 F).

We investigated if mineralization recovered in later development. Reduced mineralization

remained evident in 4-week-old juvenile zebrafish with spinal deformities. Alizarin red staining

revealed decreased mineralization of caudal vertebrae, with striking deficiency toward the

vertebral body (Supplemental Fig. 1. D).

Osteoblast differentiation is impaired in chd7$^{-/-}$ *larvae*

CHD7 has been shown to regulate osteogenesis by controlling PPARγ to promote skeletal

development and regulate expression of promoting genes. To examine if the Pparγ pathway is

directly involved in the observed bone mineralization deficiency and abnormal morphology in

chd7$^{-/-}$ larvae, we tested for Pparγ target gene expression and osteoblast differentiation marker

genes at the onset of the bone defect at 9 dpf (Fig. 1G). The key targets for Pparγ, *runx2a* and

runx2b, which are expressed mostly in craniofacial regions at this stage, were significantly

downregulated in *chd7*$^{-/-}$ larvae compared to controls. Additionally, other markers for

osteogenesis were also affected, including a significant downregulation of osteoblast markers

sp7/osterix, bglap and *postna* (Fig. 1G). The *chd7*$^{-/-}$ larvae also had significantly lower levels

of osteocyte markers while other genes regulating mineralization such as *acp5a* and *sost*, while

mgp remained unchanged compared to control larvae (Fig 1H). We also detected a significant

upregulation of the osteoclast marker *ctsk* in *chd7*$^{-/-}$ larvae.

Abnormal development of craniofacial regions and the Weberian apparatus in adult chd7$^{-/-}$

zebrafish

To assess the effects of *chd7* deficiency on bone development, including craniofacial and

vertebral growth and maturation, we analyzed bone morphology and mass using microCT in

adult zebrafish. Morphologically, adult *chd7*$^{-/-}$ zebrafish were unchanged in body length.

Since mineralization and morphology in the young larvae varied in craniofacial regions, as well

as from anterior to posterior regions of the spine, we decided to analyze the four major bony

structures of the zebrafish: the skull, Weberian apparatus, precaudal and caudal vertebrae. The

skull was notably changed in $chd7^{-/-}$ zebrafish in comparison to controls (Fig. 2A-D). In

particular, we observed significant alterations in the mandibular angle and length (Fig. 2E, F).

The face of $chd7^{-/-}$ zebrafish was also notably deformed compared to control zebrafish, with a

significantly increased craniofacial angle (Fig. 2G). Further, the angle of the mandibular arch

was wider in $chd7^{-/-}$ zebrafish. Lastly, 3 out of 5 $chd7^{-/-}$ zebrafish presented with a warped skull,

with the tip of the dentary deviating from the fish's midline.

Another key structure, the Weberian apparatus with its supraneurals, intercalarium, tripus and

parapophysis, had thinner and smaller morphology in $chd7^{-/-}$ adults compared to controls (Fig.

2I). We further performed a detailed analysis of volumetric bone mineral density (vBMD), bone

volume (BV), total volume (TV), and bone volume fraction (BV/TV) of 1-year-old $chd7^{-/-}$ and

control zebrafish. We did not detect any significant differences in vBMD, BV or TV

(Supplemental. Fig. 2). However, we found a significantly greater variance in TV of both

intercalarium and parapophysis, as well as in BV of the parapophysis in 1-year-old $chd7^{-/-}$

zebrafish, as determined by F-test analysis (Supplemental Table 1). The same results were

observed in 2-year-old $chd7^{-/-}$ compared to control zebrafish (Supplemental Table 2).

Abnormal mineralization and malformations in the body end plates of precaudal vertebrae in

adult zebrafish

Vertebrae structures were analyzed by assessing key angles of the structure (Fig. 3A) and

mineralization features. MicroCT analysis of precaudal vertebrae identified abnormal patterns

of mineralization and abnormal structures, that were especially pronounced at the arches and

transverse processes in 1-year old $chd7^{-/-}$ and control zebrafish (Fig. 3B, C). Precaudal vertebrae

showed significantly higher overall vBMD, while the distribution of mineralization between

the body and arch remained unaffected (Fig. 3D, E). However, structurally, precaudal vertebrae of $chd7^{-/-}$ mutants show an increase in bone volume towards the body end plates of vertebrae, which in most cases include major malformations manifesting as highly mineralized inclusions (Fig. 3C, arrows). Additionally, vertebrae in 1-year-old $chd7^{-/-}$ zebrafish presented with a larger bone volume fraction (BV/TV) compared to controls (Fig. 3G-I). Analysis of the vertebrae neural arch revealed distortion with a significantly larger rising angle, as well as a larger variance of the angles (Fig. 3F). Vertebrae showed a significantly changed body angle and significant variance of body angle and vertebrae length (Supplemental Table 3). While a depletion of vBMD in the arches and abnormal vertebrae body structure was observed in 2-year-old fish, the effects seen on total vBMD and BV were not apparent in the few samples that survived until this stage (Supplemental Fig. 2 and Supplemental Table 4). Notably, one of our $chd7^{-/-}$ mutants showed vertebral fusion of the analyzed precaudal vertebra, which was not observed in any of the controls (Supplemental Fig. 1). This further demonstrates the wide variation in severity of the bone phenotype in $chd7^{-/-}$ zebrafish.

Abnormal caudal vertebrae in chd7$^{-/-}$ zebrafish vary in arch angles and reduced mineralization

We next determined key structural features (Fig. 4A) and mineralization characteristics of caudal vertebrae. The $chd7^{-/-}$ zebrafish presented with warped hemal and neural arches and abnormal body structure compared to controls (Fig. 4B, C). Similar to precaudal vertebrae, the growth zones of the body end plates were enlarged and showed inclusions of highly mineralized matrix (Fig. 4C, arrows). Unlike control zebrafish, $chd7^{-/-}$ mutants had abnormal vertebrae body structure, which usually exhibited an hourglass shape and resulting in greater variance of measured body angles (Supplemental Table 3).

In line with the neural arch screened in the precaudal vertebrae, the hemal arch in the caudal vertebrae showed significantly wider arch rising angles (Fig. 4F), while the neural arch displayed significantly greater variance (Fig. 4G) in 1-year-old $chd7^{-/-}$ zebrafish. In contrast to

precaudal vertebrae data, the caudal vertebrae showed significantly larger variance of Total

vBMD and of both arches and vertebral body vBMD (Supplemental Table 3). A significant

insufficiency in mineralization of the arches compared to the vertebrae body was also measured

in $chd7^{-/-}$ zebrafish compared to controls (Fig. 4D, E). Further analysis also revealed a

significant variance in BV, TV and BV/TV in $chd7^{-/-}$ zebrafish compared to controls (Fig. 4H,

I).

While significantly distorted arches were still observed in 2-year old mutants, the effects on

vBMD and BV were not noticed in the few surviving fish that could be screened (Supplemental

Fig. 3 and Supplemental Table 4).

Col2a1 of the ECM is depleted in chd7$^{-/-}$ zebrafish

To perform a more complete analysis of the skeletal phenotype, we also examined the

extracellular matrix (ECM) of bone structures in 6 dpf larvae and 1-year-old $chd7^{-/-}$ zebrafish.

Gene expression analysis at 9 dpf revealed a significant downregulation of the ECM gene

col2a1a (Fig. 5E). Antibody staining showed that all craniofacial structures are already depleted

of the key Col2a1a protein by 5 dpf (Fig. 5A). Vacuole formation within the structures was

apparently unimpaired. In adult $chd7^{-/-}$ zebrafish, immunofluorescence showed a striking

depletion of Col2a1 in the cartilage of vertebrae seen in the Weberian and precaudal vertebrae,

with almost abolished Col2a1 signal (Fig 5B). This pattern was effectively reproduced in all

regions of the spine of $chd7^{-/-}$ zebrafish.

We tested further for cell nuclei and chondrocytes in vertebral cartilage. Hematoxylin-Eosin

(HE) staining of Weberian apparatus and precaudal vertebrae sections showed a significant

reduction in the number of nuclei measured in the vertebral cartilage of 1-year-old $chd7^{-/-}$

zebrafish and controls (Fig. 5C, F). Complementary analysis using Safranin/Fast Green O

staining, revealed a significant reduction in the number of chondrocytes in the vertebral

cartilage in *chd7*[-/-] zebrafish in line with the HE staining (Fig. 5D, G). Altogether, the findings

indicate an altered cartilage composition in addition to the observed bone deformations.

Discussion

Efficiency of model

CS is a multisystemic disorder likely involving a variety of underlying mechanisms that include

CHD7. Therefore, different preclinical models are required to individually identify the various

molecular mechanisms. Even though mouse models for CS exist, none have been reported to

show the bone phenotype observed in CS patients. Our study presents a *chd7* deficient zebrafish

that has a bone phenotype resembling that of CS patients, which we used to investigate skeletal

development in CS. CS patients commonly present with square face, broad nasal bridge, small

mouth and facial asymmetry (9). While scoliosis in CS patients has been attributed to poor

muscular development, our study proposes an additional factor involving impaired early

mineralization of bone in the vertebrae. In line with clinical features exhibited by CS patients,

our mutants display craniofacial dysmorphism with reduced mineralization and spinal

deformities indicative of scoliosis (10-12). Interestingly, our analysis revealed that craniofacial

and vertebral abnormalities are already present in young larvae of *chd7*[-/-] zebrafish. Additional

clinical studies will be required to investigate the onset of skeletal deformities in CS patients.

However, early analysis of bone density and spinal development has been suggested by CS

guidelines (45, 46). This early and detailed analysis may reveal less severe or early onset

phenotypes. Similarly, we observed that adult *chd7*[-/-] zebrafish, which appeared without spinal

deformities on visual inspection, showed signs of spinal abnormalities in microCT analysis,

suggesting that mostly mild phenotypes reach this age. This is supported by the high lethality

rate shown by Jamadagni et al. (unpublished), yet the underlying cause of lethality is still

unclear. Even though 5 dpf old larvae were significantly smaller than their control counterparts,

this is expected as part of the CHARGE phenotype. Nonetheless, overall morphology suggests

that they are not in a notable developmental delay. Unfortunately, key "landmarks" for

development as previously proposed are expectedly affected in *chd7*[-/-] zebrafish, including

overall growth, pigment migration, swimbladder inflation and mineralization (47). However,

we can conclude if the mineralization effect observed at 9dpf was in fact developmental delay,

we would expect mineralization to recover. Yet, the phenotype was still evident in later

development (Supplemental Fig 1. D).

Progressive skeletal disorders such as osteopenia and osteoporosis are characterized by a

reduction in bone mineralization over time leading to increased fracture risk (48). CS patients

have been shown to present with significantly reduced bone mineral density (13).

Correspondingly, reduced vertebrae mineralization in *chd7*[-/-] zebrafish by 12 dpf was observed

in a previous study involving morpholinos in zebrafish which had observed a reduced

mineralization of vertebral structures (27). Confirming previous data, we observed insufficient

mineralization of vertebra along the posterior spinal column by 4 weeks of age. In earlier

analysis of larvae at 6 dpf, we were not able to detect a delay in the onset of mineralization, but

already by 9 dpf we saw a significant decrease in the number of calcified vertebrae.

Furthermore, mineralization appeared to be less efficient in posterior vertebrae, which could

signify an impairment in ossification, since vertebrae in zebrafish undergo ossification from an

anterior to posterior direction (49). An exception from this rule is the Weberian vertebrae C1

and C2, which are ossified later in wildtypes but earlier in *chd7*[-/-] mutants. The craniofacial

abnormalities we observed are in line with known neural crest deficiencies in CS which are

known to affect cartilage and bone development (50).

Nutrient dependency

Our study showed that the onset of calcification was not delayed in *chd7* deficient zebrafish.

However, progressing calcification was decreased already 4 days post feeding onset, even

though access to food was identical in control and mutants. In contrast, differences in

calcification of vertebrae in control and mutants without access to food was not significant.

Previous studies have shown that feeding delay until 9 dpf does not affect future fish growth

and viability and was therefore our cut-off point(51). This data suggests that possibly an

underlying mechanism involving either reduced uptake or metabolizing nutrients may play a

part in the observed phenotype and this link remains to be investigated. Bone calcification and

mineralization is dependent on proper nutrition and is often discussed regarding optimal bone

health in humans (52). Insufficient levels of calcium and nutritional uptake are considered an

increased risk for bone injury in CS patients (11). To optimize nutritional uptake, the

supplementation of calcium and other nutrients to counteract poor bone health has been

proposed for CS patients (53). In fact, case reports show supplementation with calcium and

vitamin D significantly increased aBMD in CS patients (54). A possible nutrient deficiency in

CS, could be linked to gastrointestinal nutrient uptake difficulties in *chd7$^{-/-}$* zebrafish (55), yet

this needs to be further investigated. Thus, this possible deficiency in nutrient uptake could

further hinder ossification in *chd7$^{-/-}$* mutants. We propose using our zebrafish model of CS

model to test the effect of guidelines such as calcium and nutrient supplementation, nutrient

metabolism and mechanical loading in form of swim training to enhance bone strength in CS.

Loss of chd7 reduces osteogenesis in zebrafish

Osteoblast precursors commonly express differentiation markers such as *runx2* and *Osterix/sp7*

(56). Upon maturation of these osteoblasts, expression of mineralization related genes such as

bglap (osteocalcin) are expressed. Analysis of these three genes showed that they were all

significantly downregulated in *chd7$^{-/-}$* larvae. This suggests a deficiency in mature osteoblasts,

required for the mineralization of facial bone structures and early spinal column. Misregulation

or loss of these genes, such as *osterix/sp7* has been directly linked to poor bone mineralization

in zebrafish and humans (57-59). Notably, among all osteogenesis related gene expressions

significantly reduced in *chd7$^{-/-}$* larvae, *sost*, *ctsk, acp5* and *col2a1* have been identified as direct

regulatory targets of Chd7 by Chip-Seq (60). Furthermore, the regulation of osteogenesis has

been linked to a regulatory complex with PPARγ (23). Cell culture experiments have proposed

the role of CHD7 in osteogenesis by inhibiting PPARγ target genes by a protein complex

involving CHD7, NLK and SETDB1. Upon loss of CHD7 this complex will fail to inhibit

PPARγ target genes and therefore inhibit osteogenesis in preference over adipogenesis (23).

Strikingly, our qPCR results are consistent with downregulation of Ppar γ target genes such as

runx2a and *runx2b*. Furthermore, gene expression relevant for bone calcification and

remodelling are misregulated such as *sost* and *ctsk*. Thus, our study is the first to indicate the

conservation of the proposed pathway shown in cell culture in an *in vivo* model.

Weberian apparatus

The Weberian apparatus is a specialized structure that connects the auditory system and the

swim bladder in a complex with the first 4 vertebrae to amplify sound vibrations (33). Our

results show that key structures in the apparatus fail to efficiently form and/or mineralize. Two

of these structures, the tripus and the parapophysis, are formed to connect to the swim bladder.

Both are malformed in all tested *chd7$^{-/-}$* mutants to varying degrees. This malformation may

imply an inefficiency in its function concerning both the swim bladder and auditory system.

Ear abnormalities are a major symptom in CS patients and functionality is affected.

Furthermore, the *chd7$^{-/-}$* mutant shows swim bladder defects (Breuer et al., unpublished) that

could be additionally affecting the morphology in tripus and parapophysis or vice versa. Even

though the Weberian apparatus is extremely specialized for carp and carp-like fish

(Ostariophysi), it is likely that it underlies the same pathways for osteogenesis as other parts of

the spine.

Abnormal vertebrae and vBMD variability

We examined volumetric bone mineral density (vBMD) and other parameters (e.g. BV, TV) in

vertebral bone structures. Most strikingly, we detected large variance in density and all

parameters examined, as well as some significant changes in both precaudal and caudal

vertebrae. The large variance observed in *chd7*[-/-] mutants is in line with CS patients showing

varying severity in spinal deformities (12, 61). Furthermore, previous studies have shown that

even minor changes in the described parameters can impact the risk for spinal deformities and

fractures in disease and under treatment (62, 63).

Another interesting feature we observed in most *chd7*[-/-] mutants, were highly mineralized

inclusions, most commonly, localized towards the growth zone of the vertebrae. We did not

identify the source of these highly mineralized structures, but they could be caused by localized

clusters of osteoblasts during vertebrae mineralization. Another possibility is the occurrence of

calcified cartilage, which is more mineralized in comparison to bone. However, this is highly

speculative and histologic analysis of such small and highly localized structures will be

challenging. Since these inclusions are highly mineralized, it seems likely that their appearance

impairs the bone structurally as higher mineralized collagen is stiffer, yet less tough (64). It will

be of interest to see if future studies can take a closer look at these previously unidentified

features. While these minor defects could be involved in the structural integrity of the spine in

CS patients, these minor features may be missed during dual-energy x-ray absorptiometry

(DEXA) screens of CS patients when assessing aBMD.

Analysis revealed a decreased mineralization in the caudal vertebral arches of the mutant fish.

This likely reduces the mechanical competence of these vertebrae in comparison to healthy

vertebrae. A compromised mechanical competence alongside low BMD also increases fracture

risk and can be an underlying cause of scoliosis (65-67). Unfortunately, natural history BMD

data of CS patients throughout early development is not available, to the best of our knowledge.

However, aBMD analysis of patients with idiopathic scoliosis in CS via DEXA scans has been

studied and shows significant reduction in some cases (13, 68). In general, there is an increased

fracture risk and incidence of scoliosis corresponding to decreased BMD, such as in osteopenia

and osteoporosis (48, 66). In line with this reduced mineralization an increased fracture and

injury risk has been reported for CS patients (13). Accordingly, proper observation and focus

on early DEXA analysis is considered for CS guidelines (45, 46). Our microCT analysis of

zebrafish vertebra is not without limitations. An isotropic voxel size of 10.6 µm cannot resolve

cancellous bone structures thus overestimating bone volume. However, this limitation applies

systemically so differences between groups hold true. Other outcomes such as BMD and shape

are less affected by resolution.

Our analysis throughout development of $chd7^{-/-}$ deficient zebrafish suggests that only mild

phenotypes survive onto later stages with severity decreasing as age progresses. However, the

high lethality in both patients and our model is connected to a multisystemic phenotype in which

one mechanism is unlikely to be the exclusive underlying cause for this characteristic. The link

between Chd7 and severe spinal and vertebrae abnormalities, including kyphosis has been

observed in our samples (7, 9).

Collagen and chondrocyte depletion as novel risk factor

Our data in *chd7* deficient zebrafish indicate that reduced vBMD may not be the only

underlying risk factor for idiopathic scoliosis in CS patients. Furthermore, craniofacial

development in CS may be dependent on ECM matrix composition. Bone matrix integrity may

be weakened by a decrease in the collagen component Col2a1 in the craniofacial and vertebral

cartilage of *chd7* deficient zebrafish. Col2a1 is required in the ECM of various cell types

including osteoblasts, chondrocytes, external ligament connective tissue cells, as well as

notochord basal cells and expressed in the corresponding tissues (69, 70). Each cell type is

required for proper bone formation and could likely contribute to spinal injuries and

malformations. Furthermore, *col2a1a* expression is observed throughout ossification of the

notochord (71). Reduction in this collagen matrix in the notochord can affect proper

calcification and is known to cause spondyloepiphyseal dysplasia congenita, which affects

skeletal and spine development (72). Strikingly, *sedc* mice, which are deficient in *Col2a1*, show enlargement and malformations in the growth plates of vertebrae and degeneration of cartilage, similar to our study in zebrafish (73). *Col2a1a* is expressed in osteoblasts and chondrocytes of teleosts and regulated by *sox9*, which in turn is known to be regulated by *chd7* (71, 74-76). Sox9 is also well known to be required in craniofacial development, such as its related factor Sox10, both of which have been connected to the craniofacial phenotype in CHARGE (77). The hypertrophic morphology of the chondrocytes seen in our study may further indicate signs of an altered matrix composition and may contribute to an increased risk in osteoporosis (78, 79).

In summary, our study is the first to identify skeletal abnormalities linked to *chd7* deficiency. These abnormalities include morphological changes in the skull and vertebrae, reduced bone mineralization, impaired osteoblast activity, highly mineralized inclusions and Col2a1 deficiency in the *chd7*$^{-/-}$ zebrafish model organism. The use of this preclinical model will allow for future studies to further understand the underlying mechanisms of spinal deformities in CS.

Acknowledgements

The authors would like to thank Beatrice Steyn and Priyanka Jamadagni for experimental support, and Mark Lepik for graphic design. SP is supported by the Natural Science and Engineering Research Council (NSERC), Canadian Institutes of Health Research (CIHR), an ALS Canada-Brain Canada Career Transition Award and a FRQS Junior 1 research scholar. BMW is supported by FRQS Programme de bourses de chercheur. MR is supported by FRQS Programme de bourses de chercheur. We also thank Shriners Hospital for Children and the CIHR (165939) for financial support.

Author roles

MB, MR, BMW and SP designed the study. MB, MR and CZ performed the experiments. MB and MR drafted the manuscript. MB, MR, CZ, BMW, SP reviewed and edited the manuscript.

References

1.   Wise CA, Gao X, Shoemaker S, Gordon D, Herring JA. Understanding genetic factors in idiopathic scoliosis, a complex disease of childhood. Curr Genomics. 2008;9(1):51-9.
2.   Grauers A, Einarsdottir E, Gerdhem P. Genetics and pathogenesis of idiopathic scoliosis. Scoliosis Spinal Disord. 2016;11:45.
3.   Hall BD. Choanal atresia and associated multiple anomalies. J Pediatr. 1979;95(3):395-8.
4.   Hittner HM, Hirsch NJ, Kreh GM, Rudolph AJ. Colobomatous microphthalmia, heart disease, hearing loss, and mental retardation--a syndrome. J Pediatr Ophthalmol Strabismus. 1979;16(2):122-8.
5.   Pagon RA, Graham JM, Jr., Zonana J, Yong SL. Coloboma, congenital heart disease, and choanal atresia with multiple anomalies: CHARGE association. J Pediatr. 1981;99(2):223-7.
6.   Blake KD, Salem-Hartshorne N, Daoud MA, Gradstein J. Adolescent and adult issues in CHARGE syndrome. Clin Pediatr (Phila). 2005;44(2):151-9.
7.   Jongmans MC, Admiraal RJ, van der Donk KP, Vissers LE, Baas AF, Kapusta L, et al. CHARGE syndrome: the phenotypic spectrum of mutations in the CHD7 gene. J Med Genet. 2006;43(4):306-14.
8.   Verloes A. Updated diagnostic criteria for CHARGE syndrome: a proposal. Am J Med Genet A. 2005;133A(3):306-8.
9.   Sanlaville D, Verloes A. CHARGE syndrome: an update. Eur J Hum Genet. 2007;15(4):389-99.
10.   Doyle C, Blake K. Scoliosis in CHARGE: a prospective survey and two case reports. Am J Med Genet A. 2005;133A(3):340-3.
11.   Forward KE, Cummings EA, Blake KD. Risk factors for poor bone health in adolescents and adults with CHARGE syndrome. Am J Med Genet A. 2007;143A(8):839-45.
12.   Gao X, Gordon D, Zhang D, Browne R, Helms C, Gillum J, et al. CHD7 gene polymorphisms are associated with susceptibility to idiopathic scoliosis. Am J Hum Genet. 2007;80(5):957-65.
13.   Kumaran A, Kirk J. Endocrine manifestations of CHARGE syndrome. Endocrine Abstracts. 2013;33:P80.
14.   Lalani SR, Safiullah AM, Fernbach SD, Harutyunyan KG, Thaller C, Peterson LE, et al. Spectrum of CHD7 mutations in 110 individuals with CHARGE syndrome and genotype-phenotype correlation. Am J Hum Genet. 2006;78(2):303-14.
15.   Mitchell JA, Giangiacomo J, Hefner MA, Thelin JW, Pickens JM. Dominant CHARGE association. Ophthalmic Paediatr Genet. 1985;6(1-2):271-6.
16.   Delahaye A, Sznajer Y, Lyonnet S, Elmaleh-Berges M, Delpierre I, Audollent S, et al. Familial CHARGE syndrome because of CHD7 mutation: clinical intra- and interfamilial variability. Clin Genet. 2007;72(2):112-21.
17.   Jones KM, Saric N, Russell JP, Andoniadou CL, Scambler PJ, Basson MA. CHD7 maintains neural stem cell quiescence and prevents premature stem cell depletion in the adult hippocampus. Stem Cells. 2015;33(1):196-210.
18.   Ohta S, Yaguchi T, Okuno H, Chneiweiss H, Kawakami Y, Okano H. CHD7 promotes proliferation of neural stem cells mediated by MIF. Mol Brain. 2016;9(1):96.

19.   Okuno H, Renault Mihara F, Ohta S, Fukuda K, Kurosawa K, Akamatsu W, et al. CHARGE
syndrome modeling using patient-iPSCs reveals defective migration of neural crest cells harboring
CHD7 mutations. Elife. 2017;6.
20.   Writzl K, Cale CM, Pierce CM, Wilson LC, Hennekam RC. Immunological abnormalities in
CHARGE syndrome. Eur J Med Genet. 2007;50(5):338-45.
21.   Feng W, Shao C, Liu HK. Versatile Roles of the Chromatin Remodeler CHD7 during Brain
Development and Disease. Front Mol Neurosci. 2017;10:309.
22.   Janssen N, Bergman JE, Swertz MA, Tranebjaerg L, Lodahl M, Schoots J, et al. Mutation update
on the CHD7 gene involved in CHARGE syndrome. Hum Mutat. 2012;33(8):1149-60.
23.   Takada I, Kouzmenko AP, Kato S. Wnt and PPARgamma signaling in osteoblastogenesis and
adipogenesis. Nat Rev Rheumatol. 2009;5(8):442-7.
24.   Chen Y, Wang M, Chen D, Wang J, Kang N. Chromatin remodeling enzyme CHD7 is necessary
for osteogenesis of human mesenchymal stem cells. Biochem Biophys Res Commun.
2016;478(4):1588-93.
25.   Ogier JM, Carpinelli MR, Arhatari BD, Symons RC, Kile BT, Burt RA. CHD7 deficiency in "Looper",
a new mouse model of CHARGE syndrome, results in ossicle malformation, otosclerosis and hearing
impairment. PLoS One. 2014;9(5):e97559.
26.   Hurd EA, Capers PL, Blauwkamp MN, Adams ME, Raphael Y, Poucher HK, et al. Loss of Chd7
function in gene-trapped reporter mice is embryonic lethal and associated with severe defects in
multiple developing tissues. Mamm Genome. 2007;18(2):94-104.
27.   Patten SA, Jacobs-McDaniels NL, Zaouter C, Drapeau P, Albertson RC, Moldovan F. Role of Chd7
in zebrafish: a model for CHARGE syndrome. PLoS One. 2012;7(2):e31650.
28.   Busse B, Galloway JL, Gray RS, Harris MP, Kwon RY. Zebrafish: An Emerging Model for
Orthopedic Research. J Orthop Res. 2020;38(5):925-36.
29.   Carnovali M, Banfi G, Mariotti M. Zebrafish Models of Human Skeletal Disorders: Embryo and
Adult Swimming Together. Biomed Res Int. 2019;2019:1253710.
30.   Ofer L, Dean MN, Zaslansky P, Kult S, Shwartz Y, Zaretsky J, et al. A novel nonosteocytic
regulatory mechanism of bone modeling. PLoS Biol. 2019;17(2):e3000140.
31.   Hayes AJ, Reynolds S, Nowell MA, Meakin LB, Habicher J, Ledin J, et al. Spinal deformity in aged
zebrafish is accompanied by degenerative changes to their vertebrae that resemble osteoarthritis.
PLoS One. 2013;8(9):e75787.
32.   Bergen DJM, Kague E, Hammond CL. Zebrafish as an Emerging Model for Osteoporosis: A
Primary Testing Platform for Screening New Osteo-Active Compounds. Front Endocrinol (Lausanne).
2019;10:6.
33.   Dahdul WM, Lundberg JG, Midford PE, Balhoff JP, Lapp H, Vision TJ, et al. The teleost anatomy
ontology: anatomical representation for the genomics age. Syst Biol. 2010;59(4):369-83.
34.   Suniaga S, Rolvien T, Vom Scheidt A, Fiedler IAK, Bale HA, Huysseune A, et al. Increased
mechanical loading through controlled swimming exercise induces bone formation and mineralization
in adult zebrafish. Sci Rep. 2018;8(1):3646.
35.   Haga Y, Dominique VJ, 3rd, Du SJ. Analyzing notochord segmentation and intervertebral disc
formation using the twhh:gfp transgenic zebrafish model. Transgenic Res. 2009;18(5):669-83.
36.   Morin-Kensicki EM, Melancon E, Eisen JS. Segmental relationship between somites and
vertebral column in zebrafish. Development. 2002;129(16):3851-60.
37.   Spoorendonk KM, Hammond CL, Huitema LFA, Vanoevelen J, Schulte-Merker S. Zebrafish as a
unique model system in bone research: the power of genetics and in vivo imaging. 2010;26(2):219-24.
38.   Xi Y, Chen D, Sun L, Li Y, Li L. Characterization of zebrafish mutants with defects in bone
calcification during development. Biochem Biophys Res Commun. 2013;440(1):132-6.
39.   MacRae CA, Peterson RT. Zebrafish as tools for drug discovery. Nature reviews Drug discovery.
2015;14(10):721-31.
40.   Fleming A, Sato M, Goldsmith P. High-throughput in vivo screening for bone anabolic
compounds with zebrafish. J Biomol Screen. 2005;10(8):823-31.

41.   Boswell CW, Ciruna B. Understanding Idiopathic Scoliosis: A New Zebrafish School of Thought.
Trends Genet. 2017;33(3):183-96.
42.   Guo L, Ikegawa S, Shukunami C. Emergence of Zebrafish as a Model System for Understanding
Human Scoliosis. In: Hirata H, Iida A, editors. Zebrafish, Medaka, and Other Small Fishes: New Model
Animals in Biology, Medicine, and Beyond. Singapore: Springer Singapore; 2018. p. 217-34.
43.   Gray RS, Wilm TP, Smith J, Bagnat M, Dale RM, Topczewski J, et al. Loss of col8a1a function
during zebrafish embryogenesis results in congenital vertebral malformations. Dev Biol.
2014;386(1):72-85.
44.   Gistelinck C, Kwon RY, Malfait F, Symoens S, Harris MP, Henke K, et al. Zebrafish type I collagen
mutants faithfully recapitulate human type I collagenopathies. Proc Natl Acad Sci U S A.
2018;115(34):E8037-E46.
45.   de Geus CM, Free RH, Verbist BM, Sival DA, Blake KD, Meiners LC, et al. Guidelines in CHARGE
syndrome and the missing link: Cranial imaging. Am J Med Genet C Semin Med Genet.
2017;175(4):450-64.
46.   Trider CL, Arra-Robar A, van Ravenswaaij-Arts C, Blake K. Developing a CHARGE syndrome
checklist: Health supervision across the lifespan (from head to toe). Am J Med Genet A.
2017;173(3):684-91.
47.   Parichy DM, Elizondo MR, Mills MG, Gordon TN, Engeszer RE. Normal table of postembryonic
zebrafish development: staging by externally visible anatomy of the living fish. Dev Dyn.
2009;238(12):2975-3015.
48.   Aspray TJ, Hill TR. Osteoporosis and the Ageing Skeleton. Subcell Biochem. 2019;91:453-76.
49.   Bensimon-Brito A, Cardeira J, Cancela ML, Huysseune A, Witten PE. Distinct patterns of
notochord mineralization in zebrafish coincide with the localization of Osteocalcin isoform 1 during
early vertebral centra formation. BMC Dev Biol. 2012;12:28.
50.   Pauli S, Bajpai R, Borchers A. CHARGEd with neural crest defects. Am J Med Genet C Semin
Med Genet. 2017;175(4):478-86.
51.   Hernandez RE, Galitan L, Cameron J, Goodwin N, Ramakrishnan L. Delay of Initial Feeding of
Zebrafish Larvae Until 8 Days Postfertilization Has No Impact on Survival or Growth Through the
Juvenile Stage. Zebrafish. 2018;15(5):515-8.
52.   O'Keefe JH, Bergman N, Carrera-Bastos P, Fontes-Villalba M, DiNicolantonio JJ, Cordain L.
Nutritional strategies for skeletal and cardiovascular health: hard bones, soft arteries, rather than vice
versa. Open Heart. 2016;3(1):e000325.
53.   Blake KD, Hudson AS. Gastrointestinal and feeding difficulties in CHARGE syndrome: A review
from head-to-toe. Am J Med Genet C Semin Med Genet. 2017;175(4):496-506.
54.   Foppiani L, Maffe A, Forzano F. CHARGE syndrome as unusual cause of hypogonadism:
endocrine and molecular evaluation. Andrologia. 2010;42(5):326-30.
55.   Cloney K, Steele SL, Stoyek MR, Croll RP, Smith FM, Prykhozhij SV, et al. Etiology and functional
validation of gastrointestinal motility dysfunction in a zebrafish model of CHARGE syndrome. FEBS J.
2018;285(11):2125-40.
56.   Ortuno MJ, Susperregui AR, Artigas N, Rosa JL, Ventura F. Osterix induces Col1a1 gene
expression through binding to Sp1 sites in the bone enhancer and proximal promoter regions. Bone.
2013;52(2):548-56.
57.   Kague E, Roy P, Asselin G, Hu G, Simonet J, Stanley A, et al. Osterix/Sp7 limits cranial bone
initiation sites and is required for formation of sutures. Dev Biol. 2016;413(2):160-72.
58.   Lapunzina P, Aglan M, Temtamy S, Caparros-Martin JA, Valencia M, Leton R, et al. Identification
of a frameshift mutation in Osterix in a patient with recessive osteogenesis imperfecta. Am J Hum
Genet. 2010;87(1):110-4.
59.   Niu P, Zhong Z, Wang M, Huang G, Xu S, Hou Y, et al. Zinc finger transcription factor Sp7/Osterix
acts on bone formation and regulates col10a1a expression in zebrafish. Science Bulletin.
2017;62(3):174-84.

60.   Rouillard AD, Gundersen GW, Fernandez NF, Wang Z, Monteiro CD, McDermott MG, et al. The
harmonizome: a collection of processed datasets gathered to serve and mine knowledge about genes
and proteins. Database (Oxford). 2016;2016.
61.   Tilley MK, Justice CM, Swindle K, Marosy B, Wilson AF, Miller NH. CHD7 gene polymorphisms
and familial idiopathic scoliosis. Spine (Phila Pa 1976). 2013;38(22):E1432-6.
62.   Faulkner KG. Bone matters: are density increases necessary to reduce fracture risk? J Bone
Miner Res. 2000;15(2):183-7.
63.   Li XF, Li H, Liu ZD, Dai LY. Low bone mineral status in adolescent idiopathic scoliosis. Eur Spine
J. 2008;17(11):1431-40.
64.   Currey JD. Bones: Structure and Mechanics: Princeton University Press; 2006.
65.   Zimmermann EA, Busse B, Ritchie RO. The fracture mechanics of human bone: influence of
disease and treatment. Bonekey Rep. 2015;4:743.
66.   Sarioglu O, Gezer S, Sarioglu FC, Koremezli N, Kara T, Akcali O, et al. Evaluation of vertebral
bone mineral density in scoliosis by using quantitative computed tomography. Pol J Radiol.
2019;84:e131-e5.
67.   Wang Z, Chen H, Yu YE, Zhang J, Cheuk KY, Ng BK, et al. Unique local bone tissue characteristics
in iliac crest bone biopsy from adolescent idiopathic scoliosis with severe spinal deformity. Sci Rep.
2017;7:40265.
68.   Modi A, Piziak V. MON-355 Charge Syndrome: Unusual Cause of Hypogonadism Leading to
Osteoporosis. J Endocr Soc. 2020;4(Suppl 1):MON-355.
69.   Kaneko T, Freeha K, Wu X, Mogi M, Uji S, Yokoi H, et al. Role of notochord cells and sclerotome-
derived cells in vertebral column development in fugu, Takifugu rubripes: histological and gene
expression analyses. Cell Tissue Res. 2016;366(1):37-49.
70.   Jian QL, HuangFu WC, Lee YH, Liu IH. Age, but not short-term intensive swimming, affects
chondrocyte turnover in zebrafish vertebral cartilage. PeerJ. 2018;6:e5739.
71.   Dale RM, Topczewski J. Identification of an evolutionarily conserved regulatory element of the
zebrafish col2a1a gene. Dev Biol. 2011;357(2):518-31.
72.   Vikkula M, Metsaranta M, Ala-Kokko L. Type II collagen mutations in rare and common
cartilage diseases. Ann Med. 1994;26(2):107-14.
73.   Donahue LR, Chang B, Mohan S, Miyakoshi N, Wergedal JE, Baylink DJ, et al. A missense
mutation in the mouse Col2a1 gene causes spondyloepiphyseal dysplasia congenita, hearing loss, and
retinoschisis. J Bone Miner Res. 2003;18(9):1612-21.
74.   Akiyama H, Chaboissier MC, Martin JF, Schedl A, de Crombrugghe B. The transcription factor
Sox9 has essential roles in successive steps of the chondrocyte differentiation pathway and is required
for expression of Sox5 and Sox6. Genes Dev. 2002;16(21):2813-28.
75.   Bajpai R, Chen DA, Rada-Iglesias A, Zhang J, Xiong Y, Helms J, et al. CHD7 cooperates with PBAF
to control multipotent neural crest formation. Nature. 2010;463(7283):958-62.
76.   Eames BF, Amores A, Yan YL, Postlethwait JH. Evolution of the osteoblast: skeletogenesis in gar
and zebrafish. BMC Evol Biol. 2012;12:27.
77.   Asad Z, Pandey A, Babu A, Sun Y, Shevade K, Kapoor S, et al. Rescue of neural crest-derived
phenotypes in a zebrafish CHARGE model by Sox10 downregulation. Hum Mol Genet.
2016;25(16):3539-54.
78.   Hall AC. The Role of Chondrocyte Morphology and Volume in Controlling Phenotype-
Implications for Osteoarthritis, Cartilage Repair, and Cartilage Engineering. Curr Rheumatol Rep.
2019;21(8):38.
79.   Lian C, Wang X, Qiu X, Wu Z, Gao B, Liu L, et al. Collagen type II suppresses articular chondrocyte
hypertrophy and osteoarthritis progression by promoting integrin beta1-SMAD1 interaction. Bone Res.
2019;7:8.
620

Fig 1. Mineralization in larval *chd7^{-/-}* mutants

A. Alcian blue staining of 6 dpf control and *chd7^{-/-}* zebrafish larvae, ventral view (De: dentary; Pq: palatoquadrate; Ch: ceratohyal) B. Calcein staining of 6 dpf control and *chd7^{-/-}* zebrafish larvae, ventral view (De: dentary; Q: quadrate; Op: opercle) C. Dorsal view of control and *chd7^{-/-}* larvae at 9 dpf showing severe spinal deformities in 43 out of 110 larvae screened. Deformities were present at both precaudal and caudal region (arrows). D. Lateral view of calcein staining at 9 dpf showing mineralization of vertebrae with access to food starting at 5dpf and calcein staining at 9 dpf during starvation. Arrow indicates mineralization at the first 4 (Weberian) vertebrae E. Number of vertebrae in wildtype and *chd7^{-/-}* (fed and starved) larvae showing calcification signal as tested by calcein and F. Number of Weberian vertebrae (vertebrae 1-4) showing complete calcification (see arrows in b) G-H. RT-qPCR of osteogenesis related genes screened at 9dpf, showing relative fold change. (Significance: *p<0.05; **p<0.01; ***p<0.001)

Fig 2. Skull deformities and Weberian apparatus

A, C. Lateral view of Control and *chd7⁻/⁻* adult zebrafish skull and B, D. ventral view of Control and *chd7⁻/⁻* adult zebrafish of mineralized tissue E. Angle of mandibular arch in adult zebrafish F. Length (in mm) of mandibular bone and G. Angle of the craniofacial (skull) structure H. Representative microCT overview image of the spinal cord from a 1-year old control and a *chd7⁻/⁻* adult zebrafish I. MicroCT Image of the Weberian apparatus from a 1-year old control and a *chd7⁻/⁻* mutants, indicating structures of intercalarium (red), tripus (yellow) and parapophysis (blue). Significance of student t-test with Welch's correction is included in the graphs with: *p<0.05; **p<0.01, ***p<0.001

Fig 3. Precaudal vertebrae show inefficiency for proper mineralization

A. Sketch diagram of precaudal vertebrae (frontal and lateral) indicating structure and measured angles.

B-C. (top) Individual rendering of precaudal vertebrae, frontal view of tissue over 0.41 g HA/cm^3

threshold and (bottom) vBMD intensity map (Range blue-red; 0.41 – 1.00 g HA/cm^3) showing

morphological abnormalities and growth zone malformations with highly mineralized inclusions

(arrow) D. vBMD of whole vertebrae showing increasing density with increasing age and e. increasing

ratio of vBMD arch/vertebrae body. F-H neural arch angle, BV, TV and ratio BV/TV. (n=8/genotype

were analyzed). Significance of student t-test with Welch's correction is included in the graphs with:

*p<0.05; **p<0.01. F-test results are indicated underneath as F$_{\text{degree of freedom numerator, degree of freedom denominator}}$

(F-value); p-value: *p<0.05; **p<0.01; ***p<0.001

Fig 4. Caudal vertebrae show abnormal structure

A. Sketch diagram of precaudal vertebrae indicating structure and measured angles (frontal and lateral).
B-C. (top) Individual rendering of caudal vertebrae, frontal view of tissue over 0.41 g HA/cm^3 threshold
and (bottom) vBMD intensity map (Range blue-red; 0.41 – 1.00 g HA/cm3) showing morphological
abnormalities and growth zone malformations with highly mineralized inclusions (arrow) D. vBMD of
whole vertebrae and E. increasing ratio of vBMD arch/vertebrae body. F-H neural arch angle, BV, TV
and ratio BV/TV. (n=8/genotype were analyzed). Significance of student t-test with Welch's correction
is included in the graphs with: *p<0.05; **p<0.01. F-test results are indicated underneath as $F_{\text{degree of freedom numerator, degree of freedom denominator}}$ (F-value); p-value: *p<0.05; **p<0.01; ***p<0.001

Fig 5. ECM Collagen2a1 deficiency in larvae and 1-year-old *chd7*⁻ᐟ⁻ adults

A. Immunofluorescence of Col2a1 in 5 dpf Cntrl (n=9) and *chd7*⁻ᐟ⁻ (n=8) larvae, ventral view of craniofacial cartilage (De: dentary; Pq: palatoquadrate; Ch: ceratohyal) B. Immunofluorescence of Col2a1 in precaudal vertebrae sections of Cntrl and *chd7*⁻ᐟ⁻ mutants with Col2a1 in green and DAPI in blue. C. H&E staining of precaudal vertebrae section of Cntrl and *chd7*⁻ᐟ⁻ mutants D. Safranin O/fast green staining of Weberian and precaudal vertebrae sections of Cntrl and *chd7*⁻ᐟ⁻ mutants showing cartilage in red. E. RT-qPCR of *col2a1a* at 9dpf. F. Total nuclei count in Weberian and precaudal vertebral cartilage detected in H&E staining per 100μm² in Cntrl (N=4, n=17) and *chd7*⁻ᐟ⁻ (N=4, n=18) G. Chondrocyte nuclei count in precaudal vertebral cartilage detected in Safranin O/fast green staining per 100μm² Cntrl (N=4, n=18) and *chd7*⁻ᐟ⁻ (N=4, n=19) Scale bar presents 100μm. Significance of student t-test with: *$p<0.05$; **$p<0.01$ ***$p<0.001$