Classification: Biological Sciences; Evolution

Geography is more important than life history in the recent

diversification of the tiger salamander complex

Kathryn M. Everson[a,1], Levi N. Gray[a,1], Angela G. Jones[a], Nicolette M. Lawrence[a], Mary E.

Foley[a], Kelly L. Sovacool[a,b], Justin D. Kratovil[a,c], Scott Hotaling[a,d], Paul M. Hime[a,e], Andrew

Storfer[d], Gabriela Parra-Olea[f], X. Aguilar-Miguel[g], Eric M. O'Neill[a], Luis Zambrano[h], H.

Bradley Shaffer[i], David W. Weisrock[a,2]

[a] Department of Biology, University of Kentucky, Lexington, KY 40506

[b] Department of Computational Medicine & Bioinformatics, University of Michigan, Ann Arbor,

MI 48109

[c] Ecology and Evolutionary Biology Department, University of Connecticut, Storrs, CT 06269

[d] School of Biological Sciences, Washington State University, Pullman, WA 99164

[e] Biodiversity Institute, University of Kansas, Lawrence, KS 66045

[f] Departamento de Zoología, Instituto de Biología, Universidad Nacional Autónoma de México,

Ciudad de México, México

[g] CIRB, Facultad de Ciencias, Universidad Autónoma del Estado de México, Toluca, Edo. de

México, México

[h] Facultad de Medicina Veterinaria y Zootecnia, Universidad Autónoma de Sinaloa, Culiacán

Rosales 82260, Sinaloa, Mexico

[i]Department of Ecology and Evolutionary Biology, and La Kretz Center for California

Conservation Science, University of California, Los Angeles, CA 90095

[1] K.M.E. and L.N.G. contributed equally to this work

[2] To whom correspondence may be addressed. Email: david.weisrock@uky.edu

ABSTRACT

The North American tiger salamander species complex, including its flagship species the

axolotl, has long been a source of biological fascination. The complex exhibits a wide range of

variation in developmental life history strategies, including populations and individuals that

undergo metamorphosis and those able to forego metamorphosis and retain a larval, aquatic

lifestyle (i.e., paedomorphosis). Such disparate life history strategies are assumed to cause

populations to become reproductively isolated, but the degree to which they have actually shaped

population- and species-level boundaries is poorly understood. Using a large multi-locus dataset

from hundreds of samples across North America, we identified genetic clusters with clear signs

of admixture across the geographic range of the tiger salamander complex. Population clusters

often contain a mixture of paedomorphic and metamorphic taxa, and we conclude that geography

has played a large role in driving lineage divergence relative to obligate paedomorphosis in this

system. This conclusion is bolstered by model-based analyses demonstrating gene flow between

metamorphic and paedomorphic populations. Even the axolotl, a paedomorphic species with an

isolated native range, apparently has a history of gene flow with its neighboring populations.

This fine-scale genetic perspective on life-history variation establishes a framework for

understanding how plasticity, local adaptation, and gene flow contribute to lineage divergence.

The axolotl is currently used as the vertebrate model system in regenerative biology, and our

findings chart a course for more informed use of these and other tiger salamander species in

experimental and field research, including conservation priorities.

Keywords: *Ambystoma*, life history, phylogenetics, population genomics, salamanders

SIGNIFICANCE STATEMENT

Population structure and speciation are shaped by a variety of biotic and abiotic factors.

In the tiger salamander complex, one factor that may influence diversification is life history:

some taxa are obligately paedomorphic–a condition where adults maintain an aquatic, larval

phenotype–while others are facultatively paedomorphic or entirely metamorphic. Using a large

multi-locus dataset, we found evidence of gene flow and/or panmixia between obligately and

facultatively paedomorphic taxa, suggesting that an obligately paedomorphic life history is not a

strong driver of speciation in the tiger salamander complex. We also recovered a history of gene

flow between the critically endangered axolotl and its neighboring populations, providing

important information for its conservation and captive management.

INTRODUCTION

Life history—the complement of traits affecting survival and reproduction over an organism's lifetime—affects ecology and dispersal, and therefore plays a major role in shaping population structure and speciation (1, 2). The axolotl (*Ambystoma mexicanum*) and related salamander species of the North American *Ambystoma tigrinum* complex (Table 1) display enormous variation in life history (3). Some members of this group are considered obligate paedomorphs, where sexually mature adults retain a larval, aquatic body plan that includes external gills and an enlarged tail fin (4). Others are considered obligate metamorphs, transforming from aquatic larvae to terrestrial adults. Most populations are facultative in their developmental life history strategy, transforming under certain genetic and/or environmental conditions (5–8). This lability in life history is believed to have played an important role in the diversification of the *A. tigrinum* complex, particularly in the Trans-Mexican Volcanic Belt (TMVB) region, where several essentially paedomorphic species (including the axolotl) have been described (9–11). Obligate paedomorphosis is estimated to have evolved in this region multiple times (11–13) in association with relatively large, permanent bodies of water. Presumably, restricted gene flow between these isolated, paedomorphic populations led to speciation as well as morphological adaptations to the aquatic lifestyle (14).

The term "species complex," while not a formal taxonomic category, is often used to describe groups of closely related lineages early in the divergence process. The tiger salamander species complex has been highlighted as a potentially valuable example of a recent radiation (15) that could provide insight into the early mechanisms initiating and/or maintaining diversity (16– 21). However, such species complexes poses a number of challenges: phenotypic differences among lineages may be subtle or absent (i.e., cryptic species), ancestral polymorphisms may be

Table 1. Currently recognized species in the *Ambystoma tigrinum* complex. Life histories, habitat preferences, and conservation status were compiled from AmphibiaWeb (2019) and the IUCN (2019).

Species	Description Reference	Life History	Habitat Preference	Conservation Status
Ambystoma altamirani	Dugès, 1895	Facultative paedomorph	Streams, occasional ponds	Endangered
Ambystoma amblycephalum	Taylor, 1940	Facultative paedomorph	Ponds	Critically endangered
Ambystoma andersoni	Krebs & Brandon, 1984	Paedomorph	Lake Zacapu	Critically endangered
Ambystoma bombypellum	Taylor, 1940	Facultative paedomorph	Lakes / Ponds, occasional streams	Data deficient
Ambystoma californiense	Gray 1853	Metamorph	Vernal pools	Vulnerable
Ambystoma dumerilii	(Dugès, 1870)	Paedomorph	Lake Patzcuaro	Critically endangered
Ambystoma flavipiperatum	Dixon, 1963	Facultative paedomorph	Lakes / Ponds, occasional streams	Endangered
Ambystoma granulosum	Taylor, 1944	Facultative paedomorph	Lakes / Ponds, occasional streams	Critically endangered
Ambystoma leorae	(Taylor, 1943)	Facultative paedomorph	Streams	Critically endangered
Ambystoma lermaense	(Taylor, 1940)	Facultative; often paedomorphic	Lakes / Ponds	Endangered
*Ambystoma mavortium**	Baird, 1850	Facultative paedomorph	Lakes / Ponds	Least concern
Ambystoma mexicanum	(Shaw and Nodder, 1798)	Paedomorph	Lakes Xochimilco & Chalco	Critically endangered
Ambystoma ordinarium	Taylor, 1940	Facultative paedomorph	Streams, ponds	Endangered
Ambystoma rivulare	(Taylor, 1940)	Facultative paedomorph	Streams	Endangered
Ambystoma rosaceum	Taylor, 1941	Facultative paedomorph	Streams, occasional ponds	Least concern
Ambystoma silvense	Webb, 2004	Facultative paedomorph	Lakes / Ponds	Data deficient
Ambystoma taylori	Brandon et al., 1982	Paedomorph	Lake Alchichica (saline)	Critically endangered
Ambystoma tigrinum	Green, 1825	Rare paedomorph	Lakes / Ponds	Least concern
Ambystoma velasci	(Dugès, 1888)	Facultative paedomorph	Lakes / Ponds, Streams	Least concern

*includes 5 recognized subspecies: A. m. mavortium Baird 1850, A. m. melanostictum (Cooper 1859), A. m. diaboli Dunn 1940, A. m. nebulosum Hallowell 1853, and A. m. stebbinsi Lowe 1954

stochastically inherited (i.e., incomplete lineage sorting), and reproductive barriers may be

porous (i.e., speciation with gene flow). Furthermore, phenotypic variation may not reflect

lineage divergence or speciation processes; for example, trait variation may be the result of

plasticity or balancing selection (22). The reliance on phenotype for species diagnosis in the *A.*

tigrinum complex may require particular scrutiny, as several phenotypic traits, including

developmental state and adult color pattern, can be plastic and highly variable (23–25). Given

these challenges, fundamental questions remain concerning the evolutionary distinctiveness of

component lineages within the tiger salamander complex, calling into question the current

taxonomy as an accurate reflection of the underlying population biology.

Previous research has produced mixed results regarding levels of genetic differentiation

among lineages in the *A. tigrinum* complex, as well as the relative importance of paedomorphosis

as a driver of diversification. While population- and species-level structure is evident, an

important theme emerging across multiple studies has been that reproductive barriers are porous

or incomplete (11, 26, 27). Indeed, Shaffer and McKnight (11) noted "the striking lack of

differentiation among the 14 species of the tiger salamander complex" (p. 425). Some studies

have found an elevated degree of population structure among paedomorphic populations relative

to metamorphic populations (12, 28), but others have demonstrated that paedomorphic taxa and

neighboring metamorphic populations are not fully reproductively isolated (29) and can produce

viable hybrid offspring under laboratory conditions (30, 31). Furthermore, there is evidence that

some populations considered to be obligately paedomorphic are capable of transforming at low

rates and may be more appropriately considered facultatively paedomorphic (10, 32, 33),

although the absence of transformed individuals in nature is difficult to verify without targeted

ecological studies (34).

To illuminate the processes underlying diversification in the tiger salamander complex,

an important first step must be a range-wide assessment of population structure and the

clarification of population- and species-level boundaries. With this groundwork, insights into the

role of paedomorphosis and diversification can be addressed. For a study system of this

geographic and taxonomic scale, robust inference of population structure requires information

derived from throughout the genome combined with thorough range-wide sampling. To meet

these criteria, we expand on a large multi-locus dataset containing 95 nuclear loci for 93

individuals (35), to produce a data matrix for 347 individuals across the full geographic range of

tiger salamanders (Fig. S1). Given the complex and somewhat checkered history of the group's

taxonomy, we take a naïve approach, performing population structure analyses without *a priori*

identification of taxa to resolve geographic genetic clusters and characterize patterns of

admixture. We couple these results with phylogenetic network analyses and model-based tests of

migration between metamorphic and paedomorphic populations to provide an additional

perspective on the degree to which populations are genetically connected, now or in the recent

past. We then infer a species tree for identified genetic clusters in an attempt to provide an

updated working hypothesis for the evolution of the group. Finally, we test the overarching

hypothesis that life history evolution has driven speciation in the tiger salamander complex,

particularly through the assumed isolation and independence of small-range endemics classified

as paedomorphic species.

RESULTS AND DISCUSSION

Identification of genetic lineages and gene flow

A principal components analysis (PCA) and a discriminant analysis of principal

components (DAPC; 36) both recovered *Ambystoma californiense,* the only confirmed obligate

metamorph in the complex, as genetically distinct from all other tiger salamanders (Fig. 1A, S2,

S3), which is consistent with previous work (e.g., 11, 37). After removing *A. californiense* from

the dataset, DAPC analysis supported the recognition of three primary genetic clusters: U.S.

(including samples from southern Canada), northwestern Mexico, and central Mexico (Fig. 1B,

S4). All populations in the U.S. and northern Mexico clusters are considered facultatively

paedomorphic or primarily metamorphic (38). The central Mexico cluster, however, contains a

wealth of life history extremes: four currently recognized species are considered to be obligate

paedomorphs and ten others are facultatively paedomorphic to varying degrees (13, 39). A PCA

of these data produced similar clustering results, with ordination patterns largely mirroring

geographic sampling (Fig. S5). For discussion of the taxonomic implications of our results, see

the Supplementary Material.

Within central Mexico, both DAPC and STRUCTURE analyses identified four

geographic clusters (hereafter referred to as CM1-CM4) with signatures of admixture within and

among them (Figs. 2A, S6-S8). Network analysis (40) also recovered groups that generally

correspond to CM1-CM4, but with a large number of reticulate branches (Fig. 3). This starburst-

like pattern indicates the presence of conflicting topologies within the dataset (40), which can be

caused by biological processes including incomplete lineage sorting (a historical legacy of recent

diversification) and gene flow, either current or in the evolutionarily recent past. Phylogenetic

A) DAPC of full dataset

B) DAPC without _A. californiense_

Figure 1. Results of discriminant analysis of principal components (DAPC) on (A) the entire genomic dataset (without _A. opacum_ and _A. texanum_ outgroups, and (B) the genomic dataset without the highly divergent _A. californiense_. In both plots, points represent individuals and ellipses show the groups identified by DAPC. The first and second principal components from the DAPC are on the x and y axes, respectively. Results from additional values of _K_ are provided in Supplementary Figures 2 and 3.

Figure 2. Results of population genomic analyses on the *Ambystoma tigrinum* species complex. (A) STRUCTURE analysis of the central Mexico cluster, which identified four major groups. Each vertical bar represents an individual, while the y-axis shows the membership probability for each group. Pie charts mapped below the assignment plot show the average group membership probability for that locality. Some localities were combined to a single pie chart to reduce visual clutter; see Table S1 for specific coordinates of each sample. The range of the Trans-Mexican Volcanic Belt (TMVB) is identified by shading and is modified from Ferrari (2004). (B-E) Detailed STRUCTURE analyses of each central Mexico subgroup CM1-CM4. A geographic outlier placed in CM2 is thought to represent a geographic range introduction and is not shown here, but is discussed in supplementary text and shown in Fig. S6. The range of the Cuenca Oriental is outlined in red and is modified from Percino-Daniel et al. (2016). (F,G) STRUCTURE results from analyses of the U.S. and northern Mexico groups, respectively. Detailed results are shown in Figs. S8-S9. Black squares on the maps denote the type localities of all species included in this study. On each STRUCTURE membership plot, salamander face symbols denote the presence of obligately or facultatively paedomorphic populations (with or without gills, respectively); groups labeled with both symbols contain a mixture of life history strategies.

Figure 3. Results of phylogenetic network analyses (left) and the quartets-based phylogenetic analysis (right) of the *Ambystoma tigrinum* species complex. Clades are colored according to their cluster assignments in population genetic analyses, as shown in detail in Fig. 2. In phylogenetic networks, species names are only shown for taxa that are obligately or strongly paedomorphic; all other taxa (not shown) are considered to be facultatively paedomorphic or metamorphic. Note that the quartet tree analysis used a subset of individuals, indicated by solid circles on the phylogenetic networks. Phylogenetic results from Bayesian and maximum-likelihood concatenated analyses were largely concordant with the quartets tree (Fig. S10), although one important difference (the relationship of the eastern U.S. clade) is indicated by a dashed line. Bootstrap support (BS) values are indicated on each node of the quartets tree (BS values > 95 are not shown), while tips are labeled with individual IDs (Table S1). Branch lengths are not scaled to time or substitution rate.

analyses (41–43) recovered monophyletic CM1, CM2, and CM4 clusters (see below for

discussion of CM3), but support values were often low within the central Mexico clade (Fig. 3).

Based on type locality and range information, the CM1 cluster includes three stream-

breeding and facultatively paedomorphic taxa: *Ambystoma altamirani* Duges 1895 (44)*, A.*

leorae Taylor 1943 (45)*,* and *A. rivulare* Taylor 1940 (9)*.* We did not find evidence that these

three taxa are reproductively isolated. Subsequent analyses of this cluster identified two admixed

populations associated with eastern and western portions of its geographic distribution (Fig. 2B).

Phylogenetically, CM1 was monophyletic and sister to all other central Mexican groups (Fig. 3),

consistent with their original placement in a distinct genus, *Rhyacosiredon* Dunn 1928 (46).

Analyses of CM2 grouped *A. lermaense* Taylor 1940 (9) with samples corresponding to

the type localities and range of *A. bombypellum* Taylor 1940 (9) and *A. granulosum* Taylor 1944

(47). There was no evidence for genetic isolation among any of these three taxa (Fig. 2C).

Within CM3, population-genetic analyses first separated samples of *A. ordinarium* Taylor

1940 (5) from the rest (with no hybridization detected; Fig 2D). Other samples were

geographically structured according to the northern and southern portions of the distribution (Fig

1E). The more southern cluster included two obligately paedomorphic taxa, *A. andersoni* Krebs

& Brandon 1984 (32) and *A. dumerilii* Duges 1870 (48), each endemic to single lake systems

(Lakes Zacapú and Pátzcuaro, respectively). However, our results identified admixture between

these lakes and other nearby localities. The samples from other CM3 localities likely represent *A.*

amblycephalum Taylor 1940 (9) and *A. flavipiperatum* Dixon 1963 (49).

CM4 included the axolotl, *A. mexicanum* Shaw & Nodder 1798 (50), and *A. taylori*

Brandon et al. 1981 (10)—both primarily considered to be near-obligate paedomorphs—as well

as samples assigned to the polymorphic *A. velasci* Duges 1888 (51). Subsequent DAPC and

STRUCTURE analyses of CM4 placed all but one *A. mexicanum* individual sampled from Lakes

Xochimilco and Chapultepec (i.e., the remaining habitats of the axolotl) in a distinct cluster (Fig.

2E, S7-S8), with evidence of some admixture in samples to the northwest of Mexico City.

We used a Bayesian model-testing framework (Fig. 4A) to test for genetic isolation of

paedomorphic populations in three central Mexico groups. Specifically, we tested for the

presence or absence of gene flow between (near) obligate paedomorphs (*A. andersoni* and *A.*

dumerilii in CM3, and *A. mexicanum* and *A. taylori* in CM4) and their surrounding facultatively

paedomorphic populations. We also tested for gene flow between *A. lermaense* [which has been

described as "often neotenic [= paedomorphic]" (9, pg. 427)] and remaining populations in CM2.

In four out of the five groups, we recovered models of bidirectional gene flow or complete

panmixia between putatively obligate paedomorphs and facultative paedomorphs (Fig. 4B).

Alternative models of no migration or unidirectional migration were rejected with decisive Bayes

factors (Table S3). Migration rate estimates for full-migration scenarios consistently identified

higher levels of gene flow from facultative to obligate paedomorphic populations. We found

support for genetic isolation of one paedomorphic taxon, *A. andersoni* (Fig. 4B), a species

endemic to Lake Zacapú, Michoacán. As we are unable to account for landscape barriers or

sampling gaps in these analyses, we cannot be certain whether the lack of gene flow we

recovered for *A. andersoni* is due to its paedomorphic life history or to its geography. Another

important caveat to our migration analyses is that divergence time is not included as a model

parameter, which prevents distinguishing between historical and contemporary gene flow. Thus,

our ability to detect ongoing gene flow in these tests is limited.

Within the U.S. and Canada, DAPC and STRUCTURE identified three genetic clusters

(K = 3) excluding the allopatric, more differentiated *A. californiense,* with clear signs of

Figure 4. (A) Demographic models used to test hypotheses of gene flow. For each species considered to be obligately (*A. andersoni*, *A. dumerilii*, *A. taylori*, and *A. mexicanum*) or often (*A. lermaense*) paedomorphic, we evaluated five models of population structure and migration between that taxon and the facultatively paedomorphic taxa in the same genetic cluster (CM1-CM4). (B) Results of demographic model tests. Θ denotes the median estimated mutation-scaled effective size of each population. Values next to each arrow are median mutation-scaled migration rates ($M = m/\mu$, where m is the fraction of immigrants in each generation and μ is the mutation rate per generation per site). For both Θ and M, 95% confidence intervals are shown in brackets below the median value. Gray circles denote obligately paedomorphic lineages, while black circles denote facultatively paedomorphic lineages.

admixture along zones of contact (Figs. 2F, S9, S10). Geographically, these clusters are

associated with the eastern U.S., the central + western U.S., and the southwestern U.S. + Rocky

Mountains (hereafter, we refer to the latter two groups as U.S. central and U.S. Rocky

Mountains, respectively). Phylogenetic analyses corroborated these results, albeit with extensive

reticulations among clusters in the phylogenetic network (Fig. 3) and mixed support for species-

tree relationships among the three major groups (Fig. 3 vs. Fig. S12). Further exploration of

population structure within each of the three U.S. clusters recovered additional patterns of

differentiation (Figs. S9, S10), including northern and southern clusters in the eastern U.S. and

central U.S., and an allopatric cluster restricted to the Pacific Northwest. Overall, our U.S. results

are similar to mtDNA-based results from previous studies (11, 52), which identified haplotype

clades associated with the eastern U.S., the northern Great Plains, the southern Great Plains, and

the Pacific Northwest.

All analyses of the northern Mexico group identified two genetic clusters ($K = 2$; Figs.

2G, S11). The northern cluster corresponds to *A. rosaceum* Taylor 1941 (53), while the southern

cluster could represent either a second population of *A. rosaceum* or *A. silvense* Webb 2004 (54);

we did not include any samples from the type locality of *A. silvense* in this study. Phylogenetic

analyses recovered each of these two northern Mexico clusters as monophyletic, but not sister to

one another, and with few reticulate nodes in the phylogenetic network (Fig. 3, S12).

The influence of life history on diversification

A popular perspective on diversification in this species complex has been that the

evolution of obligate paedomorphosis promotes reproductive isolation and, ultimately, speciation

(10, 11). However, an important theme of our results is that taxa with different life history

strategies commonly cluster together by geography in population genetic and phylogeographic

analyses (Figs. 2, 3). Moreover, our migration analyses support relatively recent gene flow

and/or panmixia between facultative and paedomorphic populations (Fig. 4). Thus, the evolution

of life history extremes does not appear to be the main driver of speciation in the tiger

salamander complex. It is also worth noting that facultative paedomorphosis is the inferred

ancestral condition (8) and nearly all extant lineages in the *Ambystoma tigrinum* complex are

capable of metamorphosis, either in the wild or in the lab. Only *A. dumerilii* is not known to

survive for long after metamorphosis (55). That only one described taxon in the complex is

putatively fixed for paedomorphosis raises several questions: Are the rest of the primarily

paedomorphic populations merely *en route* to fixation, rather than actually fixed? Do facultative

populations fluctuate in paedomorphic frequency over time, dependent on prevailing

environmental and/or demographic conditions (5)? Do transforming and non-transforming

individuals readily interbreed outside of the laboratory setting? While these remain open

questions, below we provide some relevant context for exploration and future research.

Theory predicts that paedomorphosis in salamanders is most likely to become fixed in a

population when ecological conditions are favorable in the larval, aquatic environment but

unfavorable on land (e.g., when terrestrial habitats are arid or resource-limited) (12, 56, 57).

However, it is not known how long these fixations persist over evolutionary time; obligate

paedomorphic lineages may have arisen and gone extinct multiple times over tiger salamander

evolutionary history (58). One situation that would result in complete speciation is if

paedomorphosis became fixed in the population and migration with surrounding facultatively

paedomorphic populations ceased due to strong assortative mating and/or selection against

hybrids (59–61). However, assortative mating within life history morphs has not been

demonstrated in other salamander groups (62–64) and there is evidence that different tiger

salamander morphs do interbreed (65). In fact, the only known instance of positive assortative

mating in *A. tigrinum* is associated with tail length (66, 67), which might favor metamorphosed

males as paedomorphic individuals can have relatively short tails (32). The evidence for

selection against intermorph offspring is also scarce. Captive breeding using artifical fertilization

has produced *A. mexicanum* x *A. tigrinum* hybrids (7, 31, 68) as well as *A. dumerilii* x *A.*

tigrinum and *A. dumerilii* x *A. rivularis* hybrids, although the fertility of offspring was not

investigated in *A. dumerilii* crosses (31). This suggests that postzygotic barriers are weak,

although the extent to which divergent lineages would interbreed in the wild remains a key

question in need of investigation.

If biological barriers to reproduction are limited or absent, there are several potential

mechanisms by which dispersal could mediate gene flow between paedomorphic and

metamorphic populations. First, metamorphosed individuals from facultatively paedomorphic

populations could move into bodies of water containing paedomorphic populations. Our

demographic analyses support this scenario, as they recovered higher rates of migration entering

rather than exiting obligately paedomorphic groups (Fig. 4). However, demographic analyses

also estimated non-zero emigration out of paedomorphic populations; thus, "obligate"

paedomorphs might also occasionally metamorphose and disperse. Occasional metamorphosis

has been documented in several putatively paedomorphic populations (9, 10, 33, 69, 70) and

metamorphosis can occur at higher frequencies in dispersing males in facultative populations

(71). Even a low frequency of transformed adults is likely sufficient to maintain evolutionary

cohesion with surrounding populations (10, 29, 72). Finally, gene flow may not necessarily be

mediated across a terrestrial environment; genetic connectivity could also be maintained if

populations have been forced into contact by water level fluctuations in the groundwater system

of a region. The dynamic lacustrine history of the Cuenca Oriental (73), for instance, suggests

that isolation of some paedomorphic populations has been punctuated by broader aquatic

connections over short geological time scales, potentially facilitating gene flow. Collectively,

these mechanisms are likely to explain the maintenance of geographic genetic groups, now or in

the evolutionarily recent past, containing a range of life histories.

The genetic cohesion we detected across wide swaths of the TMVB is particularly

striking given other evidence for local adaptation in this region. This scenario is perhaps best

exemplified in the Cuenca Oriental (Fig. 2E), where some lakes contain paedomorphic

salamanders adapted to saline conditions, a very rare trait among amphibians (74). At one site,

the saline Lake Alchichica, the population is currently considered a distinct species (*A. taylori*)

(10). However, our results indicate that *A. taylori* is not genetically distinct from surrounding,

putatively non-saline-adapted populations, a result consistent with a previous microsatellite-

based study that found significant gene flow to and from *A. taylori* (29). While we cannot rule

out the potential for localized signatures of selection and adaptation in the *A. taylori* genome, our

results highlight that even populations adapted to aquatic conditions intolerable to most other

Ambystoma have not reached a level of isolation consistent with an independent evolutionary

trajectory (75).

Comparisons to previous work

This study included extensive geographic sampling of the tiger salamander complex,

which provided a broad spatial context to more fully understand patterns of genetic variation.

However, it is important to consider the possibility that our data are simply not sensitive enough

to detect genetic differentiation associated with lineage divergence on an extremely recent time

scale. The common ancestor of *A. californiense* and the remainder of the species complex dates

to approximately five million years (11, 37), and phylogenomic data indicate that speciation

across the remaining tiger salamander lineages occurred within the last one million years (76).

Such timing could make lineage boundaries difficult to detect (15). In this context, our genome-

wide markers, developed from transcriptomic resources (35), may not have an overall

substitution rate sufficient to detect genetic differentiation. We recommend that future research

uses a larger genomic dataset or faster-evolving loci with particular focus on the sampling of

obligate paedomorphs in this system. To this point, perhaps the strongest evidence for the genetic

divergence of obligately paedomorphic populations comes from past microsatellite-based work

indicating the genetic distinctiveness of *A. andersoni* and *A. mexicanum* from a select set of

populations across central Mexico (77) and the genetic distinctiveness of *A. taylori* from

neighboring populations in the Cuenca Oriental (29). Both of these studies recovered signatures

of gene flow, but identified greater overall population structure compared to this study. These

previous results could be due to numerous factors, including a faster microsatellite mutation rate,

methodological differences, and more limited locality sampling.

CONCLUSIONS

The extent to which populations within the tiger salamander complex exhibit phenotypic

plasticity in life history traits is remarkable and is believed to have played a role in the rapid

accumulation of lineages observed in the highlands of central Mexico (11, 61). While our results

suggest there is less species-level diversity in that region than previously recognized, there is

clearly more diversity in central Mexico compared to the US and Canada where there is (a) more

geographic space and (b) less life history variation within and between lineages. While we

cannot fully explain the greater diversity in Central Mexico, our results suggest that major

patterns of diversification are related to a complex history of geographic isolation and secondary

contact, in which life history strategy has played a less important role. We agree with previous

work (12) that the complex geological history of the TMVB, including montane uplift and

fluctuating drainage connectivity since the Miocene, has been the cornerstone of the evolutionary

history of this species complex, and that the influences of geographic isolation and

paedomorphosis may work synergistically to lead to the establishment of isolated populations.

Lingering questions notwithstanding, this large-scale genetic and geographic study has

established a framework for understanding the evolutionary history of the *Ambystoma tigrinum*

species complex. The results presented here will facilitate comparative studies of the axolotl and

its allies, provide direction for conservation prioritization and management, and strengthen the

use of the tiger salamander species complex as a model system in biology.

MATERIALS & METHODS

Geographic sampling

We generated data from a total of 254 individuals sampled from across the range of the

A. tigrinum complex (Fig. S1; Table S1). These individuals were combined with 93 individuals

sampled in O'Neill et al. (35) to produce a data set comprising 347 individuals. To represent the

large geographic range of this species complex, we chose to sample a large number of localities

(188) with limited numbers of individuals sampled per locality (mean = 1.8, min. = 1, max. = 9).

This sampling included a total of 166 individuals from the US, 2 individuals from Canada, and

178 individuals sampled from Mexico. Individuals of *A. californiense* were sampled from

localities with limited to no impacts of introgression from invasive western tiger salamanders

(52). Additional outgroup data were generated for two species outside of the *A. tigrinum*

complex: *A. opacum* and *A. texanum*. Full details regarding the generation of these data can be

found in the Supplementary Materials.

Data collection and sequencing

We generated DNA sequence data from a panel of 95 nuclear loci developed specifically

for the tiger salamander species complex. A more complete description of marker development

can be found in O'Neill et al. (35). Genomic DNA was extracted using a DNeasy Blood and

Tissue kit (Qiagen). For a small number of DNA extractions, we increased DNA quantities using

a Repli-g whole genome amplification kit (Qiagen). We used an initial round of PCR in 96-well

plate format to amplify all loci from an individual, followed by a smaller second round of PCR to

amplify loci that did not amplify in the initial PCR. See O'Neill et al. (35) for the details of PCR

conditions and primer sequences.

PCR products from all loci were pooled for each individual in roughly equal

concentrations based on the intensity of amplification as visualized on an agarose gel. Indexed

Illumina sequencing libraries were generated for each individual using an Illumina Nextera XT

DNA Library Preparation Kit (Illumina). Subsequent to library preparation, indexed libraries

were quantified using a Qubit Fluorometer, pooled in equimolar concentrations, and checked on

an Agilent Bioanalyzer to assure proper fragmentation.

Sequencing was performed in four rounds using an Illumina MiSeq. We performed an

initial round using a total of seven individuals to test the compatibility of the Illumina Nextera

XT library kit with our PCR amplicons. Three subsequent rounds of library preparation and

sequencing were performed on sets of 96 individuals each. Note that some individuals were

sequenced in multiple rounds due to initially low read counts. All sequencing was performed

with paired-end 150 bp reads. Overall, we generated a total of 29,426,894 PE reads across all

newly sequenced individuals, with an average of 309,757 PE reads, 1,148X coverage, and 7.6%

missing loci per individual (Table S2).

Bioinformatics and dataset generation

All sequence reads were processed using a newly developed bioinformatic pipeline

written for this project (https://doi.org/10.5281/zenodo.3585970) that produces multiple

sequence alignments for individual loci and genome-wide SNP matrices sampled from variable

sites. This pipeline was developed using the Snakemake workflow management system (78),

linking together multiple software tools to take sequence data from raw reads to phased sequence

alignments for each locus. Briefly, as used in this study, demultiplexed paired-end Illumina fastq

files were used as input, with separate forward and reverse read files for each individual.

Sequence data were trimmed and filtered in Trimmomatic (79) using a sliding window of 4 base-

pairs and a minimum average quality score of 15. Filtered sequence reads were then aligned to

reference sequences from the O'Neill et al. (2013) dataset (35); specifically, we used the clean

sequences of an *A. ordinarium* sample that had high coverage and low amounts of missing data.

The resulting aligned contigs were processed using SAMtools (80) to filter and prepare data for

FreeBayes (81), which was used to call variable sites. Variants were filtered with VCFTools (82)

by removing indels and setting a quality threshold of phred score > 20 and a minimum read

depth of 30. The program WhatsHap (83) was used to perform read-based phasing of the data for

each locus x individual contig. Finally, phased haplotypes from each individual (two copies,

regardless of homo- or heterozygosity) were combined into an alignment of all individuals using

MAFFT with the default auto parameter (84). We generated fasta files of SNPs using the SNP-

sites program (85) and created a SNP genotype matrix by sampling variable sites from a

concatenated sequence alignment of all loci. Through our informatic processing and manual

inspections of the data, we identified three loci (E12G1, E6A11, and E7G8) as potential paralogs

based on high alignment error and high levels of heterozygosity for all individuals. These three

loci were excluded from all analyses.

Across the remaining 92 loci, alignment lengths ranged from 123 to 630 bp (avg. = 270

bp) with a total concatenated alignment of 24,788 bp. For the full dataset, including *A. texanum*

and *A. opacum* outgroups, single-locus alignments contained an average of 55 variable sites

(min. = 15, max. = 121) and an average of 41 parsimony informative sites (min. = 11, max. =

102). Population genetic analyses restricted to the *A. tigrinum* complex including *A.*

californiense—which were further filtered for non-biallelic SNPs and a minor allele count ≥ 3—

contained a total of 2,360 SNPs.

Population structure and lineage discovery

We developed hypotheses of population-level lineages across the range of the *A. tigrinum*

complex using a bottom-up approach, starting with the identification of major geographic

patterns of differentiation, and then performing a recursive set of analyses on more restricted sets

of individuals, as identified in the previous analytical step. In our initial round of analyses we

used two non-parametric methods: principal components analysis (PCA) and discriminant

analysis of principal components (DAPC) (36). While both analyses provide a multivariate

summary of genetic data, DAPC is also used to assess the fit of data to varying numbers of

population clusters. These analyses were applied to our full genotypic dataset including *A.*

californiense and all remaining individuals from the *A. tigrinum* complex. The PCA was

calculated using the function 'prcomp' in the R package stats (86), while the DAPC was

calculated using the package adegenet (87). The optimal number of principal components to

retain for DAPC was identified using cross-validation via the xvalDapc function with default

parameter values. DAPC was performed without prior assignment of individuals to groups and

an exploratory approach was used to identify patterns of differentiation in ordination space

across a range of cluster levels ($K = 1$-20). We used two metrics to identify the best estimate of

the primary splits in our data. First, we used the Bayesian Information Criterion (BIC) calculated

in the DAPC analysis to assess the fit of the data to different levels of K. We note that the level

of K with the absolute lowest BIC may not be a better explanation of the data than a K with a

slightly higher BIC (88); therefore, we applied this measure for general guidance on a range of K

that may describe the data well. We paired this assessment with visualizations of the first and

second principal components (Fig. S2, S4) and DAPC ordination plots to identify the level of K

at which similar clustering patterns could be observed with minimal change at successively

higher levels of K. DAPC of the complete tiger salamander species complex identified a

consistent pattern beginning at $K = 5$ for high differentiation of all *A. californiense* samples (Fig.

S3). Further DAPC analysis with *A. californiense* removed identified a consistent pattern

beginning at $K = 5$ for differentiation between clusters of populations from northern and central

Mexico, and three clusters of U.S. populations (two from the Western U.S. and one from the

Eastern U.S.; Fig. S5).

Using the clusters identified in the DAPC analysis of the total data set, we then used both

DAPC and the program STRUCTURE v.2.3.4 (89) to analyze subsequent data sets comprising

smaller numbers of individuals. Recursive rounds of DAPC analyses were performed as

described above and were stopped when BIC scores showed little improvement (ΔBIC < 2) at

values of K > 1. STRUCTURE analyses were performed using an admixture model and a total of

500,000 generations following a burn-in of 100,000 generations. Analyses were performed for K

= 1-10 with 16 replicate analyses. We identified a "best" value of K using two approaches. First,

we calculated ΔK using the Evanno method (91) via the CLUMPAK web tool (90). A limitation

of the Evanno method is that it cannot identify the best value of $K = 1$ (91), thus, we also visually

inspected individual group assignments and concluded a value of $K = 1$ if the corresponding

DAPC cluster showed little improvement (ΔBIC < 2) at values of K > 1, or when visualization

showed extensive admixture without distinct clustering.

Tests of migration and population structure

We used the coalescent-based program Migrate-N v.3.2 (92) to explicitly test for

population structure and gene flow in each obligately paedomorphic species of *Ambystoma* (*A.*

andersoni, A. dumerilii, A. taylori, and *A. mexicanum,* plus the "often neotenic" species *A.*

lermaense). For each model we evaluated (Fig. 4A), we treated the obligately paedomorphic

species as one population, and the facultatively paedomorphic individuals from the same genetic

cluster (CM2-CM4) as the second population. The best-fitting model was determined via Bayes

factors (BF), which were calculated using Bézier-corrected marginal likelihoods (Table S2); the

highest-ranking models have a BF = 0. Each Migrate-N analyses ran for 50 million steps,

recording every 10 steps, with a burn-in of five million and the default heating scheme. Suitable

upper bounds for priors on population size (Θ) and migration rate (M) were determined from an

initial test run of 10 million steps for each analysis. Median values and 95% confidence intervals

of Θ and *M* are reported (Fig. 4B).

Phylogenetic reconstruction

As our dataset included a large number of admixed individuals, we estimated

phylogenetic networks to visualize reticulation events. We used SplitsTree v. 4.14.8 (40, 93) to

generate four networks: one with all tiger salamander individuals, and one each for the U.S.,

central Mexico, and northern Mexico subgroups. Networks were constructed using uncorrected

p-distances and the NeighborNet algorithm (94).

We also used three different analytical approaches to place hypothesized population

lineages in a phylogenetic framework. For all analyses we used a reduced data set containing the

concatenated data for 2-7 representative individuals from each population genetic cluster, which

limited computation time and avoided violating the coalescent-model assumption of zero gene

flow. We first inferred the phylogeny using Bayesian Inference in BEAST v.1.8.3 (42). Analyses

were run for 5 million generations, sampling every 1,000 generations after the first 500,000

generations were removed as burn-in. Run convergence was assessed with Tracer v.1.6.0 (95).

Next, we inferred a maximum-likelihood phylogeny using RAxML v.8 (43). Node support was

assessed using a rapid bootstrap analysis with 1,000 replicates, which was summarized as a 95%

rule consensus tree using the program SumTrees in the DendroPy python library (96). For both

BEAST and RAxML, PartitionFinder (97) was used to identify the number of preferred gene

partitions and their substitution models, and analyses were performed on the CIPRES Science

Gateway server (98). Finally, we inferred phylogenetic relationships using SVDquartets (41)

implemented in PAUP* version 4.0a164 (99), sampling all possible quartets and assessing node

support with 1,000 bootstrap replicates. For all phylogenetic analyses, trees were visualized

using FigTree v.1.4.2 (100).

DATA AVAILABILITY

Supplementary figures and tables are provided with the online version of this manuscript.

Input files for all population genetic and phylogenetic analyses are available via figshare

(https://figshare.com/projects/Life_history_strategy_does_not_reflect_genetic_differentiation_in

_the_tiger_salamander_species_complex/74115). Sequence data are available on the NCBI

Sequence Read Archive (BioProject accession PRJNA594660).

ACKNOWLEDGEMENTS

This project was supported by grants from the National Science Foundation (DEB-

0949532, DEB-1355000, and a Doctoral Dissertation Improvement Grant awarded to Justin

Kratovil) and by a University Research Fellowship awarded to Kathryn Everson from the

University of Kentucky. We thank numerous individuals for contributing samples to this project,

including Chris Beachy, Jim Bogart, Ken Carbale, Sheri Church, Jeff LeClere, Carol Hall, Paul

Moler, Nancy Staub, and Ken Wray. We are grateful to Jose Bocanegra, Ben Browning,

Mackenzie Humphrey, Mary Virginia Gibbs, Ricky Grewelle, Alan Lemmon, Emily Moriarty

Lemmon, Deborah Lu, Stephanie Mitchell, Alex Noble, Tolu Odukoya, Ben Tuttle, and Josh

Williams for their assistance with data collection and analyses.

REFERENCES

1. S. R. Palumbi, Marine speciation on a small planet. *Trends Ecol. Evol.* **7**, 114–118 (1992).

2. S. Mopper, S. Y. Strauss, *Genetic Structure and Local Adaptation in Natural Insect Populations: Effects of Ecology, Life History, and Behavior* (Springer US, 1998).

3. S. J. Gould, *Ontogeny and Phylogeny* (Harvard University Press, 1977).

4. D. B. Wake, Paedomorphosis. *J. Herpetol.* **14**, 80 (1980).

5. H. H. Whiteman, Evolution of facultative paedomorphosis in salamanders. *Q. Rev. Biol.* **69**, 205–221 (1994).

6. R. B. Page, *et al.*, Effect of thyroid hormone concentration on the transcriptional response underlying induced metamorphosis in the Mexican axolotl (Ambystoma). *BMC Genomics* **9**, 78 (2008).

7. S. R. Voss, Genetic Basis of Paedomorphosis in the Axolotl, Ambystoma mexicanum: A Test of the Single-Gene Hypothesis. *J. Hered.* **86**, 441–447 (1995).

8. S. R. Voss, H. B. Shaffer, Adaptive evolution via a major gene effect: paedomorphosis in the Mexican axolotl. *Proc. Natl. Acad. Sci. USA* **94**, 14185–9 (1997).

9. E. H. Taylor, New salamanders from Mexico with a discussion of certain known forms. *Univ. Kansas Sci. Bull.* **26**, 407–439 (1939).

10. R. Brandon, E. Maruska, W. Rumph, A new species of neotenic *Ambystoma* (Amphibia, Caudata) endemic to Laguna Alchichica, Puebla, Mexico. *Bull. South. Calif. Acad. Sci.* **80**, 112–125 (1981).

11. H. B. Shaffer, M. L. McKnight, The polytypic species revisited: Genetic differentiation and molecular phylogenetics of the tiger salamander *Ambystoma tigrinum* (Amphibia: Caudata) complex. *Evolution* **50**, 417–433 (1996).

514   12.   H. B. Shaffer, Evolution in a paedomorphic lineage. I. An electrophoretic analysis of the

515   Mexican ambystomatid salamanders. *Evolution* **38**, 1194–1206 (1984).

516   13.   H. B. Shaffer, Phylogenetics of model organisms: The laboratory axolotl, *Ambystoma*

517   *Mexicanum*. *Syst*. *Biol*. **42**, 508–522 (1993).

518   14.   H. B. Shaffer, Evolution in a paedomorphic lineage. II. Allometry and form in the

519   Mexican ambystomatid salamanders. *Evolution* **38**, 1207–1218 (1984).

520   15.   H. B. Shaffer, R. C. Thomson, Delimiting Species in Recent Radiations. *Syst*. *Biol*. **56**,

521   896–906 (2007).

522   16.   P. R. Grant, Speciation and the adaptive radiation of Darwin's finches. *Am*. *Sci*. **69**, 653–

523   663 (1981).

524   17.   O. Seehausen, Explosive speciation rates and unusual species richness in haplochromine

525   cichlid fishes: Effects of sexual selection. *Adv*. *Ecol*. *Res*. **31**, 237–274 (2000).

526   18.   M. Fegan, P. Prior, "How complex is the *Ralstonia solanacearum* species complex" in

527   *Bacterial Wilt Disease and the Ralstonia Solanacearum Species Complex*, C. Allen, P.

528   Prior, A. C. Hayward, Eds. (American Phytopathological Society Press, 2005), pp. 449–

529   461.

530   19.   J. Mallet, Hybrid speciation. *Nature* **446**, 279–283 (2007).

531   20.   M. C. Fontaine, *et al*., Extensive introgression in a malaria vector species complex

532   revealed by phylogenomics. *Science* **347**, 1258524 (2015).

533   21.   J. Geml, G. A. Laursen, K. O'Neill, H. C. Nusbaum, D. L. Taylor, Beringian origins and

534   cryptic speciation events in the fly agaric (*Amanita muscaria*). *Mol*. *Ecol*. **15**, 225–239

535   (2005).

22.   K. S. Jones, D. W. Weisrock, Genomic data reject the hypothesis of sympatric ecological

speciation in a clade of *Desmognathus* salamanders. *Evolution* **72**, 2378–2393 (2018).

23.   R. Newman, Adaptive plasticity in amphibian metamorphosis. *Bioscience* **42**, 671–678

(1992).

24.   H. H. Whiteman, S. A. Wissinger, W. S. Brown, Growth and foraging consequences of

facultative paedomorphosis in the tiger salamander, *Ambystoma tigrinum nebulosum*.

Evol. Ecol. **10**, 433–446 (1996).

25.   P. J. Fernandez, J. P. Collins, Effect of environment and ontogeny on color pattern

variation in Arizona tiger salamanders (*Ambystoma tigrinum nebulosum* Hallowell).

Copeia **1988**, 928 (1988).

26.   S. P. D. Riley, H. B. Shaffer, S. R. Voss, B. M. Fitzpatrick, Hybridization between a rare,

native tiger salamander (*Ambystoma californiense*) and its introduced congener. *Ecol.*

Appl. **13**, 1263–1275 (2003).

27.   D. W. Weisrock, H. B. Shaffer, B. L. Storz, S. R. Storz, S. R. Voss, Multiple nuclear gene

sequences identify phylogenetic species boundaries in the rapidly radiating clade of

Mexican ambystomatid salamanders. *Mol. Ecol.* **15**, 2489–2503 (2006).

28.   G. Parra-Olea, *et al.*, Conservation genetics of threatened Mexican axolotls (*Ambystoma*).

Anim. Conserv. **15**, 61–72 (2012).

29.   R. Percino-Daniel, E. Recuero, E. Vázquez-Domínguez, K. R. Zamudio, G. Parra-Olea,

All grown-up and nowhere to go: Paedomorphosis and local adaptation in *Ambystoma*

salamanders in the Cuenca Oriental of Mexico. *Biol. J. Linn. Soc.* **118**, 582–597 (2016).

30.   R. A. Brandon, Hybridization between the Mexican salamanders *Ambystoma dumerilii*

and *Ambystoma mexicanum* under laboratory conditions. *Herpetologica* **28**, 199–207

(1972).

31.   R. A. Brandon, Interspecific hybridization among Mexican and United States salamanders

of the genus *Ambystoma* under laboratory conditions. *Herpetologica* **33**, 133–152 (1977).

32.   S. Krebs, R. Brandon, A new species of salamander (family Ambystomatidae) from

Michoacan, Mexico. *Herpetologica* **40**, 238–245 (1984).

33.   J. B. Armstrong, G. M. Malacinski, *Developmental Biology of the Axolotl* (Oxford

University Press, 1989).

34.   L. L. Bailey, T. R. Simons, K. H. Pollock, Estimating site occupancy and species

detection probability parameters for terrestrial salamanders. *Ecol. Appl.* **14**, 692–702

(2004).

35.   E. M. O'Neill, *et al.*, Parallel tagged amplicon sequencing reveals major lineages and

phylogenetic structure in the North American tiger salamander (*Ambystoma tigrinum*)

species complex. *Mol. Ecol.* **22**, 111–129 (2013).

36.   T. Jombart, S. Devillard, F. Balloux, Discriminant analysis of principal components: A

new method for the analysis of genetically structured populations. *BMC Genet.* **11**, 94

(2010).

37.   H. B. Shaffer, G. B. Pauly, J. C. Oliver, P. C. Trenham, The molecular phylogenetics of

endangerment: Cryptic variation and historical phylogeography of the California tiger

salamander, *Ambystoma californiense*. *Mol. Ecol.* **13**, 3033–3049 (2004).

38.   J. W. Petranka, *Salamanders of the United States and Canada* (Smithsonian Books, 2010).

39. ,   AmphibiaWeb. *Univ. California, Berkeley, CA* (2020) (May 11, 2020).

40.   D. H. Huson, SplitsTree: Analyzing and visualizing evolutionary data. *Bioinformatics* **14**,

68–73 (1998).

41.   J. Chifman, L. Kubatko, Quartet inference from SNP data under the coalescent model.

Bioinformatics **30**, 3317–3324 (2014).

42.   A. J. Drummond, A. Rambaut, BEAST: Bayesian evolutionary analysis by sampling trees.

BMC Evol. Biol. **7**, 214 (2007).

43.   A. Stamatakis, RAxML version 8: A tool for phylogenetic analysis and post-analysis of

large phylogenies. *Bioinformatics* **30**, 1312–1313 (2014).

44.   A. A. Dugès, *Description d'un axolotl des montagnes de las cruces (*Amblystoma *[sic]*

altamirani, *A. Dugès)* (Imprimerie du Ministère de "Fomento," 1895).

45.   E. H. Taylor, Herpetological novelties from Mexico. *Univ. Kansas Sci. Bull.* **29**, 343–358

(1943).

46.   E. Dunn, A new genus of salamanders from Mexico. *Proc. New Engl. Zool. Club* **10**, 85–

86 (1928).

47.   E. H. Taylor, A new ambystomid salamander from the Plateau Region of Mexico. *Univ.*

Kansas Sci. Bull. **30**, 57–61 (1944).

48.   A. A. Dugès, Una nueva especie de ajolote de la Laguna de Patzcuaro. *La Nat.* **1**, 241–244

(1870).

49.   J. Dixon, A new species of salamander of the genus Ambystoma from Jalisco, Mexico.

Copeia **1**, 99–101 (1963).

50.   G. Shaw, F. Nodder, *The Naturalist's Miscellany; or Coloured Figures of Natural Objects*

Drawn and Described Immediately from Nature, 9th Ed. (Nodder & Co., 1798).

51.   A. A. Dugès, Erpetología del valle de México. *La Nat.* **2**, 97–146 (1888).

52.   J. R. Johnson, R. C. Thomson, S. J. Micheletti, H. B. Shaffer, The origin of tiger

salamander (*Ambystoma tigrinum*) populations in California, Oregon, and Nevada:

Introductions or relicts? *Conserv. Genet.* **12**, 355–370 (2011).

53. E. H. Taylor, Two new ambystomid salamanders from Chihuahua. *Copeia* **3**, 143–146

(1941).

54. R. Webb, Observations on tiger salamanders (*Ambystoma tigrinum* complex, family

Ambystomatidae) in Mexico with description of a new species. *Bull. Maryl. Herpetol.*

Soc. **40**, 122–143 (2004).

55. R. A. Brandon, Spontaneous and induced metamorphosis of *Ambystoma dumerilii*

(Dugès), a paedogenetic Mexican salamander, under laboratory conditions. *Herpetologica*

32, 429–438 (1976).

56. H. M. Wilbur, J. P. Collins, Ecological aspects of amphibian metamorphosis: Nonnormal

distributions of competitive ability reflect selection for facultative metamorphosis. *Science*

182, 1305–1314 (1973).

57. R. M. Bonett, M. A. Steffen, S. M. Lambert, J. J. Wiens, P. T. Chippindale, Evolution of

paedomorphosis in plethodontid salamanders: Ecological correlates and re-evolution of

metamorphosis. *Evolution* **68**, 466–482 (2014).

58. C. R. Marshall, E. C. Raff, R. A. Raff, Dollo's law and the death and resurrection of

genes. *Proc. Natl. Acad. Sci. USA* **91**, 12283–12287 (1994).

59. M. J. West-Eberhard, Phenotypic Plasticity and the Origins of Diversity. *Annu. Rev. Ecol.*

Syst. **20**, 249–278 (1989).

60. M. J. West-Eberhard, Alternative adaptations, speciation, and phylogeny (A Review).

Proc. Natl. Acad. Sci. USA **83**, 1388–92 (1986).

61. B. M. Fitzpatrick, Underappreciated consequences of phenotypic plasticity for ecological

speciation. *Int. J. Ecol.* **2012**, 1–12 (2012).

62. M. Denoël, P. Poncin, J.-C. Ruwet, Sexual compatibility between two heterochronic

morphs in the Alpine newt, *Triturus alpestris*. *Anim. Behav.* **62**, 559–566 (2001).

63.   J. D. Krenz, P. A. Verrell, Integrity in the midst of sympatry: Does sexual incompatibility

facilitate the coexistence of metamorphic and paedomorphic mole salamanders

(*Ambystoma talpoideum*)? *J. Zool.* **258**, S0952836902001589 (2002).

64.   H. H. Whiteman, J. D. Krenz, R. D. Semlitsch, Intermorph breeding and the potential for

reproductive isolation in polymorphic mole salamanders (*Ambystoma talpoideum*). *Behav.*

Ecol. Sociobiol. **60**, 52–61 (2006).

65.   H. Whiteman, J. Gutrich, R. Moorman, Courtship behavior in a polymorphic population of

the tiger salamander, *Ambystoma tigrinum nebulosum*. *J. Herpetol.* **33**, 348–351 (1999).

66.   R. . Howard, R. . Moorman, H. . Whiteman, Differential effects of mate competition and

mate choice on eastern tiger salamanders. *Anim. Behav.* **53**, 1345–1356 (1997).

67.   D. H. Bos, R. N. Williams, D. Gopurenko, Z. Bulut, J. A. DeWoody, Condition-dependent

mate choice and a reproductive disadvantage for MHC-divergent male tiger salamanders.

Mol. Ecol. **18**, 3307–3315 (2009).

68.   C. E. Nelson, R. R. Humphrey, Artificial interspecific hybridization among *Ambystoma*.

Herpetologica **28**, 27–32 (1972).

69.   J. M. Velasco, Anotaciones y observaciones al trabajo del Sr. D. A. Weismann sobre la

transformación del ajolote Mexicano en Amblistoma. *La Nat.* **5**, 58–84 (1880).

70.   H. M. Smith, The Mexican axolotl: Some misconceptions and problems. *Bioscience* **19**,

593–615 (1969).

71.   A. G. E. Mathiron, J.-P. Lena, S. Baouch, M. Denoël, The 'male escape hypothesis': Sex-

biased metamorphosis in response to climatic drivers in a facultatively paedomorphic

amphibian. *Proc. R. Soc. B Biol. Sci.* **284**, 20170176 (2017).

72.   C. L. Morjan, L. H. Rieseberg, How species evolve collectively: Implications of gene flow

653   and selection for the spread of advantageous alleles. *Mol. Ecol.* **13**, 1341–1356 (2004).

654   73.   J. L. Tamayo, R. C. West, "The hydrography of middle America" in *Handbook of Middle*

655   *American Indians*, R. Wauchope, R. C. West, Eds. (University of Texas Press, 1964), pp.

656   84–121.

657   74.   G. R. Hopkins, E. D. Brodie, Occurrence of amphibians in saline habitats: A review and

658   evolutionary perspective. *Herpetol. Monogr.* **29**, 1–27 (2015).

659   75.   K. De Queiroz, Species concepts and species delimitation. *Syst. Biol.* **56**, 879–886 (2007).

660   76.   P. M. Hime, *et al.*, Phylogenomics reveals ancient gene tree discordance in the amphibian

661   tree of life. *Syst. Biol.* (2020) https:/doi.org/10.1093/sysbio/syaa034 (May 12, 2020).

662   77.   G. Parra-Olea, *et al.*, Conservation genetics of threatened Mexican axolotls (*Ambystoma*).

663   *Anim. Conserv.* **15**, 61–72 (2012).

664   78.   J. Koster, S. Rahmann, Snakemake: A scalable bioinformatics workflow engine.

665   *Bioinformatics* **28**, 2520–2522 (2012).

666   79.   A. M. Bolger, M. Lohse, B. Usadel, Trimmomatic: A flexible trimmer for Illumina

667   sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

668   80.   H. Li, A statistical framework for SNP calling, mutation discovery, association mapping

669   and population genetical parameter estimation from sequencing data. *Bioinformatics* **27**,

670   2987–2993 (2011).

671   81.   E. Garrison, G. Marth, Haplotype-based variant detection from short-read sequencing.

672   *arXiv* **1207.3907** (2012).

673   82.   P. Danecek, *et al.*, The variant call format and VCFtools. *Bioinformatics* **27**, 2156–2158

674   (2011).

675   83.   M. Martin, *et al*., WhatsHap: Fast and accurate read-based phasing. *bioRxiv* **10.1101/08**

676   (2016).

677   84.   K. Katoh, D. M. Standley, MAFFT multiple sequence alignment software version 7:

678   Improvements in performance and usability. *Mol. Biol. Evol.* **30**, 772–780 (2013).

679   85.   A. J. Page, *et al*., SNP-sites: Rapid efficient extraction of SNPs from multi-FASTA

680   alignments. *Microb. genomics* **2**, e000056 (2016).

681   86.   R. C. Team, R: a language and environment for statistical computing (2012).

682   87.   T. Jombart, adegenet: A R package for the multivariate analysis of genetic markers.

683   *Bioinformatics* **24**, 1403–1405 (2008).

684   88.   T. Jombart, C. Collins, A tutorial for discriminant analysis of principal components

685   (DAPC) using adegenet 2.0.0. *adegenet.r-forge.r-project.org* (2015) (May 30, 2019).

686   89.   J. K. J. Pritchard, M. Stephens, P. Donnelly, Inference of population structure using

687   multilocus genotype data. *Genetics* **155**, 945–959 (2000).

688   90.   N. M. Kopelman, J. Mayzel, M. Jakobsson, N. A. Rosenberg, I. Mayrose, Clumpak: A

689   program for identifying clustering modes and packaging population structure inferences

690   across K. *Mol. Ecol. Resour.* **15**, 1179–1191 (2015).

691   91.   G. Evanno, S. Regnaut, J. Goudet, Detecting the number of clusters of individuals using

692   the software structure: A simulation study. *Mol. Ecol.* **14**, 2611–2620 (2005).

693   92.   P. Beerli, M. Palczewski, Unified framework to evaluate panmixia and migration direction

694   among multiple sampling locations. *Genetics* **185**, 313–326 (2010).

695   93.   D. H. Huson, D. Bryant, Application of phylogenetic networks in evolutionary studies.

696   *Mol. Biol. Evol.* **23**, 254–267 (2006).

94.   D. Bryant, V. Moulton, Neighbor-Net: An agglomerative method for the construction of

phylogenetic networks. *Mol. Biol. Evol.* **21**, 255–265 (2003).

95.   A. Rambaut, A. Drummond, D. Xie, G. Baele, M. Suchard, Tracer v1.7 (2018).

96.   J. Sukumaran, M. Holder, DendroPy: A Python library for phylogenetic computing.

Bioinformatics **26**, 1569–1571 (2010).

97.   R. Lanfear, B. Calcott, S. Y. W. Ho, S. Guindon, PartitionFinder: Combined selection of

partitioning schemes and substitution models for phylogenetic analyses. *Mol Biol Evol* **29**,

1695–1701 (2012).

98.   M. A. Miller, W. Pfeiffer, T. Schwartz, Creating the CIPRES Science Gateway for

inference of large phylogenetic trees in *2010 Gateway Computing Environments*

Workshop (GCE), (IEEE, 2010), pp. 1–8.

99.   D. Swofford, PAUP 4.0 b10: Phylogenetic analysis using parsimony (2002) (December

21, 2013).

100. A. Rambaut, FigTree, a graphical viewer of phylogenetic trees (2007) (May 8, 2014).