Effect of sunlight on the efficacy of commercial antibiotics used in agriculture

Sebastian Khan, Amanda Osborn, and Prahathees J. Eswara[*]

Department of Cell Biology, Microbiology, and Molecular Biology, University of South Florida,

Tampa, Florida, USA

*Address correspondence to Prahathees J. Eswara, eswara@usf.edu.

Keywords: Antibiotic stewardship, antibiotic resistance, Liberibacter; huanglongbing, Erwinia,

fire blight, aquaculture, streptomycin; oxytetracycline.

Running title: Effect of sunlight on commercial antibiotics

ABSTRACT

Antibiotic stewardship is of paramount importance to limit the emergence of antibiotic-

resistant bacteria in not only hospital settings, but also in animal husbandry,

aquaculture, and agricultural sectors. Currently, large quantities of antibiotics are

applied to treat agricultural diseases like citrus greening disease (CGD). The two

commonly used antibiotics approved for this purpose are streptomycin and

oxytetracycline. Although investigations are ongoing to understand how efficient this

process is to control the spread of CGD, to our knowledge, there have been no studies

that evaluate the effect of environmental factors such as sunlight on the efficacy of the

above-mentioned antibiotics. We conducted a simple disc-diffusion assay to study the

efficacy of streptomycin and oxytetracycline after exposure to sunlight for 7- or 14-day

periods using *Escherichia coli* and *Bacillus subtilis* as the representative strains of

Gram-negative and Gram-positive organisms respectively. Freshly prepared discs and

discs stored in the dark for 7 or 14 days served as our controls. We show that the

antibiotic potential of oxytetracycline exposed to sunlight dramatically decreases over

the course of 14 days against both *E. coli* and *B. subtilis*. However, the effectiveness of

streptomycin was only moderately impacted by sunlight. It is important to note that

antibiotics that last longer in the environment may play a deleterious role in the rise and

spread of antibiotic-resistant bacteria. Further studies are needed to substantively

analyze the safety and efficacy of antibiotics used for broader environmental

applications.

IMPORTANCE

Although antibiotics have been used for agricultural purposes for decades, due to the

rapid rise in antibiotic resistance this usage needs to be revisited. Questions remain on

the appropriate mode of application of antibiotics and the actual benefits of using

antibiotics for treating the infections caused by plant pathogens, especially for the ones

that are intracellular in nature. Here we show that the two commonly used commercial

antibiotics, oxytetracycline and streptomycin, lose their efficacy at different rates in the

presence of sunlight. While the former loses its potency within days the latter remains

active for many days. Thus, oxytetracycline may not be active long enough to produce

desired effect and streptomycin may persist in the environment and as a side effect due

to its selective pressure, may force the rise of streptomycin-resistant pathogens.

INTRODUCTION

Antibiotic resistance-related mortalities are expected to exceed the other leading causes

of death such as cancer worldwide by 2050 [1]. Antibiotic stewardship is therefore

promoted in all sectors including human health, animal husbandry, and agriculture [2-4].

The World Health Organization and the United States Centers for Disease Control and

Prevention have recognized antimicrobial resistance as an enormous ongoing threat to

public health [5, 6]. Runoff of antibiotics in hospital waste water [7] and intentional use in

aquaculture [8], animal husbandry [9-11], and crop management [12] contribute to the

rise and spread of antibiotic resistant bacteria. In this context, alarm was raised recently

regarding the spraying of antibiotics in open fields as an infection control strategy to

stem the spread of bacterial disease in plants [13, 14]. Specifically, the strategy

approved by the United States Environmental Protection Agency [13, 15, 16] is to use

streptomycin and oxytetracycline to control the spread of citrus greening disease

(CGD), also known as huanglongbing (yellow dragon disease). CGD is a devastating

bacterial disease caused by *Candidatus Liberibacter asiaticus* (CLas) that is transmitted

between plants by certain psyllids, which are sap-feeding insects. CLas is a fastidious,

Gram-negative, intracellular plant pathogen that belongs to the phylum of α-

proteobacteria [17, 18]. Streptomycin and oxytetracycline are also used to treat

infections caused by another bacterial plant pathogen, *Erwinia amylovora*, which causes

fire blight in apples, pears, and other related species [19]. *E. amylovora* has dual growth

modes - an epiphytic mode that is readily accessible for external antibiotics and an

endophytic mode that is less accessible to external antibiotics [19]. In addition,

tetracycline antibiotics including oxytetracycline are used in animal husbandry [20] and

aquaculture [21]. Apart from the uses described above, data also suggests that

antibiotics may find their way into and possibly persist in different animal and plant

tissues [22-25], which could be an alternate pathway that can lead to the development of

antibiotic-resistant bacteria. Thus, a comprehensive knowledge of the fate of antibiotics

used in agriculture is urgently needed to hopefully curb the rise and spread of antibiotic

resistance.

Although the application of antibiotics to treat CGD inspired us to pursue this study, the

primary objective of this report is to investigate the effect of environmental factors,

specifically sunlight, on streptomycin and oxytetracycline. To this end, we conducted a

disc-diffusion assay with Gram-negative *Escherichia coli* and Gram-positive *Bacillus*

subtilis and monitored the zones of inhibition of antibiotic-containing discs that were

exposed to sunlight for a 7- or 14-day period. Discs that were kept in the dark for

equivalent duration or that were freshly prepared served as our controls. Based on our

results, we report that sunlight significantly impairs the efficacy of oxytetracycline, but

only moderately impacts streptomycin. While short-lived antibiotics may not be active

long enough for their intended purpose, stable antibiotics may apply constant selection

pressure and create an environment conducive for the emergence of antibiotic-resistant

strains [26]. Although this study (designed for undergraduate-level students [27]) is not

comprehensive, our data provides a window into the life span of commercial antibiotics

in nature that we hope highlights the need for further rigorous safety and efficacy

investigations for the environmental use of antibiotics.

RESULTS

Oxytetracycline loses its antibiotic potential in the presence of sunlight in the span of few days.

To monitor the effect of sunlight on the efficacy of oxytetracycline, we conducted a disc-diffusion assay. Briefly, we prepared multiple discs with oxytetracycline (50 µg) dissolved in water and placed the antibiotic-laden discs in either a natural outdoor setting with abundant sunlight to simulate agricultural use, or in a dark indoor cabinet for 7 or 14 days. In addition to the discs that were kept in the dark, we also used freshly prepared discs and vehicle (water) discs as controls. The discs were then placed, as shown in **Fig. 1**, on a pre-inoculated plate containing either a lawn of *E. coli* or *B. subtilis* cells. In all cases, as expected, the blank disc (N; negative control) and the freshly prepared discs (P; positive control) showed negligible and maximum zones of inhibition (ZOI), respectively (**Figs. 1A-D**). The discs that were kept in the dark (labeled "D") for the duration of 7 or 14 days appeared to produce similar ZOI as our positive control of approximately 9 mm for *E. coli* and 8 mm for *B. subtilis* (**Figs. 1EF**). This suggests that oxytetracycline maintains its efficiency in the dark at room temperature for at least the maximum duration of this experiment (14 days). Next, we quantified the ZOI of the discs that were exposed to sunlight (labeled "L") for either a 7- or 14-day period. We observed that the efficacy of oxytetracycline gradually and significantly decreased over time to almost similar to our negative control in both *E. coli* and *B. subtilis* and only retained less than 15% activity after 14 days (**Figs. 1A-F**). This implies that in the presence of sunlight, oxytetracycline loses its antibiotic potential in a matter of few days.

Moderate negative effects of sunlight on the efficacy of streptomycin.

A similar experimental setup to the one discussed above was adopted for studying the

effects of sunlight on streptomycin. As noted earlier, blank discs and freshly prepared

discs with streptomycin (200 μg) served as our negative and positive controls

respectively. As expected, the ZOI were unobservable for our blank discs and at a

maximum for our positive controls (**Figs. 1G-L**). Similar to oxytetracycline, streptomycin

is also able to maintain its efficacy when kept in darkness for the duration of our

experiment (**Figs. 1G-L**). However, unlike oxytetracycline, streptomycin appears to be

moderately resistant to sunlight. At the 7-day mark, based on the ZOI (**Figs. 1KL**), the

discs exposed to sunlight appear to have retained almost approximately 80% and 70%

of their activity in *E.coli* and *B. subtilis* respectively, when compared to that of our

positive control. Further measurable decrease to nearly 50% efficiency compared to our

positive control was noted subsequent to 14 days of sunlight exposure for *E. coli*.

However, the decrease in efficiency for *B. subtilis* at the 14-day time point was within

the standard error when compared to that of the 7-day time point (**Figs. 1HJKL**).

DISCUSSION

Rapid rise of antibiotic resistance in bacteria is a major concern worldwide with

enormous predicted fatalities. Antibiotics are now routinely used in clinics, animal

husbandry, and agriculture. Acknowledgement of the fact that the rise of antibiotic

resistance stemming from one of those settings could potentially render antibiotics

useless and lead to the formation of a multidisciplinary collaborative initiative under the

umbrella term One Health [2, 3]. Despite this, environmental antibiotic pollution is a

growing concern that requires urgent attention [28].

Some commercial antibiotics such as oxytetracycline and streptomycin are produced by

soil-dwelling *Streptomyces* spp. However, soil bacteria do not produce antibiotics at

levels comparable to commercial applications – which can occasionally be in the scale

of thousands of kilograms [13, 15, 16]. Also, the efficiency of superficial application of

antibiotics in limiting the growth of plant bacterial pathogens, including some that are

intracellular, is unclear. Recent studies have suggested injection of oxytetracycline

produces better results [19, 29]. The spread of antibiotic resistance has been

documented from agricultural use for antibiotics like tetracycline and streptomycin [30-

32]. It has been noted that antibiotic resistance genes are naturally found in the

environment [33, 34]. Therefore, application of consistent selection pressure by

excessive and frequent use of antibiotics may enrich the population of naturally resistant

organisms. However, at least in some instances under certain conditions, it was noted

that streptomycin use did not alter the composition of soil microbial communities

appreciably [35, 36].

Several reports on degradation kinetics and mechanisms of degradation of the

antibiotics that are discussed here are available [21, 37-44]. It has been reported that the

half-life of oxytetracycline at 25 °C is approximately 7 days, at 35 °C is 3 days and at 60

°C is 0.2 day, indicating a rapid temperature-dependent degradation of oxytetracycline,

as the half-life at 4 °C is 120 days [37]. According to the same study, the half-life due to

photolysis in the presence of sunlight is in the same order of magnitude. A similar

investigation exists evaluating the photostability and temperature stability of

streptomycin [44]. Briefly, the photodegradation of streptomycin is more modest than

oxytetracycline by nearly 10-fold. The half-life of streptomycin was determined to be

nearly 105, 42 and 30 days at 15 °C, 25 °C, and 40 °C respectively, implying a

decreased rate of degradation when compared to oxytetracycline. A description of the

possible degradation products of oxytetracycline and streptomycin are available [37, 44].

Our results showing a faster loss of efficacy for oxytetracycline than streptomycin upon

sunlight exposure are therefore in agreement with the reported degradation kinetics of

these antibiotics. To our knowledge, analysis such as the one we have conducted to

monitor the biological efficacy of antibiotics subsequent to exposure to environmental

elements are either lacking or not publicly available (as recognized by this article [14]).

Our experimental conditions simulate the agricultural use of antibiotics and our results

indicate that sunlight (heat and/or ultraviolet radiation) contributes to the degradation of

oxytetracycline and streptomycin. Although our report is limited in scope, we believe it

sheds light on the fate of antibiotics in the environment. Further studies to understand

the effects of antibiotics are needed to inform the public and appropriate regulatory

agencies [2-4].

MATERIALS AND METHODS

Strains used and general methods

The *B. subtilis* strain PY79 and the *E. coli* strain K-12 were incubated in 2 ml LB at 37 °C and grown until the culture OD_{600} reached 1.0 (exponential growth phase). A 100 µl aliquot of culture was then spread onto LB agar plates using sterile beads and set to dry completely prior to the placement of discs, see section below.

Disc-diffusion assay

UV sterilized Whatman filter paper discs (7 mm) were impregnated with 5 µl of a freshly made stock antibiotic solution of either 40 mg/ml streptomycin sulfate (MilliporeSigma) in sterile distilled water or 10 mg/ml oxytetracycline hydrochloride (Alfa Aesar) in sterile distilled water to reach a concentration of 200 µg for streptomycin and 50 µg for oxytetracycline in each disc, and then set to dry completely. The concentrations selected were based on the concentration range recommended for agricultural use [45], and after empirically ensuring similar initial zones of inhibition for both antibiotics in the strains tested. To mimic the use of agricultural antibiotics, the discs were then placed outdoors (during spring months in Tampa, FL, USA where the average daytime temperature ranged from 27 to 32 °C) in direct sunlight for 7 or 14 consecutive 24-h periods (days) in parafilm-sealed sterile Petri dishes. Discs that were kept indoors in a dark cabinet at room temperature for 7 or 14 days, freshly prepared discs made the day of testing, and 5 µl of sterile water were used as controls. Discs were then transferred and pressed onto the pre-inoculated LB agar plates and incubated overnight at 37 °C.

The zone of inhibition measurements were taken from the center of the disc to the edge

of the zone of inhibition, minus disc radius (3.5 mm).

Statistical analysis

GraphPad Prism Software (version 8.3.1) was used to analyze the data. All data

represent biological triplicate data with technical replicates. Graphs show mean values

and error bars represent standard deviation (SD).

ACKNOWLEDGEMENTS

We thank our lab members for comments on the manuscript and assistance with data

visualization. This work was funded by a start-up grant from USF (PE). A preprint of this

manuscript is available on bioRxiv [46].

AUTHOR CONTRIBUTIONS

The conception and design of the study (SK, PE), data acquisition (SK, AO), analysis

and/or interpretation of the data (SK, AO, PE), and writing of the manuscript (SK, PE).

FIGURE LEGEND

Figure 1. Oxytetracycline and streptomycin lose antibiotic potential in the presence of sunlight. Shown are representative disc-diffusion assay results for the effects of oxytetracycline **(A-D)** or streptomycin **(G-J)** on growth of either Gram-positive *B. subtilis* or Gram-negative *E. coli*. Quantification of the zones of inhibition in millimeters are plotted for each 7- or 14-day cohort of oxytetracycline **(E-F)** and streptomycin **(K-L).** Significance was determined using a one-way ANOVA with Tukey's multiple comparisons analysis. Error bars represent standard deviation (SD) of the mean from three biological replicates. N: negative control (discs prepared with sterile water), P: positive control (discs prepared the day of testing), L7 or L14: 7 or 14 days in sunlight, D7 or D14: 7 or 14 days in darkness. ****: $p<0.0001$, ***: $p<0.001$, **: $p<0.01$.

REFERENCES

1.   Editors PM. Antimicrobial Resistance: Is the World UNprepared? PLoS Med. 2016;13(9):e1002130. doi: 10.1371/journal.pmed.1002130. PubMed PMID: 27618631; PubMed Central PMCID: PMCPMC5019402.

2.   McEwen SA, Collignon PJ. Antimicrobial Resistance: a One Health Perspective. Microbiol Spectr. 2018;6(2). doi: 10.1128/microbiolspec.ARBA-0009-2017. PubMed PMID: 29600770.

3.   Hernando-Amado S, Coque TM, Baquero F, Martinez JL. Defining and combating antibiotic resistance from One Health and Global Health perspectives. Nat

Microbiol. 2019;4(9):1432-42. doi: 10.1038/s41564-019-0503-9. PubMed PMID:

31439928.

4. Thanner S, Drissner D, Walsh F. Antimicrobial Resistance in Agriculture. mBio.

2016;7(2):e02227-15. doi: 10.1128/mBio.02227-15. PubMed PMID: 27094336; PubMed

Central PMCID: PMCPMC4850276.

5. Toner E, Adalja A, Gronvall GK, Cicero A, Inglesby TV. Antimicrobial resistance

is a global health emergency. Health Secur. 2015;13(3):153-5. doi:

10.1089/hs.2014.0088. PubMed PMID: 26042858; PubMed Central PMCID:

PMCPMC4486712.

6. Kadri SS. Key Takeaways From the U.S. CDC's 2019 Antibiotic Resistance

Threats Report for Frontline Providers. Crit Care Med. 2020;48(7):939-45. doi:

10.1097/CCM.0000000000004371. PubMed PMID: 32282351; PubMed Central PMCID:

PMCPMC7176261.

7. Hocquet D, Muller A, Bertrand X. What happens in hospitals does not stay in

hospitals: antibiotic-resistant bacteria in hospital wastewater systems. J Hosp Infect.

2016;93(4):395-402. doi: 10.1016/j.jhin.2016.01.010. PubMed PMID: 26944903.

8. Cabello FC, Godfrey HP, Buschmann AH, Dolz HJ. Aquaculture as yet another

environmental gateway to the development and globalisation of antimicrobial resistance.

Lancet Infect Dis. 2016;16(7):e127-e33. doi: 10.1016/S1473-3099(16)00100-6. PubMed

PMID: 27083976.

9. Martin MJ, Thottathil SE, Newman TB. Antibiotics Overuse in Animal Agriculture:

A Call to Action for Health Care Providers. Am J Public Health. 2015;105(12):2409-10.

doi: 10.2105/AJPH.2015.302870. PubMed PMID: 26469675; PubMed Central PMCID:

PMCPMC4638249.

10.   Landers TF, Cohen B, Wittum TE, Larson EL. A review of antibiotic use in food

animals: perspective, policy, and potential. Public Health Rep. 2012;127(1):4-22. doi:

10.1177/003335491212700103. PubMed PMID: 22298919; PubMed Central PMCID:

PMCPMC3234384.

11.   Van Boeckel TP, Pires J, Silvester R, Zhao C, Song J, Criscuolo NG, et al.

Global trends in antimicrobial resistance in animals in low- and middle-income

countries. Science. 2019;365(6459). doi: 10.1126/science.aaw1944. PubMed PMID:

31604207.

12.   Sundin GW, Wang N. Antibiotic Resistance in Plant-Pathogenic Bacteria. Annu

Rev Phytopathol. 2018;56:161-80. doi: 10.1146/annurev-phyto-080417-045946.

PubMed PMID: 29856934.

13.   McKenna M. Antibiotics set to flood Florida's troubled orange orchards. Nature.

2019;567(7748):302-3. doi: 10.1038/d41586-019-00878-4. PubMed PMID: 30890811.

14.   Editorial. Spraying diseased citrus orchards with antibiotics could backfire.

Nature. 2019;567(7748):283. doi: 10.1038/d41586-019-00875-7. PubMed PMID:

30890810.

15.   Collins S, Kough JL. Review of GeoLogic/Agrosource's Analysis of

Oxytetracycline's Safety with Regard to Its Microbiological Effect on Bacteria of Human

Health Concern (FDA/CVM Guidance to Industry #152) for Registration on Citrus Crop

Group 10–10 [Memorandum] Washington, D.C. US Environmental Protection Agency.

2017;https://www.regulations.gov/document?D=EPA-HQ-OPP-2015-0820-0012.

16.   Donley N. The USA lags behind other agricultural nations in banning harmful

pesticides. Environ Health. 2019;18(1):44. doi: 10.1186/s12940-019-0488-0. PubMed

PMID: 31170989; PubMed Central PMCID: PMCPMC6555703.

17.   Achor D, Welker S, Ben-Mahmoud S, Wang C, Folimonova SY, Dutt M, et al.

Dynamics of Candidatus Liberibacter asiaticus Movement and Sieve-Pore Plugging in

Citrus Sink Cells. Plant Physiol. 2020;182(2):882-91. doi: 10.1104/pp.19.01391.

PubMed PMID: 31818905; PubMed Central PMCID: PMCPMC6997701.

18.   Merfa MV, Perez-Lopez E, Naranjo E, Jain M, Gabriel DW, De La Fuente L.

Progress and Obstacles in Culturing 'Candidatus Liberibacter asiaticus', the Bacterium

Associated with Huanglongbing. Phytopathology. 2019;109(7):1092-101. doi:

10.1094/PHYTO-02-19-0051-RVW. PubMed PMID: 30998129.

19.   Acimovic SG, Zeng Q, McGhee GC, Sundin GW, Wise JC. Control of fire blight

(Erwinia amylovora) on apple trees with trunk-injected plant resistance inducers and

antibiotics and assessment of induction of pathogenesis-related protein genes. Front

Plant Sci. 2015;6:16. doi: 10.3389/fpls.2015.00016. PubMed PMID: 25717330; PubMed

Central PMCID: PMCPMC4323746.

20.   Granados-Chinchilla F, Rodriguez C. Tetracyclines in Food and Feedingstuffs:

From Regulation to Analytical Methods, Bacterial Resistance, and Environmental and

Health Implications. J Anal Methods Chem. 2017;2017:1315497. doi:

10.1155/2017/1315497. PubMed PMID: 28168081; PubMed Central PMCID:

PMCPMC5266830.

21.   Leal JF, Santos EBH, Esteves VI. Oxytetracycline in intensive aquaculture: water

quality during and after its administration, environmental fate, toxicity and bacterial

resistance. Reviews in Aquaculture. 2019;11(4):1176-94. doi: 10.1111/raq.12286.

22.   Mayerhofer G, Schwaiger-Nemirova I, Kuhn T, Girsch L, Allerberger F. Detecting

streptomycin in apples from orchards treated for fire blight. J Antimicrob Chemother.

2009;63(5):1076-7. doi: 10.1093/jac/dkp055. PubMed PMID: 19240075.

23.   Araby E, Nada HG, Abou El-Nour SA, Hammad A. Detection of tetracycline and

streptomycin in beef tissues using Charm II, isolation of relevant resistant bacteria and

control their resistance by gamma radiation. BMC Microbiol. 2020;20(1):186. doi:

10.1186/s12866-020-01868-7. PubMed PMID: 32600267; PubMed Central PMCID:

PMCPMC7325294.

24.   Poapolathep A, Poapolathep S, Jermnak U, Imsilp K, Wannapat N, Sugita-

Konishi Y, et al. Muscle tissue kinetics of oxytetracycline following intramuscular and

oral administration at two dosages to giant freshwater shrimp (Macrobrachium

rosenbergii). J Vet Pharmacol Ther. 2008;31(6):517-22. doi: 10.1111/j.1365-

2885.2008.00988.x. PubMed PMID: 19000273.

25.   Al-Rimawi F, Hijaz F, Nehela Y, Batuman O, Killiny N. Uptake, Translocation,

and Stability of Oxytetracycline and Streptomycin in Citrus Plants. Antibiotics (Basel).

2019;8(4). doi: 10.3390/antibiotics8040196. PubMed PMID: 31717884; PubMed Central

PMCID: PMCPMC6963747.

26.   Shentu JL, Zhang K, Shen DS, Wang MZ, Feng HJ. Effect from low-level

exposure of oxytetracycline on abundance of tetracycline resistance genes in arable

soils. Environ Sci Pollut Res Int. 2015;22(17):13102-10. doi: 10.1007/s11356-015-4099-

1. PubMed PMID: 25925140.

27.   Li J, Luo L. Nurturing Undergraduate Researchers in Biomedical Sciences. Cell.

2020;182(1):1-4.

28.   Kraemer SA, Ramachandran A, Perron GG. Antibiotic Pollution in the

Environment: From Microbial Ecology to Public Policy. Microorganisms. 2019;7(6). doi:

10.3390/microorganisms7060180. PubMed PMID: 31234491; PubMed Central PMCID:

PMCPMC6616856.

29.   Li J, Pang Z, Duan S, Lee D, Kolbasov VG, Wang N. The in Planta Effective

Concentration of Oxytetracycline Against 'Candidatus Liberibacter asiaticus' for

Suppression of Citrus Huanglongbing. Phytopathology. 2019;109(12):2046-54. doi:

10.1094/PHYTO-06-19-0198-R. PubMed PMID: 31369360.

30.   Cycon M, Mrozik A, Piotrowska-Seget Z. Antibiotics in the Soil Environment-

Degradation and Their Impact on Microbial Activity and Diversity. Front Microbiol.

2019;10:338. doi: 10.3389/fmicb.2019.00338. PubMed PMID: 30906284; PubMed

Central PMCID: PMCPMC6418018.

31.   Tancos KA, Villani S, Kuehne S, Borejsza-Wysocka E, Breth D, Carol J, et al.

Prevalence of Streptomycin-Resistant Erwinia amylovora in New York Apple Orchards.

Plant Dis. 2016;100(4):802-9. doi: 10.1094/PDIS-09-15-0960-RE. PubMed PMID:

30688602.

32.   Popowska M, Rzeczycka M, Miernik A, Krawczyk-Balska A, Walsh F, Duffy B.

Influence of soil use on prevalence of tetracycline, streptomycin, and erythromycin

resistance and associated resistance genes. Antimicrob Agents Chemother.

2012;56(3):1434-43. doi: 10.1128/AAC.05766-11. PubMed PMID: 22203596; PubMed

Central PMCID: PMCPMC3294877.

33.   Sundin GW, Monks DE, Bender CL. Distribution of the streptomycin-resistance

transposon Tn5393 among phylloplane and soil bacteria from managed agricultural

habitats. Can J Microbiol. 1995;41(9):792-9. doi: 10.1139/m95-109. PubMed PMID:

7585356.

34.   Schmitt H, Stoob K, Hamscher G, Smit E, Seinen W. Tetracyclines and

tetracycline resistance in agricultural soils: microcosm and field studies. Microb Ecol.

2006;51(3):267-76. doi: 10.1007/s00248-006-9035-y. PubMed PMID: 16598633.

35.   Walsh F, Smith DP, Owens SM, Duffy B, Frey JE. Restricted streptomycin use in

apple orchards did not adversely alter the soil bacteria communities. Front Microbiol.

2013;4:383. doi: 10.3389/fmicb.2013.00383. PubMed PMID: 24550889; PubMed

Central PMCID: PMCPMC3908321.

36.   Shade A, Klimowicz AK, Spear RN, Linske M, Donato JJ, Hogan CS, et al.

Streptomycin application has no detectable effect on bacterial community structure in

apple orchard soil. Appl Environ Microbiol. 2013;79(21):6617-25. doi:

10.1128/AEM.02017-13. PubMed PMID: 23974143; PubMed Central PMCID:

PMCPMC3811482.

37.   Xuan R, Arisi L, Wang Q, Yates SR, Biswas KC. Hydrolysis and photolysis of

oxytetracycline in aqueous solution. J Environ Sci Health B. 2010;45(1):73-81. doi:

10.1080/03601230903404556. PubMed PMID: 20390934.

38.   Choi S, Sim W, Jang D, Yoon Y, Ryu J, Oh J, et al. Antibiotics in coastal

aquaculture waters: Occurrence and elimination efficiency in oxidative water treatment

processes. J Hazard Mater. 2020;396:122585. doi: 10.1016/j.jhazmat.2020.122585.

PubMed PMID: 32298861.

39.   Wang Q, Yates SR. Laboratory study of oxytetracycline degradation kinetics in

animal manure and soil. J Agric Food Chem. 2008;56(5):1683-8. doi:

10.1021/jf072927p. PubMed PMID: 18257526.

40.   Liu Y, Bao Y, Cai Z, Zhang Z, Cao P, Li X, et al. The effect of aging on

sequestration and bioaccessibility of oxytetracycline in soils. Environ Sci Pollut Res Int.

2015;22(14):10425-33. doi: 10.1007/s11356-015-4190-7. PubMed PMID: 25721525.

41.   Slana M, Dolenc MS. Environmental Risk Assessment of antimicrobials applied

in veterinary medicine-A field study and laboratory approach. Environ Toxicol

Pharmacol. 2013;35(1):131-41. doi: 10.1016/j.etap.2012.11.017. PubMed PMID:

23274419.

42.   Leal JF, Esteves VI, Santos EBH. Solar photodegradation of oxytetracycline in

brackish aquaculture water: New insights about effects of Ca2+ and Mg2+. Journal of

Photochemistry and Photobiology A: Chemistry. 2019;372:218-25.

43.   Li Z-j, Qi W-n, Feng Y, Liu Y-w, Ebrahim S, Long J. Degradation mechanisms of

oxytetracycline in the environment. Journal of Integrative Agriculture. 2019;18(9):1953-

60.

44.   Shen Y, Zhao W, Zhang C, Shan Y, Shi J. Degradation of streptomycin in aquatic

environment: kinetics, pathway, and antibacterial activity analysis. Environ Sci Pollut

Res Int. 2017;24(16):14337-45. Epub 2017/04/22. doi: 10.1007/s11356-017-8978-5.

PubMed PMID: 28429270.

45.   Vidaver AK. Uses of antimicrobials in plant agriculture. Clin Infect Dis. 2002;34

Suppl 3:S107-10. Epub 2002/05/04. doi: 10.1086/340247. PubMed PMID: 11988880.

46.   Khan S, Osborn A, Eswara PJ. Effect of sunlight on the efficacy of commercial

antibiotics used in agriculture. bioRxiv. 2020;197848. doi:

https://doi.org/10.1101/2020.07.10.197848.

Figure 1