A drug repurposing screen identifies hepatitis C antivirals as inhibitors of the SARS-CoV-2 main
protease.

Jeremy D. Baker[1,2], Rikki L. Uhrich[2], Gerald C. Kraemer[3], Jason E. Love[,4], and Brian C. Kraemer[1,2,5,6*]

[1]Division of Gerontology and Geriatric Medicine, Department of Medicine, University of Washington,
Seattle, WA 98104, USA.
[2]Geriatrics Research Education and Clinical Center, Veterans Affairs Puget Sound Health Care System,
Seattle, WA 98108, USA.
[3]Thomas Jefferson High School, Auburn, WA 98001, USA.
[4]Western Washington Pathology, Tacoma, 98405, USA.
[5]Department of Psychiatry and Behavioral Sciences, University of Washington, Seattle, Washington 98195,
USA.
[6]Department of Pathology, University of Washington, Seattle, Washington 98195, USA.

*Corresponding Author (Email: kraemerb@uw.edu)

Abstract

The SARS coronavirus type 2 (SARS-CoV-2) emerged in late 2019 as a zoonotic virus highly transmissible

between humans that has caused the COVID-19 pandemic [1,2]. This pandemic has the potential to disrupt

healthcare globally and has already caused high levels of mortality, especially amongst the elderly. The

overall case fatality rate for COVID-19 is estimated to be ~2.3% overall [3] and 32.3% in hospitalized patients

age 70-79 years [4].   Therapeutic options for treating the underlying viremia in COVID-19 are presently

limited by a lack of effective SARS-CoV-2 antiviral drugs, although steroidal anti-inflammatory treatment

can be helpful. A variety of potential antiviral targets for SARS-CoV-2 have been considered including

the spike protein and replicase. Based upon previous successful antiviral drug development for HIV-1 and

hepatitis C, the SARS-CoV-2 main protease (Mpro) appears an attractive target for drug development.

Here we show the existing pharmacopeia contains many drugs with potential for therapeutic repurposing

as selective and potent inhibitors of SARS-CoV-2 Mpro. We screened a collection of ~6,070 drugs with a

previous history of use in humans for compounds that inhibit the activity of Mpro *in vitro.* In our primary

screen we found ~50 compounds with activity against Mpro (overall hit rate <0.75%). Subsequent dose

validation studies demonstrated 8 dose responsive hits with an IC50 $\leq$ 50 μM. Hits from our screen are

enriched with hepatitis C NS3/4A protease targeting drugs including Boceprevir (IC50=0.95 μM),

Ciluprevir (20.77μM). Narlaprevir (IC50=1.10μM), and Telaprevir (15.25μM). These results demonstrate

that some existing approved drugs can inhibit SARS-CoV-2 Mpro and that screen saturation of all approved

drugs is both feasible and warranted.   Taken together this work suggests previous large-scale commercial

drug development initiatives targeting hepatitis C NS3/4A viral protease should be revisited because some

previous lead compounds may be more potent against SARS-CoV-2 Mpro than Boceprevir and suitable for

rapid repurposing.

Keywords

COVID-19; SARS-CoV-2; Protease, Mpro; Antiviral; drug repurposing. Boceprevir, Narlaprevir,
Telaprevir, 3C-like protease

Introduction

The SARS virus and SARS-CoV-2, the cause of the COVID-19 pandemic, are zoonotic coronaviruses found in bats that can infect humans. Initial symptoms of SARS-CoV-2 infection include fever, myalgia, cough, and headache. Infection usually resolves without active medical intervention, but for a subset of cases infection can progress to viral pneumonia and a variety of complications including acute lung damage leading to death [5]. While complications are atypical in most cases, mortality rates increase dramatically with the age and impaired health of infected patients. To date, much of our knowledge of COVID-19 virology has been inferred from the study of similar Severe Acute Respiratory Syndrome (SARS) coronavirus and related coronaviruses including Middle East Respiratory Syndrome [reviewed in [6]]. Like all coronaviruses, SARS-CoV-2 exhibits an enveloped ribonucleoprotein helical capsid containing a single positive-stranded genomic RNA. Infection starts with receptor-mediated virus internalization, uncoating, and translation of the viral genome [7]. Polyprotein cleavage by viral proteases yields a complement of viral structural and accessory proteins. This polyprotein cleavage is mediated by the main viral protease (Mpro), a chymotrypsin-like protease responsible for endoproteolytic cleavages of viral polyproteins producing functional viral proteins. Recent structural biology work has solved the crystal structure of SARS-CoV-2 Mpro yielding structural insights into Mpro function [8,9].

Antiviral drugs effective for COVID-19 would have a broad impact on global healthcare in the current coronavirus pandemic. Existing antiviral drugs on the market target a wide variety of both RNA and DNA viruses. Probably the most successful approved antivirals are protease inhibitors such as atazanavir for HIV-1 and simeprevir for hepatitis C. [reviewed in [10] and [11]]. Other conceptual COVID-19 antiviral targets include the host ACE2 receptor, viral replicase, and viral genome encapsidation. However, previous work with other RNA viruses suggest that Mpro function is essential for viral replication and readily targetable using existing technology. Thus, while there are many potentially

targetable activities for COVID-19, the coronavirus Mpro seems a likely choice for rapid drug

development.

To accelerate drug development we employed a drug repurposing strategy, an approach of utilizing

previously approved drugs for new indications [12,13]. Previous work suggests libraries enriched with known

bioactive drug-like compounds provide the best opportunity for finding new lead compounds [14,15]. Thus

we attempted the selective optimization of side activities (SOSA) approach [16] as a rapid and cost effective

means to identify candidate hits while minimizing the number of compounds screened. The SOSA

approach proceeds by two steps. First a limited set of carefully chosen, structurally diverse, well-

characterized drug molecules are screened; as approved drugs, their bioavailability, toxicity and efficacy

in human therapy has already been demonstrated [16,17]. To screen as much of the available approved drug

space as possible in an easily accessible format we chose to screen the Broad Institute Drug Repurposing

Library (6070 compounds, see **Table S1**) [18]. This represents about half of the approximately 14,000

approved or experimental drugs known to human clinical medicine [19]. There are significant cost and time

advantages realized by drug repurposing as it can accelerate the preclinical phase of development and

streamline clinical trials to focus on efficacy rather than safety.

Repositioning existing approved drugs with the capacity to inhibit COVID-19 virus replication and

infection would be of profound utility and immediately impact health care in the current pandemic. There

are no drugs in clinical use specifically targeting coronavirus replication. The major advantage of the

approach taken here is that by screening drugs with a history of previous clinical use, we will be focusing

on compounds with known properties in terms of pharmacokinetics (PK), pharmacodynamics (PD) and

toxicity. Thus, the Broad Repurposing Library we screened consists of compounds suitable for rapid

translation to human efficacy trials.

Results

<u>Development of fluorescent Mpro Assays</u>

We began assay development by selecting potentially suitable synthetic Mpro substrates and compared

catalyzed hydrolysis curves between 5 fluorescently labeled substrates (Ac-Abu-Tle-Leu-Gln-AFC [20],

DABCYL-VKLQ-EDANS, Ac-VKLQ-AFC, DABCYL-TSAVLQSGFRKM-EDANS[21] , and MCA-

AVLQSGFR-K(Dnp)-K-NH2)[22]. We chose to use the recently published Ac-Abu-Tle-Leu-Gln-AFC

(Abu=2-Aminobutyrate, Tle=tButylglycine) synthetic non-canonical amino-acid containing peptide as

Mpro more readily cleaves this preferred sequence as compared to the native VKLQ sequence [20](Fig 1A).

Substrates   DABCYL-TSAVLQSGFRKM-EDANS   and   MCA-AVLQSGFR-K(DnP)-K-NH2   had

drastically lower rates of Mpro catalyzed hydrolysis and were not considered further in our assay

development (Fig 1A). To determine concentration ratios between Mpro and substrate, we next preformed

a two-dimensional titration and chose 625nM Mpro and 8µM substrate for a balance of relatively modest

Mpro protein requirement and a robust fluorescence intensity (Fig 2B). Before screening the Broad library,

we piloted our assay conditions against the NIH Clinical collections library (~650 compounds) and

calculated our Z'-factor for each plate at 0.780 and 0.784 (Fig 1C and D). Z'-factor is a score of suitability

of assays for high-throughput screening and is derived from the equation $Z'\text{-}factor = 1 - \frac{3(\sigma_p + \sigma_n)}{|\mu_p - \mu_n|}$, where

σ = standard deviation, µ=mean, p=positive controls, and n=negative controls. A score greater than 0.5

indicates a screenable assay. Although no promising compounds were identified from this smaller library,

it demonstrated that our assay was sufficiently robust for screening the much larger Broad Repurposing

library.

Fig 1. Mpro assay optimization
(a) Screen of selection of reported Mpro substrates: DABCYL-VKLQ-EDANS, Ac-VKLQ-AFC, Ac-Abu-Tle-Leu-Gln-AFC, DABCYL-TSAVLQSGFRKM-EDANS, MCA-AVLQSGFR-K(Dnp)-K-NH$_2$. Fold change of increase in RFU was measured over 30 minutes holding substrate constant at 12.5µM with increasing concentrations of Mpro recombinant protein as indicated. (b) Two-dimensional titration of substrate Ac-Abu-Tle-Leu-Gln-AFC against Mpro. Concentrations are indicated for substrate along the x-axis and for Mpro along the y-axis. Heat map corresponds to the change in RFU over 10 minutes and pink outline (ΔRFU 10m=20,810) indicates chosen concentration for NIH Clinical Collection screen (0.625µM Mpro and 8µM Substrate). (c) Z-score index of NIH Clinical Collection screen (640 compounds). Hit window was considered at Z-score ≤ -2 and was calculated as the Z-score of ΔRFU at 10 minutes corresponding to the linear portion of the curve. X-axis indicates arbitrary compound number arranged by increasing Z-score. (d) Z'-factor for the two NIH Clinical Collection 384-well plates. Pink circles indicate negative control (DMSO) and black circles represent positive controls (no protein). Z'factor calculated at 0.780 and 0.784 for plates 1 and 22 respectively. Y axis represents change in RFU over 10 minutes.

Drug Repurposing Strategy – screening the Broad Repurposing Library
The concept of drug repurposing is to utilize existing therapeutic drugs to treat a new disease indication.

This approach is particularly relevant for COVID-19 because of the potential for an accelerated clinical

impact as compared to *de novo* drug development. A systematic approach to facilitate drug repurposing

has recently been described ([18], http:// www.broadinstitute.org/repurposing) and has made a large collection

of drugs with previous history of use in humans available for high throughput screening. We acquired this

~6,070 compound library as an assay ready collection in 384-well format. We conducted the library screen

at 384-well density using the optimized kinetic Mpro assay described in Fig 1.   Our overall repurposing

strategy is described in Fig 2A. We conducted a single point screen at 50 μM compound concentration and

observed ~50 compounds with activity against SARS-CoV-2 Mpro for an overall hit rate <0.75%. These

compounds were screened in parallel against the natural amino acid substrate (Ac-VKLQ-AFC) as well as

a kinetically preferred substrate (Ac-Abu-Tle-Leu-Gln-AFC) (Fig 2B).   Individual compounds are shown

in Table 1.

Fig 2. Screening pipeline and Broad library screen
(a) Schematic of screening funnel. The Broad Repurposing library was screened both empirically and
virtually. Any hit from the Broad library (Z-score ≤ -2) was validated for dose-responsiveness. All suitable
compounds passing this filter with satisfactory curve fitting and potency were ordered as powder and re-
validated. Future efforts will test for selectivity and in orthogonal assays for suitability. Although outside
the scope of this report, determination of viral anti-replicative properties as well as toxic profile at required
dosage will be determined. The goal of this paradigm is to find suitable candidates for development both
as tools for probing underlying mechanisms of SARS-CoV-2 as well as for translational potential. (b)
Screen of the Broad Repurposing Library. Library was screened at a concentration of 50μM against both
Ac-VKLQ-AFC (black) and Ac-Abu-Tle-Leu-Gln-AFC (purple). Hit window was considered for
compounds falling below Z-score ≤ -2 against both substrates and consisted of 50 compounds. Compounds
ordered by average Z-score.

152 **Table 1.**
153 Hits identified from Broad Repurposing Library screen against SARS-CoV-2 Mpro.
154

Common name	Mechanism of Action	Target	Status
Mitoquinone			
Octenidine	membrane integrity inhibitor		Launched
Boceprevir	HCV inhibitor	CMA1, CTSA, CTSF, CTSK, CTSL, CTSS	Launched
STS		PROC, PROS1	Launched
NH125	EEF2 inhibitor		Preclinical
Visomitin			
Cetrimonium			Launched
Hexachlorophene	potassium channel activator	GLUD1, SDHD	Launched
Benzethonium	sodium channel blocker	SCN10A	Launched
Rose-bengal	contrast agent		Launched
LGD-6972			
Narlaprevir	HCV inhibitor		P2/P3
Obatoclax	BCL inhibitor	BCL2	Phase 3
Domiphen			Preclinical
PSB-06126	NTPDase inhibitor	ENTPD3	Preclinical
NSC-95397	CDC inhibitor	CDC25A, CDC25B	Preclinical
Deltarasin	phosphodiesterase inhibitor	KRAS	Preclinical
Calpeptin	calpain inhibitor		Preclinical
TNP-470	methionine aminopeptidase inhibitor		Phase 2
TC-LPA5-4	lysophosphatidic acid receptor antagonist	LPAR5	Preclinical
Hemin	enzyme inducer		Launched
Sennoside			Preclinical
PYR-41	ubiquitin activating enzyme inhibitor		Preclinical
Telaprevir	HCV inhibitor	CTSA, PGR	Launched
Evans-blue	glutamate receptor modulator	GRIA1, PTPN1	Launched
C646	histone acetyltransferase inhibitor	EP300	Preclinical
NSC-663284	CDC inhibitor	CDC25A, CDC25B, CDC25C	Preclinical
TCID			Preclinical
Hematoporphyrin			Launched
16-BAC	cationic surfactant		Preclinical
BMS-833923	smoothened receptor antagonist	SMO	Phase 2
AVN-492			
Ascorbyl palmitate			Preclinical
Elacestrant			
Eifuroxazide			
Aurothioglucose	PKC inhibitor	PRKCI	Launched
Nifursol	bacterial DNA inhibitor		Launched
NS-1643	voltage-gated K^+ channel activator	KCNH2, KCNH6, KCNMA1	Preclinical
Tiplaxtinin	plasminogen activator inhibitor	SERPINE1	Phase 1
Thiomersal	antibiotic	OXCT1	Launched
Indocyanine-green	contrast agent	SLCO1B1	Launched
Chlortetracycline	protein synthesis inhibitor		Launched
Carbazochrome			Launched
Altrenogest	progestogen hormone	PGR	Launched
Emricasan			
GSK2801	bromodomain inhibitor	BAZ2A, BAZ2B	Preclinical
Anthralin	DNA synthesis inhibitor		Launched
Melphalan	DNA alkylating agent, DNA inhibitor		Launched
RITA	MDM inhibitor	MDM2	Preclinical
Azeliragon	RAGE receptor antagonist	AGER	Phase 3

Analysis of potency

We validated the hits from the primary screen by conducting a 10-point dose-response analysis with a drug
concentration range from 150 μM down to 7.6 nM (3-fold dilution series). From this dose-response analysis,
IC50 values were calculated for dose-responsive hits. Several of the drugs uncovered in our screen
including, Boceprevir (IC50 = 0.95 μM), Ciluprevir (IC50 = 20.77 μM), Narlaprevir (IC50 = 1.10 μM),
Telaprevir (IC50 = 15.25 μM), are antiviral compounds targeting the hepatitis C NS3 protease.
Boceprevir, Narlaprevir, and Telaprevir are approved drugs with a track record of safe use in human patients
[23-28]. Other relatively potent dose responsive compounds emerging from our screen include calpeptin (IC50
= 4.05 μM), aurothioglucose (IC50 = 13.32 μM), PYR-41 (IC50 = 17.38 μM), and hemin (IC50= 9.68 μM)
(Fig 3).

Fig 3. Dose-response validation of compounds against SARS-CoV-2 Mpro
Dose response curves were generated using Ac-Abu-Tle-Leu-Gln-AFC substrate. Percent inhibition of
each compound was calculated at indicated 10 concentrations by comparing slope of treatment versus
DMSO control. Error bars represent standard deviation and n=3 for each concentration. IC50 values are
indicated and calculated by 4-parameter nonlinear regression curve fitting.
175

The relative utility of *in silico* and HTS repurposing screens

The recent publication of the crystal structure for Mpro has enabled computational approaches to Mpro

drug discovery [8,9]. We leveraged the existing structural data (PBD entry 6LU7) to conduct a computational

free energy calculation based *in silico* screening approach. To do this we have utilized the Schrodinger

Maestro software package [29,30] to conduct a computational docking of all compounds in the Broad

Repurposing library.   Using this approach, we derived a docking score for each compound (see Table S1

for broad repurposing library with docking scores). We observe a poor correlation (Pearson r=0.02864)

between Mpro docking score and Z-score in the protease inhibition assay (Fig 4A). Furthermore, top hits

from the screen also exhibit a weak correlation (Pearson r=-0.1503) between compound potency and

docking score (Fig 4B).

Fig 4. Correlation of Experimental and In Silico identification

(a) Correlation of Z-score for all Broad Repurposing library compounds and Docking Score. Pearson r =
0.02864 (95% CI 0.003478-0.05376), P value = 0.0257 and XY pairs = 6068. Linear equation Y=0.03452x+
0.2202 as fitted by simple linear regression, slope significantly deviated from 0, P value = 0.0016. (b)
Correlation of IC50 vs Docking Score. Pearson r = -0.1053 (95% CI -0.5136-0.3420), P value = 0.6498 and
XY Pairs = 21. Linear equation Y=-4.188x+20.41 as fitted by simple linear regression, slope not
significantly deviated from 0, P value = 0.6498.

Table S1. Broad Library SDF with Docking scores—see attached file

Discussion

Therapeutic potential of Mpro as a target

A general discussion of drug repurposing for COVID-19 suggests viral encoded proteases may be relevant

therapeutic targets for coronaviruses [31]. Relative to most other human viruses, our understanding of the

virology of SARS-CoV-2 remains incomplete. However, after decades of extensive research, we have

learned a great deal about viral proteases in general and the chemical means to inhibit them from our studies

of HIV-1, hepatitis C and rhinoviruses. Likewise protease inhibitors targeting the SARS protease have

been investigated [32,33]. Furthermore, previous work on SARS Mpro has demonstrated that it is a targetable

enzyme worth significant translational effort [34].

Perspective on hit compounds and future screening

Sequence alignment shows a high degree of homology between SARS Mpro and SARS-CoV-2 Mpro with

~95% amino acid sequence identity. Recent studies have solved the crystal structure of SARS-CoV-2 Mpro

and compared it with SARS Mpro showing that they have similar but distinct active site pockets and will

require distinct dugs for potent and highly specific inhibition. This sequence and structural information has

provided an opportunity to conduct *in silico* docking of known drugs to the COVID-19 virus Mpro active

site. However thus far, such analysis has not uncovered potent inhibitors of Mpro. Recent *in silico* work

has suggested that protease inhibitor drugs may inhibit SARS-CoV-2 Mpro [35,36].   However, our findings

suggest that *in silico* approaches alone cannot substitute for enzyme kinetic screening evaluation of Mpro

inhibitors because most identified high scoring compounds in *in silico* docking studies lack activity against

Mpro in kinetic protease assays.

Potential for Drug Repurposing and Clinical Trials

To date clinical trials for COVID-19 have not yielded potent antiviral therapies. The objective of this work

is to complete a survey of approved drugs to identify therapies that can block COVID-19 viral replication

by inhibiting the main viral protease. The advantage of this approach is that any approved drug identified

can be advanced rapidly to clinical trials without extensive multi-year preclinical development efforts. This

is also particularly germane given the limitations of animal models of COVID-19 infection and

pathogenesis.

A diverse variety of initial hits were identified in our high throughput screen of the broad library.

Of these, the most potent hits are all known protease inhibitors and there is strong representation from

protease inhibitors developed to inhibit HCV protease NS3/4A (Boceprevir, Ciluprevir, Narlaprevir, and

Telaprevir).   Clearly   as   approved   or   well-developed   clinical   candidates,   these   drugs   exhibit

pharmacological and pharmacodynamic properties well suited to repurposing as a COVID-19 antiviral

therapy.   Boceprevir and Narlaprevir appear the most potent against Mpro and may be suitable for

repurposing.

Previous clinical evaluation of Boceprevir (also known as Victrelis) showed it to be safe and

effective for treating HCV [37]. Boceprevir was approved as a first in class HCV NS3/4A serine protease

inhibitor for treatment of chronic HCV infection. Boceprevir was FDA approved for use in the USA in

2011 and Boceprevir treatment is given as a combination therapy with interferon α2b and ribavirin.

Likewise, clinical evaluation of Narlaprevir (also known as Arlansa or SCH900518) showed it to be both

safe and to exhibit antiviral activity when combined with interferon α2b [38]. Furthermore, Narlaprevir has

been show effective against HCV NS3/4A mutations causing resistance to protease inhibitors [39].

Narlaprevir was approved for use against HCV in Russia in 2016.

Our findings demonstrate Boceprevir and Narlaprevir potency against SARS-CoV-2 Mpro in the

one micromolar range. Previous work to develop Boceprevir and Narlaprevir as approved HCV

therapeutics make them drug repurposing candidates worth further evaluation. Likewise, previous

commercially developed NS3/4A inhibitor lead compounds may be suitable for further repurposing studies.

The COVID-19 pandemic has revealed an urgent unmet medical need for potent antiviral agents for

treatment of SARS-CoV-2 infection. Because antiviral therapies are frequently most effective when used

in combination [40], it may be useful to consider combining Mpro inhibition with other antiviral strategies for

treating SARS-CoV-2 infection. For instance, inhibition of the SARS-CoV-2 replicase in combination with

Mpro inhibition might exhibit synergistic antiviral activity. Remdesivir, a broad-spectrum antiviral

replicase inhibitor has shown efficacy against a wide variety of RNA viral replicases including SARS-CoV-

2 [41,42]. Thus, combination therapies using Boceprevir/Remdesivir or Narlaprevir/Remdesivir may yield a

synergistic drug repurposing strategy for treating COVID-19.   Taken together, the work presented here

supports the rapid evaluation of previous HCV NS3/4A inhibitors for repurposing as a COVID-19 therapy.

Methods:

Recombinant Protein

Recombinant Mpro was purified using constructs and methods as previously described [8]. pGEX-6p-1

plasmid containing SARS-CoV-2 Mpro was gifted from Hilgenfeld lab at Luebeck University, Germany.

Plasmid was transformed into BL21 (DE3) bacteria (NEB). A single colony was inoculated into 10mL

Terrific Broth (TB) + Carbenicillin (25ug/mL) and grown overnight to saturation. Overnight culture was

transferred into 1L of TB and grown in a shaking incubator at 37ºC until log phase (OD_{600}~0.7). Culture

was induced with IPTG (1mM final) and kept in 37ºC shaking incubator for 4 hours. Culture was spun

down at 3,400rpm for 30 min at 4ºC, and pellet resuspended in PBS with 10% sucrose then spun at previous

conditions. PBS was aspirated and bacteria pellet was snap frozen in liquid nitrogen and stored at -70ºC.

Pellet was thawed and resuspended in Lysis buffer (PBS, 0.3% lysozyme, 1mM DTT, 1.5% Sarkosyl,

RNAse A, and DNAse I) and sonicated for 10 seconds ON time, 20 seconds OFF time for 5 minutes of

total ON time at 60% amplitude. Lysate was spun at 16,000rpm for 30 minutes at 4ºC. 4mL of Ni-NTA

beads and supernatant were rotated for 2 hours at room temperature. Gravity column was used for

purification with His_6-tagged Mpro binding to Ni-NTA beads (Qiagen 30210), washed with lysis buffer +

10mM imidazole and eluted with increasing concentration of imidazole (50mM, 100mM, 150mM and

200mM). The majority of Mpro eluted at between 150-200mM imidazole and was 90%+ pure by

Coomassie stained gel analysis. Mpro fractions were pooled and buffer exchanged into 20mM Tris pH 7.8,

150mM NaCl, 1mM EDTA, 1mM DTT, and snap frozen in liquid nitrogen and stored at -70ºC. Yield and

purity were assessed via BCA (ThermoFisher 23225) and Coomassie-stained SDS-PAGE.

Fluorescent Mpro Protease Assay

Fluorescent Mpro peptides were synthesized by Anaspec (www.anaspec.com) at 90% purity and frozen in 1 mg aliquots. Stock concentration of substrates were made by reconstituting powder in 100uL DMSO (10mg/ml) and storing at -70°C. Optimal Mpro substrates were previously determined to be Ac-Val-Lys-Leu-Gln-AFC for physiological substrates and Ac-Abu-Tle-Leu-Gln-AFC, a noncanonical amino acid substrate [20]. Ac-Val-Lys-Leu-Gln-AFC and Ac-Abu-Tle-Leu-Gln-AFC fluorogenic substrates were monitored at 380/20 nm excitation and 500/20 nm emission wavelengths. FRET-based substrates Dabcyl-Val-Lys-Leu-Gln-EDANS was measured at 336/20 nm excitation and 490/20 nm emission wavelengths and MCA-Ala-Val-Lys-Gln-Ser-Gly-Phe-Lys-DNP-Lys was monitored at 325/20 nm excitation and 392/20 nm emission wavelengths. We used 20mM Tris pH 7.8, 150mM NaCl, 1mM EDTA, 1mM DTT, 0.05% Triton X-100 as the assay buffer. Assay conditions were at room temperature (25°C) for all assays.

2D Titration main screen optimization

2D titration for determining the main screen ratios was done in 96 well black opaque plates (Corning 3686 NBS). The top concentration of Mpro was 2.5μM and serial diluted to 0.0395μM along the Y-axis of the plate. The top concentration of substrate was 16μM and serial diluted along the X-axis of the plate. Fluorescence was monitored at 380/20 nm excitation and 500/20 nm emission wavelengths.

Broad Repurposing Library

The Broad Repurposing Library was ordered and plated into black opaque 384-well plates (Greiner 781209) at 100nL of 10mM (slight variations depending on compound) compound in DMSO. 10uL of diluted Mpro (625nM final concentration in reaction buffer detailed above) was added with a MultiFlo FX liquid dispenser using a 5μL cassette. Compounds were incubated with Mpro for 10 minutes at RT after which

303   10uL of substrate (8uM final concentration of either Ac-VKLQ-AFC or Ac-Abu-Tle-Leu-Gln-AFC) was

304   dispensed into the plate and read using a Cytation 5 multi-mode reader immediately at 380/20 nm excitation

305   and 500/20 nm emission wavelengths every 5 minutes for 30 minutes. Data was analyzed using Biotek

306   Gen5 software, Microsoft Excel, and GraphPad Prism 8.

Dose validation assays

309   Hit compounds were ordered from the Broad Institute pre-plated in 384-well format (Greiner 781209) as

310   10-point serial dilutions (3-fold) at 300nL per well. Mpro (80nM final concentration) and substrate (Ac-

311   Abu-Tle-Leu-Gln-AFC at 32μM final concentration) were dispensed in the same manner described above.

312   Inhibition was calculated as $1 - \frac{\Delta RFU(10m)sample}{\Delta RFU(10m)control}$ at each concentration and data fitted to 4-parameter

313   nonlinear regression model using GraphPad Prism 8.

In Silico **Docking of the Broad Repurposing Library with Mpro**

316   We utilized the Schrodinger Maestro software package [29,30] to conduct a computational docking of all

317   compounds in the Broad Repurposing library. In this approach we generated a receptor grid model of the

318   Mpro active site and serially docked each compound in the Broad Repurposing library with the active site

319   model using the physics based Glide algorithm [43,44]. We chose the Glide algorithm over the many competing

320   options because of its superior performance in head to head comparisons of algorithms [43].

Statistical analyses, IC50 Calculation, Selectivity Calculation, and figures.

323   Graphs were generated using GraphPad Prism 8. IC50 calculations were performed using GraphPad Prism

324   8 curve fitting using 4-parameter non-linear regression.

Author Contributions

J.D.B. and B.C.K. designed research; J.D.B., R.L.U. performed research; J.C.K performed *in silico* docking.

J.D.B. and B.C.K. analyzed data; B.CK. & J.D.B. wrote the first draft of the manuscript. All authors

reviewed, edited, and approved the final version of the manuscript.

Funding Sources

This work was funded by the Department of Veterans Affairs and a Postdoctoral Fellowship from the

Washington Research Foundation (WRF to J.D.B.)

Acknowledgements:

We thank Timothy Strovas, Aleen Saxton, Caleb A. Baker, Misa Baum, and Jeanna Wheeler for sharing

their expertise. We thank the Center for the Development of Therapeutics and Repurposing Hub at the

Broad for providing the compounds library (https://www.broadinstitute.org/repurposing).   We thank

James Meabon for assistance with acquiring and maintaining critical HTS infrastructure.

340 **Data Sharing:**

341 Raw data from the primary screen of the Broad library is available online in the Sage Biosystems data

342 repository, synapse.org (see: https://www.synapse.org/#!Synapse:syn22275070/files/).

345   References

1   Phelan, A. L., Katz, R. & Gostin, L. O. The Novel Coronavirus Originating in Wuhan,
China: Challenges for Global Health Governance. *JAMA*, doi:10.1001/jama.2020.1097
(2020).

2   Chang *et al.* Epidemiologic and Clinical Characteristics of Novel Coronavirus Infections
Involving 13 Patients Outside Wuhan, China. *JAMA*, doi:10.1001/jama.2020.1623 (2020).

3   Rabi, F. A., Al Zoubi, M. S., Kasasbeh, G. A., Salameh, D. M. & Al-Nasser, A. D. SARS-
CoV-2 and Coronavirus Disease 2019: What We Know So Far. *Pathogens* **9**,
doi:10.3390/pathogens9030231 (2020).

4   Zhao, H. L., Huang, Y. M. & Huang, Y. Mortality in Older Patients with COVID-19. *J Am
Geriatr Soc*, doi:10.1111/jgs.16649 (2020).

5   Zhao, W., Zhong, Z., Xie, X., Yu, Q. & Liu, J. Relation Between Chest CT Findings and
Clinical Conditions of Coronavirus Disease (COVID-19) Pneumonia: A Multicenter
Study. *AJR Am J Roentgenol*, 1-6, doi:10.2214/AJR.20.22976 (2020).

6   de Wit, E., van Doremalen, N., Falzarano, D. & Munster, V. J. SARS and MERS: recent
insights into emerging coronaviruses. *Nat Rev Microbiol* **14**, 523-534,
doi:10.1038/nrmicro.2016.81 (2016).

7   Shi, S. T. & Lai, M. M. Viral and cellular proteins involved in coronavirus replication.
Curr Top Microbiol Immunol **287**, 95-131, doi:10.1007/3-540-26765-4_4 (2005).

8   Zhang, L. *et al.* Crystal structure of SARS-CoV-2 main protease provides a basis for design
of improved alpha-ketoamide inhibitors. *Science*, doi:10.1126/science.abb3405 (2020).

9   Jin, Z. *et al.* Structure of M(pro) from COVID-19 virus and discovery of its inhibitors.
Nature, doi:10.1038/s41586-020-2223-y (2020).

10   Piliero, P. J. Atazanavir: a novel HIV-1 protease inhibitor. *Expert opinion on
investigational drugs* **11**, 1295-1301, doi:10.1517/13543784.11.9.1295 (2002).

11   You, D. M. & Pockros, P. J. Simeprevir for the treatment of chronic hepatitis C. *Expert
opinion on pharmacotherapy* **14**, 2581-2589, doi:10.1517/14656566.2013.850074 (2013).

12   Ashburn, T. T. & Thor, K. B. Drug repositioning: identifying and developing new uses for
existing drugs. *Nat Rev Drug Discov* **3**, 673-683, doi:10.1038/nrd1468 (2004).

13   Tobinick, E. L. The value of drug repositioning in the current pharmaceutical market. *Drug
News Perspect* **22**, 119-125, doi:10.1358/dnp.2009.22.2.1303818 (2009).

14 Kwok, T. C. *et al.* A small-molecule screen in C. elegans yields a new calcium channel
antagonist. *Nature* **441**, 91-95 (2006).

15 Petrascheck, M., Ye, X. & Buck, L. B. An antidepressant that extends lifespan in adult
Caenorhabditis elegans. *Nature* **450**, 553-556 (2007).

16 Wermuth, C. G. Selective optimization of side activities: the SOSA approach. *Drug Discov*
Today **11**, 160-164, doi:S1359-6446(05)03686-X [pii] 10.1016/S1359-6446(05)03686-X
(2006).

17 Wermuth, C. G. Selective optimization of side activities: another way for drug discovery.
J Med Chem **47**, 1303-1314, doi:10.1021/jm030480f (2004).

18 Corsello, S. M. *et al.* The Drug Repurposing Hub: a next-generation drug library and
information resource. *Nat Med* **23**, 405-408, doi:10.1038/nm.4306 (2017).

19 D. Wishart, C. K., V. Law. *Drug Bank: Open Data Drug & Drug Target Database*,
<http://www.drugbank.ca/stats> (2012).

20 <https://doi.org/10.1101/2020.03.07.981928> (2020).

21 Chuck, C. P. *et al.* Profiling of substrate specificity of SARS-CoV 3CL. *PLoS One* **5**,
e13197, doi:10.1371/journal.pone.0013197 (2010).

22 Dai, W. *et al.* Structure-based design of antiviral drug candidates targeting the SARS-CoV-
2 main protease. *Science* **368**, 1331-1335, doi:10.1126/science.abb4489 (2020).

23 Njoroge, F. G., Chen, K. X., Shih, N. Y. & Piwinski, J. J. Challenges in modern drug
discovery: a case study of boceprevir, an HCV protease inhibitor for the treatment of
hepatitis C virus infection. *Acc Chem Res* **41**, 50-59, doi:10.1021/ar700109k (2008).

24 Rotella, D. P. The discovery and development of boceprevir. *Expert opinion on drug*
discovery **8**, 1439-1447, doi:10.1517/17460441.2013.843525 (2013).

25 Arasappan, A. *et al.* Discovery of Narlaprevir (SCH 900518): A Potent, Second Generation
HCV NS3 Serine Protease Inhibitor. *ACS medicinal chemistry letters* **1**, 64-69,
doi:10.1021/ml9000276 (2010).

26 Hotho, D. M. *et al.* Sustained virologic response after therapy with the HCV protease
inhibitor narlaprevir in combination with peginterferon and ribavirin is durable through
long-term follow-up. *J Viral Hepat* **20**, e78-81, doi:10.1111/jvh.12012 (2013).

27 Reesink, H. W. *et al.* Rapid decline of viral RNA in hepatitis C patients treated with VX-
950: a phase Ib, placebo-controlled, randomized study. *Gastroenterology* **131**, 997-1002,
doi:10.1053/j.gastro.2006.07.013 (2006).

28 Lang, L. Combination therapy with telaprevir and pegylated interferon suppresses both
wild-type and resistant hepatitis C virus. *Gastroenterology* **132**, 5-6,
doi:10.1053/j.gastro.2006.12.011 (2007).

29 Wang, L. *et al.* Accurate and reliable prediction of relative ligand binding potency in
prospective drug discovery by way of a modern free-energy calculation protocol and force
field. *J Am Chem Soc* **137**, 2695-2703, doi:10.1021/ja512751q (2015).

30 Kuhn, B. *et al.* Prospective Evaluation of Free Energy Calculations for the Prioritization of
Cathepsin L Inhibitors. *J Med Chem* **60**, 2485-2497, doi:10.1021/acs.jmedchem.6b01881
(2017).

31 Li, G. & De Clercq, E. Therapeutic options for the 2019 novel coronavirus (2019-nCoV).
Nat Rev Drug Discov **19**, 149-150, doi:10.1038/d41573-020-00016-0 (2020).

32 Wang, H., Xue, S., Yang, H. & Chen, C. Recent progress in the discovery of inhibitors
targeting coronavirus proteases. *Virologica Sinica* **31**, 24-30, doi:10.1007/s12250-015-
3711-3 (2016).

33 Zhao, Q., Weber, E. & Yang, H. Recent developments on coronavirus main protease/3C
like protease inhibitors. *Recent patents on anti-infective drug discovery* **8**, 150-156,
doi:10.2174/1574891x113089990017 (2013).

34 Chu, C. M. *et al.* Role of lopinavir/ritonavir in the treatment of SARS: initial virological
and clinical findings. *Thorax* **59**, 252-256, doi:10.1136/thorax.2003.012658 (2004).

35 Chen, Y. W., Yiu, C. B. & Wong, K. Y. Prediction of the SARS-CoV-2 (2019-nCoV) 3C-
like protease (3CL (pro)) structure: virtual screening reveals velpatasvir, ledipasvir, and
other drug repurposing candidates. *F1000Research* **9**, 129,
doi:10.12688/f1000research.22457.2 (2020).

36 Shamsi, A. *et al.* Glecaprevir and Maraviroc are high-affinity inhibitors of SARS-CoV-2
main protease: possible implication in COVID-19 therapy. *Bioscience reports* **40**,
doi:10.1042/BSR20201256 (2020).

37 Poordad, F. *et al.* Boceprevir for untreated chronic HCV genotype 1 infection. *N Engl J
Med* **364**, 1195-1206, doi:10.1056/NEJMoa1010494 (2011).

38 de Bruijne, J. *et al.* Antiviral activity of narlaprevir combined with ritonavir and pegylated
interferon in chronic hepatitis C patients. *Hepatology* **52**, 1590-1599,
doi:10.1002/hep.23899 (2010).

39 Tong, X. *et al.* Preclinical characterization of the antiviral activity of SCH 900518
(narlaprevir), a novel mechanism-based inhibitor of hepatitis C virus NS3 protease.
Antimicrob Agents Chemother **54**, 2365-2370, doi:10.1128/AAC.00135-10 (2010).

40   Tan, S. L., He, Y., Huang, Y. & Gale, M., Jr. Strategies for hepatitis C therapeutic
intervention: now and next. *Curr Opin Pharmacol* **4**, 465-470,
doi:10.1016/j.coph.2004.07.003 (2004).

41   Brown, A. J. *et al.* Broad spectrum antiviral remdesivir inhibits human endemic and
zoonotic deltacoronaviruses with a highly divergent RNA dependent RNA polymerase.
Antiviral Res **169**, 104541, doi:10.1016/j.antiviral.2019.104541 (2019).

42   Wang, M. *et al.* Remdesivir and chloroquine effectively inhibit the recently emerged novel
coronavirus (2019-nCoV) in vitro. *Cell Res* **30**, 269-271, doi:10.1038/s41422-020-0282-0
(2020).

43   Friesner, R. A. *et al.* Glide: a new approach for rapid, accurate docking and scoring. 1.
Method and assessment of docking accuracy. *J Med Chem* **47**, 1739-1749,
doi:10.1021/jm0306430 (2004).

44   Halgren, T. A. *et al.* Glide: a new approach for rapid, accurate docking and scoring. 2.
Enrichment factors in database screening. *J Med Chem* **47**, 1750-1759,
doi:10.1021/jm030644s (2004).