Collateral Sensitivity Associated with Antibiotic Resistance Plasmids

Cristina Herencias[1,a], Jerónimo Rodríguez-Beltrán[1,2,a,*], Ricardo León-Sampedro[1,2], Aida Alonso-

del Valle[1], Jana Palkovičová[3], Rafael Cantón[1,4], Álvaro San Millán[1,2,5,*].

[1] *Servicio de Microbiología. Hospital Universitario Ramón y Cajal* and *Instituto Ramón y Cajal de*

Investigación Sanitaria. Madrid, Spain.

[2] *Centro de Investigación Biológica en Red Epidemiología y Salud Pública*, *Instituto de Salud*

Carlos III. Madrid. Spain.

[3] Department of Biology and Wildlife Diseases, Faculty of Veterinary Hygiene and Ecology,

University of Veterinary and Pharmaceutical Sciences Brno, Czech Republic.

[4] *Red Española de Investigación en Patología Infecciosa*. *Instituto de Salud Carlos III*. Madrid.

Spain.

[5] *Centro Nacional de Biotecnología-CSIC*, Madrid, Spain.

[a] These authors contributed equally to this study and should be regarded as joint first authors.

* Correspondence: asanmillan@cnb.csic.es, jeronimo.rodriguez.beltran@gmail.com

Abstract

Collateral sensitivity (CS) is a promising alternative approach to counteract the rising problem of

antibiotic resistance (ABR). CS occurs when the acquisition of resistance to one antibiotic

produces increased susceptibility to a second antibiotic. For CS to be widely applicable in clinical

practice, it would need to be effective against the different resistance mechanisms available to

bacteria. Recent studies have focused on CS strategies designed against ABR mediated by

chromosomal mutations. However, one of the main drivers of ABR in clinically relevant bacteria is

the horizontal transfer of ABR genes mediated by plasmids. Here, we report the first analysis of CS

associated with the acquisition of complete ABR plasmids, including the clinically important

carbapenem-resistance conjugative plasmid pOXA-48. In addition, we describe the conservation of

CS in clinical *E. coli* isolates and its application to the selective elimination of plasmid-carrying

bacteria. Our results provide new insights that establish the basis for developing CS-informed

treatment strategies to combat plasmid-mediated ABR.

Introduction

The rapid evolution of antibiotic resistance (ABR) in bacteria reduces the utility of clinically relevant

antibiotics, making ABR one of the major challenges facing public health (Jim, 2016; MacLean and

San Millan, 2019). There is therefore an urgent need for new treatment strategies to fight against

resistant pathogens (Baym et al., 2016). Among the most promising alternatives, much attention

has focused on collateral sensitivity (CS). CS occurs when the acquisition of resistance to one

antibiotic causes increased susceptibility to a second antibiotic (Szybalski and Bryson, 1952) and

can be exploited for the design of multi-drug strategies that select against ABR (Imamovic and

Sommer, 2013; Lázár et al., 2013). Moreover, recent evidence suggests that the development of

resistance might be prevented by treatments based on the sequential cycling of antibiotics whose

resistance mechanisms produce reciprocal CS (Imamovic and Sommer, 2013). Several studies

have cataloged CS networks emerging from ABR mutations in chromosomal genes (Imamovic and

Sommer, 2013; Lázár et al., 2013; Maltas and Wood, 2019; Podnecky et al., 2018; Roemhild et al.,

2020), but the CS effects produced by the horizontal acquisition of ABR plasmids remain poorly

understood.

Mobile genetic elements, especially plasmids, play a crucial role in the dissemination of ABR

genes between clinical pathogens and are one of the major drivers of the alarming worldwide rise

in ABR (Partridge et al., 2018). Despite the evolutionary advantage conferred by plasmids in the

presence of antibiotics, plasmids acquisition tends to produce common metabolic alterations in the

host bacterium (San Millan et al., 2018), that typically translate into a fitness cost (San Millan and

MacLean, 2017). We reasoned that the physiological alterations produced upon acquisition of ABR

plasmids might lead to exploitable CS responses. To test this hypothesis, we analyzed the

collateral effects associated with the acquisition of 6 natural plasmids carrying clinically relevant

ABR genes in *Escherichia coli* MG1655. Our results reveal that most ABR plasmids, including the

clinically important carbapenem-resistance conjugative plasmid pOXA-48, produce CS events of

moderate effect. To extend our results, we explored the degree of conservation of CS responses

associated with pOXA-48 across phylogenetically diverse wild-type *E. coli.* Our results

demonstrate that these responses can be exploited to preferentially eradicate plasmid-carrying

bacteria.

Results and discussion

Collateral sensitivity induced by ABR plasmids

To test whether ABR plasmids elicit exploitable CS to other antibiotics, we selected 6 clinically

relevant ABR plasmids (see methods, Table 1 and Supplementary Figure 1). All plasmids carried

important ß-lactamase genes, including extended-spectrum ß-lactamases (ESBLs), AmpC-type ß-

lactamases, or carbapenemases, as well as other relevant ABR genes (Table 1). These plasmids

belonged to a broad range of incompatibility groups, covered a range of different sizes, included

conjugative and non-conjugative specimens, and produced variable fitness effects, ranging from

10% to 27% reductions in relative fitness, when introduced into *E. coli* MG1655 (Figure 1A).

We performed dose-response experiments with 13 antibiotics from 8 drug families (Supplementary

Table 1). For this analysis, we determined the minimal inhibitory concentration (MIC, see methods)

using the broth microdilution method for plasmid-free MG1655 and each of its 6 plasmid-carrying

derivatives (Supplementary Table 1). We analyzed the collateral antibiotic susceptibility effects

associated with the presence of each plasmid, measured as the fold-change in the antibiotic MIC

between plasmid-carrying and plasmid-free bacteria (Figure 1B). As expected from their genetic

content, all plasmids conferred resistance to ß-lactam antibiotics and most of them conferred

additional resistance to other antibiotics from unrelated families (Figure 1B and Supplementary

Table 1). CS was defined as a minimum 2-fold reduction in MIC (calculated as the fold-change of

the median MIC value obtained from 4-5 independent MIC determinations). Using this definition,

we identified 8 instances of plasmid-induced CS. The plasmids that produced CS effects were

pOXA-48 (to colistin and azithromycin), pKAZ3 (to kanamycin and gentamicin), pKA2Q (to

tetracycline), and pCF12 (to kanamycin, azithromycin, and tetracycline).

To validate the MIC results, we conducted two further series of experiments for the 8 cases of

plasmid-induced CS: (i) complete growth curves under a range of antibiotic concentrations and (ii)

disk-diffusion antibiograms on agar plates. Analysis of the area under the growth curve, which

integrates all growth parameters (maximum growth rate, lag duration and carrying capacity)

(DelaFuente et al., 2020), confirmed significant plasmid-mediated CS effects at different antibiotic

concentrations for 6 of the combinations (pOXA-48–azithromycin, pCF12–azithromycin, pOXA-48–

colistin, pCF12–kanamycin, pKAZ3–kanamycin, and pKA2Q–tetracycline; Figure 1C,

Supplementary Figure 2 and Supplementary Table 2). The disk-diffusion results largely agreed

with the MIC and growth-curve data, showing a general trend towards an increased diameter of the

antibiotic inhibition zone in plasmid-carrying strains compared with plasmid-free MG1655 (Figure

1D). These differences were statistically significant for pCF12–kanamycin, pCF12–azithromycin,

pOXA-48–azithromycin, pOXA-48–colistin, pCF12–tetracycline, and pKA2Q–tetracycline (unpaired

t-test with Welch's correction; $P<0.044$ in all cases). Overall, 7 of the 8 CS instances identified by

MIC testing were validated with at least one of the additional methods (growth curves or disk-

diffusion assay), and 6 of them showed full agreement across all 3 methods. The disk-diffusion

assays produced notably robust results with minimum hands-on time, suggesting that this

technique is appropriate for screening CS responses in large strain collections. Together, these

results demonstrate that ABR plasmids produce moderate but significant CS to different antibiotics,

comparable to that generated by ABR chromosomal mutations in *E. coli* (Imamovic and Sommer,

2013; Lázár et al., 2013; Podnecky et al., 2018).

Conservation of pOXA-48–mediated CS

The success and applicability of CS-informed therapeutic strategies are crucially dependent on the

conservation of CS across diverse genetic contexts (Podnecky et al., 2018). To address the

phylogenetic preservation of plasmid-induced CS, we focused on the plasmid pOXA-48, which

confers resistance to last-resort carbapenem antibiotics, and whose prevalence in clinical settings

is rising alarmingly (David et al., 2019). We tested the degree of conservation of pOXA-48–

mediated CS patterns (to azithromycin and colistin) across phylogenetically diverse *E. coli* strains

using disk-diffusion assays. We determined antibiotic susceptibility in 9 diverse *E. coli* clinical

isolates (Figure 2A) and their pOXA-48–carrying derivatives. CS to azithromycin showed a striking

degree of conservation across the strains, with 8 out of 9 plasmid-carriers showing significantly

higher susceptibility than their plasmid-free counterparts. In contrast, CS to colistin was conserved

in only 2 of the tested strains (Figure 2B). Nevertheless, all strains showed CS responses to at

least one antibiotic, suggesting that pOXA-48 elicits CS responses that could potentially be

exploited therapeutically.

Exploitation of CS for the preferential extinction of pOXA-48–carrying *E. coli*

We next tested the potential of exploiting CS effects to eradicate plasmid-carrying subpopulations.

We rationally selected 4 *E. coli* isolates and their pOXA-48 carrying derivatives with defined CS

profiles: to colistin only (Ec10), to azithromycin only (Ec21 and Ec25), or to both antibiotics (Ec18).

MG1655 was included as a control. Each bacterial population was propagated in the presence of a

single antibiotic or with sequential exposure to both antibiotics over 2 days. On the first day of the

experiment, 24 bacterial populations of each strain were inoculated into media containing either

colistin (4 mg/L) or azithromycin (8 mg/L). After growth for 22 hours, surviving populations were

transferred to fresh media containing colistin or azithromycin in a full factorial experimental design.

This approach gave rise to 4 antibiotic treatments: 2 single-drug treatments (denoted Azi→ Azi and

Col→ Col) and 2 treatments in which the antibiotic changed (Azi→ Col and Col→ Azi). In 11 of the

20 possible strain-treatment combinations, survival patterns showed significantly higher mortality rates for the pOXA-48–carrying strains than for their plasmid-free counterparts (Figure 3; log-rank test P<0.001 in all cases). Crucially, the mortality patterns observed were consistent with the CS patterns obtained in the disk-diffusion technique (Figure 2B). For instance, plasmid-carrying MG1655 and Ec18 strains, which show CS to both antibiotics, exhibited significant reductions in survival in all treatments. In contrast, for strain Ec10, which shows CS exclusively to colistin, significant extinction was observed only when the treatment included colistin as the first antibiotic (Col→ Col or Col→ Azi).

Concluding remarks

Our results reveal that the acquisition of clinically-relevant ABR plasmids induces CS to unrelated antibiotics. These findings thus serve as a stepping stone toward the development of new approaches aimed at blocking plasmid-mediated horizontal spread of ABR genes. These anti-plasmid strategies could help to tackle the alarming clinical and community spread of ABR. However, the molecular basis of plasmid-associated CS is currently unknown. The feasibility of rational broad-spectrum anti-plasmid strategies will depend on gaining comprehensive knowledge about the molecular mechanisms that increase antibiotic susceptibility upon plasmid acquisition. Until these therapies are available, CS-informed empirical treatment of plasmid-carrying bacteria has the potential to help resolve antibiotic resistant infections. In this regard, our results suggest that colistin and azithromycin may be valuable antibiotics for the treatment of pOXA-48–carrying enterobacteria. Both antibiotics are already used to treat enterobacterial infections (Lübbert, 2016; Morrill et al., 2015), and colistin in particular is currently used as a last resort antibiotic against carbapenem-resistant enterobacteria (Rodriguez-Baño et al., 2018). In summary, the findings presented here open new avenues of research aimed at understanding plasmid-associated CS and pave the way for the development of new CS-based treatment strategies to combat plasmid-mediated ABR.

Material and Methods

Bacterial strains, plasmids and culture conditions

All experiments were performed in liquid Lennox lysogeny broth (LB; CONDA) or on LB agar (15 g/L) unless indicated. Mueller Hinton II broth (Oxoid) was used to verify that MIC values of *E. coli* MG1655 were comparable with those obtained in LB medium. The antibiotics used were amoxicillin-clavulanic acid, ceftazidime, cefotaxime, gentamicin (Normon), ertapenem (Merck Sharp & Dohme), kanamycin, chloramphenicol, tetracycline (Nzytech), tigecycline (Pfizer), azithromycin, colistin (Altan Pharmaceuticals), and rifampicin (Sanofi-aventis).

Plasmids pKAZ3 (Accession Number KR827392.1), pKP-M1144 (KF745070.2), pKA2Q (MT720904) and pEncl-30969cz (MG049738.1) were previously published (Flach et al., 2015; Hernández-García et al., 2018a; Papagiannitsis et al., 2015, 2018). Plasmid pCEMR is a derivative of pEncl-30969cz in which a spontaneous deletion of ~19kb had occurred (MT720903). Plasmids pOXA-48_K8 (MT441554) and pCF12 (MT720906) were isolated from rectal swabs of patients hospitalized in the Ramón y Cajal University Hospital in the framework of the R-GNOSIS project (Hernández-García et al., 2018b). pOXA-48_K8 is a pOXA-48-like plasmid that shares a 100% coverage and >99% identity with the first described pOXA-48 plasmid (León-Sampedro et al., 2020; Poirel et al., 2012). For simplicity, we refer to this plasmid as pOXA-48 throughout the text. Plasmids pKAZ3, pOXA-48 and pCF12 were conjugated into MG1655 using *E. coli* β3914 as donor strain, which is auxotrophic for diaminopimelic acid (Le Roux et al., 2007). Transconjugants were selected in LB plates containing ampicillin (100 mg/L) for plasmids pKAZ3 and pOXA-48 or aztreonam (25 mg/L) for pCF12 plasmid. Plasmids pCEMR, pKP-M1144 and pKA2Q were purified using a commercial mini-prep kit (Macherey-Nagel) and transformed into calcium chloride competent MG1655 cells. Transformant cells were selected on LB-ampicillin plates (100 mg/L). Strain constructions were verified by assessing that the antibiotic resistance profile of plasmid-bearing MG1655 matched the expected resistance of the plasmid (Supplementary Table 1) and by sequencing the complete genomes of transconjugant clones, which also revealed that no significant chromosomal mutations accumulated during the process of plasmid acquisition (0-2

mutations between plasmid-carrying and plasmid-free complete genomes, see Supplementary

Table 3).

The clinical *E. coli* strains used in this study were isolated during the R-GNOSIS project, which

included representative ESBL-producing enterobacteria from hospitalized patients in the Ramon y

Cajal University Hospital during a 2 year period (Hernández-García et al., 2018b; León-Sampedro

et al., 2020). The *E. coli* strains used here, and their corresponding transconjugants, were

described and characterized previously (Alonso-del Valle et al., 2020) and Supplementary Table

4).

Antimicrobial susceptibility testing

Single colonies of plasmid-free and plasmid-carrying bacteria were inoculated in LB starter cultures

and incubated at 37 °C for 16 hours at 225 rpm (4-5 biological replicates performed on different

days). Each culture was diluted 1:2,000 in LB medium (~$5 \cdot 10^4$ colony forming units) and 200 µl

were added to a 96-well microtiter plate containing the appropriate antibiotic concentration. MIC

values were measured after 22 hours of incubation at 37 °C. Optical density at 600 nm (OD_{600}) was

determined in a Synergy HTX (BioTek) plate reader after 30 seconds of orbital shaking. MIC

values corresponded with the lowest antibiotic concentration that resulted in $OD_{600} < 0.2$. We found

that this threshold is qualitatively consistent with standard practice (i.e. visual determination of

growth inhibition), while providing higher accuracy and repeatability. The final MIC value is the

median of 4 to 5 replicates. MIC determinations of plasmid-free and plasmid-carrying strains were

performed in parallel to ensure reproducibility. Fold change in MIC was determined as the ratio

between the median MIC form plasmid-carrying derivatives and that form the wild-type strain.

Disk-diffusion susceptibility testing

LB plates were swabbed with a 0.5 McFarland matched bacterial suspension and sterile disks

containing the antibiotic were placed on the plates. Plates were incubated at 37 °C for 22 hours

and pictures were taken using an in-house photographic system and a cell phone (Huawei Mate

20). Inhibition halos were measured using ImageJ software. Antibiotic disk content used in

diffusion assays were as follows (all from Bio-Rad): gentamicin 10 µg, azithromycin 15 µg,

kanamycin 30 µg, colistin 10 µg, and tetracycline 30 µg.

Bacterial growth curves

Starter cultures were prepared and incubated as described above. Each culture was diluted

1:2,000 in LB medium and 200 µl were added to a 96-well microtiter plate containing the

appropriate antibiotic concentration. Plates were incubated 24 hours at 37 °C with strong orbital

shaking before reading OD_{600} every 10 minutes in a Synergy HTX (BioTek) plate reader. 6

biological replicates were performed for each growth curve. The area under the growth curve was

obtained using the '*auc*' function from the '*flux'* R package. Data was represented using a R

custom script and the '*ggplot2'* package.

Competition assays

Competition assays were performed to measure the relative fitness of plasmid-carrying strains

using flow cytometry as previously described (Rodriguez-Beltran et al., 2018). Briefly, competitions

were performed between MG1655 clones carrying each of the 6 plasmids and plasmid-free

MG1655 against a MG1655 derivative carrying an arabinose inducible chromosomal copy of *gfp*

(MG1655::*gfp*). Pre-cultures were incubated at 37 °C with 225 rpm shaking overnight in 96-well

plates carrying 200 µl of LB broth per well. Pre-cultures were mixed at 1:1 proportion and diluted

1:400 in fresh media. Initial proportions of GFP and non-fluorescent competitors were confirmed in

a CytoFLEX Platform (Beckman Coulter Life Sciences) flow cytometer, recording 10,000 events

per sample. To measure these proportions, we incubated a culture aliquot in NaCl 0.9% containing

0.5% L-arabinose for 1.5 hours to induce the expression of the chromosomal GFP. Mixtures were

competed for 22 hours in LB medium at 37 ºC with shaking (225 rpm). Final proportions were

estimated again by flow cytometry as described above. The fitness of each strain relative to

MG1655::*gfp* was calculated using the formula: $w = \ln(N_{final,gfp\text{-}}/N_{initial,gfp\text{-}})/\ln(N_{final,gfp\text{+}}/N_{initial,gfp\text{+}})$ where *w*

is the relative fitness of the non GFP-tagged strain, $N_{initial,gfp\text{-}}$ and $N_{final,gfp\text{-}}$ are the numbers of non

GFP-tagged cells before and after the competition and $N_{initial,gfp\text{+}}$ and $N_{final,gfp\text{+}}$ are the numbers of

MG1655::*gfp* cells before and after the competition. Data was normalized by dividing the relative

fitness of plasmid-carrying derivative strains by the fitness calculated for plasmid-free MG1655. Six

biological replicates were performed for each competition.

CS-informed antibiotic treatments

Overnight bacterial cultures were diluted and seeded into 24 independent wells of a 96-well plate

filled with 200 µl of LB. After 16 hours of incubation at 37 ºC, these populations were diluted

1:2,000 into fresh medium containing either azithromycin (8 mg/L) or colistin (4 mg/L) and grown at

37 ºC for 22 hours. The following day, bacterial populations were diluted (1:2,000) and inoculated

into new plates containing, again, either azithromycin or colistin and allowed to grow as above.

This approach led to four antibiotic treatments, two in which the antibiotic remains constant (Azi→

Azi and Col→ Col) and two in which the antibiotic treatment alternates (Azi→ Col and Col→ Azi).

The survival of bacterial populations was assessed by measuring OD_{600} every day. Populations

that did not reach an OD_{600} of 0.2 were declared extinct.

Whole genome sequencing

Genomic DNA was isolated using the Wizard genomic DNA purification kit (Promega) and

quantified using the QuantiFluor dsDNA system (Promega) following manufacturer's instructions.

Whole genome sequencing was conducted at the Wellcome Trust Centre for Human Genetics

(Oxford, UK), using the Illumina HiSeq4000 platform with 125 base pair (bp) paired-end reads. For

plasmid pCF12, first described in this study, we performed additional long-read sequencing using

PacBio technologies. PacBio sequencing was performed at The Norwegian Sequencing Centre

PacBio RSII platform using P5-C3 chemistry. Illumina and PacBio sequence reads were trimmed

using the Trimmomatic v0.33 tool (Bolger et al., 2014). SPAdes v3.9.0 was used to generate *de*

novo assemblies from the trimmed sequence reads with the –cov-cutoff flag set to 'auto'

(Bankevich et al., 2012). Unicycler was used to generate hybrid assemblies from Illumina and

PacBio data (Wick et al., 2017). QUAST v4.6.0 was used to generate assembly statistics (Gurevich

et al., 2013). All the *de novo* assemblies reached enough quality including total size of ~4.6 Mb,

and a total number of contigs over 1 kb lower than 200. Prokka was used to annotate the *de novo*

assemblies (Seemann, 2014). The plasmid content of each genome was characterised using

PlasmidFinder 2.1 (Carattoli et al., 2014), and the antibiotic resistance gene content was

characterised with ResFinder 3.2 (Zankari et al., 2012). MOB families were characterised using MOB-typer tool included in MOB-suite (Robertson and Nash, 2018). Variant calling was performed using Snippy v4.6.0 (https://github.com/tseemann/snippy). The sequences generated and analysed during the current study are available in the Sequence Read Archive (SRA) repository, BioProject ID: PRJNA644278 (https://www.ncbi.nlm.nih.gov/bioproject/644278), and Genbank accession numbers are provided in Table 1.

To determine the distribution of the *E. coli* isolates across the phylogeny of the species, we obtained 1334 assemblies of *Escherichia coli* complete genomes from the RefSeq database (https://www.ncbi.nlm.nih.gov/assembly). Distances between genomes were established using Mash v2.0 (Ondov et al., 2016) and a phylogeny was constructed with mashtree v0.33 (Katz et al., 2019). The tree was represented with midpoint root using the phylotools package in R (https://github.com/helixcn/) and visualised using the iTOL tool (Letunic and Bork, 2011). To determine the phylotype of each genome, we used the ClermonTyping tool (Beghain et al., 2018).

Statistical analyses

Data sets were analyzed using Prism 6 software (GraphPad Software Inc.) and R. Normality of the data was assessed by visual inspection and the Shapiro-Wilk test. ANOVA test were perfomed to ascertain the effect of the 'plasmid x antibiotic concentration' interaction term in the analysis of growth curves (area under the growth curve). Statistical analyses of the disk-diffusion halos were performed using unpaired t-tests with Welch's correction. Survival curves were analyzed using the log-rank test within the '*survminer*' R package.

Acknowledgments

We are grateful to thank Costas Papagiannitsis, Carl-Frederik Flach, Fabrice Elia Graf and Martin Palm for generous gifts of plasmids. We also appreciate the technical support of Carmen de la Vega and Laura Jaraba. This work was supported by the European Research Council under the European Union's Horizon 2020 research and innovation programme (ERC grant agreement no. 757440-PLASREVOLUTION) and by the *Instituto de Salud Carlos III* (grant PI16-00860) co-funded by European Development Regional Fund "a way to achieve Europe". ASM is supported by a

287 Miguel Servet Fellowship (MS15-00012). JRB is a recipient of a *Juan de la Cierva-Incorporación*

288 Fellowship (IJC2018-035146-I) co-funded by *Agencia Estatal de Investigación del Ministerio de*

289 *Ciencia e Innovación*. CH is supported by *Comunidad Autónoma de Madrid* (PEJD-2018-

290 POST/BMD-8016).

References

Alonso-del Valle A, León-Sampedro R, Rodríguez-Beltrán J, Hernández-García M, Ruiz-Garbajosa

P, Cantón R, San Millán Á, Spain M, Peña-Miller R. 2020. The distribution of plasmid fitness

effects explains plasmid persistence in bacterial. *bioRxiv* 2020.08.01.230672.

Baym M, Stone LK, Kishony R. 2016. Multidrug evolutionary strategies to reverse antibiotic

resistance. *Science (80-)* **351**:aad3292–aad3292. doi:10.1126/science.aad3292

Beghain J, Bridier-Nahmias A, Nagard H Le, Denamur E, Clermont O. 2018. ClermonTyping: An

easy-to-use and accurate in silico method for Escherichia genus strain phylotyping. *Microb*

Genomics **4**:1–8. doi:10.1099/mgen.0.000192

Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data.

Bioinformatics **30**:2114–2120. doi:10.1093/bioinformatics/btu170

Carattoli A, Zankari E, García-Fernández A, Larsen MV, Lund O, Villa L, Aarestrup FM, Hasman H.

2014. In Silico Detection and Typing of Plasmids using PlasmidFinder and Plasmid Multilocus

Sequence Typing. *Antimicrob Agents Chemother* **58**. doi:10.1128/AAC.02412-14

David S, Reuter S, Harris SR, Glasner C, Feltwell T, Argimon S, Abudahab K, Goater R, Giani T,

Errico G, Aspbury M, Sjunnebo S. 2019. Epidemic of carbapenem-resistant Klebsiella

pneumoniae in Europe is driven by nosocomial spread. *Nature* **4**:1919–1929.

doi:10.1038/s41564-019-0492-8.Epidemic

DelaFuente J, Rodriguez-Beltran J, San Millan A. 2020. Methods to Study Fitness and

Compensatory Adaptation in Plasmid-Carrying BacteriaHorizontal Gene Transfer. pp. 371–

382. doi:10.1007/978-1-4939-9877-7_26

Flach CF, Johnning A, Nilsson I, Smalla K, Kristiansson E, Larsson DGJ. 2015. Isolation of novel

IncA/C and IncN fluoroquinolone resistance plasmids from an antibiotic-polluted lake. *J*

Antimicrob Chemother **70**:2709–2717. doi:10.1093/jac/dkv167

Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. Genome analysis QUAST: quality assessment

tool for genome assemblies. *Bioinformatics* **29**:1072–1075. doi:10.1093/bioinformatics/btt086

Hernández-García M, León-Sampedro R, Pérez-Viso B, Morosini MI, López-Fresneña N, Díaz-

Agero C, Coque TM, Ruiz-Garbajosa P, Cantón R. 2018a. First report of an OXA-48- And

CTX-M-213-producing Kluyvera species clone recovered from patients admitted in a

university hospital in Madrid, Spain. *Antimicrob Agents Chemother* **62**.

doi:10.1128/AAC.01238-18

Hernández-García M, Pérez-Viso B, Carmen Turrientes M, Díaz-Agero C, López-Fresneña N,

Bonten M, Malhotra-Kumar S, Ruiz-Garbajosa P, Cantón R. 2018b. Characterization of

carbapenemase-producing Enterobacteriaceae from colonized patients in a university hospital

in Madrid, Spain, during the R-GNOSIS project depicts increased clonal diversity over time

with maintenance of high-risk clones. *J Antimicrob Chemother* **73**:3039–3043.

doi:10.1093/jac/dky284

Imamovic L, Sommer MOA. 2013. Use of collateral sensitivity networks to design drug cycling

protocols that avoid resistance development. *Sci Transl Med* **5**:1–10.

doi:10.1126/scitranslmed.3006609

Jim O. 2016. Tackling drug-resistant infections globally: final report and recommnedations. The

review on antimicrobial resistance.

Katz L, Griswold T, Morrison S, Caravas J, Zhang S, Bakker H, Deng X, Carleton H. 2019.

Mashtree: a rapid comparison of whole genome sequence files. *J Open Source Softw* **4**:1762.

doi:10.21105/joss.01762

Lázár V, Pal Singh G, Spohn R, Nagy I, Horváth B, Hrtyan M, Busa-Fekete R, Bogos B, Méhi O,
Csörgo B, Pósfai G, Fekete G, Szappanos B, Kégl B, Papp B, Pál C. 2013. Bacterial evolution
of antibiotic hypersensitivity. *Mol Syst Biol* **9**. doi:10.1038/msb.2013.57

Le Roux F, Binesse J, Saulnier D, Mazel D. 2007. Construction of a Vibrio splendidus Mutant
Lacking the Metalloprotease Gene vsm by Use of a Novel Counterselectable Suicide Vector
†. *Appl Environ Microbiol* **73**:777–784. doi:10.1128/AEM.02147-06

León-Sampedro R, DelaFuente J, Díaz-Agero C, Musicha P, Rodriguez-Beltran J, de la Vega C,
Hernández-García M, Group S, Lopez-Fresneña N, Ruiz-Garbajosa P, Canton R, Cooper BS,
San A, Behnke M, Blok H, Brun-Buisson C, Fankhauser C, Gastmeier P, Goossens H,
Hansen S, Harbarth S, Huttner B, Kloosterman F, Kola A, Kuperberg A, Lammens C, Leus F,
Maechler F, Malhotra S, Schotsman J, Schwab F, Triay C. 2020. Dissemination routes of the
carbapenem resistance plasmid pOXA-48 in a hospital setting. *bioRxiv 20200420050476*.
doi:10.1101/2020.04.20.050476

Letunic I, Bork P. 2011. Interactive Tree of Life v2: Online annotation and display of phylogenetic
trees made easy. *Nucleic Acids Res* **39**:475–478. doi:10.1093/nar/gkr201

Lübbert C. 2016. Antimicrobial therapy of acute diarrhoea: A clinical review. *Expert Rev Anti Infect*
Ther **14**:193–206. doi:10.1586/14787210.2016.1128824

MacLean R, San Millan A. 2019. The Evolution of Antibiotic Resistance. *Science (80-)* **365**.
doi:10.1126/SCIENCE.AAX3879

Maltas J, Wood KB. 2019. Pervasive and diverse collateral sensitivity profiles inform optimal
strategies to limit antibiotic resistance. *Plos Biol* **17**.
doi:https://doi.org/10.1371/journal.pbio.3000515 October

Morrill HJ, Pogue JM, Kaye KS, LaPlante KL. 2015. Treatment Options for Carbapenem-Resistant
Enterobacteriaceae Infections. *Open forum infentious diesases* **2**. doi:DOI: 10.1093/ofid/ofv05

Ondov BD, Treangen TJ, Melsted P, Mallonee AB, Bergman NH, Koren S, Phillippy AM. 2016.

Mash: Fast genome and metagenome distance estimation using MinHash. *Genome Biol* **17**:1–14. doi:10.1186/s13059-016-0997-x

Papagiannitsis CC, Dolejska M, Izdebski R, Dobiasova H, Studentova V, Esteves FJ, Derde LPG, Bonten MJM, Hrabák J, Gniadkowski M. 2015. Characterization of pKP-M1144, a Novel ColE1-Like Plasmid Encoding IMP-8, GES-5, and BEL-1-Lactamases, from a Klebsiella pneumoniae Sequence Type 252 Isolate. doi:10.1128/AAC.00937-15

Papagiannitsis CC, Paskova V, Chudejova K, Medvecky M, Bitar I, Jakubu V, Zemlickova H, Jirsa R, Hrabak J. 2018. Characterization of pEncl-30969cz, a novel ColE1-like plasmid encoding VIM-1 carbapenemase, from an Enterobacter cloacae sequence type 92 isolate. *Diagn Microbiol Infect Dis* **91**:191–193. doi:10.1016/J.DIAGMICROBIO.2018.01.024

Partridge SR, Kwong SM, Firth N, Jensen SO. 2018. Mobile Genetic Elements Associated with Antimicrobial Resistance. *Clin Microbiol Rev* **31**.

Podnecky NL, Fredheim EGA, Kloos J, Sørum V, Primicerio R, Roberts AP, Rozen DE, Samuelsen Ø, Johnsen PJ. 2018. Conserved collateral antibiotic susceptibility networks in diverse clinical strains of Escherichia coli. *Nat Commun* **9**:3673. doi:10.1038/s41467-018-06143-y

Poirel L, Bonnin RA, Nordmann P. 2012. Genetic Features of the Widespread Plasmid Coding for the Carbapenemase OXA-48. *Antimicrob Agents Chemother* **56**:559–562. doi:10.1128/AAC.05289-11

Robertson J, Nash JHE. 2018. MOB-suite: software tools for clustering, reconstruction and typing of plasmids from draft assemblies. *Microb genomics* **4**. doi:10.6084/m9.figshare.6177188

Rodriguez-Baño J, Gutiérrez-Gutiérrez B, Machuca I, Pascual A. 2018. Treatment of Infections Caused by Extended-Spectrum-Beta-. *Clin Microbiol Rev* **31**:1–42.

Rodriguez-Beltran J, Hernandez-Beltran JCR, DelaFuente J, Escudero JA, Fuentes-Hernandez A, MacLean RC, Peña-Miller R, San Millan A. 2018. Multicopy plasmids allow bacteria to escape from fitness trade-offs during evolutionary innovation. *Nat Ecol Evol* **2**:873–881.

doi:10.1038/s41559-018-0529-z

Roemhild R, Linkevicius M, Andersson DI. 2020. Molecular mechanisms of collateral sensitivity to

the antibiotic nitrofurantoin. *PLOS Biol* **18**:e3000612. doi:10.1371/journal.pbio.3000612

San Millan A, MacLean RC. 2017. Fitness Costs of Plasmids: a Limit to Plasmid Transmission.

Microbiol Spectr **5**:1–12. doi:10.1128/microbiolspec.mtbp-0016-2017

San Millan A, Toll-Riera M, Qi Q, Betts A, Hopkinson RJ, McCullagh J, MacLean RC. 2018.

Integrative analysis of fitness and metabolic effects of plasmids in Pseudomonas aeruginosa

PAO1. *ISME J* **12**:3014–3024. doi:10.1038/s41396-018-0224-8

Seemann T. 2014. Genome analysis Prokka: rapid prokaryotic genome annotation. *Bioinformatics*

30:2068–2069. doi:10.1093/bioinformatics/btu153

Szybalski W, Bryson V. 1952. Genetic studies on microbial cross resistance to toxic agents. I.

Cross resistance of Escherichia coli to fifteen antibiotics. *J Bacteriol* **64**:489–499.

Wick RR, Judd LM, Gorrie CL, Holt KE. 2017. Unicycler: Resolving bacterial genome assemblies

from   short   and   long   sequencing   reads.   *PLoS   Comput   Biol*   **13**:1–22.

doi:10.1371/journal.pcbi.1005595

Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, Aarestrup FM, Larsen

MV. 2012. Identification of acquired antimicrobial resistance genes. *J Antimicrob Chemother*

67:2640–2644. doi:10.1093/jac/dks261

 Figures and Tables

 Figure 1. Collateral sensitivity and fitness effects associated with ABR plasmids. (**A**)

Competition-assay–determined fitness of plasmid-carrying clones relative to the plasmid-free

MG1655 strain. Bars represent the means of 6 independent experiments (represented by dots),

and error bars represent the standard error of the mean. (**B**) Heat-map representing collateral

responses to antibiotics associated with plasmid acquisition. The color code represents the fold

change of MIC in plasmid-carrying derivatives compared to plasmid-free MG1655 (see legend);

MIC values were the median of 4-5 independent determinations. Asterisks indicate a ≥ 2-fold

decrease in MIC, which we defined as significant CS. AMC, amoxicillin-clavulanic acid; CTX,

cefotaxime; CAZ, ceftazidime; ERT, ertapenem; GM, gentamicin; KM, kanamycin; TET,

tetracycline; TGC, tigecycline; CM, chloramphenicol; AZI, azithromycin; CIP, ciprofloxacin; RIF,

rifampicin; and COL, colistin. (**C**) Growth curves of MG1655 and plasmid-carrying MG1655 strains

exposed to increasing antibiotic concentrations. The antibiotic concentration, in mg/L, is indicated

at the top left corner of each panel. Lines represent the mean of 6 biological replicates, and the

shaded area indicates standard error of the mean. (**D**) Boxplot representations of the inhibition halo

diameters, in mm, obtained from disk-diffusion antibiograms of plasmid-free and plasmid-carrying

MG1655. Horizontal lines within boxes indicate median values, upper and lower hinges correspond

to the 25th and 75th percentiles, and whiskers extend to observations within 1.5 times the

interquartile range. Individual data points are also represented (11-15 biological replicates).

Asterisks indicate statistically significant differences (unpaired t-test with Welch's correction $P<$

0.044).

Table 1. Plasmids used in this study.

Plasmid	Incompatibility group	Mobility (MOB family)	Size (bp)	Resistance Genes	Reference (Genbank Ac. No.)
pOXA-48_K8	IncL	Conjugative (MOBP)	65,499	bla_{OXA-48}	(León-Sampedro et al., 2020) (MT441554)
pKAZ3	IncA/C	Conjugative (MOBH)	147,957	bla_{VEB-9}, *qnrVC-1*, *sul1*, *tetA'*, *tetC*, *dfrA1*, *dfrA23*	(Flach et al., 2015)(KR827392.1)
pKA2Q	IncQ	Mobilizable (MOBF)	8,789	bla_{FOX-8}	(Hernández-García et al., 2018a) (MT720904)
pCF12	IncX3	Conjugative (MOBP)	43,704	bla_{SHV-12}, *qnrS1*	This work (MT720906)
pCEMR	ColE-like	Mobilizable	14,478	bla_{VIM-1}, *aac(6')-Ib3 sul1*, *aadA1*, *qacEΔ1*	(Papagiannitsis et al., 2018) (MG049738.1)
pKP-M1144	ColE-like	Non-transmissible	12,029	bla_{BEL-1}, bla_{GES-5}, bla_{IMP-8}, *aac(6')-Ib3*	(Papagiannitsis et al., 2015) (KF745070.2)

[1] pOXA-48_K8 is a pOXA-48-like plasmid (see methods and Supplementary Figure 1). For simplicity, we refer to this plasmid as pOXA-48 throughout the study.

Figure 2. Phylogeny and collateral sensitivity profiles of *E. coli* clinical isolates. (**A**) Phylogenetic tree of *E. coli* species, highlighting the strains used in this study and their sequence types (ST). Colors within tree branches represent different STs. The tree depicts the phylogenetic relationships of 1344 representative *E. coli* genomes obtained from NCBI. (**B**) Representation of the inhibition halo diameters, in mm, obtained from disk-diffusion antibiograms of clinical isolates and their transconjugants carrying pOXA-48. Horizontal lines inside boxes indicate median values, upper and lower hinges correspond to the 25th and 75th percentiles, and whiskers extend to observations within 1.5 times the interquartile range. Individual data points are also represented (6-12 replicates, jittered to facilitate data visualization). Asterisks denote statistically significant differences (t-test with Welch's correction P<0.035 in all cases).

Figure 3. Plasmid-mediated CS can be exploited to preferentially eliminate plasmid-carrying bacteria. Survival curves of plasmid-free (gray solid lines) and plasmid-carrying (red dashed lines) bacterial populations in response to constant or sequential antibiotic treatments. Bacteria were cultured for 2 days with either constant antibiotic exposure or with a change of antibiotic after the first day; treatments are specified by the background colors in each panel. Log-rank test p-values assessing statistically different survival rates are shown. n.s. non-significant. For strains Ec10, Ec18, Ec21 and Ec25 n=24; for MG1665 n=48. AZI, azithromycin; COL, colistin.

 Supplementary Material

 Supplementary Figure 1. Plasmids used in this study.

Scheme depicting the 6 plasmids used in this study. Antibiotic resistance genes are labeled and

highlighted in yellow. Accession numbers are as follows: pKAZ3 (MT720905), pOXA-48_K8

(MT441554), pCF12 (MT720906), pCEMR (MT720903), pKP-1144 (MT720902), pKA2Q

(MT720904). pOXA-48_K8 is a pOXA-48-like plasmid that shares a 100% coverage and >99%

identity with the first described pOXA-48 plasmid (Poirel et al., 2012). For simplicity we refer to this

plasmid as pOXA-48 throughout the text. Plasmids are not drawn to scale.

Supplementary Figure 2. Analysis of bacterial growth curves

Comparison of the relative area under the growth curve of MG1655 (gray lines) and its plasmid-

carrying derivatives (red) under different antibiotic concentrations. Each data point corresponds to

the mean of 6-12 biological replicates and error bars correspond to the standard error of the mean.

Individual data points are also represented. P-values are provided for the significantly different

comparisons (ANOVA effect of 'plasmid x antibiotic concentration' interaction; ANOVA results are

shown in Supplementary Table 2). AZI: azithromycin, COL: colistin, GM: gentamycin, KM:

kanamycin, TET: tetracycline.

474 **Supplementary Table 1. Antibiotic susceptibility results**

| Drug class/target | Antibiotics | Minimal inhibitory concentration (mg/L) | | MG1655/ pOXA-48 | MG1655/ pKAZ3 | MG1655/ pKA2Q | MG1655/ pCF12 | MG1655/ pCEMR | MG1655/ pKP-M1144 |
| | | MG1655 | | | | | | | |
		MH	LB						
B-lactam/ Cell wall synthesis	Amoxicillin-clavulanic acid (AMC)	6	16	>512	32	256	128	512	384
	Cefotaxime (CTX)	0.03	0.375	8	8	8	8	8	8
	Ceftazidime (CAZ)	0.5	0.375	0.5	32	32	32	32	32
	Ertapenem (ERT)	0.01	0.062	0.187	0.507	0.125	0.125	0.25	0.75
Aminoglycoside/ Protein synthesis (30S)	Gentamicin (GM)	2	6	6	2	4	4	8	8
	Kanamycin (KM)	8	12	24	6	12	4	288	24
Tetracycline/ Protein synthesis (30S)	Tetracycline (TET)	2	4	4	32	2	2	4	4
	Tigecycline (TGC)	0.5	1	1	1	1	1	1	1
Amphenicol/ Protein synthesis (50S)	Chloramphenicol (CM)	8	8	16	8	8	8	8	8
Macrolide/ Protein synthesis (50S)	Azithromycin (AZI)	16	16	8	12	12	8	12	12
Quinolone/	Ciprofloxacin	0.5	0.5	0.75	8	1	8	2	3

DNA gyrase	(CIP)								
Rifamycin/ RNA polymerase	Rifampicin (RIF)	8	8	8	24	24	8	16	256
Polymyxin/ Lipopolysaccharide	Colistin (COL)	4	8	4	6	8	6	8	6

475 MIC of the 13 antibiotics for *E.* coli MG1655 in LB and MH medium and the MIC of the six plasmid-

476 carrying MG1655 strains in LB medium. Median value of 4-5 independent biological replicates.

478 **Supplementary Table 2. ANOVA results of the growth curves of plasmids producing CS**

479 **effects**

Area under the growth curve

Plasmid/Antibiotic	P-val	df	F
pCF12/AZI	<0.000001	3	46.435
pCF12/KM	0.000008	3	8.859
pCF12/TET	<0.000001	3	12.323
pKA2Q/TET	0.460776	3	0.878
pKAZ3/GM	0.264279	3	1.368
pKAZ3/KM	0.025920	3	3.383
pOXA-48/AZI	<0.000001	3	12.664
pOXA-48/COL	<0.000001	3	5.9618

480 Results of the 'plasmid x antibiotic concentration' interaction term of ANOVA analyses for each

481 plasmid-antibiotic combination.

482 **Supplementary Table 3. Chromosomal mutations accumulated during plasmid acquisition.**

Strain	Gene	Gene product	Type	Position[A]	Reference[B]	Alternative	Effect
MG1655/pKP-M1144	*gadB*	Glutamate decarboxylase beta	SNP	642/1401	G	A	Synonymous (Thr214Thr)
MG1655/pKP-M1144	*selB*	Selenocysteine-specific elongation factor	SNP	1256/1845	G	T	Missense (Trp419Leu)
MG1655/pCEMR	*ppk*	Polyphosphate kinase	SNP	1488/2067	T	C	Synonymous (Arg496Arg)
MG1655/pCEMR	*selB*	Selenocysteine-specific elongation factor	SNP	1256/1845	G	T	Missense (Trp419Leu)
MG1655/pKA2Q	*cpxA*	Sensor histidine kinase	SNP	515/1374	A	G	Missense (Met172Thr)
MG1655/pCF12	Intergenic	-	Insertion	1,212,080	AGGGGGGGGGT	AGGGGGGGGGGT	-

483 [A]Position relative to the start of the open reading frame, except for the intergenic mutation which

484 refers to the chromosomal coordinate. [B]Reference genome is plasmid-free MG1655 (deposited in

485 Sequence Read Archive [SRA], BioProject ID: PRJNA644278)

487 **Supplementary Table 4. Characteristics of the *E. coli* clinical isolates used in this study.**

Strain	Phylogroup	Sequence Type	Plasmids	Resistance genes
Ec02	B1	453	Col-like, IncFIB, IncI, IncFIC, IncY	*aadA5, blaCTX-M-1, blaTEM-1B, dfrA17, sul2, tet(B), mdf(A)*
Ec03	B2	131	Col-like, IncFIB, IncFII, IncB/O/K/Z	*blaCTX-M-14, blaTEM-1B, mdf(A)*
Ec04	B2	131	Col-like, IncFIB, IncFII, IncFIA	*aadA5, aph(3'')-Ib, aph(6)-Id, blaCTX-M-14, blaTEM-1B, dfrA17, mph(A), sul1, sul2, tet(A), mdf(A)*
Ec06	A	10	Col-like, IncFII, IncI	*aadA5, blaCTX-M-1, sul2, tet(A), mdf(A)*
Ec10	D	69	Col-like, IncFII, pEC4115	*aac(3)-Iva, aph(3'')-Ib, aph(4)-Ia, aph(6)-Id, blaCTX-M-1, mph(A), mdf(A)*
Ec18	F	648	Col-like, IncY	*aadA5, blaCTX-M-14, dfrA17, mph(A), sul1, mdf(A)*
Ec20	B2	131	IncFIB, IncI, IncFIA, IncFIC	*aac(3)-Iid, blaCTX-M-15, blaTEM-1B, mdf(A)*
Ec21	D	405	Col-like, IncFIB, IncFII, IncFIA, p0111	*aac(3)-Iia, aph(3'')-Ib, aph(6)-Id, blaCTX-M-15, floR, sul2, tet(A), tet(B), mdf(A)*
Ec25	A	10	IncFIB, IncFII, IncX, IncFIC	*aph(3')-Ia, aph(6)-Id, blaCTX-M-3, dfrA14, fosA3, sul2, mdf(A)*