The spatial proteome of the human intervertebral disc reveals architectural changes in health, ageing and degeneration

Vivian Tam[1,2*], Peikai Chen[1*], Anita Yee[1], Nestor Solis[3], Theo Klein[3,4],

Mateusz Kudelko[1], Rakesh Sharma[5], Wilson CW Chan[1,2,6], Christopher M. Overall[3],

Lisbet Haglund[7], Pak C Sham[8], Kathryn SE Cheah[1], Danny Chan[1,2]

[1] School of Biomedical Sciences, The University of Hong Kong, Hong Kong, China;

[2] The University of Hong Kong Shenzhen Institute of Research and Innovation, Shenzhen, China;

[3] Centre for Blood Research, Faculty of Dentistry, University of British Columbia, Vancouver, Canada;

[4] Present address: Triskelion BV, Zeist, The Netherlands;

[5] Proteomics and Metabolomics Core Facility, The University of Hong Kong, Hong Kong, China;

[6] Department of Orthopaedics Surgery and Traumatology, HKU-Shenzhen Hospital, Shenzhen, China;

[7] Department of Surgery, McGill University, Montreal, Canada;

[8] Centre for PanorOmic Sciences (CPOS), The University of Hong Kong, Hong Kong, China.

KSEC is a senior editor at eLife. All other authors declare that they do not have any conflict of interests

* These authors contributed equally to this manuscript

Correspondence should be addressed to:

Danny Chan
School of Biomedical Sciences
Faculty of Medicine
The University of Hong Kong
21 Sassoon Road, Hong Kong
Email: chand@hku.hk
Tel.: +852 3917 9482

Abstract

The proteome of the human intervertebral disc, particularly the extracellular matrix, underpins its integrity and function. We describe high quality spatial proteomes from young and aged lumbar discs, profiled across different regions, and identified sets of proteins that are variable or constant with respect to disc compartments, levels, and anatomic directions. From these, four protein modules defining a young disc were derived. Mapping these modules to aged disc proteomes showed that the central nucleus pulposus becomes more like the outer annulus fibrosus. Transcriptome analysis supported these proteomic findings. Concurrently, dynamic proteomic analyses indicate impaired protein synthesis and increased degradative activities. We identified a novel set of proteins that correlates with disc hydration that are predictive of MRI intensities. Signalling pathways (SHH, WNT, BMP/TGFβ) and hypertrophic chondrocyte differentiation events relating to ageing and degeneration were implicated. This work presents the foundation proteomic architectural landscapes of young and ageing human discs.

(149 words)

Key words: human, intervertebral discs, nucleus pulposus, annulus fibrosus, ageing, extracellular matrix, proteomics, TAILS, degradome, SILAC, transcriptomics, hypertrophy-like

Introduction

The 23 intervertebral discs (IVDs) in the human spine provide stability, mobility and flexibility. IVD degeneration (IDD), most common in the lumbar region (Saleem et al., 2013; Teraguchi et al., 2014), is associated with a decline in function and a major cause of back pain, affecting up to 80% of the world's population at some point in life (Rubin, 2007), presenting significant socioeconomic burdens. Multiple interacting factors such as genetics, influenced by ageing, mechanical and other stress factors, contribute to the pathobiology, onset, severity and progression of IDD (Munir et al., 2018).

IVDs are large, avascular, extracellular matrix (ECM)-rich structures comprising three compartments: a hydrated nucleus pulposus (NP) at the centre, surrounded by a tough annulus fibrosus (AF) at the periphery, and cartilaginous endplates of the adjoining vertebral bodies (Humzah and Soames, 1988). The early adolescent NP is populated with vacuolated notochordal-like cells, which are gradually replaced by small chondrocyte-like cells (Risbud et al., 2015). Blood vessels terminate at the endplates, nourishing and oxygenating the NP via diffusion, whose limited capacity mean that NP cells are constantly subject to hypoxic, metabolic and mechanical stresses (Urban et al., 2004).

With ageing and degeneration, there is an overall decline in cell "health" and numbers (Rodriguez et al., 2011; Sakai et al., 2012), disrupting homeostasis of the disc proteome. The ECM has key roles in biomechanical function and disc hydration. Indeed, a hallmark of IDD is reduced hydration in the NP, diminishing the disc's capacity to dissipate mechanical loads. Clinically, T-2 weighted magnetic resonance imaging (MRI) is the gold standard for assessing IDD, that uses disc hydration and structural features such as bulging or annular tears to measure severity (Pfirrmann et al., 2001; Schneiderman et al., 1987). The hydration and mechanical properties of the IVD are dictated by the ECM composition, which is produced and maintained by the IVD cells.

To meet specific biomechanical needs, cells in the IVD compartments synthesise different compositions of ECM proteins. Defined as the "matrisome" (Naba et al., 2012), the ECM houses the cells and facilitates their inter-communication by regulation of the availability and presentation of signalling molecules (Taha and Naba, 2019). With ageing or degeneration, the NP becomes more fibrotic and less compliant (Yee et al., 2016), ultimately affecting disc biomechanics (Newell

et al., 2017). Changes in matrix stiffness can have a profound impact on cell-matrix interactions
and downstream transcriptional regulation, signalling activity and cell fate (Park et al., 2011).

The resulting alterations in the matrisome lead to vicious feedback cycles that reinforce cellular
degeneration and ECM changes. Notably, many of the associated IDD genetic risk factors, such
as COL9A1 (Jim et al., 2005), ASPN (Song et al., 2008), and CHST3 (Song et al., 2013), are
variants in genes encoding matrisome proteins, highlighting their importance for disc function.
Therefore, knowledge of the cellular and extracellular proteome and their spatial distribution in
the IVD is crucial to understanding the mechanisms underlying the onset and progression of IDD
(Feng et al., 2006).

Current knowledge of IVD biology is inferred from a limited number of transcriptomic studies on
human (Minogue et al., 2010; Riester et al., 2018; Rutges et al., 2010) and animal (Veras et al.,
2020) discs. Studies showed that cells in young healthy NP express markers including CD24,
KRT8, KRT19 and T (Fujita et al., 2005; Minogue et al., 2010; Rutges et al., 2010), whilst NP
cells in aged or degenerated discs have different and variable molecular signatures (Chen et al.,
2006; Rodrigues-Pinto et al., 2016), such as genes involved in TGFβ signalling (TGFA, INHA,
INHBA, BMP2/6). The healthy AF expresses genes including collagens (COL1A1 and COL12A1)
(van den Akker et al., 2017), growth factors (PDGFB, FGF9, VEGFC) and signalling molecules
(NOTCH and WNT) (Riester et al., 2018). Although transcriptomic data provides valuable cellular
information, it does not faithfully reflect the molecular composition. Cells represent only a small
fraction of the disc volume, transcriptome-proteome discordance does not enable accurate
predictions of protein levels from mRNA (Fortelny et al., 2017), and the disc matrisome
accumulates and remodels over time.

Proteomic studies on animal models of IDD, including murine (McCann et al., 2015), canine
(Erwin et al., 2015), and bovine (Caldeira et al., 2017), have been reported. Nevertheless, human-
animal differences in cellular phenotypes and mechanical loading physiologies mean that these
findings might not translate to the human scenario. So far, human proteomic studies have
compared IVDs with other cartilaginous tissues (Onnerfjord et al., 2012a); and have shown
increases in fibrotic changes in ageing and degeneration (Yee et al., 2016), a role for inflammation
in degenerated discs (Rajasekaran et al., 2020), the presence of haemoglobins and

immunoglobulins in discs with spondylolisthesis and herniation (Maseda et al., 2016), and changes
in proteins related to cell adhesion and migration in IDD (Sarath Babu et al., 2016). The reported
human disc proteomes were limited in the numbers of proteins identified and finer
compartmentalisation within the IVD, and disc levels along the lumbar spine have yet to be
studied. Nor have the proteome dynamics in term of ECM remodelling (synthesis and degradation)
in young human IVDs and changes in ageing and degeneration been described.

In this study, we established a reference spatial proteome of young healthy IVDs based on 33 LC-
MS/MS (liquid chromatography mass spectrometry) profiles generated from a systematic division
of the different compartments of three whole lower lumbar discs from a 16-year-old male non-
degenerated individual. To gain insights into how the IVD proteomic landscape changes with
ageing, we analysed 33 LC-MS/MS profiles from IVDs isolated from a 59-year-old male
individual at the corresponding disc levels. We evaluated differences and similarities in the
proteome of these disc compartments, by principal component analysis (PCA), analysis of variance
(ANOVA) and identification of differentially expressed proteins (DEPs). The proteomic findings
were consolidated with 8 transcriptome profiles from 4 individuals.

We discovered modules containing specific sets of proteins that describe the landscape of a young
IVD, and the deconstruction of these modules with ageing and degeneration. We analysed the
dynamic changes in the disc proteome by SILAC (Stable Isotopic Labelling of Amino Acids in
Culture) (Ong et al., 2002) on the IVDs of 5 individuals, and by TAILS (Terminal Amine Isotopic
Labelling of Substrates) (Kleifeld et al., 2010) on 6 individuals, to determine synthesis and
degradation status, and their relationships with disc compartment and IDD. We confirmed the
presence of a basal ECM remodelling process in the NP of young IVDs that increases with ageing
and degeneration. Finally, we correlated types of proteins with tissue hydration using high
resolution MRI of the aged discs, to derive a hydration signature predictive of MRI intensities.

Results

Disc samples and their phenotypes

Lumbar disc components from 17 individuals (Table 1) were harvested for proteomic, transcriptomic, SILAC and degradome (Lopez-Otin and Overall, 2002) analyses. Reference proteomes were generated from lumbar disc segments of a young (16 years) and an aged (59 years) individual. T1- and T2-weighted MRI (3T) showed the young discs (L3/4, L4/5, L5/S1) were non-degenerated, with a Schneiderman score of 1 from T2 images (Figure 1A). The NP of young IVD were well hydrated (white) with no disc bulging, endplate changes, or observable inter-level variations (Figure 1A; Supplemental Figure S1A), consistent with healthy discs and were deemed fit to serve as a benchmarking reference. To investigate structural changes associated with ageing, high resolution (7T) MRI was taken for the aged discs (Figure 1B). All discs had irregular endplates, and annular tears were present (green arrowheads) adjacent to the lower endplate and extending towards the posterior region at L3/4 and L4/5 (Figure 1B). The NP exhibited regional variations in hydration in both sagittal and coronal images (Figure 1B). As the young and aged MRIs were obtained at different resolutions, direct comparison is difficult. However, the aged discs were less hydrated and the NP and AF structures less distinct, consistent with gross observations (Supplemental Figure S1A). Phenotypes of the disc samples used in the generation of other profiling data are described in methods and Table 1.

Quality data detecting large numbers of matrisome and non-matrisome proteins

To derive spatial reference proteomes for young and aged human IVDs, we subdivided the lumbar discs into 11 key regions (Figure 1C), spanning the outer-most (outer AF; OAF) to the central-most (NP) region of the disc, traversing both anteroposterior and lateral axes, adding valuable spatial information to our proteomic dataset. Since the disc is an oval shape, an inner AF (IAF) region was assigned in the lateral directions. A "mixed" compartment between the NP and IAF with undefined boundary was designated as the NP/IAF in all four (anteroposterior and lateral) directions. In all, this amounted to 66 specimens with different compartments, ages, directions and levels, which then underwent LC-MS/MS profiling (Supplemental Table S1). Systematic analyses of the 66 profiles are depicted in a flowchart (Figure 1D).

A median of 654 proteins per profile were identified for the young samples and 829 proteins for the aged samples, with a median of 742 proteins per profile for young and aged combined. The proteome-wide distributions were on similar scales across the profiles (Supplemental Figure S1B). Of the 3,100 proteins detected in total, 418 were matrisome proteins (40.7% of all known matrisome proteins) and 2,682 non-matrisome proteins (~14% of genome-wide non-matrisome genes) (Figure 1E; Supplemental Figure S1C), and 983 were common to all four major compartments, namely the OAF, IAF, NP/IAF, and NP (Figure 1E, upper panel). A total of 1,883 proteins were identified in young discs, of which 690 (36%) were common to all regions. Additionally, 45 proteins (2.4%) were unique to NP, whilst NP/IAF, IAF and OAF had 86 (4.6%), 54 (2.9%) and 536 (28%) unique proteins, respectively (Figure 1E, middle panel). For the aged discs, 2,791 proteins were identified, of which 803 (28.8%) were common to all regions. NP, NP/IAF and IAF had 34 (12%), 80 (28.7%) and 44 (15%) unique proteins, respectively, with the OAF accounting for the highest proportion of 1,314 unique proteins (47%) (Figure 1E, lower panel). The aged OAF had the highest number of detected proteins with an average of 1,156, followed by the young OAF with an average of 818 (Figure 1F). The quantity and spectrum of protein categories identified suggest sufficient proteins had been extracted and the data are of high quality.

Levels of matrisome proteins declines in all compartments of aged discs

We divided the detected matrisome proteins into core matrisome (ECM proteins, encompassing collagens, proteoglycans and glycoproteins), and non-core matrisome (ECM regulators, ECM affiliated and secreted factors), according to a matrisome classification database (Naba et al., 2012) (matrisomeproject.mit.edu) (Figure 1F). Despite the large range of total numbers of proteins detected (419 to 1,920) across the 66 profiles (Figure 1F), all six sub-categories of the matrisome contained similar numbers of ECM proteins (Figure 1F,G; Supplemental Figure S2A). The non-core matrisome proteins were significantly more abundant in aged than in young discs (Supplemental Figure S2A). On average, 19 collagens, 18 proteoglycans, 68 glycoproteins, 52 ECM regulators, 22 ECM affiliated proteins, and 29 secreted factors of ECM were detected per profile. The majority of the proteins in these matrisome categories were detected in all disc compartments, and in both age groups. A summary of all the comparisons are presented in

Supplemental Figure S1C-E, and the commonly expressed matrisome proteins are listed in Table
2.

Even though there are approximately three times more non-matrisome than matrisome proteins
per profile on average (Figure 1F), their expression levels in terms of label-free quantification
(LFQ) values are markedly lower (Supplemental Figure S2B). Specifically, the expression levels
of core-matrisome were the highest, with an average $\log_2$(LFQ) of 30.65, followed by non-core
matrisome at 28.56, and then non-matrisome at 27.28 (Supplemental Figure S2B). Within the core-
matrisome, the expression was higher ($p=6.4\times10^{-21}$) in young (median 30.74) than aged (median
29.72) discs (Supplemental Figure S2C,D). This difference between young and aged discs is
consistent within the sub-categories of core and non-core matrisome, with the exception of the
ECM regulator category (Figure 1H). The non-core matrisome and non-matrisome, however,
exhibited smaller cross-compartment and cross-age differences in terms of expression levels
(Supplemental Figure S2E-H). This implies that the amount of ECM protein in each compartment
of the disc declines with ageing and possibly changes in the relative composition, but the numbers
of proteins detected per matrisome sub-category remain similar. This agrees with the concept that
ECM synthesis is not sufficient to counterbalance degradation, as exemplified in a proteoglycan
study (Silagi et al., 2018).

Cellular activities inferred from non-matrisome proteins

Although 86.5% (2,682) of the detected proteins were non-matrisome, their expression levels were
considerably lower than matrisome proteins across all sample profiles (Figure 1F). A functional
categorisation according to the Human Genome Nomenclature Committee gene family annotations
(Yates et al., 2017) showed many categories containing information for cellular components and
activities, with the top 30 listed in Figure 1I. These included transcriptional and translational
machineries, post-translational modifications, mitochondrial function, protein turnover; and
importantly, transcriptional factors, cell surface markers, and inflammatory proteins that can
inform gene regulation, cell identity and response in the context of IVD homeostasis, ageing and
degeneration.

These functional overviews highlighted 77 DNA-binding proteins and/or transcription factors, 83
cell surface markers, and 175 inflammatory-related proteins, with their clustering data presented

as heatmaps (Supplemental Figure S3). Transcription factors and cell surface markers are detected

in some profiles (Supplemental Figure S3A and B). The heatmap of the inflammatory-related

proteins showed that more than half of the proteins are detected in the majority of samples, with 4

major clusters distinguished by age and expression levels (Supplemental S3C). For example, one

of the clusters in the aged samples showed enrichment for complement and coagulation cascades

(False Discovery Rate, FDR q=1.62×10^{-21}) and clotting factors (FDR q=6.05×10^{-9}), indicating

potential infiltration of blood vessels. Lastly, there are 371 proteins involved in signalling

pathways, and their detection frequency in the different compartments and heat map expression

levels are illustrated in Supplemental Figure S3D.

Histones and housekeeping genes inform cross compartment- and age-specific

variations in cellularity

Cellularity within the IVD, especially the NP, decreases with age and degeneration (Rodriguez et

al., 2011; Sakai et al., 2012). We assessed whether cellularity of the different compartments could

be inferred from the proteomic data. Quantitation of histones can reflect the relative cellular

content of tissues (Wisniewski et al., 2014). We detected 10 histones, including subunits of histone

1 (HIST1H1B/C/D/E, HIST1H2BL, HIST1H3A, HIST1H4A) and histone 2 (HIST2H2AC,

HIST2H2BE, HIST2H3A), with 4 subunits identified in over 60 sample profiles that are mutually

co-expressed (Supplemental Figure S4A). Interestingly, histone concentrations, and thus

cellularity, increased from the inner to the outer compartments of the disc, and showed a highly

significant decrease in aged discs compared to young discs across all compartments (Figure 1J),

(Wilcoxon p=5.6×10^{-4}) (Supplemental Figure S4B).

GAPDH and ACTA2 are two commonly used reference proteins, involved in the metabolic

process and the cytoskeletal organisation of cells, respectively. They are expected to be relatively

constant between cells and are used to quantify the relative cellular content of tissues (Barber et

al., 2005). They were detected in all 66 profiles. GAPDH and ACTA2 amounts were significantly

correlated with a Pearson correlation coefficient (PCC) of 0.794 (p=9.3×10^{-9}) (Supplemental

Figure S4C), and they were both significantly co-expressed with the detected histone subunits,

with PCCs of 0.785 and 0.636, respectively (Figure 1K; Supplemental Figure S4D). As expected,

expression of the histones, GAPDH and ACTA2 was not correlated with two core-matrisome

proteins, ACAN and COL2A1 (Supplemental Figure S4E); whereas ACAN and COL2A1 were
significantly co-expressed (Supplemental Figure S4F), as expected due to their related regulation
of expression and tissue function. Thus, cellularity information can be obtained from proteomic
information, and the histone quantification showing reduced cellularity in the aged IVD is
consistent with the reported changes (Rodriguez et al., 2011; Sakai et al., 2012).

Phenotypic variations revealed by PCA and ANOVA

PCA captures the information content distinguishing age and tissue types

To gain a global overview of the data, we performed PCA on a set of 507 proteins selected
computationally, allowing maximal capture of valid values, while incurring minimal missing
values, followed by imputations (Supplemental Figure S5; Methods). The first two principal
components (PCs) explained a combined 65.5% of the variance, with 39.9% and 25.6% for the
first and second PCs, respectively (Figure 2A and B). A support vector machine with polynomial
kernel was trained to predict the boundaries: it showed PC1 to be most informative to predict age,
with a clear demarcation between the two age groups (Figure 2A, vertical boundary), whereas PC2
distinguished disc sample localities, separating the inner compartments (NP, NP/IAF and IAF)
from the OAF (Figure 2A, horizontal boundary). PC3 captured only 5.0% of the variance (Figure
2C,D), but it distinguished disc level, separating the lowest level (L5/S1) from the rest of the
lumbar discs (L3/4 and L4/5) (Figure 2D, horizontal boundary). Samples in the upper level (L3/4
and L4/5) appeared to be more divergent, with the aged disc samples deviating from the young
ones (Figure 2D).

Top correlated genes with the principal components are insightful of disc homeostasis

To extract the most informative features of the PCA, we performed proteome-wide associations
with each of the top three PCs, which accounted for over 70% of total variance, and presented the
top 100 most positively and top 100 most negatively correlated proteins for each of the PCs (Figure
2E-G). As expected, the correlation coefficients in absolute values were in the order of PC1 >
PC2 > PC3 (Figure 2E-G). The protein content is presented as non-matrisome proteins (grey
colour) and matrisome proteins (coloured) that are sub-categorised as previously. For the
negatively correlated proteins, the matrisome proteins contributed to PC1 in distinguishing young

disc samples, as well as to PC2 for sample location within the disc, but less so for disc level in PC3. Further, the relative composition of the core and non-core matrisome proteins varied between the three PCs, depicting the dynamic ECM requirement and its relevance in ageing (PC1), tissue composition within the disc (PC2) and mechanical loading (PC3).

PC1 of young discs identified known chondrocyte markers, CLEC3A (Lau et al., 2018) and LECT1/2 (Zhu et al., 2019); hedgehog signalling proteins, HHIPL2, HHIP, and SCUBE1 (Johnson et al., 2012); and xylosyltransferase-1 (XYLT1), a key enzyme for initiating the attachment of glycosaminoglycan side chains to proteoglycan core proteins (Silagi et al., 2018) (Figure 2E). Most of the proteins that were positively correlated in PC1 were coagulation factors or coagulation related, suggesting enhanced blood infiltration in aged discs. PC2 implicated key changes in molecular signalling proteins (hedgehog, WNT and Nodal) in the differences between the inner and outer disc regions (Figure 2F). Notably, PC2 contains heat shock proteins (HSPA1B, HSPA8, HSP90AA1, HSPB1) which are more strongly expressed in the OAF than in inner disc, indicating the OAF is under stress (Takao and Iwaki, 2002). Although the correlations in PC3 were much weaker, proteins such as CILP/CILP2, DCN, and LUM were associated with lower disc level.

ANOVA reveals the principal phenotypes for categories of ECMs

To investigate how age, disc compartment, level, and direction affect the protein profiles, we carried out ANOVA for each of these phenotypic factors for the categories of matrisome and non-matrisome proteins (Supplemental Figure S5H-J). In the young discs, the dominant phenotype explaining the variances for all protein categories was disc compartment. It is crucial that each disc compartment (NP, IAF and OAF) has the appropriate protein composition to function correctly (Supplemental Figure S5I). This also fits the understanding that young healthy discs are axially symmetric and do not vary across disc levels. In aged discs, compartment is still relevant for non-matrisome proteins and collagen, but disc level and directions become influential for other protein categories, which is consistent with variations in mechanical loading occurring in the discs with ageing and degeneration (Supplemental Figure S5J). In the combined (young and aged) disc samples, age was the dominant phenotype across major matrisome categories, while compartment best explained the variance in non-matrisome, reflecting the expected changes in cellular (non-matrisome) and structural (matrisome) functions of the discs (Supplemental Figure S5H). This

306 guided us to analyse the young and aged profiles separately, before performing cross-age
307 comparisons.

The spatial proteome of young and healthy discs

PCA of the 33 young profiles showed a distinctive separation of the OAF from the inner disc
regions on PC1 (upper panel of Figure 3A). In PC2, the lower level L5/S1 could generally be
distinguished from the upper lumbar levels (lower panel of Figure 3A). The detected proteome of
the young discs (Figure 3B) accounts for 9.2% of the human proteome (or 1,883 out of the 20,368
on UniProt). We performed multiple levels of pairwise comparisons (summarised in Supplemental
Figure S6A) to detect proteins associated with individual phenotypes, using three approaches (see
Methods): statistical tests; proteins detected in one group only; or proteins using a fold-change
threshold. We detected a set of 671 DEPs (Supplemental Table S2) (termed the 'variable set'),
containing both matrisome and non-matrisome proteins (Figure 3D), and visualised in a heatmap
(Supplemental Figure S7), with identification of four modules (Y1-Y4).

Expression modules showing lateral and anteroposterior trends

To investigate how the modules are associated with disc components, we compared their protein
expression profiles along the lateral and anteroposterior axes. The original $\log_2$(LFQ) values were
transformed to z-scores to be on the same scale. Proteins of the respective modules were
superimposed on the same charts, disc levels combined or separated (Figure 3E-H; Supplemental
Figure S8). Module Y1 is functionally relevant to NP, containing previously reported NP and novel
markers KRT19, CD109, KRT8 and CHRD (Anderson et al., 2002), CHRDL2, SCUBE1 (Johnson
et al., 2012) and CLEC3B. Proteins levels in Y1 are lower on one side of the OAF, increases
towards the central NP, before declining towards the opposing side, forming a concave pattern in
both lateral and anteroposterior directions, with a shoulder drop occurring between IAF and OAF
(Figure 3E).

Module Y2 was enriched in proteins for ECM organisation (FDR q=8.8×10^{-24}); Y3 was enriched
for proteins involved in smooth muscle contraction processes (FDR q=8.96×10^{-19}); and Y4 was
enriched for proteins of the innate immune system (FDR q=2.93×10^{-20}). Interestingly, these
modules all showed a tendency for convex patterns with upward expression toward the OAF

regions, in both lateral and anteroposterior directions, in all levels, combined or separated (Figure
3F-H; Supplemental S8B-D). The higher proportions of ECM, muscle contraction and immune
system proteins in the OAF are consistent with the contractile function of the AF (Nakai et al.,
2016), and with the NP being avascular and "immune-privileged" in a young homeostatic
environment (Sun et al., 2020).

Inner disc regions are characterised with NP markers

The most distinctive pattern on the PCA is the separation of the OAF from the inner disc, with 99
proteins expressed higher in the OAF and 55 expressed higher in the inner disc (Figure 3I,J).
Notably, OAF and inner disc contained different types of ECM proteins. The inner disc regions
were enriched in collagens (COL3A1, COL5A1, COL5A2, and COL11A2), matrillin (MATN3),
and proteins associated with ECM synthesis (PCOLCE) and matrix remodelling (MXRA5). We
also identified in the inner disc previously reported NP markers (KRT19, KRT8) (Risbud et al.,
2015), in addition to inhibitors of WNT (FRZB, DKK3) and BMP (CHRD, CHRDL2) signalling
(Figure 3I,J). Of note, FRZB and CHRDL2 were recently shown to have potential protective
characteristics in osteoarthritis (Ji et al., 2019). The TGFβ pathway appears to be suppressed in the
inner disc where antagonist CD109 (Bizet et al., 2011; Li et al., 2016) is highly expressed, and the
TGFβ activity indicator TGFBI is expressed higher in the OAF than in the inner disc.

The OAF signature is enriched with proteins characteristic of tendon and ligament

The OAF is enriched with various collagens (COL1A1, COL6A1/2/3, COL12A1 and COL14A1),
basement membrane (BM) proteins (LAMA4, LAMB2 and LAMC1), small leucine-rich
proteoglycans (SLRP) (BGN, DCN, FMOD, OGN, PRELP), and BM-anchoring protein (PRELP)
(Figure 3I,J). Tendon-related markers such as thrombospondins (THBS1/2/4) (Subramanian and
Schilling, 2014) and cartilage intermediate layer proteins (CILP/CILP2) are also expressed higher
in the OAF. Tenomodulin (TNMD) was exclusively expressed in 9 of the 12 young OAF profiles,
and not in any other compartments (Supplemental Table S2). This fits current understanding of
the AF as a tendon/ligament-like structure (Nakamichi et al., 2018). In addition, the OAF was
enriched in actin-myosin (Figure 3I), suggesting a role of contractile function in the OAF, and in
heat shock proteins (HSPA1B, HSPA8, HSPB1, HSP90B1, HSP90AA1), suggesting a stress
response to fluctuating mechanical loads.

Spatial proteome enables clear distinction between IAF and OAF

We sought to identify transitions in proteomic signatures between adjacent compartments. The NP

and NP/IAF protein profiles were highly similar (Supplemental Figure 6B). Likewise, NP/IAF and

IAF showed few DEPs, except COMP which was expressed higher in IAF (Supplemental Figure

6C). OAF and NP (Supplemental Figure 6R), and OAF and NP/IAF (Supplemental Figure 6O)

showed overlapping DEPs, consistent with NP and NP/IAF having highly similar protein profiles,

despite some differences in the anteroposterior direction (Supplemental Figure 6P,Q). The clearest

boundary within the IVD, between IAF and OAF, was marked by a set of DEPs, of which

COL5A1, SERPINA5, MXRA5 were enriched in the IAF, whereas LAMB2, THBS1, CTSD

typified the OAF (Supplemental Figure 6D). These findings agreed with the modular patterns

(Figure 3E-H).

The constant set shows common characteristics for the young reference architecture

Of the 1,880 proteins detected, 1204 proteins were not found to vary with respect to the phenotypic

factors. The majority of these proteins were detected in few profiles (Figure 3B) and were not used

in the comparisons. We set a cutoff for a detection in >1/2 of the profiles to prioritise a set of 245

proteins, hereby referred to as the 'constant set' (Figure 3C). Both the variable and the constant

sets contained high proportions of ECM proteins (Figure 3D). Amongst the proteins in the constant

set that were detected in all 33 young profiles were known protein markers defining a young NP

or disc, including COL2A1, ACAN and A2M (Risbud et al., 2015). Other key proteins in the

constant set included CHAD, HAPLN1, VCAN, HTRA1, CRTAC1, and CLU. Collectively, these

proteins showed the common characteristics shared by compartments of young discs, and they,

alongside the variable set, form the architectural landscape of the young disc.

Diverse changes in the spatial proteome with ageing

Fewer inner-outer differences but greater variation between levels in aged discs

PCA was used to identify compartmental, directional and level patterns for the aged discs (Figure

4A). Albeit less clear than for the young discs (Figure 3A), the OAF could be distinguished from

the inner disc regions on PC1, explaining 46.7% of the total variance (Figure 4A). PC2 showed a

more distinct separation of signatures from lumbar disc levels L5/S1 to the upper disc levels
(L4/L5 and L3/4), accounting for 21.8% of the total variance (Figure 4A).

Loss of the NP signature from inner disc regions

As with the young discs, we performed a series of comparative analyses (Figure 4C; Supplemental
Figure S9 A-H; Supplemental Table S3). Detection of DEPs between the OAF and the inner disc
(Figure 4B), showed that 100 proteins were expressed higher in the OAF, similar to young discs
(Figure 3C). However, in the inner regions, only 9 proteins were significantly expressed higher, in
marked contrast to the situation in young discs. Fifty-five of the 100 DEPs in the OAF region
overlapped in the same region in the young discs, but only 3 of the 9 DEPs in the inner region were
identified in the young disc; indicating changes in both regions but more dramatic in the inner
region. This suggests that ageing and associated changes may have initiated at the centre of the
disc. The typical NP markers (KRT8/19, CD109, CHRD, CHRDL2) were not detectable as DEPs
in the aged disc; but CHI3L2, A2M and SERPING1 (Figure 4B), which have known roles in tissue
fibrosis and wound healing (Lee et al., 2011; Naveau et al., 1994; Wang et al., 2019a), were
detected uniquely in the aged discs.

Gradual modification of ECM composition and cellular responses in the outer AF

A comparative analysis of the protein profiles indicated that the aged OAF retained 55% of the
proteins of a young OAF. These changes are primarily reflected in the class of SLRPs (BGN, DCN,
FMOD, OGN, and PRELP), and glycoproteins such as CILP, CILP2, COMP, FGA/B, and FGG.
From a cellular perspective, 45 proteins enriched in the aged OAF could be classified under
'responses to stress' (FDR q=1.86×10^{-7}; contributed by CAT, PRDX6, HSP90AB1, EEF1A1,
TUBB4B, P4HB, PRDX1, HSPA5, CRYAB, HIST1H1C), suggesting OAF cells are responding
to a changing environment such as mechanical loading and other stress factors.

Convergence of the inner disc and outer regions in aged discs

To map the relative changes between inner and outer regions of the aged discs, we performed a
systematic comparison between compartments (Supplemental Figure S9 A-D). The most
significant observation was a weakening of the distinction between IAF and OAF that was seen in
young discs, with only 17 DEPs expressed higher in the OAF of the aged disc (Supplemental

Figure S9C). More differences were seen when we included the NP in the comparison with OAF

(Supplemental Figure S9D), indicating some differences remain between inner and outer regions

of the aged discs. While the protein profiles of the NP and IAF were similar, with no detectable

DEPs (Supplemental Figure S9A,B), their compositions shared more resemblance with the OAF.

These progressive changes of the protein profiles and DEPs between inner and outer compartments

suggest the protein composition of the inner disc compartments becomes more similar to the OAF

with ageing, with the greatest changes in the inner regions. This further supports a change initiating

from the inner region of the discs with ageing.

Changes in young module patterns reflect convergence of disc compartments

We investigated protein composition in the young disc modules (Y1-Y4) across the lateral and

anteroposterior axes (Figure 4C-F). For module Y1 that consists of proteins defining the NP

region, the distinctive concave pattern has flattened along both axes, but more so for the

anteroposterior direction where–the clear interface between IAF and OAF was lost (Figure 4C;

Supplemental Figure S9I). Similarly for modules Y2 and Y3, which consist of proteins defining

the AF region, the trends between inner and outer regions of the disc have changed such that the

patterns become more convex, with a change that is a continuum from inner to outer regions

(Figure 4D,E; Supplemental Figure S9J,K). These changes in modules Y1-3 in the aged disc

further illustrate the convergence of the inner and outer regions, with the NP/IAF becoming more

OAF-like. For Y4, the patterns along the lateral and anteroposterior axes were completely

disrupted (Figure 4F; Supplemental Figure S9J). As Y4 contains proteins involved in vascularity

and inflammatory processes, these changes indicate disruption of cellular homeostasis in the NP.

Disc level variations reflect spatial and temporal progression of disc changes

The protein profiles of the aged disc levels, consistent with the PCA findings, showed similarity

between L3/4 and L4/5 (Supplemental Figure S9E), but, in contrast to young discs, differences

between L5/S1 and L4/5 (Supplemental Figure S9F), and more marked differences between L5/S1

and L3/4 (Supplemental Figure S9G,H). Overall, the findings from PCA, protein profiles (Figure

2B-D) and MRI (Figure 1B) agree. As compared to young discs, the more divergent differences

across the aged disc levels potentially reflect progressive transmission from the initiating disc to

the adjacent discs with ageing. To further investigate the aetiologies underlying IDD, cross-age

comparisons are needed.

Aetiological insights uncovered by young/aged comparisons

Next, we performed extensive pair-wise comparisons between the young and aged samples under

a defined scheme (Supplemental Figure S10A; Supplemental Table S4). First, we compared all 33

young samples with all 33 aged samples, which identified 169 DEPs with 104 expressed higher in

the young and 65 expressed higher in the aged discs (Figure 5A). A simple GO term analysis

showed that the most important biological property for a young disc is structural integrity, which

is lost in aged discs (Figure 5B; Supplemental Table S5). The protein classes most enriched in the

young discs were related to cartilage synthesis, chondrocyte development, and ECM organisation

(Figure 5B). The major changes in the aged discs relative to young ones, were proteins involved

in cellular responses to an ageing environment, including inflammatory and cellular stress signals,

progressive remodelling of disc compartments, and diminishing metabolic activities (Figure 5C;

Supplemental Table S5).

Inner disc regions present with most changes in ageing

For young versus old discs, we compared DEPs of the whole disc (Figure 5A) with those from the

inner regions (Figure 5D) and OAF (Figure 5E) only. Seventy-five percent (78/104) of the down-

regulated DEPs (Figure 5F) were attributed to the inner regions, and only 17% (18/104) were

attributed to the OAF (Figure 5F). Similarly, 65% (42/65) of the up-regulated DEPs were solely

contributed by inner disc regions, and only 18.4% (12/65) by the OAF. Only 5 DEPs were higher

in the OAF, while 49 were uniquely up-regulated in the inner discs (Figure 5G). The key biological

processes in each of the compartments are highlighted in Figure 5F, G.

The changing biological processes in the aged discs

Expression of known NP markers was reduced in aged discs, especially proteins involved in the

ECM and its remodelling, where many of the core matrisome proteins essential for the structural

function of the NP were less abundant or absent. While this is consistent with previous

observations (Feng et al., 2006), of interest was the presence of a set of protein changes that were

also seen in the OAF, which was also rich in ECM and matrix remodelling proteins (HTRA1,

SERPINA1, SERPINA3, SERPINC1, SERPINF1, and TIMP3) and proteins involved in fibrotic

events (FN1, POSTN, APOA1, APOB), suggesting these changes are occurring in the aging IVDs

(Figure 5 F,G).

Proteins associated with cellular stress are decreased in the aged inner disc, with functions ranging

from molecular chaperones needed for protein folding (HSPB1, HSPA1B and HSPA9) to

modulation of oxidative stress (SOD1) (Figure 5F). SOD1 has been shown to become less

abundant in the aged IVD (Hou et al., 2014) and in osteoarthritis (Scott et al., 2010). HSPB1 is

cytoprotective and a deficiency is associated with inflammation reported in degenerative discs

(Wuertz et al., 2012). We found an increased concentration of clusterin (CLU) (Figure 5G), an

extracellular chaperone that aids the solubilisation of misfolded protein complexes by binding

directly and preventing protein aggregation (Trougakos, 2013; Wyatt et al., 2009), and also has a

role in suppressing fibrosis (Peix et al., 2018).

Inhibitors of WNT (DKK3 and FRZB), and antagonists of BMP/TGFβ (CD109, CHRDL2, DCN

FMOD, INHBA and THBS1) signalling were decreased or absent in the aged inner region (Figure

5F,G), consistent with the reported up-regulation of these pathways in IDD (Hiyama et al., 2010)

and its closely related condition osteoarthritis (Leijten et al., 2013). Targets of hedgehog signalling

(HHIPL2 and SCUBE1) were also reduced (Figure 5F), consistent with SHH's key roles in IVD

development and maintenance (Rajesh and Dahia, 2018). TGFβ signalling is a well-known

pathway associated with fibrotic outcomes. WNT is known to induce chondrocyte hypertrophy

(Dong et al., 2006), that can be enhanced by a reduction in S100A1 (Figure 5F), a known inhibitor

of chondrocyte hypertrophy (Saito et al., 2007).

To gain an overview of the disc compartment variations between young and aged discs, we

followed the same strategy as in Supplemental Figure S7 to aggregate three categories of DEPs in

all 23 comparisons (Supplemental Figure S10A), and created a heatmap from the resulting 719

DEPs. This allowed us to identify 6 major protein modules (Figure 5H). A striking feature is

module 6 (M6), which is enriched for proteins involved in the complement pathway (GSEA

Hallmark FDR q=4.9×10^{-14}) and angiogenesis (q=2.3×10^{-3}). This module contains proteins that

are all highly expressed in the inner regions of the aged disc, suggesting the presence of blood. M6

also contains the macrophage marker CD14, which supports this notion.

We visualised the relationship between the young (Y1-4) and young/aged (M1-6) modules using

an alluvial chart (Figure 5I). Y1 corresponds primarily to M1b that is enriched with fibrosis,

angiogenesis, apoptosis and EMT (epithelial to mesenchymal transition) proteins. Y2 seems to

have been deconstructed into three M modules (M2b > M3 > M4). M2b and M3 contain proteins

linked to heterogeneous functions, while proteins in M4 are associated with myogenesis and

cellular metabolism, but also linked to fibrosis and angiogenesis. Y3 primarily links to M2a with

a strong link to myogenesis, and mildly connects with M3 and M4. Y4 has the strongest connection

with M6a, which is linked to coagulation. Both the variable and the constant sets of the young disc

were also changed in ageing.

In all, the IVD proteome showed that with ageing, activities of the SHH pathways were decreased,

while those of the WNT and BMP/TGFβ pathways, EMT, angiogenesis, fibrosis, cellular stresses

and chondrocyte hypertrophy-like events were increased.

Concordant changes between the transcriptome and proteome of disc cells

The proteome reflects both current and past transcriptional activities. To investigate upstream

cellular and regulatory activities, we obtained transcriptome profiles from two IVD compartments

(NP and AF) and two sample states (young, scoliotic but non-degenerated, YND; aged individuals

with clinically diagnosed IDD, AGD) (Table 1). The transcriptome profiles of YND and AGD are

similar to the young and aged disc proteome samples, respectively. After normalisation

(Supplemental Figure S11A) and hierarchical clustering, we found patterns reflecting relationships

among IVD compartments and ages/states (Supplemental Figure S11B-D). PCA of the

transcriptome profiles showed that PC1 captured age/state variations (Figure 6A) and PC2

captured the compartment (AF or NP) differences, with a high degree of similarity to the proteomic

PCA that explained 65.0% of all data variance (Figure 2A).

Transcriptome shows AF-like characteristics of the aged/degenerated NP

We compared the transcriptome profiles of different compartments and age/state-groups. We

detected 88 DEGs (differentially expressed genes) between young AF and NP samples

(Supplemental Figure S11E); 39 were more abundant in NP (including known NP markers *CD24*,

KRT19) and 49 were more abundant in AF, including the AF markers, *COL1A1* and *THBS1*. In

the AGD samples, 11 genes differed between AF and NP (Supplemental Figure S11F), comparable

to the proteome profiles (Figure 4C). Between the YND and AGD AF, there were 45 DEGs, with

COL1A1 and *MMP1* more abundant in YND and *COL10A1*, WNT signalling (*WIF1*, *WNT16*),

inflammatory (*TNFAIP6*, *CXCL14*, *IL11*), and fibrosis-associated (*FN1*, *CXCL14*) genes more

abundant in AGD (Supplemental Figure S11G). The greatest difference was between YND and

AGD NP, with 216 DEGs (Supplemental Figure S11H), with a marked loss of NP markers

(*KRT19*, *CD24*), and gain of AF (*THBS1*, *DCN*), proteolytic (*ADAMTS5*), and EMT (*COL1A1*,

COL3A1, *PDPN*, *NT5E*, *LTBP1*) markers with age. Again, consistent with the proteomic findings,

the most marked changes are in the NP, with the transcriptome profiles becoming AF-like.

Concordance between transcriptome and proteome profiles

We partitioned the DEGs between the YND and AGD into DEGs for individual compartments

(Figure 6B). The transcriptomic (Figure 6B) Venn diagram was very similar to the proteomic one

(Figure 5F-G). For example, WNT/TGFβ antagonists and ECM genes were all down-regulated

with ageing/degeneration, while genes associated with stress and ECM remodelling were more

common. When we directly compared the transcriptomic DEGs and proteomic DEPs across

age/states and compartments (Figure 6C-F), we observed strong concordance between the two

types of datasets for a series of markers. In the young discs, concordant markers included *KRT19*

and *KRT8*, *CHRDL2*, *FRZB*, and *DKK3* in the NP, and *COL1A1*, *SERPINF1*, *COL14A1*, and

THBS4 in the AF (Figure 6C). In the AGD discs, concordant markers included *CHI3L2*, *A2M* and

ANGPTL4 in the NP and *MYH9*, *HSP90AB1*, *HBA1*, and *ACTA2* in the AF (Figure 6D). A high

degree of concordance was also observed when we compared across age/states for the AF (Figure

6E) and NP (Figure 6F).

Despite the transcriptomic samples having diagnoses (scoliosis for YND and IDD for AGD),

whereas the proteome samples were cadaver samples with no reported diagnosis of IDD, the

changes detected in the transcriptome profiles substantially support the proteomic findings. A

surprising indication from the transcriptome was the increased levels of *COL10A1* (Lu et al.,

2014), *BMP2* (Grimsrud et al., 2001), *IBSP*, defensin beta-1 (*DEFB1*), *ADAMTS5*, pro-

inflammatory (*TNFAIP6*, *CXCL*) and proliferation (*CCND1*, *IGFBP*) genes in the AGD NP

(Supplemental Figure S11E-H), reaffirming the involvement of hypertrophic-like events (Melas
et al., 2014) in the aged and degenerated NP.

The genome-wide transcriptomic data included over 20 times more genes per profile than the
proteomic data, providing additional biological information about the disc, particularly low
abundance proteins, such as transcription factors and surface markers. For example, additional
WNT antagonists were *WIF1* (Wnt inhibitory factor) and *GREM1* (Figure 6B) (Leijten et al.,
2013). Comparing the YND NP against YND AF or AGD NP (Supplemental Figure S11E,H), we
identified higher expression of three transcription factors, *T* (brachyury), *HOPX* (homeodomain-
only protein homeobox), and *ZNF385B* in the YND NP. Brachyury is a well-known marker for
the NP (Risbud et al., 2015), and *HOPX* is differentially expressed in mouse NP as compared to
AF (Veras et al., 2020), and expressed in mouse notochordal NP cells (Lam, 2013). Overall,
transcriptomic data confirmed the proteomic findings and revealed additional markers.

Changes in the active proteome in the ageing IVD

The proteomic data up to this point is a static form of measurement (static proteome) and represents
the accumulation and turnover of all proteins up to the time of harvest. The transcriptome indicates
genes that are actively transcribed, but does not necessarily correlate to translation or protein
turnover. Thus, we studied changes in the IVD proteome (dynamic proteome) that would reflect
newly synthesised proteins and proteins cleaved by proteases (degradome), and how they relate to
the static proteomic and transcriptomic findings reported above.

Aged or degenerated discs synthesise fewer proteins

We performed e*x vivo* labelling of newly synthesised proteins using the SILAC protocol (Ong et
al., 2002) (Figure 7A; Methods) on AF and NP samples from 4 YND individuals and one AGD
individual (Table 1). In the SILAC profiles, light isotope-containing signals correspond to the pre-
existing unlabelled proteome, and heavy isotope-containing signals to newly synthesised proteins
(Figure 7B). The ECM compositions in the light isotope-containing profiles (Figure 7B, middle
panel) are similar to the static proteome samples of the corresponding age groups described above
(Figure 1F). In contrast, the heavy isotope-containing profiles contained fewer proteins in the AGD

than in the YND samples (Figure 7B, left panel) and showed variable heavy to light ratio profiles
(Figure 7B, right panel).

To facilitate comparisons, we averaged the abundance of the proteins detected in the NP or AF for
which we had more than one sample, then ranked the abundance of the heavy isotope-containing
(Figure 7C) and light isotope-containing (Figure 7D) proteins. The number of proteins newly
synthesised in the AGD samples was about half that in the YND samples (Figure 7C). This is
unlikely to be a technical artefact as the total number of light isotope-containing proteins detected
in the AGD samples is comparable to the YND, in both AF and NP (Figure 7D), and the difference
is again well illustrated in the heavy to light ratios (Figure 7E).

Reduced synthesis of non-matrisome proteins was found for the AGD samples (GAPDH as a
reference point (dotted red lines) (Figures 7C,D; Figure 7C,E, grey portions). Of the 68 high
abundant non-matrisome proteins in the YND NP compartment that were not present in the AGD
NP, 28 are ribosomal proteins (Supplemental Figure S12B), suggesting reduced translational
activities. This agrees with our earlier findings of cellularity, as represented by histones, in the
static proteome (Figure 1J, K).

Changes in protein synthesis in response to the cell microenvironment affects the architecture of
the disc proteome. To understand how the cells may contribute and respond to the accumulated
matrisome in the young and aged disc, we compared the newly synthesised matrisome proteins of
AGD and YND samples rearranged in order of abundance (Figure 7E). More matrisome proteins
were synthesised in YND samples across all classes. In YND AF, collagens were synthesised in
higher proportions than in AGD AF with the exception of fibril-associated COL12A1 (Figure 7E,
top panel). Similarly, higher proportions in YND AF were observed for proteoglycans (except
FMOD), glycoproteins (except TNC, FBN1, FGG and FGA), ECM affiliated proteins (except
C1QB), ECM regulators (except SERPINF2, SERPIND1, A2M, ITIH2, PLG), and secreted
factors (except ANGPTL2). Notably, regulators that were exclusively synthesised in young AF
are involved in collagen synthesis (P4HA1/2, LOXL2, LOX, PLOD1/2) and matrix turnover
(MMP3), with enrichment of protease HTRA1 and protease inhibitors TIMP3 and ITIH in YND
AF compared to AGD AF.

In AGD NP, overall collagen synthesis was less than in YND NP (Figure 7E, lower panel);
however, there was more synthesis of COL6A1/2/3 and COL12A1. Furthermore, AGD NP
synthesised more LUM, FMOD, DCN, PRG4, and PRELP proteoglycans than YND NP. Notably,
there was less synthesis of ECM-affiliated proteins (except C1QC and SEMA3A) and regulators
– particularly those involved in collagen synthesis (P4HA1/2, LOXL2, LOX) – but an increase in
protease inhibitors. A number of newly synthesised proteins in AGD NP were similarly
represented in the transcriptome data, including POSTN, ITIH2, SERPINC1, IGFBP3, and PLG.
Some genes were simultaneously underrepresented in the AGD NP transcriptome and newly
synthesised proteins, including hypertrophy inhibitor GREM1 (Leijten et al., 2013).

Proteome of aged or degenerated discs is at a higher degradative state

The degradome reflects protein turnover by identifying cleaved proteins in a sample (Lopez-Otin
and Overall, 2002).When combined with relative quantification of proteins through the use of
isotopic and isobaric tagging and enrichment for cleaved neo amine (N)-termini of proteins before
labelled samples are quantified by mass-spectrometry, degradomics is a powerful approach to
identify the actual status of protein cleavage *in vivo.*

We employed the well validated and sensitive terminal amine isotopic labelling of substrates
(TAILS) method (Kleifeld et al., 2010; Rauniyar and Yates, 2014) to analyse and compare 6 discs
from 6 individuals (3 young or non-degenerated, YND; and 3 aged and degenerated, AGD) (Table
1) (Figure 7F) (Kleifeld et al., 2010; Rauniyar and Yates, 2014) using the 6-plex tandem mass tag
(TMT)-TAILS (labelling 6 independent samples and analysed together on the mass spectrometer),
with one of the YND samples (YND136) as the reference sample (Figure 7F). Whereas shotgun
proteomics is intended to identify the proteome components, N-terminome data is designed to
identify the exact cleavage site in proteins that also evidence stable cleavage products *in vivo*.

Here, TAILS identified 92 and 68 cleaved proteins in the AF and NP disc samples, respectively.
For each protein, we calculated the means and standard deviations in the AGD and YND samples,
and then ranked these according to the normalised differences between the two groups, for NP
(Figure 7G) and AF (Figure 7H). By setting a cut-off of absolute normalised difference of greater
than 1, we detected proteins that were either cleaved less or cleaved more in AGD samples
compared to the YND samples. Interestingly, in both AF and NP, the AGD samples had more

cleaved proteins than the YND samples. In the NP, 5 proteins were less cleaved in AGD and 10

proteins were more cleaved (Figure 7G); while in the AF, 5 were less cleaved and 17 were more

cleaved (Figure 7H).

The proteins with more cleavage sites were functionally relevant to the corresponding disc

compartments. For example, in the YND NP, there were more cleaved neo N-terminal peptides

from cleaved CHRDL2, COL2A1, immunoglobulins and LYZ; whereas the AGD NP was

enriched for semi-trypic cleaved neo N-terminal peptides corresponding to cleaved blood proteins

(A2M, ALB, HBA1) and matrix proteins FN1, COMP, and PRG4 (Figure 7G). In the AGD AF,

we quantified increased proportions of peptides relative to YND AF, that corresponded to cleaved

blood proteins (HBB, HBA1, C3, immunoglobulins), isoform 2 of clusterin (CLU), FN1, COMP,

HAPLN1, CILP, FGFBP2; whereas in the YND AF there were more peptides corresponding to

cleaved collagen types I, II and VI (Figure 7H).

The cleavage sites in the non-degenerated aged samples in both compartments did not deviate

noticeably from the reference sample, in terms of relative abundance. These results suggest that

protein degradation activity increases with age, whether or not degeneration occurs.

MRI landscape correlates with proteomic landscape

We tested for a correlation between MRI signal intensity and proteome composition. In

conventional 3T MRI of young discs, the NP is brightest reflecting its high hydration state while

the AF is darker, thus less hydrated (Figure 1A; Supplemental Figure S13B). Since aged discs

present with more MRI phenotypes, we used higher resolution MRI (7T) on them (Figure 1B;

Figure 8A), which showed less contrast between NP and AF than in the young discs. To enhance

robustness, we obtained three transverse stacks per disc level for the aged discs (Figure 8B;

Supplemental Figure S13D), and averaged the pixel intensities for the different compartments

showing that overall, the inner regions were still brighter than the outer (Figure 8C).

Next, we performed a level-compartment bi-clustering on the pixel intensities of the aged disc

MRIs, which was bound by disc level and compartment (Figure 8D). The findings resembled the

proteomic clustering and PCA patterns (Figure 2; Supplemental Figure S6V-W). We performed a

pixel intensity averaging of the disc compartments from the 3T images (Supplemental Figure 13B),

and a level-compartment bi-clustering on the pixel intensities (Supplemental Figure S13C). While

the clustering can clearly partition the inner from the outer disc compartments, the information

value from each of the compartments is less due to the lower resolution of the MRI. In all, these

results indicate a link between regional MRI landscapes and proteome profiles, prompting us to

investigate their potential connections.

Proteome-wide associations with MRI landscapes reveals a hydration matrisome

The MRI and the static proteome were done on the same specimens in both individuals, so we

could perform proteome-wide associations with the MRI intensities. We detected 85 significantly

correlated ECM proteins, hereby referred to as the hydration matrisome (Figure 8E). We found no

collagen to be positively correlated with brighter MRI, which fits current understanding as

collagens contribute to fibrosis and dehydration. Other classes of matrisome proteins were either

positively or negatively correlated, with differential components for each class (Figure 8E).

Positively correlated proteoglycans included EPYC, PRG4 (lubricin) and VCAN, consistent with

their normal expression in a young disc and hydration properties. Negatively correlated proteins

included OAF (TNC, SLRPs) and fibrotic (POSTN) markers (Figure 8E).

Given this MRI-proteome link and the greater dynamic ranges of MRI in the aged discs enabled

by the higher resolution 7T MRIs (Figure 8D), we hypothesised that the hydration matrisome

might be used to provide information about MRI intensities and thus disc hydration. To test this,

we trained a LASSO regression model (Tibshirani, 1996) of the aged MRIs using the hydration

matrisome (85 proteins), and applied the model to predict the intensity of the MRIs of the young

discs, based on the young proteome of the same 85 proteins. Remarkably, we obtained a PCC of

0.689 (p=8.9×10^{-6}; Spearman=0.776) between the actual and predicted MRI (Supplemental Figure

S13E). The predicted MRI intensities of the young disc exhibited a smooth monotonic decrease

from the NP towards IAF, then dropped suddenly towards the OAF (Figure 8F, right panel), with

an ROC AUC (receiver operating characteristics, area under the curve) of 0.996 between IAF and

OAF (Supplemental Figure S13F). In comparison, actual MRIs exhibited a linear decrease from

NP to OAF (Figure 8F, left panel). On reviewing these two patterns, we argue that the predicted

intensities may be a more faithful representation of the young discs' water contents than the actual

MRI, as it reflects the gross images (Supplemental Figure S1A), PCA (Figure 3A) and Y1 modular

trend in the young discs (Figure 3E). This exercise not only revealed the inherent connections

between regional MRI and regional proteome, but also identified a set of ECM components that is

predictive of MRI relating to disc hydration, which may be valuable for future clinical applications.

Discussion

The IVD is architecturally designed for its function. Thus, spatial information relating to the molecular make-up in different compartments is critical for understanding its biology in health and disease. This can only be obtained from intact IVDs. In our study, we provided information on the spatial proteome of entire human IVDs encompassing different compartments along the lateral and anteroposterior axes, and at different levels of the lumbar discs. This spatial proteome provides a reference map of the proteomic architecture of a young disc which can be used to provide insights into ageing and degeneration.

A discovery dataset was established from intact lumbar discs of a young and an aged individual, consisting of 66 proteomic specimens in total; with 11 regional specimens per disc for three lumbar disc levels per individual. The young discs were considered non-degenerated by disc morphologies and MRI. The aged discs displayed degeneration characteristics including reduced MRI intensities and annular tears. We extended the findings of this data set with transcriptomics, SILAC and degradome profiling from another 15 individuals.

Using a well-established protein extraction protocol (Onnerfjord et al., 2012b), and chromatographic fractionation of the peptides prior to mass spectrometry, we produced a dataset of the human intervertebral disc comprising 3,100 proteins, encompassing ~400 matrisome and 2,700 non-matrisome proteins, with 1,769 proteins detected in 3 or more profiles, considerably higher than recent studies (Maseda et al., 2016; Rajasekaran et al., 2020; Ranjani et al., 2016; Sarath Babu et al., 2016). The high quality of our data enabled the application of unbiased approaches including PCA and ANOVA to reveal the relative importance of the phenotypic factors. Particularly, age was found to be the dominant factor influencing proteome profiles.

Comparisons between different compartments of the young disc produced a reference landscape containing known (KRT8/19 for the NP; COL1A1, CILP and COMP for the AF) and novel (FRZB, CHRD, CHRDL2 for the NP; TNMD, SLRPs and SOD1 for the AF) markers. The young healthy discs were enriched for matrisome components consistent with a healthy functional young IVD. Despite morphological differences between NP and IAF, the inner disc compartments (NP, NP/IAF and IAF) display high similarities, in contrast to the large differences between IAF and OAF, which was consistent for discs from all lumbar disc levels. This morphological-molecular

discrepancy might be accounted for by subtle differences in the ECM organisation, such as differences in GAG moieties on proteoglycans, or levels of glycosylation or other modifications of ECM that diversify function (Silagi et al., 2018). Nonetheless, we partitioned the detected proteins into a variable set that captures the diversity, and a constant set that lays the common foundation of all young compartments, which work in synergy to achieve disc function.

Clustering analysis of the 671 DEPs of the variable set identified 4 key modules (Y1-Y4). Visually, Y1 and Y2 mapped across the lateral and anteroposterior axes with opposing trends. Molecularly, module Y1 (NP) contained proteins promoting regulation of matrix remodelling, such as matrix degradation inhibitor MATN3 (Jayasuriya et al., 2012) and MMP inhibitor TIMP1. Inhibitors of WNT and BMP signalling were also present. Module Y2 (OAF) included COL1A1, THBS1/2/3, CILP1/CILP2 and TNMD, consistent with the OAF's tendon-like features (Nakamichi et al., 2018). It also included a set of SLRPs that might play roles in regulation of collagen assembly (Robinson et al., 2017; Taye et al., 2020), fibril alignment (Robinson et al., 2017), maturation and crosslinking (Kalamajski et al., 2016); while others are known to inhibit or promote TGFβ signalling (Markmann et al., 2000). Notably, the composition of the IAF appeared to be a transition zone between NP and OAF rather than an independent compartment, as few proteins can distinguish it from adjacent compartments. Classes of proteins in both Y3 (smooth muscle feature) and Y4 (immune and blood) resemble Y2, which reflects the contractile property of the AF (Nakai et al., 2016) and the capillaries infiltrating or present at the superficial outer surface of the IVD.

In the aged disc, the change in the DEPs between the inner and outer regions of the discs suggests extensive changes in the inner compartment(s). Mapping the aged data onto modules Y1-Y4 allowed a visualisation of the changes. The flattening of the Y1 and Y2 modules along both the lateral and anteroposterior axes indicated a convergence of the inner and outer disc. This is supported by the observed rapid decline of NP proteins and increase of AF proteins in the inner region. Fewer changes were seen in the aged OAF, which concurs with the notion that degenerative changes originate from the NP and radiate outwards. The most marked change was seen in module Y4 (blood), where the pattern was inverted, characterised by high expression in the NP but low in OAF. While contamination cannot be excluded and there are reports that capillaries do not infiltrate the NP even in degeneration (Nerlich et al., 2007), our finding is consistent with other proteomic studies showing enrichment of blood proteins in pathological NP (Maseda et al., 2016).

The route of infiltration can be from the fissured AF or cartilage endplates. Calcified endplates are
more susceptible to microfractures, which can lead to blood infiltration into the NP (Sun et al.,
2020). Of interest is the involvement of an immune response within the inner disc. This
corroborates reports of inflammatory processes in ageing and degenerative discs, with the up-
regulation of pro-inflammatory cytokines and presence of inflammatory cells (Molinos et al.,
2015; Wuertz et al., 2012).

The SILAC and degradome studies provided important insights into age-related differences in the
biosynthetic and turnover activity in the IVD. The SILAC data indicated that protein synthesis is
significantly impaired in aged degenerated discs. These findings correlate with reports of reduced
cellularity in ageing (Rodriguez et al., 2011), which we have also ascertained by leveraging the
relationship of histones and housekeeping genes with cell numbers. From the TAILS degradome
analysis, we observed more cleaved protein fragments in aged compartments, particularly for
structural proteins important for tissue integrity such as COMP and those involved in cell-matrix
interactions such as FN1, which was coupled with the enrichment of the proteolytic process GO
terms in the aged static proteome. Collectively, this reveals a systematic modification and
replacement of the primary proteomic architecture of the young IVD with age that is associated
with diminished or failure in functional properties in ageing or degeneration.

Despite known transcriptome-proteome discordance (Fortelny et al., 2017), our identification of
concordant changes allow insights into active changes in the young and aged discs. For example,
inhibitors of the WNT pathway and antagonists of BMP/TGFβ signalling (Leijten et al., 2013)
were down regulated in the aged discs in both the proteome and transcriptome. Interestingly, the
activation of these pathways is known to promote chondrocyte hypertrophy (Dong et al., 2006),
and hypertrophy has been noted in IDD (Rutges et al., 2010). This suggests a model where the
regulatory environment suppressing cellular hypertrophy changes with ageing or degeneration,
resulting in conditions such as cellular senescence and tissue mineralisation that are part of the
pathological process. In support, S100A1, a known inhibitor of chondrocyte hypertrophy (Saito et
al., 2007) is down regulated, while chondrocyte hypertrophy markers *COL10A1* and *IBSP*
(Komori, 2010) are up-regulated in the aged disc. Similar changes have been observed in ageing
mouse NP (Veras et al., 2020) as well as in osteoarthritis (Zhu et al., 2009) where chondrocyte
hypertrophy is thought to be involved in its aetiology (Ji et al., 2019; van der Kraan and van den

Berg, 2012). Given that WNT inhibitors are already in clinical trials for osteoarthritis (Wang et

al., 2019b), this may point to a prospective therapeutic strategy for IDD.

A key finding of our study is the direct demonstration, within a single individual, of association

between the hydration status of the disc as revealed by MRI, and the matrisome composition of

the disc proteome. The remarkable correlation between predicted hydration states inferred from

the spatial proteomic data and the high-definition phenotyping of the aged disc afforded by 7T

MRI has enormous potential for understanding the molecular processes underlying IDD.

In conclusion, we have generated a comprehensive dataset of the young and aged disc spatial

proteome, that provides detailed information on the proteome in the different compartments of the

IVD and the architectural changes that accompany ageing. The data reveal the role of the spatial

organisation of the proteome that underpins the functional properties of the different compartments

of the human IVD with implications for the underlying molecular pathology of degeneration

(Figure 8G, H). The rich datasets are a valuable resource for cross referencing with animal and *in*

vitro studies to evaluate clinical relevance and guide the development of therapeutics of human

IDD.

Materials and methods

Cadaveric specimens

Two human lumbar spines were obtained through approved regulations and governing bodies, with one young (16M) provided by L.H. (McGill University) and one aged (59M) from Articular Engineering, LLC (IL, USA). The young lumbar spine was received frozen as an intact whole lumbar spine. The aged lumbar spine was received frozen, dissected into bone-disc-bone segments. The cadaveric samples were stored at -80^0C until use.

Clinical specimens

Clinical specimens were obtained with approval by the Institutional Review Board (references UW 13-576 and EC 1516-00 11/01/2001) and with informed consent in accordance with the Helsinki Declaration of 1975 (revision 1983) from another 15 patients undergoing surgery for IDD, trauma or AIS (adolescent idiopathic scoliosis) at Queen Mary Hospital (Hong Kong), and Duchess of Kent Children's Hospital (Hong Kong). Information of both the cadaveric and clinical samples are summarised in Table 1.

MRI imaging of cadaveric samples

The discs were thawed overnight at 4^0C, and then pre-equalised for scanning at room temperature. For the young IVD, these were imaged together as the lumbar spine was kept intact. T2-weighted and T1-weighted sagittal and axial MRI, T1-rho MRI and Ultrashort-time-to-echo MRI images were obtained using a 3T Philips Achieva 3.0 system at the Department of Diagnostic Radiology, The University of Hong Kong.

For the aged discs, the IVD were imaged separately as bone-disc-bone segments, at the Department of Electrical and Electronic Engineering, The University of Hong Kong. The MRS and CEST imaging were performed. The FOV for the CEST imaging was adjusted to 76.8×76.8 mm^2 to accommodate the size of human lumbar discs (matrix size = 64×64, slice thickness = 2 mm). All MRI experiments were performed at room temperature using a 7 T pre-clinical scanner (70/16 Pharmascan, Bruker BioSpin GmbH, Germany) equipped with a 370 mT/m gradient system along

835 each axis. Single-channel volume RF coils with different diameters were used for the samples

836 based on size (60 mm for GAG phantoms and human cadaveric discs).

Image assessment of the aged lumbar IVD

838 The MRI images in the transverse view were then assessed for intensity of the image (brighter

839 signifying more water content). Three transverse MRI images per IVD were overlaid with a grid

840 representing the areas that were cut for mass-spectrometry measurements as outlined previously.

841 For each region, the 'intensity' was represented by the average of the pixel intensities, which were

842 graphically visualised and used for correlative studies.

Division of cadaveric IVD for mass spectrometry analysis

The endplates were carefully cut off with a scalpel, exposing the surface of the IVD, which were

then cut into small segments spanning seven segments in the central left-right lateral axis, and

five segments in the central anteroposterior axis (Figure 1C). In all, this corresponds to a total of

11 locations per IVD. Among them, 4 are from the OAF, 2 from the IAF (but only in the lateral

axis), 1 from the central NP, and 4 from a transition zone between IAF and the NP (designated

the 'NP/IAF'). Samples were stored frozen at -80^0C until use.

SILAC by ex vivo culture of disc tissues

NP and AF disc tissues from spine surgeries (Table 1) were cultured in custom-made Arg- and

Lys-free α-MEM (AthenaES) as per formulation of Gibco α-MEM (Cat #11900-024),

supplemented with 10% dialysed FBS (10,000 MWCO, Biowest, Cat# S181D),

penicillin/streptomycin, 2.2 g/L sodium bicarbonate (Sigma), 30 mg/L L-methionine (Sigma), 21

mg/L "heavy" isotope-labelled $^{13}C_6$ L-arginine (Arg6, Cambridge Isotopes, Cat # CLM-2265-H),

146 mg/L "heavy" isotope-labelled 4,4,5,5-D4 L-Lysine (Lys4, Cambridge Isotopes, Cat # DLM-

2640). Tissue explants were cultured for 7 days in hypoxia (1% O_2 and 5% CO_2 in air) at 37^0C

before being washed with PBS and frozen until use.

Protein extraction and preparation for cadaveric and SILAC samples

The frozen samples were pulverised using a freezer mill (Spex) under liquid nitrogen. Samples

were extracted using 15 volumes (w/v) of extraction buffer (4M guanidine hydrochloride (GuHCl),

50 mM sodium acetate, 100 mM 6-aminocaproic acid, and HALT protease inhibitor cocktail

(Thermo Fischer Scientific), pH 5.8). Samples were mechanically dissociated with 10 freeze-thaw

cycles and sonicated in a cold water bath, before extraction with gentle agitation at 4°C for 48

hours. Samples were centrifuged at 15,000g for 30 minutes at 4°C and the supernatant was ethanol

precipitated at a ratio of 1:9 for 16 hours at -20°C. The ethanol step was repeated and samples were

centrifuged at 5000 g for 45 min at 4°C, and the protein pellets were air dried for 30 min.

Protein pellets were re-suspended in fresh 4M urea in 50 mM ammonium bicarbonate, pH 8, using

water bath sonication to aid in the re-solubilisation of the samples. Samples underwent reduction

with TCEP (5mM final concentration) at 60°C for 1 hr, and alkylation with iodoacetamide (500

mM final concentration) for 20 min at RT. Protein concentration was measured using the BCA

assay (Biorad) according to manufacturer's instructions. 200 μg of protein was then buffer

exchanged with 50 mM ammonium bicarbonate with centricon filters (Millipore, 30 kDa cutoff)

according to manufacturer's instructions. Samples were digested with mass spec grade

Trypsin/LysC (Promega) as per manufacturer's instructions. For SILAC-labelled samples, formic

acid was added to a final concentration of 1%, and centrifuged and the supernatant then desalted

prior to LC-MS/MS measurements. For the cadaveric samples, the digested peptides were then

acidified with TFA (0.1% final concentration) and quantified using the peptide quantitative

colorimetric peptide assay kit (Pierce, catalogue 23275) before undergoing fractionation using the

High pH reversed phase peptide fractionation kit (Pierce, catalogue number 84868) into four

fractions. Desalted peptides, were dried, re-suspended in 0.1% formic acid prior to LC-MS/MS

measurements.

883 *Mass spectrometry for cadaveric and SILAC samples*

Samples were loaded onto the Dionex UltiMate 3000 RSLC nano Liquid Chromatography coupled

to the Orbitrap Fusion Lumos Tribid Mass Spectrometer. Peptides were separated on a commercial

Acclaim C18 column (75 μm internal diameter × 50 cm length, 1.9 μm particle size; Thermo).

Separation was attained using a linear gradient of increasing buffer B (80% ACN and 0.1% formic

acid) and declining buffer A (0.1% formic acid) at 300 nL/min. Buffer B was increased to 30% B

in 210 min and ramped to 40% B in 10 min followed by a quick ramp to 95% B, where it was held

for 5 min before a quick ramp back to 5% B, where it was held and the column was re-equilibrated.

Mass spectrometer was operated in positive polarity mode with capillary temperature of 300°C.
Full survey scan resolution was set to 120 000 with an automatic gain control (AGC) target value
of 2×10^6, maximum ion injection time of 30 ms, and for a scan range of 400−1500 m/z. Data
acquisition was in DDA mode to automatically isolate and fragment topN multiply charged
precursors according to their intensities. Spectra were obtained at 30000 MS2 resolution with AGC
target of 1×10^5 and maximum ion injection time of 100 ms, 1.6 m/z isolation width, and
normalised collisional energy of 31. Preceding precursor ions targeted for HCD were dynamically
excluded of 50 s.

Label free quantitative data processing for cadaveric samples

Raw data were analysed using MaxQuant (v.1.6.3.3, Germany). Briefly, raw files were searched
using Andromeda search engine against human UniProt protein database (20,395 entries, Oct
2018), supplemented with sequences of contaminant proteins. Andromeda search parameters for
protein identification were set to a tolerance of 6 ppm for the parental peptide, and 20 ppm for
fragmentation spectra and trypsin specificity allowing up to 2 miscleaved sites. Oxidation of
methionine, carboxyamidomethylation of cysteines was specified as a fixed modification. Minimal
required peptide length was specified at 7 amino acids. Peptides and proteins detected by at least
2 label-free quantification (LFQ) ion counts for each peptide in one of the samples were accepted,
with a false discovery rate (FDR) of 1%. Proteins were quantified by normalised summed peptide
intensities computed in MaxQuant with the LFQ option enabled. A total of 66 profiles were
obtained: 11 locations × 3 disc levels × 2 individuals; with a median of 665 proteins (minimum
419, maximum 1920) per profile.

Data processing for SILAC samples

The high resolution, high mass accuracy mass spectrometry (MS) data obtained were processed
using Proteome Discoverer (Ver 2.1), wherein data were searched using Sequest algorithm against
Human UniProt database (29,900 entries, May 2016), supplemented with sequences of
contaminant proteins, using the following search parameters settings: oxidized methionine (M),
acetylation (Protein N-term), heavy Arginine (R6) and Lysine (K4) were selected as dynamic
modifications, carboxyamidomethylation of cysteines was specified as a fixed modification,
minimum peptide length of 7 amino acids was enabled, tolerance of 10 ppm for the parental

peptide, and 20 ppm for fragmentation spectra, and trypsin specificity allowing up to 2 miscleaved

sites. Confident proteins were identified using a target-decoy approach with a reversed database,

strict FDR 1% at peptide and PSM level. Newly synthesised proteins were heavy labelled with

Arg6- and Lys4 and the data was expressed as the normalised protein abundance obtained from

heavy (labelled)/light (un-labelled) ratio.

Degradome sample preparation, mass spectrometry and data processing

Degradome analyses was performed on NP and AF from three non-degenerated and three

degenerated individuals (Table 1). Frozen tissues were pulverised as described above and prepared

for TAILS as previously reported (Kleifeld et al., 2010). After extraction with SDS buffer (1%

SDS, 100 mM dithiothreitol, 1X protease inhibitor in deionised water) and sonication (three cycles,

15s/cycle), the supernatant (soluble fraction) underwent reduction at 37^0C and alkylation with a

final concentration of 15mM iodoacetamide for 30 min at RT. Samples were precipitated using

chloroform/methanol, and the protein pellet air dried. Samples were re-suspended in 1M NaOH,

quantified by nanodrop, diluted to 100mM HEPES and 4M GnHCl and pH adjusted pH6.5-7.5)

prior to 6-plex TMT labelling as per manufacturer's instructions (Sixplex TMT, Cat# 90061,

ThermoFisher Scientific). Equal ratios of TMT-labelled samples were pooled and

methanol/chloroform precipitated. Protein pellets were air-dried and re-suspended in 200mM

HEPES (pH8), and digested with trypsin (1:100 ratio) for 16 hr at 37^0C, pH 6.5 and a sample was

taken for pre-TAILS. High-molecular-weight dendritic polyglycerol aldehyde polymer (ratio of

5:1 w/w polymer to sample) and NaBH$_3$CN (to a final concentration of 80 mM) was added,

incubated at 37^0C for 16 hr, followed by quenching with 100 mM ethanolamine (30 min at 37^0C)

and underwent ultrafiltration (MWCO of 10,000). Collected samples were desalted, acidified to

0.1% formic acid and dried, prior to MS analysis.

Samples were analysed on a Thermo Scientific Easy nLC-1000 coupled online to a Bruker

Daltonics Impact II UHR QTOF. Briefly, peptides were loaded onto a 20cm x 75µm I.D. analytical

column packed with 1.8µm C18 material (Dr. Maisch GmbH, Germany) in 100% buffer A (99.9%

H$_2$O, 0.1% formic acid) at 800 bar followed by a linear gradient elution in buffer B (99.9%

acetonitrile, 0.1% formic acid) to a final 30% buffer B for a total 180 min including washing with

95% buffer B. Eluted peptides were ionized by ESI and peptide ions were subjected to tandem MS

analysis using a data-dependent acquisition method. A top17 method was employed, where the top

17 most intense multiply charged precursor ions were isolated for MS/MS using collision-induced-

dissociation, and actively excluded for 30s.

MGF files were extracted and searched using Mascot against the UniProt Homo sapiens database,

with semi-ArgC specificity, TMT6plex quantification, variable oxidation of methionine, variable

acetylation of N termini, 20 ppm MS1 error tolerance, 0.05 Da MS2 error tolerance and 2 missed

cleavages. Mascot .dat files were imported into Scaffold Q+S v4.4.3 for peptide identification

processing to a final FDR of 1%. Quantitative values were calculated through Scaffold and used

for subsequent analyses.

Transcriptomic samples: isolation, RNA extraction and data processing

AF and NP tissues from 4 individuals were cut into approximately $0.5cm^3$ pieces, and put into the

Dulbecco's modified Eagle's medium (DMEM) (Gibco) supplemented with 20 mM HEPES

(USB), 1% penicillin-streptomycin (Gibco) and 0.4% fungizone (Gibco). The tissues were

digested with 0.2% pronase (Roche) for 1 hour, and centrifuged at 200 g for 5 min to remove

supernatant. AF and NP were then digested by 0.1% type II collagenase (Worthington

Biochemical) for 14 hours and 0.05% type II collagenase for 8 hours, respectively. Cell suspension

was filtered through a 70 µm cell strainer (BD Falcon) and centrifuged at 200 g for 5 min. The cell

pellet was washed with phosphate buffered saline (PBS) and centrifuged again to remove the

supernatant.RNA was then extracted from the isolated disc cells using Absolutely RNA Nanoprep

Kit (Stratagene), following manufacturer's protocol, and stored at -80°C until further processing.

The quality and quantity of total RNA were assessed on the Bioanalyzer (Agilent) using the RNA

6000 Nano total RNA assay. cDNA was generated using Affymetrix GeneChip Two-Cycle cDNA

Synthesis Kit, followed by *in vitro* transcription to produce biotin-labelled cRNA. The sample was

then hybridised onto the Affymetrix GeneChip Human Genome U133 Plus 2.0 Array. The array

image, CEL file and other related files were generated using Affymetrix GeneChip Command

Console. The experiment was conducted as a service at the Centre for PanorOmic Sciences of the

University of Hong Kong.

CEL and other files were loaded into GeneSpring GX 10 (Agilent) software. The RMA algorithm

was used for probe summation. Data were normalised with baseline transformed to median of all

samples. A loose filtering based on the raw intensity values was then applied to remove

background noise. Consequently, transcriptomic data with a total of 54,675 probes (corresponding

to 20,887 genes) and 8 profiles were obtained.

Bioinformatics and functional analyses

The detected proteins were compared against the transcription factor (TF) database (Vaquerizas et

al., 2009) and the human genome nomenclature consortium database for cell surface markers

(CDs) (Braschi et al., 2019), where 77 TFs and 83 CDs were detected (Supplemental Figure S3A).

Excluding missing values, the LFQ levels among the data-points range from 15.6 to 41.1, with a

Gaussian-like empirical distribution (Supplemental Figure S5A). The numbers of valid values per

protein were found to decline rapidly when they were sorted in descending order (Supplemental

Figure S5B, upper panel). To perform principal component analyses (PCAs), only a subset of

genes with sufficiently large numbers of valid values (i.e. non-missing values) were used. The cut-

off for this was chosen based on a point corresponding to the steepest slope of descending order

of valid protein numbers (Supplemental Figure S5B, second panel), such that the increase of valid

values is slower than the increase of missing values beyond that point. Subsequently, the top 507

genes were picked representing 59.8% of all valid values. This new subset includes 12.4% of all

missing values. Since the subset of data still contains some missing values, an imputation strategy

was adopted employing the Multiple Imputation by Chained Equations (MICE) method and

package (van Buuren and Groothuis-Oudshoorn, 2011), with a max iteration set at 50 and the

default PMM method (predictive mean matching). To further ensure normality, Winsorisation was

applied such that genes whose average is below 5% or above 95% of all genes were also excluded

from PCA. The data was then profile-wise standardised (zero-mean and 1 standard deviation)

before PCA was applied on the R platform (Team, 2013).

To assess the impact of the spatiotemporal factors on the proteomic profiles, we performed

Analysis of Variance (ANOVA), correlating each protein to the age, compartments, level, and

directionality. To draw the soft boundaries on the PCA plot between groups of samples, support

vector machines with polynomial (degree of 2) kernel were applied using the LIBSVM package

(Chang and Lin, 2011) and the PCA coordinates as inputs for training. A meshed grid covering the

whole PCA field was created to make prediction and draw probability contours for -0.5, 0, and 0.5

from the fitted model. Hierarchical clustering was performed with (1- correlation coefficient) as

the distance metrics unless otherwise specified.

To address the problem of 'dropout' effects while avoiding extra inter-dependency introduced due

to imputations, we adopted three strategies in calculating the differentially expressed proteins

(DEPs), namely the statistical DEPs (sDEPs), exclusive DEPs (eDEPs), and the fold-change DEPs

(fDEPs). First, for the proteins that have over half valid values in both groups under comparison,

we performed t-testing with p-values adjusted for multiple testing by the false discovery rate

(FDR). Those with FDR below 0.05 were considered sDEPs. Second, for the proteins where one

group has some valid values while the other group is completely not detected, we considered the

ones with over half valid values in one group to be eDEPs. For those proteins that were expressed

in <50% in both groups, we considered the ones with fold-change greater than 2 to be fDEPs.

To fit the lateral and anteroposterior trends for the modules of genes identified in the young

samples, a Gaussian Process Estimation (GPE) model was trained using the GauPro package in R

(Team, 2013). Pathway analyses was conducted on the GSEA (Subramanian et al., 2005).

Signalling proteins was compiled based on 25 Signal transduction pathways listed on KEGG

(Kanehisa et al., 2019).

The LASSO model between MRI and proteome was trained using the R package "glmnet",

wherein the 85 ECMs were first imputed for missing values in them using MICE. Nine ECMs

were not imputed for too many missing values, leaving 76 for training and testing. The best value

for λ was determined by cross-validations. A model was then trained on the aged MRIs (dependent

variable) and aged proteome of the 76 genes (independent variable). The fitted model was then

applied to the young proteome to predict MRIs of the young discs.

Raw data depository and software availability

The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium

via the PRIDE (Vizcaino et al., 2016) repository with the following dataset identifiers for cadaver

samples   (PXD017774),   SILAC   samples   (PXD018193),   and   degradome   samples

(PXD018298000). The RAW data for the transcriptome data has been deposited on NCBI GEO

with accession number GSE147383.

1035

1036

 Tables

 Table 1. Summary of disc samples.

Samples	Age	Sex	Disc level/s	Disc regions	Reason for surgery
Cadaver samples					
Young spine	16	M	L3/4, L4/5, L5/S1	NP, NP/IAF, IAF, OAF	N/A
Aged spine	59	M	L3/4, L4/5, L5/S1	NP, NP/IAF, IAF, OAF	N/A
Transcriptome samples					
YND74	17	M	L1/2	NP, OAF	Scoliosis
YND88	16	M	L1/2	NP, OAF	Scoliosis
AGD40	62	F	L4/5	NP, OAF	Degeneration
AGD45	47	M	L4/5	NP, OAF	Degeneration
SILAC samples					
YND148	19	F	L2/3	OAF	Scoliosis
YND149	15	F	L1/2	OAF	Scoliosis
YND151	15	F	L1/2	NP, OAF	Scoliosis
YND152	14	F	L1/2	NP, OAF	Scoliosis
AGD80	63	M	L4/5	NP, OAF	Degeneration
Degradome samples					
YND136	17	F	L1/2	NP, OAF	Scoliosis
YND141	20	F	L1/2	NP, OAF	Scoliosis
YND143	53	M	L1/2	NP, OAF	Trauma
AGD62	55	F	L5/S1	NP, OAF	Degeneration
AGD65	68	F	L4/5	NP, OAF	Degeneration
AGD67	55	M	L4/5	NP, OAF	Degeneration

1040   **Table 2.** Commonly expressed ECM and associated proteins across all 66 profiles.

Categories (Number of proteins)	Protein names
Core matrisome	
Collagens (13)	COL1A1/2, COL2A1, COL3A1, COL5A1, COL6A1/2/3, COL11A1/2, COL12A1, COL14A1, COL15A1
Proteoglycans (14)	ACAN, ASPN, BGN, CHAD, DCN, FMOD, HAPLN1, HSPG2, LUM, OGN, OMD, PRELP, PRG4, VCAN
Glycoproteins (34)	ABI3BP, AEBP1, CILP, CILP2, COMP, DPT, ECM2, EDIL3, EFEMP2, EMILIN1, FBN1, FGA, FGB, FN1, FNDC1, LTBP2, MATN2/3, MFGE8, MXRA5, NID2, PCOLCE, PCOLCE2, PXDN, SMOC1/2, SPARC, SRPX2, TGFBI, THBS1/2/4, TNC, TNXB
Other matrisome	
ECM affiliated proteins (10)	ANXA1/2/4/5/6, CLEC11A, CLEC3A/B, CSPG4, SEMA3C
ECM regulators (16)	A2M, CD109, CST3, F13A1, HTRA1, HTRA3, ITIH5, LOXL2/3, PLOD1, SERPINA1/3/5, SERPINE2, SERPING1, TIMP1
Secreted factors (2)	ANGPTL2, FGFBP2

1041

Supplemental Tables

Supplemental Table 1. Raw data of the 66 LC-MS/MS static proteome profiles.

Supplemental Table 2. Differentially expressed proteins (DEPs) among pairs of sample groups
within the 33 young disc profiles.

Supplemental Table 3. Differentially expressed proteins (DEPs) among pairs of sample groups
within the 33 aged disc profiles.

Supplemental Table 4. Differentially expressed proteins (DEPs) between young and aged
sample groups.

Supplemental Table 5. Significantly enriched gene ontology (GO) terms associated with
proteins expressed higher in all young or all aged discs.
1052

 Figure legends

 Figure 1. MRI, outline of workflow, and global overview of data.

 (A) Clinical T2-weighted MRI images of the young lumbar discs in the sagittal and coronal

 plane (left and right panels), T1 MRI image of the young lumbar spine (middle panel).

 (B) High resolution T2-weighted MRI of the aged lower lumbar spine in sagittal (left panel)

 and coronal plane (right pane).

 (C) Diagram showing the anatomy of the IVD and locations from where the samples were

 taken. VB: vertebral body; NP, nucleus pulposus; AF, annulus fibrosus; IAF: inner AF;

 OAF: outer AF; NP/IAF: a transition zone between NP and IAF.

 (D) Schematic diagram showing the structure of the data, and the flow of analyses.

 (E) Venn diagrams showing the overlap of detected proteins in the four major compartments.

 Top panel, young and aged profiles; middle, young only; bottom, aged only.

 (F) Barchart showing the numbers of proteins detected per sample, categorised into

 matrisome (coloured) or non-matrisome proteins (grey).

 (G) Barcharts showing the composition of the matrisome and matrisome-associated proteins.

 Heights of bars indicate the number of proteins in each category expressed per sample.

 The N number in brackets indicate the aggregate number of proteins.

 (H) Violin plots showing the level of sub-categories of ECMs in different compartments of

 the disc. The green number on top of each violin shows its median. LFQ: Label Free

 Quantification.

 (I) Top 30 HGNC gene families for all non-matrisome proteins detected in the dataset.

 (J) Violin plots showing the averaged expression levels of 10 detected histones across the

 disc compartments and age-groups.

 (K) Scatter-plot showing the co-linearity between GAPDH and histones.

 Figure 2. Principle component analysis (PCA) of the 66 profiles based on a set of 507 genes

 selected by optimal cut-off (see Supplemental Figure S5A-C).

 (A) Scatter-plot of PC1 and PC2 color-coded by compartments, and dot-shaped by age-

 groups. Solid curves are the support vector machines (SVMs) decision boundaries

 between inner disc regions (NP, NP/IAF, IAF) and OAF, and dashed curves are soft

 boundaries for probability equal to ±0.5 and are applied to all plots in this figure.

(B) Scatter-plot of PC1 and PC2 color-coded by disc levels. The SVM boundaries are trained

between L5/S1 and upper levels (L3/4 and L4/5).

(C) Scatter-plot of PC1 and PC3, color-coded by disc compartments. The SVM boundaries

are trained between inner disc regions and OAF.

(D) Scatter-plot of PC1 and PC3, color-coded by disc levels. The SVM boundaries are

trained between L5/S1 and upper levels (L3/4 and L4/5).

(E) Top 100 positively and negatively correlated genes with PC1, color-coded by ECM

categories.

(F) Top 100 positively and negatively correlated genes with PC2, color-coded by ECM

categories.

(G) Top 100 positively and negatively correlated genes with PC3, color-coded by ECM

categories.

Figure 3. Delineating the young and healthy disc proteome.

(A) PCA plot of all 33 young profiles. Curves in the upper panel show the SVM boundaries

between the OAF and inner disc regions, those in the lower panel separate the L5/S1 disc

from the upper disc levels. L, left; R, right; A, anterior; P, posterior.

(B) A schematic illustrating the partitioning of the detected human disc proteome into

variable and constant sets.

(C) A histogram showing the distribution of non-DEPs in terms of their detected frequencies

in the young discs. Only 245 non-DEP proteins were detected in over 16 profiles, which

is thus defined to be the constant set; while the remaining ~1,000 proteins were

considered marginally detected.

(D) Piecharts showing the ECM compositions in the variable (left) and constant (right) sets.

The constant set proteins that were detected in all 33 young profiles are listed at the

bottom.

(E) Normalised expression (Z-scores) of proteins in the young module Y1 (NP signature)

laterally (top panel) and anteroposterially (bottom panel), for all three disc levels

combined. The red curve is the Gaussian Process Estimation (GPE) trendline, and the

blue curves are 1 standard deviation above or below the trendline.

(F) Lateral trends of module Y2 (AF signature) for each of the three disc levels.

(G) Lateral trends of module Y3 (Smooth muscle cell signature) for each of the three disc

levels.

(H) Lateral trends of module Y4 (Immune and blood) for each of the three disc levels.

(I) Volcano plot of differentially expressed proteins (DEPs) between OAF and inner disc (an

aggregate of NP, NP/IAF, IAF), with coloured dots representing DEPs.

(J) A functional categorisation of the DEPs in (I).

Figure 4. Characterisation of the aged disc proteome.

(A) PCA plot of all the aged profiles on PC1 and PC2, color-coded by compartments. Curves

in the left panel show the SVM boundaries between OAF and inner disc; those in the

right panel separate the L5/S1 disc from the upper disc levels. Letters on dots indicate

directions: L, left; R, right; A, anterior; P, posterior.

(B) Volcano plot showing the DEPs between the OAF and inner disc (an aggregate of NP,

NP/IAF and IAF), with the coloured dots representing statistically significant

(FDR<0.05) DEPs.

(C) Using the same 4 modules identified in young samples, we determined the trend for these

in the aged samples. Locational trends of module Y1 showing higher expression in the

inner disc, albeit they are more flattened than in the young disc samples. Top panel shows

left to right direction and bottom panel shows anterior to posterior direction. The red

curve is the Gaussian Process Estimation (GPE) trendline, and the blue curves are 1

standard deviation above or below the trendline. This also applies to (D), (E) and (F).

(D) Lateral trends for module Y2 in the aged discs.

(E) Lateral trends for module Y3 in the aged discs.

(F) Lateral trends for module Y4 in the aged discs.

Figure 5. Comparison between young and aged samples.

(A) Volcano plot showing the DEPs between all the 33 young and 33 aged profiles. Coloured

dots represent statistically significant DEPs.

(B) GO term enrichment of DEPs higher in young profiles.

(C) GO term enrichment of DEPs higher in aged profiles. Full names of GO terms in (B) and

(C) are listed in Supplemental Table S5.

(D) Volcano plot showing DEPs between aged and young inner disc regions.

(E) Volcano plot showing DEPs between aged and young OAF.

(F) Venn diagram showing the partitioning of the young/aged DEPs that were down-

regulated in aged discs, into contributions from inner disc regions and OAF.

(G) Venn diagram showing the partitioning of the young/aged DEPs that were up-regulated

in aged discs, into contributions from inner disc regions and OAF.

(H) A heat map showing proteins expressed in all young and aged disc, with the

identification of 6 modules (module 1: higher expression in young inner disc regions,

modules 2 and 4: higher expression in young OAF, module 3: highly expressing in aged

OAF, module 5: higher expression across all aged samples, and module 6: higher

expression in aged inner disc, and some OAF).

(I) An alluvial chart showing the six modules identified in (H) and their connections to the

previously identified four modules and constant set in the young reference proteome; as

well as their connections to enriched GO terms.

Figure 6. Concordance between proteomic and transcriptome data.

(A) A PCA plot of the 8 transcriptomic profiles. Curves represent SVM boundaries between

patient-groups or compartments.

(B) Venn diagrams showing the partitioning of the young/aged DEGs into contributions from

inner disc regions and OAF. Left: down-regulated in AGD samples; right: up-regulated.

(C) Transcriptome data from the NP and AF of two young individuals were compared to the

proteomic data, with coloured dots representing identified proteins also expressed at the

transcriptome level.

(D) Transcriptome and proteome comparison of aged OAF and NP.

(E) Transcriptome and proteome comparison of young and aged OAF.

(F) Transcriptome and proteome comparison of young and aged NP.

Figure 7. The dynamic proteome of the intervertebral disc shows less biosynthesis of proteins in

aged tissues.

(A) Schematic showing pulse-SILAC labelling of *ex-vivo* cultured disc tissues where heavy

Arg and Lys are incorporated into newly made proteins (heavy), and pre-existing proteins

1171   remaining unlabelled (light). NP and AF tissues from young (n=3) and aged (n=1) were

1172    cultured for 7 days in hypoxia prior to MS.

1173  (B) Barcharts showing the number of identified non-matrisome (grey) and matrisome

1174    (coloured) existing proteins (middle panel); newly synthesised proteins (left panel), and

1175    the heavy/light ratio (right panel) for each of the samples.

1176  (C) The quantities of each of the heavy labelled (newly synthesised) proteins identified for

1177    each of the four groups were averaged, and then plotted in descending order of

1178    abundance. It shows that YND AF and NP synthesise higher numbers of proteins than the

1179    AGD AF and NP. The red dotted reference line shows the expression of GAPDH.

1180  (D) The quantities of each of the light (existing) proteins identified for each group was

1181    averaged, and then plotted in descending order of abundance which shows that there are

1182    similar levels of existing proteins in the four pooled samples.

1183  (E) The matrisome proteins of (C) were singled out for display. The abundance of these

1184    proteins in YND samples were generally higher across all types of matrisome proteins

1185    than the AGD, with the exceptions of aged related proteins.

1186  (F) Schematic showing the workflow of degradome analysis by N-terminal amine isotopic

1187    labelling (TAILS) for the identification of cleaved neo N-terminal peptides.

1188  (G) Identification of cleaved proteins ranked according to tandem mass tag (TMT) isobaric

1189    labelling of N-terminal peptides in NP. The numbers of N-terminal peptides identified

1190    from corresponding proteins (x axis) were plotted in increasing ratios, with those that

1191    were less cleaved in AGD NP to the left, whereas proteins cleaved at a greater number of

1192    sites or in increased amount per site in AGD NP to the right and included a higher

1193    proportion of ECM and plasma protein N-terminal peptides that were identified. Data is

1194    expressed as the $\log_2$(ratio) of N-terminal peptides relative to the YND136 sample. n.s.:

1195    not significant.

1196  (H) Comparison of protein degradation in AF. The numbers of peptides identified from

1197    corresponding proteins (x axis) were plotted in increasing ratios, with those that were less

1198    cleaved in AGD AF to the left, whereas proteins cleaved at a greater number of sites or in

1199    increased amount per site in AGD AF to the right. A higher proportion of ECM and

1200    plasma protein N-terminal peptides were also identified in AGD samples, whilst in YND

AF, the highest amount of detected peptides came from collagens. Data is expressed as

the $\log_2$(ratio) relative to the YND136 sample.

Figure 8. MRI intensities and their correlation with the proteomic data.

(A) The middle MRI stack of each disc level in the aged cadaveric sample.

(B) Schematic of the disc showing the three stacks of MRI images per disc.

(C) Violin plots showing the pixel intensities within each location per disc level,

corresponding to the respective locations taken for mass spectrometry measurements.

Each violin-plot is the aggregate of three stacks of MRIs per disc.

(D) A heatmap bi-clustering of levels and compartments based on the MRI intensities.

(E) The hydration ECMs: the ECM proteins most positively and negatively correlated with

MRI.

(F) The 3T MRI intensities of the young discs across the compartments (left), and the

predicted MRI intensities based on a LASSO regression model trained on the hydration

ECMs (right).

(G) A water-tank model of the dynamics in disc proteomics showing the balance of the

proteome is maintained by adequate anabolism to balance catabolism.

(H) Diagram showing the partitioning of the detected proteins into variable and constant sets,

whereby four modules characterising the young healthy disc were further derived; and

showing their changes with ageing. SMC: smooth muscle cell markers.

Author contributions

VT handled IRB/IC, coordinated MRI measurements, performed tissue dissection, sample preparation, data processing (MaxQuant), analyses (Perseus) and interpretation, prepared figures. PKC performed bioinformatics analyses (ANOVA, PCA, SVM, DEGs, GPE, LASSO), and prepared figures. VT and PKC wrote the first draft of the manuscript. AY generated the microarray data. RS, NS and TK performed the mass spectrometry and processed the data, supervised and funded by CMO. MK, PS and WC were involved in interpretation. LH provided the young spine and interpreted the data. KC provided critical input, interpreted results and was involved in writing. DC conceived ideas, supervised the project, interpreted data and results, wrote the manuscript. All authors contributed to writing and approved the manuscript.

Acknowledgments

We thank Dr Ed Wu and Dr Anna Wang of the Dept of EEE at HKU for performing the high-resolution MRI on the aged discs. We thank Dr Dino Samartzis for arranging the MRI of the young lumbar spine, and Prof. Kenneth Cheung and Dr Jason Cheung for collecting surgical disc specimens. Part of this work was supported by the Theme-based Research Scheme (T12-708/12N) and Area of Excellence (AoE/M-04/04) of the Hong Kong Research Grants (RGC) Council awarded to KSEC (Project Coordinator and PI) and DC (co-PI), the RGC European Union - Hong Kong Research and Innovation Cooperation Co-funding Mechanism (E-HKU703/18) awarded to DC, and by the Ministry of Science and Technology of the People's Republic of China: National Strategic Basic Research Program ("973") (2014CB942900) awarded to DC. The TAILS analyses were supported by a Canadian Institutes of Health Research Foundation Grant FDN-148408 to CMO, who holds a Canada Research Chair in Protease Proteomics and Systems Biology.

References

Anderson, R.M., Lawrence, A.R., Stottmann, R.W., Bachiller, D., and Klingensmith, J. (2002). Chordin
and noggin promote organizing centers of forebrain development in the mouse. Development *129*,
4975-4987.
Barber, R.D., Harmer, D.W., Coleman, R.A., and Clark, B.J. (2005). GAPDH as a housekeeping gene:
analysis of GAPDH mRNA expression in a panel of 72 human tissues. Physiol Genomics *21*, 389-
395.
Bizet, A.A., Liu, K., Tran-Khanh, N., Saksena, A., Vorstenbosch, J., Finnson, K.W., Buschmann, M.D.,
and Philip, A. (2011). The TGF-beta co-receptor, CD109, promotes internalization and degradation
of TGF-beta receptors. Biochim Biophys Acta *1813*, 742-753.
Braschi, B., Denny, P., Gray, K., Jones, T., Seal, R., Tweedie, S., Yates, B., and Bruford, E. (2019).
Genenames.org: the HGNC and VGNC resources in 2019. Nucleic Acids Res *47*, D786-D792.
Chang, C.C., and Lin, C.J. (2011). LIBSVM: A Library for Support Vector Machines. Acm T Intel Syst
Tec *2*.
Chen, J., Yan, W., and Setton, L.A. (2006). Molecular phenotypes of notochordal cells purified from
immature nucleus pulposus. Eur Spine J *15 Suppl 3*, S303-311.
Dong, Y.F., Soung do, Y., Schwarz, E.M., O'Keefe, R.J., and Drissi, H. (2006). Wnt induction of
chondrocyte hypertrophy through the Runx2 transcription factor. J Cell Physiol *208*, 77-86.
Feng, H., Danfelter, M., Stromqvist, B., and Heinegard, D. (2006). Extracellular matrix in disc
degeneration. J Bone Joint Surg Am *88 Suppl 2*, 25-29.
Fortelny, N., Overall, C.M., Pavlidis, P., and Freue, G.V.C. (2017). Can we predict protein from mRNA
levels? Nature *547*, E19-E20.
Fujita, N., Miyamoto, T., Imai, J., Hosogane, N., Suzuki, T., Yagi, M., Morita, K., Ninomiya, K.,
Miyamoto, K., Takaishi, H.*, et al.* (2005). CD24 is expressed specifically in the nucleus pulposus
of intervertebral discs. Biochem Biophys Res Commun *338*, 1890-1896.
Grimsrud, C.D., Romano, P.R., D'Souza, M., Puzas, J.E., Schwarz, E.M., Reynolds, P.R., Roiser, R.N., and
O'Keefe, R.J. (2001). BMP signaling stimulates chondrocyte maturation and the expression of
Indian hedgehog. J Orthop Res *19*, 18-25.
Hiyama, A., Sakai, D., Risbud, M.V., Tanaka, M., Arai, F., Abe, K., and Mochida, J. (2010). Enhancement
of intervertebral disc cell senescence by WNT/beta-catenin signaling-induced matrix
metalloproteinase expression. Arthritis Rheum *62*, 3036-3047.
Hou, G., Lu, H., Chen, M., Yao, H., and Zhao, H. (2014). Oxidative stress participates in age-related
changes in rat lumbar intervertebral discs. Arch Gerontol Geriatr *59*, 665-669.
Humzah, M.D., and Soames, R.W. (1988). Human intervertebral disc: structure and function. Anat Rec
220, 337-356.
Jayasuriya, C.T., Goldring, M.B., Terek, R., and Chen, Q. (2012). Matrilin-3 induction of IL-1 receptor
antagonist is required for up-regulating collagen II and aggrecan and down-regulating ADAMTS-
5 gene expression. Arthritis Res Ther *14*, R197.
Ji, Q., Zheng, Y., Zhang, G., Hu, Y., Fan, X., Hou, Y., Wen, L., Li, L., Xu, Y., Wang, Y.*, et al.* (2019).
Single-cell RNA-seq analysis reveals the progression of human osteoarthritis. Ann Rheum Dis *78*,
100-110.
Jim, J.J., Noponen-Hietala, N., Cheung, K.M., Ott, J., Karppinen, J., Sahraravand, A., Luk, K.D., Yip, S.P.,
Sham, P.C., Song, Y.Q.*, et al.* (2005). The TRP2 allele of COL9A2 is an age-dependent risk factor
for the development and severity of intervertebral disc degeneration. Spine (Phila Pa 1976) *30*,
2735-2742.
Johnson, J.L., Hall, T.E., Dyson, J.M., Sonntag, C., Ayers, K., Berger, S., Gautier, P., Mitchell, C., Hollway,
G.E., and Currie, P.D. (2012). Scube activity is necessary for Hedgehog signal transduction in vivo.
Dev Biol *368*, 193-202.

Kalamajski, S., Bihan, D., Bonna, A., Rubin, K., and Farndale, R.W. (2016). Fibromodulin Interacts with
Collagen Cross-linking Sites and Activates Lysyl Oxidase. J Biol Chem *291*, 7951-7960.
Kanehisa, M., Sato, Y., Furumichi, M., Morishima, K., and Tanabe, M. (2019). New approach for
understanding genome variations in KEGG. Nucleic Acids Res *47*, D590-D595.
Kleifeld, O., Doucet, A., auf dem Keller, U., Prudova, A., Schilling, O., Kainthan, R.K., Starr, A.E., Foster,
L.J., Kizhakkedathu, J.N., and Overall, C.M. (2010). Isotopic labeling of terminal amines in
complex samples identifies protein N-termini and protease cleavage products. Nat Biotechnol *28*,
281-288.
Komori, T. (2010). Regulation of bone development and extracellular matrix protein genes by RUNX2.
Cell Tissue Res *339*, 189-195.
Lam, T.-k. (2013). Fate of notochord descendent cells in the intervertebral disc. In HKU Theses Online
(HKUTO) (The University of Hong Kong (Pokfulam, Hong Kong)).
Lau, D., Elezagic, D., Hermes, G., Morgelin, M., Wohl, A.P., Koch, M., Hartmann, U., Hollriegl, S.,
Wagener, R., Paulsson, M.*, et al.* (2018). The cartilage-specific lectin C-type lectin domain family
3 member A (CLEC3A) enhances tissue plasminogen activator-mediated plasminogen activation.
J Biol Chem *293*, 203-214.
Lee, C.G., Da Silva, C.A., Dela Cruz, C.S., Ahangari, F., Ma, B., Kang, M.J., He, C.H., Takyar, S., and
Elias, J.A. (2011). Role of chitin and chitinase/chitinase-like proteins in inflammation, tissue
remodeling, and injury. Annu Rev Physiol *73*, 479-501.
Leijten, J.C., Bos, S.D., Landman, E.B., Georgi, N., Jahr, H., Meulenbelt, I., Post, J.N., van Blitterswijk,
C.A., and Karperien, M. (2013). GREM1, FRZB and DKK1 mRNA levels correlate with
osteoarthritis and are regulated by osteoarthritis-associated factors. Arthritis Res Ther *15*, R126.
Li, C., Hancock, M.A., Sehgal, P., Zhou, S., Reinhardt, D.P., and Philip, A. (2016). Soluble CD109 binds
TGF-beta and antagonizes TGF-beta signalling and responses. Biochem J *473*, 537-547.
Lopez-Otin, C., and Overall, C.M. (2002). Protease degradomics: a new challenge for proteomics. Nat Rev
Mol Cell Biol *3*, 509-519.
Lu, Y., Qiao, L., Lei, G., Mira, R.R., Gu, J., and Zheng, Q. (2014). Col10a1 gene expression and
chondrocyte hypertrophy during skeletal development and disease. Frontiers in Biology *9*, 195-
204.
Markmann, A., Hausser, H., Schonherr, E., and Kresse, H. (2000). Influence of decorin expression on
transforming growth factor-beta-mediated collagen gel retraction and biglycan induction. Matrix
Biol *19*, 631-636.
Maseda, M., Yamaguchi, H., Kuroda, K., Mitsumata, M., Tokuhashi, Y., and Esumi, M. (2016). Proteomic
Analysis of Human Intervertebral Disc Degeneration. Journal of Nihon University Medical
Association *75*, 16-21.
Melas, I.N., Chairakaki, A.D., Chatzopoulou, E.I., Messinis, D.E., Katopodi, T., Pliaka, V., Samara, S.,
Mitsos, A., Dailiana, Z., Kollia, P.*, et al.* (2014). Modeling of signaling pathways in chondrocytes
based on phosphoproteomic and cytokine release data. Osteoarthritis Cartilage *22*, 509-518.
Minogue, B.M., Richardson, S.M., Zeef, L.A., Freemont, A.J., and Hoyland, J.A. (2010). Characterization
of the human nucleus pulposus cell phenotype and evaluation of novel marker gene expression to
define adult stem cell differentiation. Arthritis Rheum *62*, 3695-3705.
Molinos, M., Almeida, C.R., Caldeira, J., Cunha, C., Goncalves, R.M., and Barbosa, M.A. (2015).
Inflammation in intervertebral disc degeneration and regeneration. J R Soc Interface *12*, 20150429.
Munir, S., Rade, M., Maatta, J.H., Freidin, M.B., and Williams, F.M.K. (2018). Intervertebral Disc Biology:
Genetic Basis of Disc Degeneration. Curr Mol Biol Rep *4*, 143-150.
Naba, A., Clauser, K.R., Hoersch, S., Liu, H., Carr, S.A., and Hynes, R.O. (2012). The matrisome: in silico
definition and in vivo characterization by proteomics of normal and tumor extracellular matrices.
Mol Cell Proteomics *11*, M111 014647.
Nakai, T., Sakai, D., Nakamura, Y., Nukaga, T., Grad, S., Li, Z., Alini, M., Chan, D., Masuda, K., Ando,
K.*, et al.* (2016). CD146 defines commitment of cultured annulus fibrosus cells to express a
contractile phenotype. J Orthop Res *34*, 1361-1372.

Nakamichi, R., Kataoka, K., and Asahara, H. (2018). Essential role of Mohawk for tenogenic tissue
homeostasis including spinal disc and periodontal ligament. Mod Rheumatol *28*, 933-940.
Naveau, S., Poynard, T., Benattar, C., Bedossa, P., and Chaput, J.C. (1994). Alpha-2-macroglobulin and
hepatic fibrosis. Diagnostic interest. Dig Dis Sci *39*, 2426-2432.
Nerlich, A.G., Schaaf, R., Walchli, B., and Boos, N. (2007). Temporo-spatial distribution of blood vessels
in human lumbar intervertebral discs. Eur Spine J *16*, 547-555.
Newell, N., Little, J.P., Christou, A., Adams, M.A., Adam, C.J., and Masouros, S.D. (2017). Biomechanics
of the human intervertebral disc: A review of testing techniques and results. J Mech Behav Biomed
Mater *69*, 420-434.
Ong, S.E., Blagoev, B., Kratchmarova, I., Kristensen, D.B., Steen, H., Pandey, A., and Mann, M. (2002).
Stable isotope labeling by amino acids in cell culture, SILAC, as a simple and accurate approach
to expression proteomics. Mol Cell Proteomics *1*, 376-386.
Onnerfjord, P., Khabut, A., Reinholt, F.P., Svensson, O., and Heinegard, D. (2012a). Quantitative
proteomic analysis of eight cartilaginous tissues reveals characteristic differences as well as
similarities between subgroups. J Biol Chem *287*, 18913-18924.
Onnerfjord, P., Khabut, A., Reinholt, F.P., Svensson, O., and Heinegard, D. (2012b). Quantitative
proteomic analysis of eight cartilaginous tissues reveals characteristic differences as well as
similarities between subgroups. Journal of Biological Chemistry *287*, 18913-18924.
Park, J.S., Chu, J.S., Tsou, A.D., Diop, R., Tang, Z., Wang, A., and Li, S. (2011). The effect of matrix
stiffness on the differentiation of mesenchymal stem cells in response to TGF-beta. Biomaterials
32, 3921-3930.
Peix, L., Evans, I.C., Pearce, D.R., Simpson, J.K., Maher, T.M., and McAnulty, R.J. (2018). Diverse
functions of clusterin promote and protect against the development of pulmonary fibrosis. Sci Rep
8, 1906.
Pfirrmann, C.W., Metzdorf, A., Zanetti, M., Hodler, J., and Boos, N. (2001). Magnetic resonance
classification of lumbar intervertebral disc degeneration. Spine (Phila Pa 1976) *26*, 1873-1878.
Rajasekaran, S., Tangavel, C., K, S.S., Soundararajan, D.C.R., Nayagam, S.M., Matchado, M.S.,
Raveendran, M., Shetty, A.P., Kanna, R.M., and Dharmalingam, K. (2020). Inflammaging
determines health and disease in lumbar discs-evidence from differing proteomic signatures of
healthy, aging, and degenerating discs. Spine J *20*, 48-59.
Rajesh, D., and Dahia, C.L. (2018). Role of Sonic Hedgehog Signaling Pathway in Intervertebral Disc
Formation and Maintenance. Curr Mol Biol Rep *4*, 173-179.
Ranjani, V., Sreemol, G., Muthurajan, R., Natesan, S., Gnanam, R., Kanna, R.M., and Rajasekaran, S.
(2016). Proteomic Analysis of Degenerated Intervertebral Disc-identification of Biomarkers of
Degenerative Disc Disease and Development of Proteome Database. Global Spine Journal *6*,
WO025.
Rauniyar, N., and Yates, J.R., 3rd (2014). Isobaric labeling-based relative quantification in shotgun
proteomics. J Proteome Res *13*, 5293-5309.
Riester, S.M., Lin, Y., Wang, W., Cong, L., Mohamed Ali, A.M., Peck, S.H., Smith, L.J., Currier, B.L.,
Clark, M., Huddleston, P.*, et al.* (2018). RNA sequencing identifies gene regulatory networks
controlling extracellular matrix synthesis in intervertebral disk tissues. J Orthop Res *36*, 1356-1369.
Risbud, M.V., Schoepflin, Z.R., Mwale, F., Kandel, R.A., Grad, S., Iatridis, J.C., Sakai, D., and Hoyland,
J.A. (2015). Defining the phenotype of young healthy nucleus pulposus cells: recommendations of
the Spine Research Interest Group at the 2014 annual ORS meeting. J Orthop Res *33*, 283-293.
Robinson, K.A., Sun, M., Barnum, C.E., Weiss, S.N., Huegel, J., Shetye, S.S., Lin, L., Saez, D., Adams,
S.M., Iozzo, R.V.*, et al.* (2017). Decorin and biglycan are necessary for maintaining collagen fibril
structure, fiber realignment, and mechanical properties of mature tendons. Matrix Biol *64*, 81-93.
Rodrigues-Pinto, R., Berry, A., Piper-Hanley, K., Hanley, N., Richardson, S.M., and Hoyland, J.A. (2016).
Spatiotemporal analysis of putative notochordal cell markers reveals CD24 and keratins 8, 18, and
19 as notochord-specific markers during early human intervertebral disc development. J Orthop
Res *34*, 1327-1340.

Rodriguez, A.G., Slichter, C.K., Acosta, F.L., Rodriguez-Soto, A.E., Burghardt, A.J., Majumdar, S., and
Lotz, J.C. (2011). Human disc nucleus properties and vertebral endplate permeability. Spine (Phila
Pa 1976) *36*, 512-520.
Rubin, D.I. (2007). Epidemiology and risk factors for spine pain. Neurol Clin *25*, 353-371.
Rutges, J.P., Duit, R.A., Kummer, J.A., Oner, F.C., van Rijen, M.H., Verbout, A.J., Castelein, R.M., Dhert,
W.J., and Creemers, L.B. (2010). Hypertrophic differentiation and calcification during
intervertebral disc degeneration. Osteoarthritis Cartilage *18*, 1487-1495.
Saito, T., Ikeda, T., Nakamura, K., Chung, U.I., and Kawaguchi, H. (2007). S100A1 and S100B,
transcriptional targets of SOX trio, inhibit terminal differentiation of chondrocytes. EMBO Rep *8*,
504-509.
Sakai, D., Nakamura, Y., Nakai, T., Mishima, T., Kato, S., Grad, S., Alini, M., Risbud, M.V., Chan, D.,
Cheah, K.S.*, et al.* (2012). Exhaustion of nucleus pulposus progenitor cells with ageing and
degeneration of the intervertebral disc. Nat Commun *3*, 1264.
Saleem, S., Aslam, H.M., Rehmani, M.A., Raees, A., Alvi, A.A., and Ashraf, J. (2013). Lumbar disc
degenerative disease: disc degeneration symptoms and magnetic resonance image findings. Asian
Spine J *7*, 322-334.
Sarath Babu, N., Krishnan, S., Brahmendra Swamy, C.V., Venkata Subbaiah, G.P., Gurava Reddy, A.V.,
and Idris, M.M. (2016). Quantitative proteomic analysis of normal and degenerated human
intervertebral disc. Spine J *16*, 989-1000.
Schneiderman, G., Flannigan, B., Kingston, S., Thomas, J., Dillin, W.H., and Watkins, R.G. (1987).
Magnetic resonance imaging in the diagnosis of disc degeneration: correlation with discography.
Spine (Phila Pa 1976) *12*, 276-281.
Silagi, E.S., Shapiro, I.M., and Risbud, M.V. (2018). Glycosaminoglycan synthesis in the nucleus pulposus:
Dysregulation and the pathogenesis of disc degeneration. Matrix Biol *71-72*, 368-379.
Song, Y.Q., Cheung, K.M., Ho, D.W., Poon, S.C., Chiba, K., Kawaguchi, Y., Hirose, Y., Alini, M., Grad,
S., Yee, A.F.*, et al.* (2008). Association of the asporin D14 allele with lumbar-disc degeneration in
Asians. Am J Hum Genet *82*, 744-747.
Song, Y.Q., Karasugi, T., Cheung, K.M., Chiba, K., Ho, D.W., Miyake, A., Kao, P.Y., Sze, K.L., Yee, A.,
Takahashi, A.*, et al.* (2013). Lumbar disc degeneration is linked to a carbohydrate sulfotransferase
3 variant. J Clin Invest *123*, 4909-4917.
Subramanian, A., and Schilling, T.F. (2014). Thrombospondin-4 controls matrix assembly during
development and repair of myotendinous junctions. Elife *3*.
Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A., Paulovich, A.,
Pomeroy, S.L., Golub, T.R., Lander, E.S.*, et al.* (2005). Gene set enrichment analysis: a knowledge-
based approach for interpreting genome-wide expression profiles. Proc Natl Acad Sci U S A *102*,
15545-15550.
Sun, Z., Liu, B., and Luo, Z.J. (2020). The Immune Privilege of the Intervertebral Disc: Implications for
Intervertebral Disc Degeneration Treatment. Int J Med Sci *17*, 685-692.
Taha, I.N., and Naba, A. (2019). Exploring the extracellular matrix in health and disease using proteomics.
Essays Biochem *63*, 417-432.
Takao, T., and Iwaki, T. (2002). A comparative study of localization of heat shock protein 27 and heat
shock protein 72 in the developmental and degenerative intervertebral discs. Spine (Phila Pa 1976)
27, 361-368.
Taye, N., Karoulias, S.Z., and Hubmacher, D. (2020). The "other" 15-40%: The Role of Non-Collagenous
Extracellular Matrix Proteins and Minor Collagens in Tendon. J Orthop Res *38*, 23-35.
Team, R.C. (2013). R: A language and environment for statistical computing.
Teraguchi, M., Yoshimura, N., Hashizume, H., Muraki, S., Yamada, H., Minamide, A., Oka, H., Ishimoto,
Y., Nagata, K., Kagotani, R.*, et al.* (2014). Prevalence and distribution of intervertebral disc
degeneration over the entire spine in a population-based cohort: the Wakayama Spine Study.
Osteoarthritis Cartilage *22*, 104-110.
Tibshirani, R. (1996). Regression shrinkage and selection via the Lasso. J Roy Stat Soc B Met *58*, 267-288.

Trougakos, I.P. (2013). The molecular chaperone apolipoprotein J/clusterin as a sensor of oxidative stress:
implications in therapeutic approaches - a mini-review. Gerontology *59*, 514-523.
Urban, J.P., Smith, S., and Fairbank, J.C. (2004). Nutrition of the intervertebral disc. Spine (Phila Pa 1976)
29, 2700-2709.
van Buuren, S., and Groothuis-Oudshoorn, K. (2011). mice: Multivariate Imputation by Chained Equations
in R. J Stat Softw *45*, 1-67.
van den Akker, G.G.H., Koenders, M.I., van de Loo, F.A.J., van Lent, P., Blaney Davidson, E., and van der
Kraan, P.M. (2017). Transcriptional profiling distinguishes inner and outer annulus fibrosus from
nucleus pulposus in the bovine intervertebral disc. Eur Spine J *26*, 2053-2062.
van der Kraan, P.M., and van den Berg, W.B. (2012). Chondrocyte hypertrophy and osteoarthritis: role in
initiation and progression of cartilage degeneration? Osteoarthritis Cartilage *20*, 223-232.
Vaquerizas, J.M., Kummerfeld, S.K., Teichmann, S.A., and Luscombe, N.M. (2009). A census of human
transcription factors: function, expression and evolution. Nat Rev Genet *10*, 252-263.
Veras, M.A., McCann, M.R., Tenn, N.A., and Seguin, C.A. (2020). Transcriptional profiling of the murine
intervertebral disc and age-associated changes in the nucleus pulposus. Connect Tissue Res *61*, 63-
81.
Vizcaino, J.A., Csordas, A., del-Toro, N., Dianes, J.A., Griss, J., Lavidas, I., Mayer, G., Perez-Riverol, Y.,
Reisinger, F., Ternent, T.*, et al.* (2016). 2016 update of the PRIDE database and its related tools.
Nucleic Acids Res *44*, D447-456.
Wang, X.L., Hou, L., Zhao, C.G., Tang, Y., Zhang, B., Zhao, J.Y., and Wu, Y.B. (2019a). Screening of
genes involved in epithelial-mesenchymal transition and differential expression of complement-
related genes induced by PAX2 in renal tubules. Nephrology (Carlton) *24*, 263-271.
Wang, Y., Fan, X., Xing, L., and Tian, F. (2019b). Wnt signaling: a promising target for osteoarthritis
therapy. Cell Commun Signal *17*, 97.
Wisniewski, J.R., Hein, M.Y., Cox, J., and Mann, M. (2014). A "proteomic ruler" for protein copy number
and concentration estimation without spike-in standards. Mol Cell Proteomics *13*, 3497-3506.
Wuertz, K., Vo, N., Kletsas, D., and Boos, N. (2012). Inflammatory and catabolic signalling in
intervertebral discs: the roles of NF-kappaB and MAP kinases. Eur Cell Mater *23*, 103-119;
discussion 119-120.
Wyatt, A.R., Yerbury, J.J., and Wilson, M.R. (2009). Structural characterization of clusterin-chaperone
client protein complexes. J Biol Chem *284*, 21920-21927.
Yates, B., Braschi, B., Gray, K.A., Seal, R.L., Tweedie, S., and Bruford, E.A. (2017). Genenames.org: the
HGNC and VGNC resources in 2017. Nucleic Acids Res *45*, D619-D625.
Yee, A., Lam, M.P., Tam, V., Chan, W.C., Chu, I.K., Cheah, K.S., Cheung, K.M., and Chan, D. (2016).
Fibrotic-like changes in degenerate human intervertebral discs revealed by quantitative proteomic
analysis. Osteoarthritis Cartilage *24*, 503-513.
Zhu, M., Tang, D., Wu, Q., Hao, S., Chen, M., Xie, C., Rosier, R.N., O'Keefe, R.J., Zuscik, M., and Chen,
D. (2009). Activation of beta-catenin signaling in articular chondrocytes leads to osteoarthritis-like
phenotype in adult beta-catenin conditional activation mice. J Bone Miner Res *24*, 12-21.
Zhu, S., Qiu, H., Bennett, S., Kuek, V., Rosen, V., Xu, H., and Xu, J. (2019). Chondromodulin-1 in health,
osteoarthritis, cancer, and heart disease. Cell Mol Life Sci *76*, 4493-4502.
1485

FIGURE 1

FIGURE 2

FIGURE 5

FIGURE 6

FIGURE 7