Whole-proteome Tree of Insects: Grouping and phylogeny without sequence alignment

JaeJin Choi[1,2], Byung-Ju Kim[3] & Sung-Hou Kim[1,2,3,#]

[1]Department of Chemistry and Center for Computational Biology, University of California, Berkeley, CA, 94720, USA, [2] Biological Systems and Engineering Division, and Molecular Biophysics & Integrated Bioimaging Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA, and [3]Human Genome Research Center, Incheon National University, Incheon, Republic of Korea (South Korea)

ORCID: JJ Choi (0000-0002-2860-9606), BJ Kim (0000-0002-8906-5622), SH Kim (0000-0002-7293-5994)

[#]Corresponding author: Sung-Hou Kim; sunghou@berkeley.edu; 510-708-2564

419 Latimer Hall, University of California, Berkeley CA, 94720, USA

e-mail addresses of co-authors: JJC (jaejinchoi@berkeley.edu); BJK (bjk@inu.ac.kr)

Short running title: **Whole-Proteome Tree of Insects**

Keywords: Organism tree, Gene/protein tree, Feature Frequency Profile, Jensen-Shannon divergence, Evolutionary progression scale, Cumulative genomic divergence, Arthropodal burst

Abstract

An "organism tree" of insects, the largest and most species-diverse group of all

living animals, can be considered as a conceptual tree to capture a simplified narrative of

the complex evolutionary courses of the extant insects. Currently, the most common

approach has been to construct a "protein tree", as a surrogate for the organism tree, by

Multiple Sequence Alignment (MSA) of highly homologous regions of a set of select

proteins to represent each organism. However, such selected regions account for a very

small fraction of the whole-proteome of each organism.

Information Theory provides a method of comparing two sets of *all* proteins, two

whole-proteomes, without MSA: By treating each whole-proteome sequence as a "book"

of amino acid alphabets, the information contents of two whole-proteomes can be

quantitatively compared using the text comparison method of the theory, without sequence

alignment, providing an opportunity to construct a "whole-proteome tree" of insects as a

surrogate for an organism tree of insects.

A whole-proteome tree of the insects in this study shows that: (a) all the

founders of the major groups of the insects have emerged in an explosive "burst" near

the root of the tree, (b) the most basal group of all the insects is a subgroup of

Hemiptera consisting of aphids and psyllids, and (c) there are other notable

differences in the phylogeny of the groups compared to those of the recent protein

trees of insects.

Introduction

Sequence-alignment-based "protein trees": An "organism tree" of insects can be

considered as a practically useful narrative to convey a simplified evolutionary

relationship among the insects. However, it is a conceptual tree that cannot be

experimentally validated. Thus, it is expected that the effort will continue to find one or

more "surrogate trees" derived from various descriptors of the characteristics associated

with each insect and to find improved methods to estimate evolutionary distances from

the divergence of the descriptors under as few subjective assumptions as possible at the

time of investigation.

At present, the best descriptor of an insect as an organism is its entire whole-

genome sequence information, from which whole-proteome sequence can be derived.

However, for several decades, due to the technical difficulties and high cost of whole

genome sequencing, and to the difficult task of comparing unaligned sequences, the most

practically feasible and common approach to construct a surrogate tree has been to

construct a Multiple Sequence Alignment (MSA)-based "protein tree" under a few

important, but debatable assumptions: (a) a set of regions with high homology selected

among each of homologous proteins may have enough information to represent a whole

organism, and (b) the divergence of certain characteristics, most commonly, point

substitution rates within each MSA aligned-regions may be a reasonable measure to

represent the evolutionary divergence/distances among the whole organisms, without

considering possible evolutionary roles of all other proteins without high homologous

regions among them and of other mutational events including absence/presence of

proteins.

Such "alignment-based" protein trees represent the evolutionary phylogeny of the

selected regions of the selected proteins, but not full characteristics of all proteins, let

alone the whole organisms, because the aligned regions account, in general, for a very

small fraction of all proteins (Pace 2009).

Information-theory-based ("alignment-free") "whole-proteome trees": This situation

has since changed significantly in two important aspects: (a) During last decades, a large

number (over 134 species, mostly insects, as of 2020) of whole-genome sequences of

extant insect species have been accumulating in public databases, and (b) Information

Theory, developed to analyze linear electronic signals, was found to be adaptable to

analyze other linear information, such as natural languages and genomic information,

without sequence alignment ("alignment-free") (Zielezinski et al. 2017; Blaisdell 1986).

In this approach, the whole content, not selected portions of the whole content, of a whole-

proteome sequence, can be described by "*n*-Gram" or "*k*-mers" (Zielezinski et al. 2017).

N-Gram of a whole-proteome is the collection of all overlapping short subsequences of

length *n*, and it contains all information necessary to reconstruct the original sequence.

Furthermore, the information divergence (difference) between two *n*-Grams can be

estimated by, for example, Jensen-Shannon divergence (JSD) without alignment of the

whole proteome sequences (Lin 1991). Such approach has been widely tested and

validated for comparing texts and books of natural languages for latent semantic analysis

since 1990s (Deerwester et al. 1990) and gene sequences consisting of coding and non-

coding regions as well as amino acid sequences since 1986 (Blaisdell 1986).

Some of these validated methods have been adapted and optimized to handle

whole-proteome sequences in Feature Frequency Profile (FFP) method (Sims et al. 2009).

Since there is no "golden standard" for a phylogenetic evolutionary tree of a group of

organisms that can be experimentally validated, the FFP method has been tested using 26

books in English alphabets from diverse authors and genres, after removing all spaces and

delimiters as well as author names, book titles, headers, footers, etc. In general, the

method performed well in grouping the "books" by the genre and authors (see Fig. 1 of

Sims et al. 2009). In a recent bench-marking studies of 24 Alignment-free methods, FFP

method was ranked among the top 5 best-performing tools for phylogeny prediction based

on the input data of assembled whole genome sequences (Zielezinski et al. 2019).

In the FFP method, whole-genome/whole-proteome information is used under the

assumptions very different from the protein trees: (a) whole-proteome sequence of an

organism represents the organism better than the collection of short regions of highly

homologous sequence from a set of selected genes/proteins used in the protein trees and

(b) a combination of *all types of mutations*, such as point substitution, insertion/deletion of

various length, recombination, duplication, transfer or swapping of genes etc., contribute

to the evolutionary processes of the organisms, rather than only point substitution rates in

the protein trees. Thus, whole-proteome tree may provide an independent view of the

evolutionary relationship among the insect organisms.

Experiences from earlier whole-proteome trees: In the last decade, we have tested and

optimized the protocols for building whole-proteome trees using various different

populations such as Bacteria and Archaea Domains (Jun et al. 2010), Fungi Kingdom

(Choi & Kim 2017) and, most recently, all three Domains at a deep phylogenic level

(Choi & Kim 2020). From these studies we have learned that: (a) among three types of

the trees (whole-genome DNA tree, whole-transcriptome RNA tree, and whole-proteome

amino-acid tree) the whole-proteome trees produce the most topologically stable trees; (b)

for a give group of organisms, the optimal length of the sequence strings to be used in FFP

method can be empirically determined; and (c) cumulative genomic divergence (CGD) is

a useful and computable quantity for the point of the emergence for the founders of a

group in the "evolutionary progression scale".

In this study, we optimized various parameters and protocols specifically for the

population of insects and present a view of the whole-proteome tree based on whole-

proteome sequences of 134 diverse arthropod species (123 insects plus 11 non-insect

arthropods), available in the NCBI database (O'Leary et al. 2016), and discuss its

implications to phylogenic aspects of insect evolution.

Results

To compare the current protein trees with our whole-proteome Tree of Insects

(ToIn) we chose two recent and very comprehensively-analyzed protein trees: The first

one is the recent "alignment-based" tree of 144 insect taxa based on 1,478 single-copy

protein-coding nuclear genes (Fig. 1 of Misof et al. 2014).   The second is the tree for 76

arthropod taxa (Fig. 2 of Thomas et al. 2020) based on up to 4,097 single copy protein-

coding genes. In both cases, the number of aligned genes used are a small fraction of

about 10,000 to 31,000 genes among their study insects. Both protein trees agree with

each other, in general, on the branching order of the Order groups of the respective

populations, and on similar time spread of the emergence of the founders of the groups in

chronological time scale estimated based on available fossils and calibration methods

under various assumptions.

In comparing our whole-proteome ToIn to these two protein trees, we focus on

two aspects separately: grouping patterns and phylogeny of the groups. For the former,

we use two methods: First, we cluster our study population by several unsupervised

clustering algorithms using only the distances estimated from the "divergence" among

whole-proteome FFPs with no explicit constraints of the presence of the common

ancestor(s) or specific evolutionary models (see Construction of whole-proteome Tree of

Insects in Materials and Methods).   We then ask whether the "clustering pattern" is

similar to the "clading pattern" in the protein trees and in our whole-proteome ToIn,

recognizing that the both tree constructions assume the constrains of the common

ancestor(s) and specific evolutionary models.   For the phylogeny of the groups, we

compare the order of branching of the groups and their emergence points on the

evolutionary progression scale in our tree and in chronological scale in the protein trees

(see "Cumulative Genomic Divergence (CGD)" as "Evolutionary Progression Scale" in

Materials and Methods).

A. Demographic grouping pattern by clustering and clading

Clustering: We have tested the grouping pattern of the insects by several

unsupervised clustering algorithms, such as Principal Component Analysis (PCA), Multi-

Dimensional Scaling, and t-Distributed Stochastic Neighbor Embedding (t-SNE) (R Core

Team 2016; v.d. Maaten & Hinton 2008), all of which can be accomplished from the

same starting "distance" matrix constructed using the divergence of whole-proteome

sequences, as calculated by JSD (Lin 1991) of FFPs, among all pairs of the study

organisms. All three clustering methods showed the clustering pattern compatible with

the current grouping of arthropods with common and scientific names, mostly based on

morphological characteristics. Figure 1, a classical PCA clustering, which is very similar

to that of MDS method, shows that all our study population are distributed into 5

"spokes". Two long spokes (IV and V) corresponds to all the members of Diptera and

Hymenoptera of Insecta Class, respectively. The remaining three short spokes (I, II, and

III) correspond to: Members of Chelicerata and Crustacea of non-insect arthropods in

spoke I; those of Hemipters-A and Lepidoptera of Insecta Class in spoke II, and those of

Hemiptera-B, Coleoptera, and Blattodea of Insecta Class in spoke III.   The most

noticeable difference with the grouping in the current protein trees (Misof et al. 2014;

Thomas et al. 2020) is that Hemiptera is split into two separate groups, labeled in this

study as Hemiptera-A and -B. Another clustering by t-SNE (see Supplemental

information, Fig. S1) also shows a similar split of Hemiptera into two, which is

unexpected, because the assumptions and algorithms in t-SNE and PCA are completely

different.

Clading: As a second method of the grouping, we use the clading pattern of the

organisms in our whole-proteome ToIn. Figure 2 shows the topology of the ToIn,

constructed using Neighbor-Joining method implemented in BIONJ (Saitou & Nei 1987;

Gascuel 1997). In this study, we use the divergence of whole-proteome sequences of two

organisms as the estimates for the evolutionary distances between them, as calculated by

JSD of pair-wise FFPs at an optimal Feature length (see Construction of whole-proteome

Tree of Insects in Materials and Methods and Supplementary information Fig. S2). We

also assume an evolutionary model of Maximum Parsimony (minimum evolution) in a

way that the chosen neighbors to be joined are those that minimize the total sum of

branch-lengths at each stage of step-wise joining of neighbors starting from a star-like

tree (Saitou & Nei 1987). The tree shows that most of the clusters in Fig.1 can be

identified among the clades in the ToIn.

Robustness of grouping: The grouping pattern by clustering *and* clading in our

study agrees well with those of the protein trees (Misof et al. 2014; Thomas et al. 2020)

except for Hemiptera group (see Notable differences in grouping and phylogenic

positions in Discussions and Implications). Thus, it is surprising that the demographic

grouping pattern is robust, in general, regardless of not only the information type (select

protein-characteristics, or whole-proteome characteristics), but also of the methods

(clustering or clading) used in grouping. For an implication of this result, see Similarities

in grouping patterns in Discussion and Implications. However, not surprisingly, there

are significant differences from the protein trees (Misof et al. 2014; Thomas et al. 2020)

in branch-length and branching order of the groups (see Dissimilarities in branching

orders and branch-lengths in Discussions and Implications).

B. Emergence of the "Founders" of all major groups in a staged "burst"

For the following results we define "Cumulative Genomic Divergence (CGD)" for an internal node of the ToIn as the cumulative scaled-branch-length from the tree root to the node (see Cumulative Genomic Divergence (CGD) as "Evolutionary Progression Scale" in Materials and Methods) to represent the extent of the "evolutionary progression" of the node. The progression is scaled such that the root node of ToIn is set at $CGD = 0$ (see Outgroup in Discussions and Implications) and the leaf nodes of the extant organisms at $CGD = 100$, on average.

"Arthropodal burst" near the root of ToIn: Figures 3, 4 and 5 show the whole-proteome tree with CGD values. They reveal that the "founders" (for definition, see Supplemental information, Fig. S3) of all major groups of insects as well as non-insect arthropods (at Subphyla and Order levels) emerged in a staged burst within a short evolutionary progression span between CGD of 1.6 and 5.8 (marked by a small red arc in Figs. 3A and 3B), near the root of the tree. This observation is dramatically different from those of the protein trees (Misof et al. 2014; Thomas et al. 2020), where the founders of the major groups of all arthropods emerged throughout a long time-span of chronological scale.

A subgroup of Hemiptera (Hemiptera-A) is the most basal group of all Insecta: The first founders of Class Insecta to emerge is the founders of Hemiptera-A group (aphids and a psyllid) at CGD of about 3.7 (Figs. 3A, 3B). This is in stark contrast

to the protein trees, where all Hemiptera is the sister to Thysanoptera (thrips) (Misof et al.

2014) or a group of all Hemiptera, thrips and human louse is the sister to all other large

groups of insects except Blattodea group (Thomas et al. 2020) (see also Notable

differences in grouping and phylogenic positions in Discussions and Implications).

Order of emergence of the "founders" of all major groups of Insecta: Figure

4 shows a series of staged emergence of the founders of all major groups of Insecta.

After the most basal group of Hemiptera-A group (aphids and a psyllid) at CGD of

around 3.7, the founders of Diptera group emerged at CGD of 4.1, and those of the

remaining five Order-level groups (Lepidoptera, Hemiptera-B (bugs, a planthopper and a

whitefly), Coleoptera, Blattodea + a thrips, and Hymenoptera groups) at CGD of 4.4, 4.8,

5.2, 5.8, and 5.8, respectively. For possible implications see Notable differences in

grouping and phylogenic positions in Discussions and Implications below**)**

Discussions and Implications

*Similarities in grouping patterns***:** As mentioned earlier, it is surprising that the grouping

patterns at Order level between the protein trees (Misof et al. 2014; Thomas et al. 2020)

and our whole-proteome ToIn are very similar (see below for one notable exception of

Hemiptera) despite the facts that the types of input data (multiple-aligned regions of

selected proteins vs. whole-proteome) and estimation methods for evolutionary distance

used (based on point mutational rates vs. whole genomic divergences) are very different.

A possible implication is that, after the "burst", the members of each group evolved

largely "isolated" within the group without significant genomic mixing between the

groups, thus, resulting in much smaller genomic variation within the group than between

the groups, as manifested by mostly isolated clusters.

Dissimilarities in branching orders and branch-lengths: It is not surprising that the

branching orders and branch-lengths are not similar between the protein trees (Misof et al.

2014; Thomas et al. 2020) and our whole-proteome tree, because the assumptions under

which the estimations for evolutionary distances among the organisms are calculated are

very different: in the protein trees, the distances are calculated only for the aligned

portions of the selected genes using, e.g., point-substitutional mutation rates, while in our

tree they are calculated by accounting presence/absence of all amino acid short strings,

Features, for all proteins due to all types of mutations.

Evolutionary progression scale vs. Chronological time scale: It is difficult to design a

scale that quantitatively measures the degree of evolutionary progression, because it is not

clear what characteristics of an organism can best reflect the progression and also are

quantitatively measurable. Since we are using whole-proteome sequence to represent each

organism, we use the divergence of the whole-proteome sequences as the evolutionary

progression scale (Choi & Kim 2020). In contrast to linear chronological time scale, the

evolutionary progression scale is most likely not strictly linear, because any significant

geological and ecological events may accelerate or decelerate the evolutionary progression

for a given organism.   However, the direction of arrows in both scales are the same,

suggesting that the two scales may be calibrated when sufficient fossils, other independent

records, and improved calibration methods become available (see Cumulative Genomic

Divergence (CGD) as "Evolutionary Progression Scale" in Materials and Methods).

Meanwhile, we use the evolutionary progression scale to compare the order of emergence

of the founders of various major groups under the assumption that the whole-proteome

sequence divergence can be considered as informational entropy, which increases as

evolution progresses, similar to the physical entropy of universe increases as the universe

evolve.

"Burst" vs. Gradual emergence of the founders of major groups: While cognizant of

the difference and similarity of the two scales, the most dramatic difference is observed in

the span of the scales within which the founders of all major groups at Order level

emerged in the protein trees (Misof et al. 2014; Thomas et al. 2020) and in our whole-

proteome ToIn: In the protein trees, the founders of all the groups at Order level emerged

gradually during a long chronological time span of about 350 Million years (Myrs)

corresponding roughly 60% of about 570 Myrs between the tree root to the extant

arthropods (Fig. 1 of Misof et al. 2014), or about 210 Myrs corresponding to roughly 37%

of the same full chronological scale (Fig. 2 of Thomas et al. 2020). In drastic contrast, the

founders of all the major groups in our tree emerged within about 4% of the full

evolutionary progression scale in a sudden burst ("Arthropodal burst"; see Figs. 3A and

3B) near the root of our whole-proteome tree. This drastically contrasting observations

between the two types of trees may have an important qualitative evolutionary implication

in constructing the narrative for the birth of the insect diversity.

Notable differences in grouping and phylogenic positions: Despite the drastic difference

in the emergence pattern of the founders (burst vs. gradual) mentioned above, the order of

emergence of the major groups at Order and Subphylum levels agree between the two

protein trees (Misof et al. 2014; Thomas et al. 2020) and our whole-proteome tree with

some notable differences in Hemiptera and Blattodea as described in Results above.

These differences may get resolved once the whole-genome sequences of many more

relevant organisms become available. At present, we suggest some possible implications

as described below:

Hemiptera: As mentioned earlier, in our whole-proteome ToIn (Figs. 3A and

3B) as well as in PCA (Fig. 1) and t-SNE (Supplemental Fig. S1) clustering plots,

Hemiptera is divided into two separate clades/clusters, which we call Hemiptera-A

("primitive" Hemiptera, such as aphids and a psyllid) and Hemiptera-B ("bugs" such a

planthopper, a whitefly, a stink bug and a bed bug), and their phylogenetic positions are

very far apart (see Figs. 2, 3 and 5): Hemiptera-A at the basal position of all Insecta and

Hemiptera-B as sister to the group consisting of Lepidoptera, Coleoptera and

Hymenoptera. But, in the protein trees (Misof et al. 2014; Thomas et al. 2020) the both

groups form a single clade, and is at basal or sister to all other large groups of insects

except Blattodea group. This difference in clustering and phylogenetic positioning

suggest that, when viewed at whole-proteome level, which includes *both* homologous and

non-homologous proteins, the members are more similar within each subgroup than

between the two subgroups in our tree.   But, when viewed, as in the protein trees, only

for the select homologous proteins in the *absence* of the non-homologous proteins, which

are the overwhelming majority of the all proteins, they are similar among all of them to

form only one clade.

313   **Blattodea:** Two termites (Blattodea) and one thrips (Thysanoptera), both eusocial

and hemimetabolous, form a clade in our whole-proteome ToIn and the clade is sister (or

basal) to Hymenoptera group, which is also eusocial but holometabolous (see Fig. 5).

However, in one protein tree (Misof et al. 2014), Blattodea group (cockroaches and

termites, which are eusocial and hemimetabolous) is a member of a larger clade

Polyneoptera and placed at the basal position to all other Order groups of Insecta, which

are largely non-social and hemi- or holo-metabolous, while, in the other protein tree

(Thomas et al. 2020), Blattodea group forms a separate clade, and is placed near the basal

position of all other Order groups of Insecta.   This is in contrast to what we observe in our

ToIn, where Hemiptera-A, is the basal group of Insecta.

Outgroup: Since our method does not require multiple sequence alignment, we

constructed, as was described in our earlier works on whole-proteome trees (Jun et al.

2010; Choi & Kim 2017; Choi & Kim 2020), the proteome sequence of an "artificial

(faux) arthropod" by "shuffling" (Knuth 1973; Fisher & Yates 1948) the alphabets of the

whole proteome sequence of an organism in the study group. We used two such artificial

arthropods (named R28612 and r12957) to form the outgroup for this study. Each has the

same size and amino acid composition of corresponding protein of an extant arthropod,

but does not have gene sequences information for the organism's survival.

Materials and Methods

Sources and selection of proteome sequences

We downloaded the proteome sequences for 134 arthropods from NCBI RefSeq

DB using NCBI FTP depository (O'Leary et al. 2016). Protein sequences derived from

all organelles were excluded from this study. Also excluded from our study are those

derived from whole genome sequences assembled with "low" completeness based on two

criteria: (a) the genome assembly level indicated by NCBI as "contig" or lower (i.e. we

selected those with the assembly levels of 'scaffold', 'chromosome' or 'complete

genome'), and (b) the proteome size smaller than 80% of the smallest proteome size

among highly assembled arthropod genomes (*Anopheles gambiae str. PEST* with 14,089

proteins at "chromosome" assembly level; TaxID 180454).

All taxonomic names and their taxon identifiers (TaxIDs) of the organisms in this

study are from NCBI taxonomy database, and listed in Supplementary Information,

Dataset S1.

Construction of whole-proteome Tree of Insects

Based on our earlier experiences of constructing whole-proteome trees of

prokaryotes (Jun 2010), fungi (Choi & Kim 2017) and all life forms (Choi & Kim 2020)

by Feature Frequency Profile (FFP) method (Sims et al. 2009), following choices have

been made to obtain a topologically stable whole proteome ToIn of maximum parsimony

(minimum evolution) by BIONJ (Saitou & Nei 1987): a) Among three types of genomic

information (DNA sequence of the whole genome, RNA sequence of whole

transcriptome and amino acid sequence of whole proteome) whole-proteome trees are

most "topologically stable" as estimated by Robinson-Foulds metric (Robinson &Foulds

1981) at respective "optimal Feature-length"; b) For FFP as the "descriptor" of the whole

proteome of each organism, the optimal Feature-length is about 10 amino-acid string (see

Supplementary Information, Fig. S2); and c) Jensen-Shannon Divergence (JSD) (Lin

1991) is an appropriate measure of "divergence of information content", as the measure

of dissimilarity between two whole-proteome descriptors, for constructing the distance

matrix of BIONJ (Saitou & Nei 1987; Gascuel 1997).   It is important to note that such

FFP of a whole-proteome sequence of an organism has all the information necessary to

reconstruct the original whole proteome sequence.

"Cumulative Genomic Divergence (CGD)" as "Evolutionary progression scale"

In Information Theory (Shannon 1948), the Jensen-Shannon Divergence (JSD)

(Lin 1991), bound between zero and one, is commonly used as a measure of the

dissimilarity between two probability distribution of informational features. The FFP as

the descriptor for a linear sequence information of the whole proteome of an organism is

such a probability distribution. Thus, a JSD value of two FFPs, used as a measure of the

information divergence between two proteome sequences, is also bound between 0 and 1,

corresponding to the JSD value between two FFPs of identical whole proteome sequences

and two completely different whole proteome sequences, respectively. Any whole

proteome-sequence "dissimilarity" between two extant organisms accumulated during the

evolution can be considered as caused by changes of, ultimately, genomic sequences of

all protein coding genes due to all types of mutational events, such as point substitutions,

insertion/deletion of various lengths, inversion, recombination, loss/gain of genes, etc. as

well as other unknown mechanisms, and they will bring JSD somewhere between 0.0 and

1.0 depending on the degree of the sequence divergence.

In this study the collection of the JSDs for all pairs of the study organisms plus 2

out-group members (see Outgroup in Discussions and Implications) constitutes the

"distance matrix" for BIONJ (Saitou & Nei1987; Gascuel 1997).   Since all the branch-

lengths are derived from the JSD values, the cumulative branch-length of an internal

node, which we call "cumulative genomic divergence (CGD)" (to reflect the fact that the

proteomic divergence is ultimately derived by the genomic divergence during evolution)

of the node, can be considered as the point of evolutionary stage reached by the node on

an "evolutionary progression scale". For convenience of assigning the nodes on the

progression scale, CGDs are scaled, as mentioned earlier, such that the CGD value at the

root node of ToIn is set to zero and the leaf nodes of the extant organisms to 100, on

average, corresponding to the fully evolved genomic states of the organisms, which we

define as the beginning and ending point of the "evolutionary progression scale" for the

organisms (see Fig. 3A).

Clustering methods

We use two unsupervised methods to observe the clustering patterns based solely

on whole-proteome sequences: Principal Component Analysis (PCA) and t-Distributed

Stochastic Neighbor Embedding (t-SNE) (R Core Team 2016; v.d. Maaten & Hinton

2008). Both are dimensional reduction methods, but with different strengths and

weaknesses for our purposes, which help to visualize any clustering pattern in the data

distribution. Both are based only on the evolutionary distances (CGD in this study),

estimated by the divergence of whole-proteome sequences among all pairs of the study

arthropods. In PCA, the distances within a cluster as well as between two clusters are

quantitative, thus, two close clusters nearby may not resolve well. In t-SNE, which

applies Machine Learning to emphasize the resolution of nearby clusters, but the inter-

cluster distances are de-emphasized, thus, not quantitative.

Declarations

Acknowledgement: We gratefully acknowledge the comments and advices from the

arthropod experts at University of California, Berkeley, CA., Profs. Kipling Will and Peter

Oboyski, and the constructive suggestions on our application of Information Theory and

Unsupervised Clustering methods from Prof. Se-Ran Jun of College of Medicine,

University of Arkansas for Medical Sciences, Little Rock, AR, and Dr. Chao Zhang, Chief

Scientific Officer, of Plexxikon, Inc. All the silhouettes are generated by Rigel Sison.

Funding: This research was partly supported by a grant (to SHK) from World Class

University Project, Ministry of Education, Science and Technology, Republic of Korea

and a gift grant to University of California, Berkeley, CA. (to support JJC). SHK

acknowledges having appointments as Visiting Professorships at Yonsei University,

Korea Advanced Institute of Science and Technology, and Incheon National University

in South Korea during the manuscript preparation.

*Competing financial interests***:** Authors declare that there are no competing financial

interests in connection with this paper.

Author contribution: Conceptual design of the study and speculative interpretations and

implications of the results by SHK; filtering and curation of genomic and proteomic

sequence data from NCBI database, computational-algorithm design, programming and

execution by JJC and BJK; unsupervised clustering by various algorithms were

performed by BJK; interpretation of computational results by SHK, JJC and BJK;

manuscript preparation by SHK with extensive discussions with JJC and BJK; all figures

are designed by SHK, JJC and BJK.

Computer code availability: The FFP programs for this study (2v.3.0) written in

GCC(g++) is available in Github: https://github.com/jaejinchoi/FFP.

References

Blaisdell, B. E. (1986). A measure of the similarity of sets of sequences not requiring sequence

alignment. *Proceedings of the National Academy of Sciences of the United States of*

America, *83*(14), 5155–5159.

Choi, J.J., & Kim, S.-H. (2017). A genome Tree of Life for the Fungi kingdom. *Proceedings of*

the National Academy of Sciences, *114*(35), 9391–9396

Choi, J.J., & Kim, S.-H. (2020). Whole-proteome tree of life suggests a deep burst of organism

diversity. *Proceedings of the National Academy of Sciences*, *117*(7), 3678–3686

Deerwester, S., Dumais, S. T., Furnas, G. W., Landauer, T. K., & Harshman, R. (1990). Indexing

by latent semantic analysis. *Journal of the American Society for Information Science*, *41*(6),

391–407.

Fisher, R. A., & Yates, F. (1948). *Statistical tables for biological, agricultural and medical*

research. London: Oliver and Boyd.

Gascuel, O. (1997). BIONJ: an improved version of the NJ algorithm based on a simple model of

sequence data. *Molecular Biology and Evolution*, *14*(7), 685–695.

Jun, S.-R., Sims, G. E., Wu, G. A., & Kim, S.-H. (2010). Whole-proteome phylogeny of

prokaryotes by feature frequency profiles: An alignment-free method with optimal feature

resolution. *Proceedings of the National Academy of Sciences*, *107*(1), 133–138.

Knuth, D. E. (1973). *Seminumerical algorithms, in The art of computer programming* (3rd ed.).

Boston: Addison-Wesley.

Letunic, I., & Bork, P. (2019). Interactive Tree Of Life (iTOL) v4: Recent updates and new

developments. *Nucleic Acids Research*. 47(W1), W256–W259

465 Lin, J. (1991). Divergence measures based on the Shannon entropy. *IEEE Transactions on*

466   *Information Theory*, *37*(1), 145–151.

467 Misof, B., Liu, S., Meusemann, K., Peters, R. S. R., Donath, A., Mayer, C., … Zhou, X. (2014).

468   Phylogenomics resolves the timing and pattern of insect evolution. *Science*, *346*(6210),

469   763–768.

470 O'Leary, N. A., Wright, M. W., Brister, J. R., Ciufo, S., Haddad, D., McVeigh, R., … Pruitt, K.

471   D. (2016). Reference sequence (RefSeq) database at NCBI: Current status, taxonomic

472   expansion, and functional annotation. *Nucleic Acids Research*, *44*(D1), D733–D745.

473 Pace, N. R. (2009). Mapping the Tree of Life: Progress and Prospects. *Microbiology and*

474   *Molecular Biology Reviews*, *73*(4), 565–576.

475 R Core Team. (2016). *R: A language and environment for statistical computing*. Retrieved from

476   https://www.r-project.org/

477 Robinson, D. F., & Foulds, L. R. (1981). Comparison of phylogenetic trees. *Mathematical*

478   *Biosciences*, *53*(1), 131–147.

479 Saitou, N., & Nei, M. (1987). The neighbor-joining method: a new method for reconstructing

480   phylogenetic trees. *Molecular Biology and Evolution*, *4*(4), 406–425.

481 Shannon, C. (1948). A Mathematical Theory of Communication. *The Bell System Technical*

482   *Journal*, *Vol.27*(1948), 379–423.

483 Sims, G. E., Jun, S., Wu, G. A., & Kim, S. (2009). Alignment-free genome comparison with

484   feature frequency profiles (FFP) and optimal resolutions. *Proceedings of the National*

485   *Academy of Sciences of the United States of America*, *106*(8), 2677–2682.

Thomas, G. W. C., Dohmen, E., Hughes, D. S. T., Murali, S. C., Poelchau, M., Glastad, K., …

Richards, S. (2020). Gene content evolution in the arthropods. *Genome Biology*, *21*(1), 15.

1-14

Van Der Maaten, L., & Hinton, G. (2008). Visualizing data using t-SNE. *Journal of Machine*

Learning Research, *9*, 2579–2605.

Zielezinski, A., Girgis, H. Z., Bernard, G., Leimeister, C.-A., Tang, K., Dencker, T., …

Karlowski, W. M. (2019). Benchmarking of alignment-free sequence comparison methods.

Genome Biology, *20*(1), 144. 1–18

Zielezinski, A., Vinga, S., Almeida, J., & Karlowski, W. M. (2017, October 3). Alignment-free

sequence comparison: benefits, applications, and tools. *Genome Biology*, Vol. 18, pp. 1–17.

Figure legends

Fig. 1: Unsupervised clustering (grouping) of 134 extant arthropods (123 insects plus

11 non-insect arthropods) by classical PCA

Classical PCA plotted for the three principal axes reveals about 5 large clusters arranged

in 5-spokes. Two long spokes (IV and V) corresponds to all the members of Diptera and

Hymenoptera, respectively. The remaining three short spokes (I, II, and III) correspond

to: Members of Chelicerata and Crustacea in spoke I; those of Hemipters-A and

Lepidoptera in spoke II, and those of Hemiptera-B, Coleoptera, and Blattodea in spoke III.

Fig. 2: Topology of the linear representation of whole-proteome Tree of Insects (ToIn)

The colors of the first (inner) colored-band distinguish organisms in different Classes, and

those of the second (outer) band among different Orders (the names of different color-

bands are shown in Fig. 3A). Scientific names and common names, when available, of

each organism are also listed. The silhouettes of sampled organisms are shown next to

their names. To emphasize the clading pattern, all branch-lengths are ignored. The first

two items refer to two members of the outgroup (see Outgroup in Discussions and

Implications) constructed by shuffling (Knuth 1973; Fisher & Yates 1948) the whole-

proteome sequences of the two arthropods. The visualization of the ToIn was made using

iTOL (Letunic & Bork 2019).

Fig. 3A: "Pie" representation of whole-proteome ToIn with the cumulative branch-

lengths scale.

This view of the whole-proteome ToIn shows all branch-lengths to emphasize the

progression of evolution of each member in the study population from the root of the tree

at CGD = 0 to the extant forms of the members at CGD = 100, on average. The small red

arc near the root is at CGD=5.8, by which point of the evolutionary progression the

founders of all major groups (consisting of 7 Order groups and 2 Subphylum groups

shown in Fig. 1) have emerged, suggesting that the remaining 94.2 on CGD scale

corresponds to further diversification and gradual evolution of the founders and common

ancestors *within* each major group toward their extant forms. The visualization of the

ToIn was made using iTOL (Letunic & Bork 2019).

Fig. 3B: Expanded view of Fig. 3A near the root of the whole-proteome ToIn

Examples of the founders of all major groups are shown as blue dots, and the common

ancestors of extant groups within two major groups, Diptera and Hymenoptera, as red

dots.   The visualization of the ToIn was made using iTOL (Letunic & Bork 2019).

Fig. 4: Simplified whole-proteome ToIn

The vertical axis shows cumulative genomic divergence (CGD) values, which ranges from

zero to around 100, and they correspond to the extent of evolutionary progression from the

root of the ToIn to the extant leaves. For simplicity, "singletons" (that do not belong to

any named groups) are not shown, and all the leaf nodes and their branches of a common-

named group (in parenthesis) are combined into a single dotted line coming out from their

common ancestor node of the extant group shown as a blue sphere. Each internal node

represents a "pool of founding ancestors" (see Supplementary Information Fig. S3).

Dotted vertical lines are to indicate that they are arbitrarily shortened to accommodate

large jumps of CGD values within a limited space of the figure. The double-headed arrow

at bottom right indicates the short range of the CGD values, within which the founders of

all the major groups of the extant organisms in this study have emerged in a "burst". For

our interpretation of horizontal lines and vertical lines, see Supplementary Information

Fig. S3.

Fig. 1

PCA analysis on Arthropoda136

About 13+ clusters grouped into
5-spoked star

IV

(Fruit fly A)

(Fruit fly B)

Diptera

(Other flies)

V

Blattodea
(Termites, Thrips)

Hymenoptera

(Ants)

(Bees)

(Mosquitos)

III

(Sawflies/Wasps)

I

Chelicerata,
Crustacea

(Bugs)

Coleoptera
(Beetles)

(Aphids, psyllid)

Lepidoptera
(Moths, Butterflies)

Hemiptera-B

II

Hemiptera-A

PCA Dim2

0.5

-10.2

-10.4

I: Chelicerata, Crustacea
II: Hemiptera-A, Lepidoptera
III: Hemiptera-B, Coleoptera
      Blattodea
IV: Diptera
V: Hymenoptera

Fig. 2

Fig. 3A.

Fig. 3B.

"Arthropodal Burst": 🔵 Emergence of the founders of all extant major groups (Subphyla Crustacea and Chelicerata, and all Order groups of Hexapoda); 🔴 Emergence of the founders of common-named groups in Diptera and Hymenoptera, two largest groups

Fig. 4