## Independence of Cued and Contextual Components of Fear Conditioning is Gated by the Lateral Habenula

Tomas E. Sachella [1], Marina R. Ihidoype [1], Christophe Proulx [2], Jorge H. Medina [3], Pablo Mendez [4], Joaquin Piriz [1] †*.

[1] Grupo de Neurociencia de Sistemas, Instituto de Fisiología y Biofísica "Houssay" (IFIBIO "Houssay"), Universidad de Buenos Aires, CONICET, Buenos Aires, Argentina.

[2] CERVO Brain Research Center, Department of Psychiatry and Neurosciences, Université Laval.

[3] Instituto de Biología Celular y Neurociencia "Dr. Eduardo De Robertis" (IBCN), Universidad de Buenos Aires, CONICET, Buenos Aires, Argentina.

[4] Instituto Cajal, CSIC, Madrid, España.

† Present address: Instituto de Fisiología Biología Molecular y Neurociencias (IFIBYNE), Universidad de Buenos Aires, CONICET, Buenos Aires, Argentina.

* Correspondence:

Joaquin Piriz

Email: pirizjoaquin@gmail.com

22    Abstract

23           Fear is an extreme form of aversion that, if inappropriately generalized, initiates pathological

conditions such as panic or anxiety disorders. Fear conditioning (FC) is the best understood model of fear

learning. FC forms two associations that independently link the cue and the training context to fear

responses. The lateral habenula (LHb) encodes aversive information. However, how the LHb participates

in fear learning has not been intensively studied. Here we studied the role of the LHb in FC learning using

optogenetics and pharmacological tools in rats. We found that disrupting neuronal activity of the LHb

during training abolishes independent expression of contextual and cued memories, yet memory is

normally expressed when the cue is played in the training context. Our results show that neuronal activity

at the LHb regulates independent expression of sub-components of a fear memory, assigning to that

structure a previously uncharacterized role as regulator of fear memories.

Introduction

34           Fear is an extreme form of aversion characterized by an uncontrollable reaction to threatening

35    stimulus. In nature, stimuli to be fear of will never be identical, thus fear is adaptively generalized to some

36    degree (Asok et al., 2019). However, exacerbated fear generalization is supposed to be the basis to the

37    development of psychopathological conditions like panic or anxiety disorders, highlighting the relevance

38    of the study of basic mechanisms of fear learning (Dunsmoor & Paz, 2015; Lissek et al., 2010, 2014).

39           Fear conditioning (FC) is probably the most studied and best understood model of fear learning

(Maren, 2008; Maren & Quirk, 2004). In FC an association is made between a neutral stimulus and a

biologically-relevant aversive stimulus called unconditioned stimulus (US). Most frequently, FC training

involves the pairing of a tone, named cue, and an electric foot-shock US. It has long been known that such

protocol leads to two independent associations, one relating the cue to the US, and the other linking the

context where the training took place to the US (J. Kim & Fanselow, 1992; LeDoux & Phillips, 1992).

Consequently, subjects will display defensive responses, characterized by freezing behavior, when re-

exposed to the training context or to the cue presented in a novel and neutral context.

47           The lateral habenula (LHb) is a central hub for the processing of aversive information in the brain.

Aversion related information reaches the LHb from numerous structures of the limbic system and basal

ganglia. In turn, the LHb projects to the brain stem, where it is one of the few structures that control

(directly and indirectly) both serotoninergic and dopaminergic systems (Hikosaka, 2010; Hu et al., 2020;

Proulx et al., 2014). That connectivity places the LHb in a central position to control aversive associative

learning. However, how the LHb is related to fear learning has not been intensively studied yet.

In a previous work, using the Inhibitory Avoidance paradigm, we found that inactivation of the
LHb during training affects temporal stability of the memory but not its formation (Tomaiuolo et al., 2014).
To better understand how the LHb influences aversive learning, in this work we studied its role in FC
learning by means of pharmacological and optogenetics manipulations. We show that interfering with
neuronal activity of the LHb during FC training severely impairs recall of contextual and cued FC memories.
However, when the cue is played in the training context, a conserved FC memory is evidenced. Thus, our
results suggest that the formation of two independent memory traces during FC requires an intact LHb
activity, placing this structure in a central position in the neuronal circuits mediating FC learning.

## Results

### Pharmacological inactivation of the LHb impairs contextual fear conditioning

To examine the role of the LHb in contextual FC, we analyzed the effect of its inactivation during
training. For that purpose, we performed surgeries, in rats, to implant bilateral intracerebral cannulae
aimed at the LHb. After a recovery time of 10 to 12 days, animals were bilaterally infused the GABA-A
agonist muscimol or vehicle in the LHb, and were trained in contextual FC thirty minutes later. During
training rats were placed in a FC chamber and presented with four unsignaled mild foot-shocks (**Figure
1A**). We observed that the infusion of muscimol in the LHb did not modify freezing behavior during FC
training (**Figure 1B and Supplementary Figure 1**). Also, in a separate cohort of animals, we observed that
such procedure did not modify locomotion or exploratory behavior in an open field (**OF; Supplementary
Figure 2**), indicating that inactivation of the LHb does not have a general effect over motility or context
exploration. To evaluate contextual FC, animals were placed back in the conditioning chamber 7 days after
training and freezing behavior was quantified. Muscimol group displayed significantly lower levels of
freezing than the control group, suggesting that inactivation of the LHb during contextual FC training
impairs long-term memory formation (**Figure 1C**).

Long-term memories tend to decay over time, a phenomenon that could already be evident 7
days after memory formation (Bekinschtein et al., 2007; Rossato et al., 2009). To test whether an
accelerated decay of contextual FC memory could account for the memory deficits observed in the
muscimol group, we repeated our previous experiment but tested contextual FC memory 24 hours after
training. Contextual FC memory deficits were also evident 24 hours after training (**Figure 1 D**), further
suggesting that inactivation of the LHb during contextual FC training has a detrimental effect over long-
term memory formation.

Importantly, muscimol infusion in the LHb before exposure to an OF did not impact on the
decrease in exploratory behavior observed upon a subsequent re-exposure performed 48 hours later
(**Supplementary Figure 2**), which is acts as a measure of contextual memory on that test (Etkin et al.,
2006; Vianna, 2000). That observation suggests that the inactivation of the LHb by muscimol does not
induce a general deficit in context encoding.

Pharmacological inactivation of the LHb impairs cued fear conditioning

Having observed that inactivation of the LHb impairs the formation of contextual FC we extended
our experiments analyzing the effect of LHb inactivation on cued FC memory formation. Animals were
infused muscimol or vehicle in the LHb, and 30 minutes later were trained in cued FC, in which each of
the 4 foot-shocks was preceded by a 17 seconds long tone (**Figure 2A**). Cued FC memory was tested 24
hours or 7 days later by exposing the animals to the tone in a novel context (Context B). Behavior during
training confirmed that inactivation of the LHb does not affect the surge of freezing elicited by shock
exposure (**Figure 2B, Supplementary Figure 1B**). During tests, presentation of the cue elicited a robust
freezing in the vehicle groups that was markedly reduced in the muscimol groups (**Figure 2C, D**). These
results indicate that cued FC memory is also impaired by the inactivation of the LHb during training. Thus,
inactivation of the LHb during FC training impairs formation of both contextual and cued FC long-term
memories.

To control for spatial specificity of muscimol infusion we implanted cannulae targeting areas
around the LHb and infused muscimol before cued FC training (**Supplementary Figure 3A, B**). During tests,
performed 7 days later, we found that neither infusion of muscimol 1 mm lateral, dorsal or ventral to the
LHb affected freezing behavior (**Supplementary Figure 3 B, C**). Moreover, we did not see a reduction in
freezing behavior during cued FC test in muscimol infused animals implanted with cannulae aimed at the
LHb in which histological control showed cannulations bilaterally missed the LHb (**Supplementary Figure
3A, C**).

To control for non-transient effects of muscimol infusion, we trained in cued FC the animals that
had previously undergone muscimol inactivation of the LHb before exposure to an OF (**Supplementary
Figure 4A**). In that group of animals, we found no differences in freezing behavior during either cued FC
training or testing (**Supplementary Figure 4B, C**), demonstrating the transient nature of that manipulation.
Thus, deficits in contextual and cued FC in the muscimol groups should be attributed to a transient
inactivation of the LHb during training.

Context + tone test evidences a conserved fear conditioning memory

To further investigate to what extent inactivation of the LHb impairs FC learning, we performed
additional experiments in which freezing to the cue was evaluated in the training context (Context A), a
condition we called context + tone (**Figure 3A**). In that test the pre-tone period is a readout of the
contextual component of the memory after a cued FC training. Indeed, during the pre-tone period we
observed a clear reduction of freezing in the muscimol group, confirming that inactivation of the LHb
during training impairs contextual FC (**Figure 3B**). Presenting the tone further increased freezing in the
control group and, most notably, also induced a robust freezing in the muscimol group, which reached
freezing values equivalent to the control group (**Figure 3B**). Thus, under inactivation of the LHb, a FC
memory is formed that could be effectively retrieved when the cue is presented in the conditioning
context.

In a previous paper we demonstrated that the inactivation of the LHb impairs long-term stability
of the Inhibitory Avoidance memory (Tomaiuolo et al., 2014). To investigate long-term stability of FC
memory formed under inactivation of the LHb, we evaluated freezing in context + tone conditions 3 weeks
after training in a new set of animals (**Figure 3 A, C**). Freezing levels of the muscimol group in context +
tone condition were lower than in the vehicle group 3 weeks after training (**Figure 3C**), showing that FC
training under inactivation of the LHb generates a weak memory that is harder to retrieve and is
temporarily less stable.

<u>Context + tone retrieval reconsolidates cued but not contextual memory</u>

In animals that were trained in FC under inactivation of the LHb we observed conserved freezing
responses in context + tone conditions. However, it could not be known if that freezing represents the
retrieval of the contextual memory, the cued memory, both or neither of them. To get an insight on that,
we analyzed if memory retrieval in context + tone conditions induces the reconsolidation of cued and
contextual memories. Animals were infused muscimol in the LHb and trained in cued FC as described
above. Seven days later they went through a recall session in which the cue was presented in the training
context (*i.e.* as in context + tone test). Twenty-four hours later they went through a test session in which
we evaluated reconsolidation of cued and contextual memories. To evaluate reconsolidation of the cued
memory, animals were placed in Context B and exposed to the cue (reconsolidation group, **Figure 4A**). To
evaluate reconsolidation of the contextual memory, animals were re-exposed to the training context
(**Context A; Figure 4C**). In addition, at the end of this latter experiment the cue was presented, configuring
a re-test in context + tone conditions (**Figure 4C**). A "no-reconsolidation" group was also included as a
control for cued memory tested "reconsolidation" group. In the no-reconsolidation group, cued memory

was tested 8 days after training in Context B without any recall session (**Figure 4A**). Notably, during cued

memory test, the "reconsolidation" group that went through the recall session displayed higher levels of

freezing to the cue than the no-reconsolidation group (**Figure 4B**). In fact, in the reconsolidation group

levels of freezing to the tone during test session were equivalent to those shown during the recall session

(**Figure 4B**). In contrast, in the group in which we tested reconsolidation of contextual memory, we did

not observe any increase in freezing to the context between test and recall session (**Figure 4D**). Then,

levels of freezing remained low even though those animals effectively displayed freezing in context + tone

condition in the recall session (**Figure 4D**). Also, freezing in context + tone conditions was not different

between the recall and the test session and was high in both cases (**Figure 4D**). Those results suggest the

recall session reconsolidated the cued memory but not the contextual memory, arguing that freezing

observed during context + tone condition mainly reflects the reactivation of the cued memory trace.

Furthermore, they suggest that FC training under inactivation of the LHb creates a cued memory whose

retrieval depends on contextual information.

<u>Optogenetic inactivation of the LHb impairs cued fear conditioning</u>

To get a better spatial and temporal control of the interventions over the LHb we manipulated its

activity using optogenetic tools. To first confirm and support results obtained from muscimol inactivation

of the LHb, we injected an adeno-associated virus (AAV) encoding the inhibitory light-dependent proton

pump Archaerhodopsin (ArchT) fused to the fluorescent protein GFP (AAV-CamKIIα-ArchT-GFP) bilaterally

in the LHb of rats (**Figure 5A**). Control rats were injected with an AAV encoding the fluorescent protein

GFP (AAV-CamKIIα-GFP). In both groups, optic fiber cannulae were implanted with the tip immediately

above the LHb (**Figure 5A, B**). Whole cell patch-clamp recordings confirmed that LHb neurons could be

inhibited by ArchT (**Supplementary Figure 5**). Light was applied during the FC training session starting at

tone onset and stopping 5 s after shock termination (**Figure 5A**). We did not observe differences in freezing

between ArchT and GFP groups during FC training (**Supplementary Figure 6A**). In accordance with

pharmacological results, we found that freezing to the tone during cued memory test was lower in the

ArchT than in the GFP group. (**Figure 5C, E**). On the other hand, in contrast to pharmacological

manipulation, contextual memory was not affected by optogenetic inhibition of the LHb during cue and

US (**Figure 5D, E**). Additionally, as observed in pharmacological experiments, context + tone condition

evoked similar freezing in ArchT and GFP groups (**Figure 5D, E**).

<u>Artificial activation of the LHb impairs cued and contextual fear conditioning</u>

To rigorously test the involvement of the LHb in contextual FC using optogenetic tools, it would

be necessary to disturb neuronal activity at the LHb during context encoding as well (*i.e.* during whole

training period). Optogenetic inhibition mediated by light activated hyperpolarizing pumps such as ArchT

or NpHR, require sustained illumination which could create confounding effects, such as temperature

changes, when prolonged for long periods like the one involved in our protocols (**Supplementary Figure**

**7**). Also, sustained on-off patterns of optogenetic inhibition induce burst activity of LHb neurons that

induce depression-like symptoms (Yang *et al.*, 2018; **Supplementary Figure 5D**). To avoid those potential

artifacts, we decided to take a different approach. We reasoned that neuronal encoding would be

disturbed by any major intervention over neuronal activity. Thus, we opted to disrupt endogenous activity

of the LHb by imposing an artificial excitation during the whole training. Animals were injected with an

AAV encoding the fast excitatory opsin oChIEF (Lin et al., 2009), and implanted with optic fibers bilaterally

in the LHb (**Figure 6A, B**). Three weeks later they were trained in cued FC under a sustained 20 Hz

excitation drive with light pulses of 5 ms (**Figure 6A**), which has been shown before not to induce

depression-like symptoms (Yang *et al.*, 2018; **Figure 6A, B**). Modelling of light induced temperature

changes supports the idea that such protocol would result in minimal temperature changes in the

illuminated tissue (Stujenske, Spellman and Gordon, 2015; **Supplementary Figure 7**). We found that

artificial stimulation of the LHb during FC training mostly replicates results obtained by pharmacological

inhibition (**Figure 6**). oChIEF activation did not modify behavior during FC training (**Supplementary Figure**

**6B**). During cued memory test, oChIEF stimulated animals showed lower freezing to the tone than the

control group (**Figure 6C, E**). Next day, during context + tone test session, oChIEF stimulated animals

showed almost no freezing to the context but displayed freezing to the cue, reaching levels higher than in

cued memory test performed the previous day (**Figure 6D, E**). Thus, disrupting the encoding of the LHb

during whole training has a deleterious effect over FC learning equivalent to that of pharmacological

inhibition. Remarkably, AAVs did not infect medial habenula (**Figure 5B, 6B**). Thus, results from

optogenetics experiments could not be attributed to an effect over that structure, a criticism valid for

muscimol mediated inactivation.

## Discussion

Few articles had studied the LHb in the context of FC (Barrett & Gonzalez-Lima, 2018; Durieux et

al., 2020; Song et al., 2017; Wang et al., 2013). In this work we directly investigated that subject expanding

our understanding of the role of the LHb in FC learning. Our findings reveal that the LHb is a relevant

component of the FC circuit that plays a role in the formation of both cued and contextual memories, and,

most notably, in the interaction between them. In the following subsections we discuss our findings in
detail.

<u>Cued memory and the LHb</u>

During FC training, subjects infer which stimuli have predictive value over the arrival of the US to
organize proper defensive behavioral responses. In normal conditions, that processing leads to a double
association that assigns a predictive value to both the context and the cues. Prevailing models postulate
that context-US association involves context encoding circuits centered in the hippocampus which send
contextual representation to the amygdala where the association takes place (W. Bin Kim & Cho, 2020;
Maren et al., 2013), and locate cue-US association in the amygdala (Bocchio et al., 2017). Particularly, solid
evidence indicates that tone-US association takes place in cortical and thalamic auditory inputs to the
lateral amygdala (Nabavi et al., 2014; Rumpel et al., 2005; Sah et al., 2008). Thus, as postulated in that
model, cued memory is assumed to be intrinsically context-independent. However, our experiments
suggest that the formation of a cued memory that could be expressed in a context-independent manner
requires signaling by the LHb during memory formation.

Two observations lead us to reach that conclusion. First, we found that disrupting neuronal
activity at the LHb during FC training, either by inhibition (pharmacological or optogenetic), or by
sustained optogenetic activation, results in a condition in which retrieval of FC memory requires that the
cue is presented in the training context. This observation implies an interaction between the context and
the cue in guiding memory retrieval. Second, cued memory is recovered following that retrieval. Since
contextual FC memory is not restored by the same manipulation, it could be inferred that the freezing
observed when the cue is played in the training context is mostly attributable to the reactivation of the
cued memory. In addition, the latter observation argues that freezing in context + tone conditions does
not reflect the retrieval of two weak fear memories (contextual and cued) that add up to trigger freezing
behavior. Altogether, those two observations suggest that, under inactivation of the LHb, a cued FC
memory is formed whose retrieval depends on the context in which the cue is presented.

It has been shown that during FC training, activity of the LHb increases in response to cue
presentation, paralleling the appearance of conditioned responses (Trusel et al., 2019; D. Wang et al.,
2017). Though, no mechanistic information about how the LHb influences cued memory formation could
be inferred from the present work, a tempting hypothesis is that the emergence of responses of the LHb
to the cue acts as a prediction error signal over the arriving of the US that, when confirmed, emancipates
the cue-US association from the context where it was formed.

 Contextual memory and the LHb

Contextual FC is also blocked by pharmacological inactivation or artificial excitation of the LHb
during whole training. It has been proposed elsewhere that the LHb influences context encoding through
a functional interaction with the hippocampus (Baker et al., 2019; Goutagny et al., 2013). That hypothesis
has received support from physiological data generated by other authors and us showing that firing of the
neurons of the LHb is synchronized with hippocampal theta rhythm (Aizawa et al., 2013; Bertone-Cueto
et al., 2020; Goutagny et al., 2013). In addition, two recent papers from Nyiri group showed that LHb-
innervated Nucleus Incertus (Szőnyi, Sos, et al., 2019) and Dorsal Raphe neurons (Szőnyi, Zichó, et al.,
2019) modulate hippocampal theta. Thus, based on that evidence, deficits in contextual FC could be
attributed to a disrupted context encoding.

However, three results presented here suggest that deficits in contextual FC induced by disrupting
neuronal activity of the LHb during training reflect deficits in context-US association rather than in context
representation. First, freezing in context + tone conditions, evidences the retention and effective recall of
contextual information. Second, as discussed above, retrieval in context + tone conditions is not sufficient
to reconsolidate contextual FC memory, suggesting that the contextual memory retrieved is not linked to
fear response. Third, inactivation of the LHb does not affect the decrease in the exploration of an OF upon
re-exposure, showing, in a different paradigm, that inactivation of the LHb does not impede the formation
of a contextual memory. More experiments would be required to determine if the LHb participates in
associative or representational components of contextual FC.

Retrieval and temporal stability, two faces of memory strength

How easy a memory is retrieved could be read as a parameter of memory strength. Also, in
temporal dimension, the strength of a memory could be read from how long it lasts (Bekinschtein et al.,
2008). In our previous work we described that inactivation of the LHb before training in the Inhibitory
Avoidance generates a memory that is normally expressed 24 hours after training but could not be
evidenced 7 days later (Tomaiuolo et al., 2014). In that paradigm, temporal stability of the memory is
defined by shock intensity (*i.e.* by the salience of the US), thus it could be evaluated as a parameter of
memory strength. There, we also found that a series of manipulations known to extend temporal stability
of weak Inhibitory Avoidance memories did also extend temporal stability of the memory formed under
inactivation of the LHb (Tomaiuolo et al., 2014). Based on those findings, we postulated that the main
effect of the inactivation of the LHb was to reduce memory strength. Our present results support and
extend that hypothesis to the FC paradigm. In one side, we found FC memory is harder to retrieve, since

matching between training and testing conditions should be increased in order to effective retrieval. In
addition, we also found that the FC memory expressed under context + tone conditions is temporarily less
stable (Figure 3C). Thus, FC memory formed under inactivation of the LHb could be considered weaker in
those two aspects. Taken together those results indicate a general role of the LHb in defining aversive
memory strength.

Concluding remarks

Inappropriate fear generalization has been postulated as the trigger to psychopathological
conditions like panic or anxiety disorders (Lissek et al., 2010, 2014). Notably, studies on fear generalization
have focused on single components of FC, analyzing either generalization of cued or contextual
components of fear memories, but the interaction between them has not received much attention (Asok
et al., 2019). The results presented here suggest that contextual information influences retrieval of cued
memory, then, cued memory generalization could actually reflect the interaction between cue and
contextual memory generalization. Notably, FC memory formed without proper LHb signaling could be
considered an example of below-normal levels of generalization. Therefore, it could be speculated the
LHb also influences fear generalization. Particularly it would be interesting to analyze fear generalization
in described models of LHb hyperactivity induced mood disturbances (Hu et al., 2020).

## Materials and Methods

Animals

Experiments were performed in male Wistar rats obtained from the vivariums of the Faculty of
Pharmacy and Biochemistry of the Buenos Aires University, Argentina, and Janvier Labs, France. Animals
were 5-6 weeks old at the time of surgery. Animals were housed 4 to 6 per cage, with *ad libitum* access to
food and water, under 12-hour light/dark cycle (lights on at 7:00 am), at constant temperature of 22 ± 2
°C. All the procedures where performed during light hours. Experimental procedures were approved by
the Animal Care and Use Committee of the University of Buenos Aires (CICUAL), and the Autonomous
Community of Madrid (PROEX 167/18).

Surgeries

**Pharmacology:** rats under deep ketamine/xylazine anesthesia (100 and 5 mg/kg respectively)
were bilaterally implanted with 22-Gauge guide cannulae aimed 2.0 mm above the LHb (AP -3.0 mm, ML
±0.7 mm, DV -3.8 mm from Bregma), or 1.0 mm dorsal, ventral or lateral to LHb coordinates in specific

experiments. Cannulae were fixed to the skull with three surgical steel screws and dental acrylic. During

surgery, animals received a dose of analgesic (meloxicam 0.6 mg/kg) and antibiotic (gentamicin 3 mg/kg).

Behavioral procedures began 7-9 days after surgery.

**Optogenetics:** rats under deep isofluorane anesthesia (4 % induction, 1-2 % maintenance, in 0.8

L/minute oxygen) were bilaterally injected with 300 nl of viral vector (AAV8-CamKIIα-ArchT-GFP, 6.2 x $10^{12}$

viral particles/ml from UNC Vector Core, AAV8-CamKIIα-GFP, 6.3 x $10^{12}$ viral particles/ml from UNC Vector

Core, or AAV8-hSyn-oChIEF-tdTomato homemade as described in Proulx et al. (2018)) per side at the LHb

(AP -2.9 mm, ML ± 0.7 mm, DV -5.0 mm from Bregma). Subsequently, optical fiber implants (200 μm core,

0.45 NA) were bilaterally inserted with the tips aiming just above the LHb, with 4 degrees angle from the

sagittal plane (AP -3.0 mm, ML ±1.05 mm, DV -4.5 mm from Bregma), and fixed to the skull with three

surgical steel screws and dental acrylic. During surgery, animals received a dose of analgesic (meloxicam

0.6 mg/kg) and antibiotic (gentamicin 3 mg/kg). Behavioral procedures began three weeks after surgery.

<u>Contextual and Cued Fear Conditioning Training</u>

For three consecutive days before the FC training, rats were habituated to handling once a day.

There, animals were grasped by hand and slightly restrained against the chest of the investigator for 1-3

minutes a manipulation similar to the one performed during intracerebral drug infusions. On training day,

animals were transferred inside their home cage to a room contiguous to the training room where they

acclimatized for 1 hour before the beginning of any other procedure. FC training took place in Context A

(electric foot-shock delivering grid floor, black plastic walls and ceiling with one transparent plexiglass

side, 50x25x30 cm wide, width and height, illuminated by white light). Cued FC protocol consisted of a

180 s baseline period of free exploration followed by 4 tone-shock presentations (17 s, 3 kHz, 80 dB tone

followed by 3 s, 0.60 mA, square monophasic 60 Hz electric shock) with an inter-stimulus interval of 70 s.

Thirty seconds after the last shock, animals were returned to their home cage in the acclimation room.

When training session ended, home cages were left in the acclimation room for at least one hour before

returning them to the animal facility. Ethanol 50 % was used to clean the conditioning cage between

subjects. For contextual FC training procedures were equal but the tone was omitted during conditioning.

Training was video recorded for posterior analysis.

**Pharmacology experiments:** after the acclimation period, rats were randomly assigned to receive

bilateral intra LHb infusions of GABAa agonist, muscimol (Sigma, 60 ng/μl in saline solution, 0.5 μl/side),

or saline. Fluorescent green beads were added to both solutions to aid in checking the infusion site (1:1000

dilution of concentrated 1 μm diameter fluorescent green beads, Bangs Laboratories, USA). During

infusions animals were grasped by hand and slightly restrained in the lap or the arm of the investigator.

Infusions were delivered at a rate of 0.5 µl/minute through a 30-Gauge needle extending 2.0 mm beyond

the tip of implanted guide cannula and connected to a 10 µl Hamilton syringe by a polyethylene tube.

Infusion needle was left in place for an additional minute to minimize backflow. After that animals were

returned to their home cage. FC training began 30 minutes later.

**Optogenetics experiments**: after the acclimation period, rats were taken to the FC room and were

bilaterally connected to 1 to 2 optic fiber branching patch-cords (200 µm core each, Doric Lenses, Canada).

The single end of the branching patch-cord (400 µm core) was connected to a rotary joint (Doric Lenses,

Canada) that was attached to the laser source by a simple patch-cord (400 µm core, 0.45 NA, ThorLabs,

USA). FC chamber (Context A) had a modification in the ceiling (10 cm hole in the center) to allow optic

fiber patch-cord movement. For the ArchT experiment, continuous light of 532 nm at 10 mW at the tip of

the fiber implant was delivered by a laser source (CNI, China), starting at tone onset and stopping 5 s after

shock termination. For the oChIEF experiment, 5 ms pulses at 20 Hz of 447 nm light at 10 mW at the tip

of the fiber implant, were delivered by laser source (Tolket, Argentina) during the whole training. A mixed

control group composed of oChIEF injected without light stimulation and saline injected animals with light

stimulation was used in that experiment since an AAV-tdTomato was not available. At the end of FC

training, patch-cord was disconnected from the fiber implants and animals were returned to home cage.

Contextual FC Test

The contextual test was performed maintaining the same conditions of the FC training day (*e.g.*

acclimation room, FC chamber lights and investigator's nitrile gloves). During test, animals were placed in

the training context (Context A) for 180 s and then returned to its home cage. Test was video recorded

for posterior analysis.

Cued FC Test

Several precautions were taken to avoid generalization during cued test. Home cages were moved

to a different acclimation room of the one used in the training day, the investigator used latex gloves,

animals were transported between the acclimation room and the test chamber within a plastic box, and

acetic acid 1 % m/v was employed for cleaning between subjects. Test was performed in Context B, a

white acrylic box with a transparent plexiglass ceiling (25x25x40 cm wide, width and height) with floor

covered in wood shaving and lit up with red light. Cue test consisted of a pre-tone period of 180 s, followed

by 60 s tone (same as training tone) and 30 s post-shock period. After that, animals were removed from

the test cage and placed back in the home cage. Test was video recorded for posterior analysis. 1 hour

after finishing tests, home cages were taken back to the animal facility.

<u>Context + Tone Test</u>

Context + tone test proceeded as contextual FC test but after 180 s in the test cage the tone was

presented for 60 s. 30 s after tone ending animals were returned to their home cage.

<u>Open Field</u>

Animals were placed for 15 minutes into an open field (OF; square grey plastic floor and walls 50

cm side x 35 cm height) and video recorded. Behavior during OF was analyzed offline with ANY-Maze video

tracking system (v4.82, Stoelting Co., USA).

<u>Histologic Control</u>

Within a week after the end of behavioral procedures animals were euthanized with a lethal dose

of ketamine/xylazine, 50 ml of ice-cold phosphate-buffered saline (PBS) 0.1 M was transcardially perfused,

followed by 50 ml of paraformaldehyde 4 %. After that, the brain extracted and incubated at 4 °C in

paraformaldehyde 4 % overnight. Then, brain was transferred to 30 % m/v sucrose in PBS 0.1 M, incubated

for three days at 4 °C and 150 μm coronal sections were cut at a freezing microtome. Sections were

mounted with glycerol 50 % v/v in PBS 0.1 M, examined and photographed at 40X magnification with GFP

and DAPI filters. For the pharmacological experiments, injection sites were determined by the presence

of green beads. Animals in which both infusions were correct were included in the analysis. Animals from

the cohorts used in Figure 2B-C in which both infusions were outside the LHb were included in the miss

group presented in Figure S3. For the optogenetics experiments, animals with transfection and/or optic

fiber misses were not considered in the analysis.

<u>Freezing analysis</u>

Freezing was manually scored offline. Experimenter was blinded for animal´s treatment. An

Arduino custom system was used for the manual scoring. Briefly, a button was used to synchronize the

start of the video and another button was used to register the freezing state while pressed. The Arduino

system communicated time and freezing-button state at 10 Hz to the computer, and then saved as CSV

file for later analysis. For freezing over time graphs, data was binned in 5 s intervals, and inter-subjects´

mean and SEM was calculated.

     Statistical Analysis

Each animal was taken as an independent measure. During test session of cued or context + tone

experiments, average freezing before and during tone presentation were treated as repeated measures.

Thus, in most experiments analysis was done using a two-way repeated measures design in which the

analyzed factors were "test stage" (pre-tone or tone) and "treatment" (muscimol, optogenetic excitation

or inhibition, or the corresponding control group). In experiments in which subjects were tested more

than one day, "testing day" was considered an additional factor. Freezing during training session was

analyzed separately. There the stage factor comprised freezing during tone or the inter-stimulus period.

Statistical analysis of FC experiments was done by generalized linear mixed modelling (GzLMM). Typical

distribution of individuals freezing values accumulated at lower and higher values with few individuals

freezing around the group mean, indicating a non-normal distribution. We therefore compared models

with beta and gaussian distribution (rescaling freezing to y"=[y'(n-1)+0.5]/n as suggested in Smithson &

Verkuilen (2006)). For the model fitting in R (R Foundation for Statistical Computing, Vienna, Austria) we

used the 'glmmTMB' package (Brooks et al., 2017). Beta distribution provided better adjustment and less

tendencies between group residues than gaussian distribution (data not shown). Logit link function was

employed. For the GzLMM we considered every interaction between the corresponding fixed factors (test

stage, treatment, testing day) and subject was treated as a random factor with random intercepts. ANOVA

and contrasts were done applying 'car' and 'emmeans' packages (Lenth, 2020; Weisberg, 2019) where

GzLMM was used. GzLMM ANOVAs used Wald $\chi^2$ statistics for comparisons since neither likelihood ratio

tests nor *F* tests are supported. Assumptions of homoscedasticity and normal residues were met in the

open field experiment; therefore, it was analyzed via conventional two-way repeated measures ANOVA.

Each experiment was repeated at least twice in separate cohorts of animals. Each cohort included all the

treatments. Sample size calculations were done using power analysis. From pilot context test

experiments, we considered a standard deviation of aprox. 20 % aiming to detect a minimal difference

among groups of 50 %, with an 80 % power, and a type I error of 5 %. Based on these parameters and first

assuming gaussian distribution, initially we aimed at an n = 14. Later we readjusted to n = 6 on cued

memory tests, where the observed difference between vehicle and muscimol infused groups (aprox. 60

%), standard deviation (aprox. 25 %) and beta distribution at tone tests allowed to maintain about 80 %

power and 5 % type I error. For experiment in Figure 4C (reconsolidation of contextual memory) statistical

evaluation of the effect of the recall session was done following a within subject design since recall session

on 7$^{th}$ day and test session on 8$^{th}$ day were replicates. In bar graphs, freezing during training is included as

a parameter of the freezing of a naïve animal but, as mentioned above, freezing during training was
analyzed separately.

## Acknowledgements

We would like to thank Dr. Roberto Malinow for his help in the generation of AAV vectors, to
Yamila Paez for her technical assistance, and to Sadegh Nabavi, Mariano Belluscio, Sebastian Giusti and
Paula Ospital for their helpful comments. **Funding**: this work was supported by the following grants: PICT
2016-0034 from the National Agency for Scientific and Technological Promotion of Argentina (ANPCyT) to
JM; I-COOP+(CSIC) ref COOPA20198 to PM; NARSAD, Young Investigator Award 2015 (#23861) from the
Brain and Behavior Foundation, PICT 2015-2609 and PICT 2017-4594 from the ANPCyT to JP. TS was
supported by a predoctoral fellowship from the CONICET. **Authors contributions**: Design: TES, MRI, JHM,
PM, JP; Pilot experiments: CP; Experiments: TES, MRI, JP, PM; Analysis: TES, MRI, PM, JP; Writing: TES,
MRI, CP, JP. **Competing interests**: the authors declare that they have no competing interests. **Data and
materials availability**: all data needed to evaluate the conclusions in the paper are present in the paper
and/or the Supplementary Information. Additional data are available from authors upon request.

## References

Aizawa, H., Yanagihara, S., Kobayashi, M., Niisato, K., Takekawa, T., Harukuni, R., McHugh, T. J., Fukai, T.,
Isomura, Y., & Okamoto, H. (2013). The synchronous activity of lateral habenular neurons is essential
for regulating hippocampal theta oscillation. *The Journal of Neuroscience : The Official Journal of the
Society for Neuroscience*, *33*(20), 8909–8921. https://doi.org/10.1523/JNEUROSCI.4369-12.2013
Asok, A., Kandel, E. R., & Rayman, J. B. (2019). The Neurobiology of Fear Generalization. *Frontiers in
Behavioral Neuroscience*, *12*(January), 1–15. https://doi.org/10.3389/fnbeh.2018.00329
Baker, P. M., Rao, Y., Rivera, Z. M. G., Garcia, E. M., & Mizumori, S. J. Y. (2019). Selective Functional
Interaction Between the Lateral Habenula and Hippocampus During Different Tests of Response
Flexibility. *Frontiers in Molecular Neuroscience*, *12*, 245. https://doi.org/10.3389/fnmol.2019.00245
Barrett, D. W., & Gonzalez-Lima, F. (2018). Prefrontal-limbic Functional Connectivity during Acquisition
and　　　Extinction　　　of　　　Conditioned　　　Fear. *Neuroscience*, *376*, 162–171.
https://doi.org/10.1016/j.neuroscience.2018.02.022
Bekinschtein, P., Cammarota, M., Igaz, L. M., Bevilaqua, L. R. M., Izquierdo, I., & Medina, J. H. (2007).
Persistence of Long-Term Memory Storage Requires a Late Protein Synthesis- and BDNF- Dependent
Phase in the Hippocampus. *Neuron*, *53*(2), 261–277. https://doi.org/10.1016/j.neuron.2006.11.025

Bekinschtein, P., Cammarota, M., Katche, C., Slipczuk, L., Rossato, J. I., Goldin, A., Izquierdo, I., & Medina,
J. H. (2008). BDNF is essential to promote persistence of long-term memory storage. *Proceedings of*
*the National Academy of Sciences*, *105*(7), 2711–2716. https://doi.org/10.1073/pnas.0711863105

Bertone-Cueto, N. I., Makarova, J., Mosqueira, A., García-Violini, D., Sánchez-Peña, R., Herreras, O.,
Belluscio, M., & Piriz, J. (2020). Volume-Conducted Origin of the Field Potential at the Lateral
Habenula. In *Frontiers in Systems Neuroscience* (Vol. 13, p. 78).
https://doi.org/10.3389/fnsys.2019.00078

Bocchio, M., Nabavi, S., & Capogna, M. (2017). Synaptic Plasticity, Engrams, and Network Oscillations in
Amygdala Circuits for Storage and Retrieval of Emotional Memories. *Neuron*, *94*(4), 731–743.
https://doi.org/10.1016/j.neuron.2017.03.022

Brooks, M. E., Kristensen, K., van Benthem, K. J., Magnusson, A., Berg, C. W., Nielsen, A., Skaug, H. J.,
Mächler, M., & Bolker, B. M. (2017). glmmTMB balances speed and flexibility among packages for
zero-inflated generalized linear mixed modeling. *R Journal*, *9*(2), 378–400.
https://doi.org/10.32614/rj-2017-066

Dunsmoor, J. E., & Paz, R. (2015). Fear Generalization and Anxiety: Behavioral and Neural Mechanisms.
*Biological Psychiatry*, *78*(5), 336–343. https://doi.org/10.1016/j.biopsych.2015.04.010

Durieux, L., Mathis, V., Herbeaux, K., Muller, M., Barbelivien, A., Mathis, C., Schlichter, R., Hugel, S.,
Majchrzak, M., & Lecourtier, L. (2020). Involvement of the lateral habenula in fear memory. *Brain*
*Structure and Function*, *0123456789*. https://doi.org/10.1007/s00429-020-02107-5

Etkin, A., Alarcón, J. M., Weisberg, S. P., Touzani, K., Huang, Y. Y., Nordheim, A., & Kandel, E. R. (2006). A
Role in Learning for SRF: Deletion in the Adult Forebrain Disrupts LTD and the Formation of an
Immediate Memory of a Novel Context. *Neuron*, *50*(1), 127–143.
https://doi.org/10.1016/j.neuron.2006.03.013

Goutagny, R., Loureiro, M., Jackson, J., Chaumont, J., Williams, S., Isope, P., Kelche, C., Cassel, J.-C., &
Lecourtier, L. (2013). Interactions between the Lateral Habenula and the Hippocampus: Implication
for Spatial Memory Processes. *Neuropsychopharmacology*, *38*(12), 2418–2426.
https://doi.org/10.1038/npp.2013.142

Hikosaka, O. (2010). The habenula: from stress evasion to value-based decision-making. *Nature Reviews.*
*Neuroscience*, *11*(7), 503–513. https://doi.org/10.1038/nrn2866

Hu, H., Cui, Y., & Yang, Y. (2020). Circuits and functions of the lateral habenula in health and in disease.
*Nature Reviews Neuroscience*, *21*(5), 277–295. https://doi.org/10.1038/s41583-020-0292-4

Kim, W. Bin, & Cho, J.-H. (2020). Encoding of contextual fear memory in hippocampal-amygdala circuit.

479   *Nature Communications*, *11*(1), 1382. https://doi.org/10.1038/s41467-020-15121-2

480 Kim, J., & Fanselow, M. (1992). Modality-specific retrograde amnesia of fear. *Science*, *256*(5057), 675–

481   677. https://doi.org/10.1126/science.1585183

482 LeDoux, J. E., & Phillips, R. (1992). Differential contribution of amygdala and hippocampus to cued and

483   contextual fear conditioning. *Behavioral Neuroscience*, *106*(2), 274–285.

484   http://psycnet.apa.org/journals/bne/106/2/274.html

485 Lenth, R. (2020). *emmeans: Estimated Marginal Means, aka Least-Squares Means.* (R package version

486   1.4.7). https://cran.r-project.org/package=emmeans

487 Lin, J. Y., Lin, M. Z., Steinbach, P., & Tsien, R. Y. (2009). Characterization of engineered channelrhodopsin

488   variants with improved properties and kinetics. *Biophysical Journal*, *96*(5), 1803–1814.

489   https://doi.org/10.1016/j.bpj.2008.11.034

490 Lissek, S., Kaczkurkin, A. N., Rabin, S., Geraci, M., Pine, D. S., & Grillon, C. (2014). Generalized Anxiety

491   Disorder Is Associated With Overgeneralization of Classically Conditioned Fear. *Biological Psychiatry*,

492   *75*(11), 909–915. https://doi.org/10.1016/j.biopsych.2013.07.025

493 Lissek, S., Rabin, S., Heller, R. E., Lukenbaugh, D., Geraci, M., Pine, D. S., & Grillon, C. (2010).

494   Overgeneralization of Conditioned Fear as a Pathogenic Marker of Panic Disorder. *American Journal*

495   *of Psychiatry*, *167*(1), 47–55. https://doi.org/10.1176/appi.ajp.2009.09030410

496 Maren, S. (2008). Pavlovian fear conditioning as a behavioral assay for hippocampus and amygdala

497   function: cautions and caveats. *European Journal of Neuroscience*, *28*(8), 1661–1666.

498   https://doi.org/10.1111/j.1460-9568.2008.06485.x

499 Maren, S., Phan, K. L., & Liberzon, I. (2013). The contextual brain: Implications for fear conditioning,

500   extinction and psychopathology. *Nature Reviews Neuroscience*, *14*(6), 417–428.

501   https://doi.org/10.1038/nrn3492

502 Maren, S., & Quirk, G. J. (2004). Neuronal signalling of fear memory. *Nature Reviews Neuroscience*, *5*(11),

503   844–852. https://doi.org/10.1038/nrn1535

504 Nabavi, S., Fox, R., Proulx, C. D., Lin, J. Y., Tsien, R. Y., & Malinow, R. (2014). Engineering a memory with

505   LTD and LTP. *Nature*, *511*(7509), 348–352. https://doi.org/10.1038/nature13294

506 Proulx, C. D., Aronson, S., Milivojevic, D., Molina, C., Loi, A., Monk, B., Shabel, S. J., & Malinow, R. (2018).

507   A neural pathway controlling motivation to exert effort. *Proceedings of the National Academy of*

508   *Sciences*, *115*(22), 5792–5797. https://doi.org/10.1073/pnas.1801837115

509 Proulx, C. D., Hikosaka, O., & Malinow, R. (2014). Reward processing by the lateral habenula in normal and

510   depressive behaviors. *Nature Neuroscience*, *17*(9), 1146–1152. https://doi.org/10.1038/nn.3779

Rossato, J. I., Bevilaqua, L. R. M., Izquierdo, I., Medina, J. H., & Cammarota, M. (2009). Dopamine Controls

Persistence of Long-Term Memory Storage. *Science*, *325*(5943), 1017–1020.

https://doi.org/10.1126/science.1172545

Rumpel, S., LeDoux, J., Zador, A., & Malinow, R. (2005). Postsynaptic Receptor Trafficking Underlying a

Form of Associative Learning. *Science*, *308*(5718), 83–88. https://doi.org/10.1126/science.1103944

Sah, P., Westbrook, R. F., & Lüthi, A. (2008). Fear Conditioning and Long-term Potentiation in the

Amygdala. *Annals of the New York Academy of Sciences*, *1129*(1), 88–95.

https://doi.org/10.1196/annals.1417.020

Smithson, M., & Verkuilen, J. (2006). A better lemon squeezer? Maximum-likelihood regression with beta-

distributed dependent variables. *Psychological Methods*, *11*(1), 54–71.

https://doi.org/10.1037/1082-989X.11.1.54

Song, M., Jo, Y. S., Lee, Y. K., & Choi, J. S. (2017). Lesions of the lateral habenula facilitate active avoidance

learning and threat extinction. *Behavioural Brain Research*, *318*, 12–17.

https://doi.org/10.1016/j.bbr.2016.10.013

Stefanelli, T., Bertollini, C., Lüscher, C., Muller, D., & Mendez, P. (2016). Hippocampal Somatostatin

Interneurons Control the Size of Neuronal Memory Ensembles. *Neuron*, *89*(5), 1074–1085.

https://doi.org/10.1016/j.neuron.2016.01.024

Stujenske, J. M., Spellman, T., & Gordon, J. A. (2015). Modeling the Spatiotemporal Dynamics of Light and

Heat Propagation for InVivo Optogenetics. *Cell Reports*, *12*(3), 525–534.

https://doi.org/10.1016/j.celrep.2015.06.036

Szőnyi, A., Sos, K. E., Nyilas, R., Schlingloff, D., Domonkos, A., Takács, V. T., Pósfai, B., Hegedüs, P., Priestley,

J. B., Gundlach, A. L., Gulyás, A. I., Varga, V., Losonczy, A., Freund, T. F., & Nyiri, G. (2019). Brainstem

nucleus incertus controls contextual memory formation. *Science (New York, N.Y.)*, *364*(6442), 0–14.

https://doi.org/10.1126/science.aaw0445

Szőnyi, A., Zichó, K., Barth, A. M., Gönczi, R. T., Schlingloff, D., Török, B., Sipos, E., Major, A., Bardóczi, Z.,

Sos, K. E., Gulyás, A. I., Varga, V., Zelena, D., Freund, T. F., & Nyiri, G. (2019). Median raphe controls

acquisition of negative experience in the mouse. *Science*, *366*(6469), eaay8746.

https://doi.org/10.1126/science.aay8746

Tomaiuolo, M., Gonzalez, C., Medina, J. H., & Piriz, J. (2014). Lateral Habenula determines long-term

storage of aversive memories. *Frontiers in Behavioral Neuroscience*, *8*, 170.

https://doi.org/10.3389/fnbeh.2014.00170

Trusel, M., Nuno-Perez, A., Lecca, S., Harada, H., Lalive, A. L., Congiu, M., Takemoto, K., Takahashi, T.,

Ferraguti, F., & Mameli, M. (2019). Punishment-Predictive Cues Guide Avoidance through

Potentiation of Hypothalamus-to-Habenula Synapses. *Neuron*, *102*(1), 120-127.e4.

https://doi.org/10.1016/j.neuron.2019.01.025

Vianna, M. R. M. (2000). Role of Hippocampal Signaling Pathways in Long-Term Memory Formation of a

Nonassociative Learning Task in the Rat. *Learning & Memory*, *7*, 333–340.

https://doi.org/10.1101/lm.34600

Wang, D., Li, Y., Feng, Q., Guo, Q., Zhou, J., & Luo, M. (2017). Learning shapes the aversion and reward

responses of lateral habenula neurons. *ELife*, *6*, e23045. https://doi.org/10.7554/elife.23045

Wang, Z., Wang, L., Yamamoto, R., Sugai, T., & Kato, N. (2013). Role of the lateral habenula in shaping

context-dependent locomotor activity during cognitive tasks. *NeuroReport*, *24*(6), 276–280.

https://doi.org/10.1097/WNR.0b013e32835ee1e8

Weisberg, J. F. and S. (2019). *An {R} Companion to Applied Regression.* (S. Publications (ed.); Third Edit).

Yang, Y., Cui, Y., Sang, K., Dong, Y., Ni, Z., Ma, S., & Hu, H. (2018). Ketamine blocks bursting in the lateral

habenula to rapidly relieve depression. *Nature*, *554*(7692), 317–322.

https://doi.org/10.1038/nature25509

## Fig. 1

<u>Figure 1. Inactivation of the LHb blocks the formation of contextual FC long-term memory.</u>

**A)** Experiment diagram: bilateral vehicle/muscimol intra LHb infusions were performed 30 minutes before

training. Contextual FC was tested 24 hours or 7 days later. During training subjects were let to freely

explore the cage for a baseline period of 197 s. After that they were exposed to 4 foot-shocks (0.6 mA, 3

s) interspaced by 70 s. In the test session animals were re-exposed to the training context for 180 s and

freezing was quantified.

**B)** Freezing during cued FC training for the group tested 7 days after training. Left panel: freezing over

time, shocks were presented at times indicated by the orange shaded areas. Right panel: average freezing

at inter-shock periods. Freezing in both groups was equivalent (GzLMM t test: $t_{(20)}$ = 0.046, p = 0.9635;

$n_{vehicle}$ = 11, $n_{muscimol}$ = 12).

**C, D)** Seven days and 24 hours context tests. Left panels: freezing over time for groups tested at 7 days (C)

and 24 hours (D). Right panels: average freezing. In both experiments freezing in the muscimol group was

lower than in the control group (GzLMM t test for 7 days context test: $t_{(20)}$ = 4.294, p = 0.0004, $n_{vehicle}$ = 11,

$n_{muscimol}$ = 12; GzLMM t test for 24 hours context test: $t_{(16)}$ = 3.964, p = 0.0011; $n_{vehicle}$ = 9, $n_{muscimol}$ = 10.

In freezing over time plots, lines represent intersubjects' mean, and shaded area represents +SEM. In bar

plots, each dot represents a subject and bars represent mean and errors bars +SEM. ns p > 0.05, ** p <

0.01, *** p < 0.001.

**Fig. 2**

Figure 2. Inactivation of the LHb blocks the formation of cued FC long-term memory.

**A)** Experiment diagram: bilateral vehicle/muscimol intra LHb infusions were performed 30 minutes before training. Cued FC was tested 24 hour or 7 days later. During training, subjects were let to freely explore the cage for a baseline period of 180 s that was followed by 4 tone-shock pairings (17 s of tone followed by 3 s, 0.6 mA shock) interspaced by 70 s. During test, tone was presented for 60 s after a pre-tone period of 180 s.

**B)** Freezing during cued FC training for the group tested 7 days after training. Left panel: freezing over

time. Tones were presented during the times indicated by the gray shaded areas. Shocks were presented

at times indicated by the orange shaded areas. Right panel: mean freezing during tone and post-shock

period. Freezing in both groups was equivalent. GzLMM Two-way repeated measures ANOVA of

treatment, training stage, and interaction: $\chi^2$ treatment $_{(1)}$ = 0.185, p = 0.6668; $\chi^2$ stage $_{(1)}$ = 6.308, p =

0.0120; $\chi^2$ interaction $_{(1)}$ = 0.028, p = 0.8661. Sidak's multiple comparisons test: Post-shock freezing: vehicle

vs. muscimol: $t_{(28)}$ = 0.430, p = 0.8912; Tone freezing: vehicle vs. muscimol: $t_{(28)}$ = 0.297, p = 0.9463; $n_{vehicle}$

= 8, $n_{muscimol}$ = 9.

**C, D)** Seven days and 24 hours cued FC tests. Left panels: freezing over time for groups tested 7 days (C)

and 24 hours (D) after training. Gray areas indicate tone presentation. Right panels: average freezing for

pre- and tone period. Freezing during pre-tone period was not different between vehicle and muscimol

groups in either experiment. In contrast, during tone presentation, a highly significant reduction in

freezing was observed in the muscimol group in both experiments. GzLMM Two-way repeated measures

ANOVA of treatment, test stage, and interaction for 7 days experiment: $\chi^2$ treatment $_{(1)}$ = 2.157, p = 0.1420;

$\chi^2$ stage $_{(1)}$ = 19.100, p < 0.0001; $\chi^2$ interaction $_{(1)}$ = 32.939, p < 0.0001. Sidak's multiple comparisons test:

Pre-tone freezing: vehicle vs. muscimol: $t_{(28)}$ = 1.469, p = 0.2828; Tone freezing: vehicle vs. muscimol: $t_{(28)}$

= 5.396, p < 0.0001; $n_{vehicle}$ = 8, $n_{muscimol}$ = 9. GzLMM Two-way repeated measures ANOVA of treatment,

test stage, and interaction for 24 hours experiment: $\chi^2$ treatment $_{(1)}$ = 0.024, p = 0.8781; $\chi^2$ stage $_{(1)}$ = 0.108,

p = 0.7428; $\chi^2$ interaction $_{(1)}$ = 12.447, p = 0.0004. Sidak's multiple comparisons test: Pre-tone freezing:

vehicle vs. muscimol: $t_{(28)}$ = 0.153, p = 0.9854; Tone freezing: vehicle vs. muscimol: $t_{(28)}$ = 4.441, p = 0.0003;

$n_{vehicle}$ = 9, $n_{muscimol}$ = 8.

In freezing over time plots, line represents intersubjects' mean, and shaded area represents SEM. In bar

plots each dot represents a subject and bars represent mean + SEM. ns p > 0.05, *** p < 0.001, **** p <

0.0001.

# Fig. 3

Figure 3. FC memory formed under inactivation of the LHb could be retrieved in context + tone conditions and showed decreased temporal stability.

**A)** Experiment diagram: bilateral intra LHb infusions of vehicle/muscimol were performed 30 minutes before training. Animals were trained in cued FC. Test was performed 7 or 21 days later in the same context used for training (Context A). The tone was presented for 60 s after a pre-tone period of 180 s.

**B)** Freezing during context + tone test 7 days after training. Left panel: freezing over time. Gray area indicates tone presentation. Right panel: average freezing for pre- and tone period. Freezing to the context was higher in vehicle group. However, tone elicited a robust freezing in the muscimol group that reached values equivalent to the vehicle group. GzLMM Two-way repeated measures ANOVA of treatment, test stage, and interaction: $\chi^2$ treatment $_{(1)}$ = 6.830, p = 0.0090; $\chi^2$ stage $_{(1)}$ = 44.992, p < 0.0001; $\chi^2$ interaction $_{(1)}$ = 3.161, p = 0.0754. Sidak's multiple comparisons test: Pre-tone freezing: vehicle vs.

muscimol: $t_{(46)}$ = 2.613, p = 0.0474; Tone freezing: vehicle vs. muscimol: $t_{(46)}$ = 0.899, p = 0.8460; Vehicle:

pre-tone vs. tone freezing: $t_{(46)}$ = 6.014, p < 0.0001; muscimol: pre-tone vs. tone freezing: $t_{(46)}$ = 6.708, p <

0.0001; $n_{vehicle}$ = 13, $n_{muscimol}$ = 13.

**C)** Freezing during context + tone test 21 days after training. Left panel: freezing over time. Gray area

indicates tone presentation. Right panel: average freezing for pre- and tone period. 21 days after training

freezing in context + tone condition in the muscimol group was lower than in the vehicle group indicating

a diminished temporal stability of context + tone memory. GzLMM Two-way repeated measures ANOVA

of treatment, test stage, and interaction: $\chi^2$ treatment $_{(1)}$ = 7.123, p = 0.0076; $\chi^2$ stage $_{(1)}$ = 41.364, p <

0.0001; $\chi^2$ interaction $_{(1)}$ = 0.176, p = 0.6751. Sidak's multiple comparisons test: Pre-tone freezing: vehicle

vs. muscimol: $t_{(26)}$ = 2.669, p = 0.0508; Tone freezing: vehicle vs. muscimol: $t_{(26)}$ = 2.683, p = 0.0491; Vehicle:

pre-tone vs. tone freezing: $t_{(26)}$ = 7.354, p < 0.0001; muscimol: pre-tone vs. tone freezing: $t_{(26)}$ = 6.431, p <

0.0001; $n_{vehicle}$ = 8, $n_{muscimol}$ = 8.

In freezing over time plots, line represents intersubjects' mean, and shaded area represents +SEM. In bar

plots each dot represents a subject and bars represent mean + SEM. ns, p > 0.05, * p < 0.05.

## Fig. 4

**Figure 4. Retrieval of FC memory in context + tone conditions reconsolidates cued but not contextual memories.**

**A)** Diagram of reconsolidation experiment of cued FC memory: bilateral intra LHb muscimol infusions were performed 30 minutes before cued FC training as described previously. Seven days later, animals of the reconsolidation group were subjected to a recall session in which the cue was presented in the training context. The following day, reconsolidation of cued memory was tested in context "B". The no-reconsolidation group was not exposed to the recall session and was tested for cued memory in context "B" 8 days after training.

**B)** Retrieval of FC memory in context + tone conditions restores cued FC memory. Left and center: freezing over time for the recall and the cued memory test sessions, respectively. Animals of the no-reconsolidation group showed very low levels of freezing, confirming the impairment in cued memory

induced by inactivation of the LHb during training. In contrast animals of the reconsolidation group

presented a marked freezing to the tone during the test session, reaching levels of freezing equivalent to

context + tone condition during the recall session. Right: quantification of freezing. Freezing to the tone

in the test session was significantly higher in the reconsolidation group. GzLMM Two-way repeated

measures ANOVA of reconsolidation, test stage, and interaction during tone test session: $\chi^2$

reconsolidation $_{(1)}$ = 0.207, p = 0.6489; $\chi^2$ stage $_{(1)}$ = 0.009, p = 0.9232; $\chi^2$ interaction $_{(1)}$ = 9.170, p = 0.0025.

Sidak's multiple comparisons test: Pre-tone freezing: reconsolidation vs. no-reconsolidation: $t_{(18)}$ = 0.455,

p = 0.8805; Tone freezing: reconsolidation vs. no-reconsolidation: $t_{(18)}$ = 4.219, p = 0.0010; $n_{reconsolidation}$ =

6, $n_{NO-reconsolidation}$ = 6.

**C)** Diagram of the reconsolidation experiment of contextual FC memory: bilateral intra LHb muscimol

infusions were performed 30 minutes before cued FC training as described previously. Seven days later,

animals were subjected to a recall session in which the cue was presented in the training context. The

following day animals underwent the same procedure.

**D)** Retrieval of FC memory in context + tone conditions does not restore contextual FC memory. Left and

center: freezing over time for the recall and the test sessions, respectively. Right: paired comparison of

freezing for each subject on the recall and the test session. No differences in freezing were observed

between sessions indicating that retrieval of FC memory during the recall session did not restore

contextual FC memory, as it did with cued memory. GzLMM Two-way repeated measures ANOVA of

session day, test stage, and interaction: $\chi^2$ day $_{(1)}$ = 0.514, p = 0.4732; $\chi^2$ stage $_{(1)}$ = 20.398, p < 0.0001; $\chi^2$

interaction $_{(1)}$ = 0.240, p = 0.6240. Sidak's multiple comparisons test: Pre-tone freezing: recall vs. test day:

$t_{(26)}$ = 0.717, p = 0.7292; Tone freezing: recall vs. test day: $t_{(26)}$ = 0.088, p = 0.9952; n = 6.

In freezing over time plots, line represents intersubjects' mean and shaded area represents +SEM, and

gray area indicates tone presentation. In bar plots each dot represents a subject and bars represent mean

+ SEM. ns p > 0.05, ** p < 0.01.

## Fig. 5

Figure 5. Optogenetic inactivation of the LHb during cue and US, reduced cued but not contextual FC memories.

**A)** Experiment diagram: Top-Left: animals were bilaterally transfected with AAV-ArchT-GFP or AAV-GFP in the LHb and implanted with optic fibers above the LHb 4 weeks before training. Top-Right: during training, optogenetic light stimulation was delivered starting with the tone and stopping 5 seconds after the shock (tone and shock presentations were as previously described for cued FC). Bottom: diagram of training and

tests. Cued memory was tested 7 days after training in Context B. The same animals were tested the

following day in context + tone condition to evaluate contextual FC memory and freezing in context + tone

condition.

**B)** Microphotographs of the AAV-ArchT-GFP infection (top, middle, bottom: DAPI, GFP, and merge

respectively). Dashed white lines in the middle panel delimitates brain structures. * indicates the optic

fiber tract. MHb: medial habenula, sm: stria medullaris, 3V: third ventricle. Scale bars: 1 mm.

**C, D)** Freezing over time during tone test at day 7 (C) and during context + tone test at day 8 (D). Gray area

indicates tone presentation. Line represents intersubjects' mean, and shaded area represents +SEM.

**E)** Average freezing on tone test, and context + tone test sessions. During tone test ArchT group displayed

lower levels of freezing to the tone than GFP group. The following day, during context + tone test, freezing

levels of the ArchT group to both to the context and to the tone were equivalent to those of the GFP

group.

GzLMM Three-way repeated measures ANOVA of test day, test stage, optogenetic inhibition and

interactions: $\chi^2$ day $_{(1)}$ = 24.559, p < 0.0001; $\chi^2$ stage $_{(1)}$ = 23.070, p < 0.0001; $\chi^2$ inhibition $_{(1)}$ = 0.5568, p =

0.4555; $\chi^2$ interaction(Day×Stage) $_{(1)}$ = 1.659, p = 0.1978; $\chi^2$ interaction(Day×Inhibition) $_{(1)}$ = 0.294, p =

0.5879; $\chi^2$ interaction(Stage×Inhibition) $_{(1)}$ = 5.006, p = 0.0253; $\chi^2$ interaction(Day×Stage×Inhibition) $_{(1)}$ =

1.088, p = 0.2970. Sidak's multiple comparisons test: Day7: Pre-tone freezing: GFP vs. ArchT: $t_{(66)}$ = 0.746,

p = 0.9747; Day 7: Tone freezing: GFP vs. ArchT: $t_{(66)}$ = 3.410, p = 0.0067; Day 8: Pre-tone freezing: GFP vs.

ArchT: $t_{(66)}$ = 0.178, p = 1.0000; Day 8: Tone freezing: GFP vs. ArchT: $t_{(66)}$ = 1.145, p = 0.8308; ArchT: Tone

freezing: Day 7 vs. Day 8: $t_{(66)}$ = 3.366, p = 0.0076; Day 8: ArchT: Pre-tone vs. Tone freezing: $t_{(66)}$ = 3.475, p

= 0.0054; $n_{GFP}$ = 9, $n_{ArchT}$ = 10. In bar plots each dot represents a subject and bars represent mean + SEM.

ns p > 0.05, ** p < 0.01.

**Fig. 6**

Figure 6. Disruption of LHb neuronal activity during complete FC training impairs contextual and cued FC memories.

**A)** Experiment diagram: Top-Left: animals were bilaterally infected with AAV-oChIEF-tdTomato in the LHb and implanted with optic fibers above the LHb 4 weeks before training. Top-Right: during whole training session the LHb was stimulated with 5 ms light pulses at 20 Hz to disrupt endogenous neuronal activity. Bottom: diagram of training and tests. Cued memory was tested 7 days after training in Context B. The

same animals were tested the following day in context + tone condition to evaluate contextual FC memory

and freezing in context + tone condition.

**B)** Microphotographs of the AAV-oChIEF-tdTomato infection (top, middle, bottom: DAPI, tdTomato

fluorescence, and merge respectively). Dashed white lines in the middle panel delimitates brain

structures. * indicates the optic fiber tract. MHb: medial habenula, sm: stria medullaris, 3V: third ventricle.

Scale bars: 1 mm.

**C, D)** Freezing over time during cued test at day 7 (C) and during context + tone test at day 8 (D). Gray

area indicates tone presentation. Line represents intersubjects' mean, and shaded area represents + SEM.

**E)** Average freezing of tone test and context + tone test sessions. Disruption of endogenous LHb activity

during training by optogenetic stimulation induced deficits in cued and contextual memories:

GzLMM Three-way repeated measures ANOVA of test day, test stage, optogenetic stimulation and

interactions: $\chi^2$ day $_{(1)}$ = 20.439, p < 0.0001; $\chi^2$ stage $_{(1)}$ = 100.217, p < 0.0001; $\chi^2$ stimulation $_{(1)}$ = 5.135, p =

0.0234; $\chi^2$ interaction(Day×Stage) $_{(1)}$ = 5.805, p = 0.0160; $\chi^2$ interaction(Day×stimulation) $_{(1)}$ = 2.048, p =

0.15237; $\chi^2$ interaction(Stage×stimulation) $_{(1)}$ = 2.489, p = 0.1146; $\chi^2$ interaction(Day×Stage×stimulation)

$_{(1)}$ = 4.905, p = 0.0268. Sidak's multiple comparisons test: Day 7: Pre-tone freezing: Control vs. oChIEF: $t_{(42)}$

= 2.266, p = 0.1601; Day 7: Tone freezing: Control vs. oChIEF: $t_{(42)}$ = 4.454, p = 0.0004; Day 8: Pre-tone

freezing: Control vs. oChIEF: $t_{(42)}$ = 3.874, p = 0.0022; Day 8: Tone freezing: Control vs. oChIEF: $t_{(42)}$ = 2.995,

p = 0.0272; oChIEF: Tone freezing: Day 7 vs. Day 8: $t_{(42)}$ = 3.810, p = 0.0027; Day 8: oChIEF: Pre-tone vs.

Tone freezing: $t_{(42)}$ = 7.283, p < 0.0001; $n_{Control}$ = 7, $n_{oChIEF}$ = 6. In bar plots, each dot represents a subject

and bars represent mean + SEM. ns p > 0.05, * p < 0.05, ** p < 0.01 *** p < 0.001.

## Suppl. Fig. 1

735    Supplementary Figure 1. Freezing during training for experimental groups included in Figures 1-3.

**A-D)** Left: temporal course of freezing during training in the experiments specified in the plot title. Right: average freezing of those experiments. In either condition no significant differences were observed between muscimol and vehicle group. In freezing over time plots orange shaded areas indicate shock presentation, gray shaded areas indicate tone presentation, line plots represent intersubjects' mean and shaded area over line plots represents +SEM. In bar plots each dot represents a subject and bars represent mean +SEM. ns $p > 0.05$.

Statistics:

- Contextual FC training – 24 hours test (A): GzLMM t test: $t_{(16)} = 1.217$, $p = 0.2414$. $n_{vehicle} = 9$, $n_{muscimol} = 10$.

- Cued FC training – 24 hours test (B): GzLMM two-way repeated measures ANOVA of treatment, training stage, and interaction: $\chi^2$ treatment $_{(1)} = 3.796$, $p = 0.0514$; $\chi^2$ stage $_{(1)} = 0.550$, $p = 0.4584$; $\chi^2$ interaction $_{(1)} = 0.852$, $p = 0.3560$. Sidak's multiple comparisons test: Post-shock freezing: vehicle vs. muscimol: $t_{(28)} = 1.948$, $p = 0.1192$; Tone freezing: vehicle vs. muscimol: $t_{(28)} = 1.245$, $p = 0.3972$. $n_{vehicle} = 9$, $n_{muscimol} = 8$.

- Cued FC training - 7 days context + tone test (C): GzLMM two-way repeated measures ANOVA of treatment, training stage, and interaction: $\chi^2$ treatment $_{(1)} = 0.037$, $p = 0.8471$; $\chi^2$ stage $_{(1)} = 8.741$, $p = 0.0031$; $\chi^2$ interaction $_{(1)} = 8.674$, $p = 0.0032$. Sidak's multiple comparisons test: Post-shock freezing: vehicle vs. muscimol: $t_{(46)} = 0.193$, $p = 0.9769$; Tone freezing: vehicle vs. muscimol: $t_{(46)} = 2.216$, $p = 0.0624$. $n_{vehicle} = 13$, $n_{muscimol} = 13$.

- Cued FC training - 21 days context + tone test (D): GzLMM two-way repeated measures ANOVA of treatment, training stage, and interaction: $\chi^2$ treatment $_{(1)} = 3.723$, $p = 0.0537$; $\chi^2$ stage $_{(1)} = 0.023$, $p = 0.8802$; $\chi^2$ interaction $_{(1)} = 4.535$, $p = 0.0332$. Sidak's multiple comparisons test: Post-shock freezing: vehicle vs. muscimol: $t_{(26)} = 1.930$, $p = 0.1251$; Tone freezing: vehicle vs. muscimol: $t_{(26)} = 0.521$, $p = 0.8454$. $n_{vehicle} = 8$, $n_{muscimol} = 8$.

For cued FC trainings (B, C, D), stage factor refers to freezing between shock and tone, or during tone (post-shock or tone respectively).

## Suppl. Fig. 2

764 <u>Supplementary Figure 2. Inactivation of the LHb does not affect locomotion, exploration, or habituation</u>

765 <u>to an Open Field.</u>

766 **A)** Experiment diagram: bilateral intra LHb infusions of vehicle/muscimol were performed 30 minutes

767 before exposing the animals to an OF for 15 minutes. 48 hours later animals were exposed to the same

768 OF for a second time.

**B)** Distance traveled over time (60 s bins) during the first day (left) and the second day (right) in the OF. Thin lines represent individuals, thick lines represent the average. Right: total distance traveled. Distance traveled by both groups diminished the second day. No differences between groups were observed (two-way repeated measures ANOVA of treatment, day, and interaction: F treatment $_{(1, 10)}$ = 0.9320, p = 0.3571; F day $_{(1, 10)}$ = 15.92, p = 0.0026; F interaction $_{(1, 10)}$ = 0.2279, p = 0.6434. Sidak's multiple comparisons test: Day 1: vehicle vs. muscimol: t = 0.9534, p = 0.5798; Day 2: vehicle vs. muscimol: t = 0.1807, p = 0.9800).

**C)** Percentage of mobility time in the Open Field in each day. Left: percentage of mobility over time (60 s bins) during the first day (left) and the second day (right). Thin lines represent individuals, thick lines represent the average. Right: percentage of mobility time during whole OF assay. Mobility of both groups diminished the second day. No differences between groups were observed (two-way repeated measures ANOVA of treatment, day, and interaction: F treatment $_{(1, 10)}$ = 1.670, p = 0.2253; F day $_{(1, 10)}$ = 21.48, p = 0.0009; F interaction $_{(1, 10)}$ = 0.8727, p = 0.3722. Sidak's multiple comparisons test: First day: vehicle vs. muscimol: t = 1.512, p = 0.2708; Second day: vehicle vs. muscimol: t = 0.005307, p > 0.9999).

**D)** Heat map representation of spatial occupancy during OF. Left: average heat maps vehicle (top) and muscimol (bottom) groups on the first day (left), and the second day (right). The center zone is indicated by the orange square. Right: percentage of time in the center zone in both groups at each day. Time in the center did not change between days and was equivalent among groups (two-way repeated measures ANOVA of treatment, day, and interaction: F treatment $_{(1, 10)}$ = 0.01082, p = 0.9192; F day $_{(1, 10)}$ = 0.0003566, p = 0.9853; F interaction $_{(1, 10)}$ = 0.05192, p = 0.8244. Sidak's multiple comparisons test: Day 1: vehicle vs. muscimol: t = 0.1082, p = 0.9928; Day 2: vehicle vs. muscimol: t = 0.2416, p = 0.9645).

** p < 0.01, *** p < 0.001.

**Suppl. Fig. 3**

792    Supplementary Figure 3. Controls of specificity of muscimol inactivation of the LHb.

**A)** Photomicrographs of green beads fluorescence of a representative infusion on the LHb (left) or a double miss infusion (right). References: Hb: habenula; 3V: third ventricle.

**B)** Diagram and representative photomicrographs of controls of specificity of muscimol infusion. Infusions were aimed 1 mm dorsal (top right picture), ventral (bottom left picture) or lateral (bottom right picture). References: Hb: habenula; 3V: third ventricle.

**C)** Freezing during cued memory testing in miss-LHb group, dorsal, ventral, and lateral controls. Miss-LHb group was generated by selecting animals infused with muscimol in which both cannulae did not target the LHb from the same cohorts included in Figure 2C. Dorsal, ventral, and lateral groups were done as separate experiments. As in figure 2, infusions were done 30 minutes before training and test was performed 7 days later. Miss-LHb group showed higher freezing to the tone than muscimol-LHb group and equal freezing to the tone than the vehicle-LHb group. Muscimol infusion dorsal, ventral, or lateral to the LHb did not affect tone evoked freezing. In bar plots each dot represents a subject and bars represent mean + SEM. Muscimol-LHb and vehicle-LHb groups are taken from Figure 2C. ns p > 0.05, ** p < 0.001. Scale bars: 1 mm.

Statistics:

- GzLMM Two-way repeated measures ANOVA of treatment, test stage, and interaction for miss graph: $\chi^2$ treatment $_{(2)}$ = 2.5079, p = 0.2854; $\chi^2$ stage $_{(1)}$ = 85.1604, p < 0.0001; $\chi^2$ interaction $_{(2)}$ = 33.459, p < 0.0001. Sidak's multiple comparisons test: Pre-tone freezing: vehicle vs. muscimol: $t_{(40)}$ = 1.457, p = 0.6306; Pre-tone freezing: vehicle vs. miss: $t_{(40)}$ = 1.228, p = 0.7863; Pre-tone freezing: muscimol vs. miss: $t_{(40)}$ = 0.168, p = 1.0000; Tone freezing: vehicle vs. muscimol: $t_{(40)}$ = 5.191, p < 0.0001; Tone freezing: vehicle vs. miss: $t_{(40)}$ = 1.550, p = 0.5636; Tone freezing: muscimol vs. miss: $t_{(40)}$ = 3.559, p = 0.0058; $n_{vehicle}$ = 8, $n_{muscimol}$ = 9, $n_{miss}$ = 7.

- GzLMM Two-way repeated measures ANOVA of treatment, test stage, and interaction for dorsal graph: $\chi^2$ treatment $_{(1)}$ = 1.7632, p = 0.1842; $\chi^2$ stage $_{(1)}$ = 102.565, p < 0.0001; $\chi^2$ interaction $_{(1)}$ = 0.133, p = 0.7153. Sidak's multiple comparisons test: Pre-tone freezing: vehicle vs. muscimol: $t_{(26)}$ = 1.328, p = 0.3532; Tone freezing: vehicle vs. muscimol: $t_{(26)}$ = 1.916, p = 0.1283; $n_{vehicle}$ = 8, $n_{muscimol}$ = 8.

- GzLMM Two-way repeated measures ANOVA of treatment, test stage, and interaction for ventral graph: $\chi^2$ treatment $_{(1)}$ = 0.525, p = 0.4689; $\chi^2$ stage $_{(1)}$ = 165.9408, p < 0.0001; $\chi^2$ interaction $_{(1)}$ = 0.259, p = 0.6109. Sidak's multiple comparisons test: Pre-tone freezing: vehicle vs. muscimol: $t_{(32)}$ = 0.724, p = 0.7235; Tone freezing: vehicle vs. muscimol: $t_{(32)}$ = 0.360, p = 0.9224; $n_{vehicle}$ = 10, $n_{muscimol}$ = 9.

825    - GzLMM Two-way repeated measures ANOVA of treatment, test stage, and interaction for lateral

826    graph: $\chi^2$ treatment $_{(1)}$ = 0.326, p = 0.5680; $\chi^2$ stage $_{(1)}$ = 27.113, p < 0.0001; $\chi^2$ interaction $_{(1)}$ < 0.0001,

827    p = 0.9980. Sidak's multiple comparisons test: Pre-tone freezing: vehicle vs. muscimol: $t_{(26)}$ = 0.571,

828    p = 0.8176; Tone freezing: vehicle vs. muscimol: $t_{(26)}$ = 0.605, p = 0.7978; $n_{vehicle}$ = 8, $n_{muscimol}$ = 8.

# Suppl. Fig. 4

Supplementary Figure 4. Inactivation of the LHb does not permanently block FC learning.

**A)** Experiment diagram: bilateral intra LHb infusions of vehicle/muscimol were performed in the animals that went through the OF (Supplementary Figure 2). Seven days after the second exposure to the OF animals were trained in cued FC. Memory was tested 7 days later.

**B)** Cued FC training of the animals that were previously exposed to the OF. Left panel: freezing over time. Tones were presented at times indicated by the gray shaded areas. Shocks were presented at times indicated by the orange shaded areas. Right panel: mean freezing during tone and post-shock period. Freezing in both groups was equivalent. GzLMM Two-way repeated measures ANOVA of treatment, training stage, and interaction: $\chi^2$ treatment $_{(1)}$ = 0.685, p = 0.4077; $\chi^2$ stage $_{(1)}$ = 0.050, p = 0.8231; $\chi^2$ interaction $_{(1)}$ = 7.700, p = 0.0055. Sidak's multiple comparisons test: Post-shock freezing: vehicle vs. muscimol: $t_{(18)}$ = 0.828, p = 0.6620; Tone freezing: vehicle vs. muscimol: $t_{(18)}$ = 1.265, p = 0.3945. $n_{vehicle}$ = 6, $n_{muscimol}$ = 6.

**C)** Test of cued FC memory on the same animals. Left panel: freezing over time. Gray area indicates tone presentation. Right panels: average freezing for pre- and tone period. Freezing was not different between

vehicle and muscimol groups. GzLMM Two-way repeated measures ANOVA of treatment, test stage, and

interaction: $\chi^2$ treatment $_{(1)}$ = 0.0697, p = 0.7918; $\chi^2$ stage $_{(1)}$ = 15.254, p < 0.0001; $\chi^2$ interaction $_{(1)}$ = 0.103,

p = 0.7478. Sidak's multiple comparisons test: Pre-tone freezing: vehicle vs. muscimol: $t_{(18)}$ = 0.264, p =

0.9579; Tone freezing: vehicle vs. muscimol: $t_{(18)}$ = 0.190, p = 0.9780; $n_{vehicle}$ = 6, $n_{muscimol}$ = 6.

In freezing over time plots, line represents intersubjects' mean and shaded area represents SEM. In bar

plots each dot represents a subject and bars represent mean + SEM. ns p > 0.05.

**Suppl. Fig. 5**

853 Supplementary Figure 5. Optogenetic inhibition of LHb neurons by ArchT.

854 **A)** Example of membrane potential response to 600 ms current injection of an ArchT expressing LHb

855 neuron (-40, -10, 15, and 40 pA steps). Rebound spiking was observed after cessation of large

hyperpolarizing currents. Right graph: resting membrane potential of recorded neurons; n = 13 neurons from 3 rats.

**B)** Light induced photocurrent. ArchT was activated with 590 nm LED for 1 s (orange line) with a nominal power of 0.15 mW; n = 11 neurons from 3 rats.

**C)** Light induced action potential suppression in ArchT expressing LHb neurons. Light pulse (590 nm, orange line) was applied every second time neurons were depolarized with positive current injection. Left, representative recordings. Right, data summary. Light activation of ArchT significantly decreases the number of action potentials evoked by current steps in LHb neurons (paired t test, $t_{(9)}$ = 6.25, p < 0.001); n = 9 neurons from 3 rats.

**D)** Membrane potential response to a 30 s light pulse (orange line) in LHb neurons. Rebound spiking was observed after termination of light pulse, a representative trace is expanded in the middle traces. This behavior confirms the reported rebound responses of the LHb neurons following long periods of optogenetic inhibition reported previously (Yang et al., 2018). Right: quantification of the stationary light induced ΔVm during the last 5 s of the 30 s pulse; n = 10 neurons from 3 rats.

Methods details for this experiment: Surgery was similar to the optogenetic manipulation experiments with the following differences: 3 weeks old rats were employed, the viral vector injection sites were adjusted according to Bregma-Lambda distance (Bλ) of each rat (AP = -2.9 × Bλ ÷ 8.5 mm, ML = ± 0.7 × Bλ ÷ 8.5 mm from Bregma and DV -4.3 mm from dura mater), no fiber optic implants, nor screws or acrylic was used, and the skin was stitched at the end of surgery. Recordings were obtained from ArchT expressing LHb neurons in acute brain slices prepared 3 weeks after infection. General methods for slice preparation and recordings were replicated from previous publications of the group (Stefanelli et al., 2016).

## Suppl. Fig. 6

<u>Supplementary Figure 6. Freezing during training for experimental groups included in Figures 5 and 6.</u>

**A-B)** Left: temporal course of freezing during training in the ArchT (A) and oChIEF (B) stimulation

experiments. Right: average freezing on the same experiments. In freezing over time plots vertical orange

shaded areas indicate shock presentation, gray shaded areas indicate tone presentation, line plots

represent intersubjects' mean and shaded area over line plots represents +SEM. In either condition no

significant differences were observed between control and optogenetic stimulated groups.

Statistics:

- For ArchT experiment: GzLMM two-way repeated measures ANOVA of optogenetic inhibition,

training stage, and interaction: $\chi^2$ inhibition $_{(1)}$ = 0.002, p = 0.9631; $\chi^2$ stage $_{(1)}$ = 3.173, p = 0.0749; $\chi^2$

interaction $_{(1)}$ = 5.000, p = 0.0254. Sidak's multiple comparisons test: Post-shock freezing: GFP vs.

ArchT: $t_{(32)}$ = 0.046, p = 0.9987; Tone freezing: GFP vs. ArchT: $t_{(32)}$ = 2.098, p = 0.0859. $n_{GFP}$ = 9, $n_{ArchT}$

= 10.

- For oChIEF experiment: GzLMM two-way repeated measures ANOVA of optogenetic disruption,

training stage, and interaction: $\chi^2$ disruption $_{(1)}$ = 0.007, p = 0.9358; $\chi^2$ stage $_{(1)}$ = 0.018, p = 0.8946;

$\chi^2$ interaction $_{(1)}$ = 0.0650, p = 0.7988. Sidak's multiple comparisons test: Post-shock freezing: Control

vs. oChIEF: $t_{(20)}$ = 0.081, p = 0.9960; Tone freezing: Control vs. oChIEF: $t_{(20)}$ = 0.113, p = 0.9921. $n_{Control}$

= 7, $n_{oChIEF}$ = 6.

Stage factor refers to freezing between shock and tone, or during tone (post-shock or tone respectively).

## Suppl. Fig. 7

<u>Supplementary Figure 7. Modelling of temperature increase induced by optogenetic stimulation.</u>

Simulation was implemented using the toolbox developed by Stujenske, Spellman and Gordon (2015)

using parameters similar to the one used in our real experiments: optic fiber: radius = 0.2 mm, NA = 0.45,

wavelength: 532 nm for ArchT and 447 nm for oChIEF, laser intensity 10 mW. Additional parameters were

taken from Stujenske et al. (2015) (montecarlolight_function: modelnumber = 1; HeatDiffusionLight:

U_STEP = 0.03 mm, T_SAVE = 500, R_AVG = 0.25 mm).

**A**) Heat plot of predicted temperature changes at different distances from the optic fiber tip induced by

optogenetic stimulation during FC training. Left: temperature changes induced by constant light

stimulation required for inhibition with ArchT. Right: temperature changes induced by pulsed light

stimulation used for oChIEF activation.

**B**) Depth profile of temperature change in the steady state (calculated as the mean in the 250-300 s

interval). Constant light stimulation with 532 nm light would induce a significant temperature increase

that would extend over more than 1 mm and would induce non-controllable out of target changes in

neuronal activity. In contrast, pulsed activation of oChIEF would not induce physiologically relevant

temperature changes at any depth (Stujenske et al., 2015).

**C**) Temporal profile of temperature change at 0.25 and 1 mm from the fiber tip for both stimulation

protocols.