Multi-Drug-Resistance in Drug-Sensitive and Drug-Resistant Ovarian Cancer Cells Responds Differently to Cell Culture Dimensionality

Vasilij Koshkin,[†] Mariana Bleker de Oliveira,[†] Chun Peng,[†] Laurie E. Ailles,[‡] Geoffrey Liu,[§] Allan Covens,[‖] and Sergey N. Krylov*,[†]

[†]Centre for Research on Biomolecular Interactions, York University, Toronto, Ontario M3J 1P3, Canada

[‡]Princess Margaret Cancer Centre and Department of Medical Biophysics, University of Toronto, Toronto M5G 1L7, Canada Department of Medical Biophysics, University of Toronto, Toronto, Ontario N5G 1L7, Canada

[§]Department of Medicine, Medical Oncology, Princess Margaret Cancer Centre, Toronto, Ontario M5G 2M9, Canada

[‖]Sunnybrook Odette Cancer Centre, Toronto, Ontario M4N 3M5, Canada

*Corresponding author's e-mail: skrylov@yorku.ca

Abstract

Does cell clustering in ovarian cancer influence intrinsic and acquired multi-drug resistance (MDR) differently? To address this question, we studied cultured monolayers (representing individual cells) and cultured spheroids (representing clusters) formed by drug-naïve (intrinsic MDR) and drug-exposed (acquired MDR) lines of ovarian cancer A2780 cells by cytometry of reaction rate constant (CRRC). MDR efflux was characterized by accurate and robust "cell number *vs*. MDR efflux rate constant (k_{MDR})" histograms. Both drug-naïve and drug-exposed monolayer cells presented unimodal histograms; the histogram of drug-exposed cells was shifted towards higher k_{MDR} value suggesting greater MDR activity. Spheroids of drug-naïve cells presented a bimodal histogram indicating the presence of two subpopulations with different MDR activity. In contrast, spheroids of drug-exposed cells presented a unimodal histogram qualitatively similar to that of the monolayers of drug-exposed cells but with a moderate shift towards greater MDR activity.

Implications: The observed greater effect of cell clustering on intrinsic than on acquired MDR can help guide the development of new therapeutic strategies targeting clusters of circulating tumor cells in OC and other types of cancer.

Introduction

Ovarian cancer (OC) is the seventh most common cancer in women, and chemotherapy is its frontline treatment after surgery (1). However, effect of chemotherapeutic treatment is limited by chemoresistance. Approximately 30% of patients have pre-intrinsic chemoresistance and others eventually acquire chemoresistance during continuing exposure to chemotherapeutic agents (2). Because chemoresistance is among the main reasons preventing progress in OC treatment, there has been intensive research in this field. A promising new approach in elucidating the mechanisms of cancer chemoresistance is studying circulating tumor cells (CTCs). CTCs exist as individual cells or multicellular clusters that detach from the primary tumor and circulate in the bloodstream and give rise to systemic metastases which are seen in some OC patients (3). Chemoresistance of CTC clusters is greater than that of individual CTCs (4). Understanding the mechanisms of chemoresistance in CTCs in OC is important as there is a correlation between the number of CTC clusters and clinical features of OC (5).

There are several cellular processes that contribute to both intrinsic and acquired chemoresistance (6). One of such processes is active extrusion of drugs from cells by ATP-binding cassette transporters (membrane proteins) (7). This process has low drug specificity and is termed multi-drug resistance (MDR). Here, we use the term of MDR solely to describe the catalytic process of drug transport across the membrane (from inside to outside of the cell). Activity of MDR in OC tumor cells has been shown to correlate with clinical chemoresistance, and MDR is its likely driver (8).

The presumed roles of MDR transport and CTC clusters in the development of chemoresistance logically lead to a question: how does MDR transport activity of clusters differ from that of single cells in cases of intrinsic and acquired resistance? Addressing this question requires a cytometry technique capable of accurately measuring MDR activity and applicable to both single cells and intact clusters. Classical cytometry technics cannot be used for accurate measurements of MDR activity, making the above-posed question difficult to approach experimentally. In contrast, cytometry of reaction rate constant (CRRC) can be used for this purpose. In general, CRRC utilizes time-lapse fluorescence microscopy to measure a rate constant of a catalytic reaction in individual cells and, thus, facilitates accurate size determination for subpopulations of cells with distinct efficiencies of this reaction (9). Time-lapse fluorescence images are used to build kinetic traces of substrate conversion into a product. A reaction rate constant is then found for every cell using a known mechanism of the reaction. Finally, a CRRC histogram that plots the frequencies (*i.e.* number of cells) for varying ranges of the rate constant value is used to accurately measure sizes of cell subpopulations with distinct reaction activities (9).

When applied to MDR transport, CRRC is used to record kinetics of fluorescent substrate extrusion from cells. The extrusion process is governed by the Michalis-Menten mechanism, which is characterized by two parameters: the maximum velocity, V_{max}, and the Michaelis constant, K_M. A ratio between these parameters is a first order rate constant of MDR transport, $k_{MDR} = V_{max}/K_M$, which can be easily determined from time dependence of intracellular fluorescence intensity of MDR substrate. CRRC histograms that plot number of cells *vs*. k_{MDR} ranges are robust towards variations in substrate concentration and observation time (9); therefore, such histograms facilitate accurate determining the sizes of cell subpopulations with different activities of MDR (9).

The most straightforward application of CRRC is to 2D models, such as cells cultured as monolayers or cells obtained by disintegration of cell clusters (e.g. spheroids or tissue samples) and allowed to settle on the surface (10). However, if CRRC is

Figure 1. CRRC histograms of frequencies of cell subpopulations by MDR transport first order rate constant (k_{MDR}, measured in s^{-1}) in monolayer-grown cells (grey lines) and spheroid-grown cells (black lines) for drug-naïve A2780S cell line (A) and its derivative drug-exposed to cisplatin A2780CP cell line (B). Traces in panel A have been adopted with permission from Figure 4 in Analytical Chemistry 2020; 92, 9348–9355; Copyright 2020 American Chemical Society.

based on confocal microscopy, it can also be applied to intact cell clusters (10). Thus, CRRC is uniquely capable of accurately measuring MDR activity in both 2D and 3D models, making it suitable for addressing our question of how MDR activity of single cells differs from that of aggregated cells in (i) drug-naïve and (ii) drug-exposed tumor cells.

To address this question in the context of OC, we choose two sublines of A2780 OC cells, recently used to mimic OC circulating cells (11). The A2780S cell subline is derived from a patient who was not exposed to chemotherapy, and, thus, this drug-naïve cell line represents intrinsic chemoresistance (12). The A2780CP cell subline is derived from the A2780S subline and has been cultured in the presence of cisplatin to develop drug-resistance. Therefore, drug-exposed A2780CP cells represent OC tumor cells that were exposed to chemotherapy and developed acquired chemoresistance. Importantly, both cell lines can be grown as monolayers or as multicellular spheroids (13). Therefore, we can view cultured monolayers of A2780S and A2780CP cells as models of circulating single cells, while their cultured spheroids can be considered as models of CTC clusters. Then, comparison of CRRC histograms of a cell monolayer with that of cultured spheroids can answer our question.

Materials and Methods

Detailed description of experimental techniques of CRRC for cell monolayers and cells in intact spheroids can be found elsewhere (10). Briefly, monolayers and spheroids were cultured in DMEM medium supplemented with bovine serum and antibiotics under standard cell culture conditions. The culturing of small multicellular spheroids was based on the liquid overlay approach (14), adapted for ovarian cells (15). Cells were placed into wells coated with agarose to prevent adhesion and allowed to form spheroids for 2–3 days. For time-lapse imaging, spheroids were placed onto coverslips and allowed to settle and attach to the surface for 5 h (16). Imaging of MDR efflux was performed with a FV300 confocal cell imager (Olympus) in the time-lapse mode

with single and multiple optical sections taken for monolayers and spheroids, respectively. Cells were loaded with a fluorescent MDR substrate (fluorescein) and allowed to extrude it. Kinetics of substrate extrusion was monitored by measuring intracellular fluorescence intensity over time to determine k_{MDR} for individual cells, and, importantly, only outer spheroidal cells were taken into consideration. Finally, kinetic cytometry histograms were plotted to compare MDR activity (k_{MDR}) of monolayers and spheroids in A2780S and A2780CP cell lines (Figure 1).

Results and Discussion

CRRC results of the drug-naïve A2780S cell line are shown in Figure 1A; they have been adopted from our recently published paper (10). The monolayer histogram (grey line) was found to be unimodal, suggesting a single population of cells. The spheroid histogram (black line) revealed a bimodal distribution, suggesting two cell subpopulations: the first one is larger and has the same peak k_{MDR} value as the monolayer cells, while the second one is smaller and has a peak k_{MDR} value which is almost 3 times greater. This latter subpopulation has a greater MDR capacity and its appearance is caused by cell-cell interactions in the 3D spheroids (17). The presence of a drug-resistant subpopulation (presumably containing tumor initiating cells) in the spheroids is consistent with a notion that CTC clusters have a greater drug-resistance capacity than single CTCs. The unimodal right-skewed histogram of monolayer A2780S cells is similar to that reported earlier (9) Its shape is consistent with the often reported asymmetric expression of MDR transporters, when the majority of cells with a basal (low) level of transporter expression form the main peak and a much smaller subpopulation of cells with the elevated level of transporter expression forms the distribution tail towards higher k_{MDR} values (18). When A2780S cells are cultured as spheroids, the tail becomes a distinct peak, which corresponds to a distinct subpopulation of cells with its own elevated peak k_{MDR} value: the histogram becomes bimodal. This bimodality is not associated with cell position in the 3D spheroids as only the outer spheroidal cells were analysed. It is remarkable that the activation of MDR transport in spheroidal A2780S cells is not distributed equally across all spheroidal cells. Instead, the activation proceeds through enlarging the size of the subpopulation of cells with a greater MDR-transport activity. This characteristic of MDR modulation agrees well with the notion that the size and activity of the drug-resistant subpopulation determine the overall resistance of a heterogeneous cell population (19).

CRRC results of the cisplatin-exposed (drug-resistant) A2780CP cell line are depicted in Figure 1B. Monolayer cells (grey line) showed a unimodal distribution with the median (peak) k_{MRD} value exceeding that of monolayer A2780S cells (grey line in Figure 1A) approximately by a factor of 2. When cultured as spheroids, A2780CP cells also showed a unimodal distribution (black line), and the histogram of the spheroidal A2780CP cells was moderately shifted to the right with respect to that of the monolayer A2780CP cells. In addition, the peak maximum of the spheroidal A2780CP cells was at the same k_{MDR} position as the peak maximum of the drug-resistant subpopulation in the spheroidal A2780S cells (right-hand-side peak in the black line in Figure 1A). Thus, the drug-resistant subpopulation dominates even in monolayers formed by these cells and predictably dominates in spheroids resulting in a unimodal k_{MDR} distribution in the spheroid culture. Peak maximum value in the spheroid culture slightly exceeds that in monolayer (by a factor of 1.2). Lower kurtosis (indicator of distribution peakedness/flatness, −0.95 *vs* 1. 87) indicates that MDR distribution in A2780CP spheroids is more heterogeneous than in monolayers. Greater heterogeneity can be associated with the larger fraction of cells with elevated k_{MDR} able to survive chemotherapy and initiate tumor relapse.

The potential clinical implications of our findings are dual. First, considering the role of clustering of OC cells in intrinsic and

acquired chemoresistance, our results suggest that intrinsic resistance can be more sensitive to the cell clustering. In particular, elevated intrinsic chemoresistance of multicellular aggregates can be associated with the chemosensitizing effect of debulking surgery (20). Second, activation of MDR transport within spheroids was ascribed to activation of the HIF pathway caused either by hypoxia inside the spheroids or by cytotoxic agents (21). However, we found MDR-transport activation not only inside the spheroids but also on their surface; moreover, this activation is observed in both drug-naïve (intrinsic MDR) and drug-exposed (acquired MDR) cells. These observations strongly suggest that there are other mechanisms of MDR-transport activation in addition to the HIF-hypoxia pathway.

Overall, our data show that the extent of MDR-transport activation in OC cell clusters strongly depends on the previous chemotherapeutic history of spheroid-forming A2780 cells. This fact, if confirmed on primary ovarian tumor cells, will help clinicians to optimize OC treatment, since therapeutic approaches might have different outcomes for drug-naïve and drug-exposed tumors.

References

1. Reid BM, Permuth JB, Sellers TA. Epidemiology of ovarian cancer: a review. Cancer Biol Med 2017;14:9–32.
2. Cornelison R, Llaneza DC, Landen CN, Emerging therapeutics to overcome chemoresistance in epithelial ovarian cancer: a mini-review. Int J Mol Sci 2017;18:E2171.
3. Patel TS, Shah C,. Shah MC, and Shah MJ. Axillary node metastasis from primary ovarian carcinoma. J Cytol 2014; 31: 202–204.
4. Guiliano M, Shaikh A, Lo H, Arpino G, Placido S, Zhang X, et al. Perspective on circulating tumor cell clusters: why it rakes a village to metastasize. Mol Cancer Res 2018;78:845–852.
5. Lee M, Kim E, Cho Y, Kim A, Chung H, Park N et al. Predictive value of circulating tumor cells (CTCs) captured by microfluidic device in patients with epithelial ovarian cancer. Gynecol Oncol 2017;145:361–365.
6. Norouzi-Barough L, Sarookhani MR, Sharifi M, Moghbelinejad S, Jangjoo S, Salehi R. Molecular mechanisms of drug resistance in ovarian cancer. J Cell Physiol 2018; 233:4546–4562.
7. Srivastava R, Srivastava A, Chochung Y. Multidrug resistance in cancer. Int J Oncol 1996;9:879–884.
8. Zhang Y, Sriraman S, Kenny H, Luther E, Torchilin V, Lengyel E. Reversal of chemoresistance in ovarian cancer by co-delivery of a P-glycoprotein inhibitor and paclitaxel in a liposomal platform. Mol Cancer Ther 2016;15:2282–2293.
9. Koshkin V, Kochmann S, Sorupanathan A, Peng C, Ailles LE, Liu G, Krylov, S. N. Cytometry of reaction rate constant: measuring reaction rate constant in individual cells to facilitate robust and accurate analysis of cell-population heterogeneity Anal Chem 2019;91:4186–4194.
10. Koshkin V, Bleker de Oliveira M, Peng C, Ailles L, Liu G, Covens A, Krylov SN. Spheroid-based approach to assess the tissue relevance of analysis of dispersed-settled tissue cells by cytometry of the reaction rate constant. Anal Chem 2020;92: 9348–9355.
11. Li N, Zuo H, Chen L, Liu H, Zhou J, Yai Y et al. Circulating tumor cell detection in epithelial ovarian cancerusing dual-component antibodies targeting EpCAM and Frα. Cancer Manag Res 2019;11:10939–10948.
12. Mutch DG, Herzog TJ, Chen CA, Collins JL. The effects of cyclosporin A on the lysis of ovarian cancer cells by cisplatin or adriamycin. Gynecol Oncol 1992;47:28–33.
13. Moraya A, Ali J, Samadder P, Liang L, Morrison L, Werbowetski-Ogilvie T et al. Novel glycolipid agents for killing cisplatin-resistant human epithelial ovarian cancer cells. J Exp Clin Cancer Res 2017;36:67.
14. Friedrich J, Seidel C, Ebner R, Kunz-Schughart LA. Spheroid-based drug screen: considerations and practical approach. Nat Protoc 2009;4:309–324.
15. Frankel A, Buckman R, Kerbel RS. Abrogation of taxol-induced G2-M arrest and apoptosis in human ovarian cancer cells grown as multicellular tumor spheroids. Cancer Res 1997;57: 2388–2393.
16. Acker H, Carlsson J, Holtermann G, Nederman T, Nylén T, Influence of glucose and buffer capacity in the culture medium on growth and pH in spheroids of human thyroid carcinoma and human glioma origin. Cancer Res 1987;47:3504–3508.
17. Ham S, Joshi R, Thakuri P, Tavana H. Liquid-based three-dimensional tumor models for cancer research and drug discovery. Exp Biol Med 2016;241:939–954.
18. Szakács G, Annereau P, Lababidi S, Shankawaram U, Arciello A, Bussey K et al., M. Predicting drug sensitivity and resistance: profiling ABC transporter genes in cancer cells. Cancer Cell 2004;6:129–137.
19. Howard G, Johnson K, Ayala A, Yankeelov T, Brock A. A multi-state model of chemoresistance to characterize phenotypic dynamics in breast cancer. Sci Rep 2018;8:12058.
20. Eisenhauer EL, Abu-Rustum NR, Sonoda Y, Aghajanian C , Barakat RR, Chi DS. The effect of maximal surgical cytoreduction on sensitivity to platinum-taxane chemotherapy and subsequent survival in patients with advanced ovarian cancer. Gynecol Oncol 2008; 108: 276–281.
21. He M, Wu H, Jiang Q, Liu Y, Han Li, Yan Y et al. Hypoxia-inducible factor-2α directly promotes BCRP expression and mediates the resistance of ovarian cancer stem cells to adriamycin. Mol Oncol 2019;13: 403–421.