The lncRNA *APOLO* interacts with the transcription factor WRKY42 to trigger root hair cell expansion in

response to cold

Michaël Moison[*1], Javier Martínez Pacheco[*2], Leandro Lucero[*1], Camille Fonouni-Farde[1], Johan

Rodríguez-Melo[3], Aurélie Christ[4], Jérémie Bazin[4]; Moussa Benhamed[4]; Fernando Ibañez[3], Martin Crespi[4],

José M. Estevez[†,2,5] and Federico Ariel[†1]

[1] Instituto de Agrobiotecnología del Litoral, Universidad Nacional del Litoral, CONICET, FBCB, Centro

Científico Tecnológico CONICET Santa Fe, Colectora Ruta Nacional No 168 km. 0, Paraje El Pozo, Santa Fe

3000, Argentina.

[2] Fundación Instituto Leloir and IIBBA-CONICET, Av. Patricias Argentinas 435, Buenos Aires CP C1405BWE,

Argentina.

[3] Instituto de Investigaciones Agrobiotecnológicas, CONICET, Universidad Nacional de Río Cuarto, Río

Cuarto 5800, Argentina.

[4] Institute of Plant Sciences Paris-Saclay (IPS2), CNRS, INRA, University Paris-Saclay and University of Paris

Bâtiment 630, 91192 Gif sur Yvette, France.

[5] Centro de Biotecnología Vegetal (CBV), Facultad de Ciencias de la Vida (FCsV), Universidad Andrés Bello,

Santiago, Chile and Millennium Institute for Integrative Biology (iBio), Santiago, Chile.

[*]These authors contributed equally to this work

[†] Correspondence to: Federico Ariel (fariel@santafe-conicet.gov.ar) and José M. Estevez

(jestevez@leloir.org.ar)

Text Word count (without Abstract and References)**:** 4,984

ABSTRACT

Plant long noncoding RNAs (lncRNAs) have emerged as important regulators of chromatin dynamics,

impacting on transcriptional programs leading to different developmental outputs. The lncRNA *AUXIN*

REGULATED PROMOTER LOOP (*APOLO*) directly recognizes multiple independent loci across the

Arabidopsis genome and modulates their three-dimensional chromatin conformation, leading to

transcriptional shifts. Here, we show that *APOLO* recognizes the locus encoding the root hair (RH) master

regulator ROOT HAIR DEFECTIVE 6 (RHD6) and controls *RHD6* transcriptional activity leading to cold-

enhanced RH elongation, in association with the Polycomb Repressive Complexes (PRC) 1 and 2.

Additionally, we demonstrate that *APOLO* interacts with the transcription factor WRKY42 and modulates

its binding to the *RHD6* promoter. WRKY42 is required for the activation of *RHD6* by low temperatures

and *WRKY42* deregulation impairs cold-induced RH expansion. Collectively, our results indicate that a

novel ribonucleoprotein complex involving *APOLO* and WRKY42 forms a regulatory hub which activates

RHD6 by shaping its epigenetic environment and integrates different signals governing RH growth and

development.

Words: 160

Key words: root hairs; cell expansion; long noncoding RNAs; *APOLO*; transcription factors; RHD6;

WRKY42; cold

INTRODUCTION

Root hairs (RHs) are single cell projections developed from specialized epidermal trichoblast cells able to increase their size several hundred times in a polar manner to reach and promote the uptake of water-soluble nutrients, interact with microorganisms and anchor the plant to the soil. The specification of epidermal cells into RHs involves a cell fate determination process whose underlying mechanisms are only partially understood. In *Arabidopsis thaliana*, RH cell fate is controlled by a developmental program involving a complex of transcription factors (TFs) promoting the expression of the homeodomain protein GLABRA 2 (GL2)(1–4). GL2 blocks RH development by inhibiting the transcription of the master regulator *ROOT HAIR DEFECTIVE 6* (*RHD6*)(5). In the trichoblast cells that differentiate into RHs, a second TF complex suppresses *GL2* expression (3), forcing the cells to enter the RH cell fate program via the concomitant activation of RHD6 along with downstream TFs (6,7). Briefly, RHD6 together with its homolog RHD6-LIKE 1 (RSL1) induce the expression of other TFs from the bHLH family, including RSL2 and RSL4, ultimately triggering the differentiation of the RHs and their subsequent polarized tip-growth (8–10). In addition, it was proposed that RSL4 controls the expression of a small subset of nearly 125 genes (9, 11–13), including several cell wall extensins (*EXTs*) (14, 15) sufficient to promote RH growth (16).

RH expansion is regulated both by cell-intrinsic factors (e.g. endogenous phytohormones such as auxin) and external environmental signals (e.g. phosphate (Pi) availability in the soil) (17, 18). Pi starvation is one of the key environmental factors promoting rapid RH growth (9, 12, 13). In *Arabidopsis*, it triggers *RSL4* expression via an enhanced auxin production, activating downstream effector genes mediating cell growth (9, 12, 17-19). Accordingly, several auxin-related TFs have been implicated in Pi-starvation signaling in roots, including WRKY proteins that control the expression of the Pi transporter families Pi-permease PHO1 and PHOSPHATE TRANSPORTER (PHT) (20–23). Under Pi-sufficient conditions, WRKY6 and WRKY42 bind to W-boxes of the *PHO1* promoter and suppress its expression. During Pi starvation, WRKY42 is degraded by the 26S proteasome pathway, resulting in the activation of *PHO1* transcription (21, 23). In addition, WRKY42 functions as a positive regulator of *PHT1;1*, by binding to its promoter under Pi-sufficient condition (23). Overall, WRKY42 is part of the components activating root early-ly-responses to Pi starvation, although its role in controlling RH growth remains unexplored.

In recent years, plant long noncoding RNAs (lncRNAs) have emerged as important regulators of gene expression, and several among them have been functionally linked to Pi homeostasis. For instance, the lncRNA *INDUCED BY PHOSPHATE STARVATION 1* (*IPS1*) can sequester the Pi starvation-induced microRNA miR-399, attenuating miR-399-mediated repression of *PHO2*, a gene involved in Pi uptake (24). In

addition, the *cis*-natural antisense (*cis*-NAT) transcript *PHO1;2*, induced under Pi deficiency, was shown

to promote the translation of the *PHO1;2* mRNA involved in Pi loading into the xylem. The expression of

this *cis*-NAT is associated with the transport of the sense–antisense RNA pair toward the polysomes (25).

More recently, it was shown that the lncRNA *AUXIN REGULATED PROMOTER LOOP* (*APOLO*) recognizes

multiple spatially independent genes by sequence complementarity and DNA-RNA duplex formation,

known as R-loops. Upon recognition, *APOLO* shapes the three-dimensional (3D) conformation of its tar-

get regions by decoying the Polycomb Repressive Complex 1 (PRC1) component LIKE HETEROCHROMA-

TIN PROTEIN 1 (LHP1), thereby regulating their transcription (26, 27).

Here, we show that the lncRNA *APOLO* directly regulates a subset of genes involved in RH devel-

opment, including the master regulator of RH initiation *RHD6*. *APOLO* activates *RHD6* transcription by

modulating the formation of a local chromatin loop encompassing its promoter region, an epigenetic

regulatory mechanism likely involving PRC1 and PRC2 components. Furthermore, we found that *APOLO*

interacts with the TF WRKY42, forming a new hub that regulates *RHD6* to induce RH growth in response

to low temperatures.

RESULTS

***APOLO* regulates root hair cell elongation in response to low temperatures**

The lncRNA *APOLO* was previously reported to recognize a subset of independent loci enriched in

categories related to cell wall composition and organization (27). A closer look at *APOLO bona fide* targets

allowed us to identify seventeen genes involved in RH growth and expansion (**Supplementary Table 1**), a

process dependent on cell wall remodeling molecules, including EXTs and EXT-related proteins (14, 15,

28–30). Interestingly, according to single-cell RNA-seq datasets (31) *APOLO* transcripts are enriched in RH

cells (**Supplementary Figure 1**). Notably, sixteen *APOLO* direct targets were upregulated and one

downregulated (*EXT18*) upon *APOLO* over-expression (**Supplementary Table 2**). Furthermore, 52 addi-

tional RH-related genes were upregulated in *35S:APOLO* seedlings, albeit not identified as *APOLO* direct

targets (**Supplementary Table 2**) (27). Among them, the RH central TF genes *RHD6* (as a direct target),

RSL2 and *RSL4* (as indirectly regulated) were induced upon *APOLO* over-expression.

It was reported that the *APOLO* locus is targeted by the RNA-polymerase Pol V and silenced by

RNA-directed DNA Methylation (RdDM, (26)). A search in a small RNA-Seq performed in WT roots sub-

jected to different temperature treatments (32) revealed that RdDM-related 24nt siRNA accumulation

over the *APOLO* locus is less abundant at low temperatures (15°C; **Figure 1A**), suggesting that *APOLO*

transcription is regulated by cold. Accordingly, we found that *APOLO* transcriptional accumulation in-

creases in roots after 24h at 10°C (**Figure 1B**). An analysis of the promoter activity of the 5.2kb region

upstream *APOLO* (27) directing the expression of a *GFP* reporter gene, additionally revealed a higher

transcriptional activity at low temperatures in the RHs (**Figure 1C**). Strikingly, we observed that two RNAi-

APOLO and two *35S:APOLO* independent lines (26, 27) exhibit a basal increase of RH length at 22°C, and

uncovered a strong induction of RH elongation in WT and RNAi-*APOLO* at 10°C, in contrast to *35S:APOLO*

lines (**Figure 1D**). Accordingly, *RHD6* is induced in response to cold in WT roots and RNAi-*APOLO* roots

display higher *RHD6* basal levels than the WT (**Figure 1E**). Collectively, our findings suggest that *APOLO*

participates in the induction of cold-mediated RH elongation and that a deregulation of *APOLO* transcript

levels can impact RH growth.

Previous studies pointed out a key role of RHD6 (together with RSL1) in RH development, which

is mediated by RSL4 and RSL2 as downstream regulators of RH cell elongation (6, 7). Considering that

RHD6, *RSL2* and *RSL4* transcript levels were upregulated in *35S:APOLO* seedlings (**Supplementary Table**

2; (27)), we assessed if these TFs were also controlling the promotion of RH growth by low temperatures.

To this end, we tested how *rhd6/rsl1/rsl4* and *rsl2*, *rsl4* and double mutant plants *rsl2/rsl4* respond to

low temperatures in comparison with control conditions. The *rsl2* mutant was highly responsive to low

temperatures in a similar manner to WT while *rsl4* was impaired in the response to cold. The double

mutant *rsl2/rsl4* and the triple mutant *rhd6/rsl1/rsl4* did not develop RHs in either of the two conditions

(**Supplementary Figure 2A**). In addition, constitutive expression of *RSL4* (*35S:RSL4*) as well as its expres-

sion under the control of the RH specific *EXPANSIN7* promoter (*EXP7p:RSL4*) boosted basal RH growth

without further enhancement in response to cold (**Supplementary Figure 2B**). These results demonstrate

that RSL4 is largely required for RHD6-dependent activation of RH growth at low temperatures, and RSL2

might participate to a lower extent.

***APOLO* directly modulates the three-dimensional chromatin conformation of the root hair specific lo-**

cus *RHD6*

Among *APOLO* targets involved in RH development, we found the master regulator of RH initia-

tion *RHD6* (6, 7). The epigenetic profile of the *RHD6* locus corresponds to typical *APOLO* targets (**Figure**

2A, (27)), including H3K27me3 deposition (track 1), LHP1 recognition (track 2, chromatin

immunoprecipitation (ChIP)-Seq, (33)), and *APOLO* binding regions (tracks 3 to 5, chromatin isolation by

RNA purification (ChIRP)-Seq, (27)). A GAAGAA box, shown to be important for *APOLO* target recognition

(27) is located in the *RHD6* locus and coincides with *APOLO* binding site. In addition, a small peak indi-

cates the presence of an R-loop coinciding with *APOLO* recognition sites over *RHD6* (tracks 6 to 9, DNA-

RNA hybrid immunoprecipitation (DRIP)-Seq (34)). The possibility that *RHD6* recognition by *APOLO* occurs

specifically in RHs may explain the dilution of the signal coming from entire seedlings (34).

Remarkably, *APOLO* recognition and R-loop formation are also detectable over *RHD6* neighbor

gene, located 3.2 kb upstream *RHD6* transcription start site (**Figure 2A**). According to *Dpn*II Hi-C datasets

from Arabidopsis seedlings (35), a chromatin loop encompassing the intergenic region upstream *RHD6*

was detected (**Figure 2B**), exhibiting *APOLO* binding at both sides of the loop base (**Figure 2A**, ChIRP-Seq).

By performing a ChIRP-qPCR with two independent sets of biotinylated probes to purify *APOLO* (ODD and

EVEN; (27)) and one additional set used as a negative control (LacZ), we confirmed that *APOLO* RNA–

RHD6 DNA interaction occurs in wild-type (WT) and is lost in *APOLO* knockdown (RNAi) seedlings ((26);

Figure 2C). In addition, the quantification of relative *RHD6* loop formation in RNAi-*APOLO* and *APOLO*

over-expressing (*35S:APOLO*; (27)) seedlings, revealed impaired loop formation in both lines (**Figure 2D**),

hinting at a stoichiometric requirement of *APOLO* for *RHD6* chromatin loop formation. Accordingly, an

RNA-Seq dataset of *35S:APOLO* seedlings *vs.* WT (27) indicates that *RHD6* transcript levels are increased

upon *APOLO* over-expression (**Figure 2E, Supplementary Table 1**), suggesting that the chromatin loop

including *RHD6* promoter region precludes transcription. Altogether, our results indicate that *APOLO*

lncRNA directly regulates *RHD6* transcriptional activity by fine-tuning local chromatin 3D conformation.

It was previously reported that PRC2 actively participates in the regulation of RH growth (36) and

that the *RHD6* locus exhibits H3K27me3 deposition and LHP1 recognition (**Figure 2A;** (33)). Thus, we de-

cided to explore the role of PRC1 and 2 in *APOLO*-mediated *RHD6* activation at low temperatures. At

22°C, *RHD6* suffers a reduction of H3K27me3 in the PRC2 mutant *curly leaf* (*clf*), in contrast to the PRC1

mutant *lhp1* (**Supplementary Figure 3A;** (33)). Interestingly, we observed that H3K27me3 deposition and

LHP1 binding diminish in WT roots treated for 24h at 10°C compared to 22°C (**Supplementary Figure 3B**),

in agreement with the induction of *RHD6* in response to cold (**Figure 2E**). Moreover, *lhp1* and *clf* mutants

exhibit a basal decrease of RH length together with a slight decrease of cold-induced RH elongation in

lhp1, and a strong decrease of cold-induced RH elongation in *clf* (**Supplementary Figure 3D**). Consistently,

RHD6 transcriptional activation by cold is abolished in the *clf* mutant (**Supplementary Figure 3C**), hinting

at an important role of chromatin rearrangement for *RHD6* activation in response to cold.

APOLO* interacts with the transcription factor WRKY42 to coordinate the activation of *RHD6

With the aim of identifying novel actors involved in cold-induced transcriptional regulation of RH

growth, we analyzed the sequence of the promoter regions of RH-related *APOLO* targets. Notably, 13 out

of the 17 *APOLO* targets contained between 1 and 4 canonical WRKY TF binding sites (W-box) in their

promoters, including *RHD6* (**Supplementary Table 1**)*. According to the Arabidopsis eFP Browser (37), the

TF-encoding gene *WRKY42* is induced in roots when seedlings are subjected to 4°C for 24h (**Figure 3A**).

Remarkably, WRKY42 is involved in the response to Pi starvation (23), an environmental condition that

promotes RH cell expansion (18) in a similar manner to low temperatures. At 10°C, we observed that

WRKY42 transcriptional accumulation augments significantly in roots (**Figure 3B**). By using a

35S:WRKY42:GFP line, we demonstrated that WRKY42 can directly bind to the promoter region of *RHD6*

(**Figure 3C**). Accordingly, the over-expression of *WRKY42* (*35S:WRKY42:GFP* line) led to a basal increase

of *RHD6* levels (**Figure 3D**) and RH elongation (**Figure 3E**) at ambient temperature, mimicking the effect

of cold. On the contrary, cold-mediated induction of *RHD6* is abolished in the *wrky42* mutant (23; **Figure**

3D), which consistently exhibits shorter RHs at 22°C and partially impaired RH elongation at low temper-

atures (**Figure 3E**). Taken together, our results suggest that WRKY42 is an important regulator of *RHD6*-

mediated RH growth in response to cold.

We thus wondered to what extent WRKY42 regulates the epigenetic landscape of the *RHD6* lo-

cus. We first observed that H3K27me3 deposition over *RHD6* is significantly augmented in the *wrky42*

mutant background, in contrast to *AZG2*, an *APOLO* target non-related to WRKY42 (**Figure 4A**), con-

sistent with reduced *RHD6* basal levels reported in *wrky42* (**Figure 3D**). Considering that other TFs were

shown to directly interact with lncRNAs in animals (38-40), we wondered if WRKY42 can recognize

APOLO in planta. Thus, *APOLO*-WRKY42 interaction was assessed and validated by RNA immuno-

precipitation (RIP-qPCR) in tobacco leaves and in *Arabidopsis* plants transitory or stably transformed with

35S:*WRKY42:GFP*, respectively (**Figure 4B**). Therefore, we evaluated the mutual contribution of *APOLO*

and WRKY42 to their respective recognition of the *RHD6* locus. *APOLO* ChIRP-qPCR in the WT and *wrky42*

mutant (**Figure 4C**) revealed similar binding to *RHD6*, indicating that WRKY42 does not participates in

APOLO-target recognition. Reciprocally, we assessed the control of *APOLO* over WRKY42 recognition of

the *RHD6* locus. To this end, chromatin was extracted from *35S:WRKY42:GFP* seedlings and increasing

amounts of *in vitro* transcribed *APOLO* were added before cross-link and regular WRKY42 ChIP over

RHD6. Strikingly, increasing concentrations of *APOLO* gradually decoy WRKY42 away from the *RHD6* lo-

cus (**Figure 4D**), hinting at a stoichiometric regulation of *APOLO* over the activity of its partner TF.

Collectively, our results indicate that the regulation of *RHD6* expression in response to cold

depends on Polycomb-dependent H3K27me3 dynamic deposition. The WRKY42-*APOLO* complex

modulates the epigenetic environment of *RHD6*, activating its transcription and promoting RH growth at

low temperatures. *RHD6* activation further triggers the expression of *RSL2* and *RSL4* that control the

transcriptional RH program inducing cell expansion in response to cold (**Figure 4E**).

DISCUSSION

Cell fate determination in the epidermis has been extensively studied (1–4). Once trichoblast

cells differentiate in the root epidermis, RHs develop as fast polar growing protuberances in response to

endogenous and environmental signals (6, 7, 9, 19). RHs are one of the main entry points in the roots for

water-soluble macronutrients, such as Pi and nitrates. Pi is an essential element for plant growth and

development, and the availability of this macronutrient is a factor limiting plant productivity. Low Pi in

the soil triggers auxin synthesis and transport in Arabidopsis roots, enhancing RH elongation to promote

Pi uptake (18). Thus, auxin mediates low Pi-induced promotion of RH cell expansion. Under low soil Pi,

auxin synthesis is enhanced specifically in the root cap (41) and transported (mostly by AUX1, PIN2, and

PGP4) from the apex to the differentiation zone, specifically leading to an increase of auxin levels in

trichoblasts (18, 42, 43). In response to the high-auxin microenvironment, RHs protrude from the root

epidermis controlled by RHD6 and RSL1 (6, 7). High levels of auxin in trichoblasts trigger a signaling cas-

cade mediated by TIR1-ARF19 (and possibly also ARF7) that directly induce the expression of *RSL4* (and

likely of *RSL2*) and promote RH elongation (9, 17, 18, 44). ARF7 and ARF19 also activate other RH genes

independently of RSL4 (45). Interestingly, our results indicate that the lncRNA *APOLO* participates in the

response to low temperatures. *APOLO* is directly activated by ARF7 and regulates the transcriptional

activity of its neighboring gene *PINOID* (*PID*) by shaping local 3D chromatin conformation (26, 27). *PID*

encodes a kinase responsible for accurate auxin polar transport by localizing PIN2 in the root cell mem-

brane (46). More recently, it was shown that *APOLO* can recognize a subset of distant genes across the

Arabidopsis genome, most of them being related to auxin synthesis and signaling (27). In this work, we

demonstrate that a group of RH related genes are directly regulated by *APOLO* in response to cold, in-

cluding the RH master regulator *RHD6*. Collectively, our results uncover a lncRNA-mediated epigenetic

link between environmental signals and auxin homeostasis modulating RH growth.

As mentioned above, nutrient availability is known to activate RH expansion through a transcrip-

tional reprogramming governed by RHD6 and downstream TFs. The quantification of RH growth of WT

plants in response to increasing concentrations of nutrients (0.5X to 2.0X MS (Murashige and Skoog) me-

dium) indicates that high concentrations of nutrients completely abolish RH growth triggered by low

temperatures **(Supplementary Figure 4A)**. In a similar way, by increasing agar concentration in the MS

medium (from 0.8% to 2.5%) restraining nutrient mobility, cold-induced RH is clearly blocked **(Supple-**

mentary Figure 4B). Altogether, these observations suggest that low temperatures may restrict nutrient

mobility and availability in the culture medium, leading to the promotion of polar RH growth. Further

research will be needed to determine what is the limiting nutrient mediating the effect of cold on RH

growth.

Although substantial progress has been achieved in the identification of the molecular actors

that control RH development, the impact of chromatin conformation in the transcriptional regulation of

central TFs remains poorly understood. In this study, we have revealed a new mechanism of gene regula-

tion in RHs by which the lncRNA *APOLO* integrates chromatin-associated ribonucleoprotein complexes

together with the TF WYRK42, participating in the transcriptional activation of *RHD6* and the down-

stream RH gene network (**Figure 4D**). *APOLO* directly regulates the chromatin 3D conformation of the

genomic region encompassing the *RHD6* locus and stoichiometrically recruits WYRK42, previously linked

to Pi-starvation (21, 23). Our results suggest that an *APOLO*-WRKY42 hub regulates RH cell elongation

through the master regulator RHD6, although the *APOLO*-WYRKY42 hub potentially targets several addi-

tional cell wall related genes (**Supplementary Table 1**) at the end of the pathway controlled by RHD6 and

the RHD6-downstream TFs RSL2/RSL4 (17, 47).

Participation of epigenetic factors in root cell identity determination strongly suggests that the

default pattern for epidermal cell fate can be overridden by environmental stimuli (48). Interestingly, it

was reported that the expression of *GLABRA2* (*GL2*), a gene encoding a TF repressing *RHD6* in

atrichoblasts, is tightly regulated at the epigenetic level. By using 3D fluorescence *in situ* hybridization, it

was shown that alternative states of chromatin organization around the *GL2* locus are required to con-

trol position-dependent cell-type specification in the root epidermis (49). Furthermore, *GL2* epigenetic

regulation was proposed to be responsive to salt stress (50). In addition, a comprehensive characteriza-

tion of alternative mutant lines uncovered the role of PRC2 in the regulation of RH development (36).

Loss-of-function mutants in different PRC2 subunits develop unicellular RHs but fail to retain the differ-

entiated state, generating a disorganized cell mass from each single RH. It was shown that the resulting

RHs are able to undergo a normal endoreduplication program, increasing their nuclear ploidy, although

they subsequently reinitiate mitotic division and successive DNA replication. It was proposed that aber-

rant RH development in PRC2-related mutants is due to the epigenetic deregulation of key genes such as

WOUND INDUCED DEDIFFERENTIATION 3 (*WIND3*) and *LEAFY COTYLEDON 2* (*LEC2*) (36). Here, we

showed that the single mutants *clf* (PRC2) and *lhp1* (PRC1) are affected in RH growth. Moreover, we

showed that H3K27me3 deposition throughout the *RHD6* locus is partially impaired in the *clf* back-

ground, which is affected in RH elongation promoted by cold. Altogether, our results suggest that

Polycomb proteins participate in the control of RH-related genes transcriptional reprogramming at low
temperatures.

Notably, CLF and LHP1 were shown to interact with a subset of lncRNAs in *Arabidopsis*, modulat-
ing the activity of PRC target genes (51). Interestingly, several lncRNAs have been linked to the control of
transcription in response to cold. *FLOWERING LOCUS C* is regulated by at least three lncRNAs. First, the
alternative splicing of a set of antisense transcripts, collectively named as *COOLAIR*, depends on the pro-
longed exposure to cold, epigenetically repressing *FLC* (52). The use of the *COOLAIR* proximal poly(A) site
results in down-regulation of *FLC* expression in a process involving FLOWERING LOCUS D (FLD), an
H3K4me2 demethylase (53). A second lncRNA called *COLD ASSISTED INTRONIC NONCODING RNA*
(*COLDAIR*) is fully encoded in the first intron of *FLC.* Similar to *COOLAIR*, its accumulation oscillates in
response to low temperatures. It was proposed that *COLDAIR* recruits the PRC2 component CLF to target
FLC for H3K27me3 deposition (54). More recently, a third lncRNA modulating *FLC* transcription was iden-
tified (55). The cold-responsive lncRNA *COLDWRAP* is derived from the *FLC* proximal promoter and it also
interacts with PRC2. It was suggested that *COLDWRAP* functions in cooperation with the lncRNA
COLDAIR to retain Polycomb at the *FLC* promoter through the formation of a repressive intragenic chro-
matin loop (55). Another lncRNA named *SVALKA* was shown to mediate the response to low tempera-
tures (56). Interestingly, the activation of *SVALKA* by cold triggers the transcription of a cryptic down-
stream lncRNA, which overlaps the antisense locus of the *C-repeat/dehydration-responsive element Bind-
ing Factor 1* (*CBF1*), involved in the early response to cold in Arabidopsis. Antisense transcription causes
Pol II head-to-head collision modulating transcriptional termination of *CBF1* (56). Here, we showed that
the auxin-responsive lncRNA *APOLO* is also transcriptionally modulated by cold. The differential accumu-
lation of 24nt siRNAs across the *APOLO* locus at low temperatures indicates that this activation is related
to a decrease in RdDM. Moreover, we showed here that the intergenic region between *PID* and *APOLO*
acting as a divergent promoter is also activated at low temperatures in RHs, as revealed by the reporter
gene *GFP*. Thus, the lncRNA *APOLO* integrates external signals into auxin-dependent developmental out-
puts in Arabidopsis.

In the last decade, lncRNAs have emerged as regulators of gene expression at different levels,
ranging from epigenetics to protein modifications and stability (57). Notably, it has been shown in ani-
mals that noncoding transcripts can be recognized by TFs. In humans, it was proposed that the interac-
tion with the lncRNA *SMALL NUCLEOLAR RNA HOST GENE 15* (*SNHG15*) stabilizes the TF Slug in colon
cancer cells. It was shown that *SNHG15* is recognized by the zinc finger domain of Slug preventing its
ubiquitination and degradation in living cells (38). Also, the transcriptional activity of the human gene

DIHYDROFOLATE REDUCTASE (*DHFR*) is regulated by a lncRNA encoded in its proximal promoter. It was

proposed that the nascent noncoding transcript forms a hybrid with its parent DNA and decoys the regu-

latory TF IIB away from the *DHFR* promoter, dissociating the transcriptional pre-initiation complex in

quiescent cells (40). The lncRNA *P21 ASSOCIATED ncRNA DNA DAMAGE ACTIVATED* (*PANDA*) was identi-

fied in human cancer and it was activated in response to DNA damage (39). *PANDA* is transcribed from

the promoter region of the *CDKN1A* gene and interacts with the TF NF-YA to limit the expression of pro-

apoptotic genes. The activity of *PANDA* has been linked to the progression of different tumors (58, 59).

Interestingly, it was shown that in addition to NF-YA, *PANDA* interacts with the scaffold-attachment-

factor A (SAFA) as well as PRC1 and PRC2 to modulate cell senescence. In proliferating cells, SAFA and

PANDA recruit Polycomb components to repress the transcription of senescence-promoting genes. Con-

versely, the loss of SAFA–PANDA–PRC interactions allows expression of the senescence program (60). In

this work, we showed that the PRC1-interacting lncRNA *APOLO* can also be recognized by the TF

WRKY42, hinting at general lncRNA-mediated mechanisms linking Polycomb complexes with the tran-

scriptional machinery across kingdoms. Furthermore, our observations indicate that the deregulation of

WRKY42 affects the epigenetic environment of *RHD6*. It was previously shown that the addition of *in*

vitro transcribed *APOLO* to RNAi-*APOLO* chromatin extracts was able to partially restore R-loop for-

mation over *APOLO* target genes, and that high levels of *APOLO* may titer LHP1 away from chromatin

(27). Here we showed that the relative accumulation of the lncRNA *APOLO* can modulate the binding

activity of its partner TF to common target genes. Collectively, our results strongly support that environ-

mentally controlled cell fate in *Arabidopsis* relies on a transcriptional reprogramming governed by a

network of epigenetic regulatory complexes, lncRNAs, TFs and effector proteins.

MATERIALS AND METHODS

Plant Material and Growth Conditions. All the *Arabidopsis thaliana* lines used were in the Columbia-0

(Col-0) background. *WRKY42* over expression transgenic plants were generated through *Agrobacterium*

tumefaciens (strain EHA105)-mediated transformation (61). *35S:WRKY42:GFP* transformant lines were

selected on MS/2 medium supplemented with kanamycin (40μg/mL) and *WRKY42* expression levels were

measured by RT-qPCR (primers used are listed in **Supplementary Table 3**). The *wrky42* mutant line be-

longs to the SALK collection (SALK_121674C), as the one previously characterized (23). Homozygous

plants were obtained in our laboratory and genotyped using the oligonucleotides indicated in **Supple-**

mentary Table 3. Seeds were surface sterilized and stratified at 4°C for 2d before being grown under

long day conditions (16h light, 140µE.m^{-2}.sec^{-1}/8h dark), on ½-strength Murashige and Skoog media (1/2

MS) (Duchefa, Netherlands) with 0.8% plant agar (Duchefa, Netherlands).

Cloning procedure. The coding region of *WRKY42* (AT4G04450) excluding the STOP codon was amplified

by PCR, cloned into the Gateway entry vector pENTR/D-TOPO (Invitrogen) and recombined by Gateway

technology (LR reaction) into the pK7FWG2,0 vector containing a *p35S-GFP* cassette

(http://www.psb.ugent.be/gateway/index.php).

Root hair phenotype characterization. For quantitative analyses of RH phenotypes, 100 fully elongated

RHs were measured (using the ImageJ software) from 10-20 roots grown on vertical agar plates for 5

days at 22° and 3 days at 10°C. Measurements were made after 8 days. Images were captured with an

Olympus SZX7 Zoom Stereo microscope (Olympus, Japan) equipped with a Q-Colors digital camera and

software.

Confocal microscopy analysis of root hairs. Confocal laser scanning microscopy was performed using

Zeiss LSM5 Pascal (Zeiss, Germany) and a 40x water-immersion objective, N/A=1.2. Fluorescence was

analyzed by using 488 nm laser for GFP excitation (Laser Intensity: 70%, Detector Gain:550, Amplifier

Offset:0.1, Amplifier Gain:1), and emitted fluorescence was recorded between 490 and 525nm for GFP

tag. Z stacks were done with an optical slice of 1µm, and fluorescence intensity was measured in 15µm

ROI (Region Of Interest) at the RH tip and summed for quantification of fluorescence using ImageJ. Five

replicates for each of ten roots and 15 hairs per root were observed. Col-0 wild type root hairs were used

as a negative control, to check autofluorescence signal occurrence and no signal were detected in the

wavelengths range stated above.

RNA extraction and RT-qPCR. Total RNA was extracted using TRIZol (Invitrogen) and 2 µg were subjected

to DNase treatment according to the manufacturer's protocol (Thermo Scientific). One µg of DNase-free

RNA was reverse transcribed using Maxima H Minus Reverse Transcriptase (Thermo Scientific). RT-qPCR

were performed using the LightCycler 480 SYBR Green I Master Kit on a LightCycler480 apparatus

(Roche) using standard protocols (40 cycles, 60°C annealing). *PP2A* (AT1G13320; primers are listed in

Supplementary Table 3) was used as reference.

RNA Immunoprecipitation. RNA immunoprecipitation (RIP) assays were performed on transiently transformed *N. benthamiana* leaves as described in (62), or in 10-day-old *A. thaliana 35S:WRKY42:GFP* seedlings as described in (63), using anti GFP (Abcam ab290) and anti-IgG (Abcam ab6702). RIP was performed using Invitrogen Protein A Dynabeads. Precipitated RNAs were prepared using TRI Reagent (Sigma-Aldrich), treated with DNase (Fermentas) and subjected to RT-qPCR (High Capacity cDNA Reverse Transcription Kit (Thermo); primers used are listed in **Supplementary Table 3**). Total RNAs were processed in parallel and considered the input sample.

Chromatin Immunoprecipitation. Chromatin immunoprecipitation (ChIP) assays were performed on 10-day-old WT seedlings treated or not during 24h at 10 °C, using anti H3K27me3 (Diagenode pAb-195-050), anti LHP1 (Covalab pab0923-P) and anti-IgG (Abcam ab6702), as described in (27). Crosslinked chromatin was sonicated using a water bath Bioruptor Pico (Diagenode; 30sec ON/30sec OFF pulses; 10 cycles; high intensity). ChIP was performed using Invitrogen Protein A Dynabeads. Precipitated DNA was recovered using Phenol:Chloroform:Isoamilic Acid (25:24:1; Sigma) and subjected to RT-qPCR (primers used are listed in **Supplementary Table 3**). Untreated sonicated chromatin was processed in parallel and considered the input sample. For *in vitro* competition assays, *APOLO* was transcribed using the T7 transcription kit (Promega; (27)). After regular chromatin isolation from 10-day-old *35S:WRKY42:GFP* seedlings, the sample was split in 4 independent tubes and diluted to 1ml in Nuclei Lysis Buffer without SDS. 0 µg, 0.1 µg, 1 µg and 10 µg of *APOLO* were added to each sample respectively, and incubated in rotation at 4 °C for 3h. Then, cross-linking was performed with 1% formaldehyde for 5 min at 4 °C, followed by 5 min with a final concentration of 50 mM glycine. SDS was added to a final concentration of 0.1% prior to sonication and the subsequent steps of a regular ChIP protocol.

Chromatin Isolation by RNA Purification followed by qPCR. A method adapted from the ChIRP protocol (64) was developed to allow the identification of plant DNA associated to specific lncRNAs, as described in (26, 27). Briefly, plants were *in vivo* crosslinked and cell nuclei were purified and extracted through sonication. The resulting supernatant was hybridized against biotinylated complementary oligonucleotides that tile the lncRNA of interest and lncRNA-associated chromatin was isolated using magnetic streptavidin beads. One hundred pmol of probes against *APOLO* (ODD and EVEN set of probes (26, 27)) and the corresponding negative set against LacZ were used for independent purification. Co-purified ribonucleoprotein complexes were eluted and used to purify RNA or DNA, which were later subjected to downstream assays for quantification as previously described (27).

Chromatin Conformation Capture. Chromosome conformation capture (3C) was performed basically as previously described in (27) starting with 2g of seedlings. Digestions were performed overnight at 37°C with 400U *Dpn*II (NEB). DNA was ligated by incubation at 16°C for 5h in 4 ml volume using 100U of T4 DNA ligase (NEB). After reverse crosslinking and Proteinase K treatment (Invitrogen), DNA was recovered by Phenol:Chloroform:Isoamilic Acid (25:24:1; Sigma) extraction and ethanol precipitation. Relative interaction frequency was calculated by qPCR (primers used are listed in **Supplementary Table 3**). A region free of *Dpn*II was used to normalize the amount of DNA.

Acknowledgments

We thank Chang Liu for providing the Hi-C plot of Figure 2B. This work was supported by grants from ANPCyT (PICT2016-0132 and PICT2017-0066) and Instituto Milenio iBio – Iniciativa Científica Milenio, MINECON to JME; ANPCyT (PICT2016-0007 and PICT2016-0289) to FA; UNL (CAI+D 2016) to LL; Laboratoire d'Excellence (LABEX) Saclay Plant Sciences (SPS; ANR-10-LABX-40) to MC; and CNRS (Laboratoire International Associé NOCOSYM) to MC and FA. LL, FI, JME and FA are researchers of CONICET; MM, JMP, CFF and JRM are fellows of the same institution.

Author contributions

MM, JMP, LL, CFF, JRM, AC, MB and FA performed the experiments. JB, MB, FI, MC, JE and FA analyzed the data. JE and FA conceived the project. FA, JE and CFF wrote the manuscript with the contribution of all authors.

References

1.   K. H. Ryu, *et al.*, The WEREWOLF MYB protein directly regulates CAPRICE transcription during cell fate specification in the Arabidopsis root epidermis. *Development* **132**, 4765–4775 (2005).

2.   S. K. Song, *et al.*, Cell fate in the Arabidopsis root epidermis is determined by competition between WEREWOLF and CAPRICE. *Plant Physiol.* **157**, 1196–1208 (2011).

3.   J. Schiefelbein, L. Huang, X. Zheng, Regulation of epidermal cell fate in Arabidopsis roots: The importance of multiple feedback loops. *Front. Plant Sci.* **5** (2014).

4.   D. Balcerowicz, S. Schoenaers, K. Vissenberg, Cell fate determination and the switch from diffuse growth to planar polarity in arabidopsis root epidermal cells. *Front. Plant Sci.* **6**, 1163 (2015).

5.   Q. Lin, *et al.*, GLABRA2 directly suppresses basic helix-loop-helix transcription factor genes with

428   diverse functions in root hair development. *Plant Cell* **27**, 2894–2906 (2015).

429 6. B. Menand, *et al.*, An ancient mechanism controls the development of cells with a rooting func-
430   tion in land plants. *Science (80-.).* **316**, 1477–1480 (2007).

431 7. N. D. Pires, *et al.*, Recruitment and remodeling of an ancient gene regulatory network during land
432   plant evolution. *Proc. Natl. Acad. Sci. U. S. A.* **110**, 9571–9576 (2013).

433 8. B. Karas, *et al.*, Conservation of Lotus and Arabidopsis basic helix-loop-helix proteins reveals new
434   players in root hair development. *Plant Physiol.* **151**, 1175–1185 (2009).

435 9. K. Yi, B. Menand, E. Bell, L. Dolan, A basic helix-loop-helix transcription factor controls cell growth
436   and size in root hairs. *Nat. Genet.* **42**, 264–267 (2010).

437 10. A. Bruex, *et al.*, A gene regulatory network for root epidermis cell differentiation in Arabidopsis.
438   *PLoS Genet.* **8** (2012).

439 11. S. K. Won, *et al.*, cis-element- and transcriptome-based screening of root hair-specific genes and
440   their functional characterization in Arabidopsis. *Plant Physiol.* **150**, 1459–1473 (2009).

441 12. S. Datta, H. Prescott, L. Dolan, Intensity of a pulse of RSL4 transcription factor synthesis deter-
442   mines Arabidopsis root hair cell size. *Nat. Plants* **1**, 1–6 (2015).

443 13. P. Vijayakumar, S. Datta, L. Dolan, ROOT HAIR DEFECTIVE SIX-LIKE4 (RSL4) promotes root hair
444   elongation by transcriptionally regulating the expression of genes required for cell growth. *New*
445   *Phytol.* **212**, 944–953 (2016).

446 14. C. Ringli, The hydroxyproline-rich glycoprotein domain of the Arabidopsis LRX1 requires Tyr for
447   function but not for insolubilization in the cell wall. *Plant J.* **63**, 662–669 (2010).

448 15. S. M. Velasquez, *et al.*, O-glycosylated cell wall proteins are essential in root hair growth. *Science*
449   *(80-.).* **332**, 1401–1403 (2011).

450 16. Y. Hwang, H. S. Choi, H. M. Cho, H. T. Cho, Tracheophytes contain conserved orthologs of a basic
451   helix-loop-helix transcription factor that modulate ROOT HAIR SPECIFIC genes. *Plant Cell* **29**, 39–
452   53 (2017).

453 17. S. Mangano, *et al.*, Molecular link between auxin and ROS-mediated polar growth. *Proc. Natl.*
454   *Acad. Sci. U. S. A.* **114**, 5289–5294 (2017).

455 18. R. Bhosale, *et al.*, A mechanistic framework for auxin dependent Arabidopsis root hair elongation
456   to low external phosphate. *Nat. Commun.* **9** (2018).

457 19. E. Marzol, C. Borassi, S. P. Denita Juárez, S. Mangano, J. M. Estevez, RSL4 Takes Control: Multiple
458   Signals, One Transcription Factor. *Trends Plant Sci.* **22**, 553–555 (2017).

459 20. B. N. Devaiah, A. S. Karthikeyan, K. G. Raghothama, WRKY75 transcription factor is a modulator of

phosphate acquisition and root development in Arabidopsis. *Plant Physiol.* **143**, 1789–1801

(2007).

21.   Y. F. Chen, *et al.*, The WRKY6 transcription factor modulates PHOSPHATE1 expression in response

to low pi stress in arabidopsis. *Plant Cell* **21**, 3554–3566 (2009).

22.   H. Wang, *et al.*, Arabidopsis WRKY45 transcription factor activates Phosphate transporter1;1 ex-

pression in response to phosphate starvation. *Plant Physiol.* **164**, 2020–2029 (2014).

23.   T. Su, *et al.*, WRKY42 Modulates phosphate homeostasis through regulating phosphate transloca-

tion and acquisition in Arabidopsis. *Plant Physiol.* **167**, 1579–1591 (2015).

24.   J. M. Franco-Zorrilla, *et al.*, Target mimicry provides a new mechanism for regulation of microRNA

activity. *Nat. Genet.* **39**, 1033–1037 (2007).

25.   M. Jabnoune, *et al.*, A Rice cis-Natural antisense RNA acts as a translational enhancer for its cog-

nate mRNA and contributes to phosphate homeostasis and plant fitness. *Plant Cell* **25**, 4166–4182

(2013).

26.   F. Ariel, *et al.*, Noncoding transcription by alternative rna polymerases dynamically regulates an

auxin-driven chromatin loop. *Mol. Cell* **55**, 383–396 (2014).

27.   F. Ariel, *et al.*, R-Loop Mediated trans Action of the APOLO Long Noncoding RNA. *Mol. Cell* **77**,

1055-1065.e4 (2020).

28.   D. T. A. Lamport, M. J. Kieliszewski, Y. Chen, M. C. Cannon, Role of the Extensin Superfamily in

Primary Cell Wall Architecture. *Plant Physiol.* **156**, 11–19 (2011).

29.   S. M. Velasquez, *et al.*, Low sugar is not always good: Impact of specific o-glycan defects on tip

growth in arabidopsis. *Plant Physiol.* **168**, 808–813 (2015).

30.   E. Marzol, *et al.*, Filling the Gaps to Solve the Extensin Puzzle. *Mol. Plant* **11**, 645–658 (2018).

31.   T.Q. Zhang, Z.G. Xu, G.D. Shang, J.W. Wang, A Single-Cell RNA Sequencing Profiles the Develop-

mental Landscape of Arabidopsis Root. *Mol Plant*. 12, 648-660 (2019).

32.   P. Gyula, *et al.*, Ambient temperature regulates the expression of a small set of sRNAs influencing

plant development through NF-YA2 and YUC2. *Plant Cell Environ.* **41**, 2404–2417 (2018).

33.   A. Veluchamy, *et al.*, LHP1 Regulates H3K27me3 Spreading and Shapes the Three-Dimensional

Conformation of the Arabidopsis Genome. *PLoS One* **11**, e0158936 (2016).

34.   W. Xu, *et al.*, The R-loop is a common chromatin feature of the Arabidopsis genome. *Nat. Plants*

3, 704–714 (2017).

35.   C. Liu, *et al.*, Genome-wide analysis of chromatin packing in Arabidopsis thaliana at single-gene

resolution. *Genome Res.* **26**, 1057–1068 (2016).

36.   M. Ikeuchi, *et al.*, PRC2 represses dedifferentiation of mature somatic cells in Arabidopsis. *Nat.*
Plants **1**, 1–7 (2015).

37.   J. Waese, *et al.*, ePlant: Visualizing and Exploring Multiple Levels of Data for Hypothesis Genera-
tion in Plant Biology. *Plant Cell* **29**, 1806-1821 (2017).

38.   H. Jiang, *et al.*, Long non-coding RNA SNHG15 interacts with and stabilizes transcription factor
Slug and promotes colon cancer progression. *Cancer Lett.* **425**, 78–87 (2018).

39.   T. Hung, *et al.*, Extensive and coordinated transcription of noncoding RNAs within cell-cycle pro-
moters in *Nature Genetics*, (NIH Public Access, 2011), pp. 621–629.

40.   I. Martianov, *et al.*, Repression of the human dihydrofolate reductase gene by a non-coding inter-
fering transcript. *Nature* **445**, 666-670 (2007).

41.   A. N. Stepanova, *et al.*, TAA1-Mediated Auxin Biosynthesis Is Essential for Hormone Crosstalk and
Plant Development. *Cell* **133**, 177–191 (2008).

42.   A. R. Jones, *et al.*, Auxin transport through non-hair cells sustains root-hair development. *Nat. Cell*
Biol. **11**, 78–84 (2009).

43.   Q. Wang, *et al.*, A phosphorylation-based switch controls TAA1-mediated auxin biosynthesis in
plants. *Nat. Commun.* **11**, 679 (2020).

44.   S. Mangano, S. P. Denita-Juarez, E. Marzol, C. Borassi, J. M. Estevez, High auxin and high phos-
phate impact on rsl2 expression and ros-homeostasis linked to root hair growth in Arabidopsis
thaliana. *Front. Plant Sci.* **9** (2018).

45.   S. Schoenaers, *et al.*, The Auxin-Regulated CrRLK1L Kinase ERULUS Controls Cell Wall Composition
during Root Hair Tip Growth. *Curr. Biol.* **28**, 722-732.e6 (2018).

46.   J. Friml, *et al.*, A PINOID-dependent binary switch in apical-basal PIN polar targeting directs auxin
efflux. *Science (80-.).* **306**, 862–865 (2004).

47.   S. Mangano, J. Martínez Pacheco, C. Marino-Buslje, J. M. Estevez, How Does pH Fit in with Oscil-
lating Polar Growth? *Trends Plant Sci.* **23**, 479–489 (2018).

48.   S. Guimil, C. Dunand, Patterning of Arabidopsis epidermal cells: epigenetic factors regulate the
complex epidermal cell fate pathway. *Trends Plant Sci.* **11**, 601–609 (2006).

49.   S. Costa, P. Shaw, Chromatin organization and cell fate switch respond to positional information
in Arabidopsis. *Nature* **439**, 493–496 (2006).

50.   C. C. Beyrne, N. D. Iusem, R. M. González, Effect of salt stress on cytosine methylation within GL2,

an Arabidopsis thaliana gene involved in root epidermal cell differentiation. Absence of inher-

itance in the unstressed progeny. *Int. J. Mol. Sci.* **20** (2019).

51 L. Lucero, *et al.*, Long noncoding RNAs shape transcription in plants. *Transcription* (2020).

52. S. Marquardt, *et al.*, Functional Consequences of Splicing of the Antisense Transcript COOLAIR on

FLC Transcription. *Mol. Cell* **54**, 156–165 (2014).

53. F. Liu, S. Marquardt, C. Lister, S. Swiezewski, C. Dean, Targeted 3' processing of antisense tran-

scripts triggers arabidopsis FLC chromatin silencing. *Science (80-.).* **327**, 94–97 (2010).

54. J. B. Heo, S. Sung, Vernalization-mediated epigenetic silencing by a long intronic noncoding RNA.

Science (80-.). **331**, 76–79 (2011).

55. D. H. Kim, S. Sung, Vernalization-Triggered Intragenic Chromatin Loop Formation by Long Noncod-

ing RNAs. *Dev. Cell* **40**, 302-312.e4 (2017).

56. P. Kindgren, R. Ard, M. Ivanov, S. Marquardt, Transcriptional read-through of the long non-coding

RNA SVALKA governs plant cold acclimation. *Nat. Commun.* **9**, 1–11 (2018).

57. F. Ariel, *et al.*, Battles and hijacks: noncoding transcription in plants. *TiPS* **20**, 362–371 (2015).

58. Y. Kotake, *et al.*, Long Non-coding RNA, PANDA, Contributes to the Stabilization of p53 Tumor

Suppressor Protein. *Anticancer Res.* **36**, 1605–11 (2016).

59. W. Shi, *et al*., Long noncoding RNA PANDA promotes esophageal squamous carcinoma cell pro-

gress by dissociating from NF-YA but interact with SAFA. *Pathol. Research and Practice* **215**,

152604 (2019).

60. P. K. Puvvula, *et al.*, Long noncoding RNA PANDA and scaffold-attachment-factor SAFA control

senescence entry and exit. *Nat. Commun.* **5** (2014).

61. S.J. Clough, A.F. Bent, Floral dip: a simplified method for Agrobacterium -mediated transformation

of Arabidopsis thaliana. *The Plant Journal* **16**, 735-743 (1998).

62. R. Sorenson, J. Bailey-Serres, Rapid immunopurification of ribonucleoprotein complexes of plants.

Methods Mol. Biol. **1284**, 209–219 (2015).

63. F. Bardou, *et al.*, Long Noncoding RNA Modulates Alternative Splicing Regulators in Arabidopsis.

Dev. Cell **30**, 166–176 (2014).

64. C. Chu, J. Quinn, H. Y. Chang, Chromatin isolation by RNA purification (ChIRP). *J. Vis. Exp.* (2012)

https:/doi.org/10.3791/3912.

65. J. Dong, C. Chen, Z. Chen, Expression profiles of the Arabidopsis WRKY gene superfamily during

plant defense response. *Plant Mol. Biol.* **51**, 21–37 (2003).

Figure 1. *APOLO* regulates root hair elongation in response to low temperatures

A. 24nt siRNA accumulation over the *APOLO* locus revealed by small RNA-Seq of roots from WT seedlings grown at 15 °C or 23 °C (32).

B. *APOLO* transcript levels measured by RT-qPCR in roots of 10-day-old plants grown at 22 °C treated or not for 24h at 10 °C.

C. The *APOLO* promoter region (full intergenic region between *PID* and *APOLO*) is active in root hairs at 22 °C and 10 °C, using the *GFP* reporter gene. Images are representative of the lowest and highest GFP signal detected in the observation of several roots. The quantification of the fluorescence is shown in the right panel.

D. Quantification of RH length of 8-day-old WT, RNAi-*APOLO* and *35S:APOLO* plants, respectively, at 22 °C and 10 °C. Each point is the mean of the length of the 10 longest RHs identified in a single root. Repre-

sentative images of each genotype are shown below the graph. Asterisks indicate significant differences
between WT and the corresponding genotype at the same temperature (one way ANOVA followed by a
Tukey-Kramer test; "*"<0.05, "**"<0.01, "***"<0.001, NS: no statistically significant).
E. *RHD6* transcript levels measured by RT-qPCR in roots of WT plants vs. two independent RNAi-*APOLO*
lines grown at 22 °C treated or not for 24h at 10 °C.
In B and E, the error bar represents the SD of 3 biological replicates. The asterisk indicates that the dif-
ference is significant (t test p<0.05).

Figure 2. *APOLO* directly modulates chromatin three-dimensional conformation of the *RHD6* locus

A. Epigenomic landscape of the *RHD6* locus. Lane 1: H3K27me3 deposition by ChIP-Seq (33). Lane 2: LHP1 deposition by ChIP-Seq (33). Lane 3 to 5: *APOLO* recognition by ChIRP-Seq (Lane 3 and 4, using ODD and EVEN sets of probes against *APOLO*, respectively; Lane 5, negative control using LacZ probes)(27). Lane 6 to 9: R-loop formation by DRIP-Seq (34), on Watson strand (Lane 6), Crick strand (Lane 7) or unstranded sequencing (Lane 8). DRIP negative control after RNAseH treatment is shown in Lane 9. Gene annotation is shown at the bottom.

B. Chromatin loops identified in the *RHD6* region by *Dpn*II HiC (35). Colors of the loops are related to the corresponding q-values indicated below. Black boxes in A and B indicate the same genomic locations, where the bases of the chromatin loop correlate with R-loop formation and *APOLO* recognition (compared to gene annotations).

C. *APOLO* association to DNA of the *RHD6* locus by ChIRP-qPCR in WT and RNAi-*APOLO* plants. The background level was determined using a set of probes against LacZ RNA.

588   **D.** Relative chromatin loop formation by 3C-qPCR deduced from **B**, in WT plants vs. *35S:APOLO* and RNAi
589   lines for the region upstream *RHD6*.
590   **E.** *RHD6* transcript levels revealed by RNA-Seq of WT vs. *35S:APOLO* plants(27).
591   In C and D, the error bar represents the SD of 3 biological replicates. The asterisk indicates that the dif-
592   ference is significant (t test "*"<0.05). In E, p-value from RNA-Seq (27) "***"<0.001.
593

Figure 3. The transcription factor WRKY42 directly regulates *RHD6* and participates in the response to cold

A. Arabidopsis eFP Browser (37) plot representing the increase of *WRKY42* transcript levels in roots of seedlings treated for 24h at 4°C.

B. *WRKY42* transcript levels measured by RT-qPCR in roots of plants grown at 22°C treated or not for 24h at 10°C.

C. Chromatin Immunoprecipitation (ChIP)-qPCR assay revealing regulation by WRKY42 of *RHD6* by direct recognition of its promoter region. Probes amplifying *PP2A* were used as a negative control of the experiment. Anti-IgG antibodies were used as a negative control for each pair of probes.

D. *RHD6* transcript levels measured by RT-qPCR in roots of WT plants vs. *wrky42* mutants and *35S:WRKY42:GFP* lines grown at 22°C treated or not for 24h at 10°C.

E. Quantification of RH length of plants WT, *wrky42* and *35S:WRKY42:GFP* at 22°C and 10°C. Each point is the mean of the length of the 10 longest RHs identified in a single root. Representative images of each

609   genotype are shown on the right. Asterisks indicate significant differences between WT and the corre-
610   sponding genotype at the same temperature (one way ANOVA followed by a Tukey-Kramer test;
611   "*"<0.05, "***"<0.001, NS: no statistically significant).
612   In B, C and D, the error bar represents the SD of 3 biological replicates. The asterisk indicates that the
613   difference is significant (t test $p<0.05$, NS: no statistically significant).
614

Figure 4. The WRKY42-*APOLO* hub activates *RHD6* transcription in response to low temperatures, pro-moting root hair expansion

A. H3K27me3 deposition in WT vs. *wrky42* mutant 10-day-old seedlings grown at 23°C over *RHD6* and a WRKY42-independent *APOLO* target (*AZG2*; (27)). An immunoprecipitation (ChIP-qPCR) was performed using an anti-IgG antibody as a negative control.

B. *APOLO*-WRKY42 interaction *in vivo* demonstrated by RNA Immunoprecipitation (RIP)-qPCR in transi-ently transformed tobacco leaves and stably transformed Arabidopsis plants. In tobacco, WRKY42:GFP and *APOLO* were co-transformed under the control of the 35S constitutive promoter. In Arabidopsis, *35S:WRKY42:GFP* transformed plants were used to detect the interaction with endogenous *APOLO*. Anti-

IgG antibodies were used as a negative control. The RNA *PP2A* was considered as a RIP negative control
in Arabidopsis plants.
C. *APOLO* association to DNA of the *RHD6* locus by ChIRP-qPCR in WT and *wrky42* mutant plants. The
background level was determined using a set of probes against LacZ RNA.
D. WRKY42 immunoprecipitation over *RHD6* with the addition of increasing concentrations of *in vitro*
transcribed *APOLO* prior to cross-linking.
E. Proposed model for the action of *APOLO* in RH growth under low temperatures. In response to cold,
APOLO levels increase, recognizing the *RHD6* locus by sequence complementarity. Then, LHP1 is decoyed
away from chromatin and H3K27me3 deposition decreases in a process involving WRKY42 and leading to
the opening of the *RHD6* promoter region. Additionally, *APOLO* and WRKY42 jointly activate *RHD6* tran-
scription. RHD6 activates the transcription of downstream TF genes *RSL2* and *4* promoting RH cell expan-
sion.
In A, B, C and D, the error bar represents the SD of 3 biological replicates. The asterisk indicates that the
difference is significant (t test "*"<0.05, "**"<0.01).

639
Supplementary Figure 1: *RHD6* **and** *APOLO* **transcripts are enriched in root hair cells**

640
Uniform Manifold Approximation and Projection (UMAP) clustering map of Arabidopsis root single cell
641
RNA-seq ((31; http://wanglab.sippe.ac.cn/rootatlas/) data. *RHD6*, *APOLO* and the trichoblast cell marker
642
COBL9 are respectively indicated in red in each plot.
643

Supplementary Figure 2. Root hair expansion in response to cold is mediated by *RSL2* and *RSL4*

Quantification of RH length of plants WT, *rhd6-3, rsl1/4; rsl2; rsl4; rsl2/4*; *35S:RSL4* and *pEXP7:RSL4* at 22°C and 10°C. Each point is the mean of the length of the 10 longest RHs identified in a single root. Representative images of each genotype are shown below the graph. Asterisks indicate significant differences between WT and the corresponding genotype at the same temperature (one way ANOVA followed by a Tukey-Kramer test; "*"<0.05, "**"<0.01, "***"<0.001, NS: no statistically significant), unless indicated differently.

654
655
Supplementary Figure 3: Induction of root hair elongation by low temperatures requires epigenetic
Polycomb-mediated reprogramming
A. H3K27me3 deposition across the *RHD6* locus in WT, *clf* and *lhp1* mutants, revealed by ChIP-seq from
seedlings grown at 23°C (33).
B. H3K27me3 deposition measured by ChIP-qPCR over the *RHD6* locus in WT plants grown at 22°C treat-
ed or not for 24h at 10°C. The negative control was performed using an anti-IgG antibody.
C. *RHD6* transcript levels measured by RT-qPCR in roots of WT plants vs. *clf* mutants, expressed as the
ratio between plants treated or not for 24h at 10°C over the mock plants grown at 22°C.
D. Quantification of RH length of plants WT, *lhp1* and *clf* at 22°C and 10°C. Each point is the mean of the
length of the 10 longest RHs identified in a single root. Representative images of each genotype are
shown on the right. Asterisks indicate significant differences between WT and the corresponding geno-

type at the same temperature (one way ANOVA followed by a Tukey-Kramer test; "**"<0.01,
"***"<0.001).
In B and C, the error bar represents the SD of 3 biological replicates. The asterisk indicates that the dif-
ference is significant (t test p<0.05).

671
Supplementary Figure 4. Root hair expansion in response to cold seems to be related to nutrient avail-
ability
A. Quantification of RH length of WT plants at 22°C and 10°C at growing concentrations of Murashige and
Skoog (MS) culture media. Each point is the mean of the length of the 10 longest RHs identified in a sin-
gle root. Representative images of each genotype are shown on the right. Asterisks indicate significant
differences between WT cultivated in the same growth conditions at different temperatures (one way
ANOVA followed by a Tukey-Kramer test). NS stands for no statistically significant difference.
B. Quantification of RH length of WT plants expressed as the ratio between 10°C and 22°C at growing
concentrations of MS culture media, for given concentrations of agar. Each point is the mean of the
length of the 10 longest RHs identified in a single root. Representative images of each genotype are
shown below the graph.
Asterisks indicate significant differences between the ratio of WT cultivated at different temperatures in
different growth conditions (one way ANOVA followed by a Tukey-Kramer test; "*"<0.05, "**"<0.01,
"***"<0.001, NS: no statistically significant).

LEGENDS TO SUPPLEMENTARY TABLES

Supplementary Table 1: *APOLO* targets related to root hair growth

The list of *APOLO bona fide* targets was determined in (27). The search for root hairs-related genes was

performed based on (9, 11, 13, 16). LHP1 binding and H3K27me3 deposition was investigated in previ-

ously published datasets (33). W-boxes (TTGACY; (65)) present in the first 2500 bp upstream the ATG of

each gene was searched manually. The position of the W-box is given only for *RHD6*, directly recognized

by WRKY42 (**Figure 4B**).

Supplementary Table 2: Root hair-related genes deregulated in *35S:APOLO* seedlings

The list of deregulated genes was determined by RNA-Seq (27). The search for root hairs-related genes

was performed based on (9, 11, 13, 16). Genes are classified as directly (same as in **Supplementary Table**

1) or indirectly regulated by *APOLO*. *RHD6*, *RSL2* and *RSL4* are highlighted in red.

Supplementary Table 3: List of oligonucleotides used in this work

Supplementary Table 1

APOLO target	Gene name	LHP1 binding	H3K27me3 deposition	W-boxes in promoter	WRKY42 target (position)
AT1G66470	RHD6	YES	YES	1	-2521
AT1G21310	EXT3	YES	YES	4	
AT1G23720	EXT15	YES	YES	3	
AT1G26250	EXT18	YES	YES	0	
AT1G62440	LRX2	YES	YES	1	
AT2G04780	FLA7	NO	NO	2	
AT2G24980	EXT6	weak	weak	2	
AT2G26420	PIP5K3	NO	NO	2	
AT3G14310	PME3	NO	NO	0	
AT3G54580	EXT17	weak	NO	1	
AT4G08400	EXT7	weak	NO	2	
AT4G08410	EXT8	weak	NO	0	
AT4G13340	LRX3	weak	NO	3	
AT4G13390	EXT12	weak	weak	0	
AT4G40090	AGP3	YES	YES	2	
AT5G06630	EXT9	YES	YES	1	
AT5G06640	EXT10	YES	YES	0	

Supplementary Table 2

	gene	name	locus	sample 1	sample 2	status	value 1	value 2	log2(FC)	test stat	p value	q value	significant?
APOLO direct targets (Supplementary Table 1)	AT1G66470	RHD6	chr1:24795213-24796988	WT	35S:APOLO	OK	0,376783	1,05922	1,4912	2,64116	0,00015	0,00097	yes
	AT1G21310	EXT3	chr1:7453235-7455069	WT	35S:APOLO	OK	152,749	461,038	1,59372	6,83739	5,00E-05	0,00035	yes
	AT1G23720	EXT15	chr1:8388654-8391540	WT	35S:APOLO	OK	3,09641	19,245	2,63582	10,0477	0,00005	0,00035	yes
	AT1G26250	EXT18	chr1:9083837-9085464	WT	35S:APOLO	OK	3,22367	1,51056	-1,09362	-2,90259	0,00005	0,00035	yes
	AT1G62440	LRX2	chr1:23111817-23115508	WT	35S:APOLO	OK	0,292088	2,59795	3,1529	6,79781	5,00E-05	0,00035	yes
	AT2G04780	FLA7	chr2:1676741-1678455	WT	35S:APOLO	OK	7,18037	16,6557	1,21388	4,52231	0,00005	0,00035	yes
	AT2G24980	EXT6	chr2:10622452-10624132	WT	35S:APOLO	OK	0,837653	6,96907	3,05654	7,82993	5,00E-05	0,00035	yes
	AT2G26420	PIP5K3	chr2:11239433-11242239	WT	35S:APOLO	OK	0,077305	0,34491	2,15759	2,67767	0,0003	0,00181	yes
	AT3G14310	PME3	chr3:4771901-4775119	WT	35S:APOLO	OK	12,0768	25,3357	1,06894	4,63885	0,00005	0,00035	yes
	AT3G54580	EXT17	chr3:20200687-20203782	WT	35S:APOLO	OK	2,80301	22,185	2,98453	11,514	5,00E-05	0,00035	yes
	AT4G08400	EXT7	chr4:5323799-5325341	WT	35S:APOLO	OK	0,166086	2,09646	3,65795	5,75847	5,00E-05	0,00035	yes
	AT4G08410	EXT8	chr4:5339085-5341209	WT	35S:APOLO	OK	0,242031	2,59029	3,41985	6,63579	5,00E-05	0,00035	yes
	AT4G13340	LRX3	chr4:7758609-7761159	WT	35S:APOLO	OK	61,2495	139,492	1,18741	5,4	5,00E-05	0,00035	yes
	AT4G13390	EXT12	chr4:7783802-7785332	WT	35S:APOLO	OK	1,28902	9,4516	2,87429	7,90266	5,00E-05	0,00035	yes
	AT4G40090	AGP3	chr4:18580837-18581585	WT	35S:APOLO	OK	0,465013	6,11837	3,71781	6,1024	0,00005	0,00035	yes
	AT5G06630	EXT9	chr5:2036360-2037843	WT	35S:APOLO	OK	0,47843	4,28202	3,16191	6,69628	5,00E-05	0,00035	yes
	AT5G06640	EXT10	chr5:2039858-2041928	WT	35S:APOLO	OK	0,704936	6,53659	3,21297	8,39508	5,00E-05	0,00035	yes
RH-related genes **NOT** identified as direct *APOLO* targets	AT1G02900	RALF1	chr1:653764-654343	WT	35S:APOLO	OK	6,43265	11,7847	0,87343	2,3996	0,0001	0,00067	yes
	AT1G12040	LRX1	chr1:4070123-4072567	WT	35S:APOLO	OK	0,254515	2,57012	3,33601	6,95711	5,00E-05	0,00035	yes
	AT1G12560	EXPA7	chr1:4276518-4277899	WT	35S:APOLO	OK	0,373865	2,37379	2,6666	4,34024	5,00E-05	0,00035	yes
	AT1G12950	RSH2	chr1:4419769-4422638	WT	35S:APOLO	OK	0,312439	1,50467	2,2678	4,34933	5,00E-05	0,00035	yes
	AT1G16440	RSH3	chr1:5615814-5617672	WT	35S:APOLO	OK	0,16653	0,405508	1,28395	1,77649	0,00455	0,01897	yes
	AT1G27740	RSL4	chr1:9654687-9656089	WT	35S:APOLO	OK	0,105364	0,682308	2,69504	3,06432	0,0005	0,00284	yes
	AT1G32640	MYC2	chr1:11798809-11800988	WT	35S:APOLO	OK	22,203	52,6239	1,24496	5,66997	5,00E-05	0,00035	yes
	AT2G35150	PHI-1-like	chr2:14817046-14818211	WT	35S:APOLO	OK	0,264466	1,09189	2,04567	3,08494	5,00E-05	0,00035	yes
	AT1G54970	PRP1	chr1:20504999-20506488	WT	35S:APOLO	OK	1,07342	8,36667	2,96244	7,31633	5,00E-05	0,00035	yes
	AT1G55330	AGP21	chr1:20648464-20648891	WT	35S:APOLO	OK	36,0237	97,1117	1,4307	5,34932	5,00E-05	0,00035	yes
	AT1G63450	RHS8	chr1:23531408-23534569	WT	35S:APOLO	OK	0,03899	0,331918	3,08967	3,72534	5,00E-05	0,00035	yes
	AT1G70460	PERK13	chr1:26555845-26559284	WT	35S:APOLO	OK	0,258225	1,21635	2,23585	4,60468	5,00E-05	0,00035	yes
	AT2G25260	HPAT2	chr2:10755523-10757772	WT	35S:APOLO	OK	0,529251	1,33354	1,33324	2,44251	5,00E-05	0,00035	yes
	AT2G43150	EXT21	chr2:17945769-17947032	WT	35S:APOLO	OK	52,0807	114,204	1,1328	5,20396	5,00E-05	0,00035	yes
	AT2G47540	SHV2	chr2:19505859-19506598	WT	35S:APOLO	OK	1,0193	4,80441	2,23678	4,04873	5,00E-05	0,00035	yes
	AT3G07340	bHLH62 TF	chr3:2340898-2343470	WT	35S:APOLO	OK	3,61588	7,57866	1,06759	3,73419	5,00E-05	0,00035	yes
	AT3G10710	RHS12	chr3:3352288-3354237	WT	35S:APOLO	OK	0,11403	1,03108	3,17667	4,47911	5,00E-05	0,00035	yes
	AT3G21180	ACA9	chr3:7425518-7432200	WT	35S:APOLO	OK	0,058342	0,415952	2,83382	4,24417	5,00E-05	0,00035	yes
	AT3G28550	EXT16	chr3:10688322-10703976	WT	35S:APOLO	OK	3,33615	28,0079	3,06958	5,7735	5,00E-05	0,00035	yes
	AT3G43960	RDL3	chr3:15774054-15775657	WT	35S:APOLO	OK	1,63093	3,72675	1,19223	3,00585	5,00E-05	0,00035	yes
	AT3G48340	CEP2	chr3:17897738-17899208	WT	35S:APOLO	OK	1,69325	11,8683	2,80925	7,74503	5,00E-05	0,00035	yes
	AT3G54580	EXT17	chr3:20200687-20203782	WT	35S:APOLO	OK	2,80301	22,185	2,98453	11,514	5,00E-05	0,00035	yes
	AT3G60330	HA7	chr3:22297406-22303673	WT	35S:APOLO	OK	0,75475	1,83265	1,27986	3,54359	5,00E-05	0,00035	yes
	AT3G62680	PRP3	chr3:23182721-23183994	WT	35S:APOLO	OK	1,90231	17,1835	3,1752	9,02884	5,00E-05	0,00035	yes
	AT4G02270	RHS13	chr4:992174-993038	WT	35S:APOLO	OK	0,956108	7,16393	2,90551	5,80735	5,00E-05	0,00035	yes
	AT4G17215	Pollen Ole e 1 allergen and extensin	chr4:9655836-9656577	WT	35S:APOLO	OK	1,17841	4,76508	2,01566	3,84981	5,00E-05	0,00035	yes
	AT4G22080	RHS14	chr4:11700554-11702666	WT	35S:APOLO	OK	0,059309	0,757044	3,67406	3,68273	0,00025	0,00154	yes
	AT4G25400	bHLH TF	chr4:12981294-12982468	WT	35S:APOLO	OK	1,35187	4,32603	1,67809	3,18257	5,00E-05	0,00035	yes
	AT4G29180	RHS16	chr4:14385592-14389689	WT	35S:APOLO	OK	0,099263	0,521194	2,39249	3,82728	5,00E-05	0,00035	yes
	AT4G29930	bHLH TF	chr4:14644008-14647591	WT	35S:APOLO	OK	2,30654	7,32455	1,66701	4,31108	5,00E-05	0,00035	yes
	AT4G33880	RSL2	chr4:16239362-16241137	WT	35S:APOLO	OK	0,038579	0,306499	2,99001	2,43441	0,0083	0,03137	yes
	AT4G36930	SPT	chr4:17414126-17416294	WT	35S:APOLO	OK	1,36121	2,40537	0,82137	2,05934	0,00055	0,00309	yes
	AT5G05500	MOP10	chr5:1629666-1630396	WT	35S:APOLO	OK	0,56784	5,15021	3,18108	5,39453	5,00E-05	0,00035	yes
	AT5G19560	ROPGEF10	chr5:6603290-6606448	WT	35S:APOLO	OK	0,058391	0,406499	2,79944	2,91293	0,0004	0,00233	yes
	AT5G22410	RHS18	chr5:7426313-7427964	WT	35S:APOLO	OK	0,342718	3,78832	3,46647	6,35329	5,00E-05	0,00035	yes
	AT5G35190	EXT13	chr5:13434180-13435167	WT	35S:APOLO	OK	1,52769	15,7388	3,3649	8,58919	5,00E-05	0,00035	yes
	AT5G41315	GL3	chr5:16529454-16532879	WT	35S:APOLO	OK	0,235867	0,44781	0,92491	1,51906	0,0112	0,04019	yes
	AT5G51060	RHD2	chr5:20752999-20762431	WT	35S:APOLO	OK	0,213913	1,38566	2,69548	3,1665	0,0001	0,00067	yes
	AT5G51790	bHLH120 TF	chr5:21039510-21041446	WT	35S:APOLO	OK	2,80331	7,68882	1,45563	2,87659	5,00E-05	0,00035	yes
	AT5G54380	THE1	chr5:22077059-22080338	WT	35S:APOLO	OK	19,729	40,4626	1,03627	4,68266	5,00E-05	0,00035	yes
	AT5G58010	LRL3	chr5:23483544-23484889	WT	35S:APOLO	OK	0,163107	1,32569	3,02286	3,78898	5,00E-05	0,00035	yes
	AT5G67400	RHS19	chr5:26894855-26896488	WT	35S:APOLO	OK	0,319731	2,96103	3,21117	5,60471	5,00E-05	0,00035	yes

Supplementary Table 3

			3C-qPCR	Experiment
AT1G66470 *RHD6*		F	TCCCGATTCTTCCCTTAAAAA	3C
		R	AAAGCAAACCTTGGCTGCTA	
		F	AATGACCATCCCAACGAGAC	ChIRP, RT-qPCR
		R	CATAAGCTGCATCTGGCTGA	
		F	GGATTAGAAAGACCAGCTGCTTTAG	ChIP
		R	CCAGTTTGTGAACGACGACA	
AT2G34655 *APOLO*		F	CTTCGAGGCGCTAAACAATC	RIP, RT-qPCR
		R	ACAGCGGTGCCACCTATTAC	
AT5G50300 *AZG2*		F	GGGAGTTGCAAAATGGCTTA	ChIP
		R	CCGACCATGTGAAAAATTCC	
AT4G04450 *WRKY42*		F	CCGCTAAACGCTGGTGAGA	RT-qPCR
		R	CGGCAAAGTTGGGATTGG	
		SALK_049063_LP2	AAGCACGCTTCGATATCTGAG	genotyping
		SALK_049063_RP2	AGTTGCGACAATCAATGGATC	
		SALK_049063_LP4	TTTGTGCGTCTGTTACGTACG	
		SALK_049063_RP4	GTACTTGCTTGCGAACAGGAC	
		F	CACCATGTTTCGTTTTCCGGTAAGTC	cloning
		R	TTGCCTATTGTCAACGTTGCTC	
AT1G13320 Housekeeping 1		F	TAACGTGGCCAAAATGATGC	RIP, RT-qPCR
		R	GTTCTCCACAACCGCTTGGT	

ChIRP 3'BIO probes

ODD probes

number	sequence
APOLO 1	aaccagccaatgaacagatg
APOLO 3	gacaagtcacacctacactc
APOLO 5	agttccaaggaatccatacc
APOLO 7	gacagcggtgccacctatta
APOLO 9	ttacaacaagccactccgta
APOLO 11	accaaaacaaccaaaatttt
APOLO 13	ccttacaaacagagcaaagt
APOLO 15	ggaagcaaagcccaaaggaa
APOLO 17	caagtaaccccagaaaacta
APOLO 19	gattccggtgaaatacaagg

EVEN probes

number	sequence
APOLO 2	gattgtttagcgcctcgaag
APOLO 4	cgagaagaactaggccaaag
APOLO 6	tgaagactaccttacataga
APOLO 8	acaaggaacttccaaccaaa
APOLO 10	caacatctcgtcaaccacat
APOLO 12	cgaaactaaaacaaagaagc
APOLO 14	ggaaataacaaggcaaaaca
APOLO 16	ccgacgattaaaaggataat
APOLO 18	gaaatacaaagccggcggtt
APOLO 20	tcgtctgaaagtttattata

EVEN LacZ probes

number	sequence
LacZ 1	ccagtgaatccgtaatcatg
LacZ 5	aatgtgagcgagtaacaacc
LacZ 7	aataattcgcgtctggcctt
LacZ 9	aattcagacggcaaacgact
LacZ 11	atcttccagataactgccgt
LacZ 13	gctgatttgtgtagtcggtt
LacZ 15	aactgttacccgtaggtagt
LacZ 17	tttcgacgttcagacgtagt
LacZ 19	accattttcaatccgcacct
LacZ 21	ttcatcagcaggatatcctg