The lncRNA *APOLO* interacts with the transcription factor WRKY42 to trigger root hair cell expansion in

response to cold

Michaël Moison[*1], Javier Martínez Pacheco[*2], Leandro Lucero[*1], Camille Fonouni-Farde[1], Johan

Rodríguez-Melo[3], Natanael Mansilla[1], Aurélie Christ[4], Jérémie Bazin[4]; Moussa Benhamed[4]; Fernando

Ibañez[3], Martin Crespi[4], José M. Estevez[†,2,5] and Federico Ariel[†1]

[1] Instituto de Agrobiotecnología del Litoral, Universidad Nacional del Litoral, CONICET, FBCB/FHUC,

Centro Científico Tecnológico CONICET Santa Fe, Colectora Ruta Nacional No 168 km. 0, Paraje El Pozo,

Santa Fe 3000, Argentina.

[2] Fundación Instituto Leloir and IIBBA-CONICET, Av. Patricias Argentinas 435, Buenos Aires CP C1405BWE,

Argentina.

[3] Instituto de Investigaciones Agrobiotecnológicas, CONICET, Universidad Nacional de Río Cuarto, Río

Cuarto 5800, Argentina.

[4] Institute of Plant Sciences Paris-Saclay (IPS2), CNRS, INRA, University Paris-Saclay and University of Paris

Bâtiment 630, 91192 Gif sur Yvette, France.

[5] Centro de Biotecnología Vegetal (CBV), Facultad de Ciencias de la Vida (FCsV), Universidad Andres Bello,

Santiago, Chile and Millennium Institute for Integrative Biology (iBio), Santiago, Chile.

[*]These authors contributed equally to this work

[†] Correspondence to: Federico Ariel (fariel@santafe-conicet.gov.ar) and José M. Estevez

(jestevez@leloir.org.ar)

Text Word count (without Abstract and References)**:** 6,287

ABSTRACT

Plant long noncoding RNAs (lncRNAs) have emerged as important regulators of chromatin dynamics,

impacting on transcriptional programs leading to different developmental outputs. The lncRNA *AUXIN*

REGULATED PROMOTER LOOP (*APOLO*) directly recognizes multiple independent loci across the

Arabidopsis genome and modulates their three-dimensional chromatin conformation, leading to

transcriptional shifts. Here, we show that *APOLO* recognizes the locus encoding the root hair (RH) master

regulator ROOT HAIR DEFECTIVE 6 (RHD6) and controls *RHD6* transcriptional activity leading to cold-

enhanced RH elongation. Furthermore, we demonstrate that *APOLO* interacts with the transcription

factor WRKY42 and modulates its binding to the *RHD6* promoter. WRKY42 is required for the activation

of *RHD6* by low temperatures and *WRKY42* deregulation impairs cold-induced RH expansion. Collectively,

our results indicate that a novel ribonucleoprotein complex involving *APOLO* and WRKY42 forms a

regulatory hub which activates *RHD6* by shaping its epigenetic environment and integrates signals

governing RH growth and development.

Words: 148

Key words: root hairs; cell expansion; long noncoding RNAs; *APOLO*; transcription factors; RHD6;

WRKY42; cold temperature

INTRODUCTION

Root hairs (RHs) are single cell projections developed from specialized epidermal trichoblast cells able to increase their size several hundred times in a polar manner to reach and promote the uptake of water-soluble nutrients, interact with soil microorganisms and support anchor to the plant. The specification of epidermal cells into RHs is a complex process whose underlying mechanisms are partially understood. In *Arabidopsis thaliana*, RH cell fate is controlled by a developmental program involving a complex of transcription factors (TFs) promoting the expression of the homeodomain protein GLABRA 2 (GL2) (Ryu et al., 2005; Song et al., 2011; Schiefelbein et al., 2014; Balcerowicz et al., 2015). GL2 blocks RH development by inhibiting the transcription of the master regulator *ROOT HAIR DEFECTIVE 6* (*RHD6*) (Lin et al., 2015). In the cells that differentiate into RHs (known as trichoblasts), a second TF complex suppresses *GL2* expression (Schiefelbein et al., 2014), forcing the cells to enter the RH cell fate program via the concomitant activation of RHD6 along with downstream TFs (Menand et al., 2007; Pires et al., 2013). Briefly, RHD6 together with its homolog RSL1 (ROOT HAIR DEFECTIVE 6 LIKE 1) induce the expression of TFs from the bHLH family, including RSL2 (ROOT HAIR DEFECTIVE 6 LIKE 2) and RSL4 (ROOT HAIR DEFECTIVE 6 LIKE 4), ultimately triggering the differentiation of the RHs and their subsequent polarized tip-growth (Karas et al., 2009; Yi et al., 2010; Bruex et al., 2012). In addition, it was proposed that RSL4 controls the expression of a small subset of nearly 125 genes (Won et al., 2009; Yi et al., 2010; Datta et al., 2015; Vijayakumar et al., 2016), including several cell wall extensins (EXTs) (Ringli, 2010; Velasquez et al., 2011) sufficient to promote RH growth (Hwang et al., 2017).

RH expansion is regulated both by cell-intrinsic factors (e.g. endogenous phytohormones such as auxin) and external environmental signals (e.g. phosphate (Pi) availability in the soil) (Mangano et al., 2017; Bhosale et al., 2018). Pi starvation is one of the key environmental factors promoting rapid RH growth (Yi et al., 2010; Datta et al., 2015; Vijayakumar et al., 2016). In *Arabidopsis*, it triggers *RSL4* expression via an enhanced auxin production, activating downstream effector genes mediating cell growth (Yi et al., 2010; Datta et al., 2015; Mangano et al., 2017; Marzol et al., 2017; Bhosale et al., 2018). Accordingly, several auxin-related TFs have been implicated in Pi-starvation signaling in roots, including WRKY proteins that control the expression of the Pi transporter families Pi-permease PHO1 and PHOSPHATE TRANSPORTER (PHT) (Devaiah et al., 2007; Chen et al., 2009; Wang et al., 2014; Su et al., 2015). Under Pi-sufficient conditions, WRKY6 and WRKY42 bind to W-boxes of the *PHO1* promoter and suppress its expression. During Pi starvation, WRKY42 is degraded by the 26S proteasome pathway, resulting in the activation of *PHO1* transcription (Chen et al., 2009; Su et al., 2015). In addition, WRKY42 functions as

a positive regulator of *PHT1;1*, by binding to its promoter under Pi-sufficient condition (Su et al., 2015).

Overall, WRKY42 is part of the components activating root early-responses to Pi starvation, although its

role in controlling RH growth remains unexplored.

In recent years, plant long noncoding RNAs (lncRNAs) have emerged as important regulators of

gene expression, and several among them, have been functionally linked to Pi homeostasis. For instance,

the lncRNA *INDUCED BY PHOSPHATE STARVATION 1* (*IPS1*) can sequester the Pi starvation-induced mi-

croRNA miR-399, attenuating miR-399-mediated repression of *PHO2*, a gene encoding an E3 ligase af-

fecting Pi uptake (Franco-Zorrilla et al., 2007). In addition, the *cis*-natural antisense (*cis*-NAT) transcript

PHO1;2, induced under Pi deficiency, was shown to promote the translation of the *PHO1;2* mRNA in-

volved in Pi loading into the xylem. The expression of this *cis*-NAT is associated with the transport of the

sense–antisense RNA pair toward the polysomes (Jabnoune et al., 2013). More recently, it was shown

that the lncRNA *AUXIN REGULATED PROMOTER LOOP* (*APOLO*) recognizes multiple spatially independent

genes by sequence complementarity and DNA-RNA duplex formation, known as R-loops. Upon recogni-

tion, *APOLO* shapes the three-dimensional (3D) conformation of its target regions by decoying the

Polycomb Repressive Complex 1 (PRC1) component LIKE HETEROCHROMATIN PROTEIN 1 (LHP1), thereby

regulating their transcription (Ariel et al., 2014; Ariel et al., 2020).

Here, we show that the lncRNA *APOLO* directly regulates a subset of genes involved in RH devel-

opment, including the master regulator of RH initiation *RHD6*. *APOLO* activates *RHD6* transcription by

modulating the formation of a local chromatin loop encompassing its promoter region, an epigenetic

regulatory mechanism likely involving PRC1 and PRC2 components. Furthermore, we found that *APOLO*

interacts with the TF WRKY42, forming a new hub that regulates *RHD6* to induce RH growth in response

to low temperatures.

RESULTS

***APOLO* regulates root hair cell elongation in response to low temperatures**

Based on Chromatin Isolation by RNA Purification (ChIRP-Seq) performed in wild-type (WT)

plants, it was previously reported that the lncRNA *APOLO* recognizes a subset of independent loci en-

riched in categories related to cell wall composition and organization (Ariel et al., 2020). A closer look at

APOLO bona fide targets allowed us to identify seventeen genes involved in RH growth and expansion

(**Supplementary Table 1**), a process dependent on cell wall remodeling molecules, including EXTs and

EXT-related proteins (Ringli, 2010; Lamport et al., 2011; Velasquez et al., 2011; Velasquez et al., 2015;

Marzol et al., 2018). Interestingly, according to single-cell RNA-seq datasets (Zhang et al., 2019) *APOLO*

transcripts are enriched in RH cells (**Supplementary Figure 1A**). Notably, sixteen *APOLO* direct targets

were upregulated upon *APOLO* over-expression (**Supplementary Table 2**). Furthermore, 52 additional RH-

related genes were upregulated in *35S:APOLO* seedlings (**Supplementary Table 2**) (Ariel et al., 2020).

Among them, the RH central TFs *RHD6* (as a direct target), *RSL2* and *RSL4* (as indirectly regulated) were

induced upon *APOLO* over-expression.

It was reported that the *APOLO* locus is targeted by the RNA-polymerase Pol V and silenced by

RNA-directed DNA Methylation (RdDM,(Ariel et al., 2014)). A search in a small RNA-Seq performed in WT

roots subjected to different temperature treatments (Gyula et al., 2018) revealed that RdDM-related 24nt

siRNA accumulation over the *APOLO* locus is less abundant at low temperatures (**Supplementary Figure

1B**), suggesting that *APOLO* transcription is regulated by cold. Accordingly, we found that *APOLO* tran-

scriptional accumulation increases in roots after 24h at 10°C (**Figure 1A**). An analysis of the promoter

activity of the 5.2kb region upstream *APOLO* (Ariel et al., 2020) directing the expression of a *GFP* reporter

gene, additionally revealed a higher transcriptional activity at low temperatures in the RHs (**Figure 1B**).

Strikingly, we observed that two RNAi-*APOLO* (repression of approximately 90%; Ariel et al., 2014) and

35S:APOLO independent lines (over expression of 30-fold and 60-fold, respectively; Ariel et al., 2020)

exhibit a basal increase of RH length at 22°C, and uncovered a strong induction of RH elongation in WT

and RNAi-*APOLO* at 10°C, in contrast to *35S:APOLO* lines (**Figure 1C**). Accordingly, *RHD6* is induced in re-

sponse to cold in WT roots, whereas RNAi-*APOLO* and *35S:APOLO* roots display higher *RHD6* basal levels

than the WT (**Figure 1D**). Notably, *RHD6* transcript levels are not further induced by cold in *35S:APOLO*

roots (**Figure 1D**), in agreement with the RH phenotype (**Figure 1C**). Collectively, our findings suggest that

APOLO participates in the induction of cold-mediated RH elongation and a deregulation of *APOLO* tran-

script levels can impact RH growth.

Previous studies pointed out a key role of RHD6 (together with RSL1) in RH development, which

is mediated by RSL4 and RSL2 as downstream regulators of RH cell elongation (Menand et al., 2007; Pires

et al., 2013). Notably, *RHD6*, *RSL2* and *RSL4* transcript levels were upregulated in *35S:APOLO* seedlings

(**Supplementary Table 2**; (Ariel et al., 2020)), although only *RHD6* was identified as a direct target of

APOLO (Ariel et al, 2020). In agreement with *RHD6* transcriptional behavior (**Figure 1D**), *RSL2* and *RSL4*

basal transcript levels are higher in RNAi-*APOLO* and *35S:APOLO* compared to WT. Interestingly, *RSL2*

and *RSL4* are induced by cold in WT and RNAi-*APOLO*, but not in *35S:APOLO* roots (**Supplementary Fig-

ure 2A**), suggesting that low temperatures can activate these two genes still in the absence of *APOLO*

and bypassing RHD6 for RH expansion (**Figure 1C**). Thus, we assessed if these TFs were also controlling

the promotion of RH growth by low temperatures. To this end, we tested how *rhd6/rsl1/rsl4* and *rsl2*,

rsl4 and double mutant plants *rsl2/rsl4* respond to low temperatures in comparison with control condi-

tions **(Supplementary Figure 2A)**. The *rsl2* mutant was highly responsive to low temperatures in a similar

manner to WT while *rsl4* was impaired in the response to cold. The double mutant *rsl2/rsl4* and the triple

mutant *rhd6/rsl1/rsl4* did not develop RHs in either of the two conditions. In addition, constitutive ex-

pression of *RSL4* (*35S:RSL4*) as well as its expression under the control of the RH specific *EXPANSIN7*

promoter (*EXP7p:RSL4*) boosted basal RH growth without further enhancement in response to cold

(**Supplementary Figure 2B**). These results demonstrate that RSL4 is largely required for RHD6-dependent

activation of RH growth at low temperature, and RSL2 might participate to a lower extent.

Nutrient unavailability is known to activate RH expansion through a transcriptional reprogram-

ming governed by RHD6 and downstream TFs. The quantification of RH growth of WT plants in response

to increasing concentrations of nutrients (0.5X to 2.0X MS (Murashige and Skoog) medium) indicates that

high concentrations impair RH growth triggered by low temperatures **(Supplementary Figure 3A)**. In a

similar way, an increase in agar concentration in the MS medium (from 0.8% to 2.5%), which likely re-

strains nutrient mobility (Singha, S., Townsend, E.C. and Oberly, 1985; Nonami and Boyer, 1989;

Ghashghaie et al., 1991; Buah et al., 1999) blocks cold-induced RH expansion **(Supplementary Figure 3B)**.

Altogether, these observations suggest that low temperatures restrict nutrient mobility and availability in

the culture medium, leading to the promotion of polar RH growth.

APOLO directly modulates the three-dimensional chromatin conformation of the root hair specific lo-

cus RHD6

Among *APOLO* targets involved in RH development, we found the master regulator of RH initia-

tion *RHD6* (Menand et al., 2007; Pires et al., 2013). The epigenetic profile of the *RHD6* locus corresponds

to typical *APOLO* targets (**Figure 2A**, (Ariel et al., 2020)), including H3K27me3 deposition (track 1), LHP1

recognition (track 2, chromatin immunoprecipitation (ChIP)-Seq, (Veluchamy et al., 2016)), and *APOLO*

binding regions (tracks 3 to 5, chromatin isolation by RNA purification (ChIRP)-Seq, (Ariel et al., 2020)). A

GAAGAA box, shown to be important for *APOLO* target recognition (Ariel et al., 2020) is located in the

RHD6 locus and coincides with *APOLO* binding site. In addition, a peak of DNA-RNA hybrid

immunoprecipitation (DRIP)-Seq from root samples indicates the presence of an R-loop coinciding with

APOLO recognition sites over *RHD6* (tracks 6 to 8, (Xu et al., 2020)).

Remarkably, *APOLO* recognition and R-loop formation are also detectable over *RHD6* neighbor

gene, located 3.2 kb upstream *RHD6* transcription start site (**Figure 2A**). According to *Dpn*II Hi-C datasets

from Arabidopsis seedlings (Liu et al., 2016), a chromatin loop encompassing the intergenic region up-

stream *RHD6* was detected (**Figure 2B**), and coincides with *APOLO* binding-sites (**Figure 2A**, ChIRP-Seq).

By performing a ChIRP-qPCR with two independent sets of biotinylated probes to purify *APOLO* (ODD and

EVEN; (Ariel et al., 2020)) and one additional set used as a negative control (LacZ), we confirmed that

APOLO RNA–*RHD6* DNA interaction occurs in WT and is lost in *APOLO* knockdown (RNAi) seedlings ((Ariel

et al., 2014); **Figure 2C**). In addition, the quantification of relative *RHD6* loop formation in RNAi-*APOLO*

and *35S:APOLO* (Ariel et al., 2020) seedlings, revealed impaired loop formation in both lines (**Figure 2D**),

hinting at a stoichiometric requirement of *APOLO* for *RHD6* chromatin loop formation. Chromatin loop

formation (**Figure 2D**) is in agreement with *RHD6* basal levels in *35S:APOLO* and RNAi-*APOLO* lines (**Figure**

1D), suggesting that the chromatin loop including *RHD6* promoter region precludes transcription. Alto-

gether, our results indicate that *APOLO* lncRNA directly regulates *RHD6* transcriptional activity by fine-

tuning local chromatin 3D conformation.

It was previously reported that PRC2 actively participates in the regulation of RH growth (Ikeuchi

et al., 2015) and that the *RHD6* locus exhibits H3K27me3 deposition and LHP1 recognition (**Figure 2A;**

(Veluchamy et al., 2016)). Considering that the lncRNA *APOLO* interacts with the PRC1 component LHP1

in vivo (Ariel et al., 2014; Ariel et al., 2020), we decided to explore the role of PRC1 and PRC2 in *APOLO*-

mediated *RHD6* activation at low temperatures. At 22°C, *RHD6* suffers a reduction of H3K27me3 in the

PRC2 mutant *curly leaf* (*clf*), in contrast to the PRC1 mutant *lhp1* (**Supplementary Figure 4A;** (Veluchamy

et al., 2016)). Interestingly, we observed that H3K27me3 deposition and LHP1 binding diminish in WT

roots treated for 24h at 10°C compared to 22°C (**Supplementary Figure 4B**), consistent with the induction

of *RHD6* in response to cold (**Figure 2E**). Moreover, *lhp1* and *clf* mutants exhibit a basal decrease of RH

length together with a slight decrease of cold-induced RH elongation in *lhp1*, and a strong decrease of

cold-induced RH elongation in *clf* (**Supplementary Figure 4D**). Consistently, although the decrease in

H3K27me3 deposition results in higher basal transcript levels of *RHD6* in the *clf* background, *RHD6* tran-

scriptional activation by cold is abolished in the *clf* and *lhp1* mutants (**Supplementary Figure 4C**), hinting

at an important role of chromatin rearrangement for *RHD6* activation in response to cold.

APOLO* interacts with the transcription factor WRKY42 to coordinate the activation of *RHD6

In order to uncover novel actors involved in cold-induced transcriptional regulation of RH

growth, we aimed at identifying *APOLO* protein partners. To this end, we performed an exploratory

ChIRP on WT seedlings using two independent set of biotinylated probes to purify *APOLO* (ODD and

EVEN; (Ariel et al., 2014; Ariel et al., 2020)), and one additional set used as a negative control (against

LacZ), as recently described (Rigo et al., 2020). Co-purified proteins were precipitated and analyzed by

mass spectrometry. Among the potential *APOLO* partners (i.e. identified with at least two hits in ODD

and EVEN samples, but absent in LacZ-ChIRP), we found the WRKY42 protein, a TF involved in the re-

sponse to Pi starvation (Su et al., 2015), an environmental condition that promotes RH cell expansion

(Bhosale, *et al.*, 2018) in a similar manner to low temperatures. Thus, *APOLO*-WRKY42 interaction was

validated by RNA immuno-precipitation (RIP-qPCR) in tobacco leaves and in *Arabidopsis* plants transitory

or stably transformed with 35S:*WRKY42:GFP*, respectively (**Figure 3A**). Interestingly, according to the

Arabidopsis eFP Browser (Waese et al., 2017), *WRKY42* is induced in roots when seedlings are subjected

to 4°C for 24h (**Figure 3B**). At 10°C, we observed that *WRKY42* transcriptional accumulation augments

significantly in roots (**Figure 3C**). Notably, 13 out of the 17 *APOLO* targets contained between 1 and 4

canonical WRKY TF binding sites (W-box) in their promoters, including *RHD6* (**Supplementary Table 1**).

By using a *35S:WRKY42:GFP* line, we determined that WRKY42 can directly bind to the promoter region

of *RHD6* (**Figure 3D**). Accordingly, the over-expression of *WRKY42* (*35S:WRKY42:GFP* line) led to a basal

increase of *RHD6* levels (**Figure 3E**) and RH elongation (**Figure 3F**) at ambient temperature, mimicking the

effect of cold. On the contrary, cold-mediated induction of *RHD6* is abolished in the *wrky42* mutant ((Su

et al., 2015); **Figure 3E**), which consistently exhibits shorter RHs at 22°C and almost no RH elongation at

low temperatures (**Figure 3F**). Taken together, these results suggest that the *APOLO*-interacting TF

WRKY42 is an important regulator of *RHD6*-mediated RH growth in response to cold.

We thus wondered to what extent WRKY42 regulates the epigenetic landscape of the *RHD6* lo-

cus. We first observed that H3K27me3 deposition over *RHD6* is significantly augmented in the *wrky42*

mutant background, in contrast to *AZG2*, an *APOLO* target non-related to WRKY42 (**Figure 4A**), con-

sistent with reduced *RHD6* basal levels reported in *wrky42* (**Figure 3E**). Therefore, we evaluated the mu-

tual contribution of *APOLO* and WRKY42 to their respective recognition of the *RHD6* locus. *APOLO* ChIRP-

qPCR in the WT and *wrky42* mutant (**Figure 4B**) revealed similar binding to *RHD6*, indicating that WRKY42

does not participate in *APOLO*-target recognition. Reciprocally, we assessed the control of *APOLO* over

WRKY42 recognition of the *RHD6* locus. To this end, we transformed transiently *A. thaliana* leaves of WT,

35S:APOLO and RNAi-*APOLO* plants with the construct *35S:WRKY42:GFP*. Observation using confocal

microscopy indicated that WRKY42:GFP is localized in the nucleus in the three genetic backgrounds

(**Supplementary Figure 5**). Remarkably, WRKY42:GFP ChIP-qPCR revealed that WRKY42 binding to *RHD6*

promoter is impaired both in *35S:APOLO* and RNAi-*APOLO* plants compared to WT (**Figure 4C**), hinting at

a stoichiometric role of *APOLO* in the modulation of TF-chromatin interaction. To further confirm our

observations, chromatin was extracted from *35S:WRKY42:GFP* seedlings and increasing amounts of *in*

vitro transcribed *APOLO* were added before cross-link and regular WRKY42 ChIP over *RHD6*. Strikingly,

increasing concentrations of *APOLO* gradually decoy WRKY42 away from the *RHD6* locus (**Figure 4D**),

further supporting the hypothesis of a stoichiometric regulation of *APOLO* over the activity of its partner

TF.

Collectively, our results indicate that the regulation of *RHD6* expression in response to cold

depends on Polycomb-dependent H3K27me3 dynamic deposition. The WRKY42-*APOLO* complex

modulates the epigenetic environment of *RHD6*, activating its transcription and promoting RH growth at

low temperatures. *RHD6* activation further triggers the expression of *RSL2* and *RSL4* that control the

transcriptional RH program inducing cell expansion in response to cold (**Figure 4E**).

DISCUSSION

Cell fate determination in the epidermis has been extensively studied (Ryu et al., 2005; Song et

al., 2011; Schiefelbein et al., 2014; Balcerowicz et al., 2015). Once trichoblast cells differentiate in the

root epidermis, RHs develop as fast polar growing protuberances in response to endogenous and envi-

ronmental signals (Menand et al., 2007; Yi et al., 2010; Pires et al., 2013; Marzol et al., 2017). RHs are

one of the main entry points in the roots for water-soluble macronutrients, such as Pi and nitrates. Pi is

an essential element for plant growth and development, and the availability of this macronutrient is a

factor limiting plant productivity. In Arabidopsis roots, low Pi in the soil triggers auxin synthesis and

transport, enhancing RH elongation to promote Pi uptake (Bhosale et al., 2018). Thus, auxin mediates

low Pi-induced promotion of RH cell expansion. Under low soil Pi, auxin synthesis is enhanced specifically

in the root cap (Stepanova et al., 2008) and transported (mostly by AUX1, PIN2, and PGP4) from the apex

to the differentiation zone, specifically leading to an increase of auxin levels in trichoblasts (Jones et al.,

2009; Bhosale et al., 2018; Wang et al., 2020). In response to the high-auxin microenvironment, RHs pro-

trude from the root epidermis controlled by RHD6 and RSL1(Menand et al., 2007; Pires et al., 2013). High

levels of auxin in trichoblasts trigger a signaling cascade mediated by TIR1-ARF19 (and possibly also

ARF7) which directly induces the expression of *RSL4* (and likely of *RSL2*) and promote RH elongation (Yi et

al., 2010; Mangano et al., 2017; Mangano et al., 2018b; Bhosale et al., 2018). ARF7 and ARF19 also acti-

vate other RH genes independently of RSL4 (Schoenaers et al., 2018). Interestingly, our results indicate

that the lncRNA *APOLO* participates in the response to low temperatures. *APOLO* is directly activated by

ARF7 and regulates the transcriptional activity of its neighboring gene *PINOID* (*PID*) by shaping local 3D

chromatin conformation (Ariel et al., 2014; Ariel et al., 2020). *PID* encodes a kinase responsible for accu-

rate auxin polar transport by localizing PIN2 in the root cell membrane (Friml et al., 2004). More recently,

it was shown that *APOLO* can recognize a subset of distant genes across the Arabidopsis genome, most of them being related to auxin synthesis and signalling (Ariel et al., 2020). In this work, we demonstrate that a group of RH related genes is directly regulated by *APOLO* in response to cold, including the RH master regulator *RHD6*. Interestingly, *RSL2* and *RSL4* are still activated by cold in RNAi-*APOLO* roots in contrast to *RHD6* (**Figure 1C** and **Supplementary Figure 2A**), suggesting that additional yet uncovered factors, which might include ARF TFs, may trigger RH expansion in an *APOLO*/*RHD6*-independent manner. Furthermore, we found that in addition to *RHD6*, *APOLO* directly regulates 16 RH-related genes downstream *RHD6* and *RSLs* (**Supplementary Table 1**), hinting at an intricate regulatory network controlling RH expansion. Collectively, our results uncover a lncRNA-mediated epigenetic link between environmental signals and auxin homeostasis modulating RH growth. Moreover, our observations suggest that low temperatures restrict nutrient mobility and availability in the culture medium, leading to the promotion of polar RH growth. Further research will be needed to determine what is the limiting nutrient mediating the effect of cold on RH growth.

Although substantial progress has been achieved in the identification of the molecular actors controlling RH development, the impact of chromatin conformation in the transcriptional regulation of central TFs remains poorly understood. In this study, we have revealed a new mechanism of gene regulation in RHs by which the lncRNA *APOLO* integrates chromatin-associated ribonucleoprotein complexes together with the TF WYRK42, participating in the transcriptional activation of *RHD6* and the downstream RH gene network (**Figure 4D**). *APOLO* directly regulates the chromatin 3D conformation of the genomic region encompassing the *RHD6* locus and stoichiometrically recruits WYRK42, previously linked to Pi-starvation (Chen et al., 2009; Su et al., 2015). Low levels of *APOLO* fail to recruit WRKY42 to *RHD6* promoter region, whereas high levels of *APOLO* likely decoy WRKY42 from target chromatin, as it was shown for *APOLO* regulation over LHP1 binding activity (Ariel et al., 2020). Notably, low and high levels of *APOLO* result in higher *RHD6* basal transcriptional accumulation (through LHP1 activity and change in chromatin status), whereas both extremes impair *RHD6* activation by cold (through WRKY42 binding modulation). Our results suggest that an WRKY42-*APOLO* hub regulates RH cell elongation through the master regulator RHD6, although the *APOLO*-WYRKY42 hub potentially targets several additional cell wall related genes (**Supplementary Table 1**) at the end of the pathway controlled by RHD6 and the RHD6-downstream TFs RSL2/RSL4 (Mangano et al., 2017; Mangano et al., 2018a) .

Participation of epigenetic factors in root cell identity determination strongly suggests the default pattern for epidermal cell fate that can be overridden by environmental stimuli (Guimil and Dunand, 2006). Interestingly, it was reported that the expression of *GLABRA2* (*GL2*), a gene encoding a

TF repressing *RHD6* in atrichoblasts, is tightly regulated at the epigenetic level. By using 3D fluorescence

in situ hybridization, it was shown that alternative states of chromatin organization around the *GL2* locus

are required to control position-dependent cell-type specification in the root epidermis (Costa and Shaw,

2006) . Furthermore, *GL2* epigenetic regulation was proposed to be responsive to salt stress (Beyrne et

al., 2019). In addition, a comprehensive characterization of alternative mutant lines uncovered the role

of PRC2 in the regulation of RH development (Ikeuchi et al., 2015). Loss-of-function mutants in different

PRC2 subunits develop unicellular RHs but fail to retain the differentiated state, generating a disor-

ganized cell mass from each single RH. It was shown that the resulting RHs are able to undergo a normal

endoreduplication program, increasing their nuclear ploidy, although they subsequently reinitiate mitotic

division and successive DNA replication. It was proposed that aberrant RH development in PRC2 related

mutants is due to the epigenetic deregulation of key regulatory genes such as *WOUND INDUCED DEDIF-*

FERENTIATION 3 (*WIND3*) and *LEAFY COTYLEDON 2* (*LEC2*) (Ikeuchi et al., 2015). Here, we showed that

the single mutants *clf* (PRC2) and *lhp1* (PRC1) are affected in RH growth. In addition, we found that

H3K27me3 deposition and LHP1 binding to the *RHD6* locus is modulated by cold. Moreover, we showed

that in the *clf* background, H3K27me3 deposition throughout the *RHD6* locus is partially impaired, and

that the *clf* mutant is affected in RH elongation promoted by cold. Notably, basal transcriptional levels of

RHD6 are higher in the *clf* mutant (Supplementary Figure 4C). A phenotypic characterization revealed

that RH density is not altered in *clf* or *lhp1* mutants, nor in *APOLO* and *WRK42* deregulated lines (**Sup-**

plementary Figure 6), suggesting that *RHD6* is not ectopically expressed in epidermic cells. Thus, *RHD6*

over-accumulation may occur in inner cell layers of the root or in RHs, although their elongation in the *clf*

mutant may be blocked by additional perturbations of Polycomb-associated regulation of downstream

genes. Altogether, our results suggest that Polycomb proteins participate in the control of RH-related

genes transcriptional reprogramming at low temperatures.

Notably, CLF and LHP1 were shown to interact with a subset of lncRNAs in *Arabidopsis*, modulat-

ing the activity of PRC target genes (Lucero et al., 2020). Interestingly, several lncRNAs have been linked

to the control of transcription in response to cold. *FLOWERING LOCUS C* is regulated by at least three

lncRNAs. First, the alternative splicing of a set of antisense transcripts, collectively named as *COOLAIR*,

depends on the prolonged exposure to cold, epigenetically repressing *FLC* (Marquardt et al., 2014). The

use of the *COOLAIR* proximal poly(A) site results in down-regulation of *FLC* expression in a process involv-

ing FLOWERING LOCUS D (FLD), an H3K4me2 demethylase (Marquardt et al., 2014). A second lncRNA

called *COLD ASSISTED INTRONIC NONCODING RNA* (*COLDAIR*) is fully encoded in the sense strand of the

first intron of *FLC.* Similar to *COOLAIR*, its accumulation oscillates in response to low temperatures. It was

proposed that *COLDAIR* recruits the PRC2 component CLF to target *FLC* for H3K27me3 deposition (Heo

and Sung, 2011). More recently, a third lncRNA modulating *FLC* transcription was identified (Kim and

Sung, 2017). The cold-responsive lncRNA *COLDWRAP* is derived from the *FLC* proximal promoter and it

also interacts with PRC2. It was suggested that *COLDWRAP* functions in cooperation with the lncRNA

COLDAIR to retain Polycomb at the *FLC* promoter through the formation of a repressive intragenic chro-

matin loop (Kim and Sung, 2017). Another lncRNA named *SVALKA* was shown to mediate the response to

low temperatures (Kindgren et al., 2018). Interestingly, the activation of *SVALKA* by cold triggers the

transcription of a cryptic downstream lncRNA, which overlaps the antisense locus of the *C-*

repeat/dehydration-responsive element Binding Factor 1 (*CBF1*), involved in the early response to cold in

Arabidopsis. Antisense transcription causes Pol II head-to-head collision modulating transcriptional ter-

mination of *CBF1* (Kindgren et al., 2018). Here, we show that the auxin-responsive lncRNA *APOLO* is also

transcriptionally modulated by cold. The differential accumulation of 24nt siRNAs across the *APOLO* lo-

cus at low temperatures indicates that this activation is related to a decrease in RdDM. Moreover, we

showed here that the intergenic region between *PID* and *APOLO* acting as a divergent promoter is also

activated at low temperatures in RHs, as revealed by the reporter gene *GFP*. Thus, the lncRNA *APOLO*

integrates external signals into auxin-dependent developmental outputs in Arabidopsis.

In the last decade, lncRNAs have emerged as regulators of gene expression at different levels,

ranging from epigenetics to protein modifications and stability (Ariel et al., 2015). Notably, it has been

shown in animals that noncoding transcripts can be recognized by TFs. In humans, it was proposed that

the interaction with the lncRNA *SMALL NUCLEOLAR RNA HOST GENE 15* (*SNHG15*) stabilizes the TF Slug

in colon cancer cells. It was shown that *SNHG15* is recognized by the zinc finger domain of Slug prevent-

ing its ubiquitination and degradation in living cells (Jiang et al., 2018). Also, the transcriptional activity of

the human gene *DIHYDROFOLATE REDUCTASE* (*DHFR*) is regulated by a lncRNA encoded in its proximal

promoter. It was proposed that the nascent noncoding transcript forms a hybrid with its parent DNA and

decoys the regulatory TF IIB away from the *DHFR* promoter, dissociating the transcriptional pre-initiation

complex in quiescent cells (Martianov et al., 2007). The lncRNA *P21 ASSOCIATED ncRNA DNA DAMAGE*

ACTIVATED (*PANDA*) was identified in human cancer and it was activated in response to DNA damage

(Hung et al., 2011). *PANDA* is transcribed from the promoter region of the *CDKN1A* gene and interacts

with the TF NF-YA to limit the expression of pro-apoptotic genes. The activity of *PANDA* has been linked

to the progression of different tumors (Kotake et al., 2016; Shi et al., 2019). Interestingly, it was shown

that in addition to NF-YA, *PANDA* interacts with the scaffold-attachment-factor A (SAFA) as well as PRC1

and PRC2 to modulate cell senescence. In proliferating cells, SAFA and *PANDA* recruit Polycomb compo-

nents to repress the transcription of senescence-promoting genes. Conversely, the loss of SAFA–PANDA–

PRC interactions allows expression of the senescence program (Puvvula et al., 2014). In this work, we

showed that the PRC1-interacting lncRNA *APOLO* can also be recognized by the TF WRKY42, hinting at

general lncRNA-mediated mechanisms linking Polycomb complexes with the transcriptional machinery

across kingdoms. Furthermore, our observations indicate that the deregulation of *WRKY42* affects the

epigenetic environment of *RHD6*. It was previously shown that the addition of *in vitro* transcribed *APOLO*

to RNAi-*APOLO* chromatin extracts was able to partially restore R-loop formation over *APOLO* target

genes, and that high levels of *APOLO* may titer LHP1 away from chromatin (Ariel et al., 2020). Here we

show that the relative accumulation of the lncRNA *APOLO* can modulate the binding activity of its part-

ner TF to common target genes. Collectively, our results strongly support that environmentally con-

trolled cell fate in *Arabidopsis* relies on a transcriptional reprogramming governed by a network of epi-

genetic regulatory complexes, lncRNAs, TFs and effector proteins.

MATERIALS AND METHODS

Plant Material and Growth Conditions. All the *Arabidopsis thaliana* lines used were in the Columbia-0

(Col-0) background*. WRKY42* over expression transgenic plants were generated through *Agrobacterium*

tumefaciens (strain EHA105)-mediated transformation (Clough and Bent, 1998). *35S:WRKY42:GFP*

transformant lines were selected on MS/2 medium supplemented with kanamycin (40μg/mL) and

WRKY42 expression levels were measured by RT-qPCR (primers used are listed in **Supplementary Table**

3). The *wrky42* mutant line belongs to the SALK collection (SALK_121674C), as the one previously charac-

terized (Su et al., 2015).The *rhd6-3/rsl1-1/rsl4-1*; *rsl2*; *rsl4*; *rsl2rsl4*, *35S:RSL4 and EXP7p:RSL4* transgenic

lines were previously described and characterized (Yi et al., 2010; Hwang et al., 2017) Homozygous

plants were obtained in our laboratory and genotyped using the oligonucleotides indicated in **Supple-**

mentary Table 3. Seeds were surface sterilized and stratified at 4°C for 2d before being grown under

long day conditions (16h light, 140μE.m^{-2}.sec^{-1}/8h dark), on ½-strength Murashige and Skoog media (1/2

MS) (Duchefa, Netherlands) with 0.8% plant agar (Duchefa, Netherlands).

Cloning procedure. The coding region of *WRKY42* (AT4G04450) excluding the STOP codon was amplified

by PCR, cloned into the Gateway entry vector pENTR/D-TOPO (Invitrogen) and recombined by Gateway

technology (LR reaction) into the pK7FWG2,0 vector containing a *p35S-GFP* cassette

(http://www.psb.ugent.be/gateway/index.php).

Root hair phenotype characterization. For quantitative analyses of RH phenotypes, 10 fully elongated RH from the root elongation zone of 15-20 roots, were measured on the same conditions for each particular case and grown on vertical plates with ½-strength Murashige and Skoog media (1/2 MS) (Duchefa, Netherlands) and 0.8% plant agar (Duchefa, Netherlands) for 5 days at 22° and 3 days at 10°C. Measurements were made after 8 days. Images were captured with an Olympus SZX7 Zoom Stereo Microscope (Olympus, Japan) equipped with a Q-Colors digital camera and software Q Capture Pro 7(Olympus, Japan). RH density was determined as the number of hairs in a representative area of the root elongation zone using the same setup stated above and ImageJ software. Results were expressed as the mean ± standard error (SE) and in the case of RH density, values were expressed per mm^2. All measurements indicate the average of three independent experiments, each involving 15-20 seedlings.

Confocal microscopy analysis of root hairs. Confocal laser scanning microscopy was performed using Zeiss LSM5 Pascal (Zeiss, Germany) and a 40x water-immersion objective, N/A=1.2. Fluorescence was analyzed by using 488 nm laser for GFP excitation (Laser Intensity: 70%, Detector Gain:550, Amplifier Offset:0.1, Amplifier Gain:1), and emitted fluorescence was recorded between 490 and 525nm for GFP tag. Z stacks were done with an optical slice of 1µm, and fluorescence intensity was measured in 15µm ROI (Region Of Interest) at the RH tip and summed for quantification of fluorescence using ImageJ. Five replicates for each of ten roots and 15 hairs per root were observed. Col-0 wild type root hairs were used as a negative control, to check autofluorescence signal occurrence and no signal were detected in the wavelengths range stated above.

RNA extraction and RT-qPCR. Total RNA was extracted using TRIZol (Invitrogen) and 2 µg were subjected to DNase treatment according to the manufacturer's protocol (Thermo Scientific). One µg of DNase-free RNA was reverse-transcribed using Maxima H Minus Reverse Transcriptase (Thermo Scientific). RT-qPCR were performed using the LightCycler 480 SYBR Green I Master Kit on a LightCycler480 apparatus (Roche) using standard protocols (40 cycles, 60°C annealing). *PP2A* (AT1G13320; primers are listed in **Supplementary Table 3**) was used as reference.

RNA Immunoprecipitation. RNA immunoprecipitation (RIP) assays were performed on transiently transformed *N. benthamiana* leaves as described in (Sorenson and Bailey-Serres, 2015), or in 10-day-old *A. thaliana 35S:WRKY42:GFP* seedlings as described in (Bardou et al., 2014), using anti GFP (Abcam ab290)

and anti-IgG (Abcam ab6702). RIP was performed using Invitrogen Protein A Dynabeads. Precipitated

RNAs were prepared using TRI Reagent (Sigma-Aldrich), treated with DNase (Fermentas) and subjected

to RT-qPCR (High Capacity cDNA Reverse Transcription Kit (Thermo); primers used are listed in **Supple-**

mentary Table 3). Total RNAs were processed in parallel and considered the input sample.

Chromatin Immunoprecipitation. Chromatin immunoprecipitation (ChIP) assays were performed on 10-

day-old WT seedlings treated or not during 24h at 10 °C, using anti H3K27me3 (Diagenode pAb-195-050),

anti LHP1 (Covalab pab0923-P) and anti-IgG (Abcam ab6702), as described in (Ariel et al., 2020).

Crosslinked chromatin was sonicated using a water bath Bioruptor Pico (Diagenode; 30sec ON/30sec OFF

pulses; 10 cycles; high intensity). ChIP was performed using Invitrogen Protein A Dynabeads. Precipitated

DNA was recovered using Phenol:Chloroform:Isoamilic Acid (25:24:1; Sigma) and subjected to RT-qPCR

(primers used are listed in **Supplementary Table 3**). Untreated sonicated chromatin was processed in

parallel and considered the input sample. For *in vitro* competition assays, *APOLO* was transcribed using

the T7 transcription kit (Promega; (Ariel et al., 2020)). After regular chromatin isolation from 10-day-old

35S:WRKY42:GFP seedlings, the sample was split in 4 independent tubes and diluted to 1ml in Nuclei

Lysis Buffer without SDS. 0 µg, 0.1 µg, 1 µg and 10 µg of *APOLO* were added to each sample respectively,

and incubated in rotation at 4 °C for 3h. Then, cross-linking was performed with 1% formaldehyde for 5

min at 4 °C, followed by 5 min with a final concentration of 50 mM glycine. SDS was added to a final con-

centration of 0.1% prior to sonication and the subsequent steps of a regular ChIP protocol.

For ChIP in transiently transformed leaves, 3-week-old *A. thaliana* were transformed as previously de-

scribed (Zhang et al., 2020) In brief, *Agrobacterium tumefaciens* strain GV3101 carrying *35S:WRKY42:GFP*

construct were grown for 2 days in YEB-induced medium plates at 28 °C. Agrobacterium cells were

scraped and resuspended in washing solution (10 mM MgCl$_2$, 100 µM acetosyringone). Infiltration solu-

tion (¼MS [pH = 6.0], 1% sucrose, 100 µM acetosyringone, 0.005% [v/v, 50 µl/l] Silwet L-77) was pre-

pared with the previous solution, adjusting the OD600=0.5. The infiltration was carried out in all leaves >

0.5cm in length of between 10 and 15 plants per genotype (WT, *35S:APOLO* and RNAi-*APOLO* lines). After

infiltration, plants were kept in light for 1h and then in darkness for 24h. Finally they were transferred

back to light. Images and samples were obtained 3 days after infiltration. For image acquisition, infiltrat-

ed leaves were imaged with a Leica TCS SP8 confocal laser scanning microscope with excitation at 488

nm (Intensity=8%) and detection at 495-530 nm for GFP and 610-670 (gain 650) nm for chlorophyll fluo-

rescence. Images were captured using 10X and 20X lenses, and processed using Fiji software (Schindelin

et al., 2012).

Chromatin Isolation by RNA Purification followed by qPCR or mass spectrometry. A method adapted from the ChIRP protocol (Chu et al., 2012) was developed to allow the identification of plant DNA associated to specific lncRNAs, as described in (Ariel et al., 2014; Ariel et al., 2020). Briefly, plants were *in vivo* crosslinked and cell nuclei were purified and extracted through sonication. The resulting supernatant was hybridized against biotinylated complementary oligonucleotides that tile the lncRNA of interest and lncRNA-associated chromatin was isolated using magnetic streptavidin beads. One hundred pmol of probes against *APOLO* (ODD and EVEN set of probes (Ariel et al., 2014; Ariel et al., 2020)) and the corresponding negative set against LacZ were used for independent purification. Co-purified ribonucleoprotein complexes were eluted and used to purify RNA or DNA, which were later subjected to downstream assays for quantification as previously described (Ariel et al., 2020).

The exploratory identification of plant nuclear proteins bound to *APOLO* was performed as described in (Rigo et al., 2020). Plant samples were prepared as for ChIRP-qPCR. For protein extraction, approximately 250 g of 7-day-old Col-0 plants grown on solid half-strength MS medium was irradiated three times with UV using a CROSSLINKERCL-508 (Uvitec) at 0.400 J/ cm^2. Protein purification samples for protein extraction were DNase-treated according to the manufacturer (Thermo Scientific). After addition of 1.8ml of TCA-acetone (5ml 6.1N TCA + 45ml acetone + 35μl β-mercaptoethanol), samples were incubated overnight at -80°C. After centrifugation at 20000rpm for 20min at 4°C, the supernatant was discarded and 1.8ml of acetone wash buffer (120ml acetone, 84μl β-mercaptoethanol) was added to the samples. Then, samples were incubated 1h at -20°C and centrifuged again at 20000 rpm for 20min at 4°C. The supernatant was discarded, and the dry pellet was used for mass spectrometry analyses.

Chromatin Conformation Capture. Chromosome conformation capture (3C) was performed basically as previously described in (Ariel et al., 2020) starting with 2g of seedlings. Digestions were performed overnight at 37°C with 400U *Dpn*II (NEB). DNA was ligated by incubation at 16°C for 5h in 4 ml volume using 100U of T4 DNA ligase (NEB). After reverse crosslinking and Proteinase K treatment (Invitrogen), DNA was recovered by Phenol:Chloroform:Isoamilic Acid (25:24:1; Sigma) extraction and ethanol precipitation. Relative interaction frequency was calculated by qPCR (primers used are listed in **Supplementary Table 3**). A region free of *Dpn*II was used to normalize the amount of DNA.

Acknowledgments

We thank Chang Liu for providing the Hi-C plot of Figure 2B. This work was supported by grants from
ANPCyT (PICT2016-0132 and PICT2017-0066) and Instituto Milenio iBio – Iniciativa Científica Milenio,
MINECON to JME; ANPCyT (PICT2016-0007 and PICT2016-0289) and Fima Leloir Award to FA; UNL (CAI+D
2016) to LL; Laboratoire d'Excellence (LABEX) Saclay Plant Sciences (SPS; ANR-10-LABX-40) to MC; and
CNRS (Laboratoire International Associé NOCOSYM) to MC and FA. LL, FI, JME and FA are researchers of
CONICET; MM, JMP, CFF and JRM are fellows of the same institution.
498

Author contributions

MM, JMP, LL, CFF, JRM, NM, AC, MB and FA performed the experiments. JB, MB, FI, MC, JE and FA ana-
lyzed the data. JE and FA conceived the project. FA, JE and CFF wrote the manuscript with the contribu-
tion of all authors.

References

Ariel, F., Jegu, T., Latrasse, D., Romero-Barrios, N., Christ, A., Benhamed, M., and Crespi, M. (2014).
Noncoding transcription by alternative rna polymerases dynamically regulates an auxin-driven
chromatin loop. *Mol. Cell* **55**:383–396.

Ariel, F., Romero-Barrios, N., Jégu, T., Benhamed, M., and Crespi, M. (2015). Battles and hijacks:
Noncoding transcription in plants. *Trends Plant Sci.* **20**:362–371.

**Ariel, F., Lucero, L., Christ, A., Mammarella, M. F., Jegu, T., Veluchamy, A., Mariappan, K., Latrasse, D.,
Blein, T., Liu, C., et al.** (2020). R-Loop Mediated trans Action of the APOLO Long Noncoding RNA.
Mol. Cell **77**:1055-1065.e4.

Balcerowicz, D., Schoenaers, S., and Vissenberg, K. (2015). Cell Fate Determination and the Switch from
Diffuse Growth to Planar Polarity in Arabidopsis Root Epidermal Cells. *Front. Plant Sci.* **6**:1–13.

**Bardou, F., Ariel, F., Simpson, C. G., Romero-Barrios, N., Laporte, P., Balzergue, S., Brown, J. W. S., and
Crespi, M.** (2014). Long Noncoding RNA Modulates Alternative Splicing Regulators in Arabidopsis.
Dev. Cell **30**:166–176.

Beyrne, C. C., Iusem, N. D., and González, R. M. (2019). Effect of salt stress on cytosine methylation
within GL2, an Arabidopsis thaliana gene involved in root epidermal cell differentiation. Absence of
inheritance in the unstressed progeny. *Int. J. Mol. Sci.* **20**:1–13.

**Bhosale, R., Giri, J., Pandey, B. K., Giehl, R. F. H., Hartmann, A., Traini, R., Truskina, J., Leftley, N.,
Hanlon, M., Swarup, K., et al.** (2018). A mechanistic framework for auxin dependent Arabidopsis
root hair elongation to low external phosphate. *Nat. Commun.* **9**:1–9.

Bruex, A., Kainkaryam, R. M., Wieckowski, Y., Kang, Y. H., Bernhardt, C., Xia, Y., Zheng, X., Wang, J. Y.,

Lee, M. M., Benfey, P., et al. (2012). A Gene Regulatory Network for Root Epidermis Cell

Differentiation in Arabidopsis. *PLoS Genet.* **8**:1–20.

Buah, J. N., Kawamitsu, Y., Sato, S., and Murayama, S. (1999). Effects of Different Types and

Concentrations of Gelling Agents on the Physical and Chemical Properties of Media and the Growth

of Banana (Musa spp.) in Vitro. *Plant Prod. Sci.* **2**:138–145.

Chen, Y. F., Li, L. Q., Xu, Q., Kong, Y. H., Wang, H., and Wu, W. H. (2009). The WRKY6 transcription factor

modulates PHOSPHATE1 expression in response to low pi stress in arabidopsis. *Plant Cell* **21**:3554–

3566.

Chu, C., Quinn, J., and Chang, H. Y. (2012). Chromatin isolation by RNA purification (ChIRP). *J. Vis. Exp.*

Advance Access published March 25, 2012, doi:10.3791/3912.

Clough, S. J., and Bent, A. F. (1998). Floral dip: a simplified method for Agrobacterium-mediated

transformation of Arabidopsis thaliana. *Plant J.* **16**:735–43.

Costa, S., and Shaw, P. (2006). Chromatin organization and cell fate switch respond to positional

information in Arabidopsis. *Nature* **439**:493–496.

Datta, S., Prescott, H., and Dolan, L. (2015). Intensity of a pulse of RSL4 transcription factor synthesis

determines Arabidopsis root hair cell size. *Nat. Plants* **1**:1–6.

Devaiah, B. N., Karthikeyan, A. S., and Raghothama, K. G. (2007). WRKY75 transcription factor is a

modulator of phosphate acquisition and root development in Arabidopsis. *Plant Physiol.* **143**:1789–

1801.

Dong, J., Chen, C., and Chen, Z. (2003). Expression profiles of the Arabidopsis WRKY gene superfamily

during plant defense response. *Plant Mol. Biol.* **51**:21–37.

Franco-Zorrilla, J. M., Valli, A., Todesco, M., Mateos, I., Puga, M. I., Rubio-Somoza, I., Leyva, A., Weigel,

D., García, J. A., and Paz-Ares, J. (2007). Target mimicry provides a new mechanism for regulation

of microRNA activity. *Nat. Genet.* **39**:1033–1037.

Friml, J., Yang, X., Michniewicz, M., Weijers, D., Quint, A., Tietz, O., Benjamins, R., Ouwerkerk, P. B. F.,

Ljung, K., Sandberg, G., et al. (2004). A PINOID-dependent binary switch in apical-basal PIN polar

targeting directs auxin efflux. *Science* **306**:862–5.

Ghashghaie, J., Brenckmann, F., and Saugier, B. (1991). Effects of agar concentration on water status

and growth of rose plants cultured in vitro. *Physiol. Plant.* **82**:73–78.

Guimil, S., and Dunand, C. (2006). Patterning of Arabidopsis epidermal cells: epigenetic factors regulate

the complex epidermal cell fate pathway. *Trends Plant Sci.* **11**:601–609.

Gyula, P., Baksa, I., Tóth, T., Mohorianu, I., Dalmay, T., and Szittya, G. (2018). Ambient temperature regulates the expression of a small set of sRNAs influencing plant development through NF-YA2 and YUC2. *Plant Cell Environ.* **41**:2404–2417.

Heo, J. B., and Sung, S. (2011). Vernalization-mediated epigenetic silencing by a long intronic noncoding RNA. *Science (80-.).* **331**:76–9.

Hung, T., Wang, Y., Lin, M. F., Koegel, A. K., Kotake, Y., Grant, G. D., Horlings, H. M., Shah, N., Umbricht, C., Wang, P., et al. (2011). Extensive and coordinated transcription of noncoding RNAs within cell-cycle promoters. In *Nature Genetics*, pp. 621–629. NIH Public Access.

Hwang, Y., Choi, H. S., Cho, H. M., and Cho, H. T. (2017). Tracheophytes contain conserved orthologs of a basic helix-loop-helix transcription factor that modulate ROOT HAIR SPECIFIC genes. *Plant Cell* **29**:39–53.

Ikeuchi, M., Iwase, A., Rymen, B., Harashima, H., Shibata, M., Ohnuma, M., Breuer, C., Morao, A. K., De Lucas, M., De Veylder, L., et al. (2015). PRC2 represses dedifferentiation of mature somatic cells in Arabidopsis. *Nat. Plants* **1**:1–7.

Jabnoune, M., Secco, D., Lecampion, C., Robaglia, C., Shu, Q., and Poirier, Y. (2013). A Rice cis-Natural antisense RNA acts as a translational enhancer for its cognate mRNA and contributes to phosphate homeostasis and plant fitness. *Plant Cell* **25**:4166–4182.

Jiang, H., Li, T., Qu, Y., Wang, X., Li, B., Song, J., Sun, X., Tang, Y., Wan, J., Yu, Y., et al. (2018). Long non-coding RNA SNHG15 interacts with and stabilizes transcription factor Slug and promotes colon cancer progression. *Cancer Lett.* **425**:78–87.

Jones, A. R., Kramer, E. M., Knox, K., Swarup, R., Bennett, M. J., Lazarus, C. M., Leyser, H. M. O., and Grierson, C. S. (2009). Auxin transport through non-hair cells sustains root-hair development. *Nat. Cell Biol.* **11**:78–84.

Karas, B., Amyot, L., Johansen, C., Sato, S., Tabata, S., Kawaguchi, M., and Szczyglowski, K. (2009). Conservation of Lotus and Arabidopsis basic helix-loop-helix proteins reveals new players in root hair development. *Plant Physiol.* **151**:1175–1185.

Kim, D. H., and Sung, S. (2017). Vernalization-Triggered Intragenic Chromatin Loop Formation by Long Noncoding RNAs. *Dev. Cell* **40**:302-312.e4.

Kindgren, P., Ard, R., Ivanov, M., and Marquardt, S. (2018). Transcriptional read-through of the long non-coding RNA SVALKA governs plant cold acclimation. *Nat. Commun.* **9**:1–11.

Kotake, Y., Kitagawa, K., Ohhata, T., Sakai, S., Uchida, C., Niida, H., Naemura, M., and Kitagawa, M. (2016). Long Non-coding RNA, PANDA, Contributes to the Stabilization of p53 Tumor Suppressor

Protein. *Anticancer Res.* **36**:1605–11.

Lamport, D. T. A., Kieliszewski, M. J., Chen, Y., and Cannon, M. C. (2011). Role of the Extensin

Superfamily in Primary Cell Wall Architecture. *Plant Physiol.* **156**:11–19.

Lin, Q., Ohashi, Y., Kato, M., Tsuge, T., Gu, H., Qu, L. J., and Aoyama, T. (2015). GLABRA2 directly

suppresses basic helix-loop-helix transcription factor genes with diverse functions in root hair

development. *Plant Cell* **27**:2894–2906.

Liu, C., Wang, C., Wang, G., Becker, C., Zaidem, M., and Weigel, D. (2016). Genome-wide analysis of

chromatin packing in Arabidopsis thaliana at single-gene resolution. *Genome Res.* **26**:1057–1068.

Lucero, L., Fonouni-Farde, C., Crespi, M., and Ariel, F. (2020). Long noncoding RNAs shape transcription

in plants. *Transcription* Advance Access published May 14, 2020,

doi:10.1080/21541264.2020.1764312.

Mangano, S., Denita-Juarez, S. P., Choi, H. S., Marzol, E., Hwang, Y., Ranocha, P., Velasquez, S. M.,

Borassi, C., Barberini, M. L., Aptekmann, A. A., et al. (2017). Molecular link between auxin and

ROS-mediated polar growth. *Proc. Natl. Acad. Sci. U. S. A.* **114**:5289–5294.

Mangano, S., Martínez Pacheco, J., Marino-Buslje, C., and Estevez, J. M. (2018a). How Does pH Fit in

with Oscillating Polar Growth? *Trends Plant Sci.* **23**:479–489.

Mangano, S., Denita-Juarez, S. P., Marzol, E., Borassi, C., and Estevez, J. M. (2018b). High Auxin and

High Phosphate Impact on RSL2 Expression and ROS-Homeostasis Linked to Root Hair Growth in

Arabidopsis thaliana. *Front. Plant Sci.* **9**:1–8.

Marquardt, S., Raitskin, O., Wu, Z., Liu, F., Sun, Q., and Dean, C. (2014). Functional Consequences of

Splicing of the Antisense Transcript COOLAIR on FLC Transcription. *Mol. Cell* **54**:156–165.

Martianov, I., Ramadass, A., Serra Barros, A., Chow, N., and Akoulitchev, A. (2007). Repression of the

human dihydrofolate reductase gene by a non-coding interfering transcript. *Nature* **445**:666–670.

Marzol, E., Borassi, C., Denita Juárez, S. P., Mangano, S., and Estevez, J. M. (2017). RSL4 Takes Control:

Multiple Signals, One Transcription Factor. *Trends Plant Sci.* **22**:553–555.

Marzol, E., Borassi, C., Bringas, M., Sede, A., Rodríguez Garcia, D. R., Capece, L., and Estevez, J. M.

(2018). Filling the Gaps to Solve the Extensin Puzzle. *Mol. Plant* **11**:645–658.

Menand, B., Yi, K., Jouannic, S., Hoffmann, L., Ryan, E., Linstead, P., Schaefer, D. G., and Dolan, L.

(2007). An ancient mechanism controls the development of cells with a rooting function in land

plants. *Science (80-.).* **316**:1477–1480.

Nonami, H., and Boyer, J. S. (1989). Turgor and Growth at Low Water Potentials. *Plant Physiol.* **89**:798–

804.

Pires, N. D., Yi, K., Breuninger, H., Catarino, B., Menand, B., and Dolan, L. (2013). Recruitment and
remodeling of an ancient gene regulatory network during land plant evolution. *Proc. Natl. Acad. Sci.*
U. S. A. **110**:9571–9576.

Puvvula, P. K., Desetty, R. D., Pineau, P., Marchio, A., Moon, A., Dejean, A., and Bischof, O. (2014). Long
noncoding RNA PANDA and scaffold-attachment-factor SAFA control senescence entry and exit.
Nat. Commun. **5**:1–16.

Rigo, R., Bazin, J., Romero-Barrios, N., Moison, M., Lucero, L., Christ, A., Benhamed, M., Blein, T.,
Huguet, S., Charon, C., et al. (2020). *An Arabidopsis long noncoding RNA modulates the*
transcriptome through interactions with a network of splicing factors. Cold Spring Harbor
Laboratory.

Ringli, C. (2010). The hydroxyproline-rich glycoprotein domain of the Arabidopsis LRX1 requires Tyr for
function but not for insolubilization in the cell wall. *Plant J.* **63**:662–669.

Ryu, K. H., Kang, Y. H., Park, Y. H., Hwang, I., Schiefelbein, J., and Lee, M. M. (2005). The WEREWOLF
MYB protein directly regulates CAPRICE transcription during cell fate specification in the
Arabidopsis root epidermis. *Development* **132**:4765–4775.

Schiefelbein, J., Huang, L., and Zheng, X. (2014). Regulation of epidermal cell fate in Arabidopsis roots:
the importance of multiple feedback loops. *Front. Plant Sci.* **5**:1–4.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden,
C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source platform for biological-image analysis.
Nat. Methods **9**:676–682.

Schoenaers, S., Balcerowicz, D., Breen, G., Hill, K., Zdanio, M., Mouille, G., Holman, T. J., Oh, J., Wilson,
M. H., Nikonorova, N., et al. (2018). The Auxin-Regulated CrRLK1L Kinase ERULUS Controls Cell
Wall Composition during Root Hair Tip Growth. *Curr. Biol.* **28**:722-732.e6.

Shi, W., Wang, Q., Bian, Y., Fan, Y., Zhou, Y., Feng, T., Li, Z., and Cao, X. (2019). Long noncoding RNA
PANDA promotes esophageal squamous carcinoma cell progress by dissociating from NF-YA but
interact with SAFA. *Pathol. Res. Pract.* **215**:1–11.

Singha, S., Townsend, E.C. and Oberly, G. H. (1985). Mineral nutrient status of crabapple and pear
shoots cultured in vitro on varying commercial agars. *J. Am. Soc. Hortic. Sci.* **110**:407-411.

Song, S. K., Ryu, K. H., Kang, Y. H., Song, J. H., Cho, Y. H., Yoo, S. D., Schiefelbein, J., and Lee, M. M.
(2011). Cell fate in the Arabidopsis root epidermis is determined by competition between
WEREWOLF and CAPRICE. *Plant Physiol.* **157**:1196–1208.

Sorenson, R., and Bailey-Serres, J. (2015). Rapid immunopurification of ribonucleoprotein complexes of

plants. *Methods Mol. Biol.* **1284**:209–219.

Stepanova, A. N., Robertson-Hoyt, J., Yun, J., Benavente, L. M., Xie, D. Y., Doležal, K., Schlereth, A.,

Jürgens, G., and Alonso, J. M. (2008). TAA1-Mediated Auxin Biosynthesis Is Essential for Hormone

Crosstalk and Plant Development. *Cell* **133**:177–191.

Su, T., Xu, Q., Zhang, F. C., Chen, Y., Li, L. Q., Wu, W. H., and Chen, Y. F. (2015). WRKY42 Modulates

phosphate homeostasis through regulating phosphate translocation and acquisition in Arabidopsis.

Plant Physiol. **167**:1579–1591.

Velasquez, S. M., Ricardi, M. M., Dorosz, J. G., Fernandez, P. V., Nadra, A. D., Pol-Fachin, L., Egelund, J.,

Gille, S., Harholt, J., Ciancia, M., et al. (2011). O-glycosylated cell wall proteins are essential in root

hair growth. *Science (80-.).* **332**:1401–1403.

Velasquez, S. M., Marzol, E., Borassi, C., Pol-Fachin, L., Ricardi, M. M., Mangano, S., Juarez, S. P. D.,

Salter, J. D. S., Dorosz, J. G., Marcus, S. E., et al. (2015). Low Sugar Is Not Always Good: Impact of

Specific O -Glycan Defects on Tip Growth in Arabidopsis. *Plant Physiol.* **168**:808–813.

Veluchamy, A., Jégu, T., Ariel, F., Latrasse, D., Mariappan, K. G., Kim, S.-K., Crespi, M., Hirt, H.,

Bergounioux, C., Raynaud, C., et al. (2016). LHP1 Regulates H3K27me3 Spreading and Shapes the

Three-Dimensional Conformation of the Arabidopsis Genome. *PLoS One* **11**:1–25.

Vijayakumar, P., Datta, S., and Dolan, L. (2016). ROOT HAIR DEFECTIVE SIX-LIKE4 (RSL4) promotes root

hair elongation by transcriptionally regulating the expression of genes required for cell growth.

New Phytol. **212**:944–953.

Waese, J., Fan, J., Pasha, A., Yu, H., Fucile, G., Shi, R., Cumming, M., Kelley, L. A., Sternberg, M. J.,

Krishnakumar, V., et al. (2017). ePlant: Visualizing and exploring multiple levels of data for

hypothesis generation in plant biology. *Plant Cell* **29**:1806–1821.

Wang, H., Xu, Q., Kong, Y. H., Chen, Y., Duan, J. Y., Wu, W. H., and Chen, Y. F. (2014). Arabidopsis

WRKY45 transcription factor activates Phosphate transporter1;1 expression in response to

phosphate starvation. *Plant Physiol.* **164**:2020–2029.

Wang, Q., Qin, G., Cao, M., Chen, R., He, Y., Yang, L., Zeng, Z., Yu, Y., Gu, Y., Xing, W., et al. (2020). A

phosphorylation-based switch controls TAA1-mediated auxin biosynthesis in plants. *Nat. Commun.*

11:1–10.

Won, S. K., Lee, Y. J., Lee, H. Y., Heo, Y. K., Cho, M., and Cho, H. T. (2009). cis-element- and

transcriptome-based screening of root hair-specific genes and their functional characterization in

Arabidopsis. *Plant Physiol.* **150**:1459–1473.

Xu, W., Li, K., Li, S., Hou, Q., Zhang, Y., Liu, K., and Sun, Q. (2020). The R-Loop Atlas of Arabidopsis

Development and Responses to Environmental Stimuli. *Plant Cell* **32**:888–903.

Yi, K., Menand, B., Bell, E., and Dolan, L. (2010). A basic helix-loop-helix transcription factor controls cell

growth and size in root hairs. *Nat. Genet.* **42**:264–267.

Zhang, T. Q., Xu, Z. G., Shang, G. D., and Wang, J. W. (2019). A Single-Cell RNA Sequencing Profiles the

Developmental Landscape of Arabidopsis Root. *Mol. Plant* **12**:648–660.

Zhang, Y., Chen, M., Siemiatkowska, B., Toleco, M. R., Jing, Y., Strotmann, V., Zhang, J., Stahl, Y., and

Fernie, A. R. (2020). A Highly Efficient Agrobacterium-Mediated Method for Transient Gene

Expression and Functional Studies in Multiple Plant Species. *Plant Commun.* **1**:100028.

Figure 1. *APOLO* regulates root hair elongation in response to low temperatures

A. *APOLO* transcript levels measured by RT-qPCR in roots of 10-day-old plants grown at 22 °C treated or not for 24h at 10 °C.

B. The *APOLO* promoter region (full intergenic region between *PID* and *APOLO*) is active in root hairs at 22 °C and 10 °C, using the *GFP* reporter gene. Images are representative of the lowest and highest GFP signal detected in the observation of several roots. The quantification of the fluorescence is shown in the right panel.

C. Quantification of RH length of 8-day-old WT, RNAi-*APOLO* and *35S:APOLO* plants, respectively, at 22 °C and 10 °C. Each point is the mean ±error of the length of the 10 longest RHs identified in the root elonga-tion zone in a single root. Representative images of each genotype are shown below the graph. The scale bas represent 500μm. Asterisks indicate significant differences between WT and the corresponding geno-type at the same temperature or significant differences between temperature treatments on the same genotype at the same temperature (one way ANOVA followed by a Tukey-Kramer test; "*"<0.05, "**"<0.01, "***"<0.001). NS stands for no statistically significant difference.

D. *RHD6* transcript levels measured by RT-qPCR in roots of WT plants vs. two independent RNAi-*APOLO* lines and two independent *35S:APOLO* lines grown at 22 °C treated or not for 24h at 10 °C.

In A and D, the error bars represent the SD of 3 biological replicates. The asterisks indicate that the dif-ferences are significant (t test p<0.05). Values are normalized using the constant housekeeping transcript *PP2A*.

Figure 2. *APOLO* **directly modulates chromatin three-dimensional conformation of the** *RHD6* **locus**

A. Epigenomic landscape of the *RHD6* locus. Lane 1: H3K27me3 deposition by ChIP-Seq (Veluchamy et al., 2016). Lane 2: LHP1 deposition by ChIP-Seq (Veluchamy et al., 2016). Lane 3 to 5: *APOLO* recognition by ChIRP-Seq (Lane 3 and 4, using ODD and EVEN sets of probes against *APOLO*, respectively; Lane 5, negative control using LacZ probes)(Ariel et al., 2020). Lane 6 to 8: R-loop formation by DRIP-Seq (R-loop Atlas, root samples,(Xu et al., 2020), on Watson strand (Lane 6) and Crick strand (Lane 7). DRIP negative control after RNAseH treatment is shown in Lane 8. Gene annotation is shown at the bottom.

B. Chromatin loops identified in the *RHD6* region by *Dpn*II HiC (Liu et al., 2016). Colors of the loops are related to the corresponding q-values indicated below. Black boxes in A and B indicate the same genomic locations, where the bases of the chromatin loop correlate with R-loop formation and *APOLO* recognition (compared to gene annotations). In red the arrows indicate the position of the probes used for 3C-qPCR shown in **D**.

C. *APOLO* association to DNA of the *RHD6* locus by ChIRP-qPCR in WT and RNAi-*APOLO* plants. The background level was determined using a set of probes against LacZ RNA.

D. Relative chromatin loop formation by 3C-qPCR deduced from **B**, in WT plants vs. *35S:APOLO* and RNAi lines for the region upstream *RHD6*. The probes used for 3C-qPCR are indicated in red in panel **B**.

In C and D, the error bars represent the SD of 3 biological replicates. The asterisks indicate that the differences are significant (t test "*"<0.05, "***"<0.001).

Figure 3. The transcription factor WRKY42 interacts with *APOLO*, directly regulates *RHD6* and participates in the response to cold

A. *APOLO*-WRKY42 *in vivo* interaction demonstrated by RNA Immunoprecipitation (RIP)-qPCR in transiently transformed tobacco leaves and stably transformed Arabidopsis plants. In tobacco, WRKY42:GFP and *APOLO* were co-transformed under the control of the 35S constitutive promoter. In Arabidopsis, *35S:WRKY42:GFP* transformed plants were used to detect the interaction with endogenous *APOLO*. Anti-IgG antibodies were used as a negative control. The RNA *PP2A* was considered as a RIP negative control in Arabidopsis plants.

B. Arabidopsis eFP Browser (Waese et al., 2017) plot representing the increase of *WRKY42* transcript levels in roots of seedlings treated for 24h at 4°C.

C. *WRKY42* transcript levels measured by RT-qPCR in roots of plants grown at 22°C treated or not for 24h at 10°C.

D. Chromatin Immunoprecipitation (ChIP)-qPCR assay revealing regulation by WRKY42 of *RHD6* by direct
recognition of its promoter region. Probes amplifying *PP2A* were used as a negative control of the exper-
iment. Anti-IgG antibodies were used as a negative control for each pair of probes.
E. *RHD6* transcript levels measured by RT-qPCR in roots of WT plants vs. *wrky42* mutants and
35S:WRKY42:GFP lines grown at 22°C treated or not for 24h at 10°C.
F. Quantification of RH length of WT, *wrky42* and *35S:WRKY42:GFP* plants at 22°C and 10°C. Each point is
the mean of the length of the 10 longest RHs identified in a single root. Representative images of each
genotype are shown on the right. The scale bas represent 500µm. Asterisks indicate significant differ-
ences between WT and the corresponding genotype at the same temperature (one way ANOVA followed
by a Tukey-Kramer test; "*"<0.05, "***"<0.001).
In A, C, D and E, the error bars represent the SD of 3 biological replicates. The asterisks indicate that the
differences are significant (t test p<0.05). NS stands for no statistically significant difference. In C and E,
values are normalized using the constant housekeeping transcript *PP2A*.

Figure 4. The WRKY42-*APOLO* hub activates *RHD6* transcription in response to low temperatures, pro-moting root hair expansion

A. H3K27me3 deposition in WT vs. *wrky42* mutant 10-day-old seedlings grown at 23°C over *RHD6* and a WRKY42-independent *APOLO* target (*AZG2*; (Ariel et al., 2020)). An immunoprecipitation (ChIP-qPCR) was performed using an anti-IgG antibody as a negative control.

B. *APOLO* association to DNA of the *RHD6* locus by ChIRP-qPCR in WT and *wrky42* mutant plants. The background level was determined using a set of probes against LacZ RNA.

C. WRKY42 binding to *RHD6* promoter in transiently transformed leaves of WT, RNAi-*APOLO* and *35S:APOLO* plants. WRKY42:GFP was immunoprecipitated using the anti-GFP antibody (ChIP-qPCR). An immunoprecipitation was performed using an anti-IgG antibody as a negative control.

D. WRKY42 immunoprecipitation over *RHD6* with the addition of increasing concentrations of *in vitro* transcribed *APOLO* prior to cross-linking.

E. Proposed model for the action of *APOLO* in RH growth under low temperatures. At 22 °C, *RHD6* moderate transcription is regulated by *APOLO*, H3K27me3 and LHP1, maintaining a repressive chromatin loop encompassing the *RHD6* promoter region. In response to cold, *APOLO* levels increase, still recognizing the *RHD6* locus by sequence complementarity (R-loops). Then, LHP1 (brown balls) is decoyed away from chromatin and PRC2-dependent H3K27me3 (green balls) deposition decreases in a process involving WRKY42 and leading to the opening of the *RHD6* promoter region. Additionally, *APOLO* and WRKY42 jointly activate *RHD6* transcription. RHD6 activates the transcription of downstream TF genes *RSL2* and *4* promoting RH cell expansion. Yet unknown factors activate *RSL2* and *4* independently from *APOLO* and RHD6 in response to cold.

In A to D the error bar represents the SD of 3 biological replicates. The asterisk indicates that the difference is significant (t test "*"<0.05, "**"<0.01).

783
Supplementary Figure 1: *RHD6* and *APOLO* transcripts are enriched in root hair cells
A. Uniform Manifold Approximation and Projection (UMAP) clustering map of Arabidopsis root single cell
RNA-seq ((Zhang et al., 2019); http://wanglab.sippe.ac.cn/rootatlas/) data. *RHD6*, *APOLO* and the
trichoblast cell marker *COBL9* are respectively indicated in red in each plot.
B. 24nt siRNA accumulation over the *APOLO* locus revealed by small RNA-Seq of roots from WT seedlings
grown at 15 °C or 23 °C (Gyula et al., 2018). On the right, a zoom of the 3' region is shown, with the over-
lapped signal at both temperatures.

A

B

Supplementary Figure 2. Root hair expansion in response to cold is mediated by *RSL2* and *RSL4*

A. *RSL2* and *RSL4* transcript levels measured by RT-qPCR in roots of WT plants vs. two independent RNAi-*APOLO* lines and two independent *35S:APOLO* lines grown at 22 °C treated or not for 24h at 10 °C. The asterisks indicate that the differences are significant (t test p<0.05). NS stands for no statistically signifi-cant difference. Values are normalized using the constant housekeeping transcript *PP2A*.

B. Quantification of RH length of plants WT, *rhd6-3, rsl1/4; rsl2; rsl4; rsl2/4*; *35S:RSL4* and *pEXP7:RSL4* at 22°C and 10°C. Each point is the mean ±error of the length of the 10 longest RHs identified in the root elongation zone in a single root. Representative images of each genotype are shown below the graph. The scale bas represent 500μm. Asterisks indicate significant differences between WT and the corre-sponding genotype at the same temperature or significant differences between temperature treatments

802 on the same genotype at the same temperature (one way ANOVA followed by a Tukey-Kramer test;

803 "*"<0.05, "**"<0.01, "***"<0.001), unless indicated differently. NS stands for no statistically significant

804 difference.

Supplementary Figure 3. Root hair expansion in response to cold seems to be related to nutrient availability

A. Quantification of RH length of WT plants at 22°C and 10°C at growing concentrations of Murashige and Skoog (MS) culture media. Each point is the mean ±error of the length of the 10 longest RHs identified in the root elongation zone in a single root. Representative images of each genotype are shown below. Asterisks indicate significant differences between temperature treatments on a specific nutrient media condition (one way ANOVA followed by a Tukey-Kramer test; "*"<0.05, "***"<0.001). NS stands for no statistically significant difference.

B. Quantification of RH length of WT plants expressed as the ratio between 10°C and 22°C at growing concentrations of MS culture media, for given concentrations of agar. Representative images of each genotype are shown below the graph. Each point is the mean ±error of the length of the 10 longest RHs identified in the root elongation zone in a single root. Note that at higher concentrations of agar, the ΔRH between conditions is lower, but the RHs are longer (as shown in the representative photos below). Asterisks indicate significant differences between agar percentage conditions (one way ANOVA followed by a Tukey-Kramer test; "*"<0.05, "**"<0.01, "***"<0.001). NS stands for no statistically significant difference. The dotted box indicates the same results as in panel A, expressed as ΔRH growth instead of absolute RH length at 22°C and 10°C independently. In the photos in A and B, the scale bas represent 500μm.

823
824
Supplementary Figure 4: Induction of root hair elongation by low temperatures requires epigenetic Polycomb-mediated reprogramming

A. H3K27me3 deposition across the *RHD6* locus in WT, *clf* and *lhp1* mutants, revealed by ChIP-seq from seedlings grown at 23°C (Veluchamy et al., 2016).

B. H3K27me3 deposition measured by ChIP-qPCR over the *RHD6* locus in WT plants grown at 22°C treated or not for 24h at 10°C. The negative control was performed using an anti-IgG antibody.

C. *RHD6* transcript levels measured by RT-qPCR in roots of WT plants vs. *clf* and *lhp1* mutants grown at 22 °C treated or not for 24h at 10 °C.

D. Quantification of RH length of plants WT, *lhp1* and *clf* at 22°C and 10°C. Each point is the mean ±error of the length of the 10 longest RHs identified in the root elongation zone in a single root. Representative

835 images of each genotype are shown on the right. The scale bas represent 500μm. Asterisks indicate sig-
836 nificant differences between WT and the corresponding genotype at the same temperature or significant
837 differences between temperature treatments on the same genotype at the same temperature (one way
838 ANOVA followed by a Tukey-Kramer test; "**"<0.01, "***"<0.001).
839 In B and C, the error bars represent the SD of 3 biological replicates. The asterisks indicate that the dif-
840 ference are significant (t test p<0.05).
841
842

843
844
Supplementary Figure 5: WRKY42:GFP is localized in the cell nucleus, regardless the expression levels
of *APOLO*.
Confocal observation of the GFP tag of WRKY42, transiently expressed in leaves of WT, *35S:APOLO* and
RNAi-*APOLO* plants. Samples were excited at 488 nm (Intensity=8%) and detected at 495-530 nm for GFP

and 610-670 (gain 650) nm for chlorophyll fluorescence. Images were captured using 10X and 20X lenses

and the scale bars are indicated accordingly. The images show a high efficiency of leaf transformation for

every genetic background.

852
Supplementary Figure 6: Root hair density is not altered by the deregulation of *APOLO*, *WRKY42*, *LHP1*
or *CLF*.
Root hair density was determined as described in the Materials and Methods section. All the lines used in
this work exhibit a similar density (one way ANOVA followed by a Tukey-Kramer test; not significant) un-
der control conditions as well as in response to cold treatment. Each point is the mean of the length of
the 10 longest RHs identified in a single root.

859
860

LEGENDS TO SUPPLEMENTARY TABLES

Supplementary Table 1: *APOLO* targets related to root hair growth.

The list of *APOLO bona fide* targets was determined in (Ariel et al., 2020). The search for root hairs-

related genes was performed based on (Won et al., 2009; Yi et al., 2010; Vijayakumar et al., 2016; Hwang

et al., 2017). LHP1 binding and H3K27me3 deposition was investigated in previously published datasets

(Veluchamy et al., 2016). W-boxes (TTGACY;(Dong et al., 2003)) present in the first 2500 bp upstream

the ATG of each gene was searched manually. The position of box is given only for *RHD6*, directly recog-

nized by WRKY42 (**Figure 4C**).

Supplementary Table 2: Root hair-related genes deregulated in *35S:APOLO* seedlings.

The list of deregulated genes was determined by RNA-Seq (Ariel et al., 2020). The search for root hairs-

related genes was performed based on (Won et al., 2009; Yi et al., 2010; Vijayakumar et al., 2016; Hwang

et al., 2017). Genes are classified as directly (same as in **Supplementary Table 1**) or indirectly regulated

by *APOLO*. *RHD6*, *RSL2* and *RSL4* are highlighted in red.

Supplementary Table 3: List of oligonucleotides used in this work.