A highly expressing, soluble, and stable plant-made IgG fusion vaccine strategy enhances

antigen immunogenicity in mice without adjuvant

Andrew G. Diamos*, Mary D. Pardhe, Haiyan Sun, Joseph G. L. Hunter, Jacquelyn Kilbourne,

Qiang Chen, and Hugh S. Mason*

Center for Immunotherapy, Vaccines, & Virotherapy, Biodesign Institute at ASU; and School of

Life Sciences, Arizona State University, Tempe, AZ 85287

*To whom correspondence should be addressed:

Email: Hugh.Mason@asu.edu; adiamos@asu.edu

Phone: 480-727-8228

Highlights

- A modified immune complex has high expression, solubility, stability, and immunogenicity.
- Antigen immunogenicity is improved up to 150-fold by fusion to plant-made IgGs.
- High serum IgG titers >1:500,000 were achieved with only two doses without adjuvant.

Summary

Therapeutics based on fusing a protein of interest to the IgG Fc domain have been enormously successful, though fewer studies have investigated the vaccine potential of IgG fusions. In this study, we systematically compared the key properties of seven different plant-made human IgG1 fusion vaccine candidates using Zika virus (ZIKV) envelope domain III (ZE3) as a model antigen. Complement protein C1q binding of the IgG fusions was enhanced by: 1) antigen fusion to the IgG N-terminus; 2) removal of the IgG light chain or Fab regions; 3) addition of hexamer-inducing mutations in the IgG Fc; 4) adding a self-binding epitope tag to create recombinant immune complexes (RIC); or 5) producing IgG fusions in plants that lack plant-specific β1,2-linked xylose and α1,3-linked fucose N-linked glycans. We also characterized the expression, solubility, and stability of the IgG fusions. By optimizing immune complex formation, a potently immunogenic vaccine candidate with improved solubility and high stability was produced at 1.5 mg IgG fusion per g leaf fresh weight. In mice, the IgG fusions elicited high titers of Zika-specific antibodies which neutralized ZIKV using only two doses without adjuvant, reaching up to 150-fold higher antibody titers than ZE3 antigen alone. We anticipate these findings will be broadly applicable to the creation of other vaccines and antibody-based therapeutics.

Keywords

Zika virus; vaccine; envelope; domain III; IgG fusion; recombinant immune complex; C1q;

transient expression; plant-made

Abbreviations

RIC, recombinant immune complex; ZIKV, Zika virus; ZE3, Zika virus envelope domain III; Fc,

the C-terminal fragment of crystallization of IgG; Fcγ, Fc from immunoglobulin G; Fab, the

antigen-binding fragment of IgG; 6D8, a human IgG1 monoclonal antibody targeting a linear

epitope on Ebola virus glycoprotein 1; C1q, the first component of the classical complement

pathway; ADCC, antibody-dependent cellular cytotoxicity; CDC, complement dependent

cytotoxicity; FcRn, neonatal Fc receptors.

1. Introduction

Subunit vaccines consisting of recombinant protein antigens are very promising due to

their safety, ease of production, and capacity to elicit targeted immune responses directed

towards desired epitopes. However, when delivered by themselves, these antigens often fail to

generate robust and long-lasting immune responses, necessitating strategies to enhance their

immunogenicity (Reed et al., 2013). Protein fusions to a variety of bioengineered IgG-based

scaffolds have demonstrated tremendous potential as therapeutic candidates, acting to enhance

the solubility and stability of the fusion partner while also allowing simple and cost-effective

purification via protein A/G affinity chromatography (Carter, 2011). Furthermore, by interacting

with neonatal Fc receptors (FcRn) in the body, IgG fusions can escape lysosomal degradation,

thereby extending the serum half-life of the Fc-fusion (Roopenian and Akilesh, 2007).

Though less well explored, IgG fusions also have many desirable characteristics for

vaccine development. Antigen-presenting cells can uptake and process IgG-bound antigen via

FcRn receptors (Baker et al., 2011; Qiao et al., 2008), complement receptor C1q (Fletcher et al.,

2018; McCloskey et al., 2011; West et al., 2018) and Fcγ receptors (Bournazos and Ravetch,

2017). However, not all IgG subclasses interact equally with immune receptors. IgG1 and IgG3

are immunostimulatory, while IgG2 and IgG4 have weak effector functions (Vidarsson et al.,

2014). IgG3 mediates the strongest complement activation and Fc receptor binding, however it

has significantly reduced serum half-life, and thus most IgG-fusions vaccine strategies use

human IgG1 (Czajkowsky et al., 2012). In addition to these factors, the polymeric state of an

antibody also affects its immunological properties. Both C1q, the first component involved in

complement activation, and the low affinity Fcγ receptors require high avidity binding for

activation. For this reason, monovalent Fc-antigen fusions cannot efficiently utilize these

pathways. By contrast, larger antigen-antibody immune complexes with multivalent Fc domains

can efficiently bind C1q and cross-link Fc receptors, resulting in greatly improved uptake and

presentation by dendritic cells, and subsequently improved activation of T-cells (Fletcher et al.,

2018; Getahun and Heyman, 2006; Ho et al., 2017; van Montfoort et al., 2012).

Capitalizing on the benefits of antigen-antibody immune complexes, a variety of vaccine

strategies have been developed to improve antigen immunogenicity. Immune complexes have

been used to focus the immune response towards favorable antigenic sites (Hioe et al., 2018;

Tsouchnikas et al., 2015) as well as enhance the potency and breadth of the immune response by

selecting for high affinity B-cells (Maamary et al., 2017). By engineering antigen-IgG fusions to

form polymeric structures via inclusion of the IgM tailpiece, both humoral and cellular immune

responses were enhanced in the absence of adjuvant (Kim et al., 2018; Webster et al., 2018).

While traditional immune complexes consisting of antigen mixed with antibody will form

variable structures depending on the oligomeric status of the antigen, we and others have

developed a recombinant immune complex (RIC) strategy that allows consistent formation of

large immune complexes which can efficiently engage complement (Mason, 2016). In the RIC

system, a well-characterized antibody is tagged with its own binding site and is fused to an

antigen of choice. The resulting molecules self-interact and form highly immunogenic antigen-

antibody clusters. Past research has shown the potential for RIC-like molecules to produce

promising vaccine candidates for *Clostridium tetani* (Chargelegue et al., 2005), Ebola virus

(Phoolcharoen et al., 2011a, 2011b), *Mycobacterium tuberculosis* (Pepponi et al., 2014), dengue

virus (Kim et al., 2015), human papillomavirus (Diamos et al., 2019) and Zika virus (Diamos et

al., 2020b).

Zika virus (ZIKV) envelope domain III (ZE3) is a promising subunit vaccine candidate

due to its ability to generate neutralizing antibodies without potential for antibody-dependent

enhancement (Shukla et al., 2020; Stettler et al., 2016; Yang et al., 2017). However, like many

subunit vaccines consisting of viral antigens, it is not strongly immunogenic on its own,

necessitating high doses with adjuvant and repeated immunizations (Yang et al., 2017). We have

previously shown that RIC carrying ZE3 elicits strong immune responses (Diamos et al., 2020b).

However, traditional RIC may be undesirable, as the large complex size renders them poorly

soluble and limits their viability as a vaccine candidate (Diamos *et al.*, 2019). Therefore, in this

study we built on recent advances in antibody engineering to design and explore a panel of

modified IgG fusion vaccines carrying ZE3 as a model antigen. These constructs contained

modifications designed to alter their polymeric and C1q binding characteristics. We then

compared the C1q binding, expression, solubility, stability, and immunogenicity of the IgG

fusions.

2. Results

2.1 Modified plant-made IgG-ZE3 fusions exhibit variable binding to complement receptor

C1q

We designed a panel of human IgG1 variants based on the previously characterized

humanized anti-Ebola monoclonal antibody 6D8 (Phoolcharoen et al., 2011a) fused to ZE3 (**Fig.**

1A). A RIC construct was created by fusing 6D8 epitope-tagged ZE3 via a flexible linker to the

C-terminus of the 6D8 heavy chain. When in solution, the 6D8 Fab arms of one 6D8-ZE3 fusion

molecule bind the epitope-tagged ZE3 on another fusion molecule, forming antigen-antibody

clusters (Mason, 2016). This RIC construct is referred to as "HLZe" as it contains the 6D8

Heavy chain, 6D8 **L**ight chain, C-terminal **Z**E3 fusion, and 6D8 **e**pitope binding tag. As a

control for immune complex formation, an otherwise identical construct was created without the

epitope tag (construct "HLZ"). This construct would not be expected to form immune complexes

as it cannot specifically bind other HLZ molecules. The 6D8 antibody without ZE3 or epitope

tag is referred to as construct "HL," and the corresponding RIC with epitope tag but lacking ZE3

is referred to as "HLe." Since C1q mediates immune complex uptake into antigen presenting

cells and complement-coated immune complexes play a key role in activating and maintaining

long-term immunity (Croix et al., 1996; McCloskey et al., 2011; Phan et al., 2007; West et al.,

2018), the constructs were analyzed for C1q binding. Additionally, C1q binding provides insight

into the polymeric status of the constructs, as RIC and other polymeric IgGs have stronger

binding to C1q when compared with monomeric IgG (Chargelegue et al., 2005; Webster et al.,

2018). To improve C1q binding, all constructs were expressed in glycoengineered *Nicotiana*

benthamiana that have suppressed fucosyl- and xylosyl transferase activities, as these glycans

inhibit immune receptor binding (Kallolimath and Steinkellner, 2015). HL or HLe made in

glycoengineered plants showed highly improved C1q binding compared to constructs expressed

in wildtype plants (**Fig. S1A**). Consistent with the formation of densely clustered antigen-

antibody complexes, HLZe showed greatly improved C1q binding compared to HLZ (p < 0.001)

(**Fig. 1B**). A small increase in C1q binding was noted with HLZ compared to HL (p < 0.05) (**Fig.**

1B), suggesting low-level aggregation of the construct or slight favorable alteration of Fc

conformation due to the ZE3 fusion. HLZe also efficiently bound FcγRIIIa, exceeding the

binding of HLZ and HL (**Fig. S1B**).

Hexamers are the ideal substrate for C1q binding (Diebolder et al., 2014) and it has been

suggested that the antibody Fab arms play a regulatory role in complement activation by

inhibiting C1q binding (Wang et al., 2016). Upon addition of certain soluble antigens or removal

of the light chains, C1q binding is dramatically improved, suggesting conformational changes are

conveyed into the Fc domain from the Fab arms to induce hexamer formation (Wang et al.,

2016). In agreement with this model, we find that the addition of soluble antigen carrying the

6D8 epitope tag modestly improved C1q binding (**Fig. S2**, compare HL vs HL + Ag) and C1q

binding was strongly enhanced by removal of the antibody light chain (**Fig. S2**, compare HL and

H). Interestingly, antigen fusion to the heavy chain N-terminus also strongly improved C1q

binding at a level similar to removing the light chain (**Fig. S2**, compare HL, H, and ZHL).

Therefore, we created a construct with ZE3 fused to the N-terminus of the heavy chain (construct

"ZH") in the absence of the light chain. ZH efficiently bound C1q similar to HLZe; however, the

effects of light chain removal and N-terminal ZE3 fusions were not additive (**Fig. 1B**). In an

attempt to further improve interaction with C1q, we made a construct that was similar to ZH

except for mutations E345R, E430G, and S440Y in the Fc region which favor formation of

hexamers (Diebolder et al., 2014). This construct (ZHx) had modestly improved C1q binding

compared to ZH (p < 0.001) (**Fig. 1B**). Past studies have shown that IgG1 Fc fusions (lacking

VH and CH1 domains) produce variable levels of C1q binding depending on the fusion partner

(Lagassé et al., 2019) . However, Fab arm removal can favor hexamer formation with strong

complement activation (Wang et al., 2016). ZE3 N-terminally fused to the 6D8 Fc (**Fig. 1A**

construct "ZFc") showed very strong C1q binding (**Fig. 1B**). Finally, as RIC require co-

expression of heavy and light chains, we constructed a simplified RIC that consisted of a single

chain antibody with the variable light (VL) domain of 6D8 inserted between the variable heavy

(VH) and CH1 domains of 6D8 to yield construct "HVL". This construct bound antigen tagged

with the 6D8 epitope, albeit at a somewhat reduced level compared to unaltered 6D8 (**Fig. S3**).

This single-chain Fab configuration was fused to ZE3 to create the single chain RIC HVLZe

(**Fig. 1A**), which displayed strong C1q binding (**Fig. 1B**). Together, these data demonstrate that

RIC formation, removal or rearrangement of light chain/Fab region, Fc hexamer mutations, and

ZE3 fusion to the IgG N-terminus all increase C1q binding of plant-made IgG1.

2.2 Reducing RIC self-binding improves solubility without reducing C1q binding

RIC suffer from low yield of soluble product (Diamos et al., 2019). While unmodified

6D8 antibody is highly soluble, addition of the short epitope tag (construct HLe) renders the

antibody mostly insoluble (**Fig. S4**). However, this is prevented by removal of the light chain

(construct He), which is needed for epitope binding (Kim et al., 2015), suggesting that the

insolubility arises from large complexes of antibody bound to the epitope tag (**Fig. S4**). To

improve RIC solubility, we shortened the 6D8 epitope tag in order to reduce the strength of self-

binding. Construct "HLa" contains the minimal reported binding region for 6D8 "VYKLDISEA"

(Wilson et al., 2000). Removal of a single amino acid from the C-terminus (construct "HLb,"

epitope VYKLDISE) had no effect on solubility (**Fig. S5A**). However, further removal of a

single amino acid from the N-terminus (construct "HLc," YKLDISE), and additional truncation

in construct "HLd" (YKLDIS) resulted in greatly improved solubility, yielding 20-fold more

soluble antibody (**Fig. S5B**). Thus, the truncated epitope present in HLd was introduced to HLZe

(construct HLZd) and characterized. Despite reducing epitope binding compared to full length

epitope tag by approximately 25-fold by ELISA (**Fig. S5C**), HLZd still maintained very strong

C1q binding, suggesting efficient complex formation still remained (**Fig. 1B**). Note that the

otherwise identical construct HLZ, which lacks the YKLDIS tag, had 10-fold lower C1q binding

(**Fig. S5D**). These data suggest that RIC solubility can be enhanced by reducing self-binding

without a loss of C1q activation.

2.3 Highly variable expression and solubility of IgG fusions in plants

Since a viable vaccine candidate must be soluble and highly expressing, we measured the

soluble yield of fully assembled IgG fusions by employing an ELISA assay that first captured

ZE3 and then detected human IgG. As a standard, purified and quantified HLZ was used. To

detect any proteolytic cleavage of ZE3, an ELISA measuring total IgG irrespective of the

presence of ZE3 was also used as a comparison, using purified and quantified unfused HL as

standard. When soluble extracts were probed for the presence of both ZE3 and IgG, construct ZH

yielded 0.83 mg fully formed product per gram leaf fresh weight (mg/g LFW), and the similar

ZFc yielded 0.58 mg/g LFW (**Fig. 2A**). However, when measuring the total IgG content, the

yield of ZH was roughly 1.2-fold higher and, for ZFc, 2.5-fold higher than ZE3 ELISA, which

suggests that ZH and especially ZFc are susceptible to proteolytic cleavage. This finding was

confirmed by visualization of the ZFc cleavage products on SDS-PAGE (**Fig. 2B**). HLZ

accumulated only 0.17 mg/g LFW fully assembled product, with roughly 40% ZE3 lost to

cleavage, suggesting general instability of the construct (**Fig. 2A, 2B**). By ELISA, the polymeric

RIC constructs seemingly had lower yields than the monomeric constructs: HLZe and HLZd

only accumulated 0.04 mg/g LFW and 0.30 mg/g LFW respectively, and the hexameric ZHx

accumulated only 0.08 mg/g LFW (**Fig. 2A**). However, when visualized under reducing SDS-

PAGE conditions, HLZd was the highest expressing construct, accumulating an estimated 1.5

mg/g LFW, over twice the highest yield of the next best construct, with no visible degradation

products (**Fig. 2A, 2B**). The discrepancy between ELISA and gel quantification likely arises due

to complexed HLZd being less accessible to the antibody probe by ELISA. Importantly, the gel

quantification results agreed with the ELISA results for all the monomeric constructs, but not for

any of the polymeric constructs (**Fig. 2A**). Thus, the total yield of HLZe and ZHx was also

higher when measured by gel quantification (**Fig. 2A, 2B**). While polymeric constructs were

partially inaccessible to the antibody probe by traditional ELISA (**Fig. 2A**), they were strongly

detected using the same antibody probe in the C1q assay (**Fig. 1B**). This likely occurred due to

C1q disaggregating the polymeric constructs, as C1q functions to solubilize immune complexes

in vivo (Thielens et al., 2017). Construct HVLZe yielded only 0.02 mg/g LFW soluble product

by ELISA and was not visible by SDS-PAGE (**Fig. 2A**). However, we detected very high levels

when extracting with 7.5 M urea (data not shown) suggesting that the construct insolubility

resulted in the low yield. In total, these results show that the different IgG fusions display

expression levels which vary by several orders of magnitude, highlighting the large differences

in their capacity to be used as viable, affordable vaccines.

2.4 Purification and aggregation of IgG fusion constructs

IgG fusions were purified to >95% homogeneity using a simple one-step purification via

protein G affinity chromatography. In agreement with the expression data, the more polymeric

constructs showed less degradation than the other constructs, and the ZFc fusion had particularly

high levels of degradation (**Fig. 3**). To investigate the polymeric characteristics of each construct,

purified IgG fusions were analyzed by sucrose gradient sedimentation. Consistent with the

formation of large immune complexes, HLZe and HVLZe were found mostly in the bottom of

the gradient while HLZ, ZH, and ZFc were found mostly at the top of the gradient (**Fig. 4A, 4B**).

A small shift in density was noted with construct HLZ (**Fig. 4A**) compared to unfused HL, which

may explain the increased C1q binding (**Fig. 1B**). Soluble extracts of ZHx contained both low-

density material as well as some very high-density material, suggesting that the combination of

light chain removal and RGY mutations contributes to the formation of larger aggregates (**Fig.**

4B). Extracts of HLZd showed intermediate density compared to HLZ and HLZe (**Fig. 4A**)

which, when taken together with the expression data and C1q binding, are consistent with HLZd

forming smaller, more soluble immune complexes compared to HLZe.

2.5 Polymeric constructs have improved stability

Producing vaccines which are stable at refrigerated temperatures, ambient temperatures,

and after freeze/thawing, is a major challenge of the vaccine industry (Kumru et al., 2014). First,

the stability of each construct was analyzed by comparing fully formed products and degradation

products on SDS-PAGE gels after treatment with various temperature conditions. The initial

level of degradation, which corresponds to any degradation that occurred during expression,

purification, or the initial freeze/thaw, was substantially reduced (20-40% more fully formed

product) in polymeric constructs when compared to monomeric constructs (**Fig. 5A**). After five

freeze-thaw cycles or two weeks at 4°C, small amounts of degradation (2-5%) were observed

with all constructs (**Fig. 5A**). High concentrations (>1 mg/ml) of HLZe and HVLZe precipitated

after several days at 4°C (data not shown), probably due to the formation of very large insoluble

complexes, thereby highlighting the undesirability of these constructs. At room temperature,

most constructs had 10-15% degradation after two weeks (**Fig. 5A**). Overall, the polymeric

constructs retained the highest stability, while the monomeric constructs, and especially the Fc

fusion, degraded more rapidly (**Fig. 5A**).

Second, we determined whether storage resulted in an impairment of C1q binding. Most

constructs retained strong C1q binding after storage; however, after room temperature storage,

repeated freeze-thaw cycles, but not 4°C storage, construct HLZ unexpectedly displayed

increased C1q binding (**Fig. 5B**). This may be due to aggregation, degradation of the Fab

regions, or loss of light chain as numerous degradation products are visible on SDS-PAGE (**Fig.**

3). Conversely, construct ZFc lost C1q binding ability as it became more degraded, especially

after freeze-thaw cycles, suggesting degradation of the Fc binding region (**Fig. 5B**). Taken

together with the expression data, constructs with intermediate levels of self-binding (e.g. HLZd

and ZHx) showed the highest stability during both extraction and storage while maintaining high

solubility. Of these, only construct HLZd could be produced at high levels.

2.6 IgG fusion enhances ZE3 immunogenicity

IgG fusions have strong immunological properties even in the absence of adjuvant due to

their improved binding of C1q and uptake via FcRn and Fc receptors on antigen presenting cells,

(Czajkowsky et al., 2012). To investigate the immunogenicity of the IgG fusions created here,

BALB/c mice (n = 6) were immunized subcutaneously, without adjuvant, with two doses of each

IgG fusion construct such that each dose of ZE3 delivered was 8 μg. As a control, mice were also

immunized with 8 μg His-tagged plant-expressed ZE3. As expected, all IgG fusions very

strongly enhanced the production of ZE3-specific IgG, producing 20-fold to 150-fold higher total

IgG titers than ZE3 alone (**Fig. 6A**, p < 0.01 compared to ZE3). 5-fold to 10-fold higher mean

titers were reached by the ZHx and ZFc constructs compared to the other constructs, though

these differences were not statistically significant. The level of IgG2a antibodies were also

measured because they have important antiviral effector functions (Lu et al., 2018) and are

correlated with T-cell activation (Huber et al., 2006) and complement activity (West et al.,

2018). All IgG fusions significantly enhanced the production of IgG2a compared to ZE3 alone

(**Fig. 6B**, $p < 0.01$ compared to ZE3). The construct HLZ, which displayed the lowest C1q

binding (**Fig. 1B**), had a significantly reduced production of IgG2a compared to most other

fusions (**Fig. 6B**, $p < 0.05$ compared to HLZ). Compared to total IgG, the relative ratio of IgG2a

was lowest in the ZE3, HLZ, and ZFc groups (**Fig. 6B**, see table). Sera from mice immunized

with all IgG fusions neutralized ZIKV (**Fig 7,** $p < 0.0001$ compared to sera from PBS-injected

mice). All IgG fusions elicited significantly higher neutralizing responses than His-tagged ZE3

(**Fig 7,** $p < 0.024$ compared to ZE3).

3. Discussion

Previous work has shown that IgG fusion strategies have the potential to enhance antigen

immunogenicity (Kim et al., 2018; Konduru et al., 2011; Loureiro et al., 2011; Webster et al.,

2018; Zhao et al., 2018). However, the key characteristics to produce an optimal IgG fusion

vaccine candidate have not been directly compared. An optimal vaccine candidate must be

accessible, affordable, and stable, as well as immunogenic. Impressively, especially given the

low doses and lack of adjuvant used in this study, the IgG fusions were potently immunogenic

with many of the best candidates eliciting 100-fold higher antibody responses compared to

antigen alone. While the highest immunological readouts were obtained by constructs ZHx and

ZFc, these constructs have limited use as vaccine candidates, as ZHx suffered from low yield and

aggregation, while ZFc was extraordinarily unstable prior to and during storage. These findings

underscore the importance of evaluating multiple aspects of vaccine viability. By contrast,

construct HLZd had nearly double the yield of the next closest construct (**Fig. 2**) while

maintaining high stability (**Fig. 5**), solubility (**Fig. S5B**), and immunological properties (**Figs.**

1B, 6, 7). All of the HLZd produced in this study was made in a single small *N. benthamiana*

plant (data not shown), demonstrating the excellent potential of plant-based vaccine development

to address vaccine affordability and accessibility. The overall properties of each construct are

summarized in Table 1. These findings warrant further development of HLZd-like constructs as a

vaccine platform which we anticipate will be broadly applicable to the development of vaccines

targeting other pathogens.

While most current antibody fusion vaccines use constructs similar to ZFc, many

properties of Fc fusions, including C1q binding, vary based on the individual fusion partner and

thus must be determined empirically for each fusion (Lagassé et al., 2019). Interestingly, some

Fc fusion constructs form hexamers (Zhang et al., 2019), which may explain the strong C1q

binding and immunogenicity of ZFc observed here. Despite its strong immunogenicity, ZFc was

remarkably unstable: ~50% of the ZE3 was proteolytically cleaved either before or during

extraction (**Fig. 2A**, **2B**), even when directly extracted in SDS sample buffer or with protease

inhibitors (data not shown). Further, less than 25% of full-size ZFc molecules remained intact

upon repeated freeze-thaw or storage (**Fig. 5A**). This degradation was associated with a loss in

C1q binding (**Fig. 5B**), probably due to impairment of the Fc receptor binding domains. ZFc has

fewer disulfide bonds than whole IgG, and in general Fc fusions often suffer from instability or

undesirable aggregation (Zeng et al., 2018). Further protein engineering, including optimization

of linkers, may improve the stability and expression of these constructs. In contrast, IgG fusions

displaying self-binding properties, such as HLZd, HLZe, and ZHx, were much more stable

during extraction and storage. Consistent with their reduced binding to antibody probes during

ELISA (**Fig. 2A**), the intramolecular and intermolecular associations of these constructs may

protect them from proteolysis or other degradative processes.

The large complex size generated by traditional RIC (i.e. HLZe) renders them poorly

soluble upon extraction (**Figs. 2A/S2,** Diamos et al., 2019) with concentrations above 1-2 mg/ml

precipitating during storage (data not shown). Therefore, we reduced the binding affinity of the

RIC by mutating the 6D8 epitope tag (construct HLZd). Using this strategy of modifying the

strength of self-binding, RIC can be customized to create complexes of different sizes. These

findings will allow future investigation of the immunological and functional properties conferred

by variably sized immune complexes. While traditional RIC require co-expression of both the

heavy and light chains, this can be simplified by creating a single chain antibody with the

variable light domain fused to the variable heavy domain, as in construct HVLZe. This construct

had high stability (**Fig. 5**) and immunogenicity (**Fig. 6**), but was poorly soluble, resulting in low

recovery (**Fig. 2**). By further reducing the epitope binding of HVLZe or by altering the relative

positions of the light chain or antigen fusions, or by using the shorter "d" epitope tag, it is likely

the solubility can be improved.

The glycosylation state of the Fc strongly controls its function and can enhance or inhibit

binding to immune receptors by modulating the stability, conformation, and aggregation of the

Fc (Mastrangeli et al., 2019). These alterations result in important differences in antibody

effector functions, including antibody-dependent cellular cytotoxicity, antibody-dependent cell-

mediated phagocytosis, complement-dependent cytotoxicity, and antibody-dependent

enhancement of viral infection (Sun et al., 2018). Advances in glycoengineering have allowed

targeted optimization of the Fc glycosylation state in a variety of recombinant expression

systems (Gupta and Shukla, 2018; Kallolimath and Steinkellner, 2015; Wang et al., 2018). For

example, an anti-CD20 antibody produced in glycoengineered plants lacking core xylose and

fucose N-linked glycans showed improved binding to FcγRI, FcγRIIIa, and C1q (Marusic et al.,

2018). Similarly, anti-DENV antibodies produced in glycoengineered plants have been shown to

have diminished ADE activity and, consequently, have superior efficacy and safety profiles than

their mammalian cell-produced counterparts (Dent et al., 2016). Antibody therapeutics made in

glycoengineered plants have also been used to successfully treat rhesus macaques and humans

with Ebola (Lyon et al., 2014; Olinger et al., 2012; Qiu et al., 2014; Zeitlin et al., 2011), HIV

(Forthal et al., 2010; Ma et al., 2015; Stelter et al., 2020), and Chikungunya virus disease

(Hurtado et al., 2019). In agreement with these studies, we found that C1q binding was enhanced

in glycoengineered plants in a manner which acted synergystically with immune complex

formation (**Fig. S1A**). As the entire mamallian sialylation pathway has been reconsistuted in

glycoengineered plants (Castilho et al., 2012), and sialylated immune complexes yield higher

affinity antibodies (Maamary et al., 2017), the IgG fusion vaccines used in this study may be

improved by further glycan modification.

Despite its theoretically monomeric nature, construct HLZ elicited high titers and

comparable neutralization to the more polymeric constructs. HLZ produced a small but

repeatable improvement in C1q binding (**Fig. 1B**) and a shift in small shift in density (**Fig. 4A**,

compare HL and HLZ) which may suggest some low-level aggregation due to the ZE3 fusion.

Nonetheless, the expression of HLZ was generally low, suggesting C-terminal ZE3 fusion may

interfere with proper folding or otherwise cause instability (**Fig. 2A**). Many degradation products

are visible by SDS-PAGE following purification (**Fig. 3**) and, interestingly, the C1q binding of

HLZ continued to increase upon further degradation by incubation for 2 weeks at RT or repeated

freeze-thaw cycles (**Fig. 5B**). These data suggest that the degradation products may be more

highly immunogenic than the original construct, as light chain or Fab removal strongly improves

C1q binding (**Fig. S2**). This finding may have also contributed to the relatively high

immunogenicity produced by HLZ (**Fig. 6A**). However, despite similar total IgG titers and ZIKV

neutralization, the relative IgG2a ratio was significantly reduced for HLZ, ZE3, and ZFc when

compared to the more polymeric constructs (**Fig. 6B**). Since IgG2a antibodies are a general

indicator of a Th1 response (Huber et al., 2006) and complement activation is involved in T-cell

immunity (West et al., 2018), these constructs may have reduced T-cell activation compared to

more polymeric constructs. This is important because T-cell activation and complement

activation are vital components to generate long-term B-cell immunity (Akkaya et al., 2020;

West et al., 2018). *In vitro* neutralization (**Fig. 7**) does not account for antibody subclass effector

functions (e.g. IgG2a) or cellular immunity, thus further investigations are warranted.

Another point to consider is that, at least for certain antibody/antigen combinations,

antigen binding induces conformational changes in IgG, resulting in improved hexamer

formation and subsequently improved C1q binding (Wang et al., 2016). Mixing 6D8 with an

antigen containing the 6D8 epitope produced only a small increase in C1q binding (**Fig. S2**).

However, antigen fusion to the 6D8 N-terminus had a more pronounced effect on C1q binding

(**Fig. 1**), perhaps by strongly inducing conformational changes similar to antigen binding which

allow hexamer formation (Diebolder et al., 2014). In general, we have found that antigen fusion

to the 6D8 N-terminus greatly enhances C1q binding for a variety of large and small antigens

(data not shown). Furthermore, removal of the 6D8 light chain (construct ZH) also substantially

enhanced C1q binding (**Fig. 1**). These findings agree with the hypothesis that the Fab portions of

the antibody plays a regulatory role in C1q binding. Additionally, removal of the light chain may

also impact glycosylation and thus C1q binding (He et al., 2014).

In summary, we have developed a cheap, fast, and efficient plant expression system to

produce and purify high levels of IgG fusion vaccine candidates. Plant recombinant expression

systems are particularly well suited to make IgG fusion vaccines, such as the ones described in

this study, since they have inherent safety, high scalability, and low production costs when

compared to mammalian cell systems (Alam et al., 2018; Buyel, 2019; Chen and Davis, 2016;

Gleba et al., 2014). These benefits can be of special value for vaccine production in developing

countries (Ma et al., 2013). By directly comparing many different IgG fusion strategies, we have

identified a self-assembling immune complex construct HLZd that is highly expressing, stable,

soluble, and immunogenic. These studies warrant future research in animal models, and we

anticipate our findings will be broadly applicable to other vaccine antigens or antibody-based

therapeutics.

4. Experimental Procedures

4.1 Vector construction

The construction of a geminiviral replicon plant expression vector for ZE3, as well as its

fusion to the 6D8 C-terminus (pBYR11eM-h6D8ZE3, referred to here as construct "HLZe") or

N-terminus with epitope tag (pBYR11eMa-BAZE3-Hgp371) or without epitope tag

(pBYR11eMa-BAZE3-H) have been previously described (Diamos et al., 2020b, 2020a) A

vector pBYKEMd2-6D8 expressing the full 6D8 mAb without ZE3 fusion (construct "HL") has

been previously described (Diamos et al., 2020b, 2020a). To create a vector expressing only the

light chain of 6D8, pBYKEMd2-6D8 was digested with XhoI and the vector was self-ligated to

yield pBYKEMd-6D8K. A vector expressing only the heavy chain of 6D8 (construct "H") was

created by digesting pBYKEMd2-6D8 with SacI and self-ligating the vector, to yield

pBYKEMd-6D8H. The 6D8 epitope binding tag was added to pBYKEMd-6D8H by digesting

pBYR11eMa-BAZE3-Hgp371 with BsaI-SacI and inserting the tag-containing fragment into

pBYKEMd-6D8H digested with BsaI-SacI, yielding pBYKEMd-6D8Hgp371 (construct "HLe"

when coexpressed with light chain). To remove the epitope tag from HLZe, pBYR11eM-

h6D8ZE3 was digested with BamHI-SacI and ligated with a fragment containing ZE3 obtained

via amplification with primers ZE3-Bam-F (5'-gcgggatccaagggcgtgtcatactcc) and ZE3-Sac-R

(5'-acagagctcttaagtgctaccactcctgtg) and subsequent digestion with BamHI-SacI. The resulting

vector, pBYKEMd-HZE3, was coinfiltrated with pBYKEMd-6D8K to produce construct "HLZ."

To produce ZE3 fused to the 6D8 N-terminus without light chain, pBYR11eMa-BAZE3-H was

digested with SacI and the vector vector was self-ligated, yielding pBYKEMd-ZE3H (construct

"ZH"). To introduce hexamer mutations, a region of the 6D8 heavy chain constant region was

synthesized (Integrated DNA Technologies, Iowa, USA) containing the E345R, E430G, and

S440Y mutations, then digested with BsaI-SacI and used to replace the BsaI-SacI region of 6D8

in pBYKEMd-ZE3H, yielding pBYKEMd-ZE3Hx (construct "ZHx"). RIC epitope tag mutant

"a" was generated by annealing oligos 6D89-F (5'-ctagtgtttacaagctggacatatctgaggcataagagct) and

6D89-R (5'- cttatgcctcagatatgtccagcttgtaaaca) and ligating them into pBYR11eM-h6D8ZE3

digested SpeI-SacI; mutant "b" was generated by first amplifying mutant "a" with primers

gpDISE-Sac-R: (5'-tttgagctcttactcagatatgtccagcttgtaaac) and 35S-F (5'aatcccactatccttcgc), then

digesting the product with SpeI-SacI and ligating it into pBYR11eM-h6D8ZE3 digested with

SpeI-SacI. Mutants "c" and "d" were created similarly to mutant "a" using overlapping oligos

6D87-F (5'-ctagttacaagctggacatatctgagtaagagct) and 6D87-R (5'-cttactcagatatgtccagcttgtaa) for

"c" and 6D86-F (5'-ctagttacaagctggacatatcttaagagct) and 6D86-R (5'-cttaagatatgtccagcttgtaa).

In order to make a construct in which the variable heavy (VH) domain is linked to a

variable light chain (VL) domain that, in turn, is directly fused to the constant region of the 6D8

antibody, the variable regions were first obtained through PCR amplification and end-tailoring of

segments from pBYR11eM-h6D8ZE3. For the VH domain, the primers LIR-H3A (5'-

aagcttgttgttgtgactccgag) and 6D8VH-Spe-R (5'- cggactagtagctgaagacactgtgac) were used. The

VL region was obtained through PCR amplification of pBYR11eM-h6D8ZE3 with primers 35S-

F (5'-aatcccactatccttcgc) and 6D8VK-Nhe-R (5'-cgtgctagccttgatctccactttggtc). In order to fuse

VL region to the constant region of a human IgG antibody, a subclone was created by digesting

the PCR fragment with XhoI-NheI and inserting it into a vector, pKS-HH-gp371, that contained

the 6D8 heavy chain (Kim et al., 2015). This subclone was named pKS-VL. Next, pBYKEM-

6D8K was digested with SbfI-SacI, the PCR product that amplified the variable heavy chain

fragment was digested SbfI-SpeI, and the variable light chain subclone was digested SpeI-SacI.

These fragments were assembled to create pBYKEMd2-VHLVK (construct "HVL"). Finally,

this construct was used to create pBYKEMd2-HVLZe by a two-fragment ligation. The

pBYKEMd2-VHLVK construct was digested BsaI and SacI to obtain the vector fragment along

with the variable regions of the heavy and light chains. To obtain the ZE3 antigen segment and

the epitope tag, pBYR11eM-h6D8ZE3 was also digested BsaI-SacI. The resulting construct,

which was used to produce HVLZe, was named pBYKEMd-HVLZe.

4.2 Agroinfiltration of *Nicotiana benthamiana* leaves

Binary vectors were separately introduced into *Agrobacterium tumefaciens* EHA105 by

electroporation. The resulting strains were verified by restriction digestion or PCR, grown

overnight at 30°C, and used to infiltrate leaves of 5- to 6-week-old *N. benthamiana* maintained at

23-25°C. Briefly, the bacteria were pelleted by centrifugation for 5 min at 5,000g and then

resuspended in infiltration buffer (10 mM 2-(N-morpholino)ethanesulfonic acid (MES), pH 5.5

and 10 mM MgSO4) to OD600=0.2, unless otherwise described. The resulting bacterial

suspensions were injected by using a syringe without needle into leaves through a small puncture

(Huang and Mason, 2004). To evaluate the effects of glycosylation, transgenic plants silenced for

xylosyltransferase and fucosyltransferase were employed (Castilho and Steinkellner, 2012). Plant

tissue was harvested at 5 days post infiltration (DPI).

4.3 Protein extraction, expression and purification

Crude protein was extracted by homogenizing agroinfiltrated leaf samples with 1:5 (*w:v*)

ice cold extraction buffer (25mM Tris-HCl, pH 8.0, 125mM NaCl, 3mM EDTA, 0.1% Triton X-

100, 10 mg/mL sodium ascorbate, 0.3 mg/mL phenylmethylsulfonyl fluoride) using a Bullet

Blender machine (Next Advance, Averill Park, NY) following the manufacturer's instruction.

Homogenized tissue was rotated at 4°C for 30 min. The crude plant extract was clarified by

centrifugation at 13,000g for 15 min at 4°C and the supernatant was analyzed by SDS-PAGE or

ELISA. Alternatively, to evaluate solubility of proteins in the original homogenate, the pellet

was designated the insoluble fraction and treated with SDS sample buffer at 100˚C for 10 min

before loading on SDS-PAGE.

IgG variants were purified by protein G affinity chromatography. Agroinfiltrated leaves

were blended with 1:3 (w:v) ice cold extraction buffer (25mM Tris-HCl, pH 8.0, 125mM NaCl,

3mM EDTA, 0.1% Triton X-100, 10 mg/mL sodium ascorbate, 0.3 mg/mL

phenylmethylsulfonyl fluoride), stirred for 30 min at 4°C, and filtered through miracloth. To

precipitate endogenous plant proteins, the pH was lowered to 4.5 with 1M phosphoric acid for 5

min while stirring, then raised to 7.6 with 2M Tris base. Following centrifugation for 20 min at

16,000g, the clarified extract was loaded onto a Protein G column (Thermo Fisher Scientific,

Waltham, MA, USA) following the manufacturer's instructions. Purified proteins were eluted

with 100mM glycine, pH 2.5, directly into collection tubes containing 1M Tris-HCl pH 8.0 to

neutralize the elution buffer and stored at -80˚C. Purified protein concentration was measured by

A280 absorbance, ELISA, and gel quantification.

ZE3-His expressed from pBYe3R2K2Mc-BAZE3 was purified by metal affinity

chromatography. Protein was extracted as described above, but without acid precipitation. The

clarified extract was loaded onto a column containing TALON Metal Affinity Resin (BD

Clontech, Mountain View, CA) according to the manufacturer's instructions. The column was

washed with PBS and eluted with elution buffer (PBS, 150mM imidazole, pH 7.4). Peak ZE3

elutions were pooled, dialyzed against PBS, and stored at -80˚C.

4.5 SDS-PAGE and Western blot

Plant protein extracts or purified protein samples were mixed with SDS sample buffer (50

mM Tris-HCl, pH 6.8, 2% SDS, 10% glycerol, 0.02 % bromophenol blue) and separated on 4-

15% stain-free polyacrylamide gels (Bio-Rad, Hercules, CA, USA). For reducing conditions,

0.5M DTT was added, and the samples were boiled for 10 min prior to loading. Polyacrylamide

gels were visualized and imaged under UV light, then transferred to a PVDF membrane. For IgG

detection, the protein transferred membranes were blocked with 5% dry milk in PBST (PBS with

0.05% tween-20) overnight at 4°C and probed with goat anti-human IgG-HRP (Sigma-Aldrich,

St. Louis, MO, USA diluted 1:5000 in 1% PBSTM). Bound antibody was detected with ECL

reagent (Amersham, Little Chalfont, United Kingdom).

4.6 ELISA quantification of construct expression levels

For ZE3 ELISA, a 96-well high-binding polystyrene plate (Corning Inc, Corning, NY,

USA) was coated with a 1:500 dilution of unlabeled goat anti-human IgG (Southern Biotech,

Birmingham, AL, USA) and incubated at 37°C for 1 h. After being washed with PBST, the

plates were blocked with 5% non-fat dry milk in PBST for 30 min and washed with PBST.

Multiple dilutions, ranging from 1:100 to 1:25,000, of clarified plant extracts containing the

different constructs were added to the plate. Previously purified and quantified plant-produced

HLZ was included as a standard and positive control while samples of uninfiltrated crude extract

were added for a negative control. After a 1-h incubation, the plate was washed three times in 1x

PBST. Plates were then incubated with a 1:1000 dilution of mouse serum raised against ZE3-6H

followed by goat anti-mouse IgG HRP conjugate (Southern Biotech, Birmingham, AL, USA) for

1 h at 37°C. The plate was then washed five times with PBST, developed with TMB substrate

(Thermo Fisher Scientific, Waltham, MA, USA), and the absorbance read at 450 nm. IgG ELISA

was performed similarly, except constructs were bound directly to the plate and detected with

goat anti-mouse IgG HRP conjugate (Southern Biotech, Birmingham, AL, USA).

4.6 C1q binding

96-well high-binding polystyrene plates (Corning Inc, Corning, NY, USA) were coated

with 15 μg/ml human complement C1q (PFA, MilliporeSigma, MA) in PBS for 2h at 37°C. The

plates were washed 3 times with PBST, and then blocked with 5% dry milk in PBST for 15

minutes. After washing 3 times with PBST, purified human IgG (Southern Biotech, Birmingham,

AL, USA) and purified IgG-ZE3 fusions were added at 0.1 mg/ml with 10-fold serial dilutions

and incubated for 1.5 hours at 37°C. After washing 3 times with PBST, bound IgG was detected

by incubating with 1:1000 polyclonal goat anti human IgG-HRP (Southern Biotech,

Birmingham, AL, USA) for 1h at 37°C. The plates were washed 4 times with PBST, developed

with TMB substrate (Thermo Fisher Scientific, Waltham, MA, USA), stopped with 1M HCl, and

the absorbance was read at 450nm.

4.7 6D8 epitope binding

To test the ability of HVL to bind to the 6D8 epitope tag, 900 ng of purified dengue

consensus envelope domain III tagged with 6D8 epitope (Kim et al., 2015) were bound to a 96-

well high-binding polystyrene plate (Corning Inc, Corning, NY, USA). After a 1-hour incubation

at 37°C, the plate was washed thrice with PBST and blocked with 5% dry milk in PBST for 30

minutes. Then, the plate was washed thrice with PBST and various dilutions of either purified

HLV or full-length 6D8 antibody were added to the plate. The plate was incubated at 37°C for 1-

hour, washed thrice with PBST and detected with HRP-conjugated mouse anti-human IgG (Fc

only) (Southern Biotech, Birmingham, AL, USA) antibody at a 1:2000 dilution. Then, the plate

was thoroughly washed with PBST and developed with TMB substrate (Thermo Fisher

Scientific, Waltham, MA, USA). The absorbance was read at 450nm.

4.8 Sucrose gradient density centrifugation

Purified samples of each IgG fusion (100 µl) were loaded onto discontinuous sucrose

gradients consisting of 350 µl layers of 5, 10, 15, 20, and 25% sucrose in PBS in a 2.0 ml

microcentrifuge tubes and centrifuged at $21,000g$ for 16 h at 4 °C. Fractions were collected from

the top and analyzed by SDS-PAGE, followed by visualization on stain-free gels (Bio-Rad,

Hercules, CA, USA). The relative band intensity of each fraction was determined using ImageJ

software, with the peak band arbitrarily assigned the value of 1.

4.9 Stability of IgG fusions during storage

After purification, samples of purified IgG fusions were frozen at -80˚C. Samples were

either untreated (referred to as "initial"), or thawed and then subjected to either five additional

freeze/thaw cycles, or incubated for 2 weeks at 4°C, or incubated for 2 weeks at 23˚C. Initial and

treated samples were visually inspected for any signs of precipitation, and then analyzed by

reducing and non-reducing SDS-PAGE to observe cleavage of ZE3 or other degradation

products. ImageJ analysis was used to compare the band intensity of the fully formed product to

that of any degradation products. Each sample was also analyzed as described in section 4.6 for

C1q binding.

4.10 Immunization of mice and sample collection

All animals were handled in accordance to the Animal Welfare Act and Arizona State

University IACUC under study number 18-1616R. Female BALB/C mice, 6-8 weeks old, were

immunized subcutaneously with purified IgG fusion variants. In all treatment groups, the total

weight of antigen was set to deliver an equivalent 8 µg of ZE3. Doses were given on days 0 and

14. Serum collection was done as described (Santi et al., 2008) by submandibular bleed on days

0, 14, and 28.

4.11 Antibody measurements

Mouse antibody titers were measured by ELISA. Plant-expressed 6-His tagged ZE3 at 50

ng/well was bound to 96-well high-binding polystyrene plates (Corning Inc, Corning, NY, USA),

and the plates were blocked with 5% nonfat dry milk in PBST. After washing the wells with

PBST (PBS with 0.05% Tween 20), the mouse sera were diluted with 1% PBSTM (PBST with

1% nonfat dry milk) and incubated. Mouse antibodies were detected by incubation with

polyclonal goat anti-mouse IgG-horseradish peroxidase conjugate (Sigma-Aldrich, St. Louis,

MO, USA). The plate was developed with TMB substrate (Thermo Fisher Scientific, Waltham,

MA, USA), stopped with 1M HCl, and the absorbance was read at 450nm. Endpoint titers were

taken as the reciprocal of the lowest dilution which produced an OD_{450} reading twice the

background produced using PBS as the sample. IgG2a antibodies were measured from sera

diluted 1:100 in 1% PBSTM and detected with IgG2a horseradish peroxidase conjugate (Santa

Cruz Biotechnology, Dallas, TX, USA). IgG2a values were normalized by total IgG, and the

value obtained for ZE3 was arbitrarily defined as 1.0. The ratios for each construct were

calculated as IgG2a:IgG values relative to that of ZE3.

4.12 Plaque reduction neutralization assay

Serum samples from terminal blood collection were pooled for each mouse group and

heat inactivated. As described previously (Dent et al., 2016), the PRNT assay was carried out

using mouse sera diluted in Opti-Mem media (Invitrogen) at ratio of 1:10. Each sample was

incubated with 100 pfu ZIKV (PRVABC59, ATCC# VR-1843) in equal volume for 1hr at 37°C

before the virus/serum mixture was added to each well of VERO cells (ATCC # CCL-81) in a 24

well plate. The virus/serum mixture was aspirated after a 1.5 hour incubation at 37°C.

Subsequently, VERO cells were overlaid with 0.8% agarose in DMEM medium containing 5%

FBS (Invitrogen, CA) and incubated at 37°C for three days. Finally, VERO cells were fixed with

4% paraformaldehyde (PFA, MilliporeSigma, MA) overnight and stained with 0.2% crystal

violet. Plaques from each well were counted and neutralization % was calculated by using the

formula: [(number of ZIKV plaque per well in virus only control wells)-(number of ZIKV plaque

per well of diluted serum) / (number of ZIKV plaque per well in virus only control wells) x 100].

Neutralizing titers >10 refer to constructs with >50% neutralization at a 1:10 serum dilution.

Acknowledgements

This work was supported by funds provided by the School of Life Sciences, the Graduate and

Professional Student Association, and the Center for Immunotherapy, Vaccines, and Virotherapy

of the Biodesign Institute at Arizona State University. This work was also supported in part by a

grant from National Institute of Allergy and Infectious Diseases (NIAID) # R33AI101329 to QC.

Conflict of Interest

The authors have no conflicts of interest to declare.

Author Contributions

AD and HM designed experiments and analyzed data. AD, HM, and MP constructed vectors. AD

performed C1q binding, solubility, sucrose gradient, and expression experiments. AD, JH, and

MP performed purification experiments. AD and JH performed stability experiments. JK

performed mouse immunization and bleeds. AD and MP performed antibody titer experiments.

HS performed ZIKV neutralization experiments. AD wrote the manuscript. AD, HM, and QC

critically revised the MS.

References

Akkaya, M., Kwak, K., and Pierce, S. K. (2020). B cell memory: building two walls of

protection against pathogens. *Nat. Rev. Immunol.* 20, 229–238. doi:10.1038/s41577-019-

0244-2.

Alam, A., Jiang, L., Kittleson, G. A., Steadman, K. D., Nandi, S., Fuqua, J. L., et al. (2018).

Technoeconomic Modeling of Plant-Based Griffithsin Manufacturing. *Front. Bioeng.*

Biotechnol. 6, 102. doi:10.3389/fbioe.2018.00102.

Baker, K., Qiao, S. W., Kuo, T. T., Aveson, V. G., Platzer, B., Andersen, J. T., et al. (2011).

Neonatal Fc receptor for IgG (FcRn) regulates cross-presentation of IgG immune

complexes by CD8-CD11b+ dendritic cells. *Proc. Natl. Acad. Sci. U. S. A.* 108, 9927–9932.

doi:10.1073/pnas.1019037108.

Bournazos, S., and Ravetch, J. V. (2017). Fcγ Receptor Function and the Design of Vaccination

Strategies. *Immunity* 47, 224–233. doi:10.1016/j.immuni.2017.07.009.

Buyel, J. F. (2019). Plant molecular farming – Integration and exploitation of side streams to

achieve sustainable biomanufacturing. *Front. Plant Sci.* 9. doi:10.3389/fpls.2018.01893.

Carter, P. J. (2011). Introduction to current and future protein therapeutics: A protein engineering

perspective. *Exp. Cell Res.* 317, 1261–1269. doi:10.1016/j.yexcr.2011.02.013.

Castilho, A., Neumann, L., Daskalova, S., Mason, H. S., Steinkellner, H., Altmann, F., et al.

(2012). Engineering of sialylated mucin-type O-glycosylation in plants. *J. Biol. Chem.* 287,

36518–36526. doi:10.1074/jbc.M112.402685.

Castilho, A., and Steinkellner, H. (2012). Glyco-engineering in plants to produce human-like N-

glycan structures. *Biotechnol. J.* 7, 1088–1098. doi:10.1002/biot.201200032.

Chargelegue, D., Drake, P. M. W., Obregon, P., Prada, A., Fairweather, N., and Ma, J. K. C.

(2005). Highly immunogenic and protective recombinant vaccine candidate expressed in

transgenic plants. *Infect. Immun.* 73, 5915–5922. doi:10.1128/IAI.73.9.5915-5922.2005.

Chen, Q., and Davis, K. R. (2016). The potential of plants as a system for the development and

production of human biologics. *F1000Research* 5, 912. doi:10.12688/f1000research.8010.1.

Croix, D. A., Ahearn, J. M., Rosengard, A. M., Han, S., Kelsoe, G., Ma, M., et al. (1996).

Antibody response to a T-dependent antigen requires B cell expression of complement

receptors. *J. Exp. Med.* 183, 1857–1864. doi:10.1084/jem.183.4.1857.

Czajkowsky, D. M., Hu, J., Shao, Z., and Pleass, R. J. (2012). Fc-fusion proteins: New

developments and future perspectives. *EMBO Mol. Med.* 4, 1015–1028.

doi:10.1002/emmm.201201379.

Dent, M., Hurtado, J., Paul, A. M., Sun, H., Lai, H., Yang, M., et al. (2016). Plant-produced anti-

dengue virus monoclonal antibodies exhibit reduced antibody-dependent enhancement of

infection activity. *J. Gen. Virol.* 97, 3280–3290. doi:10.1099/jgv.0.000635.

Diamos, A. G., Hunter, J. G. L., Pardhe, M. D., Rosenthal, S. H., Sun, H., Foster, B. C., et al.

(2020a). High Level Production of Monoclonal Antibodies Using an Optimized Plant

Expression System. *Front. Bioeng. Biotechnol.* 7, 472. doi:10.3389/fbioe.2019.00472.

Diamos, A. G., Larios, D., Brown, L., Kilbourne, J., Kim, H. S., Saxena, D., et al. (2019).

Vaccine synergy with virus-like particle and immune complex platforms for delivery of

human papillomavirus L2 antigen. *Vaccine* 37, 137–144.

doi:10.1016/j.vaccine.2018.11.021.

Diamos, A. G., Pardhe, M. D., Sun, H., Hunter, J. G. L., Mor, T., Meador, L., et al. (2020b).

Codelivery of improved immune complex and virus-like particle vaccines containing Zika

virus envelope domain III synergistically enhances immunogenicity. *Vaccine* 38, 3455–

3463. doi:10.1016/j.vaccine.2020.02.089.

Diebolder, C. A., Beurskens, F. J., de Jong, R. N., Koning, R. I., Strumane, K., Lindorfer, M. A.,

et al. (2014). Complement Is Activated by IgG Hexamers Assembled at the Cell Surface.

Science (80-.). 343, 1260–1263. doi:10.1126/science.1248943.

Fletcher, E. A. K., van Maren, W., Cordfunke, R., Dinkelaar, J., Codee, J. D. C., van der Marel,

G., et al. (2018). Formation of Immune Complexes with a Tetanus-Derived B Cell Epitope

Boosts Human T Cell Responses to Covalently Linked Peptides in an Ex Vivo Blood Loop

System. *J. Immunol.* 201, 87–97. doi:10.4049/jimmunol.1700911.

Forthal, D. N., Gach, J. S., Landucci, G., Jez, J., Strasser, R., Kunert, R., et al. (2010). Fc-

Glycosylation Influences Fcγ Receptor Binding and Cell-Mediated Anti-HIV Activity of

Monoclonal Antibody 2G12. *J. Immunol.* 185, 6876–6882. doi:10.4049/jimmunol.1002600.

Getahun, A., and Heyman, B. (2006). How antibodies act as natural adjuvants. *Immunol. Lett.*

104, 38–45. doi:10.1016/j.imlet.2005.11.005.

Gleba, Y. Y., Tusé, D., and Giritch, A. (2014). Plant viral vectors for delivery by Agrobacterium.

Curr. Top. Microbiol. Immunol. 375, 155–92. doi:10.1007/82_2013_352.

Gupta, S. K., and Shukla, P. (2018). Glycosylation control technologies for recombinant

therapeutic proteins. *Appl. Microbiol. Biotechnol.* 102, 10457–10468. doi:10.1007/s00253-

018-9430-6.

He, J., Lai, H., Engle, M., Gorlatov, S., Gruber, C., Steinkellner, H., et al. (2014). Generation and

Analysis of Novel Plant-Derived Antibody-Based Therapeutic Molecules against West Nile

Virus. *PLoS One* 9, e93541. doi:10.1371/journal.pone.0093541.

Hioe, C. E., Kumar, R., Upadhyay, C., Jan, M., Fox, A., Itri, V., et al. (2018). Modulation of

Antibody Responses to the V1V2 and V3 Regions of HIV-1 Envelope by Immune Complex

Vaccines. *Front. Immunol.* 9, 2441. doi:10.3389/fimmu.2018.02441.

Ho, N. I., Camps, M. G. M., de Haas, E. F. E., Trouw, L. A., Verbeek, J. S., and Ossendorp, F.

(2017). C1q-Dependent Dendritic Cell Cross-Presentation of In Vivo-Formed Antigen-

Antibody Complexes. *J. Immunol.* 198, 4235–4243. doi:10.4049/jimmunol.1602169.

Huang, Z., and Mason, H. S. (2004). Conformational analysis of hepatitis B surface antigen

fusions in an Agrobacterium-mediated transient expression system. *Plant Biotechnol. J.* 2,

241–9. doi:10.1111/j.1467-7652.2004.00068.x.

Huber, V. C., McKeon, R. M., Brackin, M. N., Miller, L. A., Keating, R., Brown, S. A., et al.

(2006). Distinct contributions of vaccine-induced immunoglobulin G1 (IgG1) and IgG2a

antibodies to protective immunity against influenza. *Clin. Vaccine Immunol.* 13, 981–990.

doi:10.1128/CVI.00156-06.

Hurtado, J., Acharya, D., Lai, H., Sun, H., Kallolimath, S., Steinkellner, H., et al. (2019). In vitro

and in vivo efficacy of anti-chikungunya virus monoclonal antibodies produced in wild-type

and glycoengineered Nicotiana benthamiana plants. *Plant Biotechnol. J.*

doi:10.1111/pbi.13194.

Kallolimath, S., and Steinkellner, H. (2015). Glycosylation of plant produced human antibodies.

Hum. Antibodies 23, 45–48. doi:10.3233/HAB-150283.

Kim, M.-Y., Reljic, R., Kilbourne, J., Ceballos-Olvera, I., Yang, M.-S., Reyes-del Valle, J., et al.

(2015). Novel vaccination approach for dengue infection based on recombinant immune

679 complex universal platform. *Vaccine* 33, 1830–1838. doi:10.1016/j.vaccine.2015.02.036.

680 Kim, M. Y., Copland, A., Nayak, K., Chandele, A., Ahmed, M. S., Zhang, Q., et al. (2018).

681  Plant-expressed Fc-fusion protein tetravalent dengue vaccine with inherent adjuvant

682  properties. *Plant Biotechnol. J.* 16, 1283–1294. doi:10.1111/pbi.12869.

683 Konduru, K., Bradfute, S. B., Jacques, J., Manangeeswaran, M., Nakamura, S., Morshed, S., et

684  al. (2011). Ebola virus glycoprotein Fc fusion protein confers protection against lethal

685  challenge in vaccinated mice. *Vaccine* 29, 2968–2977. doi:10.1016/j.vaccine.2011.01.113.

686 Kumru, O. S., Joshi, S. B., Smith, D. E., Middaugh, C. R., Prusik, T., and Volkin, D. B. (2014).

687  Vaccine instability in the cold chain: Mechanisms, analysis and formulation strategies.

688  *Biologicals* 42, 237–259. doi:10.1016/j.biologicals.2014.05.007.

689 Lagassé, H. A. D., Hengel, H., Golding, B., and Sauna, Z. E. (2019). Fc-Fusion Drugs Have

690  FcγR/C1q Binding and Signaling Properties That May Affect Their Immunogenicity. *AAPS*

691  *J.* 21, 62. doi:10.1208/s12248-019-0336-8.

692 Loureiro, S., Ren, J., Phapugrangkul, P., Colaco, C. A., Bailey, C. R., Shelton, H., et al. (2011).

693  Adjuvant-Free Immunization with Hemagglutinin-Fc Fusion Proteins as an Approach to

694  Influenza Vaccines. *J. Virol.* 85, 3010–3014. doi:10.1128/jvi.01241-10.

695 Lu, L. L., Suscovich, T. J., Fortune, S. M., and Alter, G. (2018). Beyond binding: antibody

696  effector functions in infectious diseases. *Nat. Rev. Immunol.* 18, 46–61.

697  doi:10.1038/nri.2017.106.

698 Lyon, G. M., Mehta, A. K., Varkey, J. B., Brantly, K., Plyler, L., McElroy, A. K., et al. (2014).

699  Clinical care of two patients with Ebola virus disease in the United States. *N. Engl. J. Med.*

700  371, 2402–9. doi:10.1056/NEJMoa1409838.

Ma, J. K. C., Christou, P., Chikwamba, R., Haydon, H., Paul, M., Ferrer, M. P., et al. (2013).

Realising the value of plant molecular pharming to benefit the poor in developing countries

and emerging economies. *Plant Biotechnol. J.* 11, 1029–1033. doi:10.1111/pbi.12127.

Ma, J. K. C., Drossard, J., Lewis, D., Altmann, F., Boyle, J., Christou, P., et al. (2015).

Regulatory approval and a first-in-human phase I clinical trial of a monoclonal antibody

produced in transgenic tobacco plants. *Plant Biotechnol. J.* 13, 1106–1120.

doi:10.1111/pbi.12416.

Maamary, J., Wang, T. T., Tan, G. S., Palese, P., and Ravetch, J. V. (2017). Increasing the

breadth and potency of response to the seasonal influenza virus vaccine by immune

complex immunization. *Proc. Natl. Acad. Sci. U. S. A.* 114, 10172–10177.

doi:10.1073/pnas.1707950114.

Marusic, C., Pioli, C., Stelter, S., Novelli, F., Lonoce, C., Morrocchi, E., et al. (2018). N-glycan

engineering of a plant-produced anti-CD20-hIL-2 immunocytokine significantly enhances

its effector functions. *Biotechnol. Bioeng.* 115, 565–576. doi:10.1002/bit.26503.

Mason, H. S. (2016). Recombinant immune complexes as versatile and potent vaccines. *Hum.*

Vaccin. Immunother. 12, 988–9. doi:10.1080/21645515.2015.1116655.

Mastrangeli, R., Palinsky, W., and Bierau, H. (2019). Glycoengineered antibodies: towards the

next-generation of immunotherapeutics. *Glycobiology* 29, 199–210.

doi:10.1093/glycob/cwy092.

McCloskey, M. L., Curotto De Lafaille, M. A., Carroll, M. C., and Erlebacher, A. (2011).

Acquisition and presentation of follicular dendritic cell-bound antigen by lymph node-

resident dendritic cells. *J. Exp. Med.* 208, 135–148. doi:10.1084/jem.20100354.

Olinger, G. G., Pettitt, J., Kim, D., Working, C., Bohorov, O., Bratcher, B., et al. (2012).

Delayed treatment of Ebola virus infection with plant-derived monoclonal antibodies

provides protection in rhesus macaques. *Proc. Natl. Acad. Sci. U. S. A.* 109, 18030–5.

doi:10.1073/pnas.1213709109.

Pepponi, I., Diogo, G. R., Stylianou, E., van Dolleweerd, C. J., Drake, P. M. W., Paul, M. J., et

al. (2014). Plant-derived recombinant immune complexes as self-adjuvanting TB

immunogens for mucosal boosting of BCG. *Plant Biotechnol. J.* 12, 840–850.

doi:10.1111/pbi.12185.

Phan, T. G., Grigorova, I., Okada, T., and Cyster, J. G. (2007). Subcapsular encounter and

complement-dependent transport of immune complexes by lymph node B cells. *Nat.*

Immunol. 8, 992–1000. doi:10.1038/ni1494.

Phoolcharoen, W., Bhoo, S. H., Lai, H., Ma, J., Arntzen, C. J., Chen, Q., et al. (2011a).

Expression of an immunogenic Ebola immune complex in Nicotiana benthamiana. *Plant*

Biotechnol. J. 9, 807–16. doi:10.1111/j.1467-7652.2011.00593.x.

Phoolcharoen, W., Dye, J. M., Kilbourne, J., Piensook, K., Pratt, W. D., Arntzen, C. J., et al.

(2011b). A nonreplicating subunit vaccine protects mice against lethal Ebola virus

challenge. *Proc. Natl. Acad. Sci. U. S. A.* 108, 20695–700. doi:10.1073/pnas.1117715108.

Qiao, S. W., Kobayashi, K., Johansen, F. E., Sollid, L. M., Andersen, J. T., Milford, E., et al.

(2008). Dependence of antibody-mediated presentation of antigen on FcRn. *Proc. Natl.*

Acad. Sci. U. S. A. 105, 9337–9342. doi:10.1073/pnas.0801717105.

Qiu, X., Wong, G., Audet, J., Bello, A., Fernando, L., Alimonti, J. B., et al. (2014). Reversion of

advanced Ebola virus disease in nonhuman primates with ZMapp. *Nature* 514, 47–53.

doi:10.1038/nature13777.

Reed, S. G., Orr, M. T., and Fox, C. B. (2013). Key roles of adjuvants in modern vaccines. *Nat.*

Med. 19, 1597–1608. doi:10.1038/nm.3409.

Roopenian, D. C., and Akilesh, S. (2007). FcRn: the neonatal Fc receptor comes of age. *Nat.*

Rev. Immunol. 7, 715–725. doi:10.1038/nri2155.

Santi, L., Batchelor, L., Huang, Z., Hjelm, B., Kilbourne, J., Arntzen, C. J., et al. (2008). An

efficient plant viral expression system generating orally immunogenic Norwalk virus-like

particles. *Vaccine* 26, 1846–54. doi:10.1016/j.vaccine.2008.01.053.

Shukla, R., Shanmugam, R. K., Ramasamy, V., Arora, U., Batra, G., Acklin, J. A., et al. (2020).

Zika virus envelope nanoparticle antibodies protect mice without risk of disease

enhancement. *EBioMedicine* 54, 102738. doi:10.1016/j.ebiom.2020.102738.

Stelter, S., Paul, M. J., Teh, A. Y. H., Grandits, M., Altmann, F., Vanier, J., et al. (2020).

Engineering the interactions between a plant-produced HIV antibody and human Fc

receptors. *Plant Biotechnol. J.* 18, 402–414. doi:10.1111/pbi.13207.

Stettler, K., Beltramello, M., Espinosa, D. A., Graham, V., Cassotta, A., Bianchi, S., et al.

(2016). Specificity, cross-reactivity, and function of antibodies elicited by Zika virus

infection. *Science (80-.).* 353, 823–826. doi:10.1126/science.aaf8505.

Sun, H., Chen, Q., and Lai, H. (2018). Development of antibody therapeutics against

flaviviruses. *Int. J. Mol. Sci.* 19. doi:10.3390/ijms19010054.

Thielens, N. M., Tedesco, F., Bohlson, S. S., Gaboriaud, C., and Tenner, A. J. (2017). C1q: A

fresh look upon an old molecule. *Mol. Immunol.* 89, 73–83.

doi:10.1016/j.molimm.2017.05.025.

Tsouchnikas, G., Zlatkovic, J., Jarmer, J., Strauß, J., Vratskikh, O., Kundi, M., et al. (2015).

Immunization with Immune Complexes Modulates the Fine Specificity of Antibody

Responses to a Flavivirus Antigen. *J. Virol.* 89, 7970–8. doi:10.1128/JVI.00938-15.

van Montfoort, N., Mangsbo, S. M., Camps, M. G. M., van Maren, W. W. C., Verhaart, I. E. C.,

Waisman, A., et al. (2012). Circulating specific antibodies enhance systemic cross-priming

by delivery of complexed antigen to dendritic cells in vivo. *Eur. J. Immunol.* 42, 598–606.

doi:10.1002/eji.201141613.

Vidarsson, G., Dekkers, G., and Rispens, T. (2014). IgG subclasses and allotypes: From structure

to effector functions. *Front. Immunol.* 5. doi:10.3389/fimmu.2014.00520.

Wang, G., de Jong, R. N., van den Bremer, E. T. J., Beurskens, F. J., Labrijn, A. F., Ugurlar, D.,

et al. (2016). Molecular Basis of Assembly and Activation of Complement Component C1

in Complex with Immunoglobulin G1 and Antigen. *Mol. Cell* 63, 135–145.

doi:10.1016/J.MOLCEL.2016.05.016.

Wang, Q., Chung, C.-Y., Chough, S., and Betenbaugh, M. J. (2018). Antibody glycoengineering

strategies in mammalian cells. *Biotechnol. Bioeng.* 115, 1378–1393. doi:10.1002/bit.26567.

Webster, G. R., van Dolleweerd, C., Guerra, T., Stelter, S., Hofmann, S., Kim, M. Y., et al.

(2018). A polymeric immunoglobulin—antigen fusion protein strategy for enhancing

vaccine immunogenicity. *Plant Biotechnol. J.* 16, 1983–1996. doi:10.1111/pbi.12932.

West, E. E., Kolev, M., and Kemper, C. (2018). Complement and the Regulation of T Cell

Responses. *Annu. Rev. Immunol.* 36, 309–338. doi:10.1146/annurev-immunol-042617-

053245.

Wilson, J. A., Hevey, M., Bakken, R., Guest, S., Bray, M., Schmaljohn, A. L., et al. (2000).

Epitopes involved in antibody-mediated protection from Ebola virus. *Science (80-.).* 287,

1664–1666. doi:10.1126/science.287.5458.1664.

Yang, M., Dent, M., Lai, H., Sun, H., and Chen, Q. (2017). Immunization of Zika virus envelope

protein domain III induces specific and neutralizing immune responses against Zika virus.

Vaccine 35, 4287–4294. doi:10.1016/j.vaccine.2017.04.052.

Zeitlin, L., Pettitt, J., Scully, C., Bohorova, N., Kim, D., Pauly, M., et al. (2011). Enhanced

potency of a fucose-free monoclonal antibody being developed as an Ebola virus

immunoprotectant. *Proc. Natl. Acad. Sci.* 108, 20690–20694.

doi:10.1073/pnas.1108360108.

Zeng, F., Yang, C., Gao, X., Li, X., Zhang, Z., and Gong, R. (2018). Comprehensive elucidation

of the structural and functional roles of engineered disulfide bonds in antibody Fc fragment.

J. Biol. Chem. 293, 19127–19135. doi:10.1074/jbc.RA118.005367.

Zhang, Y., Zhou, Z., Zhu, S. L., Zu, X., Wang, Z., Zhang, L. ke, et al. (2019). A novel RSV F-Fc

fusion protein vaccine reduces lung injury induced by respiratory syncytial virus infection.

Antiviral Res. 165, 11–22. doi:10.1016/j.antiviral.2019.02.017.

Zhao, B., Zhang, X., Krummenacher, C., Song, S., Gao, L., Zhang, H., et al. (2018).

Immunization with Fc-based recombinant Epstein-Barr virus gp350 elicits potent

neutralizing humoral immune response in a BALB/c mice model. *Front. Immunol.* 9, 932.

doi:10.3389/fimmu.2018.00932.

Supporting Info

Figure S1. C1q binding comparison between wildtype and glycoengineered plants

Figure S2. C1q binding comparison between IgG fusions

Figure S3. Epitope binding of single chain 6D8

Figure S4. RIC insolubility

Figure S5. Solubility and binding of 6D8 epitope tag mutants

Tables

Construct	Yield	Solubility	Sedimentation	Stability	C1q	IgG	IgG2a	Neutralization
ZE3	N.D.	N.D.	N.D.	N.D.	N.D	+	+	++
HLZe	0.22 mg/g	Low	+++++	High	+++	+++	++	+++
HLZd	1.50 mg/g	High	+++	High	+++	+++	+++	+++
HLZ	0.17 mg/g	High	++	Medium	+	+++	++	+++
ZH	0.83 mg/g	High	+	Medium	++	+++	+++	+++
ZHx	0.21 mg/g	Low	+++*	High	+++	+++	++++	+++
VHLZe	<0.1 mg/g	Low	+++++	High	+++	+++	+++	+++
ZFc	0.57 mg/g	High	+	Low	+++	+++	++++	+++

Table 1. Summary of characteristics of IgG fusions

A summary of the characteristics of each IgG fusion vaccine candidate is given. For expression,

only the yield (mg/g LFW) of the fully assembled product is shown. A greater number of "+"

symbols indicates either a statistically significant increase in the mean value for that property

(for C1q, IgG, IgG2a, and neutralization), or a repeatably observed difference (for

sedimentation). For ZHx (*), peaks of both low- and high-velocity sedimentation were observed.

Figure Legends

Figure 1. **Various IgG fusion strategies enhance C1q binding**

(A) Schematic representation of IgG fusion constructs used in this study. Fusion constructs are

built on the mAb 6D8 human IgG1 backbone with the shown modifications. ZE3; the Zika

envelope domain III containing amino acids K301 to T406; e, an epitope tag containing the

"VYKLDISEA" 6D8 binding motif for RIC formation; e with lightning bolt, an epitope tag

truncated to include only "YKLDIS" to reduce RIC formation; VH, the variable heavy domain

from 6D8 which participates in binding the epitope tag; VL, the variably light domain from 6D8

which participates in binding the epitope tag; H, the heavy chain constant CH1 domain from

6D8; L, the light chain constant domain from 6D8; Fc, the heavy chain constant CH2 or CH3

domains from 6D8; Fc with lightning bolt, same as Fc but with E345R, E430G, and S440Y

mutations to induce hexamer formation. (B) C1q binding ELISA of purified IgG fusion

constructs made in plants that lack plant-specific β1,2-linked xylose and α1,3-linked fucose N-

linked glycans. ELISA plates were coated with 10 µg/ml human C1q and incubated with 10

µg/ml each molecule, using 6D8 with no fusion as a negative control. Constructs were detected

using polyclonal goat anti-human IgG-HRP. Mean OD_{450} values from three samples are shown ±

standard error with one star (*) indicating $p < 0.05$ and three stars (***) indicating $p < 0.001$ as

measured by one-way ANOVA comparing each column to the control column HL, unless

otherwise indicated with a bracket.

Figure 2. **Modified IgG fusions have improved expression and solubility**

(A) ELISA and gel quantification of IgG fusion construct expression. Clarified protein extracts

("soluble" fraction) from leaf spots agroinfiltrated with each IgG fusion construct were analyzed

by either ELISA, or SDS-PAGE followed by gel image quantification. For ZE3 ELISA, plates

were coated with polyclonal mouse anti-ZE3, incubated with serial dilutions of extracts from

each IgG fusion using purified HLZ as a standard, and probed with goat anti-human IgG-HRP.

For IgG ELISA, plates were coated with serial dilutions of extracts or human IgG standard and

probed with goat anti-human IgG-HRP. For gel quantification, ImageJ software was used to

compare the IgG fusion band intensity visualized on stain-free polyacrylamide gels using

purified 6D8 antibody as standard. Columns represent means ± standard error from three

independently infiltrated leaf samples. (B) Clarified leaf extracts were separated by reducing

SDS-PAGE and a representative gel image is shown. The general band position corresponding to

each respective heavy chain/ZE3 fusion is indicated "ZH/HZ." The small shift in size in HLZe

(~67 kDa of the reduced heavy chain fusion before glycosylation) and HLZd (~65 kDa) or HLZ

(~64.5 kDa) is due to epitope tag truncation. ZH and ZHx have a predicted size of ~64.5 kDa.

The "<" indicates ZE3-Fc (~40 kDa) and Fc (~29 kDa) fragments. The large subunit of Rubisco

is abbreviated "RbcL."

Figure 3. Purification of IgG Fusions

Agroinfiltrated leaf material from between 1-3 plants per construct was homogenized, clarified,

and purified by protein G affinity chromatography. The peak elutions were pooled and separated

on nonreducing (NR) and reducing (R) SDS-PAGE using stain-free polyacrylamide gels.

Representative lanes for each construct are compiled from multiple gels here.

Figure 4. Sucrose gradient centrifugation of IgG fusions

Purified IgG fusions were separated by sucrose gradient sedimentation using 5/10/15/20/25%

discontinuous sucrose layers. Gradient fractions were analyzed by SDS-PAGE and

representative results are shown; direction of sedimentation is left to right. The relative band

intensity was quantified using ImageJ software and the peak band was arbitrarily assigned the

value of 1.

Figure 5. Stability and C1q binding of purified IgG fusions

Samples of purified IgG fusions were frozen and thawed once after purification (initial), or

additionally subjected to either additional 5 freeze/thaw cycles, incubation for 2 weeks at 4°C, or

incubation for 2 weeks at 23˚C. (A) After each treatment, samples were separated on reducing

and nonreducing SDS-PAGE gels, and the relative proportion of fully assembled product was

analyzed using ImageJ software. The initial measurement reflects degradation which occurred

during expression or purification, whereas the other measurements reflect degradation which

occurred during each respective treatment. (B) Samples from each treatment were analyzed by

C1q binding ELISA. Columns represent the mean OD_{450} value ± standard error from three

samples.

Figure 6. Mouse immunization and serum titers

BALB/c mice (6 per group) were immunized twice two weeks apart subcutaneously with a dose

that would deliver 8 µg ZE3 for each IgG fusion or with PBS as a control. Mouse serum samples

were collected two weeks after the final dose. (A) Serially diluted mouse serum was analyzed for

total IgG production by ELISA. The endpoint was taken as the reciprocal of the greatest dilution

that gave an OD_{450} reading at least twice the background. Two stars (**) indicates $p < 0.01$ by

one-way ANOVA comparing the indicated columns to ZE3. (B) Mouse serum samples were

diluted 1:100 and analyzed for IgG2a production by ELISA. The table shows the ratio of IgG2a

to total IgG for each group, calculated as relative to ZE3, with the ZE3 value set arbitrarily at

1.0. Values with one star (*) indicates $p < 0.05$ and (**) indicates $p < 0.01$ by one-way ANOVA

comparing the indicated columns to HLZ.

Figure 7. Neutralization of ZIKV

Plaque reduction neutralization assays (PRNT) assays were performed using pooled mouse sera

with dilution ratio of 1:10 to evaluate ZIKV-specific neutralizing antibodies in the sera. Data are

presented as mean neutralization % and SD from three independent experiments with technical

triplicates for each sample. Statistical analyses were performed using one-way ANOVA, p values

from comparison between vaccine treatments and PBS were indicated with **** (<0.0001) or

from comparison between vaccine treatments and His-tagged ZE3 were indicated with * (p <

0.024).

A

HLZ HLZe HLZd ZH

ZHx HVLZe ZFc

B

C1q binding

A

Total IgG

B

Total IgG2a

	IgG2a:IgG
ZE3	1.00
HLZe	1.64
HLZd	1.77
HLZ	1.13
ZH	1.68
ZHx	1.64
HVLZe	1.79
ZFc	0.92