Optogenetic control of small GTPases reveals RhoA-mediated intracellular calcium

signaling

Hironori Inaba[1,2], Qianqian Miao[1,2], Takao Nakata[1,2,3]

[1]Department of Cell Biology, Tokyo Medical and Dental University (TMDU), Bunkyo-

ku, Tokyo, 113-8510, Japan.

[2]The Center for Brain Integration Research (CBIR), Tokyo Medical and Dental

University (TMDU), Bunkyo-ku, Tokyo, 113-8510, Japan.

[3]Correspondence should be addressed to T.N. (E-mail: info.cbio@tmd.ac.jp)

Takao Nakata, M.D., Ph.D.

Department of Cell Biology, and The Center for Brain Integration Research (CBIR),

Tokyo, Medical and Dental University (TMDU), 1-5-45 Yushima, Bunkyo-ku, Tokyo

113-8510, Japan.

TEL: +81-3-5803-5143

Running title: Optogenetic RhoA activation induces Ca^{2+} signaling

Abstract

Rho/Ras family small GTPases are known to regulate numerous cellular processes,

including cytoskeletal reorganization, cell proliferation, and cell differentiation. These

processes are also controlled by Ca^{2+}, and consequently, crosstalk between these signals

is considered likely. However, systematic quantitative evaluation is not yet reported. Thus,

we developed optogenetic tools to control the activity of small GTPases (RhoA, Rac1,

Cdc42, Ras, Rap, and Ral) using an improved light-inducible dimer system (iLID). Using

these optogenetic tools, we investigated calcium mobilization immediately after small

GTPase activation. Unexpectedly, we found that only RhoA activation induced a transient

intracellular calcium elevation in RPE1 and HeLa cells. Transients were also observed in

MDCK and HEK293T cells by RhoA activation, but interestingly, molecular mechanisms

were identified to be different among cell types. In RPE1 and HeLa cells, RhoA directly

activated phospholipase C (PLC)ε at the plasma membrane, which in turn induced Ca^{2+}

release from the endoplasmic reticulum (ER). The RhoA-PLCε axis induced calcium-

dependent NFAT nuclear translocation, suggesting it does activate intracellular calcium

signaling.

1 Introduction

Small GTPases of the Ras superfamily have been identified as molecular switches

because they exist in two states, a GTP-binding state ("ON") and a GDP-binding state

("OFF") (Mitin et al., 2005; Wennerberg et al., 2005). These states are known to be

regulated by activators – guanine nucleotide exchange factors (GEFs) – and inactivators,

that is, GTPase-activating proteins (GAPs). Rho and Ras subfamily small GTPases

localize at the plasma membrane (PM), respond to extracellular stimuli, and are

responsible for a variety of biological processes, including cytoskeletal reorganization,

cell proliferation, and cell differentiation (Mitin et al., 2005; Wennerberg et al., 2005).

Several of these biological processes are also regulated by a universal second messenger,

that is, calcium ion (Ca^{2+}) (Berridge et al., 2000; Clapham, 2007; Cullen and Lockyer,

2002). Thus, functional links must exist between these signaling pathways regulated by

Rho/Ras family small GTPases and Ca^{2+}. Small GTPases and Ca^{2+} share some

downstream factors that are coordinately regulated. For example, RhoA and Ca^{2+} regulate

myosin II activity via myosin light chain phosphorylation (Amano et al., 1996; Ikebe and

Hartshorne, 1985), and Ras and Ca^{2+} coordinate extracellular signal-regulated kinase

(ERK)/mitogen-activated kinase (MAPK) signaling pathway (Chao et al., 1992; Kolch,

2000). Besides, small GTPases and Ca^{2+} are known to regulate each other's functions:

many GEFs and GAPs are regulated both positively and negatively by Ca^{2+} (Aspenström,

2004; Cullen and Lockyer, 2002), and some small GTPases regulate intracellular calcium

signaling via activating phospholipase C (PLC) (Harden et al., 2009; Smrcka et al., 2012).

PLC converts phosphatidylinositol 4,5-bisphosphate $[PI(4,5)P_2]$ to two second

messengers: diacylglycerol (DAG) and inositol-trisphosphate (IP_3). IP_3 reportedly binds

to the IP_3 receptor (IP3R) to release Ca^{2+} from the endoplasmic reticulum (ER). This

PLC-mediated calcium influx is the major calcium signaling pathway in non-excitable

cells.

9   Despite the importance of crosstalk between small GTPases and intracellular

10   calcium, details of the processes remain to be poorly understood. Especially, assessment

11   of small GTPases' influences on intracellular calcium concentrations immediately after

12   activation has been difficult because such activity could not be directly controlled in cells.

13   However, optogenetics has changed this situation over the last decade.

14   Optogenetics is a pivotal tool for advancing cell biology because it enables

15   control of specific signaling molecules at high spatiotemporal resolution both *in vitro* and

16   *in vivo* (Guglielmi et al., 2016; Tischer and Weiner, 2014; Toettcher et al., 2011).

17   Optogenetic control of small GTPases was first reported by Hahn's group (Wu et al.,

18   2009). In the study, constitutively active mutants of Rac1 and Cdc42 were fused to the

blue-light-excited light-oxygen-voltage-sensing domain 2 (LOV2) of phototropin from

Avena sativa (Christie et al., 1999). Photoactivatable (PA)–Rac1 and PA–Cdc42 were

inactive in the dark due to steric hindrance of effector binding sites by the LOV2 domain.

Blue-light irradiation made conformation changes to alpha-helix (Jα) that connects LOV2

domains to small GTPases, allowing them to bind effectors. However, this approach was

difficult to optimize between "ON" and "OFF" states for other small GTPases. Therefore,

PM translocation of their specific GEFs with light-induced heterodimeric systems, such

as CRY2-CIBN (Kennedy et al., 2010), iLID (Guntas et al., 2015), TULIP (Strickland et

al., 2012) and PhyB-PIF (Levskaya et al., 2009) systems, has been broadly used for

regulating small GTPase activity including Rac1 (de Beco et al., 2018; Guntas et al.,

2015; Levskaya et al., 2009), Cdc42 (de Beco et al., 2018; Guntas et al., 2015; Levskaya

et al., 2009), RhoA (O'Neill et al., 2018; Valon et al., 2017; Wagner and Glotzer, 2016),

Ras (Toettcher et al., 2013), and Ral (Zago et al., 2018).

    We have developed optogenetic tools to control the activity of six members of

the Rho and Ras subfamily GTPases (RhoA, Rac1, Cdc42, Ras, Rap, and Ral) by light-

inducing GEF translocation to the PM using the iLID system. Using these optogenetic

tools, we were able to examine small GTPase-mediated intracellular calcium mobilization

for the first time. Unexpectedly, transient elevation of intracellular calcium concentrations

1   was only induced by optogenetic RhoA activation. These RhoA-mediated calcium

2   transients were observed in all cell types examined, but the molecular mechanisms were

3   different among the cell types. Furthermore, we found that RhoA directly activated PLCε

4   in RPE1 and HeLa cells, which induced intracellular calcium signaling.

Results

Development of optogenetic tools for controlling small GTPase activity

Specific control of Rho/Ras family small GTPase activity at high spatiotemporal

resolution was achieved using optogenetic tools. Among the several light-inducible

heterodimerization systems, we chose the iLID system because of the reasons that follow:

(1) it is based on the *As*LOV2 domain that can work without exogenously adding a

chromophore to mammalian cells; (2) iLID-SspB heterodimerization can be controlled

by blue-light with rapid on/off kinetics (seconds), which is suitable in controlling small

GTPase activity at high spatiotemporal resolution; and (3) the molecular weight of

proteins is small, allowing high-level expression in cells and relative ease of establishing

a lentivirus vector for stable cell lines.

To evaluate the system, we developed an optogenetic tool to control RhoA

activity (opto-RhoA; **Figure 1A**). Similar to previous studies (O'Neill et al., 2018; Valon

et al., 2017; Wagner and Glotzer, 2016), a DH domain of the RhoA-specific GEF LARG

(LARG-DH) was fused to SspB and fluorescence protein mVenus. The CAAX motif of

K-Ras (CAAX^{Kras}) was fused to the C-terminus of iLID to localize on the PM. Further,

the nuclear export signal (NES) was inserted between mVenus and SspB to avoid

localizing to the nucleus, as originally reported (Guntas et al., 2015). Cells were

transfected with a single plasmid, mVenus-SspB-LARG-DH and iLID-CAAX[KRas] linked

with the cDNA of the self-cleaved P2A peptide. In the dark, mVenus-SspB-LARG-DH

was diffusely localized in the cytoplasm (**Figure 1B, 458 nm light OFF**). Blue-light

irradiation with a 458 nm laser has induced iLID-SspB heterodimerization, which in turn

translocated LARG-DH to the PM where it could activate RhoA (**Figure 1B, 458 nm**

light ON). RhoA activation by opto-RhoA was tested using a RhoA activity reporter:

mCherry fused Rho-binding domain derived from the RhoA-specific effector Rhotekin

(mCherry-RBD[Rhotekin]) (Reid et al., 1996). Opto-RhoA and mCherry-RBD[Rhotekin] were

transfected into HeLa cells, and cells were locally irradiated with a 458 nm laser (**Figure**

1C and Movie 1). During irradiation, mVenus-SspB-LARG-DH rapidly accumulated in

the irradiated area, while mCherry-RBD[Rhotekin] did so, however, in a gradual manner

(**Figure 1D**). Accumulation disappeared after irradiation was terminated. These data

suggest that RhoA activity can be controlled spatiotemporally using the opto-RhoA as

previously reported (O'Neill et al., 2018; Valon et al., 2017; Wagner and Glotzer, 2016).

We next developed optogenetic tools for another five members of Rho and Ras

family GTPases (Rac1, Cdc42, Ras, Rap, and Ral) implicated in regulating intracellular

calcium signaling (Harden et al., 2009; Smrcka et al., 2012) (**Supplementary Figure 1**).

The LARG-DH of opto-RhoA was replaced with GEF domains of specific GEFs for each

small GTPase (see details in the Materials and Methods section); most were already

reported for application in optogenetic tools (Guntas et al., 2015; Toettcher et al., 2013;

Zago et al., 2018). We termed these optogenetic tools as opto-X (X: target GTPases).

Screening using opto-GTPases for induction of changes in intracellular calcium concentrations

We monitored changes of intracellular calcium concentrations with a genetically encoded

red fluorescent calcium indicator R-GECO1 that is known to detect physiological calcium

changes as found during neural activity and muscle contraction (Zhao et al., 2011). Opto-

GTPases and the calcium reporter R-GECO1 were transfected into human non-

transformed (RPE1) and cancer (HeLa) cells. Changes of intracellular calcium

concentrations induced by persistent activation of small GTPases were observed via

confocal microscopy. Only RhoA was identified to induce transient elevation of

intracellular calcium concentrations both in RPE1 and HeLa cells **(Figure 2,**

Supplementary Figure 2, and Movie 2). We also tested a YA mutant of opto-RhoA

(opto-RhoA[YA]) that possessed a LARG Y940A mutation that abolished GEF activity

(Martz et al., 2013). Opto-RhoA[YA] rarely induced calcium transients (1 out of 136 RPE1

cells and 0 out of 133 HeLa cells; **Figure 2C**), confirming that calcium transients were

1 induced by RhoA activation and not by blue-light irradiation or by photoswitch

2 translocation to the PM.

3  Further, we examined RhoA-mediated calcium transients in MDCK and

4 HEK293T cells. These cells also exhibited the calcium transients immediately after RhoA

5 activation (30 % and 41 % of cells in MDCK (n = 50) and HEK293T (n = 52) cells,

6 respectively; **Supplementary Figure 2 and Movie 2**). Thus, all cells examined exhibited

7 RhoA-mediated calcium transients. Of note, membrane blebbing, produced by

8 actomyosin-mediated cellular contraction induced by RhoA activation (Charras et al.,

9 2005), was seen in opto-RhoA-transfected HEK293T cells even before blue-light

10 irradiation (**Supplementary Figure 2 and Movie 2**). This observation suggests that opto-

11 RhoA was somewhat leaky, that is, opto-RhoA exerts background activity in the dark.

12 The number of blebs was dramatically increased after the blue-light irradiation, indicating

13 that the blue-light irradiation did increase RhoA activity.

14  Time-course analysis showed that initiation of an increase in intracellular

15 calcium concentration was observed within 10 seconds in RPE1, HeLa, and HEK293T

16 cells; meanwhile, maximum concentration was observed about 20 seconds after light

17 irradiation (**Figure 2D**). In contrast, initiation of this increase was observed within 20–

18 30 seconds, and the maximum increase was reached about 2 minutes after light irradiation

in MDCK cells (**Figure 2D**). Cellular responses differed among cell types.

Mechanisms of RhoA-induced calcium transients are different among cell types

We investigated molecular mechanisms of RhoA-induced intracellular calcium transients

using small molecule inhibitors (see **Figure 7** for summary). Several calcium channels

are activated by RhoA and its downstream factors (Li and Brayden, 2017; Mehta et al.,

2003; Wing et al., 2003). We initially examined the RhoA-ROCK-myosin II axis, the

major pathway of the RhoA signaling pathway. Both ROCK inhibitor, Y-276322, and

myosin II inhibitor, Blebbistatin, were observed to efficiently inhibit RhoA-induced

calcium transients in MDCK and HEK293T cells, but surprisingly not in RPE1 and HeLa

cells (**Figure 3**). Actomyosin-mediated cellular contraction activates mechanosensitive

(MS) calcium channels such as Piezo1 and transient receptor potential (TRP) family

channels in MDCK and HEK293T cells.

Further, we tested the non-selective calcium channel blockers SKF96365 and 2-

APB (**Figure 3**). SKF96365 was observed to inhibit the RhoA-induced calcium transients

in MDCK and HEK293T cells (**Figures 3C, D**). Conversely, 2-APB blocked transients

in RPE1 and HeLa cells (**Figures 3A, B**). Mechanisms of RhoA-induced calcium

transients are clearly different among cell types. Our further studies focused on molecular

mechanisms of RhoA-induced calcium transients in RPE1 and HeLa cells.

RhoA-PLCε axis induces calcium transients in RPE1 and HeLa cells

2-APB inhibits IP3R, and RhoA directly activates a PLC isoform PLCε (Wing et al.,

2003). Thus, we hypothesized that the RhoA-PLCε axis has induced calcium transients

in RPE1 and HeLa cells; we also further examined the effect of the PLC inhibitor U73122

(**Figures 3A, B**). Initially, calcium transients were not affected by U73122 in L-15

medium. However, U73122 is reported to spontaneously form conjugates with chemical

components of cell culture medium, such as L-glutamine, glutathione, and bovine serum

albumin (BSA). This conjugation causes loss of inhibitory activity toward PLC (Wilsher

et al., 2007). Calcium transients were abolished by U71322 both in RPE1 and HeLa cells

in this condition when incubated in Ringer's solution. As hypothesized, RhoA activates

PLC, which in turn induces calcium transients.

We also tested RhoA-induced calcium transients in calcium-free buffers

(**Figures 3A, B**). The percentage of cells exhibiting calcium transients significantly

decreased in both cell types. However, about 10 % of cells still showed transients, and

intracellular calcium stores have been identified as a source of these calcium transients.

Extracellular calcium could be required for proper calcium homeostasis.

We also knocked down PLCε in RPE1 and HeLa cells with small interference

RNA (siRNA) and examined calcium transients (**Figures 4A, B**). Blue-light irradiation

of PLCε-depleted cells did not induce calcium transients. Further, ECFP-fused mouse

PLCε (msPLCε), not ECFP alone, rescued calcium transient phenotypes in PLCε-

depleted cells (**Figures 4C–E, Supplementary Figure 3**). PLCε is thus essential for

calcium transients induced by RhoA in RPE1 and HeLa cells.

We performed additional rescue experiments with several PLCε mutants in

PLCε-depleted HeLa cells to further confirm the RhoA-PLCε functions. PLCε displays a

lipase domain that includes a PH domain, four EF-hand domains, a catalytic core X-Y

domain, and a C2 domain and also a CDC25 homology domain which activates Rap1

(Citro et al., 2007; Jin et al., 2001; Lopez et al., 2001) (**Figure 4C**). This information led

us to examine the function of PLCε essential for RhoA-induced calcium transients. The

GEF-inactive mutant, F702A PLCε, which corresponds to the F929A GEF-inactive

mutant of a RasGEF SOS (Hall et al., 2001), has rescued calcium phenotype in PLCε-

depleted cells. Conversely, the lipase-inactive mutant, H1433L (Lopez et al., 2001), did

not rescue the phenotype (**Figures 4D, E**). Lipase activity appears essential for calcium

transients.

PLCε is also regulated by Ras (Kelley et al., 2001; Song et al., 2001), Rap1 (Jin

et al., 2001), Ral (Kelley et al., 2004), and heterotrimeric G-protein β and γ subunits ($G_{\beta\gamma}$)

(Wing et al., 2001). A unique insert in the Y domain (Yins in **Figure 4C**) of PLCε is

required for its activation by RhoA (Wing et al., 2003). In contrast, Ras and Rap1 are seen

to bind directly to the second Ras association domain (RA2 domain in **Figure 4C**)

(Bunney et al., 2006) of PLCε. Then, we performed rescue experiments in PLCε-depleted

HeLa cells with a Yins-deleted PLCε mutant (ΔYins) and the R2130L PLCε mutant which

abolishes interactions with Ras (Bunney et al., 2006) (**Figure 4D**). As a result, the

R2130L mutant rescued the calcium phenotype in PLCε-depleted cells, but the ΔYins

mutant did not (**Figure 4E**). RhoA directly activates the phospholipase activity of PLCε,

which in turn induces intracellular calcium transients.

RhoA activation at the PM but not in the Golgi is essential for the calcium transients

PLCε was recently reported to be activated at the Golgi, and not at the PM in cardiac cells,

to hydrolyze phosphatidylinositol 4-phosphate [PI(4)P] (Malik et al., 2015; Nash et al.,

2019; Zhang et al., 2013). Our optogenetic system, iLID-CAAX^{KRas}, largely localizes to

the PM but is also partially localized in the Golgi. Thus, opto-RhoA could activate Golgi

RhoA. However, PI(4)P hydrolysis produces inositol bisphosphate (IP_2) but not IP_3, and

PLCε activation at the Golgi is not likely to induce IP3R-dependent calcium release from

the ER. We examined this issue by constructing opto-RhoA-TGN (*trans*-Golgi network)

using an N-terminus transmembrane domain of 2,6-sialyltransferase (Shiwarski et al.,

2017), and opto-RhoA-Golgi with KDELr D193N dominant-negative mutant (KDELr

DN) (Irannejad and Wedegaertner, 2010), instead of the $CAAX^{KRas}$ motif (**Figures 5A,**

B). Both in RPE1 and HeLa cells, only about 10 % of cells expressing opto-RhoA-TGN

exhibited calcium transients, and cells expressing opto-RhoA-Golgi did not (**Figure 5C**).

RhoA activation at the PM appears essential for calcium transients.

We next explored $PI(4,5)P_2$ dynamics on the PM using the $PI(4,5)P_2$ marker, a

PH domain of PLCδ (PLCδ-PH) (Stauffer et al., 1998) and an optogenetic tool for

controlling the inositol 5-phosphatase (opto-5-ptase) with the inositol 5-phosphatase

domain of OCRL ($OCRL_{cat}$) and the CRY2-CIBN heterodimerization system, as

previously reported (Idevall-Hagren et al., 2012) (**Figure 5D**). This latter construct served

as a positive control. CRY2-CIBN heterodimerization is induced by blue-light irradiation.

Cytosolic ECFP-CRY2-$OCRL_{cat}$ then translocates to the PM in functional form. When 5-

phosphatase was recruited to the PM, the fluorescence intensity of cytosolic mCherry-

PLCδ-PH increased to about 40 % within 30 seconds (**Figures 5E, F and Movie 3)**,

indicating that membrane $PI(4,5)P_2$ decreased. In contrast, when RhoA was activated by

opto-RhoA, the intensity of cytosolic mCherry-PLCδ-PH increased only about 10 % even

after 5 minutes. This result is comparable to intensity changes of mCherry alone (**Figures**

5E, F and Movie 4). RhoA-mediated cellular contraction might also cause this increase

of mCherry intensity since we quantified mCherry intensity in a constant square area. The

RhoA-PLCε axis functions at the PM, but only marginally affects membrane

phosphoinositide dynamics.

RhoA-PLCε axis induces calcium signaling

Finally, we examined RhoA-PLCε-mediated intracellular calcium transients for

activation of intercellular calcium signaling with an NFAT nuclear translocation assay.

NFAT is identified as a transcription factor, and, after dephosphorylation by calmodulin-

calcineurin, it translocates from the cytoplasm to the nucleus (Hogan et al., 2003). About

30 % of cells exhibited mCherry-NFAT nuclear translocation after RhoA activation with

opto-RhoA (**Figures 6A, B, and Movie 5**). This response is comparable to the percentage

of cells that exhibited calcium transients. Translocation was abolished by treatment with

PLCε-specific siRNA (**Figures 6C, D**), indicating that the RhoA-PLCε axis induces

intracellular calcium signaling.

Discussion

In this study, we have developed optogenetic tools to control Rho and Ras family GTPase activity; we then performed cell-based systematic functional screening (**Figure 2**). We focused on intracellular calcium signaling induced by Rho and Ras signaling. This approach for cell-based screening is applicable for other signaling pathways as well if both optogenetic tools and biosensors are available. Before the introduction of optogenetic tools, single cell-based enzymology was difficult due to the lack of sufficient spatiotemporal resolution and specificity. Optogenetics has resolved this issue by exerting control over specific signaling molecules using a light stimulus with high spatiotemporal resolution. This tool provides a powerful platform for cell-based enzymology.

Surprisingly, only RhoA activation was seen to induce calcium transients in screening using opto-GTPases (**Figure 2**); other GTPases are also reported to induce calcium signaling (Harden et al., 2009; Smrcka et al., 2012). The sensitivity of our experiments may not have been sufficient to detect calcium transients by such GTPase activation, or leaks in our optogenetic systems (**Supplementary Figure 2**) may have affected signaling responses. However, RhoA activation has induced calcium transients under the same conditions. Another possibility is that activation of only individual small GTPases does not induce calcium transients. Other signals, such as PI3K, could be

required for activating calcium signaling. Our study is the first to induce temporal

activation of specific GTPases and monitor subsequent changes of intracellular calcium.

This approach is expected to have much higher specificity and time resolution compared

with previous studies. Further, our systems can simultaneously stimulate other signals

using multiple optogenetic tools, and we can examine dual activation of small GTPases

and other signals for induction of calcium transients. Such experiments could explain

many discrepancies in signal transduction.

RhoA-induced calcium transients were observed in all cell lines examined in this

study (**Figure 2 and Supplementary Figure 2**) and have also been reported previously

in still other cell lines (Li and Brayden, 2017; Mehta et al., 2003; Wing et al., 2003). RhoA

appears to generally induce calcium signaling. However, only up to about 50 % of cells

expressing opto-RhoA show calcium transients. This observation could be explained in

part by transfection-related issues, such as damage to cells. There were cells with varying

levels of the expression of opto-RhoA and R-GECO1. Most of the cells with high

expression of opto-RhoA and/or R-GECO1 did not show calcium transients, suggesting

that the overproduction of exogenous proteins could be cytotoxic. Also, such high

expression could cause RhoA activation even in the dark, with concomitant background

effects. However, there were still cell populations that appeared healthy, did not highly

express exogenous proteins, but still did not exhibit calcium transients.

RhoA-induced calcium transients might also be dependent on cellular contexts,

such as cell-cycle stage; expression and posttranslational modifications of downstream

molecules, including RhoA and PLCε; and intracellular calcium concentrations. Most

importantly, almost all cells initially exhibited calcium transients also responded

repeatedly to blue-light irradiation after a few minutes of rest (data not shown), ensuring

reproducibility of RhoA-mediated calcium transients.

Unexpectedly, molecular mechanisms underlying calcium transients were found

to be notably different among cell types (**Figures 3, 7**). In MDCK and HEK293 cells, the

RhoA-ROCK-Myosin II axis induced calcium transients (**Figures 3C, D**). Actomyosin-

mediated contractile force is also reported to induce membrane tension changes that

activate MS channels (Jin et al., 2020) (**Figure 7, lower scheme**). One candidate is the

Piezo channel that is reported in both MDCK (Gudipaty et al., 2017) and HEK293T cells

(Cox et al., 2016). However, Piezo channel is not sensitive to SKF96365 (Weng et al.,

2018), which inhibited transient induction (**Figures 3C, D**). Thus, other MS calcium

channels, such as a member of the TRP family of channels, may be needed for RhoA-

induced calcium transients. Further studies are also required to elucidate molecular

mechanisms for the RhoA-ROCK-Myosin II-mediated calcium transients.

In contrast, the RhoA-PLCε axis was determined to be functional in RPE1 and

HeLa cells (**Figures 3, 4 and 7, upper scheme**). Calcium transients were totally blocked

by 2-APB and partially observed in Ca^{2+}-free buffer, and transients were apparently

induced by the IP_3-IP3R pathway that promotes Ca^{2+} release from the ER. In the Ca^{2+}-

free buffer, the percentage of cells exhibiting transients has decreased compared with cells

in Ca^{2+}-containing buffer. This finding may reflect changes in cellular responsiveness

since cell processes, such as cell-substrate adhesion, membrane lipid composition, and

intracellular calcium homeostasis, are largely dependent on extracellular Ca^{2+}, for

example, purified PLCε exhibits Ca^{2+}-dependent phospholipase activity (Kelley et al.,

2001).

PLCε has been identified to be activated by other small GTPases, including Ras,

Rap, and Ral (Jin et al., 2001; Kelley et al., 2004, 2001), which is inconsistent with the

present results (**Figure 2**). This discrepancy is explained by the fact that there is no

evidence that small GTPases directly activate the PLCε. In previous studies,

phospholipase activity was typically examined by IP_3 accumulation in COS-7 cells co-

expressing constitutively active mutants of small GTPases and PLCε exogenously (Ada-

Nguema et al., 2006; Kelley et al., 2004, 2001; Madukwe et al., 2018; Martins et al.,

2012; Wing et al., 2003). This assay cannot exclude side effects of Ras signaling. Also,

the purified KRas G12V constitutively active mutant activated phospholipase activity of

N-terminal deleted PLCε (Seifert et al., 2008), but did not that of the full-length intact

PLCε (Song et al., 2001) in an *in vitro* reconstitution assay. Active Ras and Rap likely

interact with the RA2 domain of PLCε and induce translocation from cytoplasm to the

PM and Golgi, respectively (Jin et al., 2001; Song et al., 2001). Thus, PLCε is identified

as an effector protein of Ras and Rap GTPases, but this present study suggests that Ras

and Rap do not directly activate its phospholipase activity. Instead, other factors,

including RhoA, activate PLCε that is recruited to the membrane by Ras and Rap (Bunney

et al., 2006). We attempted to determine whether RhoA also recruits PLCε to the PM, but

definitive results were not obtained. However, membrane targeting for PLCε with the

CAAX motif is required for RhoA-dependent activation for RA2 domain-depleted PLCε

(Madukwe et al., 2018), and membrane recruitment of PLCε by RA2 domain-interacting

proteins, Ras and Rap, could be necessary for activation by RhoA. Yins of PLCε are

required for RhoA-dependent activation, yet the Y domain does not interact with RhoA

(Wing et al., 2003). A recent study has reported that RhoA interacted with the PH and

RA1-RA2 domains of PLCε (Yu et al., 2020). However, further studies are needed to

understand molecular mechanisms of PLCε activation by RhoA, including whether

translocation of PLCε by Ras or Rap1 is essential.

We have shown that RhoA-mediated calcium transients induce NFAT

translocation to the nucleus (**Figure 6**). NFAT translocation was used because of ease of

assess, but little is known about the physiological function of the RhoA-NFAT pathway

(Henstidge et al., 2009; Xu et al., 2004; Zhu et al., 2013). Instead, our findings indicate

that the RhoA-induced calcium transients activate intracellular calcium signaling. Both

RhoA and calcium signaling commonly regulate actin cytoskeleton reorganization, cell

proliferation, and gene expression (Berridge et al., 2000; Bong and Monteith, 2018;

Clapham, 2007; Cullen and Lockyer, 2002; Heasman and Ridley, 2008), and coordination

of signaling seems reasonable. To date, upstream factors and physiological roles of RhoA-

PLCε have been reported in a few cell types including kidney podocytes (Kalwa et al.,

2015; Yu et al., 2020), astrocytes (Citro et al., 2007), and cardiac cells (Xiang et al., 2013).

PLC activates not only calcium signaling but also DAG-PKC signaling. Such signaling

reportedly affects various cellular pathways, and physiological roles of the RhoA-PLCε

axis may be different among cell types. This axis should be examined in more cell types

in the future.

Materials and Methods

Antibodies

Commercial antibodies and their dilution were indicated as follows; mouse anti-GFP

antibody diluted at 1:1000 (clone mFX75, FUJIFILM Wako Pure Chemical Corp., Osaka,

Japan); anti-vinculin antibody diluted at 1:1000 (clone 2B5A7, Proteintech Group,

Chicago, IL); rabbit anti-PLCE1 antibody diluted at 1:1000 (HPA015597, Atlas

Antibodies, Bromma, Sweden); peroxidase-conjugated mouse anti-β-actin antibody

diluted at 1:10000 (clone 2F3, FUJIFILM Wako Pure Chemical Corp.); peroxidase-

conjugated sheep anti-mouse IgG antibody diluted in 1:4000 (NA931, GE Healthcare,

Little Chalfont, UK); and peroxidase-conjugated donkey anti-rabbit IgG antibody diluted

in 1:2000–1:4000 (NA934, GE Healthcare).

Cell culture and transfection

hTERT-immortalized human retinal pigment epithelial (RPE1) cells (ATCC CRL-400)

were cultured in DMEM and F12 nutrient mix (1:1), supplemented with 10 % (v/v) fetal

bovine serum (FBS: Biowest, Nuaillé, France). HeLa (RCB0007: RIKEN BRC through

the National BioResource Project of the MEXT/AMED, Japan), MDCK (kindly gifted

from Dr. N. Yui), and HEK293T (obtained from TaKaRa, Shiga, Japan) cells were

cultured in DMEM supplemented with 10 % FBS. Plasmid transfection was performed

using Lipofectamine™ 2000 (Thermo Fisher Scientific, Waltham, MA) following the

manufacturer's instructions. Reagents were obtained from commercial sources as

follows: Y-27632 dihydrochloride (Focus, Biomolecules, Plymouth Meeting, PA); (-)-

Blebbistatin (Cayman Chemical, Ann Arbor, MI); U73122 (Cayman Chemical); 2-APB

(FUJIFILM Wako Pure Chemicals); and SKF96365 (FUJIFILM Wako Pure Chemicals).

DNA construction

pLL7.0: Venus-iLID-CAAX (from KRas4B) and pLL7.0: tgRFPt-SSPB wild-type (WT)

(Guntas et al., 2015) were gifts from Brian Kuhlman (Addgene plasmid #60411 and

#60415); PR_GEF (2XPDZ-mCherry-Larg (DH)) from Michael Glotzer (Addgene

plasmid #80407); CMV-R-GECO1 (Kalko et al., 2011) from Robert Campbell (Addgene

plasmid #32444); pcDNA3-HA-human OCRL (Erdmann et al., 2007) from Pietro De

Camilli (Addgene plasmid #22207); GFP-C1-PLCdelta-PH (Stauffer et al., 1998) from

Tobias Meyer (Addgene plasmid #21179); pTriEx-RhoA FLARE.sc Biosensor WT (Pertz

et al., 2006) from Klaus Hahn (Addgene plasmid #12150); and human C3G cDNA from

Hiroki Tanaka (Kyoto University). HA-NFAT1(1-460)-mCherry was constructed

previously (Ishii et al., 2015). Total RNA was obtained from RPE1 and HeLa cells using

QIAzol Lysis Reagent (QIAGEN; Hilden, Germany) following the manufacturer's

protocol, and first-strand cDNA was synthesized with 1 µg total RNA using a SuperScript

III First-Strand Synthesis System for RT-PCR (TaKaRa) following the manufacturer's

protocol.

msPLCε cDNA was amplified by PCR from mouse liver first-strand cDNA

library (kindly gifted from Tomohiro Ishii) using Tks Gflex polymerase (TaKaRa) and

was cloned between *Nhe*I and *Kpn*I sites of pECFP-N1 (Clontech, Palo Alto, CA).

msPLCε sequences were sensitive to siPLCε seq1, and a siRNA-resistant msPLCε mutant

was used in this study. All point mutations were generated using site-directed mutagenesis

methods by Tks Gflex polymerase. msPLCε ΔYins mutant was generated with overlap

extension PCR.

To create pmCherry-PLCδ-PH, EGFP was removed from pEGFP-C1 PLCdelta-

PH and replaced with mCherry from pmCherry-C1 (TaKaRa). To create pmCherry-C1

RBD$_{rhotekin}$, RBD$_{rhotekin}$ was amplified from pTriEx-RhoA FLARE.sc Biosensor WT and

cloned between *Bgl*II and *Eco*RI site of pmCherry-C1.

Then, the DNA coding sequence was cloned between *Nhe*1 and *Not*1 of pEGFP-

N1 to create opto-RhoA (mVenus-NES-SspB-LARG-DH-P2A-iLIDcaax). The NES

sequence (FGIDLSGLTL) was directly linked to mVenus. SspB and LARG-DH (DH

domain of LARG, aa 766–986) were linked by LDSAGGSAGGSAGGLE. The P2A

peptide sequence (GSG)ATNFSLLKQAGDVEENPGP was directly linked to LARG-

DH and followed by the linker APGS to locate the 2A product protein to the precise

subcellular domain (Kakumoto and Nakata, 2013).

The transmembrane domain of 2,6-sialyltransferase (aa 1–32) was then

amplified from the HeLa cDNA library to create opto-RhoA-TGN, and KDELr was

amplified from the RPE1 cDNA library by PCR to create opto-RhoA-Golgi. These

fragments were connected to iLID by overlap extension PCR with a GSGSGS (3×GS)

linker. iLID-KDELr was first cloned into a pECFP-C1 vector (Clontech), and KDELr

D193N dominant-negative mutant (KDELr DN) was then generated by site-directed PCR.

iLID-CAAX^{Kras} of opto-RhoA was replaced with iLID-TGN or iLID-KDELr DN.

The LARG-DH of opto-RhoA was replaced with a DH-PH domain of Tiam

(Tiam-DH-PH, aa 1012–1591), a DH-PH domain of ITSN (ITSN-DH-PH, aa 1230–1580),

a catalytic domain of SOS2 (SOS2cat, aa 563–1048), a catalytic domain of C3G (C3Gcat,

aa 687–1077), and a GEF domain of RGL2 (RGL2-GEF, aa 1–518) to create opto-Rac1,

-Cdc42, -Ras, -Rap1, and -RalA/B, respectively.

Opto-5-ptase was constructed as previously reported for mCherry-PHR-iSH-2A-

CIBNcaax (Kakumoto and Nakata, 2013), in which the vector backbone was changed to

pEGFP-N1, and mCherry and iSH were replaced with ECFP and the inositol 5-phosphate

domain of OCRL (OCRLcat, aa 234–539), respectively.

All cloned fragments were verified by sequencing.

Small interference RNAs

Transfection of siRNA duplexes used Lipofectamine^{TM} RNAiMAX reagent following the

manufacturer's protocol (Thermo Fisher Scientific). Each duplex was used at a final

concentration of 10 nM. Double-strand RNAs were purchased from Ambion, Thermo

Fisher Scientific. Target nucleotide sequences were 5′–GCAAGGAGCTGATCGATCT–

3′ (s27658, siPLCε seq1), 5′–GGACATAGGCTGACAACCA–3′ (s27659, siPLCε seq2),

and 5′–TACTGCGATATTGAAGTCC–3′ (s27660, siPLCε seq3), respectively. #2 siRNA

(Silencer Select, Thermo Fisher Scientific) was used as a negative control.

Live cell imaging

When only the plasmid transfection was performed, cells were seeded on glass-bottomed

dishes (Greiner Bio-one, Kremsmünster, Austria) coated with collagen (Cellmatrix Type

IC; Nitta Gelatin Inc., Osaka, Japan) 2 days before the observations. On the next day,

cells were transfected with the plasmid and were analyzed a day later. When both siRNA

and plasmid transfection were performed, cells were seeded 3 days before the

observations. On the next day, siRNA transfection was performed. After 24 hours since

siRNA transfection, plasmid transfection was performed, and cells were analyzed a day

later. Before experiments, the medium was changed to Leibovitz's L-15 (Thermo Fisher

Scientific), Ringer's solution (138 mM NaCl, 5.6 mM KCl, 2mM $CaCl_2$, 2 mM $MgCl_2$, 4

mM D-glucose, 5 mM HEPES, and 2 mM sodium pyruvate (pH 7.4, adjusted with

NaOH)), or calcium-free Ringer's solution (sans $CaCl_2$). Cells were analyzed under

serum-starved conditions, and a closed heated chamber was used during live cell imaging

at 37 °C without CO_2. Images were obtained with a confocal laser scanning microscope

(FV1200, Olympus, Tokyo, Japan) on an IX83 microscope (Olympus) equipped with

40×/0.95 NA and 20×/0.7 NA dry objective lenses and FV10-ASW software (Olympus).

A laser line (458 nm) was used to photo-activated photoswitches. Local irradiation was

performed using multi-area time-lapse software (Olympus). Image analysis used

ImageJ/Fiji software (NIH) (Schindelin et al., 2012). Adobe Illustrator CS6 (Adobe

Systems, San Jose, CA) was used for final figure preparation.

Immunoblotting

Cells were lysed with 2 × SDS sample buffer (125 mM Tris-HCl (pH6.8), 4 % (w/v) SDS,

10 % (w/v) sucrose, 0.01 % (w/v) BPB, 5 % (v/v) 2-mercaptoethanol). Proteins were

separated with SDS-PAGE and then later transferred to PVDF membranes by

electrophoresis using a Bio-Rad Trans-Blot device (Bio-Rad, Richmond, CA). Blots were

blocked for an hour at room temperature or overnight at 4 °C with 5 % skim milk in Tris-

buffer saline with Tween-20 (TBS-T, 50 mM Tris-HCl (pH 7.4), 138 mM NaCl, 2.7 mM

KCl, and 0.1 % (v/v) Tween-20). Membranes were washed with TBS-T and incubated

with indicated antibodies diluted in Can Get Signal Solution 1 (TOYOBO, Osaka, Japan)

for 1 h at room temperature. Membranes were washed in TBS-T, incubated with

appropriate horseradish peroxidase-conjugated secondary antibodies diluted in Can Get

Signal Solution 2 (TOYOBO) for 1 h at room temperature, and washed again in TBS-T.

Blots were developed with ImmunoStar Zeta (FUJIFILM Wako Pure Chemical), and

chemiluminescence signals were visualized via a ChemiDoc MP system (Bio-Rad).

Statistics

Statistical analyses were performed using Excel (Microsoft) and R software v.3.5.1 (R

Development Core Team, 2016). All graphs were drawn using Excel software.

Acknowledgments

The authors are grateful to Dr. Naofumi Yui (TMDU) for the kind gift of MDCK cells

and Toshiyuki Kakumoto for plasmid construction. We also thank Drs. Tomohiro Ishii

and Toshifumi Asano for the helpful discussion and Satoko Nakamura for secretarial

expertise.

Funding

This work was supported in part by JSPS KAKENHI (18K15002 and 20K16105 to H.I.,

and 18K19328 to H.I. and T.N.), by Research Grants from the Takeda Science Foundation

(to H.I.), and the Kanehara Ichiro Memorial Foundation (to H.I.).

Authors' contributions

H.I. and T.N. designed the experiments and wrote the manuscript. H.I. and Q.M.

performed the experiments, and H.I. analyzed the data.

Conflict of interests

The authors declare that they have no conflict of interests.

References

Ada-Nguema AS, Xenias H, Sheetz MP, Keely PJ. 2006. The small GTPase R-Ras

regulates organization of actin and drives membrane protrusions through the

activity of PLCε. *J Cell Sci*. doi:10.1242/jcs.02835

Amano M, Ito M, Kimura K, Fukata Y, Chihara K, Nakano T, Matsuura Y, Kaibuchi K.

1996. Phosphorylation and activation of myosin by Rho-associated kinase (Rho-

kinase). *J Biol Chem*. doi:10.1074/jbc.271.34.20246

Aspenström P. 2004. Integration of signalling pathways regulated by small GTPases

and calciumBiochimica et Biophysica Acta - Molecular Cell Research.

doi:10.1016/j.bbamcr.2004.09.029

Berridge MJ, Lipp P, Bootman MD. 2000. The versatility and universality of calcium

signalling. *Nat Rev Mol Cell Biol*. doi:10.1038/35036035

Bong AHL, Monteith GR. 2018. Calcium signaling and the therapeutic targeting of

cancer cells. *Biochim Biophys Acta - Mol Cell Res*.

doi:10.1016/j.bbamcr.2018.05.015

Bunney TD, Harris R, Gandarillas NL, Josephs MB, Roe SM, Sorli SC, Paterson HF,

Rodrigues-Lima F, Esposito D, Ponting CP, Gierschik P, Pearl LH, Driscoll PC,

Katan M. 2006. Structural and mechanistic insights into ras association domains of

phospholipase C epsilon. *Mol Cell*. doi:10.1016/j.molcel.2006.01.008

Chao TSO, Byron KL, Lee KM, Villereal M, Rosner MR. 1992. Activation of MAP

kinases by calcium-dependent and calcium-independent pathways. Stimulation by

thapsigargin and epidermal growth factor. *J Biol Chem*.

Charras GT, Yarrow JC, Horton MA, Mahadevan L, Mitchison TJ. 2005. Non-

equilibration of hydrostatic pressure in blebbing cells. *Nature* **435**:365–369.

doi:10.1038/nature03550

Christie JM, Salomon M, Nozue K, Wada M, Briggs WR. 1999. LOV (light, oxygen, or

voltage) domains of the blue-light photoreceptor phototropin (nph1): Binding sites

for the chromophore flavin mononucleotide. *Proc Natl Acad Sci U S A* **96**:8779–

8783. doi:10.1073/pnas.96.15.8779

Citro S, Malik S, Oestreich EA, Radeff-Huang J, Kelley GG, Smrcka A V., Brown JH.

2007. Phospholipase Cε is a nexus for Rho and Rap-mediated G protein-coupled

receptor-induced astrocyte proliferation. *Proc Natl Acad Sci U S A* **104**:15543–

15548. doi:10.1073/pnas.0702943104

Clapham DE. 2007. Calcium Signaling. *Cell*. doi:10.1016/j.cell.2007.11.028

Cox CD, Bae C, Ziegler L, Hartley S, Nikolova-Krstevski V, Rohde PR, Ng CA, Sachs

F, Gottlieb PA, Martinac B. 2016. Removal of the mechanoprotective influence of

the cytoskeleton reveals PIEZO1 is gated by bilayer tension. *Nat Commun*.

doi:10.1038/ncomms10366

Cullen PJ, Lockyer PJ. 2002. Integration of calcium and Ras signalling. *Nat Rev Mol*

Cell Biol. doi:10.1038/nrm808

de Beco S, Vaidžiulytė K, Manzi J, Dalier F, di Federico F, Cornilleau G, Dahan M,

Coppey M. 2018. Optogenetic dissection of Rac1 and Cdc42 gradient shaping. *Nat*

Commun. doi:10.1038/s41467-018-07286-8

Erdmann KS, Mao Y, McCrea HJ, Zoncu R, Lee S, Paradise S, Modregger J,

Biemesderfer D, Toomre D, De Camilli P. 2007. A Role of the Lowe Syndrome

Protein OCRL in Early Steps of the Endocytic Pathway. *Dev Cell* **13**:377–390.

doi:10.1016/j.devcel.2007.08.004

Gudipaty SA, Lindblom J, Loftus PD, Redd MJ, Edes K, Davey CF, Krishnegowda V,

Rosenblatt J. 2017. Mechanical stretch triggers rapid epithelial cell division

through Piezo1. *Nature*. doi:10.1038/nature21407

Guglielmi G, Falk HJ, De Renzis S. 2016. Optogenetic Control of Protein Function:

From Intracellular Processes to Tissue Morphogenesis. *Trends Cell Biol* **26**:864–

874. doi:10.1016/j.tcb.2016.09.006

Guntas G, Hallett RA, Zimmerman SP, Williams T, Yumerefendi H, Bear JE, Kuhlman

B. 2015. Engineering an improved light-induced dimer (iLID) for controlling the

localization and activity of signaling proteins. *Proc Natl Acad Sci U S A* **112**:112–

117. doi:10.1073/pnas.1417910112

Hall BE, Yang SS, Boriack-Sjodin PA, Kuriyan J, Bar-Sagi D. 2001. Structure-based

Mutagenesis Reveals Distinct Functions for Ras Switch 1 and Switch 2 in Sos-

catalyzed Guanine Nucleotide Exchange. *J Biol Chem*.

doi:10.1074/jbc.M101727200

Harden TK, Hicks SN, Sondek J. 2009. Phospholipase C isozymes as effectors of Ras

superfamily GTPases. *J Lipid Res*. doi:10.1194/jlr.R800045-JLR200

Heasman SJ, Ridley AJ. 2008. Mammalian Rho GTPases: New insights into their

functions from in vivo studies. *Nat Rev Mol Cell Biol*. doi:10.1038/nrm2476

Henstidge CM, Balenga NAB, Ford LA, Ross RA, Waldhoer M, Irving AJ. 2009. The

GPR55 ligand L-α-lysophosphatidylinositol promotes RhoA-dependent Ca 2+

signaling and NFAT activation . *FASEB J*. doi:10.1096/fj.08-108670

Hogan PG, Chen L, Nardone J, Rao A. 2003. Transcriptional regulation by calcium,

calcineurin, and NFAT. *Genes Dev*. doi:10.1101/gad.1102703

Idevall-Hagren O, Dickson EJ, Hille B, Toomre DK, De Camilli P. 2012. Optogenetic

control of phosphoinositide metabolism. *Proc Natl Acad Sci U S A* **109**.

doi:10.1073/pnas.1211305109

Ikebe M, Hartshorne DJ. 1985. Phosphorylation of smooth muscle myosin at two

distinct sites by myosin light chain kinase. *J Biol Chem*.

Irannejad R, Wedegaertner PB. 2010. Regulation of constitutive cargo transport from

the trans-Golgi network to plasma membrane by Golgi-localized G protein βγ

subunits. *J Biol Chem* **285**:32393–32404. doi:10.1074/jbc.M110.154963

Ishii T, Sato K, Kakumoto T, Miura S, Touhara K, Takeuchi S, Nakata T. 2015. Light

generation of intracellular Ca2+ signals by a genetically encoded protein BACCS.

Nat Commun **6**. doi:10.1038/ncomms9021

Jin P, Jan LY, Jan Y-N. 2020. Mechanosensitive Ion Channels: Structural Features

Relevant to Mechanotransduction Mechanisms. *Annu Rev Neurosci*.

doi:10.1146/annurev-neuro-070918-050509

Jin TG, Satoh T, Liao Y, Song C, Gao X, Kariya KI, Hu CD, Kataoka T. 2001. Role of

the CDC25 Homology Domain of Phospholipase Cε in Amplification of Rap1-

dependent Signaling. *J Biol Chem* **276**:30301–30307.

doi:10.1074/jbc.M103530200

Kakumoto T, Nakata T. 2013. Optogenetic Control of PIP3: PIP3 Is Sufficient to Induce

the Actin-Based Active Part of Growth Cones and Is Regulated via Endocytosis.

PLoS One **8**:1–17. doi:10.1371/journal.pone.0070861

Kalko EK V, Dukas R, Ratcliffe JM, Teeling EC, Haven N, Fattu JM, Bates ME,

Simmons J a, Riquimaroux H, Surlykke A, Bouffard FH, Lee DN, Dear SP,

Horiuchi TK, Krishnaprasad PS, Moss CF, Schuller G, Brudzynski SM, Syme D a,

Hollingworth S, Lindstedt SL, Baylor SM, Mead   a F, Rome LC, Goller F, Spiers

ILY, Leeuwen JL Van, Schachat F, Rossmanith GH, Hoh JFY, Kelley DB, Vater

M a, Zhao Y, Kalko EK V, Dukas R, Ratcliffe JM, Teeling EC, Haven N, Fattu

JM, Bates ME, Simmons J a, Riquimaroux H, Surlykke A, Bouffard FH, Lee DN,

Dear SP, Horiuchi TK, Krishnaprasad PS, Moss CF, Schuller G, Brudzynski SM,

Syme D a, Hollingworth S, Lindstedt SL, Baylor SM, Mead   a F, Rome LC,

Goller F, Spiers ILY, Leeuwen JL Van, Schachat F, Rossmanith GH, Hoh JFY,

Kelley DB, Vater M a. 2011. An Expanded Palette of Genetically. *Science (80-)*

557:1888–1891. doi:10.1126/science.1208592

Kalwa H, Storch U, Demleitner J, Fiedler S, Mayer T, Kannler M, Fahlbusch M, Barth

H, Smrcka A, Hildebrandt F, Gudermann T, Dietrich A. 2015. Phospholipase C

Epsilon (PLCε) Induced TRPC6 Activation: A Common but Redundant

Mechanism in Primary Podocytes. *J Cell Physiol* **230**:1389–1399.

doi:10.1002/jcp.24883

Kelley GG, Reks SE, Ondrako JM, Smrcka A V. 2001. Phospholipase Cε: A novel Ras

effector. *EMBO J*. doi:10.1093/emboj/20.4.743

Kelley GG, Reks SE, Smrcka A V. 2004. Hormonal regulation of phospholipase Cε

through distinct and overlapping pathways involving G12 and Ras family G-

proteins. *Biochem J*. doi:10.1042/BJ20031370

Kennedy MJ, Hughes RM, Peteya LA, Schwartz JW, Ehlers MD, Tucker CL. 2010.

Rapid blue-light-mediated induction of protein interactions in living cells. *Nat

Methods* **7**:973–975. doi:10.1038/nmeth.1524

Kolch W. 2000. Meaningful relationships: the regulation of the Ras/Raf/MEK/ERK

pathway by protein interactions. *Biochem J*. doi:10.1042/bj3510289

Levskaya A, Weiner OD, Lim WA, Voigt CA. 2009. Spatiotemporal control of cell

signalling using a light-switchable protein interaction. *Nature*.

doi:10.1038/nature08446

Li Y, Brayden JE. 2017. Rho kinase activity governs arteriolar myogenic

depolarization. *J Cereb Blood Flow Metab* **37**:140–152.

doi:10.1177/0271678X15621069

Lopez I, Mak EC, Ding J, Hamm HE, Lomasney JW. 2001. A Novel Bifunctional

Phospholipase C That Is Regulated by Gα 12 and Stimulates the Ras/Mitogen-

activated Protein Kinase Pathway. *J Biol Chem* **276**:2758–2765.

doi:10.1074/jbc.M008119200

Madukwe JC, Garland-Kuntz EE, Lyon AM, Smrcka A V. 2018. G protein subunits

directly interact with and activate phospholipase C. *J Biol Chem*.

doi:10.1074/jbc.RA118.002354

Malik S, De Rubio RG, Trembley M, Irannejad R, Wedegaertner PB, Smrcka A V.

2015. G protein βγ subunits regulate cardiomyocyte hypertrophy through a

perinuclear Golgi phosphatidylinositol 4-phosphate hydrolysis pathway. *Mol Biol

Cell* **26**:1188–1198. doi:10.1091/mbc.E14-10-1476

Martins M, Warren S, Kimberley C, Margineanu A, Peschard P, McCarthy A, Yeo M,

Marshall CJ, Dunsby C, French PMW, Katan M. 2012. Activity of PLCε

contributes to chemotaxis of fibroblasts towards PDGF. *J Cell Sci*.

doi:10.1242/jcs.110007

Martz MK, Grabocka E, Beeharry N, Yen TJ, Wedegaertner PB. 2013. Leukemia-

associated RhoGEF (LARG) is a novel RhoGEF in cytokinesis and required for the

proper completion of abscission. *Mol Biol Cell* **24**:2785–2794.

doi:10.1091/mbc.E12-07-0533

Mehta D, Ahmmed GU, Paria BC, Holinstat M, Voyno-Yasenetskaya T, Tiruppathi C,

Minshall RD, Malik AB. 2003. RhoA interaction with inositol 1,4,5-trisphosphate

receptor and transient receptor potential channel-1 regulates Ca2+ entry: Role in

signaling increased endothelial permeability. *J Biol Chem* **278**:33492–33500.

doi:10.1074/jbc.M302401200

Mitin N, Rossman KL, Der CJ. 2005. Signaling interplay in ras superfamily function.

Curr Biol **15**:563–574. doi:10.1016/j.cub.2005.07.010

Nash CA, Wei W, Irannejad R, Smrcka A V. 2019. Golgi localized βi-adrenergic

receptors stimulate golgi PI4P hydrolysis by PLCε to regulate cardiac hypertrophy.

Elife **8**:1–22. doi:10.7554/eLife.48167

O'Neill PR, Castillo-Badillo JA, Meshik X, Kalyanaraman V, Melgarejo K, Gautam N.

2018. Membrane Flow Drives an Adhesion-Independent Amoeboid Cell Migration

Mode. *Dev Cell* **46**:9-22.e4. doi:10.1016/j.devcel.2018.05.029

Pertz O, Hodgson L, Klemke RL, Hahn KM. 2006. Spatiotemporal dynamics of RhoA

activity in migrating cells. *Nature* **440**:1069–1072. doi:10.1038/nature04665

Reid T, Furuyashiki T, Ishizaki T, Watanabe G, Watanabe N, Fujisawa K, Morii N,

Madaule P, Narumiya S. 1996. Rhotekin, a new putative target for Rho bearing

homology to a serine/threonine kinase, PKN, and rhophilin in the Rho-binding

domain. *J Biol Chem* **271**:13556–13560. doi:10.1074/jbc.271.23.13556

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch

S, Rueden C, Saalfeld S, Schmid B, Tinevez JY, White DJ, Hartenstein V, Eliceiri

K, Tomancak P, Cardona A. 2012. Fiji: An open-source platform for biological-

image analysis. *Nat Methods*. doi:10.1038/nmeth.2019

Seifert JP, Zhou Y, Hicks SN, Sondek J, Harden TK. 2008. Dual activation of

phospholipase C-ε by rho and ras GTPases. *J Biol Chem*.

doi:10.1074/jbc.M805038200

Shiwarski DJ, Darr M, Telmer CA, Bruchez MP, Puthenveedu MA. 2017. PI3K class II

α regulates δ-opioid receptor export from the trans-Golgi network. *Mol Biol Cell*

28:2202–2219. doi:10.1091/mbc.E17-01-0030

Smrcka A V., Brown JH, Holz GG. 2012. Role of phospholipase Cε in physiological

phosphoinositide signaling networks. *Cell Signal*.

doi:10.1016/j.cellsig.2012.01.009

Song C, Hu CD, Masago M, Kariya KI, Yamawaki-Kataoka Y, Shibatohge M, Wu D,

Satoh T, Kataoka T. 2001. Regulation of a Novel Human Phospholipase C, PLCε,

through Membrane Targeting by Ras. *J Biol Chem*. doi:10.1074/jbc.M008324200

Stauffer TP, Ahn S, Meyer T. 1998. Receptor-induced transient reduction in plasma

membrane Ptdlns(4,5)P2 concentration monitored in living cells. *Curr Biol* **8**:343–

346. doi:10.1016/s0960-9822(98)70135-6

Strickland D, Lin Y, Wagner E, Hope CM, Zayner J, Antoniou C, Sosnick TR, Weiss EL, Glotzer M. 2012. TULIPs: Tunable, light-controlled interacting protein tags for cell biology. *Nat Methods* **9**:379–384. doi:10.1038/nmeth.1904

Team RDC. 2016. R: A language and environment for statistical computing.

Tischer D, Weiner OD. 2014. Illuminating cell signalling with optogenetic tools. *Nat Rev Mol Cell Biol* **15**:551–558. doi:10.1038/nrm3837

Toettcher JE, Voigt CA, Weiner OD, Lim WA. 2011. The promise of optogenetics in cell biology: Interrogating molecular circuits in space and time. *Nat Methods* **8**:35–38. doi:10.1038/nmeth.f.326

Toettcher JE, Weiner OD, Lim WA. 2013. Using optogenetics to interrogate the dynamic control of signal transmission by the Ras/Erk module. *Cell*. doi:10.1016/j.cell.2013.11.004

Valon L, Marín-Llauradó A, Wyatt T, Charras G, Trepat X. 2017. Optogenetic control of cellular forces and mechanotransduction. *Nat Commun* **8**. doi:10.1038/ncomms14396

Wagner E, Glotzer M. 2016. Local RhoA activation induces cytokinetic furrows independent of spindle position and cell cycle stage. *J Cell Biol* **213**:641–649.

doi:10.1083/jcb.201603025

Weng Y, Yan F, Chen R, Qian M, Ou Y, Xie S, Zheng H, Li J. 2018. PIEZO channel

protein naturally expressed in human breast cancer cell MDA-MB-231 as probed

by atomic force microscopy. *AIP Adv*. doi:10.1063/1.5025036

Wennerberg K, Rossman KL, Der CJ. 2005. The Ras superfamily at a glance. *J Cell Sci*

118:843–846. doi:10.1242/jcs.01660

Wilsher NE, Court WJ, Ruddle R, Newbatt YM, Aherne W, Sheldrake PW, Jones NP,

Katan M, And SAE, Raynaud FI, Therapeutics E. 2007. The Phosphoinositide-

Specific Phospholipase C Inhibitor Conjugates with Common Components of Cell

Culture. *Drug Metab Dispos*. doi:10.1124/dmd.106.014498.activity

Wing MR, Houston D, Kelley GG, Der CJ, Siderovski DP, Harden TK. 2001.

Activation of Phospholipase C-ε by Heterotrimeric G Protein βγ-Subunits. *J Biol*

Chem. doi:10.1074/jbc.C100574200

Wing MR, Snyder JT, Sondek J, Harden TK. 2003. Direct Activation of Phospholipase

C-ε by Rho. *J Biol Chem* **278**:41253–41258. doi:10.1074/jbc.M306904200

Wu YI, Frey D, Lungu OI, Jaehrig A, Schlichting I, Kuhlman B, Hahn KM. 2009. A

genetically encoded photoactivatable Rac controls the motility of living cells.

Nature **461**:104–108. doi:10.1038/nature08241

Xiang SY, Ouyang K, Yung BS, Miyamoto S, Smrcka A V., Chen J, Brown JH. 2013.

PLCε, PKD1, and SSH1L transduce RhoA signaling to protect mitochondria from

oxidative stress in the heart. *Sci Signal*. doi:10.1126/scisignal.2004405

Xu FP, Chen MS, Wang YZ, Yi Q, Lin SB, Chen AF, Luo JD. 2004. Leptin induces

hypertrophy via endothelin-1-reactive oxygen species pathway in cultured neonatal

rat cardiomyocytes. *Circulation*. doi:10.1161/01.CIR.0000140766.52771.6D

Yu S, Choi W Il, Choi YJ, Kim HY, Hildebrandt F, Gee HY. 2020. PLCE1 regulates

the migration, proliferation, and differentiation of podocytes. *Exp Mol Med*.

doi:10.1038/s12276-020-0410-4

Zago G, Veith I, Singh MK, Fuhrmann L, De Beco S, Remorino A, Takaoka S, Palmeri

M, Berger F, Brandon N, El Marjou A, Vincent-Salomon A, Camonis J, Coppey

M, Parrini MC. 2018. Ralb directly triggers invasion downstream ras by

mobilizing the wave complex. *Elife* **7**:1–23. doi:10.7554/eLife.40474

Zhang L, Malik S, Pang J, Wang H, Park KM, Yule DI, Blaxall BC, Smrcka A V. 2013.

Phospholipase Cε Hydrolyzes Perinuclear Phosphatidylinositol 4-Phosphate to

Regulate Cardiac Hypertrophy. *Cell* **153**:216–227. doi:10.1016/j.cell.2013.02.047

Zhao Y, Araki S, Wu J, Teramoto T, Chang YF, Nakano M, Abdelfattah AS, Fujiwara

M, Ishihara T, Nagai T, Campbell RE. 2011. An expanded palette of genetically

1   encoded Ca2+ indicators. *Science (80-)*. doi:10.1126/science.1208592

2   Zhu L, Qi XY, Aoudjit L, Mouawad F, Baldwin C, Nattel S, Takano T. 2013. Nuclear

3   factor of activated T cells mediates RhoA-induced fibronectin upregulation in

4   glomerular podocytes. *Am J Physiol - Ren Physiol*.

5   doi:10.1152/ajprenal.00495.2012

Figure legends

Figure 1. Development of an optogenetic tool to control RhoA activity. (A) Schematic

of opto-RhoA. Spatiotemporal control of RhoA activity is achieved using an optogenetic

probe to recruit RhoA-specific GEF LARG to the PM. An iLID molecule is anchored to

the PM via the CAAX motif, while a protein consisting of SspB fused to the DH domain

of LARG is distributed throughout the cytosol. When irradiated with blue-light, iLID

undergoes a conformational change exposing a binding site for SspB, and driving LARG-

DH to the PM, where it activates RhoA. **(B)** Representative images of HeLa cells

expressing opto-RhoA before (458 nm light OFF) and after (458 nm light ON) light

irradiation with a 458 nm laser. **(C)** Representative images of HeLa cells expressing opto-

RhoA and mCherry-RBD$_{rhotekin}$. Opto-RhoA was locally activated with a 458 nm laser

within 300–600 sec every 20 sec. Blue boxes in images indicate the activated area, and

lower panels showed high magnification images for this area. **(D, E)** Quantification of

local intensity increases of mVenus-SspB-LARG-DH **(D)** and mCherry-RBD$_{rhotekin}$ **(E)**

from the images shown in **(C)**. Activation period is indicated by a blue background.

Experiments were repeated seven times.

Figure 2. RhoA activation induces intracellular calcium transients in various cell

types. (A, B) Screening for small GTPases that induce intracellular calcium mobilization.

RPE1 (A) and HeLa (B) cells transiently express opto-GTPases, and the calcium reporter

R-GECO1 were observed with confocal microscopy in Leibovitz's L-15 medium. Opto-

GTPases were activated by a multi-argon 458 nm laser every 10 sec for 2 minutes.

Percentages of cells exhibiting intracellular calcium transients ($F/F_0 > 2$) in response to

opto-GTPases activation are shown. Data are presented as means ± SEM from more than

three independent experiments. In total, >150 cells were analyzed for each photoswitch.

(C) Representative images of RPE1 cells expressing opto-RhoA or GEF-inactive mutant

opto-RhoA^{YA} and calcium reporter, R-GECO1. From time 0 sec, opto-RhoA was

activated by a multi-argon 458 nm laser every 10 sec. Scale bar = 20 μm. (D) Intensity

changes of R-GECO1 during RhoA activation in various cell types. Only cells exhibiting

intracellular calcium transients were analyzed. Data are presented as means ± SEM from

indicated numbers of cells.

Figure 3. Mechanisms of intracellular calcium transients induced by RhoA are

different among cell types. RPE1 (A), HeLa (B), MDCK (C), and HEK293T (D) cells

expressing opto-RhoA and R-GECO1 were treated with DMSO, 50 μM Y-27632 (ROCK

inhibitor), 20 µM Blebbistatin (myosin II inhibitor), 10 µM SKF96365, 100 µM 2-APB

(calcium channel inhibitors), or 10 µM U73122 (PLC inhibitor) and were observed using

confocal microscopy (see Figure 2). Leibovitz's L-15 medium and Ringer's solution with

or without Ca^{2+} were used for the experiments. Percentages of cells exhibiting

intracellular calcium transients ($F/F_0 > 2$) in response to RhoA activation are shown. Data

are shown as means $\pm$ SEM from more than three independent experiments. In total, >200

cells were analyzed for each condition. ***, $p < 0.001$; **, $p < 0.01$; and n.s., not

significant; two-tailed unpaired Student's t-test.

Figure 4. RhoA directly activates PLCε and induces intracellular calcium transients

in RPE1 and HeLa cells. (A) RPE1 and HeLa cells were treated with PLCε-specific or

control siRNAs for 48 h. Proteins in total cell extracts were separated with SDS-PAGE

and immunoblotted with anti-PLCε and anti-vinculin antibodies. Vinculin was used as a

loading control. **(B)** siRNA-mediated PLCε-depleted RPE1 and HeLa cells express opto-

RhoA and R-GECO1 (see Figure 2). **(C)** Schematic of mouse PLCε (msPLCε). CDC25,

CDC25 homology guanine nucleotide exchange factor (GEF) domain; PH, pleckstrin

homology domain; EF, EF-hand motif; X and Y, X and Y domains fold to form the

catalytic core of the phospholipase; C2, C2 domain; RA, ras (RA) association homology

domain. **(D)** Proteins in total extracts from HEK293T cells expressing ECFP, msPLCε-

ECFP wild-type (WT), or indicated msPLCε-ECFP mutants were separated with SDS-

PAGE and immunoblotted with anti-GFP and anti-β-actin antibodies. β-actin was used as

a loading control. **(E)** siRNA-mediated PLCε-depleted HeLa cells transiently expressing

ECFP, msPLCε-ECFP WT or indicated msPLCε-ECFP mutants, and R-GECO1 (see

Figure 2). Data are presented as means ± SEM from three independent experiments. In

total, >150 cells were analyzed for each condition. ***, $p < 0.001$; and n.s., not

significant; two-tailed unpaired Student's t-test.

Figure 5. RhoA activates PLCε at the PM and marginally affects membrane

phosphoinositide dynamics. (A) Schematic of opto-RhoA-TGN and opto-RhoA-Golgi.

TGN, trans-Golgi network localized N-terminus transmembrane domain of 2,6-

sialyltransferase; KDELr DN, dominant-negative (DN) KDELr D193N. **(B)**

Representative images of HeLa cells expressing opto-RhoA-TGN or opto-RhoA-Golgi

before (458 nm light OFF) and after (458 nm light ON) light irradiation with a 458 nm

laser. **(C)** RPE1 and HeLa cells expressing opto-RhoA, opto-RhoA-TGN, or opto-RhoA-

Golgi, and R-GECO1 (see Figure 2). Data are presented as means ± SEM from three

independent experiments (in total >150 cells were analyzed for each photoswitch). ***,

$p < 0.001$; **, $p < 0.01$; and n.s., not significant; one-way ANOVA and Tukey's test. **(D)**

Schematic of opto-5-ptase. OCRLcat, inositol 5-phosphatase catalytic domain of OCRL.

(E) Representative images of cells expressing opto-5-ptase or opto-RhoA and $PI(4,5)P_2$

marker, mCherry-PLCδ-PH, before and after 458 nm light irradiation. **(F)** Quantification

of cytosolic mCherry-PLCδ-PH intensity. HeLa cells expressing indicated photoswitches

(opto-5-ptase or opto-RhoA), and markers (mCherry-PLCδ-PH or mCherry alone) were

observed by confocal microscopy with or without 458 nm light irradiation every 10 s (458

nm ON or 458 nm OFF, respectively). Cells were first observed without 458 nm light

irradiation, and then the same cells were observed with light irradiation. Data are

presented as means ± SEM from the indicated number of experiments.

Figure 6. Optogenetic RhoA activation induces intracellular calcium signaling (A)

Time-lapse imaging of NFAT-mCherry translocation in HeLa cells. Cells expressing opto-

RhoA and NFAT-mCherry were observed by confocal microscopy with a 458 nm laser

irradiation every 10 sec. Bar = 20 μm. **(B)** Time-course of the relative fluorescence

intensity of nuclear NFAT-mCherry. Cells indicated in **A** were analyzed. **(C, D)** Nuclear

to cytosol ratio (**C**) and changes of this ratio (**D**) of NFAT-mCherry before (dark) and after

5 min (Lit). HeLa cells were treated with individual siRNA duplexes for 48 h and

transiently expressed opto-RhoA and NFAT-mCherry. Data from three independent

experiments is shown in box-and-whisker plots. Boxes represent the first to third quartiles

(interquartile range: IQR). The horizontal line inside the box represents the median—

vertical lines above and below the box span 1.5×IQR. Dots represent outliers that are

above or below 1.5×IQR. In total, 61, 58, and 72 cells treated with si control, si PLCε

seq1, and seq3 were analyzed, respectively. ***, $p < 0.001$; **, $p < 0.01$; and n.s., not

significant; two-tailed paired Student's t-test (**C**) and one-way ANOVA followed by

Wilcoxon signed-rank test (**D**).

Figure 7. Summary of the study. See details in the discussion.

A RPE1

B HeLa

C

mVenus-SspB-LARG-DH | R-GECO1

D

A

opto-RhoA-TGN | mVenus | SspB | LARG-DH | iLID | TGN
opto-RhoA-Golgi | mVenus | SspB | LARG-DH | iLID | KDELr DN

NES, P2A, TGN labels

B
458 nm light OFF | 458 nm light ON
opto-RhoA-TGN
opto-RhoA-Golgi
20 μm

C
- opto-RhoA
- opto-RhoA-TGN
- opto-RhoA-Golgi

Cells with calcium transients (%)

RPE1, HeLa

D
Opto-5-ptase | ECFP | PHR | OCRLcat | CIBN | -CAAX^{KRas}

T2A

E
458 nm OFF | 458 nm light ON
mCherry-PLCδ-PH
opto-5-ptase: 0s, 20s
opto-RhoA: 0s, 200s

F
- opto-5-ptase, mCherry-PLCδ-PH, 458 nm ON (n = 7)
- opto-RhoA, mCherry-PLCδ-PH, 458 nm ON (n = 33)
- opto-RhoA, mCherry-PLCδ-PH, 458 nm OFF (n = 33)
- opto-RhoA, mCherry, 458 nm ON (n = 24)
- opto-RhoA, mCherry, 458 nm OFF (n = 24)

Cytosolic mCherry-PLCδ-PH

RPE1, HeLa

MDCK, HEK293T