Integration of plant primary and secondary metabolites as fitness-relevant foraging cues for a specialist herbivore

Ricardo A. R. Machado[a,b]*, Vanitha Theepan[a], Christelle A.M. Robert[a], Tobias Züst[a], Lingfei Hu[a,c,d], Qi Su[a], Bernardus C. J. Schimmel[a], Matthias Erb[a]*

[a] Institute of Plant Sciences, University of Bern, Altenbergrain 21, 3013 Bern, Switzerland

[b] Institute of Biology, University of Neuchâtel, Rue Emile-Argand 11, 2000 Neuchâtel, Switzerland

[c] Institute of Soil and Water Resources and Environmental Science, College of Environmental and Resource Sciences, Zhejiang University, Yuhangtang road 866, 310058 Hangzhou, China

[d] Zhejiang Provincial Key Laboratory of Agricultural Resources and Environment, Zhejiang University, Yuhangtang road 866, 310058 Hangzhou, China

* Corresponding authors: Matthias Erb & Ricardo A. R. Machado.

Email: matthias.erb@ips.unibe.ch; ricardo.machado@unine.ch

Keywords: Chemical ecology, plant-insect interactions, animal behavior, plant defense.

Author Contributions: R.A.R.M and M.E. conceived the original project. R.A.R.M., C.A.M.R., L.H. and M.E. designed experiments. R.A.R.M., V.T., C.A.M.R., L.H., Q.S. and B.C.J.S. performed experiments. R.A.R.M., C.A.M.R., T.Z., L.H. and M.E. analyzed experiments. M.E. and R.A.R.M. wrote the first draft of the manuscript. All authors contributed to the final version of the manuscript.

Abstract

Plants produce complex mixtures of primary and secondary metabolites. Herbivores use these
metabolites as behavioral cues to increase their fitness. However, how herbivores integrate primary
and secondary metabolites into fitness-relevant foraging decisions *in planta* is poorly understood.
We developed a molecular manipulative approach to modulate the availability of sugars and
benzoxazinoid secondary metabolites as foraging cues for a specialist maize herbivore, the western
corn rootworm. By disrupting benzoxazinoid biosynthesis in maize and sugar perception in the
western corn rootworm, we show that sugars and benzoxazinoids act as distinct and dynamically
integrated mediators of short-distance host finding and acceptance. While sugars improve the
capacity of rootworm larvae to find a host plant and to distinguish post-embryonic from less
nutritious embryonic roots, benzoxazinoids are specifically required for the latter. Host acceptance
in the form of root damage is increased by benzoxazinoids and sugars in an additive manner. This
pattern is driven by increasing damage to post-embryonic roots in the presence of benzoxazinoids
and sugars. Benzoxazinoid- and sugar-mediated foraging directly improves western corn rootworm
growth and survival. Interestingly, western corn rootworm larvae retain a substantial fraction of
their capacity to feed and survive on maize plants in the absence of both cues. This study unravels
fine-grained differentiation and integration of primary and secondary metabolites into herbivore
foraging and documents how the capacity to compensate for the absence of important chemical
cues enables a specialist herbivore to survive within unpredictable metabolic landscapes.

Introduction

Herbivore foraging behavior contributes to the distribution and performance of herbivores and plants in natural and agricultural ecosystems (1–3). Insect herbivores often exhibit pronounced oviposition and feeding preferences for specific plant species, genotypes within species physiological states within genotypes (1, 4, 5). Most insect herbivores also show characteristic preferences for specific plant organs and tissues (6–8).

Herbivores establish preferences through different types of host-cues (9–12). Chemical cues, including plant primary and secondary metabolites, are particularly important for herbivores, as they provide specific information about the identity, physiological status and nutritional value of host plants and tissues (13–15). The overarching view in the field of chemical ecology is that primary metabolites are used as cues to identify nutritious hosts and tissues, while volatile and non-volatile secondary metabolites are used as indicators of toxicity and defense status, and as signature cues of specific host plant lineages and species (16–18). Specialized herbivores in particular are often attracted and stimulated by host-specific secondary metabolites (19). Over the past decades, herbivores have been found to respond to a multitude of plant primary and secondary metabolites in artificial diet experiments (13, 16, 17). An increasing number of studies now also document the importance of these metabolites *in planta* through molecular manipulative approaches (20–26).

Plants produce diverse sets of primary and secondary metabolites (27), and herbivores likely integrate many of these metabolites into their foraging behavior (16, 28, 29). Chemical cue integration is thought to allow herbivores to obtain more accurate information about the nutritional value and toxicity of complex host plant metabolomes (17) and thus to increase the robustness of their foraging decisions (12, 29). Many insect herbivores are known to be attracted to specific combinations of volatile chemicals, for instance (14, 29). Furthermore, some herbivores avoid combinations of secondary metabolites more strongly than individual compounds (30–32). A limited number of studies also indicate that herbivores may be able to integrate primary and secondary metabolites into their foraging strategies (17). Using artificial diet experiments, it was found that tannins reduce food intake by locusts at low protein:carbohydrate ratios, but not at high ratios, a behavior which mirrored the conditional impact of tannins on locust performance (33, 34). Modulation of prior exposure to secondary metabolites on subsequent food choice was observed for both insect herbivores (35) and mammals (36, 37). Conversely, sugars were found to mask the aversive taste of secondary metabolites (38). Despite these advances, we currently lack a detailed

understanding of how primary and secondary metabolites interact to determine herbivore behavior

under biologically realistic conditions (9). The paucity of manipulative experiments that test for

interactions between host-derived chemical foraging cues *in planta* limits our capacity to assess

the concerted impact of different metabolites on herbivore feeding preferences, and, more

generally, our understanding of the role of plant metabolic complexity in herbivore behavior and

plant-herbivore interactions.

An implicit assumption of herbivore foraging theory is that herbivore behavior improves herbivore

fitness (39). With some notable exceptions, herbivores generally prefer to oviposit and feed on

plants and tissues that increase their performance (40–42). Host plant chemicals likely play an

important role in these preference-performance relationships (12, 17). However, the contribution

of individual behavioral cues to herbivore fitness in the context of the full chemical complexity of

a given host plant has remained difficult to quantify (30, 43, 44). Targeted molecular manipulation

of the production and perception of chemical cues provides new opportunities in this context,

including the evaluation of the benefits of the integration of multiple chemical cues into herbivore

foraging (29), and the assessment of the importance of the flexible use of multiple foraging cues

with redundant information content (11, 12).

Insect herbivores use chemosensory receptors to detect volatile and non-volatile plant chemicals

(45, 46). Gustatory receptors (GRs) are required for the detection of a variety of chemicals (47–

49), including non-volatile plant chemicals such as alkaloids (24) and sugars (50). Knocking out

specific GRs reduces oviposition of swallowtail (*Papilio xuthus*) butterflies (24) and host

recognition by silkworm (*Bombyx mori*) larvae (23), thus demonstrating the functional importance

of individual GRs for herbivore behavior. Work in Drosophila (*Drosophila melanogaster*) revealed

that some GRs are broadly tuned and can mediate avoidance to many different compounds (51),

while others are narrowly tuned and confer responsiveness to specific compounds (52). Gr43a is a

highly conserved insect taste receptor that specifically responds to D-fructose in Drosophila (49),

the diamondback moth (*Plutella xylostella*) (53) and the cotton bollworm (*Helicoverpa armigera*)

(54). As Gr43a is responsible for the detection D-fructose in the hemolymph as a proxy for

carbohydrate supply, *Gr43a* silenced flies also become unresponsive to other dietary sugars (55).

Interestingly, different GRs can interact dynamically through competition, inhibition and activation

(47, 56, 57), resulting in substantial potential for GR-mediated integration of multiple chemical

cues into behavioral responses.

Here, we developed a manipulative approach to evaluate the importance of maize primary and
secondary metabolites for the foraging and foraging-dependent performance of the western corn
rootworm (*Diabrotica virgifera virgifera*). The western corn rootworm is an economically
damaging maize pest (58). Its larvae are highly specialized on maize roots (59). Over the last
decades, the chemical ecology of the western corn rootworm has been studied in detail (13, 58, 59).
Field and laboratory studies revealed tight associations between maize root chemistry and western
corn rootworm behavior. Western corn rootworm larvae respond behaviorally to a wide variety of
chemical cues, including CO_2 (60), sugars and fatty acids (61, 62), aromatic and terpene volatile
organic compounds (63, 64), conjugated phenolic acids (65) and benzoxazinoids (66). The
emerging picture is that these chemicals likely allow western corn rootworm larvae to locate plant
roots from a distance (67), to discriminate between plants of different quality (63–65, 68), and to
identify and feed on the most nutritious roots (66, 69). Several *in vitro* experiments also suggest
that the western corn rootworm can integrate multiple chemical cues for host finding and
acceptance (61), hinting at the substantial sensory capacity of this specialist root feeder. By
independently manipulating the availability of benzoxazinoids and sugars as foraging cues of the
western corn rootworm through plant genetics and insect RNA interference (RNAi), we
demonstrate that sugars and benzoxazinoids serve both specific and integrated roles as
determinants of the behavior and behaviorally driven performance of this specialist herbivore.

Results
The western corn rootworm prefers to feed on root tissues that are rich in benzoxazinoids and
soluble sugars
The root system of young maize plants consists of embryonic and post-embryonic roots (Fig. 1*A*).
Embryonic roots emerge directly from the embryo and comprise primary and seminal roots. Post-
embryonic roots emerge from the hypocotyl and stem and comprise crown roots, and, at later
developmental stages of the plant, internode-derived brace roots (70). To determine feeding
preferences of the western corn rootworm within the root system of young maize plants, we infested
soil-grown maize plants with western corn rootworm larvae for 7 days and then scored the damage
on the different root types. In line with earlier studies (66, 69), we found low amounts of damage
on embryonic roots and substantial damage on post-embryonic roots (Fig. 1*B*). While embryonic
roots showed scattered bite marks, post-embryonic roots were often partially or even fully removed
(Fig. S1). To test whether this feeding preference is reflected in the distribution of the larvae within
the root system, we carried out a series of behavioral experiments. First, we laid out intact maize

root systems on a filter paper and then recorded the position of second instar larvae at different

time points after their release. Five times more larvae were found on post-embryonic than

embryonic roots 30 min after the release of the larvae (Fig. 1*C*). This preference persisted over the

duration of the experiment. To test whether the preference of the larvae for post-embryonic roots

may be due to differences in abundance or the relative position of the two root types within the root

system (Fig. 1*A*), we offered post-embryonic and embryonic root pieces of equal size to the larvae.

Similar to what was observed for entire root systems, significantly more larvae were found on the

post-embryonic root pieces than the embryonic root pieces after 30 min (Fig. 1*D*). Thus, the

western corn rootworm preferentially stays and feeds on post-embryonic roots of young maize

plants.

To gain insights into the phytochemical differences that may drive the preference of the western

corn rootworm for post-embryonic roots, we performed untargeted metabolomics on methanolic

extracts of both root types. Using UHPLC-Q-TOF-MS and ESI-, 4512 mass features were detected,

358 and 250 of which were enriched in post-embryonic or embryonic roots based on more than

two-fold difference in normalized signal intensity at a statistical significance of $p<0.05$ (Fig. 1*E*).

Using exact masses and fragmentation patterns, 28 metabolites could be putatively identified

within the dataset. Identified metabolites included benzoxazinoids, phenolic acid derivatives,

amino acids and sugars (Fig. 1*F*, table S1). Seventeen of these metabolites showed root-type

specific accumulation patterns (Fig. 1*F*). Among the secondary metabolites, benzoxazinoids and

phenolic acids showed pronounced shifts in their profiles, with some metabolites accumulating in

higher amounts in post-embryonic roots and others accumulating in higher amounts in embryonic

roots (Fig. 1*F*). Among the detected primary metabolite features, amino acids and sugars were

more abundant in post-embryonic- than in embryonic roots, with the most pronounced shifts

observed for features matching leucin and monosaccharides (Fig. 1*F*). Thus, the preference of the

western corn rootworm for post-embryonic roots is associated with distinct primary and secondary

metabolite accumulation patterns in these tissues.

Fig. 1. *The western corn rootworm prefers to feed on root tissues that are rich in benzoxazinoids and soluble sugars.*
(*A*) Left: young maize plants produce embryonic primary and seminal roots (orange arrows) and post-embryonic crown roots (blue arrows). Right: Western corn rootworm larvae are highly specialized maize root feeders that are very mobile in the second and third instar. A third instar larvae is shown. (*B*) Average damage score observed on post-embryonic and embryonic roots of soil-grown maize plants after 7 days of infestation by western corn rootworm larvae (***$p<0.001$, Wilcoxon Signed Rank test, n=20 plants with 15 larvae each, data from Fig. 4*D*). Dots represent damage scores on individual plants (averaged within root types). For frequency distributions of individual roots, refer to Fig. S1. (*C*) Preference of western corn rootworm larvae for post-embryonic and embryonic roots within the root system

(**p<0.01, ***p<0.001, FDR-corrected Least Square Mean post hoc tests, n=16 dishes with 6 larvae each). (*D*)
Preference of western corn rootworm larvae for root pieces of equal size of post-embryonic and embryonic roots
(***p<0.001; FDR-corrected Least Square Mean post hoc tests, n=18 dishes with 5 larvae each). (*E*) Metabolomics
profiles of methanolic extracts of post-embryonic and embryonic roots. Orange features are more abundant in
embryonic roots, blue features are more abundant in post-embryonic roots (min. 2-fold difference, p<0.05, FDR-
corrected Student's *T*-tests, n=9-10). Numbers denote features that were tentatively assigned to structures based on
exact mass and fragment information (Table S2). (*F*) Relative abundance differences of tentatively identified
metabolites between post-embryonic and embryonic roots (fold change of peak areas; *p<0.05, FDR-corrected
Student's *T*-tests, n=9-10). Picture credits: Ricardo Machado, Lingfei Hu, Cyril Hertz. Error bars denote standard
errors of means (SEM).

To narrow down the list of metabolites that may prompt the western corn rootworm to feed on post-

embryonic roots, we performed a literature search on western corn rootworm attractants and

feeding stimulants. Thirteen different compounds known to elicit behavioral responses in the

western corn rootworm *in vitro* were found (Table S2). Cross-referencing this table with the list of

chemicals that accumulated in post-embryonic roots at higher concentrations resulted in the

identification of four candidate compounds that may mediate post-embryonic root preference: The

monosaccharides glucose and fructose, which act as feeding stimulants (61) and the

benzoxazinoids DIMBOA-Glc and DIMBOA, which form iron complexes in the rhizosphere that

act as short-distance host preference and acceptance cues (66). Based on these results and earlier

studies (66, 69), we hypothesized that benzoxazinoids and monosaccharides may interact to

determine the preference of the western corn rootworm for post-embryonic roots. Given the

observed overlap between post-embryonic root preference and overall plant attractiveness (66), we

also hypothesized that the same compounds may mediate general host recognition and acceptance

in this specialist herbivore.

Independent manipulation of benzoxazinoids and sugars as foraging cues

To test for the individual and combined roles of benzoxazinoids and sugars in mediating western

corn rootworm behavior, we sought to independently manipulate their availability as foraging cues.

The first dedicated step in benzoxazinoid biosynthesis is mediated by the indole-3-glycerol

phosphate lyase *Bx1* (71). By consequence, *bx1* mutants are benzoxazinoid deficient (71, 72). To

test if *bx1* mutant plants can be used to manipulate benzoxazinoids independently of root sugars,

we performed a series of targeted analyses of wild type (WT) B73 and *bx1* roots. In accordance

with our metabolomics screen and earlier studies (66, 69), post-embryonic roots of WT plants

contained 4-fold higher DIMBOA levels and 2- to 3-fold higher DIMBOA-Glc and DIM_2BOA-

Glc levels, but lower HDMBOA-Glc levels than embryonic roots (Fig. 2*A*). Benzoxazinoid levels

in the *bx1* mutant were reduced by more than 97%, and residual benzoxazinoid concentrations were

not significantly different between crown and primary roots (Fig. 2*A*). As expected from the

metabolomic analyses, post-embryonic roots of WT plants contained higher levels of the

monosaccharides glucose and fructose than embryonic roots (Fig 2*A*). Sucrose levels were also

higher in post-embryonic roots (Fig 2*A*). Sugar profiles were similar in WT and *bx1* roots (Fig.

2*A*), demonstrating that benzoxazinoid availability can be manipulated independently of sugar

availability. The differential abundance between and crown and primary roots of the other 17

identified metabolic features was conserved in the *bx1* mutant as well (Fig. S2). We only detected

a significant interaction between root type and genotype for tryptophan, which is also derived from

indole-3-glycerol phosphate (71). Tryptophan accumulated in higher amounts in post-embryonic

roots of both wild type and *bx1* mutant plants, and this difference was more pronounced in the *bx1*

mutant (Fig. S2). Thus, we conclude that apart from benzoxazinoids, the metabolic differences

between crown and primary roots are conserved in the *bx1* mutant, thus validating the use of this

genotype to assess the behavioral impact of benzoxazinoids on western corn rootworm behavior.

Fig. 2. *A mutation in the Bx1-gene suppresses root-type specific benzoxazinoid accumulation independently of root*
sugars. (*A*) Concentrations of benzoxazinoids in embryonic and post-embryonic roots of wild type B73 and *bx1* mutant
plants (n=11-18). (B) Concentrations of glucose, fructose and sucrose in embryonic and post-embryonic roots of wild
type B73 and *bx1* mutant plants (n=8). Letters indicate significant differences in total amounts between root types and
genotypes (p<0.05, Holm-Sidak post hoc tests). Asterisks indicate significant differences in the concentrations of
individual compounds between post-embryonic and embryonic roots (p<0.05, Holm-Sidak post hoc tests). Error bars
denote standard errors of means (SEM).

To manipulate the availability of sugars as foraging cues independently of benzoxazinoids, we
targeted sugar perception in the western corn rootworm. This approach was chosen because
manipulation of primary metabolism in plants results in pleiotropic effects (73, 74). Sugar
perception in insects is highly conserved and mediated by *Gr43a*-like genes (49, 50, 53, 54, 75–
78). BLAST search of publicly available nucleotide sequences revealed a single putative *Gr43a*
gene in the western corn rootworm, here named *DvvGr43a* (Fig. 3*A*). The full-length sequence of
this gene can be retrieved from NCBI [Accession: XM_028279548.1]. Similarity between
DvvGr43a and putative *Gr43a* genes from different insects, many of which have been functionally
characterized, was found to be between 42% and 64% at the protein level. No other closely related
genes were found in western corn rootworm genome. Structural prediction indicated that the
DvvGr43a gene encodes a 7-transmemrane domain protein, consistent with its putative role as a
gustatory receptor (Fig. 3*B*). Gene-expression profiling revealed stronger expression of *DvvGr43a*
in the head than the body of western corn rootworm larvae (Fig. 3C). To manipulate *DvvGr43a*,
we targeted its expression through RNA interference (RNAi) by feeding the larvae with a 240 bp
dsRNA targeting *DvvGr43a*. Compared to control larvae that were fed with dsRNA targeting green
fluorescent protein (GFP), this treatment resulted in a >70% reduction in *DvvGr43a* transcript
abundance (Fig. 3*D*). To assess silencing specificity, we measured the transcription of the putative
CO_2-receptor *DvvGr2* (67). Feeding dsRNA targeting *DvvGr43a* did not change the transcript
abundance of *DvvGr2* (Fig. 3*E*). Accordingly, the responsiveness of *DvvGr43a*-silenced larvae to
volatile CO_2 remained intact (Fig. S3). To evaluate whether *DvvGr43a* silencing changes the ability
of western corn rootworm to respond to benzoxazinoids, we conducted choice assays with the
behaviorally active benzoxazinoid iron complex $Fe(III)(DIMBOA)_3$ (66). After 1 hour, the
majority of western corn rootworm larvae were found on filter paper discs spiked with
$Fe(III)(DIMBOA)_3$, independently of *DvvGr43a* silencing. Next, we tested whether *DvvGr43a*
silencing changes the ability of the larvae to respond to sugars. Three hours after the start of the
experiment, control larvae preferred to locate on filter discs spiked with glucose, fructose, sucrose
or a mixture of the three compounds (Fig. 3*G-J*). These preferences were absent in larvae fed with
DvvGr43a dsRNA (Fig. 3 *G-J*), demonstrating that silencing *DvvGr43a* eliminates the capacity of
western corn rootworm larvae to respond behaviorally to three major plant sugars. *DvvGr43a*-
dependent sugar preference was similar in non-starved and starved western corn rootworm larvae
(Fig. S4). Thus, combining *bx1* plant mutants and *DvvGr43a*-silenced larvae can be used to
evaluate the individual and combined impact of benzoxazinoids and sugars on herbivore foraging.

Fig. 3. *DvvGr43a mediates sugar preference of the western corn rootworm without influencing responsiveness to benzoxazinoids.* (*A*) Phylogenetic relationships between gustatory sugar receptors of different insects and *DvvGr43a*

of the western corn rootworm. The tree is based on protein sequences and drawn to scale, with branch lengths measured
in the number of substitutions per site. Asterisks indicate functionally characterized receptors. (*B*) Protein tertiary
structure of *DvvGr43a* as predicted with the Phyre2 algorithm. (*C*) *DvvGr43a* expression in the bodies (thorax and
abdomen) or heads of second instar western corn rootworm larvae (*p<0.05, Student's *T*-test n=10). (*D*) *DvvGr43a*
expression in western corn rootworm larvae fed with dsRNA targeting green fluorescent protein (control, GFP) or
DvvGr43a (Gr43a, ***p<0.001, Student's *T*-test, n=11). (*E*) *DvvGr2* expression in western corn rootworm larvae fed
with dsRNA targeting green fluorescent protein (control, GFP) or *DvvGr43a* (n=8). (*F*) Preference of GFP and
DvvGr43a dsRNA fed larvae for buffer or $Fe(III)(DIMBOA)_3$ on filter discs after 1 h (***p<0.001, FDR-corrected
Least Square Mean post hoc tests, n=10 dishes with 6 larvae each). (*G-J*) Preference of GFP or *DvvGr43a* dsRNA fed
larvae for buffer or glucose, fructose, sucrose or a mixture of the three on filter discs at different time points (*p<0.05;
**p<0.01; ***p<0.001, FDR-corrected Least Square Mean post hoc tests, n=15 dishes with 6 larvae each). Error bars
denote standard errors of means (SEM).

Benzoxazinoids and sugars play distinct roles in root herbivore foraging

To understand the importance of benzoxazinoids and sugars in mediating host and tissue finding

of the western corn rootworm in a neutral environment, we observed the behavior and feeding of

control and *DvvGr43a*-silenced larvae on wild type and *bx1* mutant root systems laid out on moist

filter paper. Ninety % of control larvae were found on the roots rather than the filter paper 4 hours

after their release (Fig. 4*A*). This number dropped to 70% for *DvvGr43a* silenced larvae,

independently of plant genotype. A different pattern was found when looking at the distribution of

the larvae on the different root types within root systems (Fig. 4*B*). On wild type plants,

significantly more control larvae were found on post-embryonic than embryonic roots. This

difference was absent for control larvae feeding on *bx1* mutant plants. For *DvvGr43a* silenced

larvae, more larvae were found on post-embryonic than embryonic roots, even though this

preference was attenuated relative to control larvae. No preference was found for *DvvGr43a*

silenced larvae feeding on *bx1* mutant roots (Fig. 4*B*). These results show that sugars contribute to

root system finding and post-embryonic root preference but are not strictly required for either

process. Benzoxazinoids on the other hand do not contribute to root system finding but are

specifically required for the capacity of the western corn rootworm to distinguish post-embryonic

from embryonic roots.

To evaluate whether benzoxazinoids and sugars affect host acceptance in the form of active

feeding, we infested roots of soil-grown wild type and *bx1* mutant plants with control or *DvvGr43a*-

silenced western corn rootworm larvae and inspected the roots for characteristic feeding marks and

damage after 7 days. As feeding by the western corn rootworm cannot be observed directly in the

soil, and root biomass is a poor proxy for root consumption due to confounding root regrowth

effects and the disproportionate impact of severed roots (79), this approach was considered most

suitable for the question at hand. The experiment revealed that root damage across different roots

was increased by the availability of benzoxazinoids and sugars as foraging cues in an additive

fashion (Fig. 4*C*). Frequent root removal was observed in wild type plants infested with control

larvae. Intermediate root damage in the form of frequent biting marks and brown discoloration was

observed for *bx1* mutant roots infested with control larvae and wild type roots infested with

DvvGr43a-silenced larvae. Infrequent biting marks and/or no visible damage was observed for *bx1*

mutant roots infested with *DvvGr43a*-silenced larvae. The same additive pattern was also visible

when looking at the distribution of damage patterns between the different root types (Fig. 4*D*).

Post-embryonic root damage was strongest and significantly more pronounced than embryonic root

damage in wild type plants infested with control larvae. Post-embryonic root damage and damage

differences with embryonic roots were intermediate for *bx1* mutants infested with control larvae

and wild type plants infested with *DvvGr43a*-silenced larvae. Post-embryonic root damage was

lower and no longer different from damage to embryonic roots for *bx1* mutants infested with

DvvGr43a-silenced larvae (Fig. 4*D*). Thus, both benzoxazinoids and sugars increase root damage

in an additive manner, and this pattern is driven by post-embryonic root damage. Nevertheless, bite

marks were still observed on 69% of *bx1* mutant roots infested with *DvvGr43a*-silenced larvae,

showing that the larvae attack maize roots even in the absence of these cues. Taken together, these

results reveal distinct and additive roles of benzoxazinoids and sugars in root and tissue location

and feeding preferences of the western corn rootworm.

Fig. 4. *Benzoxazinoids and sugars play distinct roles in root herbivore foraging in vivo*. (*A*) Proportion of control (GFP) or *DvvGr43a*-silenced (Gr43a) larvae found on the roots of wild type B73 or *bx1* mutant roots 4 hours after their release. Letters indicate statistically significant differences between treatments (Holm-Sidak post-hoc tests, $p < 0.05$, n=15 dishes with 6 larvae each). (*B*) Proportion of control (GFP) or *DvvGr43a*-silenced (Gr43a) larvae found on embryonic (E) and post-embryonic (PE) roots within the same experiment. Asterisks indicate significant differences between root types (***$p < 0.001$, FDR-corrected Least Square Mean post hoc tests, n=15 dishes with 6 larvae each). (*C*) Average feeding damage by control (GFP) or *DvvGr43a*-silenced larvae per root of soil-grown wild type B73 or *bx1* mutant plants 7 days after infestation. Letters indicate statistically significant differences ($p < 0.05$, Tukey's post hoc tests, n=20 plants with 15 larvae each). Bubble plots are shown for illustrative purposes. The sizes of the circles are proportional to the relative frequency (% within each treatment) of the different types of observed damage (each column sums up to 100%). For damage scale, refer to Fig. 1. (*D*) Average feeding damage on embryonic (E) or post-embryonic (PE) roots within the same experiment. Asterisks indicate significant differences in root damage between root types (*$p < 0.05$, ***$p < 0.001$ Wilcoxon Signed Rank tests, n=20 plants with 15 larvae each). Bubble plots are shown for illustrative purposes. The sizes of the circles are proportional to the relative frequency (% within each root type) of the different types of observed damage (each column sums up to 100%). Error bars denote standard errors of means (SEM).

Using benzoxazinoids and sugars as foraging cues improves herbivore growth and survival

Herbivore foraging theory predicts that herbivores use plant chemical cues to make fitness-relevant

foraging decisions. To test whether the integration of primary and secondary metabolites into

foraging improves herbivore performance and survival, we first evaluated whether the absence of

benzoxazinoids or the inability to detect sugars may have direct performance effects on the western corn rootworm (as opposed to the effects mediated by changes in foraging behavior). Benzoxazinoids improve western corn rootworm performance under iron-limiting conditions (66) and upon attack by entomopathogenic nematodes (80, 81). When bioavailable iron is abundant and natural enemies are absent, larval performance under no-choice conditions is not altered by benzoxazinoids (66, 69). Thus, growing maize plants with bioavailable EDTA-Fe as iron source allowed us to exclude direct effects of benzoxazinoids on larval performance. To test whether silencing *DvvGr43a* has direct negative effects on western corn rootworm performance, we reared control and *DvvGr43a*-silenced larvae on young maize seedlings that produce embryonic, but no post-embryonic roots. Larval weight gain and mortality of control and *DvvGr43a*-silenced larvae after 6 days was similar (Fig. S5), demonstrating that silencing *DvvGr43a* has no direct effects on larval performance in a no-choice situation. Note that larval mortality in this experiment was >40%, which is within the normal range for early instars of this species (82, 83).

Having established the validity of our experimental approach, we proceeded to evaluate the impact of benzoxazinoid- and sugar-dependent foraging behavior on western corn rootworm performance. Highest larval weight and lowest mortality was observed for control larvae feeding on wild type plants (Fig. 5*A-B*). Intermediate weight gain and mortality was observed for control larvae feeding on *bx1* mutant roots and for *DvvGr43a*-silenced larvae feeding on wild type roots. Lowest weight gain and highest mortality was observed for *DvvGr43a*-silenced larvae feeding on *bx1* mutant roots. Larval performance thus mirrored root damage patterns (Fig. 4*C-D*). This was confirmed by correlation analysis which revealed a positive relationship between total larval weight and survival on a given plant and the total damage observed on post-embryonic roots (Fig. 5*C-F*). In conclusion, sugars and benzoxazinoids act together to improve western corn rootworm performance by acting as foraging cues that guide the herbivore to feed on the most profitable host tissue.

Fig. 5. *Using benzoxazinoids and sugars as foraging cues improves herbivore growth and survival.* (*A*) Larval weight
of control (GFP) and DvvGr43a-silenced dsRNA fed western corn rootworm larvae feeding on wild type B73 or *bx1*
mutant plants for 7 days. Note that in a no-choice situation, neither the *bx1* mutation nor DvvGr43a silencing reduce
larval performance ((69) and Fig. S5). Letters indicate significant differences (p<0.05, Holm-Sidak post hoc tests,
n=20 pots with 15 larvae each). (*B*) Larval mortality within the same experiment. Letters indicate significant
differences (p<0.05, Holm-Sidak post hoc tests, n=20 pots with 15 larvae each). (*C-F*) Correlations between
cumulative damage per plant and larval performance parameters for post-embryonic roots (*C-D*) and embryonic roots
(*E-F*). Linear regressions are shown for significant correlations (p<0.05). *P*-values are shown for Spearman Rank
Order correlations.

Discussion

Plant primary and secondary metabolites can act together to determine the behavior of generalist herbivores on artificial diet (33, 34). The present work expands these findings by demonstrating that primary and secondary metabolites can have distinct, overlapping, and additive roles as determinants of the behavior, and, consequently, the performance of a specialist herbivore *in planta*.

Herbivores are thought to use primary and secondary metabolites to make foraging decisions within complex host metabolomes (16), but how they combine these two types of cues during different stages of host finding and acceptance is not well understood. We find that the western corn rootworm employs sugars, but not benzoxazinoids for host finding. Sugars increase the successful localization of post-embryonic roots, but only in the presence of benzoxazinoids, which are essential for the latter. In contrast to these distinct roles, sugars and benzoxazinoids have similar roles in enhancing overall root damage and tissue-specific post-embryonic root damage, and act in an additive manner in this context. Thus, western corn rootworm larvae employ primary and secondary metabolites in varying combinations and hierarchies for different behavioral patterns. This finding differs from current hypotheses and observations for polyphagous insects that suggest sequential use of different foraging cues (84, 85). The explanation for this discrepancy is likely found in the distribution patterns of plant primary and secondary metabolites in maize roots and rhizosphere. Benzoxazinoids and sugars constantly accumulate in maize roots, with stable concentration differences between post-embryonic and embryonic roots (66, 69). Both compound classes are also exuded into the rhizosphere (86, 87). Current work using enzyme-inhibiting and non-inhibiting extraction solvents (66, 69, 86) suggests that benzoxazinoids are likely exuded as glucosides before being deglycosylated at the root surface, resulting in the release of aglucones and glucose. Aglucones such as DIMBOA then form attractive complexes with free iron. (66). Thus, both sugars and benzoxazinoids are present inside and outside of the roots, with outside accumulation likely being more dynamic than inside accumulation. Taking into account that benzoxazinoids are plant-specific, while sugars are more general cues (88, 89), one can envisage a scenario where the western corn rootworm relies on sugars as general short-distance cues for the presence of roots (with potentially high concentrations on the surface of maize roots due to benzoxazinoid deglycosylation), followed by benzoxazinoids as specific within-plant tissue selection cues. Benzoxazinoids may also be used as specific short-distance host recognition cues in case roots of multiple plant species are intermingled. Following host and tissue finding, the

herbivore would then use both sugars and benzoxazinoids as stable host acceptance and feeding
cues. High resolution metabolite imaging (90) together with the manipulation of gustatory
receptors (23) could help to test these hypotheses and explore the connections between the
distribution dynamics of different plant metabolites and herbivore foraging.

Plants produce complex chemical mixtures, and herbivores likely combine many of these
chemicals to make appropriate foraging decisions (11, 12, 29). In line with this idea, recent *in vivo*
work shows that, while individual compound classes and insect gustatory receptors can be of major
importance for specific behavioral patterns, their disruption is not sufficient to fully suppress these
behaviors (22–24). In the case of the western corn rootworm, we find that even the suppression of
two major foraging cues does not fully abolish its ability to find and accept a host plant, at least
under laboratory conditions. Several other plant-derived chemicals, including different volatiles
and fatty acids, are known to attract, arrest and stimulate feeding by the western corn rootworm
(61, 62, 64). These cues are likely to enable basic behavioral patterns in the absence of
benzoxazinoids. We conclude that the flexible integration of complex blends of chemical cues
provides partial resistance to chemically mediated behavioral disruption. Recent work highlights
the importance of learning in herbivore foraging (12, 91), which likely increases the capacity of
herbivores to respond to dynamic chemical landscapes and to substitute between different cues
with redundant information content. How previous experiences influence the foraging behavior of
the western corn rootworm remains to be determined. All larvae used in our experiments were
reared on maize, reflecting their high degree of specialization on this host plant (59). Western corn
rootworm adults are more flexible with their diet (59, 92), and it would be interesting to understand
how their experience influences the responses of the next larval generation to host plant cues.

The capacity of an herbivore to select a host plant and feed on specific tissues generally improves
its fitness (40, 42, 93). In line with these preference-performance relationships, diet experiments
often document strong associations between chemically-mediated herbivore feeding preferences
and herbivore performance (17). In many of these cases, herbivore feeding is directly elicited by
substances that improve performance (17). However, plant metabolomes are likely to complex to
be fully assessed by an herbivore, who thus need to rely on digestive feedbacks or subsets of
chemical cues to make decisions. To what extent the use of plant chemical cues enhances herbivore
performance by guiding herbivores to nutritious tissues is poorly understood, and interactions
between primary and secondary metabolite chemical cues have, to the best of our knowledge, not

been tested *in planta*. Our experimental approach rules out direct effects of benzoxazinoids and sugar perception on western corn rootworm performance (66, 69) and thus allows us to quantify their net impact as foraging cues. We find that the availability of sugars and benzoxazinoids during foraging improves herbivore weight gain and survival individually by 8%, and in combination by 15%, benefits which are most likely driven by increased post-embryonic root feeding. Post-embryonic roots of young maize plants are growing more vigorously than embryonic roots, receive a higher share of photosynthates and are richer in carbohydrates, amino acids and total soluble protein and increase western corn rootworm growth in no-choice experiments (69). In combination, these findings demonstrate that the combined use of primary and secondary metabolites as foraging cues enhance the performance of a specialist insect herbivore by guiding it to its most profitable feeding niche.

The gustatory receptor Gr43a is a highly conserved insect sugar receptor that is narrowly tuned to D-fructose (49), but mediates feeding preferences for different sugars through dietary feedbacks (55). Given the conserved function, predicted protein structure, tissue-specific expression pattern and effect on sugar preferences, *DvvGr43a* is very likely to act as a sugar sensor in the western corn rootworm as well. An interesting observation in this context is that *DvvGr43a*-dependent larval feeding preferences for sugars and sugar mixtures were only established after 2-3 hours, while feeding preferences for Fe(III)(DIMBOA)$_3$ were established more rapidly (66). In Drosophila, Gr43a increases sugar intake under food limiting conditions and suppresses sugar intake under satiation (49). We thus hypothesized that the delayed *DvvGr43a*-dependent sugar feeding preference of western corn rootworm larvae for sugars may be the result of a starvation response. However, we observed a similar lag time in sugar preference in pre-starved larvae. This suggests that *DvvGr43a* likely functions as a sugar sensor rather than as a nutrient uptake regulator in the western corn rootworm. The observed lag phase may be explained by the potential role of *DvvGr43a* as an internal sugar sensor that triggers behavioral responses once sugars are taken up and elicit changes in hemolymph sugar levels. A deeper molecular characterization could shed light on the precise neurological and physiological roles of *DvvGr43a* in the western corn rootworm in the future.

In conclusion, the experiments presented here reveal how primary and secondary metabolites interact to guide the foraging of a specialist herbivore. The distinct individual and combined roles of these metabolites reveal finely tuned herbivore foraging strategies that are directly linked to

improved herbivore performance and survival. At the same time, foraging is remarkably robust to
perturbation, most likely due to the flexible use of redundant host plant cues. Our mechanistic
approach thus offers a glimpse into the complex, multilayered interplay between plant
metabolomes and chemically guided herbivore foraging.

Materials and Methods

<u>Plants and insects</u>

Maize seeds (*Zea mays* L., inbred line B73) were provided by Delley Semences et Plantes SA
(Delley, CH). The near-isogenic benzoxazinoid deficient *bx1* mutant line in a B73 background was
obtained by backcrossing the original *bx1* mutant five times into B73 (94). Seedlings were grown
under greenhouse conditions (23 ± 2˚C, 60% relative humidity, 16:8 h L/D, and 250 mmol/m^2/s^1
additional light supplied by sodium lamps). Plantaaktiv$^®$ 16+6+26 Typ K fertilizer (Hauert HBG
Dünger AG, Grossaffoltern, Switzerland) was added twice a week after plant emergence following
the manufacturer's recommendations. When plants were used to feed insects, seedlings were
germinated in vermiculite (particle size: 2-4 mm; tabaksamen, Switzerland) and used within four
days after germination. Eggs of the non-diapausing *Diabrotica virgifera virgifera* strain were
originally supplied by USDA-ARS-NCARL, Brookings, SD. Insect colonies were subsequently
established and are maintained in the University of Bern and the University of Neuchatel. Upon
egg hatching, insects were maintained in organic soil (Selmaterra, Bigler Samen AG, Thun,
Switzerland) and fed freshly germinated maize seedlings.

<u>Herbivore behavior</u>

To evaluate the preference of western corn rootworm larvae for different plant genotypes and root
types, we use three different experimental setups: potted plants, root systems laid out in petri dishes
and detached roots in petri dishes.

For western corn rootworm feeding preference experiments using whole plants, three-week-old
maize plants grown in 200 mL cylindrical plastic pots filled with sand and topped off with potting
soil were infested with fifteen second instar western corn rootworm larvae. Twenty plants per plant
genotype/type of larvae combination were evaluated (n=20). Seven days after infestation, plants
were gently excavated and inspected for root damage. Damage was recorded for each root
individually using the following scale. 0: no visual damage, 1: one-three bite marks, 2: more than
three bite marks, 3: several bite marks and brown discoloration, 4: root partially removed, and 5:
root fully removed.

Western corn rootworm feeding preference assays using petri dishes were conducted as described

(95, 96). Briefly, two-week-old maize plants grown as described above were gently excavated. The

root systems were washed and laid out onto a moist filter paper embedded in a petri dish (14 cm

diameter, Greiner Bio-one, Austria). A circular whole in the rim of the dishes accommodated the

plant stems, with the shoots remaining outside of the dishes. Five third instar western corn

rootworm larvae were released into each petri dish. Petri dishes were sealed and then covered with

aluminum foil. The number of western corn rootworm larvae feeding on each root type was

determined at different time points after their release.

To evaluate western corn rootworm feeding preference using detached roots, pieces of equal sizes

of two post-embryonic roots and two embryonic roots from two-week-old maize plants were placed

next to each other onto a moist filter paper (diameter 90 mm, GE Healthcare, UK) embedded into

a petri plate (90 mm diameter, Greiner Bio-one, Austria). Five third instar western corn rootworm

larvae were released into each petri dish. Petri plates were sealed and then covered with aluminum

foil. The number of western corn rootworm larvae feeding on each root type was determined at

different time points.

To evaluate the attractiveness of CO_2 for western corn rootworm larvae, we conducted choice

experiments using belowground olfactometers (64). CO_2 levels were increased in one arm of the

olfactometer by delivering CO_2-enriched synthetic air (1% CO_2, Carbagas, Switzerland).

Olfactometer arms were closed on top using parafilm during CO_2 delivery. A manometer connected

to the synthetic air bottle allowed to fine-tune CO_2 delivery rates. CO_2 levels were measured using

a gas analyzer (Li7000, Li-Cor Inc., Lincoln, Nebraska, USA). Once desired CO_2 concentrations

were reached, CO_2 delivery was terminated, olfactometer arms were connected to olfactometer

central connectors and six second to third instar western corn rootworm larvae were release

immediately and their choice were evaluated within ten minutes of release. Six olfactometers per

larval type were assayed (n=6).

To evaluate the attractive effects of $Fe(III)(DIMBOA)_3$ for western corn rootworm larvae, we

release six second to third instar larvae in the middle of a moist sand-filled petri dish (9 cm

diameter, Greiner Bio-One GmbH, Frickenhausen, DE) where they encountered a filter paper disc

treated with 10 μl of $Fe(III)(DIMBOA)_3$ (1 μg/ml of water) or, on the opposite side, a filter paper

disc treated with water only. Ten independent petri dishes per larval type were evaluated (n=10).

$Fe(III)(DIMBOA)_3$ was prepared fresh by mixing $FeCl_3$ and DIMBOA at a 1:2 ratio as described

(96). Larval preferences were recorded 1 h after releasing the larvae.

To evaluate attractive effects of soluble sugars for western corn rootworm larvae, we followed the

procedure described by 97 with minor modifications. Briefly, we released six second to third instar

western corn rootworm larvae in the middle of a moist sand-filled petri dish (6 cm diameter, Greiner

Bio-One GmbH, Frickenhausen, DE) where they encountered a filter paper disc treated with 10 μl

of either glucose (30 mg/ml), fructose (30 mg/ml), sucrose (30 mg/ml), or a mixture of the three

(10 μl containing each sugar at a concentration of 30 mg/ml), or, on the opposite side, a filter paper

treated with water only. Fifteen independent petri dishes per treatment were assayed (n=15). Larval

preferences were recorded regularly for 3 h. In a second experiment, larvae were pre-fed on maize

roots or starved for 12 h before being released into petri dishes with filter paper discs treated with

a sugar mixture or water, as described above. Eight independent petri dishes with six larvae each

were assayed (n=8).

<u>Herbivore performance</u>

To evaluate the impact of benzoxazinoids and sugars on western corn rootworm performance,

three-week-old maize plants grown in 200 mL cylindric plastic pots (11 cm depth and 4 cm

diameter) filled with sand were infested with fifteen second instar western corn rootworm larvae

(n=20). Seven days after infestation, the larvae were collected, counted, and weighted using a

microbalance. Note that weight and survival was determined on the same plants as root damage

(see section "Herbivore behavior").

To evaluate the impact of *DvvGr43a*-silencing on western corn rootworm performance in a no-

choice setting, GFP and *DvvGr43a* fed first instar larvae were reared on 3-4-day old maize

seedlings, which only produce embryonic roots, in soil-filled plastic cups (n=40). Nine larvae were

used per cup. Fresh seedlings were provided to the larvae every three days for a total of six days,

after which the larvae were collected, counted, and weighted using a microbalance.

<u>Root metabolite measurements</u>

Untargeted root metabolomics was conducted as described (81). Embryonic and post-embryonic

roots of B73 plants were harvested separately, flash-frozen and ground in liquid nitrogen. Sixty to

eighty mg of plant tissues were then extracted in 800 μL of acidified H_2O/MeOH (50:50 v/v; 0.1%

formic acid). Resulting root extracts were analysed using an Acquity UHPLC system coupled to a

G2-XS QTOF mass spectrometer equipped with an electrospray source (Waters). Gradient elution

was performed on an Acquity BEH C18 column (2.1 × 50 mm i.d., 1.7 μm particle size) at 1-27.5%

B over 3.5 min, 27.5-100% B over 1 min, holding at 100% B for 1 min, and reconditioning at 1%

B for 1 min, where A =0.1% formic acid/water and B = 0.1% formic acid/acetonitrile. The flow

rate was 0.4 mL/min. The temperature of the column was maintained at 40°C, and the injection

volume was 1 μL. The QTOF MS was operated in negative mode. The data were acquired over an

m/z range of 100–1200 with scans of 0.15 s at collision energy of 4 V and 0.2 s with a collision

energy ramp from 10 to 40 V. The capillary and cone voltages were set to 2 kV and 20 V,

respectively. The source temperature was maintained at 140°C, the desolvation gas was 400°C at

1000 L h-1 and cone gas flow was 50 L/hr. Accurate mass measurements (<2 ppm) were obtained

by infusing a solution of leucin encephalin at 200 ng/mL at a flow rate of 10 μL/min through the

Lock Spray probe (Waters). The chromatograms were processed using Progenesis QI (Nonlinear

Dynamics, Newcastle UK) with default settings for spectral alignment, peak picking, and

deconvolution, resulting in a total of 4956 mass features. Of these, 444 mass features eluting

between 0 and 0.3 min were discarded, since their lack of retention and excessive coelution made

reliable identification impossible.

Abundances for each remaining mass feature were normalised by ArcSin transformation, and mean

normalized abundances for post-embryonic and embryonic roots were compared by Student's t-

test. P-values were adjusted by Benjamini–Hochberg false discovery rate multiple testing

correction and a fold-change cut-off of 2.0 was applied to generate a list of compounds that

significantly differed between treatments. Volcano plots were used to visualize treatment

differences, and the list of differentially expressed features was tested against an in-house mass-

fragmentation database using Progenesis QI (Progenesis MetaScope v1.0.6253.26544, Nonlinear

Dynamics, Newcastle, UK). In addition, chromatograms were manually scanned for exact masses

of known plant metabolites, specifically amino acids, sugars, organic acids, and flavonols. Potential

matches were likewise tested against the mass fragmentation database, and putative identifications

were accepted based on exact mass, mass fragmentation patterns, and relative retention times.

Root benzoxazinoid quantification was conducted as described (98). For this, embryonic and post-

embryonic roots of B73 or *bx1* plants were harvested separately, flash-frozen and ground in liquid

nitrogen. Approximately 100mg of plant tissues were then extracted in 1 mL of acidified

H_2O/MeOH (50:50 v/v; 0.1% formic acid), and analyzed with an Acquity UHPLC system coupled

to a UV detector and a Waters QDa mass spectrometer equipped with an electrospray

source(Waters). Compounds were separated on an Acquity BEH C18 column (2.1×50 mm i.d., 1.7

μm particle size). Water (0.1% formic acid) and acetonitrile (0.1% formic acid) were employed as

mobile phases A and B. The elution profile was: 0-1 min, 10% B; 1-4 min, 10-30% B; 4-5 min,

30- 9740% B; 5-6 min, 40-100% B; 6-8.5 min, 100% B; followed by 1.5 min reconditioning at

10% B. The mobile phase flow rate was 0.4 mL/min. The column temperature was maintained at

40°C, and the injection volume was 3 μL. The MS was operated in negative mode and data were

acquired in single ion recording (SIR) mode, using $[M-H]^-$ and $[M+HCOO^-]^-$ adducts of

benzoxazinoid compounds. All other MS parameters were left at their default values as suggested

by the manufacturer. Benzoxazinoid peaks were identified from MS signals and quantified from

integrated UV signals at 265nm. External standards for HMBOA, DIMBOA, DIMBOA-Glc,

HDMBOA-Glc, and MBOA were used for absolute quantification. Where concentrations were too

low to result in reliable UV signals, a compound-specific MS-to-UV signal conversion factor was

estimated to calculate 'predicted' UV signals from the more sensitive MS signals before

quantification.

Root sugar levels were measured by an enzymatic/spectrophotometric method as described (99).

Embryonic and post-embryonic roots of B73 and *bx1* plants were harvested separately, flash-frozen

and ground in liquid nitrogen. Soluble sugars were extracted from 100 mg root tissue using 80%

(v/v) ethanol, followed by an incubation step (10 min at 78°C) with constant shaking at 800 rpm.

Pellets were re-extracted twice with 50% (v/v) ethanol (10 min at 78°C with constant shaking at

800 rpm). Supernatants from all extraction steps were pooled together, and sucrose, glucose and

fructose were quantified as described (100).

<u>Identification of *DvvGr43a*</u>

To identify the *Gr43a* receptor orthologues in the western corn rootworm, we used the *Gr43a*

receptor-encoding gene sequence of *Drosophila melanogaster* as query against publicly available

nucleotide sequences and against the western corn rootworm genome sequences (NCBI accession:

PXJM00000000.2) using the National Center for Biotechnology Information Basic Local

Alignment Search Tool (NCBI BLAST). The full gene sequence can be retrieved from the National

Center for Biotechnology Information (NCBI) data bank using the following accession number:

XM_028279548.1. Then, we reconstructed evolutionary relationships between the identified

DvvGr43a gene and several other putative Gr43a genes that have been functionally characterized

as glucose, fructose, and/or sucrose receptors (49, 50, 53, 54, 75–78). The evolutionary

relationships were inferred using the Maximum Likelihood method based on the Le Gascuel 2008

model in MEGA7 (101, 102). The tree with the highest log likelihood (-14287.71) is shown. The

percentage of trees in which the associated taxa clustered together is shown next to the branches.

Initial tree(s) for the heuristic search were obtained automatically by applying Neighbor-Join and

BioNJ algorithms to a matrix of pairwise distances estimated using a JTT model, and then selecting

the topology with superior log likelihood value. A discrete Gamma distribution was used to model

evolutionary rate differences among sites (5 categories (+G, parameter = 1.0811)). The tree is

drawn to scale, with branch lengths measured in the number of substitutions per site. There was a

total of 569 positions in the final dataset. Graphical representation and edition of the phylogenetic

tree were performed with the Interactive Tree of Life (version 3.5.1) (103). Protein tertiary

structures and topologies were predicted using Phyre2 (104). The protein sequences used to

reconstruct evolutionary relationships can be retrieved from the NCBI under the following

accession numbers: *Heliothis virescens* Gr4 (CAD31946.1), *Danaus plexippus* Gr4 (EHJ77681.1),

Bombyx mori Gr9(NP_001124345.1), *Bombus impatiens* Gr43a (XP_003486787.1), *Nasonia*

vitripennis Gr3 (NP_001164386.1), *Apis mellifera* Gr3 (XP_016768876.1), *Drosophila*

melanogaster Gr43a (NP_523650.2), *Tribolium castaneum* Gr20 (EFA05758.1), *Anopheles*

gambiae Gr43a (XP_318100.4), *Camponotus floridanus* Gr43a (EFN61344.1), *Aedes aegypti*

Gr43a (XP_001658898.3), *Culex quinquefasciatus* Gr43a (XP_001842305.1), *Orussus abietinus*

Gr43a (XP_012270798.1), *Aethina tumida* Gr43a (XP_019876958.1), *Anoplophora glabripennis*

Gr43a (XP_018574194.1), *Helicoverpa armigera* Gr4 (AGK90011.1), *Helicoverpa armigera* Gr9

(JX970522.1), *Trichogramma chilonis* Gr43a (QAY30709.1), and *Diabrotica virgifera* Gr43a

(XP_028135349.1). For *Plutella xylostella* GR43a-1 and *Plutella xylostella* GR43a-2, and *D.*

melanogaster gustatory receptors refer to: (53, 105), as these sequences have not been deposited

in a public repository.

Double stranded RNA production

Double stranded RNA targeting *DvvGr43a* was produced by *Escherichia coli* HT115. To this end,

electrocompetent *E. coli* HT115 cells were transformed with recombinant L4440 plasmids that

contained either a 240 bp long *DvvGr43a* gene fragment, or a 240 bp long green fluorescent protein

(GFP) gene fragment flanked by two T7 promotors in opposite directions. Gene fragments were

synthetized *de novo* (Eurofins, Germany). To induce the production of dsRNA, an overnight

bacterial culture was used to inoculate fresh Luria-Berthani broth (25 g/L, Luria/Miller, Carl Roth

GmbH, Karlsruhe, Germany). Once the bacterial culture reached an OD_{600} of 0.6-0.8, it was

supplemented with isopropyl β-D-1-thiogalactopyranoside, IPTG) (Sigma-aldrich, Switzerland) at

a final concentration of 2mM. Bacterial cultures were incubated at 37°C in an orbital shaker at 130

rpm for 16h. Bacteria were harvested by centrifugation and stored at -20°C for further use (106).

Gene silencing experiments

To induce gene silencing in western corn rootworm, six to ten western corn rootworm larvae were

released in solo cups (30 mL, Frontier Scientific Services, Inc., Germany) containing approx. 2g

of autoclaved soil (Selmaterra, Bigler Samen AG, Thun, Switzerland) and 2-3 freshly germinated

maize seedling. Maize seedlings were coated with 1 ml of bacterial solution containing

approximately 200-500 ng of dsRNA targeting *DvvGr43a*. As controls, larvae were fed with

bacteria producing dsRNA targeting green fluorescent protein genes (GFP). Seedlings coated with

bacteria were provided every other day for three consecutive times. Two days after, larvae were

collected and used for experiments.

Gene expression measurements

Total RNA was isolated from approximately 10 mg of frozen, ground and homogenized western

corn rootworm larval tissue (3-7 larvae per biological replicate, n=8-11) using the GenElute

Universal Total RNA Purification Kit (Sigma-Aldrich, St. Louis, MO, USA). A NanoDrop

spectrophotometer (ND-1000, Thermo Fisher Scientific, Waltham, MA, USA) was used to

estimate RNA purity and quantity. DNase-treated RNA was used as template for reverse

transcription and first strand cDNA synthesis with PrimeScript Reverse Transcriptase (Takara Bio

Inc., Kusatsu, Japan). DNase treatment was carried out using the gDNA Eraser (Perfect Real Time)

following manufacturer's instructions (Takara Bio Inc.). For gene expression analysis, 2 μl of

undiluted cDNA (i.e. the equivalent of 100 ng total RNA) served as template in a 20 μl qRT-PCR

using the TB Green Premix Ex Taq II (Tli RNaseH Plus) kit (Takara Bio Inc.) and the Roche

LightCycler 96 system (Roche, Basel, Switzerland), according to manufacturer's instructions.

Transcript abundance of the following western corn rootworm genes were analysed: *DvvGr43a*

and *DvvGr2*. *Actin* was used as reference gene to normalize expression data across samples.

Relative gene expression levels were calculated by the $2^{-\Delta\Delta Ct}$ method (107).The following primers

were used: DvvGr43a-F GTCACATTCACCACGGGTCT, DvvGr43a-R

CGTTCGGTCTCTAACTTTGGC, DvvGr2-F GAACTAAGCGAGCTCCTCCA, DvvGr2-R

CAGAAGCACCATGCAATACG, DvvActin-F TCCAGGCTGTACTCTCCTTG, and DvvActin-

R CAAGTCCAAACGAAGGATTG.

<u>Statistical analyses</u>

Differences in root benzoxazinoid and root sugar levels as well as gene expression levels were

analyzed by One- and Two-Way Analyses of Variance (ANOVAs). Normality and equality of

variance were verified using Shapiro–Wilk and Levene's tests, respectively. Holm–Sidak post hoc

tests were used for multiple comparisons. Datasets from experiments that did not fulfill the

assumptions for ANOVA were natural log-, root square-, or rank-transformed before analysis.

Differences in damage scores between embryonic and post-embryonic roots were assessed using

Wilcoxon Signed Rank Tests. Differences in average root damage scores between treatments were

assessed using Kruskal-Wallis ANOVAs on ranks followed by Tukey's post hoc tests. The above

analyses were performed in Sigma Plot 12.0 (SystatSoftware Inc., San Jose, CA, USA) with the

support of the built-in Advisor Wizard function. Differences in larval preference were assessed

through Generalized Linear Models (GLM) with binomial distribution and corrected for

overdispersion with quasi-binomial function when necessary followed by analysis of variance

(ANOVA) and FDR-corrected Least Square Means post hoc tests. These analyses were followed

by residual analysis to verify the suitability of the error distribution and model fitting. These

Analyses were conducted using R 3.2.2 (43) using the packages "lme4", "car", "lsmeans" and

"RVAideMemoire" (108–112). Mass features were normalized using the function *arcsinh_x* in the

R package "bestNormalize" and compared between root types using functions *t.test* in base R and

foldchange in the package "gtools". P-values were FDR-corrected using the function *p.adjust*.

Acknowledgements: We thank Elisa J. Bernklau (Colorado State University) for advice on sugar

preference experiments.

Funding: This project was supported by a European Union Horizon 2020 Marie Sklodowska-Curie

Action (MSCA) Individual Fellowship (Grant # 794947 to B.C.J.S.) and the Swiss National

Science Foundation (Grants # 155781, 160786 and 157884 to M.E.).

Competing interests: The Authors declare no competing interests.

Data and materials availability: The data generated for this manuscript is available on Dryad

[DOI to be inserted].

References

1. A. Kempel, M. Razanajatovo, C. Stein, S. B. Unsicker, H. Auge, W. W. Weisser, M. Fischer, D. Prati, Herbivore preference drives plant community composition. *Ecology* **96**, 2923–2934 (2015).

2. E. S. Bakker, J. L. Gill, C. N. Johnson, F. W. M. Vera, C. J. Sandom, G. P. Asner, J.-C. Svenning, Combining paleo-data and modern exclosure experiments to assess the impact of megafauna extinctions on woody vegetation. *PNAS* **113**, 847–855 (2016).

3. S. B. Heard, L. C. Remer, Travel costs, oviposition behaviour and the dynamics of insect–plant systems. *Theor. Ecol.* **1**, 179–188 (2008).

4. Y. Xiao, Q. Wang, M. Erb, T. C. J. Turlings, L. Ge, L. Hu, J. Li, X. Han, T. Zhang, J. Lu, G. Zhang, Y. Lou, Specific herbivore-induced volatiles defend plants and determine insect community composition in the field. *Ecol. Lett.* **15**, 1130–1139 (2012).

5. N. M. van Dam, K. Hadwich, I. T. Baldwin, Induced responses in Nicotiana attenuata affect behavior and growth of the specialist herbivore *Manduca sexta*. *Oecologia* **122**, 371–379 (2000).

6. R. C. Smallegange, J. J. A. van Loon, S. E. Blatt, J. A. Harvey, N. Agerbirk, M. Dicke, Flower vs. leaf feeding by Pieris brassicae: glucosinolate-rich flower tissues are preferred and sustain higher growth rate. *J. Chem. Ecol.* **33**, 1831–1844 (2007).

7. S. N. Johnson, D. B. Read, P. J. Gregory, Tracking larval insect movement within soil using high resolution X-ray microtomography. *Ecol. Entomol.* **29**, 117–122 (2004).

8. J. Jiménez, E. Garzo, J. Alba-Tercedor, A. Moreno, A. Fereres, G. P. Walker, The phloem-pd: a distinctive brief sieve element stylet puncture prior to sieve element phase of aphid feeding behavior. *Arth.-Plant Interact.* **14**, 67–78 (2020).

9. S. Pentzold, A. Burse, W. Boland, Contact chemosensation of phytochemicals by insect herbivores. *Nat. Prod. Rep.* **34**, 478–483 (2017).

10. T. J. A. Bruce, L. J. Wadhams, C. M. Woodcock, Insect host location: a volatile situation. *Trends Plant Sci.* **10**, 269–274 (2005).

11. Y. Aartsma, A. Cusumano, M. Fernández de Bobadilla, Q. Rusman, I. Vosteen, E. H. Poelman, Understanding insect foraging in complex habitats by comparing trophic levels: insights from specialist host-parasitoid-hyperparasitoid systems. *Curr. Opin. Insect Sci.* **32**, 54–60 (2019).

12. D. Carrasco, M. C. Larsson, P. Anderson, Insect host plant selection in complex environments. *Curr. Opin. Insect Sci.* **8**, 1–7 (2015).

13. M. Erb, M. Huber, C. A.M. Robert, A. P. Ferrieri, R. A.R. Machado, C. C.M. Arce, "The Role of Plant Primary and Secondary Metabolites in Root-Herbivore Behaviour, Nutrition and Physiology" in *Behaviour and Physiology of Root Herbivores*, S. N. Johnson, I. Hiltpold, T. C. J. Turlings, Eds. (Elsevier Science, 2013), pp. 53–95.

741   14.   A. Reinecke, M. Hilker, "Plant Semiochemicals - Perception and Behavioural Responses
742   by Insects" in *Annual plant reviews*, J. A. Roberts, Ed. (Wiley, 2018-), pp. 115–153.

743   15.   S. N. Johnson, P. J. Gregory, Chemically-mediated host-plant location and selection by
744   root-feeding insects. *Physiol. Entomol.* **31**, 1–13 (2006).

745   16.   R. F. Chapman, Contact chemoreception in feeding by phytophagous insects. *Annu. Rev.*
746   *Entomol.* **48**, 455–484 (2003).

747   17.   S. T. Behmer, Insect herbivore nutrient regulation. *Annu. Rev. Entomol.* **54**, 165–187
748   (2009).

749   18.   M. D. Dearing, W. J. Foley, S. McLean, The Influence of Plant Secondary Metabolites on
750   the Nutritional Ecology of Herbivorous Terrestrial Vertebrates. *Annu. Rev. Ecol. Evol. Syst.*
751   **36**, 169–189 (2005).

752   19.   R. Nishida, Chemical ecology of insect-plant interactions: ecological significance of plant
753   secondary metabolites. *Biosci. Biotechnol. Biochem.* **78**, 1–13 (2014).

754   20.   W. Zhou, A. Kügler, E. McGale, A. Haverkamp, M. Knaden, H. Guo, F. Beran, F. Yon,
755   R. Li, N. Lackus, T. G. Köllner, J. Bing, M. C. Schuman, B. S. Hansson, D. Kessler, I. T.
756   Baldwin, S. Xu, Tissue-Specific Emission of (E)-α-Bergamotene Helps Resolve the Dilemma
757   When Pollinators Are Also Herbivores. *Curr. Biol.* **27**, 1336–1341 (2017).

758   21.   M. de Vos, K. L. Kriksunov, G. Jander, Indole-3-acetonitrile production from indole
759   glucosinolates deters oviposition by *Pieris rapae*. *Plant Physiol.* **146**, 916–926 (2008).

760   22.   R. A. Fandino, A. Haverkamp, S. Bisch-Knaden, J. Zhang, S. Bucks, T. A. T. Nguyen, K.
761   Schröder, A. Werckenthin, J. Rybak, M. Stengl, M. Knaden, B. S. Hansson, E. Große-Wilde,
762   Mutagenesis of odorant coreceptor Orco fully disrupts foraging but not oviposition behaviors
763   in the hawkmoth *Manduca sexta*. *PNAS* **116**, 15677–15685 (2019).

764   23.   Z.-J. Zhang, S.-S. Zhang, B.-L. Niu, D.-F. Ji, X.-J. Liu, M.-W. Li, H. Bai, S. R. Palli, C.-
765   Z. Wang, A.-J. Tan, A determining factor for insect feeding preference in the silkworm,
766   *Bombyx mori*. *PLoS Biol.* **17**, e3000162 (2019).

767   24.   K. Ozaki, M. Ryuda, A. Yamada, A. Utoguchi, H. Ishimoto, D. Calas, F. Marion-Poll, T.
768   Tanimura, H. Yoshikawa, A gustatory receptor involved in host plant recognition for
769   oviposition of a swallowtail butterfly. *Nat. Commun.* **2**, 542 (2011).

770   25.   M. Huber, J. Epping, C. Schulze Gronover, J. Fricke, Z. Aziz, T. Brillatz, M. Swyers, T.
771   G. Köllner, H. Vogel, A. Hammerbacher, D. Triebwasser-Freese, C. A. M. Robert, K.
772   Verhoeven, V. Preite, J. Gershenzon, M. Erb, A Latex Metabolite Benefits Plant Fitness under
773   Root Herbivore Attack. *PLoS Biol.* **14**, e1002332 (2016).

774   26.   Z. Bont, C. Arce, M. Huber, W. Huang, A. Mestrot, C. J. Sturrock, M. Erb, A Herbivore
775   Tag-and-Trace System Reveals Contact- and Density-Dependent Repellence of a Root Toxin.
776   *J. Chem. Ecol.* **43**, 295–306 (2017).

777   27.   T. Hartmann, From waste products to ecochemicals: fifty years research of plant
778   secondary metabolism. *Phytochemistry* **68**, 2831–2846 (2007).

28.   S. N. Johnson, U. N. Nielsen, Foraging in the dark - chemically mediated host plant location by belowground insect herbivores. *J. Chem. Ecol.* **38**, 604–614 (2012).

29.   T. J. A. Bruce, J. A. Pickett, Perception of plant volatile blends by herbivorous insects-- finding the right mix. *Phytochemistry* **72**, 1605–1611 (2011).

30.   L. A. Richards, A. E. Glassmire, K. M. Ochsenrider, A. M. Smilanich, C. D. Dodson, C. S. Jeffrey, L. A. Dyer, Phytochemical diversity and synergistic effects on herbivores. *Phytochem. Rev.* **15**, 1153–1166 (2016).

31.   M. P. Calcagno, J. Coll, J. Lloria, F. Faini, M. E. Alonso-Amelot, Evaluation of synergism in the feeding deterrence of some furanocoumarins on *Spodoptera littoralis*. *J. Chem. Ecol.* **28**, 175–191 (2002).

32.   Y. Akhtar, M. B. Isman, Binary mixtures of feeding deterrents mitigate the decrease in feeding deterrent response to antifeedants following prolonged exposure in the cabbage looper, *Trichoplusia ni* (Lepidoptera: Noctuidae). *Chemoecol.* **13**, 177–182 (2003).

33.   S. J. Simpson, D. Raubenheimer, The geometric analysis of nutrient–allelochemical interactions: a case study using locusts. *Ecology* **82**, 422–439 (2001).

34.   S. T. Behmer, S. J. Simpson, D. Raubenheimer, Herbivore foraging in chemically heterogeneous environments: nutrients and secondary metabolites. *Ecology* **83**, 2489–2501 (2002).

35.   M. S. Singer, E. A. Bernays, Y. Carrière, The interplay between nutrient balancing and toxin dilution in foraging by a generalist insect herbivore. *Anim. Behav.* **64**, 629–643 (2002).

36.   E. Baraza, J. J. Villalba, F. D. Provenza, Nutritional context influences preferences of lambs for foods with plant secondary metabolites. *Appl. Anim. Behav. Sci.* **92**, 293–305 (2005).

37.   J. J. Villalba, F. D. Provenza, J. P. Bryant, Consequences of the interaction between nutrients and plant secondary metabolites on herbivore selectivity: benefits or detriments for plants? *Oikos* **97**, 282–292 (2002).

38.   N. Cocco, J. I. Glendinning, Not all sugars are created equal: some mask aversive tastes better than others in an herbivorous insect. *J. Exp. Biol.* **215**, 1412–1421 (2012).

39.   G. H. Pyke, Optimal Foraging Theory: A Critical Review. *Annu. Rev. Ecol. Syst.* **15**, 523–575 (1984).

40.   S. Gripenberg, P. J. Mayhew, M. Parnell, T. Roslin, A meta-analysis of preference-performance relationships in phytophagous insects. *Ecol. Lett.* **13**, 383–393 (2010).

41.   C. A. Kelly, M. D. Bowers, Preference and performance of generalist and specialist herbivores on chemically defended host plants. *Ecol. Entomol.* **41**, 308–316 (2016).

42.   F. R. Badenes-Pérez, M. Reichelt, J. Gershenzon, D. G. Heckel, Phylloplane location of glucosinolates in Barbarea spp. (Brassicaceae) and misleading assessment of host suitability by a specialist herbivore. *New Phytol.* **189**, 549–556 (2011).

43.   C. A. M. Robert, M. Erb, I. Hiltpold, B. E. Hibbard, M. D. P. Gaillard, J. Bilat, J.
Degenhardt, X. Cambet-Petit-Jean, T. C. J. Turlings, C. Zwahlen, Genetically engineered
maize plants reveal distinct costs and benefits of constitutive volatile emissions in the field.
Plant Biotechnol. J. **11**, 628–639 (2013).

44.   R. A. Machado, M. McClure, M. R. Hervé, I. T. Baldwin, M. Erb, Benefits of jasmonate-
dependent defenses against vertebrate herbivores in nature. *Elife* **5 (**2016).

45.   A. R. Agnihotri, A. A. Roy, R. S. Joshi, Gustatory receptors in Lepidoptera:
chemosensation and beyond. *Insect molecular biology* **25**, 519–529 (2016).

46.   A. de Fouchier, W. B. Walker, N. Montagné, C. Steiner, M. Binyameen, F. Schlyter, T.
Chertemps, A. Maria, M.-C. François, C. Monsempes, P. Anderson, B. S. Hansson, M. C.
Larsson, E. Jacquin-Joly, Functional evolution of Lepidoptera olfactory receptors revealed by
deorphanization of a moth repertoire. *Nat. Commun.* **8**, 15709 (2017).

47.   J. Shim, Y. Lee, Y. T. Jeong, Y. Kim, M. G. Lee, C. Montell, S. J. Moon, The full
repertoire of Drosophila gustatory receptors for detecting an aversive compound. *Nat.*
Commun. **6**, 8867 (2015).

48.   W. D. Jones, P. Cayirlioglu, I. G. Kadow, L. B. Vosshall, Two chemosensory receptors
together mediate carbon dioxide detection in Drosophila. *Nature* **445**, 86–90 (2007).

49.   T. Miyamoto, J. Slone, X. Song, H. Amrein, A fructose receptor functions as a nutrient
sensor in the Drosophila brain. *Cell* **151**, 1113–1125 (2012).

50.   X.-L. Liu, Q. Yan, Y.-L. Yang, W. Hou, C.-L. Miao, Y.-C. Peng, S.-L. Dong, A
Gustatory Receptor GR8 Tunes Specifically to D-Fructose in the Common Cutworm
Spodoptera litura. *Insects* **10 (**2019).

51.   S. J. Moon, Y. Lee, Y. Jiao, C. Montell, A Drosophila gustatory receptor essential for
aversive taste and inhibiting male-to-male courtship. *Curr. Biol.* **19**, 1623–1627 (2009).

52.   Y. Lee, S. J. Moon, Y. Wang, C. Montell, A Drosophila Gustatory Receptor Required for
Strychnine Sensation. *Chemical senses* **40**, 525–533 (2015).

53.   X.-L. Liu, S.-J. Sun, W. Hou, J. Zhang, Q. Yan, S.-L. Dong, Functional characterization
of two spliced variants of fructose gustatory receptor in the diamondback moth, *Plutella*
xylostella. *Pestic. Biochem. Physiol.* **164**, 7–13 (2020).

54.   X.-J. Jiang, C. Ning, H. Guo, Y.-Y. Jia, L.-Q. Huang, M.-J. Qu, C.-Z. Wang, A gustatory
receptor tuned to D-fructose in antennal sensilla chaetica of *Helicoverpa armigera*. *Insect*
Biochem. Mol. Biol. **60**, 39–46 (2015).

55.   D. Mishra, T. Miyamoto, Y. H. Rezenom, A. Broussard, A. Yavuz, J. Slone, D. H.
Russell, H. Amrein, The molecular basis of sugar sensing in Drosophila larvae. *Curr. Biol.* **23**,
1466–1471 (2013).

56.   R. Delventhal, J. R. Carlson, Bitter taste receptors confer diverse functions to neurons.
Elife **5 (**2016).

57. S. Fujii, A. Yavuz, J. Slone, C. Jagge, X. Song, H. Amrein, Drosophila sugar receptors in sweet taste perception, olfaction, and internal nutrient sensing. *Curr. Biol.* **25**, 621–627 (2015).

58. M. Schumann, Z. S. Ladin, J. M. Beatens, I. Hiltpold, Navigating on a chemical radar: Usage of root exudates by foraging *Diabrotica virgifera virgifera* larvae. *J. Appl. Entomol.* **142**, 911–920 (2018).

59. J. L. Spencer, B. E. Hibbard, J. Moeser, D. W. Onstad, Behaviour and ecology of the western corn rootworm (*Diabrotica virgifera virgifera* LeConte). *Agric. For. Entomol.* **11**, 9–27 (2009).

60. E. J. Bernklau, L. B. Bjostad, Reinvestigation of Host Location by Western Corn Rootworm Larvae (Coleoptera: Chrysomelidae): CO_2 Is the Only Volatile Attractant. *J. Econ. Entomol.* **91**, 1331–1340 (1998).

61. E. J. Bernklau, L. B. Bjostad, Identification of Feeding Stimulants in Corn Roots for Western Corn Rootworm (Coleoptera: Chrysomelidae) Larvae. *J. Econ. Entomol.* **101**, 341–351 (2008).

62. E. J. Bernklau, B. E. Hibbard, D. L. Dick, C. D. Rithner, L. B. Bjostad, Monogalactosyldiacylglycerols as Host Recognition Cues for Western Corn Rootworm Larvae (Coleoptera: Chrysomelidae). *J. Econ. Entomol.* **108**, 539–548 (2015).

63. E. J. Bernklau, B. E. Hibbard, A. P. Norton, L. B. Bjostad, Methyl Anthranilate as a Repellent for Western Corn Rootworm Larvae (Coleoptera: Chrysomelidae). *J. Econ. Entomol.* **109**, 1683–1690 (2016).

64. C. A. M. Robert, M. Erb, M. Duployer, C. Zwahlen, G. R. Doyen, T. C. J. Turlings, Herbivore-induced plant volatiles mediate host selection by a root herbivore. *New Phytol.* **194**, 1061–1069 (2012).

65. M. Erb, C. A. M. Robert, G. Marti, J. Lu, G. R. Doyen, N. Villard, Y. Barrière, B. W. French, J.-L. Wolfender, T. C. J. Turlings, J. Gershenzon, A Physiological and Behavioral Mechanism for Leaf Herbivore-Induced Systemic Root Resistance. *Plant Physiol.* **169**, 2884–2894 (2015).

66. L. Hu, P. Mateo, M. Ye, X. Zhang, J. D. Berset, V. Handrick, D. Radisch, V. Grabe, T. G. Köllner, J. Gershenzon, C. A. M. Robert, M. Erb, Plant iron acquisition strategy exploited by an insect herbivore. *Science* **361**, 694–697 (2018).

67. V. Theepan, B. C. J. Schimmel, C. C. M. Arce, M. Erb, R. A. R. Machado, *Plant-derived CO_2 mediates long-distance host location and quality assessment by a root herbivore* (2020).

68. C. A. M. Robert, M. Erb, B. E. Hibbard, B. Wade French, C. Zwahlen, T. C. J. Turlings, A specialist root herbivore reduces plant resistance and uses an induced plant volatile to aggregate in a density-dependent manner. *Funct. Ecol.* **26**, 1429–1440 (2012).

69. C. A. M. Robert, N. Veyrat, G. Glauser, G. Marti, G. R. Doyen, N. Villard, M. D. P. Gaillard, T. G. Köllner, D. Giron, M. Body, B. A. Babst, R. A. Ferreri, T. C. J. Turlings, M.

Erb, A specialist root herbivore exploits defensive metabolites to locate nutritious tissues.
Ecol. Lett. **15**, 55–64 (2012).

70.   F. Hochholdinger, R. Tuberosa, Genetic and genomic dissection of maize root
development and architecture. *Curr. Opin. Plant Biol.* **12**, 172–177 (2009).

71.   M. Frey, P. Chomet, E. Glawischnig, C. Stettner, S. Grün, A. Winklmair, W. Eisenreich,
A. Bacher, R. B. Meeley, S. P. Briggs, K. Simcox, A. Gierl, Analysis of a chemical plant
defense mechanism in grasses. *Science* **277**, 696–699 (1997).

72.   D. Maag, A. Köhler, C. A. M. Robert, M. Frey, J.-L. Wolfender, T. C. J. Turlings, G.
Glauser, M. Erb, Highly localized and persistent induction of Bx1-dependent herbivore
resistance factors in maize. *Plant J.* **88**, 976–991 (2016).

73.   T. L. Slewinski, R. Meeley, D. M. Braun, Sucrose transporter1 functions in phloem
loading in maize leaves. *J. Exp. Bot.* **60**, 881–892 (2009).

74.   R. A. R. Machado, C. C. M. Arce, A. P. Ferrieri, I. T. Baldwin, M. Erb, Jasmonate-
dependent depletion of soluble sugars compromises plant resistance to *Manduca sexta*. *New
Phytol.* **207**, 91–105 (2015).

75.   J. Liu, H. Wu, J. Yi, D. Jiang, G. Zhang, Identification and functional characterization of
D-fructose receptor in an egg parasitoid, *Trichogramma chilonis*. *PLoS ONE* **14**, e0217493
(2019).

76.   K. Sato, K. Tanaka, K. Touhara, Sugar-regulated cation channel formed by an insect
gustatory receptor. *PNAS* **108**, 11680–11685 (2011).

77.   T. Takada, T. Sasaki, R. Sato, S. Kikuta, M. N. Inoue, Differential expression of a
fructose receptor gene in honey bee workers according to age and behavioral role. *Arch.*
Insect. Biochem. Physiol. **97** (2018).

78.   W. Xu, H.-J. Zhang, A. Anderson, A sugar gustatory receptor identified from the foregut
of cotton bollworm *Helicoverpa armigera*. *J. Chem. Ecol.* **38**, 1513–1520 (2012).

79.   M. R. Hervé, M. Erb, Distinct defense strategies allow different grassland species to cope
with root herbivore attack. *Oecologia* **191**, 127–139 (2019).

80.   X. Zhang, C. van Doan, C. C. M. Arce, L. Hu, S. Gruenig, C. Parisod, B. E. Hibbard, M.
R. Hervé, C. Nielson, C. A. M. Robert, R. A. R. Machado, M. Erb, Plant defense resistance in
natural enemies of a specialist insect herbivore. *PNAS* **116**, 23174–23181 (2019).

81.   C. A.M. Robert, X. Zhang, R. A. Machado, S. Schirmer, M. Lori, P. Mateo, M. Erb, J.
Gershenzon, Sequestration and activation of plant toxins protect the western corn rootworm
from enemies at multiple trophic levels. *Elife* **6** (2017).

82.   D. W. Onstad, B. E. Hibbard, T. L. Clark, D. W. Crowder, K. G. Carter, Analysis of
Density-Dependent Survival of Diabrotica (Coleoptera: Chrysomelidae) in Cornfields.
Environ. Entomol. **35**, 1272–1278 (2006).

83. E. Y. Yu, A. J. Gassmann, T. W. Sappington, Effects of larval density on dispersal and fecundity of western corn rootworm, *Diabrotica virgifera virgifera* LeConte (Coleoptera: Chrysomelidae). *PLoS ONE* **14**, e0212696 (2019).

84. R. Silva, A. R. Clarke, The "sequential cues hypothesis": a conceptual model to explain host location and ranking by polyphagous herbivores. *Insect Sci.* 10.1111/1744-7917.12719 (2019).

85. K. Tsuneto, H. Endo, F. Shii, K. Sasaki, S. Nagata, R. Sato, *The two-step chemosensory system underlying the oligophagy of silkworm larvae* (2019).

86. A. L. Neal, S. Ahmad, R. Gordon-Weeks, J. Ton, Benzoxazinoids in root exudates of maize attract *Pseudomonas putida* to the rhizosphere. *PLoS ONE* **7**, e35498 (2012).

87. L. C. Carvalhais, P. G. Dennis, D. Fedoseyenko, M.-R. Hajirezaei, R. Borriss, N. von Wirén, Root exudation of sugars, amino acids, and organic acids by maize as affected by nitrogen, phosphorus, potassium, and iron deficiency. *J. Plant Nutr. Soil Sci.* **174**, 3–11 (2011).

88. R. Dick, T. Rattei, M. Haslbeck, W. Schwab, A. Gierl, M. Frey, Comparative analysis of benzoxazinoid biosynthesis in monocots and dicots: independent recruitment of stabilization and activation functions. *Plant Cell* **24**, 915–928 (2012).

89. P. Voothuluru, D. M. Braun, J. S. Boyer, An *in Vivo* Imaging Assay Detects Spatial Variability in Glucose Release from Plant Roots. *Plant Physiol.* **178**, 1002–1010 (2018).

90. R. Shroff, F. Vergara, A. Muck, A. Svatos, J. Gershenzon, Nonuniform distribution of glucosinolates in *Arabidopsis thaliana* leaves has important consequences for plant defense. *PNAS* **105**, 6196–6201 (2008).

91. P. L. Jones, A. A. Agrawal, Learning in Insect Pollinators and Herbivores. *Annu. Rev. Entomol.* **62**, 53–71 (2017).

92. C.-C. Chu, J. L. Spencer, M. J. Curzi, J. A. Zavala, M. J. Seufferheld, Gut bacteria facilitate adaptation to crop rotation in the western corn rootworm. *PNAS* **110**, 11917–11922 (2013).

93. R. Karban, A. A. Agrawal, Herbivore Offense. *Annu. Rev. Ecol. Syst.* **33**, 641–664 (2002).

94. D. Maag, A. Köhler, C. Am Robert, M. Frey, J.-L. Wolfender, T. C. J. Turlings, G. Glauser, M. Erb, Highly localized and persistent induction of bx1-dependent herbivore resistance factors in maize. *The Plant Journal* **88**, 976–991 (2016).

95. C. A. M. Robert, N. Veyrat, G. Glauser, G. Marti, G. R. Doyen, N. Villard, M. D. P. Gaillard, T. G. Köllner, D. Giron, M. Body, A specialist root herbivore exploits defensive metabolites to locate nutritious tissues. *Ecology letters* **15**, 55–64 (2012).

96. L. Hu, P. Mateo, M. Ye, X. Zhang, J. D. Berset, V. Handrick, D. Radisch, V. Grabe, T. G. Köllner, J. Gershenzon, Plant iron acquisition strategy exploited by an insect herbivore. *Science* **361**, 694–697 (2018).

964   97.   E. J. Bernklau, B. E. Hibbard, L. B. Bjostad, Sugar preferences of western corn rootworm
965   larvae in a feeding stimulant blend. *Journal of applied entomology* **142**, 947–958 (2018).

966   98.   L. Hu, C. A. M. Robert, S. Cadot, X. Zhang, M. Ye, B. Li, D. Manzo, N. Chervet, T.
967   Steinger, M. G. A. van der Heijden, K. Schlaeppi, M. Erb, Root exudate metabolites drive
968   plant-soil feedbacks on growth and defense by shaping the rhizosphere microbiota. *Nat.*
969   *Commun.* **9**, 2738 (2018).

970   99.   R. A. R. Machado, C. C. M. Arce, M. A. McClure, I. T. Baldwin, M. Erb, Aboveground
971   herbivory induced jasmonates disproportionately reduce plant reproductive potential by
972   facilitating root nematode infestation. *Plant Cell Environ.* **41**, 797–808 (2018).

973   100.   J. S. Velterop, F. Vos, A rapid and inexpensive microplate assay for the enzymatic
974   determination of glucose, fructose, sucrose, L-malate and citrate in tomato (Lycopersicon
975   esculentum) extracts and in orange juice. *Phytochem. Anal.* **12**, 299–304 (2001).

976   101.   S. Kumar, G. Stecher, K. Tamura, MEGA7: Molecular Evolutionary Genetics Analysis
977   Version 7.0 for Bigger Datasets. *Mol. Biol. Evol.* **33**, 1870–1874 (2016).

978   102.   S. Q. Le, O. Gascuel, An improved general amino acid replacement matrix. *Mol. Biol.*
979   *Evol.* **25**, 1307–1320 (2008).

980   103.   I. Letunic, P. Bork, Interactive tree of life (iTOL) v3: an online tool for the display and
981   annotation of phylogenetic and other trees. *Nucleic. Acids. Res.* **44**, W242-5 (2016).

982   104.   L. A. Kelley, S. Mezulis, C. M. Yates, M. N. Wass, M. J. E. Sternberg, The Phyre2 web
983   portal for protein modeling, prediction and analysis. *Nat. Protoc.* **10**, 845–858 (2015).

984   105.   H. M. Robertson, C. G. Warr, J. R. Carlson, Molecular evolution of the insect
985   chemoreceptor gene superfamily in *Drosophila melanogaster*. *PNAS* **100 Suppl 2**, 14537–
986   14542 (2003).

987   106.   E. Kim, Y. Park, Y. Kim, A Transformed Bacterium Expressing Double-Stranded RNA
988   Specific to Integrin β1 Enhances Bt Toxin Efficacy against a Polyphagous Insect Pest,
989   *Spodoptera exigua*. *PLoS ONE* **10**, e0132631 (2015).

990   107.   K. J. Livak, T. D. Schmittgen, Analysis of relative gene expression data using real-time
991   quantitative PCR and the 2− ∆∆CT method. *Methods* **25**, 402–408 (2001).

992   108.   M. Hervé, RVAideMemoire: diverse basic statistical and graphical functions. *R package*
993   *version 0.9-50 (*2015).

994   109.   D. Bates, M. Mächler, B. Bolker, S. Walker, *Fitting Linear Mixed-Effects Models using*
995   *lme4* (2014).

996   110.   R. V. Lenth, Least-Squares Means: The R Package lsmeans. *J. Stat. Soft.* **69 (**2016).

997   111.   R Development Core Team, RFFSC, *R: A language and environment for statistical*
998   *computing* (R foundation for statistical computing Vienna, Austria, 2011).

999   112.   J. Fox, S. Weisberg, Multivariate linear models in R. *An R Companion to Applied*
1000   *Regression. Los Angeles: Thousand Oaks (*2011).

1001   113.   L. B. Bjostad, B. E. Hibbard, 6-Methoxy-2-benzoxazolinone: A semiochemical for host
1002   location by western corn rootworm larvae. *J. Chem. Ecol.* **18**, 931–944 (1992).
1003

1004 **Table S1.** Identified metabolites in maize post-embryonic and embryonic roots. Metabolites with retention times
1005 highlighted by asterisks (*) eluted together with many weakly-retained compounds that could not be separated by
1006 reversed-phase liquid chromatography (RP-UHPLC). Their identification is therefore tentative and based on accurate
1007 masses only.

Nr	Compound class	Compound Name	Molecular formula	Retention time [min]	MS adducts	Progenesis QI fragmentation score	Characteristic fragments [m/z]
1	Benzoxazinoid	DHBOA-Glc	$C_{14}H_{17}NO_9$	1.30	M-H, M+FA-H, 2M-H	16.5	180.0303, 162.0200
2	Benzoxazinoid	DIBOA-Glc	$C_{14}H_{17}NO_9$	1.91	M-H, M+FA-H, 2M-H	2.6	180.0303, 162.0200, 134.0247
3	Benzoxazinoid	HMBOA-Glc	$C_{15}H_{19}NO_9$	2.11	M-H, M+FA-H, 2M-H	10.3	194.0459, 166.0511
4	Benzoxazinoid	DIMBOA-Glc	$C_{15}H_{19}NO_{10}$	2.18	M-H, M+FA-H, 2M-H	5.9	210.0406, 192.0302, 164.0356, 149.0119
5	Benzoxazinoid	DIM$_2$BOA-Glc	$C_{16}H_{21}NO_{11}$	2.22	M-H, M+FA-H, 2M-H	21.7	222.0408, 194.0458
6	Benzoxazinoid	HMBOA	$C_9H_9NO_4$	2.30	M-H	n/a	-
7	Benzoxazinoid	DIMBOA	$C_9H_9NO_5$	2.42	M-H	n/a	164.0353, 149.0118
8	Benzoxazinoid	HDMBOA-Glc	$C_{16}H_{21}NO_{10}$	2.74	M-H, M+FA-H, 2M-H	11.5	224.0564, 194.0460
9	Benzoxazinoid	HDM$_2$BOA-Glc	$C_{17}H_{23}NO_{11}$	2.74	M+FA-H	n/a	254.0789
10	Organic acid	Chlorogenic acid	$C_{16}H_{18}O_9$	1.77	M-H, 2M-H	24.3	191.0 562
11	Organic acid	Feruloylglucose	$C_{16}H_{20}O_9$	2.05	M-H	81.0	175.0 404
12	Organic acid	Sinapoylglucose	$C_{17}H_{22}O_{10}$	2.09	M-H	36.7	223.0615, 205.0507
13	Organic acid	Quinic acid	$C_{17}H_{12}O_6$	0.33*	M-H, 2M-H	70.1	-
14	Organic acid	Coumaroylquinic acid isomer	$C_{16}H_{18}O_8$	1.70	M-H	98.0	163.0409
15	Organic acid	Coumaroylquinic acid isomer	$C_{16}H_{18}O_8$	2.13	M-H	51.6	191.0561, 173.0455
16	Organic acid	Feruloylquinic acid isomer	$C_{17}H_{20}O_9$	2.35	M-H	46.1	191.0562
17	Organic acid	Feruloylquinic acid isomer	$C_{17}H_{20}O_9$	2.61	M-H	n/a	-
18	Flavonol	Quercetin rhamnosylglucoside	$C_{27}H_{30}O_{16}$	2.66	M-H	65.7	301.0353
19	Flavonol	Quercetin glucoside	$C_{21}H_{20}O_{12}$	2.75	M-H	26.0	301.0347, 271.0252
20	Flavonol	Quercetin malonylglucoside	$C_{24}H_{22}O_{15}$	2.91	M-H	75.1	505.0986, 301.0345

21	Flavonol	Kaempferol rhamnosylglucoside	$C_{27}H_{30}O_{15}$	2.94	M-H	98.9	285.0412
22	Flavonol	Kaempferol glucoside	$C_{21}H_{20}O_{11}$	3.04	M-H	88.6	285.0395, 255.0304
23	Flavonol	Kaempferol malonylglucoside	$C_{24}H_{22}O_{14}$	3.25	M-H	96.5	489.1037, 285.0400
24	Amino acid	Leucine	$C_6H_{13}NO_2$	0.80	M-H	n/a	-
25	Amino acid	Phenylalanine	$C_9H_{11}NO_2$	1.10	M-H	n/a	-
26	Amino acid	Tryptophan	$C_{11}H_{12}N_2O_2$	1.50	M-H, M+FA-H	n/a	-
27	Sugar	Monosaccharide	$C_6H_{12}O_6$	0.31*	M-H	n/a	-
28	Sugar	Disaccharide	$C_{12}H_{22}O_{11}$	0.34*	M-H, M+FA-H, 2M-H	3.2	-

1008

1009 **Table S2.** Root metabolites with behavioral activity towards western corn rootworm larvae.

Nr	Compound class	Compound Name	Behavioral effect	Reference.
1	Benzoxazinoid	MBOA	Host location (long-distance)	(113)
2	Benzoxazinoid	Fe-DIMBOA	Host location & acceptance (short-distance)	(66)
3	Benzoxazinoid	Fe-DIMBOA-Glc	Host location & acceptance (short-distance)	(66)
4	Sugar	Glucose	Feeding stimulation (within sugar blend)	(61)
5	Sugar	Fructose	Feeding stimulation (within sugar blend)	(61)
6	Sugar	Sucrose	Feeding stimulation (within sugar blend)	(61)
7	Fatty acid	Linoleic acid	Arrestment, feeding stimulation (within sugar blend)	(61)
8	Fatty acid	Oleic acid	Feeding stimulation (within sugar blend)	(61)
9	Fatty acid	Monoga-lactosyldiacylglyc erol	Host recognition (tight turning behavior)	(62)
10	Aminobenzole	Methyl anthranilate	Host avoidance (repellent)	(63)
11	Terpene	(*E*)-β-caryophyllene	Host location and discrimination (long distance)	(43, 68)
12	Alkene	Ethylene	Host location and discrimination (long-distance	(64)
13	Inorganic compound	CO_2	Host location (long-distance)	(60)

1010

1011

1012

Fig. S1. *Frequency of different types of damage on individual embryonic and post embryonic roots.* Data is from experiment shown in Fig. 1*B* (n=20 plants). P1-P2 refer to individual embryonic roots, C1-C6 to individual post-embryonic roots per plant. Numbers were assigned randomly to embryonic and post embryonic roots within plants. The sizes of the circles are proportional to the relative frequency (% within each root) of the different types of observed damage.

1018

Fig. S2. *Metabolic differences between embryonic and post-embryonic roots are conserved in the bx1 mutant.*
Relative abundances (signal intensities) of identified metabolic features in embryonic and post-embryonic roots of
wild type B73 and *bx1* mutant plants (n=7-10). For sugars and benzoxazinoids, refer to Fig. 2. Results of Two-way
ANOVAs for genotype effects (G), root type effects (R) and their interaction (GxR) are shown for each compound
(***p<0.001, **p<0.01, *p<0.05). Error bars denote standard errors of means (SEM).

1024

1025 **Fig. S3.** *DvvGr43a does not influence the attraction of the western corn rootworm to CO_2.* Proportion of control
1026 (GFP) or *DvvGr43a*-silenced (Gr43a) larvae found on each arm of belowground olfactometers. Asterisks indicate
1027 significant differences between treatments (*$p < 0.05$, FDR-corrected Least Square Mean post hoc tests, six two-arm
1028 olfactometers with 6 larvae each were evaluated, n=6).

1029

1030 **Fig. S4.** *DvvGr43a mediates sugar preference independently of the feeding state of the western corn rootworm.*
1031 Preference of satiated (left) and starved (right) control or *DvvGr43a*-silenced larvae for buffer or a glucose, fructose,
1032 sucrose mixture on filter discs at different time points (*$p<0.05$, FDR-corrected Least Square Mean post hoc tests,
1033 eight petri plates with six larvae each were assayed, n=8). Error bars denote standard errors of means (SEM).

1034

1035

1036   **Fig. S5.** *DvvGr43a does not directly impair larval performance*. (*A*) Weight of western corn rootworm larvae fed on
1037   GFP and DvvGr43a dsRNA on young maize seedlings that produce embryonic roots only (no-choice setting) for 7
1038   days (n=40 cups with 9 larvae each). (*B*) Larval mortality within the same experiment.