Peptide Gaussian accelerated molecular dynamics (Pep-GaMD): Enhanced sampling and free energy calculations of peptide binding

Jinan Wang and Yinglong Miao*

Center for Computational Biology and Department of Molecular Biosciences, University of

Kansas, Lawrence, KS 66047, USA

* Corresponding email: miao@ku.edu

Abstract

Peptides mediate up to 40% of known protein-protein interactions in higher eukaryotes and play an important role in cellular signaling. However, it is challenging to simulate both binding and unbinding of peptides and calculate peptide binding free energies through conventional molecular dynamics, due to long biological timescales and extremely high flexibility of the peptides. Based on the Gaussian accelerated molecular dynamics (GaMD) enhanced sampling technique, we have developed a new computational method "Pep-GaMD", which selectively boosts essential interaction potential energy of the peptide to accelerate its dissociation. In addition, another boost potential is applied to the remaining potential energy of the entire system in a dual-boost algorithm to facilitate rebinding of the peptide. Pep-GaMD has been demonstrated on binding of three model peptides to the SH3 domains. Independent 1 μs Pep-GaMD simulations have captured repetitive peptide dissociation and binding events, which enable us to calculate peptide binding free energies. The calculated binding free energies agreed very well with available experimental data. Furthermore, the all-atom Pep-GaMD simulations have provided important insights into the mechanism of peptide binding to proteins that involves long-rang electrostatic interactions and mainly conformational selection. In summary, Pep-GaMD provides a highly efficient, easy-to-use approach for unconstrained enhanced sampling and free energy calculations of peptide binding.

Keywords: Peptide Gaussian accelerated molecular dynamics (Pep-GaMD), Peptide dissociation and binding, Enhanced sampling, Binding free energy, Conformational selection.

Significance Statement

We have developed a new computational method "Pep-GaMD" for enhanced sampling of peptide-protein interactions based on the Gaussian accelerated molecular dynamics (GaMD) technique. Pep-GaMD works by selectively boosting the essential interaction potential energy of the peptide to accelerate its dissociation. In addition, another boost potential can be applied to the remaining potential energy of the entire system in a dual-boost algorithm to facilitate rebinding of the peptide. Pep-GaMD has been demonstrated on binding of three model peptides to the SH3 domains. Pep-GaMD has captured repetitive peptide dissociation and binding events within significantly shorter simulation time (microsecond) than conventional molecular dynamics. Compared with previous enhanced sampling methods, Pep-GaMD is much easier to use for unconstrained enhanced sampling, which provides a novel physics-based approach to calculating peptide binding free energies using the 3D free energy surface.

Introduction

Peptides mediate up to 40% of known protein-protein interactions in higher eukaryotes[1]. They play a key role in cellular signaling, protein trafficking, immune response and oncology[1, 2]. Protein-peptide interactions have emerged as attractive drug targets for both small molecules and designed inhibitory peptides[3-5]. Increasing number of peptide-based drugs is being licensed to market in recent years[6-9]. Therefore, understanding molecular mechanism of protein-peptide recognition has important applications in the fields of biology, medicine, and pharmaceutical sciences.

It is important to characterize protein-peptide binding conformations for rational structure-based design of peptide drugs. Experimental techniques including X-ray crystallography, nuclear magnetic resonance (NMR) and cryo-electron microscopy (cryo-EM) have been utilized to determine the structures of protein-peptide complexes[2]. Recent years have seen dramatic increase in the number of experimental protein-peptide structures. However, only static snapshots of the protein-peptide interactions could usually be captured in these structures. The thermodynamics and dynamic mechanism of peptide recognition by proteins remains poorly understood[10].

Computational methods including peptide docking have been developed to model peptide-protein structures[11, 12]. In this regard, modeling of peptide binding to proteins has been shown to be distinct from that of extensively studied protein-ligand binding and protein-protein interactions. Notably, small-molecule ligands are able to bind deeply buried sites in proteins, but peptides normally bind to protein surface, especially in the large pockets. On the other hand, peptide-protein interactions are typically weaker than protein-protein interactions, because of the small interface between peptides and their target proteins. Moreover, protein partners usually have well defined 3D structures before forming protein-protein complexes, despite possible conformational changes during association. In contrast, most peptides do not have stable structures before forming

complexes with proteins. It is difficult to incorporate extremely high flexibility of peptides and their large conformational changes (folding and unfolding) into computational modeling, especially for peptide docking[11].

Molecular dynamics (MD) is a powerful technique for all-atom simulations of biomolecules[13]. MD has been used to refine binding poses of peptides in proteins obtained from docking[14-18]. In addition, MD simulations are able to account for the flexibility of peptides and have been applied to explore their mechanisms of binding to proteins[8, 19-26]. Notably, MD simulations have successfully revealed the pathway of fast association of a proline-rich motif (PRM) peptide to the SH3 domain[19]. Supervised MD simulations have captured binding pathways of a natural peptide BAD to the Bcl-X_L protein and the p53 peptide and SAH-p53-8 stapled peptidomimetic to the MDM2 protein[24]. Weighted ensemble of a total amount of ~120 μs MD simulations has been obtained to investigate binding of the p53 peptide to the MDM2 protein in implicit solvent[23]. The simulation predicted binding rate constant agrees very well with the experiments. Very recently, MD simulation performed for 200 μs using the Anton specialized supercomputer has captured 70 binding and unbinding events between an intrinsically disordered protein fragment of the measles virus nucleoprotein and the X domain of the measles virus phosphoprotein complex, which enables detailed understanding of the peptide "folding-upon-binding" mechanism[26]. Despite these remarkable advances, these studies based on conventional MD (cMD) often suffer from high computational cost for simulations of peptide-protein interactions. It remains challenging to sufficiently sample peptide-protein interactions through cMD simulations, especially for slow dissociation of various peptides that could take place over milliseconds and longer timescales[27].

Furthermore, enhanced sampling MD methods, including steered MD[28], Modeling by Employing Limited Data (MELD) using temperature and Hamiltonian replica exchange MD[29, 30], temperature-accelerated MD[31], multi-ensemble Markov state models[27] and metadynamics[32], have been applied to improve sampling of peptide-protein interactions. Steered MD have allowed simulations of antigenic peptide unbinding from the T-cell receptor based on a dissociation reaction coordinate[28]. MELD-accelerated MD simulations have been performed to predict binding poses and relative binding free energies of the p53-derived peptides and stapled α-helical peptides to the MDM2 and MDMX proteins[31, 32]. Multi-ensemble Markov models combining hundreds-of-microsecond cMD and Hamiltonian replica exchange simulations have been implemented to characterize both dissociation and binding of the PMI peptide inhibitor to the MDM2 oncoprotein fragment[27]. While cMD was able to simulate fast events such as peptide binding, enhanced sampling simulations could capture rare events such as peptide unbinding beyond the seconds timescale. The simulation derived dissociation and association rates were found in good agreement with the experimental measurements[27]. Recently, bias-exchange metadynamics simulations with 8 defined collective variables (CVs) performed for a total of 27 μs have been used to calculate binding free energy of the p53 peptide to the MDM2 protein. Infrequent metadynamics simulations with 3 CVs have also been successfully applied to predict peptide binding and dissociation rates for the same system[32]. Therefore, enhanced MD simulations have greatly expanded our capabilities in studies of peptide-protein interactions. Nevertheless, enhanced sampling of peptide binding to proteins are still underexplored compared with more extensive studies of protein-ligand binding and protein-protein interactions[33-39].

Here, we will develop a new, easy-to-use enhanced sampling method based on the robust Gaussian accelerated molecular dynamics (GaMD) technique for highly efficient and accurate

simulations of peptide-protein interactions. GaMD works by adding a harmonic boost potential to smooth biomolecular potential energy[40]. It greatly reduces system energy barriers and accelerates biomolecular simulations by orders of magnitude[40-43]. GaMD provides unconstrained enhanced sampling without the requirement of predefined reaction coordinates or CVs. Compared with enhanced sampling methods that rely on careful selection of CVs, GaMD could be much easier to use for studying complex biological processes such as peptide binding to proteins[39]. Moreover, because the boost potential follows a Gaussian distribution, biomolecular free energy profiles can be properly recovered through cumulant expansion to the second order.[40]

Based on GaMD, we have recently developed a new ligand GaMD or "LiGaMD" method, which allows us to characterize both thermodynamics and kinetics of ligand binding quantitatively[44]. In LiGaMD, we selectively boost the ligand non-bonded interaction potential energy to enable ligand dissociation. Another boost potential is applied to the remaining potential energy of the entire system in a dual-boost algorithm to facilitate ligand rebinding. LiGaMD has been demonstrated on host-guest and protein-ligand binding model systems. Repetitive guest binding and unbinding in the β-cyclodextrin host were observed in hundreds-of-nanosecond LiGaMD simulations. The calculated guest binding free energies agreed excellently with experimental data with < 1.0 kcal/mol errors. Compared with converged microsecond-timescale cMD simulations, the sampling errors of LiGaMD_Dual simulations were also < 1.0 kcal/mol. Accelerations of ligand kinetic rate constants in LiGaMD simulations were properly estimated using Kramers' rate theory. Furthermore, LiGaMD allowed us to capture repetitive dissociation and binding of the benzamidine inhibitor in trypsin within 1 μs simulations. The calculated ligand binding free energy and kinetic rate constants compared very well with the experimental data[44].

Building upon GaMD and LiGaMD, we present a new computational method called peptide GaMD ("Pep-GaMD"), which enables us to simulate repetitive peptide binding and unbinding within microsecond simulations. In Pep-GaMD, the essential interaction potential energy of the peptide is selectively boosted to enable peptide dissociation and large conformational changes of the peptide. Selective acceleration has been found useful in previous enhanced sampling techniques, including the LiGaMD[44], selective aMD[45], selectively scaled MD[46] and essential energy space random walk[47, 48]. In addition, another boost potential is applied on the protein and solvent in a dual-boost Pep-GaMD algorithm to facilitate peptide rebinding. To demonstrate the new Pep-GaMD method, the SH3 domains with important biological functions[49, 50] is selected as our model system. Pep-GaMD has been testing on the binding of three PRM peptides, including "PAMPAR" (PDB: 1SSH), "PPPALPPKK" (PDB: 1CKA) and "PPPVPPRR" (PDB: 1CKB). Repetitive binding and unbinding of the three peptides to the SH3 domains have been observed in microsecond Pep-GaMD simulations. The peptide binding free energies calculated from Pep-GaMD simulates agree very well with available experimental data. Furthermore, the Pep-GaMD simulations have provided important insights into the binding mechanism of peptide to their target proteins at an atomistic level.

Methods

Peptide Gaussian accelerated molecular dynamics (Pep-GaMD)

Gaussian accelerated molecular dynamics (GaMD) is an enhanced sampling technique that works by adding a harmonic boost potential to smooth biomolecular potential energy surface and reduce the system energy barriers[40]. Details of the GaMD method have been described in previous

studies[40-42]. A brief summary is provided in **Supplementary Material**. Here, we developed a new

Pep-GaMD method for more efficient sampling of peptide binding to proteins.

We consider a system of peptide L binding to a protein P in a biological environment E. The system comprises of N atoms with their coordinates $r \equiv \{\vec{r}_1, \cdots, \vec{r}_N\}$ and momenta $p \equiv \{\vec{p}_1, \cdots, \vec{p}_N\}$. The system Hamiltonian can be expressed as:

$$H(r,p) = K(p) + V(r), \qquad (1)$$

where $K(p)$ and $V(r)$ are the system kinetic and total potential energies, respectively. Next, we decompose the potential energy into the following terms:

$$V(r) = V_{P,b}(r_P) + V_{L,b}(r_L) + V_{E,b}(r_E)$$

$$+ V_{PP,nb}(r_P) + V_{LL,nb}(r_L) + V_{EE,nb}(r_E)$$

$$+ V_{PL,nb}(r_{PL}) + V_{PE,nb}(r_{PE}) + V_{LE,nb}(r_{LE}), \qquad (2)$$

where $V_{P,b}$, $V_{L,b}$ and $V_{E,b}$ are the bonded potential energies in protein P, peptide L and environment E, respectively. $V_{PP,nb}$, $V_{LL,nb}$ and $V_{EE,nb}$ are the self non-bonded potential energies in protein P, peptide L and environment E, respectively. $V_{PL,nb}$, $V_{PE,nb}$ and $V_{LE,nb}$ are the non-bonded interaction energies between P-L, P-E and L-E, respectively. According to classical molecular mechanics force fields[51, 52], the non-bonded potential energies are usually calculated as:

$$V_{nb} = V_{elec} + V_{vdW}, \qquad (3)$$

where V_{elec} and V_{vdW} are the system electrostatic and van der Waals potential energies. Presumably, peptide binding mainly involves in both the bonded and non-bonded interaction energies of the peptide since peptides often undergo large conformational changes during binding to the target proteins. Thus, the essential peptide potential energy is $V_L(r) = V_{LL,b}(r_L) +$

$V_{LL,nb}(r_L) + V_{PL,nb}(r_{PL}) + V_{LE,nb}(r_{LE})$. In Pep-GaMD, we add boost potential selectively to the

essential peptide potential energy according to the GaMD algorithm:

$$\Delta V_L(r) = \begin{cases} \frac{1}{2}k_L\big(E_L - V_L(r)\big)^2, & V_L(r) < E_L \\ \qquad\qquad 0, & V_L(r) \geq E_L, \end{cases} \tag{4}$$

where E_L is the threshold energy for applying boost potential and k_L is the harmonic constant. The

Pep-GaMD simulation parameters are derived similarly as in the previous GaMD algorithm (see

details in **Supplementary Material**). When E is set to the lower bound as the system maximum

potential energy ($E=V_{max}$), the effective harmonic force constant k_0 can be calculated as:

$$k_0 = \min(1.0, k_0') = \min\left(1.0, \frac{\sigma_0}{\sigma_V}\frac{V_{max}-V_{min}}{V_{max}-V_{avg}}\right), \tag{5}$$

where $V_{max}, V_{min}, V_{avg}$ and σ_V are the maximum, minimum, average and standard deviation of the

boosted system potential energy, and σ_0 is the user-specified upper limit of the standard deviation

of ΔV (e.g., $10k_BT$) for proper reweighting. The harmonic constant is calculated as $k = k_0 \cdot$

$\frac{1}{V_{max}-V_{min}}$ with $0 < k_0 \leq 1$. Alternatively, when the threshold energy E is set to its upper bound

$E = V_{min} + \frac{1}{k}$, k_0 is set to:

$$k_0 = k_0'' \equiv \left(1 - \frac{\sigma_0}{\sigma_V}\right)\frac{V_{max}-V_{min}}{V_{avg}-V_{min}}, \tag{6}$$

if k_0'' is found to be between 0 and 1. Otherwise, k_0 is calculated using Eqn. (5).

In addition to selectively boosting the peptide, another boost potential is applied on the

protein and solvent to enhance conformational sampling of the protein and facilitate peptide

rebinding. The second boost potential is calculated using the total system potential energy other

than the essential peptide potential energy as:

$$\Delta V_D(r) = \begin{cases} \frac{1}{2}k_D\big(E_D - V_D(r)\big)^2, & V_D(r) < E_D \\ 0, & V_D(r) \geq E_D \end{cases} \qquad (7)$$

Where V_D is the total system potential energy other than the essential peptide potential energy, E_D is the corresponding threshold energy for applying the second boost potential and k_D is the harmonic constant. This leads to dual-boost Pep-GaMD with the total boost potential $\Delta V(r) = \Delta V_L(r) + \Delta V_D(r)$. Pep-GaMD is currently implemented in the GPU version of AMBER 20[53], but should be transferable to other MD programs as well (see **Supplementary Material**).

Peptide binding free energy calculations from 3D potential of mean force

We calculate peptide binding free energy from 3D potential of mean force (PMF) of peptide displacements from the target protein as the following[54, 55]:

$$\Delta G^0 = -\Delta W_{3D} - RTLn\frac{V_b}{V_0}, \qquad (9)$$

where V_0 is the standard volume, $V_b = \int_b e^{-\beta W(r)}\,dr$ is the average sampled bound volume of the peptide with $\beta = 1/k_B T$, k_B is the Boltzmann constant, T is the temperature, and ΔW_{3D} is the depth of the 3D PMF. ΔW_{3D} can be calculated by integrating Boltzmann distribution of the 3D PMF $W(r)$ over all system coordinates except the x, y, z of the peptide:

$$\Delta W_{3D} = -RTLn\frac{\int_u e^{-\beta W(r)}dr}{\int_u dr}, \qquad (10)$$

where $V_u = \int_u dr$ is the sampled unbound volume of the peptide. The exact definitions of the bound and unbound volumes V_b and V_u are not important as the exponential average cut off contributions far away from the PMF minima[55]. A python script "PyReweighting-3D.py" in the *PyReweighting* tool kit (http://miao.compbio.ku.edu/PyReweighting/)[44, 56] was applied for

reweighting Pep-GaMD simulations (see Methods in **Supplementary Material**) to calculate the 3D reweighted PMF and associated peptide binding free energies.

Protein-peptide binding simulations

Pep-GaMD simulations using the dual-boost scheme were performed on binding of three model peptides to the SH3 domains. X-ray crystal structures of the SH3 domains bound by "PAMPAR" (PDB: 1SSH), "PPPALPPKK" (PDB: 1CKA)[57] and "PPPVPPRR" (PDB: 1CKB)[57] were used. The N- and C-termini of the peptides were capped with the acetyl (ACE) and primary amide (NHE) neutral groups, respectively. The missing hydrogen atoms were added using the *tleap* module in AMBER[58]. The AMBER ff14SB force field[59] were used for the peptides and protein. Each system was neutralized by adding counter ions and immersed in a cubic TIP3P water box[60], which was extended 13 Å from the protein-peptide complex surface.

Each simulation system was first energy minimized with 1.0 kcal/mol/Å^2 constraints on the heavy atoms of the protein and peptide, including the steepest descent minimization for 50,000 steps and conjugate gradient minimization for 50,000 steps. The system was then heated from 0 K to 300 K for 200 ps. It was further equilibrated using the NVT ensemble at 300K for 200 ps and the NPT ensemble at 300 K and 1 bar for 1 ns with 1 kcal/mol/Å^2 constraints on the heavy atoms of the protein and peptide, followed by 2 ns short cMD without any constraint. The Pep-GaMD simulations proceeded with 14 ns short cMD to collect the potential statistics, 46 ns Pep-GaMD equilibration after adding the boost potential and then three independent 1000 ns production runs.

It provided more powerful sampling to set the threshold energy for applying the boost potential to the upper bound (i.e., $E = V_{min}+1/k$) in our previous study ligand dissociation and binding using LiGaMD[44]. Therefore, the threshold energy for applying the peptide boost potential was also set to the upper bound in the Pep-GaMD simulations. For the second boost potential

applied on the system total potential energy other than the essential peptide potential energy, sufficient acceleration was obtained by setting the threshold energy to the lower bound. In order to observe peptide dissociation during Pep-GaMD equilibration while keeping the boost potential as low as possible for accurate energetic reweighting, the $(\sigma_{0P}, \sigma_{0D})$ parameters were finally set to (2.2 kcal/mol, 6.0 kcal/mol), (4.0 kcal/mol, 6.0 kcal/mol) and (4.0 kcal/mol, 6.0 kcal/mol) for the Pep-GaMD simulations of the 1SSH, 1CKA and 1CKB systems, respectively. Pep-GaMD production simulation frames were saved every 0.2 ps for analysis.

The VMD[61] and CPPTRAJ[62] tools were used for simulation analysis. The 1D, 2D and 3D PMF profiles, as well as the peptide binding free energy, were calculated through energetic reweighting of the Pep-GaMD simulations. Root-mean square deviations (RMSDs) of the peptides relative to X-ray structures with the protein aligned, the peptide radius of gyration (R_g), distances between two protein residues that are located in the peptide-binding sites (denoted d_{D-W} for Asp19-Trp40 in the 1SSH structure and Asp150-Trp169 in the 1CKA and 1CKB structures) and a salt bridge distance between the protein and peptides (denoted $d_{D-R/K}$ for Asp19 (SH3)-Arg10 (peptide) in the 1SSH structure, Asp150 (SH3)-Lys8 (peptide) in the 1CKA structure and Asp150 (SH3)-Arg7 (peptide) in the 1CKB structure) were chosen as reaction coordinates for calculating the PMF profiles. The RMSD of peptides with protein aligned was chosen as reaction coordinate for calculating the 1D PMF profile. The bin size was set to 1.0 Å. 2D PMF profiles of d_{D-W}/$d_{D-R/K}$/R_g and peptide backbone RMSD were calculated to analyze conformational changes of the protein and important interactions upon peptide binding. The bin size was set to 1.0 Å for these reaction coordinates. The cutoff for the number of simulation frames in one bin was set to 500 for reweighting 1D and 2D PMF profiles. The 3D PMF profiles of peptide center-of-mass displacements from the SH3 domains in the X, Y and Z directions were further calculated from

the Pep-GaMD simulations. The bin sizes were set to 1.0 Å in the X, Y and Z directions. The cutoff of simulation frames in one bin for 3D PMF reweighting was set to the minimum number below which the calculated global minimum of 3D PMF will be shifted. The peptide binding free energies (ΔG) were calculated using the reweighted 3D PMF profiles.

Results and Discussions

Microsecond Pep-GaMD simulations captured repetitive dissociation and binding of peptides to the SH3 domains.

All-atom Pep-GaMD simulations were performed on X-ray crystal structures of the SH3 domains bound by three peptides, including "PAMPAR" (PDB: 1SSH), "PPPALPPKK" (PDB: 1CKA)[57] and "PPPVPPRR" (PDB: 1CKB)[57] (**Figs. 1A-1C**). Three independent 1000 ns Pep-GaMD production trajectories were obtained on each of the three peptide systems (**Table 1**). The Pep-GaMD simulations of the 1SSH system recorded an average boost potential of 12.83-13.01 kcal/mol with 3.75-3.80 kcal/mol standard deviation. In comparison, the average boost potential was 25.59-26.56 kcal/mol with 4.91-4.95 kcal/mol standard deviation in three simulations of the 1CKA system. The boost potential applied in simulations of the 1CKB system was similar to that of the 1CKA system, with average of 27.85-28.30 kcal/mol and 4.96-5.14 kcal/mol standard deviation (**Table 1**).

The peptide backbone RMSDs with the SH3 protein domain aligned were calculated as a function of simulation time (**Figs. 1D-1F**) to record the number of peptide dissociation and binding events (N_D and N_B) in each of the 1 μs Pep-GaMD simulations. With close examination of the peptide binding trajectories, cutoffs of the peptide backbone RMSD for the unbound and bound states were set to >20 Å and <3.5 Å, respectively. Repetitive dissociation and binding of the three

peptides were successfully captured in each of the 1 μs Pep-GaMD simulation (**Figs. 1D-1F**). In each simulation of the 1SSH system, about 3-4 binding and 3 dissociation events were observed for the peptide. In simulations of the 1CKA system, about 2-5 dissociation and 3-5 binding events were observed. A slightly greater number of peptide dissociation (3-5) and binding (4-5) events were observed in simulations of the 1CKB system (**Table 1**). The improved sampling appeared to correlate with relatively higher boost potential applied in the 1CKB system simulations, as well as lower peptide binding affinity in the 1CKB system (5.2 μM) than that in the 1CKA system (1.9 μM)[57].

Next, we computed 1D PMF free energy profiles from the Pep-GaMD simulations to quantitatively characterize peptide binding to the SH3 domains. The RMSD of each peptide relative to the native X-ray structure with the protein domain aligned was chosen as the reaction coordinate. The PMF profiles were calculated for three individual 1 μs Pep-GaMD simulations and averaged for each system (**Figs. 1G-1I**). The global free energy minima were all identified in the "Bound" state with ~2 Å peptide backbone RMSD. Error bars (standard deviations) of free energy values in the PMF profiles were mostly <1.0 kcal/mol, suggesting reasonable convergence of the Pep-GaMD simulations. Notably, differences were identified among the three peptide binding systems in the magnitudes of PMF values near the "Intermediate", "Unbound" and energy barrier regions. The energy barriers for peptide dissociation were approximately 6.0 kcal/mol, 7.1 kcal/mol and 6.5 kcal/mol in the 1SSH, 1CKA and 1CKB system simulations, respectively. Therefore, the energy barrier for peptide dissociation in the 1CKA system was the highest, followed by the 1CKB and then 1SSH systems. The "Intermediate" low-energy wells were identified for the 1SSH system with the peptide backbone RMSD centered around ~12.1 Å (**Fig. 1G**) and ~15.0 Å peptide backbone RMSD for the 1CKA (**Fig. 1H**). In comparison, no clear low-

energy well was identified in the "Intermediate" state for the 1CKB system, which showed rather a plateau in the region of 7.0 Å-15.0 Å peptide backbone RMSD (**Fig. 1I**). PMF values of the "Unbound" state relative to the "Bound" state were approximately 8.0 kcal/mol, 7.1 kcal/mol and 7.5 kcal/mol for three peptides in the 1SSH, 1CKA and 1CKB systems, respectively.

Peptide binding free energies calculated from Pep-GaMD simulations agreed well with available experimental data

We computed binding free energies of three peptides to the SH3 domains based on their 3D PMF profiles (see details in **Methods**). The 3D PMF was calculated from each individual 1 μs Pep-GaMD simulation of peptide binding to the SH3 domain in terms of displacements of the Cα atom in the central peptide proline residue (Pro8 in the 1SSH structure, Pro6 in the 1CKA structure and Pro5 in the 1CKB structure) from the Cα atom in a protein residue (Asn55 in the 1SSH structure, Pro185 in the 1CKA and 1CKB structures) in the X, Y and Z directions. The 3D PMF was energetically reweighted through cumulant expansion to the second order (**Supplementary Material**). We then calculated the peptide binding free energies using the 3D reweighted PMF profiles. For the 1CKA system, the average of calculated peptide binding free energy values was -7.72 kcal/mol and the standard deviation was 0.54 kcal/mol, which was highly consistent with the experimental value of -7.84 kcal/mol[57] (**Table 1**). For the 1CKB system, the resulting binding free energy was -6.84±0.14 kcal/mol, being closely similar to the experimental value of -7.24 kcal/mol[57] (**Table 1**). In this context, not only the relative ranks of the peptide binding free energies were consistent with experimental data, the absolute binding free energy values calculated from Pep-GaMD simulations were also in excellent agreements with those from the experiments. The differences between the simulation predicted and experimentally determined binding free energies

were less than 1.0 kcal/mol. For the 1SSH system, the calculated peptide binding free energy was -7.41±0.12 kcal/mol. However, experimental peptide binding free energy of this system was not available in the literature for comparison. According to the simulation prediction, the peptide binding free energy in the 1SSH system was between those of the 1CKA and 1CKB systems.

In summary, peptide binding free energies in the SH3 domains calculated from microsecond Pep-GaMD simulations agreed well with available experimental data. Compared with the experimental values, errors in the peptide binding free energies predicted from the Pep-GaMD simulations were smaller than 1.0 kcal/mol. Therefore, both efficient enhanced sampling and accurate free energy calculations of peptide binding were achieved through the Pep-GaMD simulations.

Peptide binding to the SH3 domains was mediated by electrostatic interactions

Next, we examined key residue interactions during peptide binding to the SH3 domains (**Fig. 2**). In the X-ray structures (**Figs. 1A-1C**), a conserved salt-bridge was found between the SH3 domains and bound peptides, i.e., SH3:Asp19 – peptide:Arg10 in the 1SSH structure, SH3:Asp150 – peptide:Lys8 in the 1CKA structure and SH3:Asp150 – peptide:Arg7 in the 1CKB structure. Therefore, the salt-bridge distance between these protein and peptide residues (denoted $d_{D-R/K}$) and the peptide backbone RMSD relative to the X-ray structure with the protein aligned were chosen as reaction coordinates to compute 2D PMF profiles. Two low-energy "Bound" and "Intermediate" states were identified from the 2D PMF profiles of all three systems (**Figs. 2A-2C**). In the 1SSH system, (peptide backbone RMSD, $d_{D-R/K}$) of the "Bound" and "Intermediate" states were centered around (2.0 Å, 4.2 Å) and (20.0 Å, 5.0 Å), respectively (**Fig. 2A**). In the 1CKA system, (peptide backbone RMSD, $d_{D-R/K}$) of the two states were centered around (2.7 Å, 4.4 Å) and (21.5 Å, 5.2Å) (**Fig. 2B**). In the 1CKB system, (peptide backbone RMSD, $d_{D-R/K}$) of the two states were centered

around (1.7 Å, 3.7 Å) and (25.0 Å, 8.0 Å) (**Fig. 2C**). Therefore, the salt bridge distances in the "Intermediate" state of the 1SSH and 1CKA structures were similar to those in their "Bound" state, but the salt-bridge distance increased to 8.0 Å in the "Intermediate" state of the 1CKB structure.

Figs. **2D-2F** depict the "Intermediate" conformations of the three peptide systems compared with their native "Bound" conformations. While the salt bridge became partially open in the "Intermediate" conformation of the 1CKB system (**Fig. 2F**), it remained fully closed in the "Intermediate" conformations of the 1SSH and 1CKA systems (**Figs. 2D-2E**). In summary, long-range electrostatic interactions played a key role in peptide binding to the SH3 domains. The salt bridge described here appeared to be an anchor that pulled the peptides to their target binding site in the SH3 domains. Similar findings were obtained from previous cMD simulations[19].

Peptides bound to the SH3 domains via conformational selection

In order to further explore the mechanism of peptide binding to the SH3 domains, we computed 2D PMF free energy profiles to characterize conformational changes of the three systems upon peptide binding. By closely examining the Pep-GaMD simulation trajectories, we observed the largest conformational changes in protein residue Trp40 in the 1SSH structure or residue Trp169 in the 1CKA and 1CKB structures, which was located at the peptide binding site of the SH3 domains (**Figs. 2D-2F**). Therefore, we calculated 2D reweighted PMF profiles from the Pep-GaMD simulations regarding the peptide backbone RMSD relative to the X-ray structure with the protein aligned and the $d_{\text{D-W}}$ distance between Asp19-Trp40 in the 1SSH structure or Asp150-Trp169 in the 1CKA and 1CKB structures (**Figs. 3A-3C**). For the 1SSH system, two low-energy states were identified from the 2D PMF profile, including the "Bound" and "Intermediate", in which the protein residue Trp40 at the peptide-binding site adopted primarily the "Closed" and

"Open" conformations, respectively (**Figs. 3A** and **2D**). The peptide backbone RMSD and $d_{\text{D-W}}$ were centered around (2.0 Å, 6.2 Å) in the "Bound/Closed" state and (9.8 Å, 9.0 Å) in "Intermediate/Open" state (**Fig. 3A**). For the 1CKA system, three low-energy states were identified from the 2D PMF profile, including the "Bound", "Intermediate" and "Unbound", in which the protein residue Trp169 adopted primarily the "Closed", "Open" and "Open" conformations, respectively (**Figs. 3B** and **2E**). The peptide backbone RMSD and $d_{\text{D-W}}$ were centered around (2.7 Å, 5.9 Å), (13.5 Å, 9.3 Å) and (33.5 Å, 9.3 Å) in the "Bound/Closed", "Intermediate/Open" and "Unbound/Open" states, respectively (**Fig. 3B**). Three similar low-energy conformational states were identified in the 1CKB system as in the 1CKA system. The peptide backbone RMSD and $d_{\text{D-W}}$ were centered around (1.7 Å, 5.8 Å), (11.5 Å, 9.3 Å) and (36.0 Å, 8.8 Å) in the "Bound/Closed", "Intermediate/Open" and "Unbound/Open" states, respectively (**Fig. 3C**).

In addition to the target protein, we analyzed conformational changes of the peptides during binding. In this regard, the radius of gyration (R_g) was calculated and monitored for possible conformational changes and even folding and unfolding of the peptides. Then the peptide R_g and backbone RMSDs relative to X-ray structures with SH3 domain aligned were used as reaction coordinates to calculate 2D PMF profiles. From the reweighted 2D PMF profiles (**Fig. 3D-3F**), we identified a clear low-energy "Bound" state in all three systems, for which the peptide backbone RMSD and R_g in the 1SSH, 1CKA and 1CKB structures were centered around (2.0 Å, 5.9 Å), (2.7 Å, 7.9 Å) and (1.7 Å, 7.5 Å), respectively. Another "Intermediate" low-energy state was also identified in the 1SSH, 1CKA and 1CKB systems, for which the peptide backbone RMSD and R_g were centered around (11.2 Å, 4.7 Å), (8.5 Å, 5.6 Å) and (8.8 Å, 5.6 Å), respectively. Furthermore, a low energy "Unbound" state appeared in the reweighted 2D PMF profiles of the 1CKA (**Fig. 3E**)

and 1CKB (**Fig. 3F**), but not in the reweighted 2D PMF profile of the 1SSH system (**Fig. 3A**). Nevertheless, the "Unbound" state was found in the 2D PMF profiles without energetic reweighting for all three systems, similarly for the "Bound" and "Intermediate" states (**Fig. S1**). This suggested that complete peptide binding processes were successfully sampled in the Pep-GaMD simulations, despite differences in the free energy profiles. More importantly, the peptides sampled a large conformational space with a wide range of R_g in the "Unbound" state, but only a subset of the conformations were chosen upon binding to the target protein. Therefore, peptide binding to the SH3 domains followed predominantly a "conformational selection" mechanism.

Conclusions

A new computational method called Pep-GaMD has been developed for efficient enhanced sampling and free energy calculations of peptide binding. Pep-GaMD works by selectively boosting the essential interaction potential energy of the peptide. Microsecond Pep-GaMD simulations have allowed us to capture repetitive peptide dissociation and binding as demonstrated on binding of three model peptides to the SH3 domains. Pep-GaMD has thus enabled peptide binding free energy calculations with significantly shorter simulations compared with cMD. Without the requirement to set predefined CVs, Pep-GaMD provides a very easy-to-use technique for unconstrained enhanced sampling of peptide binding, by which the peptide binding free energy can be calculated from 3D potential of mean force through a novel physics-based approach.

Pep-GaMD shall be of wide applicability for studies of many other peptide-protein interactions[1] other than peptide binding to the SH3 domains. Beyond the peptide binding free energies (thermodynamics), we shall derive the dissociation and binding kinetic rates of peptide binding to the proteins in follow-up studies. To facilitate peptide rebinding, one can include

multiple peptides in the system within their solubility limits, an approach that has been shown to accelerate ligand binding in the recently developed LiGaMD method[44]. These developments are expected to further improve the Pep-GaMD method for applications in enhanced sampling of peptide-protein interactions.

Acknowledgements

We appreciate the help of Prof. David Case for accessing the AMBER git repository to develop our new simulation algorithms. This work used supercomputing resources with allocation award TG-MCB180049 through the Extreme Science and Engineering Discovery Environment (XSEDE), which is supported by National Science Foundation grant number ACI-1548562, and project M2874 through the National Energy Research Scientific Computing Center (NERSC), which is a U.S. Department of Energy Office of Science User Facility operated under Contract No. DE-AC02-05CH11231, and the Research Computing Cluster at the University of Kansas. This work was supported in part by the National Institutes of Health (R01GM132572) and the startup funding in the College of Liberal Arts and Sciences at the University of Kansas.

Supplementary Material

Detailed description of the GaMD and energetic reweighting methods and **Fig. S1** are provided in the **Supplementary Material**.

Data Availability Statement

The data that supports the findings of this study are available within the article and its supplementary material. The all-atom simulation files are available from the corresponding author upon reasonable request.

References

[1] E. Petsalaki, and R. B. Russell, Curr. Opin. Biotechnol. **19** (2008) 344.

[2] A. A. Das, O. P. Sharma, M. S. Kumar, R. Krishna, and P. P. Mathur, Genomics, Proteomics Bioinf. **11** (2013) 241.

[3] P. M. Watt, Nat. Biotechnol. **24** (2006) 177.

[4] L. T. Vassilev, B. T. Vu, B. Graves, D. Carvajal, F. Podlaski, Z. Filipovic, N. Kong, U. Kammlott, C. Lukacs, C. Klein, N. Fotouhi, and E. A. Liu, Science **303** (2004) 844.

[5] T. Matthews, M. Salgo, M. Greenberg, J. Chung, R. DeMasi, and D. Bolognesi, Nat. Rev. Drug Discovery **3** (2004) 215.

[6] V. M. Ahrens, K. Bellmann-Sickert, and A. G. Beck-Sickinger, Future Med. Chem. **4** (2012) 1567.

[7] K. Fosgerau, and T. Hoffmann, Drug Discov. Today **20** (2015) 122.

[8] U. Kahler, J. E. Fuchs, P. Goettig, and K. R. Liedl, J. Biomol. Struct. Dyn. **36** (2018) 4072.

[9] A. C. Lee, J. L. Harris, K. K. Khanna, and J. H. Hong, Int. J. Mol. Sci. **20** (2019) 2383.

[10] D. Kilburg, and E. Gallicchio, in *Adv. Protein Chem. Struct. Biol.*, edited by C. Z. Christov (Academic Press, 2016), pp. 27.

[11] M. Ciemny, M. Kurcinski, K. Kamel, A. Kolinski, N. Alam, O. Schueler-Furman, and S. Kmiecik, Drug Discov. Today **23** (2018) 1530.

[12] G. Weng, J. Gao, Z. Wang, E. Wang, X. Hu, X. Yao, D. Cao, and T. Hou, J. Chem. Theory Comput. **16** (2020) 3959.

[13] M. Karplus, and J. A. McCammon, Nat Struct Biol **9** (2002) 646.

[14] I. Antes, Proteins **78** (2010) 1084.

[15] S. J. de Vries, J. Rey, C. E. M. Schindler, M. Zacharias, and P. Tuffery, Nucleic Acids Res. **45** (2017) W361.

[16] C. E. M. Schindler, S. J. de Vries, and M. Zacharias, Structure **23** (2015) 1507.

[17] A. Ben-Shimon, and M. Y. Niv, Structure **23** (2015) 929.

[18] J. Wang, A. Alekseenko, D. Kozakov, and Y. Miao, Front. Mol. Biosci. **6** (2019) 112.

[19] M. Ahmad, W. Gu, and V. Helms, Angew. Chem., Int. Ed. Engl. **47** (2008) 7626.

[20] B. Knapp, S. Demharter, R. Esmaielbeiki, and C. M. Deane, Brief Bioinform. **16** (2015) 1035.

[21] S. Wan, B. Knapp, D. W. Wright, C. M. Deane, and P. V. Coveney, J. Chem. Theory Comput. **11** (2015) 3346.

[22] J. Chen, J. Wang, Q. Zhang, K. Chen, and W. Zhu, Sci Rep-Uk **5** (2015) 17421.

[23] M. C. Zwier, A. J. Pratt, J. L. Adelman, J. W. Kaus, D. M. Zuckerman, and L. T. Chong, J. Phys. Chem. Lett. **7** (2016) 3440.

[24] V. Salmaso, M. Sturlese, A. Cuzzolin, and S. Moro, Structure **25** (2017) 655.

[25] S. Yadahalli, J. Li, D. P. Lane, S. Gosavi, and C. S. Verma, Sci Rep-Uk **7** (2017) 15600.

[26] P. Robustelli, S. Piana, and D. E. Shaw, J. Am. Chem. Soc. **142** (2020) 11092.

[27] F. Paul, C. Wehmeyer, E. T. Abualrous, H. Wu, M. D. Crabtree, J. Schöneberg, J. Clarke, C. Freund, T. R. Weikl, and F. Noé, Nat. Commun. **8** (2017) 1095.

[28] M. A. Cuendet, V. Zoete, and O. Michielin, Proteins **79** (2011) 3007.

[29] J. A. Morrone, A. Perez, J. MacCallum, and K. A. Dill, J. Chem. Theory Comput. **13** (2017) 870.

[30] J. A. Morrone, A. Perez, Q. Deng, S. N. Ha, M. K. Holloway, T. K. Sawyer, B. S. Sherborne, F. K. Brown, and K. A. Dill, J. Chem. Theory Comput. **13** (2017) 863.

[31] G. Lamothe, and T. E. Malliavin, Bmc Struct Biol **18** (2018) 4.

[32] R. Zou, Y. Zhou, Y. Wang, G. Kuang, H. Agren, J. Wu, and Y. Tu, J. Chem. Inf. Model **60** (2020) 1551.

[33] C. Abrams, and G. Bussi, Entropy-Switz **16** (2014) 163.

[34] V. Spiwok, Z. Sucur, and P. Hosek, Biotechnol. Adv. **33** (2015) 1130.

[35] C. Dellago, and P. G. Bolhuis, in *Advanced Computer Simulation Approaches for Soft Matter Sciences III*, edited by C. Holm, and K. Kremer (Springer Berlin Heidelberg, Berlin, Heidelberg, 2009), pp. 167.

[36] Y. Q. Gao, L. J. Yang, Y. B. Fan, and Q. Shao, Int Rev Phys Chem **27** (2008) 201.

[37] A. Liwo, C. Czaplewski, S. Oldziej, and H. A. Scheraga, Curr. Opin. Struct. Biol. **18** (2008) 134.

[38] M. Christen, and W. F. van Gunsteren, J. Comput. Chem. **29** (2008) 157.

[39] Y. Miao, and J. A. McCammon, Mol. Simul. **42** (2016) 1046.

[40] Y. Miao, V. A. Feher, and J. A. McCammon, J. Chem. Theory Comput. **11** (2015) 3584.

[41] Y. T. Pang, Y. Miao, Y. Wang, and J. A. McCammon, J. Chem. Theory Comput. **13** (2017) 9.

[42] Y. Miao, and J. A. McCammon, Annu. Rep. Comput. Chem. **13** (2017) 231.

[43] Y. Miao, J. Chem. Phys. **149** (2018) 072308.

[44] Y. Miao, A. Bhattarai, and J. Wang, bioRxiv (2020)

[45] J. Wereszczynski, and J. A. McCammon, J. Chem. Theory Comput. **6** (2010) 3285.

[46] I. Deb, and A. T. Frank, J. Chem. Theory Comput. **15** (2019) 5817.

[47] L. Zheng, and W. Yang, J. Chem. Phys. **129** (2008) 014105.

[48] H. Li, D. Min, Y. Liu, and W. Yang, J. Chem. Phys. **127** (2007) 094101.

[49] M. Ahmad, and V. Helms, Chem. Cent. J. **3** (2009) O22.

[50] L. J. Ball, R. Kühne, J. Schneider-Mergener, and H. Oschkinat, Angew. Chem., Int. Ed. Engl. **44** (2005) 2852.

[51] K. Vanommeslaeghe, and A. D. MacKerell, Jr., Biochim. Biophys. Acta. **1850** (2015) 861.

[52] Y. Duan, C. Wu, S. Chowdhury, M. C. Lee, G. Xiong, W. Zhang, R. Yang, P. Cieplak, R. Luo, T. Lee, J. Caldwell, J. Wang, and P. Kollman, J. Comput. Chem. **24** (2003) 1999.

[53] D. A. Case, K. Belfon, I. Y. Ben-Shalom, S. R. Brozell, D. S. Cerutti, I. T.E. Cheatham, V. W. D. Cruzeiro, T. A. Darden, R. E. Duke, G. Giambasu, M. K. Gilson, H. Gohlke, A. W. Goetz, R. Harris, S. Izadi, K. K.-. jhala, A. Kovalenko, R. Krasny, T. Kurtzman, T. S. Lee, S. LeGrand, P. Li, C. Lin, J. Liu, T. Luchko, R. Luo, V. Man, K. M. Merz, Y. Miao, O. Mikhailovskii, G. Monard, H. Nguyen, A. Onufriev, F. Pan, S. Pantano, R. Qi, D. R. Roe, A. Roitberg, C. Sagui, S. Schott-Verdugo, J. Shen, C. L. Simmerling, N. Skrynnikov, J. Smith, J. Swails, R. C. Walker, J. Wang, L. Wilson, R. M. Wolf, X. Wu, D. M. York, and P. A. Kollman, (2020)

[54] S. Doudou, N. A. Burton, and R. H. Henchman, J. Chem. Theory Comput. **5** (2009) 909.

[55] I. Buch, T. Giorgino, and G. De Fabritiis, Proc. Natl. Acad. Sci. USA **108** (2011) 10184.

[56] Y. Miao, W. Sinko, L. Pierce, D. Bucher, R. C. Walker, and J. A. McCammon, J. Chem. Theory Comput. **10** (2014) 2677.

[57] X. Wu, B. Knudsen, S. M. Feller, J. Zheng, A. Sali, D. Cowburn, H. Hanafusa, and J. Kuriyan, Structure **3** (1995) 215.

[58] D. A. Case, T. E. Cheatham, 3rd, T. Darden, H. Gohlke, R. Luo, K. M. Merz, Jr., A. Onufriev, C. Simmerling, B. Wang, and R. J. Woods, J. Comput. Chem. **26** (2005) 1668.

[59] J. A. Maier, C. Martinez, K. Kasavajhala, L. Wickstrom, K. E. Hauser, and C. Simmerling, J. Chem. Theory Comput. **11** (2015) 3696.

[60] W. L. Jorgensen, J. Chandrasekhar, J. D. Madura, R. W. Impey, and M. L. Klein, J. Chem. Phys. **79** (1983) 926.

[61] W. Humphrey, A. Dalke, and K. Schulten, J. Mol. Graph. **14** (1996) 33.

[62] D. R. Roe, and T. E. Cheatham, 3rd, J. Chem. Theory Comput. **9** (2013) 3084.

Table 1 Summary of Pep-GaMD simulations performed on binding of three model peptides to the SH3 domains. $\varDelta V$ is the total boost potential. N_D and N_B are the number of observed peptide dissociation and binding events, respectively. $\varDelta G_{sim}$ and $\varDelta G_{exp}$ are the peptide binding free energies obtained from Pep-GaMD simulations and experiments, respectively.

PDB	ID	Length (ns)	$\varDelta V$ (kcal/mol)	N_D	N_B	$\varDelta G_{sim}$ (kcal/mol)	$\varDelta G_{exp}$ (kcal/mol)
1SSH	Sim1	1000	13.01±3.79	3	4	-7.41±0.12	-
	Sim2	1000	12.88±3.75	3	3		
	Sim3	1000	12.83±3.80	3	3		
1CKA	Sim1	1000	25.65±4.91	4	5	-7.72±0.54	-7.84
	Sim2	1000	26.56±4.95	3	3		
	Sim3	1000	25.59±4.89	2	3		
1CKB	Sim1	1000	27.98±5.02	3	4	-6.84±0.14	-7.24
	Sim2	1000	28.30±5.14	5	4		
	Sim3	1000	27.85±4.96	5	5		

Figure Captions

Fig. 1 Pep-GaMD simulations have captured repetitive dissociation and binding of three model peptides to the SH3 domains: (A-C) X-ray structures of the SH3 domains bound by peptides (A) "PAMPAR" (PDB: 1SSH), (B) "PPPALPPKK" (PDB: 1CKA) and (C) "PPPVPPRR" (PDB: 1CKB). The SH3 domains and peptides are shown in green cartoon. Key protein residues Asp19 and Trp40 in the 1SSH structure and Asp150 and Trp169 in the 1CKA and 1CKB structures, and peptide residues Arg10 in the 1SSH structure, Lys8 in the 1CKA structure and Arg7 in the 1CKB structure are highlighted in sticks. The "N" and "C" labels denote the N-terminus and C-terminus of the peptides. (D-F) time courses of peptide backbone RMSDs relative to X-ray structures with the protein aligned calculated from three independent 1 μs Pep-GaMD simulations of the (D) 1SSH, (E) 1CKA and (F) 1CKB structures. (G-I) The corresponding PMF profiles of the peptide backbone RMSDs averaged over three Pep-GaMD simulations of the (G) 1SSH, (H) 1CKA and (I) 1CKB structures. Error bars are standard deviations of the free energy values calculated from three Pep-GaMD simulations.

Fig. 2 Free energy profiles and low-energy conformational states of peptide binding to the SH3 domains: (A) 2D PMF profiles regarding the peptide backbone RMSD and the distance between protein Asp19 and peptide Arg10 in the 1SSH structure. (B) 2D PMF profiles regarding the peptide backbone RMSD and the distance between protein Asp150 and peptide Lys8 in the 1CKA structure. (C) 2D PMF profiles regarding the peptide backbone RMSD and the distance between protein Asp150 and peptide Arg7 in the 1CKB structure. (D-F) Low-energy intermediate conformations (red) as identified from the 2D PMF profiles of the (D) 1SSH, (E) 1CKA and (F) 1CKB structures, respectively. X-ray structures of the peptide-bound complexes are shown in green. Protein residues Asp19 and Trp40 in the 1SSH structure and Asp150 and Trp169 in the

1CKA and 1CKB structures, and peptide residues Arg10 in the 1SSH structure, Lys8 in the 1CKA structure and Arg7 in the 1CKB structure are highlighted in sticks. The "N" and "C" labels denote the N-terminus and C-terminus of the peptides.

Fig. 3. (A) 2D PMF profiles of peptide backbone RMSD and the distance between protein Asp19 and Trp40 calculated from Pep-GaMD simulations of the 1SSH structure. (B-C) 2D PMF profiles of the peptide backbone RMSD and distance between protein Asp150 and Trp169 calculated from Pep-GaMD simulations of the (B) 1CKA and (C) 1CKB structures. (D-F) 2D PMF profiles regarding the peptide backbone RMSD and peptide R_g in the (D) 1SSH, (E) 1CKA and (F) 1CKB structures.

Fig. 1

Fig. 2

Fig. 3