1   HIGH-RESOLUTION CHARACTERIZATION OF MALE ORNAMENTATION AND

2   REEVALUATION OF SEX LINKAGE IN GUPPIES

3   **Jake Morris[1*], Iulia Darolti[2], Wouter van der Bijl[2], Judith E. Mank[1,2]**
4

5   [1] Department of Genetics, Evolution and Environment, University College London, London, United Kingdom

6   [2] Department of Zoology and Biodiversity Research Centre, University of British Columbia, Canada

8   **Keywords:** Colour; Y-linkage; sex-linked; Sexual dimorphism; sexual selection.

Abstract

Colouration plays a key role in the ecology of many species, influencing how an organism interacts with its environment, other species and conspecifics. Guppies are sexually dimorphic, with males displaying sexually selected colouration resulting from female preference. Previous work has suggested that much of guppy colour pattern variation is Y-linked. However, it remains unclear how many individual colour patterns are Y-linked in natural populations as much of the previous work has focused on phenotypes either not found in the wild, or aggregate measures such as total colour area. Moreover, ornaments have traditionally been identified and delineated by hand, and computational methods now make it possible to extract pixels and identify ornaments more automatedly, reducing the potential for human bias. Here we developed a pipeline for automated ornament identification and high-resolution image analysis of male guppy colour patterns and applied it to a multigenerational pedigree. Our results show that loci controlling the presence or absence of individual male ornaments in our population are not predominantly Y-linked. However, we find that ornaments of similar colour are not independent of each other, and modifier loci that affect whole animal colouration appear to be at least partially Y-linked. Considering these results, Y-linkage of individual ornaments may not be important in driving colour changes in natural populations of guppies, or in expansions of the non-recombining Y region, while Y-linked modifier loci that affect aggregate traits may well play an important role.

Introduction

Colouration is an important morphological character in many species. It has the potential to greatly influence how an organism interacts with its environment via thermoregulation, UV protection [1–3], with other species via aposematism, mimicry and crypsis [4–7], and with conspecifics via intra- and intersexual selection [8,9]. Moreover, some colour phenotypes under sexual selection can incur significant metabolic production costs [10,11] or increased predation risks due to increased conspicuousness [12,13]. In these cases, the costs affect both sexes, but only one sex gains a mating advantage, resulting in sexual conflict [14,15]. In these cases, colour traits are likely to be dimorphic when the genetic architecture can be separated between females and males [15].

For species with male heterogametic sex chromosomes, one important way to establish separate male and female genetic architecture, and thereby resolve sexual conflict, is Y-linkage. Moreover, sexual conflict is thought to accelerate the evolution of the Y chromosome and its divergence from the X [16] when a sexually antagonistic allele is in close linkage with the sex-determining (Y) locus. In males, this will lead selection against recombinants to maintain linkage, ultimately resulting in a non-recombining Y chromosome region [17–19]. Theoretical models [16,17] predict an accumulation of sexually antagonistic genes on Y chromosomes, particularly in nascent sex chromosome systems before large-scale gene loss has occurred.

Male guppies (*Poecilia reticulata*) are brightly coloured and highly variable in their patterning [20]. These patterns are composed of spots and bars that can be categorized into three types, orange and red carotenoid and pterin ornaments, melanic black ornaments, and green and

blue structural colours [21,22]. In contrast, females lack any of these ornaments and are

instead a uniform colour. Although colouration leads to increased predation [23], male

guppy colouration is the result of sexual selection, with females showing a preference for

more colourful, brightly patterned males [20,21,24–26]. Guppies face fewer predators in the

upper reaches of the streams and in these more recently colonised upstream populations,

guppies have evolved more conspicuous colouration due to increased female preference

[27,28]. Male guppy colouration is therefore a sexually antagonistic trait, being selectively

disadvantageous via predation if it were found in females, but being sexually advantageous

for males.

Previous work has suggested that a sizable proportion of guppy male colouration is Y-

linked, and a summary by Lindholm and Breden (2002) identified 16 male ornament

characters as fully Y-linked, 24 that recombine between the X and Y and two that are X

linked. Indeed, Y-linkage of male colour traits in this system has been suggested as a main

factor in the evolution of recombination suppression [29] between both the small region of

the sex chromosomes that never recombines, as well as a larger region of the sex

chromosomes where recombination occurs infrequently [30]. However, as Lindholm and

Breden (2002) point out, much of the evidence for this Y-linkage has come from studies on

phenotypes not typically found in the wild [31,32], or from studies that do not or cannot

account for potential for variation in female sensitivity to hormone treatments [32–34],

and/or from studies using aggregate measures of total colour area rather than specific

ornaments [21,34,35]. Moreover, ornaments have traditionally been identified and

delineated by hand, leading to the potential for human bias if samples are not analysed

blind to parentage or place of origin [36].

Here we developed a pipeline for high resolution image analysis of male guppy colour

patterns. We use this pipeline along with two datasets from our outbred stock population

derived from the Quare river system to; i) identify and characterize male colour ornaments

agnostically with computational clustering methods, ii) determine how these ornaments

relate to each other, and iii) use pedigreed families to compare and contrast inheritance

patterns and possible Y-linkage of both individual ornament phenotypes and aggregate

pattern phenotypes of total colour area.

Methods

Our stock population of guppies originates from the Quare River in Trinidad, originally

collected in 1998 and subsequently maintained in large numbers (>500) in benign freely-

mating conditions [37]. We established pedigrees from this population in 2016 by pairing

four males and virgin females randomly in the first generation. We then paired offspring in

subsequent generations from different families to maximise the number of grandparents in

each brood, to minimize inbreeding which can make it difficult to differentiate different

modes of inheritance, and to help increase the diversity of different phenotypes segregating

in different families. This led to a minimum of six great-grandparents per family in

generation three (see Supplementary Fig. 1). In all, our pedigree stretched across three

generations and 19 crosses, and produced 214 male offspring across generations two and

three (average per family = 14). We allowed between 4 and 8 months for each generation to

permit multiple clutches in most pairs. We maintained both the stock population and

pedigree families on a 12:12 light cycle at 26C and fed a standard hobby guppy diet and

artemia daily.

Imaging

We photographed males from the controlled pedigrees, described above, as well as 171

males taken from the main stock population and not used in pedigrees as a population

baseline for ornamentation. For all males, fish were photographed as soon as possible after

sexual maturation in a specially designed tank (photarium) with internal dimensions of

Length 100mm x Width 100mm x Depth 20mm. Photos were taken with a tripod-mounted

Canon EOS 100D and an 18-55m lens, using ISO 400, shutter speed 1/640 and F4.5. Lighting

was standardized using a Polaroid PLLED312 312 LED light set to 5600K on full power. The

standardized photographic conditions limited the need for pre-processing normalisation,

however, we used the R function histMatch() from the RStoolbox v0.2.3 package [38] to

correct the small amount of variation we did observe in brightness . In addition, we also

used the R function focal() from the Raster v2.6-7 package [39] to slightly blur images for

downstream colour block extraction, thereby eliminating any potential problems with glare

or reflectance.

Following pre-processing, we cropped each photo to a 1801x931 (XY) pixel image using

custom functions. We then employed a k-means clustering algorithm with function kmeans()

from the package Stats v3.5.0 from the R core team [40], which partitions the image based on

the RGB values of pixels into k clusters, so that the sum of squares from pixels to the

assigned cluster RGB centres is minimized. Each image was partitioned using both k=20 and

k=40, so that the image was split into 20 or 40 clusters of different pixels to highlight patches

of same-coloured pixels that could be considered ornaments. k=20 and k=40 were chosen as

these values provided a level of granularity that enabled reasonable extraction of ornaments

across all fish in our dataset. With images now split into *k* clusters of different colours, we

were then able to choose clusters that allowed us to separately extract orange and black

pixels. For both orange and black pixels, we used *k*-means images to reduce the complexity

of the colour on the fish to ease pixel cluster selection for the user. This was done from the *k*-

image that gave the most distinct and defined pixel patterns. We did however extract the

original RGB values from the non *k*-means image.

Following the extraction of orange and black pixels for each fish, we used geometric

morphometric methods to align both main population and pedigree fish against a reference

shape. This method has previously been used in colour analysis R packages such as

Patternize v0.0.1 [41]. We placed 99 semi-landmarks (equally spaced points along a curve)

around the body outlines, ignoring caudal, dorsal and other fins, in the original photos

using the TPS suite (tpsDig 2.17 and tpsUtil 1.56) [42,43]. The gpagen() function from the

Geomorph v3.0.7 package [44] was used to calculate the mean shape of the 171 male fish

from the main population, and we used this as the reference shape against which the images

of extracted pixels from all pedigree males were aligned. This alignment was done using

Thin Plate Spline warping as implemented by the computeTransform() and

applyTransform() functions from the package Morpho v2.6 [45]. See Supplementary Fig. 1

for pipeline overview.

Defining ornaments and ornament inheritance

After aligning the orange and black extracted images of all 171 males from the stock

population, we had RGB scores for each pixel in each ornament in each fish. We then used

these data to define and delimit the male ornaments in our stock population. Pixels were

marked as part of an ornament based on the proportion of presence across fish. For a pixel to

be marked as ornamental it had to be found in > 11% of all assayed fish for orange pixels

and > 10% of all assayed fish for black pixels. This slightly higher threshold for orange

pigmentation was needed in order to separate orange ornaments OII and OIII which were

otherwise joined by a few black pixels (Fig 1a; 1b). We then used the function hclust() from

the package Stats v3.5.0 to cluster pixels into a predefined number of distinct groups,

defined as ornaments. For orange ornaments we used a predefined number of four

ornaments, as this was very clearly the number of distinct orange regions found in fish. For

black ornaments we tested clustering using 6,7 and 8 ornaments, with one of the black

regions being the eye. From these results, it was clear that there were five main patches of

black pigmentation, plus the eye and one very small region that we did not consider in

downstream analyses. From this we were able to measure the minimum and maximum XY

coordinates or four orange and five black ornaments, and determine the size, hue and

saturation of each ornament in each individual.

To better understand the correlational relationships among ornaments, referred to as

modulation, we calculated Pearson's correlations between ornaments for size (where

ornaments with 10 pixels or less were marked as NA) and saturation in R using the

rcorr.adjust() function (with use= "pairwise.complete.obs") from the package RcmdrMisc

v2.7-1 [46] which uses Holm's method to apply a correction to P-values calculated using the

rcorr() function from Hmisc [47]. We also calculated Spearman's correlations for

presence/absence (with ornaments with 10 pixels or less marked as absent), again using

rcorr.adjust().

With the ornaments we defined from our stock populations, we used functions from the R

packages pedigreemm v0.3-3, sommer v4.1.0, and MCMCglmm v2.29 [48–51] to estimate

heritability for size (the number of pixels), and presence/absence for individual ornaments

from our pedigrees. For size, heritability was calculated using the raw data for each

ornament (in the number of pixels) for all fish that had that ornament, and with fish missing

that ornament marked as NA. A matrix with the relatedness between all individuals in the

pedigree was calculated using the functions pedigree() and getA(). We then used the mmer()

function to fit a univariate mixed effect model to estimate heritability, with a random effect -

'animal', which is linked to the relatedness matrix included in the model formula

(ornament.size ~ (1|animal)). For binary presence/absence data, we used a threshold of 10

pixels for presence with all sizes smaller than this marked as absent. We used the function

MCMCglmm() to estimate heritability as implemented for binary traits [52]. We used

MCMCglmm() for binary traits as it has been found to be more reliable for binary data than

other methods available [53]. We used the priors (R = list(V = 1, fix = 1), G = list(G1 = list(V =

1, nu = 1,000, alpha.mu = 0, alpha.V = 1) with the residual variance fixed to 1 as

recommended for binary data [52], and used a burnin of 100,000 and the number of

iterations set to 2,500,000.

We also calculated heritability for four aggregate traits: total orange area, total black area,

average orange saturation and average black saturation. Total areas were calculated

respectively by summing the areas of all orange or all black ornaments. Average orange and

black saturation were calculated by first finding the mean saturation of pixels in each

ornament, and then calculating the weighted mean of these means (weighted by the size of

each ornament). This resulted in measures for average orange and average black saturation.

This method described above was used for estimating heritability of our aggregate traits.

Strength of Y-effects on ornamentation

We tested for the potential for Y-link effects on all four aggregate traits, total orange area,

total black area, average orange saturation and average black saturation, by looking at the

relationship between the phenotypes of our generation three offspring and the phenotypes

of their paternal and maternal grandfathers. Because the Y chromosome is passed down the

patriline, grandsons will have the Y of their paternal grandfathers but not their maternal

grandfathers. Therefore, we can hypothesise that if there are Y linked loci affecting total

black or black area, then the slope of the paternal grandfather total black or black area

should be greater than the slope of maternal grandfather phenotype on male total black or

black area.

Unfortunately, we were unable to photograph and phenotype the grandfathers of our

generation two offspring with our phenotype analysis pipeline. We therefore have a sample

of 73 males across seven families from generation three . We carried out analyses of both

maternal and paternal effects for all four aggregate traits, using mmer(). As per our

heritability analysis we used a random effect - 'animal', which is linked to a relatedness

matrix, and which controls for relatedness between individuals. We also included a fixed

effect, which was either the maternal or paternal grandfather's phenotype. The coefficients

of this model represent the slope of these grandfather phenotypes (paternal and maternal)

on all four aggregate traits in our generation three offspring. The formula of this analysis

was: (aggregate trait ~ Pat/Mat-GF phenotype + (1|animal)).

Results

Following Thin Plate Spline alignment of orange and black extracted images, we identified

four orange (OI - OIV) and five black ornaments (BI – BV, Fig. 1) found in our stock

population. Although all orange ornaments overlapped with at least one black ornament,

the overlap regions were small, and 59% of potential ornament pixels in Fig 1. were orange,

35% black, and only 6% either orange or black.

Fig 1. Male ornamentation. A) Heatmap showing the proportion of individuals from the
full stock population with orange pigmentation at each pixel, B) Heatmap showing the
proportion of individuals from the full stock population with black pigmentation at each
pixel, with the eye masked, C) Mosaic plot showing the four orange and five black
ornaments defined from the stock population. Grey pixels represent overlap between
orange and black ornaments.

We determined correlations among ornaments in terms of size and saturation to understand

the genetic relationship among different modules in our stock population. We observed

generally positive correlations for both size and presence/absence among many of the

orange ornaments, although many were not significant after adjusting for multiple testing

(Fig. 2A, Table 1 below the diagonal, Supplementary Table 1). These correlations suggest

that there may be modifier loci that globally affect the size of orange ornaments found in

males, and thus also the total amount of orange pigmentation. Many previous studies that

have thus used the total orange or black area as a trait may therefore have been at least in

part measuring the heritability or inheritance of these loci rather than any loci related to the

presence or absence of any individual ornament. In contrast, many of the correlations in size

between adjacent orange and black ornaments were found to be negative (Table 1,

Supplementary table 2), although few of these were significant. This perhaps indicates

epistatic interactions between loci controlling ornaments, increasing the size of one

ornament while reducing the size of others that are adjacent. Alternatively, it may reflect

that where orange and black ornaments overlap, only the black area is visible.

Fig 2. Significant correlations for presence and absence among orange (A) and black (B)
ornaments. Correlations are significant to adjusted P< 0.05.

Table 1. Strength of correlations for ornament size (below the diagonal) and saturation (above the diagonal) among orange and black ornaments. Adjusted P-value <0.05 in bold

Ornament	OI	OII	OIII	OIV	BI	BII	BIII	BIV	BV
OI	-	**0.53**	0.250	-0.05	-0.05	0.07	-0.28	0.04	-0.24
OII	-0.21	-	**0.720**	**0.40**	0.01	-0.03	0.20	0.21	0.21
OIII	0.08	0.19	-	**0.41**	-0.01	0.03	0.21	0.17	0.02
OIV	-0.25	0.28	0.23	-	0.27	-0.10	0.13	-0.01	0.13
BI	-0.27	-0.29	-0.15	0.34	-	0.35	0.37	0.41	0.33
BII	0.13	0.12	0.01	-0.14	**-0.47**	-	0.21	**0.54**	**0.48**
BIII	-0.05	0.03	0.22	0.10	0.17	-0.24	-	0.52	**0.62**
BIV	0.05	**-0.44**	-0.14	-0.07	-0.14	0.02	-0.25	-	**0.57**
BV	-0.10	-0.24	-0.14	0.17	0.52	-0.03	0.12	0.24	-

We also determined the correlation in saturation among ornaments (Table 1 above the

diagonal, Supplementary Table 3). For both black and orange pigmentation, we observed

generally positive correlations in saturation between adjacent ornaments of the same

pigmentation type. Thus again, ornaments in the same pathway are not independent of each

other, suggesting that modifier loci control saturation levels affecting multiple ornaments

simultaneously. We found the correlation in saturation between orange and black

ornaments was on average much lower (average = 0.049) than between ornaments of the

same colour (orange average = 0.377, black average = 0.440), suggesting that modifier loci do

not transcend biochemical pathways to affect both ornament types.

We used our pedigreed males to investigate the pattern of inheritance from father to son

(Table 2). If the presence or absence of an ornament is exclusively Y-linked, all male

offspring of a male will share their father's phenotype. However, in cases where the

presence or absence of an ornament is genetically partially or fully under autosomal, X or

pseudoautsomal control, sons will frequently have different phenotypes than their fathers.

Our results show that no ornament displays the inheritance pattern expected for a purely

male specific Y-linked trait. For two ornaments (OII and BV), we observe a majority of male

offspring have the opposite phenotype to their father. Overall these results suggest that

individual orange and black ornaments are not linked to the Y chromosome in our

population.

Table 2. Father-son heritability of presence/absence and area for each ornament, and for
total orange and total black area.

Ornament	OI	OII	OIII	OIV	BI	BII	BIII	BIV	BV	Total Or.	Total Bl.
Present	57%	89%	96%	57%	24%	98%	27%	64%	36%		
Same: different to father										-	-
Same as father	59%	44%	73%	90%	60%	87%	73%	52%	48%	-	-
h^2 (pres./abs)	0.669	0.456	0.255	0.776	0.121	0.414	0.519	0.581	0.414	0.539	0.410
h^2 (size)	0.508	0.461	0.459	0.484	0.000	0.000	0.000	0.186	0.000	0.678	0.038

Most ornaments show high heritability except for ornaments OIII and BI (Table 2). In

contrast, heritability for total area differed markedly between orange (0.678) and black

(0.038) ornaments. We also found heritable genetic variation for average (across ornaments)

orange saturation (h2 = 0.232) and for average black saturation (h2 = 0.607).

Our analysis of paternal and maternal grandfather effects (Table 3) identified a stronger

relationship between males and their paternal grandfather compared to their maternal

grandfather for total orange area and orange saturation. This indicates that there may

indeed be Y-linked modifier loci that affect total orange area and average orange saturation

in guppies, and is consistent with our previous analyses which show positive correlations in

size and saturation among orange ornaments. Although there was a greater coefficient for

total black area between males and their paternal compared to maternal grandfather, the

coefficients for black saturation were negative for both grandfathers.

Table 3. Coefficients from our analysis of paternal and maternal grandfather effects on
aggregate offspring phenotypes.

	Paternal GF Effect	Maternal GF Effect
Total Orange Area	0.807	0.232
Total Black Area	0.452	-0.214
Average Orange Saturation	0.313	-0.186
Average Black Saturation	-0.465	-0.056

Discussion

Our study uses high resolution image analysis to define individual ornaments while

reducing human bias, in order to examine inheritance patterns across a dataset of 214

offspring from 19 families. By computationally clustering pixels automatically in order to

define individual ornaments [36], we provide a comprehensive and agnostic diagnostic

method for colouration patterning in a semi-natural outbred population. Using this method,

we identified four orange and five black ornaments (Fig. 1) in our population that we then

used to study modulation, heritability and mode of inheritance.

Modulation of ornamentation

The colourful ornaments found in animals are often treated as independent phenotypes,

with the relationships among different ornaments less well considered [54]. However, in

many cases the functions of these ornaments are not independent, but as whole patterns in

conjunction [55]. Furthermore, some ornaments, particularly those that share the same

biochemical basis, may have shared aspects of genetic architecture through the pathways

that control them, and so treating them as individual units may not always be appropriate.

The colour patterns of guppies are used as a sexual signal. It has been proposed that such

signals might be made more conspicuous, and therefore attractive to females, by greater

differences between adjacent patches of colour [54]. If this is the case for guppies, we might

expect selection for linkage between adjacent black and orange patches. We tested this by

examining correlations in presence and absence between different male ornaments (Table 1).

Although we found some strong positive correlations for size within ornaments of the same

colour, there was little correlation among adjacent ornaments of different colours

(Supplementary Table 2). Our results therefore do not provide evidence of linkage between

adjacent and contrasting male ornaments in our population of guppies.

Although we find little correlation between ornaments from different biochemical pathways,

we do find several lines of evidence that guppy ornaments of the same colour are not

independent of each other. Interestingly, orange and black ornaments displayed different

genetic characteristics, with high heritability for size (Table 2) and high correlations in size

(Table 1, Fig. 2) between many of the orange ornaments, but not black ornaments. However,

for both black and orange pigmentation, we observed generally positive correlations in

saturation between adjacent ornaments of the same pigmentation type (Table 1). This

suggests the presence of one or more global modifier loci affecting saturation levels and total

size for multiple ornaments simultaneously, while other independent loci encode for the

presence or absence of individual ornaments. Furthermore, the higher similarity between

grandsons and their paternal grandfathers compared to maternal grandfathers in ornament

size and orange saturation (Table 3) suggests a Y-linked component to this modifier locus,

discussed below.

Sex linkage

Many guppy colour traits have been shown to be Y-linked, reviewed by Lindholm and

Breden [56] and Kottler and Schartl [57]. However, much of this work has either been done

on ornaments in domestic guppy breeds and therefore unlikely to be found in the wild

[31,32], or on composite traits, such as aggregate colour area [21,33,34]. We assessed the

potential for Y-linkage in individual ornaments from our semi-natural populations under

the expectation that any Y-linked ornament displayed by a father would be also present in

all his sons. Despite this previous work, we did not observe Y inheritance patterns for

presence or absence of any single one of our nine ornaments. This suggests that individual

male colour patterns in guppies from our population are not Y-linked. This is also reflected

in the heritability for presence/absence of these ornaments, with more than half of

ornaments with heritability <0.5. Even for the ornaments inherited with the highest

heritability (OIV), 9.7% of offspring males display a different phenotype to their father.

Recent studies have suggested that the guppy Y chromosome is divided into a relatively

small region where recombination between the X and Y never occurs [58–61], and a larger

adjacent region where recombination is rare, occurring in between 1% and 0.1% of males

[59,62,63]. However, because the vast majority of recombination in males is confined to the

telomeric ends of the sex chromosome, nearly the entire chromosome is passed on as a Y

chromosome in >99% of male offspring. Therefore, rare X-Y recombination events cannot

explain the difference between paternal and filial phenotypes we observe in our ornaments.

These results are at first perplexing in light of previous work, and there are several potential

explanations. First, it is possible that our population has lost Y-linkage of ornaments in the

>20 years since it was collected from the wild. However, we think this is not terribly likely

given the large size of the founding colony and the fact that it has been subsequently

maintained as an outbred, freely mating population. Both of these strategies have

maximized genetic diversity and minimized potential bottlenecks. Moreover, Y-linkage has

been observed in other lab and inbred populations [21,64], and it is not clear how or why Y-

linkage would erode.

Second, considerable variation in Y-linkage exists among wild Trinidadian populations [65],

with notable differences between high predation and low predation populations in

particular [33,34,59,66]. It is therefore possible that difference between our results and others

may simply be due to differences between populations. Although we also think this is

unlikely given that the Quare population upon which our stock population is founded has

been studied using aggregate methods, described below [34], it is however is important for

future studies to quantify Y-linkage of individual ornaments in multiple populations.

However, our results are consistent with studies which have characterized aggregate

ornamental area, which often identify significant Y contributions. For example, Postma *et al*

[35] and Brooks and Endler [21] both identified significant levels of Y-linked genetic

variation when assessing total ornamental areas. Our analysis suggests that although the

presence or absence of individual ornaments are not often Y-linked in natural populations,

the Y chromosome does carry one or more modifier loci that affect many ornaments in

aggregate (Table 3). Therefore when studies use the total orange or black area as a trait they

are in large part measuring the mode of inheritance of these loci rather than any loci related

to the presence or absence of any individual ornament [21,34].

It may be that these Y-linked modifier loci work through a simple mechanism, such as

testosterone level or testosterone sensitivity, which has been shown to affect total ornament

area [32–34]. Indeed, the extensive variation exhibited in total ornament area among natural

[34] and introduced [33] populations hints at a simple genetic architecture, as the alternative

of rapid and repeated movement of many underlying genes to the relatively small Y

chromosome, is somewhat improbable. Given the importance of testosterone level and

sensitivity in fish growth rates, maturation and colouration [67–69], and the variation

observed in natural guppy populations in these traits [70], it is likely that testosterone and/or

testosterone sensitivity vary across populations, and this may interact with colouration. The

nature of these interactions, and the identification of the loci responsible for these Y-linked

effects remain interesting areas of future research.

Conclusion

Surprisingly, we did not observe inheritance patterns for presence and absence for any of

our nine ornaments that were consistent with Y-linkage. However, we did find evidence of

Y-linked modifier loci that affect the aggregate phenotypes of total orange and black area.

Given the results of our study, the Y-linkage of individual male colour traits may therefore

not be as important in the rapid evolution of colour changes in natural populations of

guppies [33] or in the expansions of the non-recombining Y region [30] as previously

thought. However, Y-linked modifier loci that affect aggregate traits may well play a role.

Alternatively other mechanisms to resolve sexual conflict, such as changes in gene

regulation [71] may alone be enough to explain colour dimorphism in this species.

Acknowledgements: This work was supported by a grant from the European Research

Council (grant agreement N0. 680951) to JEM, who also gratefully acknowledges support

from a Canada 150 Research Chair and NSERC. We thank Natasha Bloch for her early input,

and Ben Furman, Jacelyn Shu, David Metzger, Ben Sandkam, Yuying Lin, Lydia Fong, Pedro

Almeida and Alberto Corral-Lopez for their helpful comments.

Bibliography

1. Williams P. 2007 The distribution of bumblebee colour patterns worldwide: Possible significance for thermoregulation, crypsis, and warning mimicry. *Biol. J. Linn. Soc.* **92**, 97–118. (doi:10.1111/j.1095-8312.2007.00878.x)

2. Cuthill IC *et al.* 2017 The biology of color. *Science (80-.).* **357**. (doi:10.1126/science.aan0221)

3. Roulin A. 2014 Melanin-based colour polymorphism responding to climate change. *Glob. Chang. Biol.* **20**, 3344–3350. (doi:10.1111/gcb.12594)

4. Merrill RM *et al.* 2015 The diversification of Heliconius butterflies: What have we learned in 150 years? *J. Evol. Biol.* **28**, 1417–1438. (doi:10.1111/jeb.12672)

5. Langham GM. 2005 Rufous-tailed jacamars and aposematic butterflies: do older birds attack novel prey? *Behav. Ecol.* **17**, 285–290. (doi:10.1093/beheco/arj027)

6. Summers K, Clough ME. 2001 The evolution of coloration and toxicity in the poison frog family (Dendrobatidae). *Proc. Natl. Acad. Sci. U. S. A.* **98**, 6227–6232. (doi:10.1073/pnas.101134898)

418 7. van't Hof AE, Edmonds N, Dalikova M, Marec F, Saccheri IJ. 2011 Industrial

419   Melanism in British Peppered Moths Has a Singular and Recent Mutational Origin.

420   *Science (80-.).* **332**, 958–960. (doi:10.1126/science.1203043)

421 8. Steffen JE, Guyer CC. 2014 Display behaviour and dewlap colour as predictors of

422   contest success in brown anoles. *Biol. J. Linn. Soc.* **111**, 646–655. (doi:10.1111/bij.12229)

423 9. Hughes KA, Houde AE, Price AC, Rodd FH. 2013 Mating advantage for rare males in

424   wild guppy populations. *Nature* **503**, 108–110. (doi:10.1038/nature12717)

425 10. Garratt M, Brooks RC. 2012 Oxidative stress and condition-dependent sexual signals:

426   More than just seeing red. *Proc. R. Soc. B Biol. Sci.* **279**, 3121–3130.

427   (doi:10.1098/rspb.2012.0568)

428 11. Svensson PA, Wong BBM. 2011 Carotenoid-based signals in behavioural ecology: A

429   review. *Behaviour* **148**, 131–189. (doi:10.1163/000579510X548673)

430 12. Ruell EW, Handelsman CA, Hawkins CL, Sofaer HR, Ghalambor CK, Angeloni L.

431   2013 Fear, food and sexual ornamentation: Plasticity of colour development in

432   Trinidadian guppies. *Proc. R. Soc. B Biol. Sci.* **280**. (doi:10.1098/rspb.2012.2019)

433 13. Morehouse NI, Rutowski RL. 2010 In the eyes of the beholders: Female choice and

434   avian predation risk associated with an exaggerated male butterfly color. *Am. Nat.*

435   **176**, 768–784. (doi:10.1086/657043)

436 14. Rowe L, Chenoweth SF, Agrawal AF. 2018 The genomics of sexual conflict. *Am. Nat.*

437   **192**, 274–286. (doi:10.1086/698198)

438 15. Cox RM, Calsbeek R. 2009 Sexually antagonistic selection, sexual dimorphism, and

439   the resolution of intralocus sexual conflict. *Am. Nat.* **173**, 176–187. (doi:10.1086/595841)

440 16. Rice WR. 1987 The Accumulation of Sexually Antagonistic Genes as a Selective Agent

441   Promoting the Evolution of Reduced Recombination between Primitive Sex

442   Chromosomes. *Evolution (N. Y).* **41**, 911–914. (doi:10.2307/2408899)

443   17.   Rice WR. 1984 Sex chromosomes and the evolution of sexual dimorphism. *Evolution*

444   *(N. Y).* **38**, 735–742. (doi:10.2307/2408385)

445   18.   Charlesworth B, Jordan CY, Charlesworth D. 2014 The evolutionary dynamics of

446   sexually antagonistic mutations in pseudoautosomal regions of sex chromosomes.

447   *Evolution (N. Y).* **68**, 1339–1350. (doi:10.1111/evo.12364)

448   19.   Furman BLS, Metzger DCH, Darolti I, Wright AE, Sandkam BA, Almeida P, Shu JJ,

449   Mank JE. 2020 Sex chromosome evolution: So many exceptions to the rules. *Genome*

450   *Biol. Evol.* , 1–32. (doi:10.1093/gbe/evaa081)

451   20.   Endler JA. 1983 Natural and sexual selection on color patterns in poeciliid fishes.

452   *Environ. Biol. Fishes* **9**, 173–190. (doi:10.1007/BF00690861)

453   21.   Brooks R, Endler JA. 2001 Direct and Indirect Sexual Selection and Quantitative

454   Genetics of Male Traits in Guppies (Poecilia Reticulata). *Evolution (N. Y).* **55**, 1002.

455   (doi:10.1554/0014-3820(2001)055[1002:daissa]2.0.co;2)

456   22.   Pitcher TE, Rodd FH, Rowe L. 2007 Sexual colouration and sperm traits in guppies. *J.*

457   *Fish Biol.* **70**, 165–177. (doi:10.1111/j.1095-8649.2006.01292.x)

458   23.   Endler JA. 1980 Natural Selection on Color Patterns in Poecilia reticulata. *Evolution (N.*

459   *Y).* **34**, 76. (doi:10.2307/2408316)

460   24.   Houde AE. 1992 Sex-linked heritability of a sexually selected character in a natural

461   population of Poecilia reticulata (Pisces: Poeciliidae) (guppies). *Heredity (Edinb).* **69**,

462   229–235. (doi:10.1038/hdy.1992.120)

463   25.   Lindholm A, Breden F. 2002 Sex Chromosomes and Sexual Selection in Poeciliid

464   Fishes. *Am. Nat.* **160**, S214–S224. (doi:10.1086/342898)

465   26.   Houde AE. 1987 Mate choice based upon naturally occurring color-pattern variation

466   in a guppy population. *Evolution (N. Y).* **41**, 1–10. (doi:10.1111/j.1558-

467   5646.1987.tb05766.x)

27. Reznick D, Endler JA. 1982 The Impact of Predation on Life History Evolution in

Trinidadian Guppies (Poecilia reticulata). *Soc. Study Evol.* **36**, 160–177.

28. Breden F, Stoner G. 1987 Male predation risk determines female preference in the

Trinidad guppy. *Nature* **329**, 831–833.

29. Fisher RA. 1931 The evolution of dominance. *Biol. Rev.* **6**, 345–368. (doi:10.1111/j.1469-

185X.1931.tb01030.x)

30. Wright AE *et al.* 2017 Convergent recombination suppression suggests role of sexual

selection in guppy sex chromosome formation. *Nat. Commun.* **8**, 14251.

(doi:10.1038/ncomms14251)

31. Winge Ö. 1927 The location of eighteen genes inLebistes reticulatus. *J. Genet.* **18**, 1–43.

(doi:10.1007/BF03052599)

32. Haskins CP, Young P, Hewitt RE, Haskins EF. 1970 Stabilised heterozygosis of

supergenes mediating certain y-linked colour patterns in populations. *Heredity*

(Edinb). **25**, 575–589.

33. Gordon SP, López-Sepulcre A, Rumbo D, Reznick DN. 2017 Rapid Changes in the Sex

Linkage of Male Coloration in Introduced Guppy Populations. *Am. Nat.* **189**, 000–000.

(doi:10.1086/689864)

34. Gordon SP, López-Sepulcre A, Reznick DN. 2012 Predation associated differences in

sex linkage in wild guppy coloration. *Evolution (N. Y).* **66**, 912–918. (doi:10.1111/j.1558-

5646.2011.01495.x)

35. Postma E, Spyrou N, Rollins LA, Brooks RC. 2011 Sex-dependent selection

differentially shapes genetic variation on and off the guppy Y chromosome. *Evolution*

(N. Y). **65**, 2145–2156. (doi:10.1111/j.1558-5646.2011.01314.x)

36. Holman L, Head ML, Lanfear R, Jennions MD. 2015 Evidence of experimental bias in

the life sciences: Why we need blind data recording. *PLoS Biol.* **13**, 1–12.

(doi:10.1371/journal.pbio.1002190)

37.   Kotrschal A *et al.* 2013 Artificial Selection on Relative Brain Size in the Guppy Reveals
Costs and Benefits of Evolving a Larger Brain. *Curr. Biol.* **23**, 168–171.
(doi:10.1016/j.cub.2012.11.058)

38.   Leutner B, Horning N, Schwalb-Willmann J. In press. RStoolbox: Tools for remote
sensing data analysis. R package version 0.2.6.

39.   Hijmans RJ, van Etten J. 2012 raster: Geographic analysis and modeling with raster
data. R package version 2.0-12.

40.   R Core Team. 2018 R: A language and environment for statistical computing. R
Foundation for Statistical Computing.

41.   Van Belleghem SM, Papa R, Ortiz-zuazaga H, Hendrickx F, Jiggins CD, Mcmillan
WO, Counterman BA. 2018 Patternize: an R package for quantifying color pattern
variation. *Methods Ecol. Evol.* **9**, 390–398.

42.   Rohlf JF. 2013 tpsDig, digitize landmarks and outlines.

43.   Rohlf JF. 2013 tpsUtil, file utility program.

44.   Adams DC, Otárola-Castillo E. 2013 geomorph: an R package for the collection and
analysis of geometric morphometric shape data. *Methods Ecol. Evol.* **4**, 393–399.

45.   Schlager S. 2017 Morpho and Rvcg - Shape analysis in R: R-packages for geometric
morphometrics, shape analysis and surface manipulations.

46.   Fox J. 2020 RcmdrMisc: R Commander Miscellaneous Functions. R package version
2.7-1.

47.   Harrell Jr FE, with contributions from Charles Dupont and many others. 2019 Hmisc:
Harrell Miscellaneous.

48.   Vazquez AI, Bates DM, Rosa GJM, Gianola D, Weigel KA. 2010 Technical note: An R
package for fitting generalized linear mixed models in animal breeding. *J. Anim. Sci.*

88, 497–504.

49.   Covarrubias-Pazaran G. 2016 Genome-Assisted prediction of quantitative traits using
the r package sommer. *PLoS One* **11**, 1–15. (doi:10.1371/journal.pone.0156744)

50.   Covarrubias-Pazaran G. 2018 Software update: Moving the R package sommer to
multivariate mixed models for genome-assisted prediction. *bioRxiv* , 354639.

51.   Hadfield JD. 2010 MCMC Methods for Multi-Response GLMMs in R. *J. Stat. Softw.* **33**,
1–22.

52.   de Villemereuil P. 2012 Estimation of a biological trait heritability using the animal
model.

53.   De Villemereuil P, Gimenez O, Doligez B. 2013 Comparing parent-offspring
regression with frequentist and Bayesian animal models to estimate heritability in
wild populations: a simulation study for Gaussian and binary traits. *Methods Ecol.*
Evol. **4**, 260–275.

54.   Endler JA. 2012 A framework for analysing colour pattern geometry: adjacent colours.
Biol. J. Linn. Soc. **107**, 233–253.

55.   Endler JA, Cole GL, Kranz AM. 2018 Boundary strength analysis: Combining colour
pattern geometry and coloured patch visual properties for use in predicting
behaviour and fitness. *Methods Ecol. Evol.* **9**, 2334–2348.

56.   Lindholm A, Breden F. 2002 Sex chromosomes and sexual selection in poeciliid fishes.
Am. Nat. **160**, S214–S224.

57.   Kottler VA, Schartl M. 2018 The colorful sex chromosomes of teleost fish. *Genes* **9**, 233

58.   Yamamoto T. 1975 The medaka, *Oryzias latipes*, and the guppy, *Lebistes*. In *Handbook of*
Genetics (ed RC King), pp. 133–149. Boston, MA: Spinger.

59.   Haskins CP, Haskins EF, Mclaughlin JJA, Hewitt RE. 1961 Polymorphism and

population structure in *Lebistes reticulatus*, an ecological study. In *Vertebrate Speciation*

(ed WF Blair), pp. 320–395. Austin: University of Texas Press.

60. Tripathi N, Hoffmann M, Willing E-M, Lanz C, Weigel D, Dreyer C. 2009 Genetic

linkage map of the guppy, *Poecilia reticulata*, and quantitative trait loci analysis of

male size and colour variation. *Proc. Biol. Sci.* **276**, 2195–208.

61. Darolti I, Wright AE, Mank JE. 2020 Guppy Y chromosome integrity maintained by

incomplete recombination suppression. *Genome Biol. Evol.* , 1–30.

62. Lisachov AP, Zadesenets KS, Rubtsov NB, Borodin PM. 2015 Sex chromosome

synapsis and recombination in male guppies. *Zebrafish* **00**, 1–7.

63. Bergero R, Gardner J, Bader B, Yong L, Charlesworth D. 2019 Exaggerated

heterochiasmy in a fish with sex-linked male coloration polymorphisms. *Proc. Natl.*

Acad. Sci. U. S. A. **116**, 6924–6931.

64. Khoo G, Lim TM, Chan W, Phang VPE, Khoo G, Lim TM, Chan W, Phang VPE. 1999

Linkage analysis and mapping of three sex-linked color pattern genes in the Guppy,

Poecilia reticulata. Zoolog. Sci. **16**, 893–903.

65. Fraser BA, Künstner A, Reznick DN, Dreyer C, Weigel D. 2015 Population genomics

of natural and experimental populations of guppies (*Poecilia reticulata*). *Mol. Ecol.* **24**,

389–408.

66. Hughes KA, Rodd FH, Reznick DN. 2005 Genetic and environmental effects on

secondary sex traits in guppies (*Poecilia reticulata*). *J. Evol. Biol.* **18**, 35–45.

67. Yue M, Zhao J, Tang S, Zhao Y. 2018 Effects of estradiol and testosterone on the

expression of growth-related genes in female and male nile tilapia, *Oreochromis*

niloticus. J. World Aquac. Soc. **49**, 216–228.

68. Bayley M, Junge M, Baatrup E. 2002 Exposure of juvenile guppies to three

antiandrogens causes demasculinization and a reduced sperm count in adult males.

Aquat. Toxicol. **56**, 227–239.

69.   Matty AJ, Lone KP. 2009 The effect of androgenic steroids as dietary additives on the
growth of carp (*Cyprinus carpio*). *Proc. World Maric. Soc.* **10**, 735–745.

70.   Zandonà E *et al.* 2011 Diet quality and prey selectivity correlate with life histories and
predation regime in Trinidadian guppies. *Funct. Ecol.* **25**, 964–973.

71.   Mank JE. 2017 The transcriptional architecture of phenotypic dimorphism. *Nat. Ecol.*
Evol. **1**, 0006.