Title: Parietal Mechanisms for Transsaccadic Spatial Frequency Perception: An fMRI Study

Abbreviated Title: Cortical mechanisms for transsaccadic feature processing

Author(s) and Affiliations: B. R. Baltaretu[1, 2], B. T. Dunkley[3,4,5], W. Dale Stevens[1,6] and J. D. Crawford*[1, 2,6,7]

[1] Centre for Vision Research and Vision: Science to Applications (VISTA) program, York University, Toronto, Ontario M3J 1P3, Canada

[2] Department of Biology, York University, Toronto, Ontario M3J 1P3, Canada

[3] Department of Diagnostic Imaging, Hospital for Sick Children, Toronto, Ontario M5G 1X8, Canada

[4] Neurosciences & Mental Health, Hospital for Sick Children Research Institute, Toronto, Ontario M5G 1X8, Canada

[5] Department of Medical Imaging, University of Toronto, Toronto, Ontario M5S, Canada

[6] Department of Psychology and Neuroscience Graduate Diploma Program, York University, Toronto, Ontario M3J 1P3, Canada

[7] School of Kinesiology and Health Sciences, York University, Toronto, Ontario M3J 1P3, Canada

*Corresponding Author: Bianca R. Baltaretu

b.baltaretu@gmail.com

Number of Figures: 5

Number of Supplementary Figures: 2

Number of Words (Abstract): 175

Number of Words (Introduction): 528

Number of Words (Discussion): 1192

Conflict of Interest: The authors declare no conflict of interest.

Acknowledgements: The authors thank Joy Williams, Dr. Xiaogang Yan, and Saihong Sun for technical assistance. This work was supported by a Natural Sciences and

38 Engineering Council (NSERC) of Canada Discovery grant. B.R. Baltaretu was
39 supported by the NSERC Brain-in-Action CREATE program, and J.D. Crawford is
40 supported by the Canada Research Chair Program.
41
42 **Author Contributions:** B.R.B. adapted the paradigm, collected and analyzed the data,
43 and wrote the manuscript. B.T.D. developed the original paradigm and edited the
44 manuscript. W.D.S. supervised data analysis and edited the manuscript. J.D.C.
45 supervised overall project development and edited the manuscript.
46

Abstract

Posterior parietal cortex (PPC), specifically right supramarginal gyrus, is involved in

transsaccadic memory of object orientation for both perception and action. Here, we

investigated whether PPC is involved in transsaccadic memory of other features,

namely spatial frequency. We employed a functional magnetic resonance imaging

paradigm where participants briefly viewed a grating stimulus with a specific spatial

frequency that later reappeared with the same or different frequency, after a saccade or

continuous fixation. Post-saccadic frequency modulation activated a region in the right

hemisphere spanning medial PPC (ventral precuneus) and posterior cingulate cortex.

Importantly, the site of peak precuneus activation showed saccade-specific feature

modulation (compared to fixation) and task-specific saccade modulation (compared to a

saccade localizer task). Psychophysiological interaction analysis revealed functional

connectivity between this precuneus site and the precentral gyrus (M1), lingual gyrus

(V1/V2), and medial occipitotemporal sulcus. This differed from the transsaccadic

orientation network, perhaps because spatial frequency signaled changes in object

identity. Overall, this experiment supports a general role for PPC in transsaccadic

vision, but suggests that different networks are employed for specific features.

Introduction

The visual system tracks both low-level (e.g., orientation, spatial frequency) and high-level (e.g., objects, faces) components of our visual surroundings through space and time [1], despite the interruption of several saccades (rapid eye movements) per second [2,3]. To do this, visual features must be encoded, retained, updated, and integrated across saccades [4,5], through a process called transsaccadic perception [2,6,7,8]. As argued elsewhere [9-11], transsaccadic perception likely incorporates mechanisms for both visual working memory [12,13] and spatial updating [14,15]. However, the specific *neural* mechanisms for transsaccadic feature perception are not well understood.

When saccades occur, they cause both object locations and their associated features to shift relative to eye position. It is well established that human posterior parietal cortex (PPC; specifically, the mid-posterior parietal sulcus), is involved in transsaccadic spatial updating, i.e., the updating of object *location* relative to each new eye position [16-19]. Recently, we found that inferior PPC (specifically, right supramarginal gyrus; SMG) is also modulated by transsaccadic comparisons of object orientation [20]. This area is located immediately lateral to the parietal eye field / lateral intraparietal sulcus, so might be an evolutionary expansion of the monkey intraparietal eye fields, which show modest spatial updating of object information across saccades [21]. Consistent with these findings, transcranial magnetic stimulation (TMS) of PPC (just posterior to SMG) disrupted transsaccadic memory of multiple object orientations [22,23]. SMG activity is also modulated during transsaccadic updating of object orientation for grasp, along with other parietal sensorimotor areas [24]. These findings implicate PPC in the transsaccadic updating of *both* object location and orientation.

It is not known if these neural mechanisms generalize to other stimulus features.

One might expect PPC to be involved in other aspects of transsaccadic feature memory

and integration, because of its general role in spatial updating [17,25,26]; however, the

specific mechanisms might differ. SMG seems to play a specialized role for high-level

object orientation in various spatial tasks [27,28]. Thus, while SMG might play a general

role in transsaccadic updating of all visual features, it is equally possible that the brain

engages different cortical networks for transsaccadic processing of different features, as

it does during prolonged visual fixations [29,30].

To address this question, we used an event-related fMRI paradigm, similar to

Dunkley et al. [20], where participants briefly viewed a 2D spatial frequency grating

stimulus, either while continually fixating the eyes, or while making a saccade to the

opposite side, and then judged whether a re-presented grating was the same or

different. But here, we modulated spatial frequency, rather than orientation (Fig. 1a,b).

As in Dunkley et al. [20], we first localized clusters that were activated by changes in

stimulus frequency following saccades; then, as in Baltaretu et al. [24] we determined if

the site(s) of peak PPC activation passed two additional criteria: their feature

modulations must be saccade-specific (Fig. 2, prediction 1), and 2) they must show

task-specific saccade modulations relative to a simple saccade motor task (Fig. 2,

prediction 2). Finally, we analyzed the functional connectivity of this area. Our findings

confirm the involvement of PPC in processing transsaccadic feature interactions, but

demonstrate that different PPC areas/networks are involved in transsaccadic

processing of spatial frequency versus orientation.

111   **--- Figure 1 Here ---**

112   **--- Figure 2 Here ---**

Results

Overall, our task (Fig. 1) produced widespread and progressive activation in brain areas

related to vision, visual memory and eye movements during the initial *Sensory/Memory*

Phase (Fig. S1) and *Visual/Oculomotor Updating Phase* (Fig. S2). Here, we focus on

the specific analysis pipeline that we used to identify and test if any parietal sites

showed saccade- and task-specific feature modulations for spatial frequency (Fig. 2).

For this analysis, we only analyzed the *Visual/Oculomotor Updating* phase of our task,

i.e. the period after the first stimulus presentation, starting at the time when a saccade

occurred and a second stimulus (either Same or Different) appeared (Fig. 1a). An *a*

priori power analysis suggested that 14 participants were required for the voxelwise

contrasts used in this pipeline (see Methods: Power analysis). To obtain this level, we

continued testing participants (21 in total), until 15 of these passed our behavioural

inclusion criteria for fMRI analysis (See Methods: Behavioural data and exclusion

criteria).

Selection of parietal site for hypothesis testing.

Our overall aim was to identify parietal site(s) that showed post-saccadic feature

modulations and then use these sites to test the predictions shown in Figure 2. As a first

step in this analysis, we performed a whole-brain voxelwise contrast (*Different* Spatial

Frequency > *Same* Spatial Frequency) on fMRI data derived from the

Visual/Oculomotor Updating phase of our saccade trials. The resulting group data are

shown in Fig. 3a, overlaid on a representative anatomic scan in 'inflated brain'

coordinates (note that while visually convenient, this convention results in small spatial

distortions). This contrast revealed two regions: First, a continuous region of change-

dependent activation in the right hemisphere spanning medial PPC (ventral precuneus

and the posterior cingulate gyrus, PCu/pCG). The other region (right medial frontal

gyrus, MFG), showed the opposite pattern, i.e., change suppression, but was not further

analyzed. Notably, unlike similar contrasts in studies of transsaccadic *orientation*

processing [20], SMG was not significantly modulated.

The second step in this analysis was to determine specific parietal coordinates

for our hypothesis tests. To do this, we examined the medial PCu/pCG region shown in

Fig. 3a, but raised the statistical threshold until only the peak site of parietal activation

remained, corresponding to ventral precuneus (vPCu). We then used BrainVoyager

(BrainVoyager QX v2.8, Brain Innovation) to determine the cluster of activation around

the peak voxel and extract data for prediction testing [24,31,32]. This site is shown in

Fig. 3b, overlaid on sagittal anatomic slice (which is less susceptible to distortions).

Coordinates and supporting references for this site (and the other regions mentioned)

are shown Table 1.

153   **--- Figure 3 Here ---**

Hypothesis 1: Saccade-specific feature modulations.

According to prediction 1 (Fig. 2), cortical sites involved in transsaccadic feature

comparisons should show positive (Different > Same) feature modulations following

saccades, but not during sustained fixation. Figure 3a suggests that vPCu is modulated

by frequency changes after saccades, but does not show if this modulation is saccade-

specific. To test this directly (and independently from the contrast used to localize the

site), we extracted β-weights for all participants from a small cubic region surrounding

the peak voxel coordinates for vPCu (Table 1). We did this under four conditions for the

same site: Saccade Different frequency (SD), Saccade Same frequency (SS), Fixation

Different frequency (FD) and Fixation Same frequency (FD). To test hypothesis 1, we

then compared *Different - Same* feature for *Saccade* (SD-SS) versus *Fixation* data (FD-

FS), as shown in Fig. 3c. In short, vPCu followed the predicted pattern: it showed a

significantly greater *Different-Same* feature modulation after saccades compared to

sustained fixation (t(11)= -2.07, p= 0.031, effect size= 0.60).

Hypothesis 2: Task-specific saccade modulations.

According to prediction 2 (Fig. 2), cortical sites that are specifically involved in

transsaccadic feature comparisons should be modulated by saccades in a task-specific

fashion, i.e. not simply by the motor act of producing a saccade. Figure 4 provides this

test using similar conventions to Figure 3. Figure 4a provides a general overview of the

data used in this calculation, showing Saccade > Fixation modulations during the

updating portion of our task and for reference, the same contrast from our saccade

localizer task, where participants simply looked back and forth between two points (see

Methods for details). As in our previous study [24], saccade modulations were generally

more widespread when they accompanied a more complex task. Figure 4b shows that

the location of the vPCu site used for our hypothesis testing bordered on the edge of a

major cluster of saccade modulation.

Figure 4c provides the statistical test of prediction 2, using data derived from our

vPCU site in a similar fashion to that described above. i.e. *Saccade-Fixation* difference

in the experimental task (TS-TF) versus the same difference from the saccade localizer

(S-F). Note that we could only do this for the 12 participants that passed

behavioural/imaging criteria in both tasks. Once again these data followed the predicted

pattern, showing task-specific saccade modulations in our main experiment compared

to the saccade localizer data (t(11)= 2.57, p=0.013, effect size= 0.68). Overall, these

findings suggest that activity in our ventral precuneus site reflects changes in spatial

frequency in both a saccade-specific and task-specific manner.

192   **-Figure 4 here-**

Functional network for spatial frequency-specific transsaccadic perception.

Our previous study of transsaccadic orientation updating showed a functional network

connecting right SMG to both saccade and grasp motor areas [24]. Here, we provide a

similar analysis of precuneus activity for the current perceptual task. To do this, we used

a *Saccade > Fixation* contrast in a psychophysiological interaction (PPI) analysis,

wherein right precuneus served as the seed region. Figure 5a shows these data

overlaid on a series of transverse brain slices. We observed statistically significant

functional connectivity between precuneus (cyan region) and three other regions

(orange): 1) left lingual gyrus (LG), 2) a caudal portion of the left medial

occipitotemporal sulcus (MOtS), and 3) right precentral gyrus (likely primary motor

cortex, M1).

--- Figure 5 Here ---

Discussion

The goal of this study was to determine whether the involvement of PPC in

transsaccadic updating of visual location and orientation generalizes to other object

features, such as spatial frequency. We found that medial PPC and a lateral frontal

region were modulated by spatial frequency changes across saccades. Further

hypothesis testing on PPC showed that ventral precuneus passed our predefined

criteria for putative transsaccadic updating [24]. Finally, our functional connectivity

analysis suggests that this area interacts with other visual and motor areas to perform

the task that we required of our participants.

Putative sites for transsaccadic integration of spatial frequency.

It has previously been shown that PPC, specifically intraparietal sulcus and adjacent

portions of inferior PPC, are involved in transsaccadic processing of object location and

orientation [17,20,24]. The current results extend this role to spatial frequency

processing, but implicate a different portion of PPC: ventral precuneus, along with other

sites in cingulate and frontal cortex. The latter three sites have several common

characteristics. First, although they modulate with saccades, they do not overlap with

the classic saccade motor areas (Fig. S2) [33-35]. Second, they were all located in the

right hemisphere, consistent with the lateralization observed previously for spatial

attention [36,37], spatial updating [38], and transsaccadic orientation processing [20,24].

Finally, they are all relatively 'high level' (as opposed to early visual cortex), presumably

because our participants performed a top-down, instruction-based task, in contrast to

automatic, bottom-up feature integration [11].

Since the intent of this experiment was to explore the role of PPC in

transsaccadic integration, we focused our analysis on right ventral precuneus, which

met all of our criteria for transsaccadic feature processing. In general, precuneus is

associated with complex visuospatial transformations [39-41]. Dorsal precuneus is

associated with reaching [42-44], saccades [45,46], and visual memory [47]. Ventral

precuneus is involved in change detection [47] and egocentric processing [48], and

shows connectivity with the parahippocampus, cuneus, LG (positive connections), and

MFG (negative connections) [49]. In short, ventral precuneus has functions and

connectivity generally consistent with the task and observations in the current study.

In the current study, precuneus showed extensive functional connectivity with

other areas (see next section), and fell within a broad region of right medial posterior

cortex activation, that also included another peak in the posterior cingulate gyrus. The

posterior cingulate gyrus region is thought to contribute to several saccade-related

functions that might be related to transsaccadic perception, including attention orienting

[50], and retainment of information in working memory [51]. Saccade-related attention

has been closely linked with transsaccadic perception [52].

Finally, we found the opposite pattern of post-saccade activation (*Different <*

Same frequency) in the right middle frontal gyrus. This is consistent with negative

functional connectivity between precuneus and MFG [49]. MFG is located within the

dorsolateral prefrontal cortex (dlPFC), which contains various subregions, some

saccade-sensitive [53], critical for spatial working memory and top-down control [54-

56]. This area has previously been implicated in high-level aspects of transsaccadic

perception: e.g., online TMS to dlPFC disrupted memory of multiple objects during

fixation, but enhanced transsaccadic memory [57]. In general, dlPFC, has been

implicated in various other relevant visual functions, including monochromatic

perception of texture [58-60] (although see [61]), attention orientation [62], and retention

of perceived information in working memory [55,63,64].

Putative functional network for transsaccadic perception of spatial frequency.

In our previous study of transsaccadic updating of grasp orientation, SMG mainly

showed connectivity with sensorimotor areas, including the frontal and supplementary

eye fields, anterior intraparietal cortex, and adjacent portions of superior parietal cortex

involved in grasp control [24]. Here we observed a very different pattern of connectivity:

our seed region (right ventral precuneus) showed task-related functional connectivity

with right precentral gyrus (likely M1) [65], left LG, and left MOtS. Although PPI analysis

is limited in the sense that it does not provide directionality [66], this analysis suggests a

bilateral network for transsaccadic perception of spatial frequency involving both lower

and higher-level visual areas. Based on these results, we propose the network model

illustrated in Figure 5b, where (as discussed above) ventral precuneus is the central

(PPC) hub for spatial frequency processing in this task, and contiguous activity in

posterior cingulate contributes saccade-related attention signals [52].

LG (likely corresponding with visual areas V2/V3) processes low-level

information about spatial frequency in this network [67-70], and has also been

implicated in spatial updating [16,71]. MOtS, located within occipitotemporal cortex, is

thought to be involved in higher-level aspects of spatial frequency processing related to

objects and scenes [72]. Involvement of precentral gyrus most likely reflects preparatory

activity for the final response, the button press [73-75]. Finally, although MFG/dlPFC did

not reappear in our connectivity analysis, it was suppressed by feature changes that

followed saccades (Fig. 2a) and has been shown to be functionally connected with

ventral precuneus in previous studies [49]; a blue dotted line in the model (Fig. 5b) from

this area denotes its putative role in top-down control [57,76,77].

Feature- and task-related aspects of transsaccadic perception.

As noted above, our previous experiments showed transsaccadic modulations for object

orientation in inferior parietal cortex (specifically SMG) along with some task-dependent

extrastriate and superior parietal areas [20,24]. Why would the sites and networks

observed for transsaccadic spatial frequency processing in the current study be so

different? In part, it is likely that transsaccadic perception builds on computations

already used during fixation, which are themselves feature- and task-dependent

[21,78,79,80], especially in the higher-level visual areas activated in our tasks [48,65].

At early levels like V1, there is clear multiplexing of orientation and spatial frequency

[67,81,82], but at higher levels these may tap into entirely different processes. For

example, in our experiment, a change in orientation might influence how one can orient

to an object [24,83], but a change in spatial frequency can denote more fundamental

changes in perceived identity and affordance, evoking very different cortical

mechanisms [84]. Moreover, changes in motor aspects of the task can readily explain

why the functional connectivity in the current task (with an abstract button press

response) was very different from that in our previous experiment where the perceived

object was then grasped [24]. Finally, it is likely that more automatic bottom-up aspects

of transsaccadic integration (such as the automatic integration of motion signals across

saccades) [85] involve different mechanisms again, perhaps involving earlier visual

centres [11,16,71]. Thus, the notion of a dedicated 'transsaccadic integration centre' is

likely naïve: transsaccadic vision, not prolonged fixation, is normal biological vision, and

has likely developed different and nuanced mechanisms, depending on feature and task

details.

Conclusion

In this fMRI study, we set out to explore the cortical mechanism for transsaccadic

updating of spatial frequency, with emphasis on the role of PPC. We found that PPC

(and likely other cortical areas) play a key role, but in contrast to involvement of SMG

for transsaccadic orientation perception [20], the key parietal hub for transsaccadic

spatial frequency processing was ventral precuneus. We also found different patterns of

functional connectivity compared to our previous grasp updating experiment, i.e., where

SMG activity correlated with saccade and grasp motor areas [24]. Overall, these

findings support the role of parietal cortex in transsaccadic vision, but suggest that

specific network recruitment is feature- and task-dependent.

Materials and Methods

Participants.

We tested 21 (human) participants from York University (Toronto, Canada) of

whom 15 met our inclusion criteria for analyses (see Behavioural analysis and exclusion

criteria), thus exceeding the requirement for sufficient statistical power (see Power

analysis) [86]. These 15 individuals (average age: 26.6 +/- 4.3 years; age range: 21-37;

11 females and 4 males; all right-handed) had no neurological disorders and normal or

corrected-to-normal vision. Informed consent was obtained from each participant; all

participants were remunerated for their time. We confirm that the experiment protocol

involving human participants was approved by and in accordance with guidelines of the

York University Human Participants Review Subcommittee.

Experimental set-up.

Participants passed initial MRI safety screening and were then informed about the

experimental task. Participants practiced before taking part in the experiment. Once

they felt comfortable with the task, they assumed a supine position on the MRI table,

with their head resting flat within a 32-channel head coil. An apparatus, holding a mirror

to reflect images from the screen in the MRI bore, was attached to the head coil. An

MRI-compatible eye-tracker (iViewX, SensoMotoric Instruments) was also attached to

the apparatus in order to record the position of the right eye. Participants held an MRI-

compatible button box in their right hand. Their right index and middle fingers rested on

two buttons, used to provide the task responses when presented with the 'go' cue (see

General paradigm).

Stimuli.

During the experiment, participants were presented with an 18° stimulus that contained

a vertical sine-wave grating pattern, averaged to the mean luminance of the screen.

Stimuli were presented in the centre of the screen on a light gray background (MATLAB;

Mathworks, Inc.). The two spatial frequencies that were tested were 0.7 or 1.1

cycles/degree (cpd) (see Fig. 1a).

General paradigm.

In order to identify the cortical activity correlates of transsaccadic perception of spatial

frequency, we used a modified 2 (Gaze: Fixate or Saccade) x 2 (Spatial Frequency: 0.7

or 1.1 cpd) slow event-related fMRI design [20]. Here, we modulated the spatial

frequency (repeated or changed within the same trial; *'Same'* and *'Different'* conditions,

respectively) and/or position of the eyes (continued fixation of gaze or a saccade was

produced; *'Fixation'* and *'Saccade'* conditions). This resulted in four main conditions: 1)

Fixation/Same, 2) Fixation/Different, 3) Saccade/Same, and 4) Saccade/Different.

These were randomly intermingled and repeated four times within each run; there were

six runs in total for each participant.

Trial sequence.

Each trial was composed of three main phases: 1) an initial presentation of the stimulus,

which requires sensory processing and working memory storage (*'Sensory/Memory'*

phase); 2) a second presentation of the stimulus, which requires sensory processing

and (oculomotor) updating (for saccades) (*'Visual/Oculomotor Updating'* phase); and 3)

a response period, where a button press was made to indicate if the spatial frequency of

the stimulus presentations was the same or different (*'Response'* phase). The sequence

of events in each trial (Fig. 1a) began with a 2 s fixation period, with the fixation cross

presented at one of two possible positions (13.5° to the left or right of centre, or 4.5° to

the left or right of the central stimulus). The stimulus was then presented in the centre

for 5.8 s, followed by a 200 ms static noise mask. Once the mask disappeared, a

fixation cross appeared for 200 ms at the same position (*Fixation* condition) or the other

fixation position (*Saccade* condition) (see Fig. 1b for eye position trace). The central

stimulus was presented a second time for 5.8 s at the same spatial frequency (*Same*

condition) as in the *Sensory/Memory* phase or at the other spatial frequency (*Different*

condition). Finally, after the fixation cross and stimulus disappeared, a written prompt

('R or N?') appeared for 8 s in the same location as the fixation cross, prompting

participants to indicate via button press whether the spatial frequency of the two

stimulus presentations was the same (R) or different (N).

Each trial sequence lasted 22 s in total. The four main conditions were repeated

four times per run, resulting in 16 trials per run. Each run started and ended with a 16 s

period of central fixation, serving as a baseline. Overall, each run lasted 6 min 24 s.

Saccade localizer.

We ran a separate saccade localizer in order to identify regions of the brain that

respond to saccade production. The sequence of events began with an 8 s fixation

period of a central white cross on a black background. This was followed by a period of

directed saccadic eye movements between two fixation crosses, randomly from left to

right across trials, for 8 s. The pattern of fixation followed by saccades was repeated 16

times per localizer run, lasting ~4 min 16 s for each of two runs.

MRI parameters.

A 3T Siemens Magnetom TIM Trio MRI scanner at the York MRI Facility was used to

acquire fMRI data. An echo-planar imaging (EPI) sequence (repetition time [TR] = 2000

ms; echo time [TE] = 30 ms; flip angle [FA] = 90°; field of view [FOV] = 240 x 240 mm,

matrix size = 80 x 80, in-plane resolution = 3 mm × 3 mm; slice thickness = 3 mm) was

acquired in ascending, interleaved order for each of the six functional runs and for the

two separate saccade localizer runs. Thirty-three contiguous slices were acquired per

volume for a total of 192 volumes of functional data in each experimental run, and 128

volumes of data for each saccade localizer run. A T1-weighted anatomical reference

volume was obtained for each participant using an MPRAGE sequence (TR= 1900 ms;

FA= 256 mm x 256 mm; voxel size= 1 x 1 x 1 mm^3). 192 slices were acquired per

volume of anatomical data.

Analysis

Power analysis.

To determine the appropriate number of participants to provide a sufficient effect size

and level of power, we used results from the most recent, relevant findings [24].

Specifically, we used the effect size (0.887, Cohen's d) from the most relevant region of

activation in parietal cortex (i.e., SMG) and applied the following properties: 1) two-tailed

t-test option, 2) an α value of 0.05, and 3) a power value of 0.85. Using G*Power 3.1

[87], we determined that a minimum of 14 participants would be required to achieve an

actual power value of 0.866. As noted above, we tested a total of 21 participants in

order to exceed this minimal number, after application of exclusion criteria in the next

section.

Behavioural data and exclusion criteria.

In our previous studies [20,24], we found that experiments such as this are highly

sensitive to head motion. Due to excessive head motion, data were excluded from

analyses on the basis of two criteria: 1) presence of head motion over 1 mm/degree

and/or 2) any abrupt motion of the head over 0.5 mm/degrees. If more than 50% of all

data (i.e., at least 3 runs out of the total 6 runs for a given participant) were removed

from analysis, the entire data set from that participant was removed. On this basis, data

from six participants were removed. From the remaining 15 participants, one run was

removed from data analysis for each of two participants, and two runs were removed for

each of another two participants, for a total of six runs (6.7%).

Eye tracking and button-press data were analyzed post-image acquisition in

order to determine whether the task was completed correctly or not. Eye position data

(e.g., Fig. 1b) were inspected visually to confirm that the eye fixated on the fixation

crosses and/or moved to the correct saccade location when prompted to do so in all

trials. Button press responses were also inspected offline to ensure that participants

responded correctly to the *Same/Different* condition trials. On these bases, 41 trials

were excluded from further analysis (3.1% of the remaining data). Overall, accuracy of

the 15 participants included in the final data analysis was 97.4% ± 3.1%. Only data for

correct trials were included in all further analyses.

Functional imaging data: experimental.

To model the fMRI BOLD response, we used a general linear model (GLM) analysis. In

this model, a standard two-gamma haemodynamic response function (BrainVoyager QX

2.8, Brain Innovation) was convolved with predictor variables [20]. We had five major

classes of predictors for each trial: 1) a baseline predictor ("Baseline"), corresponding to

the first and last 16 s of each run; 2) "Fixate", which represented initial trial fixation

(either left or right); 3) "Adapt", which modeled the activity in response to the first

stimulus presentation; 4) "Fixate/Same", "Fixate/Different", "Saccade/Same", or

"Saccade/Different" to model activity in response to the second stimulus presentation in

one of the four main conditions; and 5) "Response", which modeled the activity for the

button press response period. GLMs were generated using the eight predictors per run

for each participant (BrainVoyager QX 2.8, Brain Innovation).

Preprocessing of functional data from each run for all participants included slice

scan-time correction (cubic spline), temporal filtering (for removal of frequencies < 2

cycles/run), and 3D motion correction (trilinear/sinc). Anatomical data were transformed

to a standard Talairach template [88]. Functional data were coregistered using gradient-

based affine alignment (translation, rotation, scale affine transformation) to the raw

anatomical data. Lastly, the functional data were spatially smoothed using a Gaussian

kernel with a FWHM of 8 mm.

Using the random-effects (RFX) GLM with all of the runs of all of the remaining

15 participants, we performed two major types of analyses: 1) parietal site-of-interest

prediction testing and 2) voxelwise contrasts (see Hypothesis testing: Analysis and

statistical considerations and Voxelwise map contrasts: Analysis and statistical

considerations for details).

Functional imaging data: saccade localizer.

Each run of the saccade localizer had 16 repetitions of the fixation trials (8 s of central

fixation) and saccade trials (8 s of directed saccades). Each trial type was coded by an

8 s predictor ("Fixation" for the fixation trials, and "Saccade" for the saccade trials).

These predictors were convolved with the standard two-gamma haemodynamic

response function (BrainVoyager QX 2.8, Brain Innovation). Preprocessing of functional

and anatomical data occurred as for experimental data (see previous).

On the basis of behavioural data analysis (i.e., excessive motion > 1 mm), data

from three participants were removed, leaving saccade localizer data for 12 participants.

These data were used in the prediction testing in order to identify regions related to the

production of saccades.

Voxelwise map contrasts: Analysis and statistical considerations.

We conducted voxelwise contrasts on data from the 15 participants included in the

analysis (for task, and on the remaining 12 participants for the localizer). For analysis

purposes, we divided the experimental task data into the *Sensory/Memory* phase (when

participants saw and initially remembered the stimulus) and *Visual/Oculomotor Updating*

phase (when saccades occurred and the stimulus reappeared). For all (voxelwise)

contrasts, we first applied a False Discovery Rate (FDR) of $q < 0.05$, followed by cluster

threshold correction (BrainVoyager QX v2.8, Brain Innovation) to our contrasts. This

included the Saccade > Fixation contrast shown in Fig. 4a/b, and the contrasts shown in

supplementary figures (Sensory/Memory phase activity > Baseline, Fig. S1;

Visual/Oculomotor Updating > Sensory/Memory contrast, Fig. S2).

Hypothesis testing: Analysis and statistical considerations.

In order to localize our site-of-interest testing, we used a similar approach to site

localization as in Song and Jiang [31]; Baltaretu et al. [24]; and Tsushima et al. [32] by

first applying a whole-brain contrast of interest: Saccade Different – Saccade Same at a

liberal p-value ($p < 0.05$) with cluster threshold correction (Fig. 3a). We then used

reduced the p-value in order to localize the centre of our *a priori* predicted parietal

activation (i.e., within the larger pCU/pCG area, Fig. 3b), and then used BrainVoyager

(BrainVoyager QX v2.8, Brain Innovation) to determine a cluster within a 1000 mm^3

volume ($p < 0.05$, cluster corrected; Table 1). Finally, we extracted the β-weights from

associated voxels around the peak voxel for the experimental data and saccade

localizer data (separately) in order to test our hypotheses within the parietal, ventral

precuneus (i.e., first, to determine the presence of a transsaccadic feature-specific

effect and, second, whether transsaccadic effects are task-specific). These analyses

were carried out for the 12 participants who survived our behavioural criteria for the task

and localizer.

Using the β-weights, we looked for specific directionality in the two predictions

that we tested (Fig. 2), so we used one-tailed repeated measures t-tests for each of the

two predictions (one to identify saccade/feature-related specificity and another to

determine transsaccadic task specificity). For the results of these analyses, we provided

all relevant t-statistics, p-values, and effect sizes (Cohen's d, determined using

G*Power) [87].

Psychophysiological interaction.

Lastly, we were interested in uncovering the functional network for saccade-related

updating during transsaccadic perception. In order to do this, we conducted

psychophysiological interaction (PPI) analysis [24,66,89,90] on data extracted from the

posterior parietal region that showed transsaccadic effects from the *Visual/Oculomotor*

Updating phase. To perform this analysis, we used three predictors: 1) a physiological

component (which is accounted for by z-normalized time courses obtained from the

seed region for each participant for each run); 2) a psychological component (task

predictors, i.e. for the Saccade and Fixation conditions, were convolved with a

hemodynamic response function), and 3) a psychophysiological interaction component

(multiplication of seed region z-normalized time courses with task model in a volume-by-

volume manner). For the psychological component, the *Saccade* predictors were set to

a value of '+1', whereas the Fixation predictors were set to a value of '-1'; all other

predictors were set to a value of '0' in order to determine the responses to Saccades as

compared to Fixation. (For greater detail, please see O'Reilly et al. [66] and Baltaretu et

al. [24].) We created single design matrices for each run for each of the 15 participants,

which were ultimately included in an RFX GLM. Then, we used the third,

psychophysiological interaction predictor to determine the regions that comprise the

functional network that show saccade-related modulations for feature updating. We

applied the p-value threshold (0.05) to the volumetric map and cluster threshold

correction to identify clusters of activation. We used similar region localization as Song

and Jiang [31], Baltaretu et al. [24], and Tsushima et al. [32] to pinpoint specific regions

of activation for this analysis.

References

1. Suzuki, S. & Cavanagh, P. Facial organization blocks access to low-level
features: An object inferiority effect. *J. Exp. Psychol. Hum. Percept. Perform.* **21**,
901-913 (1995).
2. Irwin, D. Information integration across saccadic eye movements. *Cog. Psychol*.
23, 420-456 (1991).
3. Melcher, D. Spatiotopic transfer of visual-form adaptation across saccadic eye
movements. *Curr. Biol.* **15**, 1745-1748 (2005).
4. Rayner, K. Eye movements in reading and information processing: 20 years of
research. *Psychol. Bull*. **124**, 372-422 (1998).
5. Melcher, D. Selective attention and the active remapping of object features in
trans-saccadic perception. *Vision Res*. **49**, 1249-1255 (2009).
6. Rayner, K., McConkie, G. & Zola, D. Integrating information across eye
movements. *Cogn. Pyschol*. **12**, 206-226 (1980).
7. Irwin, D., Yantis, S. & Jonides, J. Evidence against visual integration across
saccadic eye movements. *Percept. Psychophys*. **34**, 49-57 (1983).
8. Henderson, J., Pollatsek, A. & Rayner, K. Effects of foveal priming and
extrafoveal preview on object identification. *J. Exp. Psychol*. **13**, 449-463 (1987).
9. Prime, S., Niemeier, M. & Crawford, J. Transsaccadic integration of visual
features in a line intersection task. *Exp. Brain. Res*. **169**, 532-548 (2006).
10. Prime, S., Vesia, M. & Crawford, J. Cortical mechanisms for trans-saccadic
memory and integration of multiple object features. *Phil. Trans. R. Soc. B*. **366**,
540-553 (2011).

11. Melcher, D. & Colby, C. Spatiotopic transfer of visual-form adaptation across saccadic eye movements. *Curr. Biol*. **15**, 1745-1748 (2008).

12. Luck, S. & Vogel, E. The capacity of visual working memory for features and conjunctions. *Nature* **390**, 279-281 (1997).

13. Courtney, S., Ungerleider, L., Keil, K. & Haxby, J. Object and spatial visual working memory activate separate neural systems in human cortex. *Cereb. Cortex* **6**, 39-49 (1996).

14. Klier, E. & Angelaki, D. Spatial updating and the maintenance of visual constancy. Neuroscience **156**, 801-818 (2008).

15. Funahashi, S. Thalamic mediodorsal nucleus and its participation in spatial working memory processes: comparison with the prefrontal cortex. *Front. Syst. Neurosci*. **7**, 1-13 (2013).

16. Merriam, E., Genovese, C. & Colby, C. Spatial updating in human parietal cortex. *Neuron* **39**, 361-373 (2003).

17. Medendorp, W., Goltz, H., Vilis, T. & Crawford, J. Gaze-centered updating of visual space in human parietal cortex. *J. Neurosci*. **23**, 6209-6214, (2003).

18. Khan, A. *et al.* Optic ataxia errors depend on rempped, not viewed, target location. *Nat. Neurosci*. **8**, 418-420.

19. Morris, A., Chambers, D. & Mattingley, J. Parietal stimulation destabilizes spatial updating across saccadic eye movements. *Proc. Natl. Acad. Sci.* **104**, 9069-9074 (2007).

20. Dunkley, B., Baltaretu, B. & Crawford, J. Trans-saccadic interactions in human parietal and occipital cortex during the retention and comparison of object orientation. *Cortex* **82**, 263-276 (2016).

21. Subramanian J., & Colby, C. Shape selectivity and remapping in dorsal stream visual area LIP. *J. Neurophysiol*. **111**, 613-627 (2014).

22. Prime, S., Vesia, M. & Crawford, J. Transcranial magnetic stimulation over posterior parietal cortex disrupts transsaccadic memory of multiple objects. *J. Neurosci*. **28**, 6938-6949 (2008).

23. Prime, S., Vesia, M. & Crawford, J. TMS over human frontal eye fields disrupts trans-saccadic memory of multiple objects. *Cereb. Cortex* **20**, 750-772, (2010).

24. Baltaretu, B., Monaco, S., Velji-Ibrahim, J., Luabeya, G. & Crawford, J. Parietal cortex integrates saccade and object orientation signals to update grasp plans. *J. Neurosci.* **40**, 4525-4535 (2020).

25. Bellebaum, C., Hoffman, K.-P. & Daum, I. Post-saccadic updating of visual space in the posterior parietal cortex in humans. *Behav. Brain Res.* 163, 194-203 (2005).

26. Merriam, E. & Colby, C. Active vision in parietal and extrastriate cortex. *Neuroscientist* **11**, 484-493 (2005).

27. Taylor, P., Muggleton,N., Kalla, R., Walsh, V. & Eimer, M. TMS of the right angular gyrus modulates priming of pop-out in visual search: combined TMS-ERP evidence. *J. Neurophysiol.* 106, 3001-3009 (2011).

28. Kheradmand, A., Lasker, A. & Zee, D. Transcranial magnetic stimulation (TMS) of the supramarginal gyrus: a window to perception of upright. *Cereb. Cortex* **25**, 765-771 (2013).

29. Faillenot, I., Sunaert, S., Van Hecke, P. & Orban, G. Orientation discrimination of objects and gratings compared: an fMRI study. *Eur. J. Neurosci.* **13**, 585-596 (2001).

30. Baumann, O., Endestad, T., Magnussen, S. & Greenlee, M. Delayed discrimination of spatial frequency for gratings of different orientation: behavioral and fMRI evidence for low-level perceptual memory stores in early visual cortex. *Exp. Brain Res.* **188**, 363-369 (2008).

31. Song, J.-H. & Jiang, Y. Visual working memory for simple and complex features: An fMRI study. *Neuroimage* **30**, 963-972 (2006).

32. Tsushima, Y., Sawahata, Y. & Kazuteru, K. Task-dependent fMRI decoder with the power to extend Gabor patch results to natural images. *Sci. Rep.* **10**, 1-7 (2020).

33. Dassonville, P., Schlag, J. & Schlag-Rey, M. The frontal eye field provides the goal of saccadic eye movement. *Exp. Brain Res*. **89**, 300-310 (1992).

34. Gaymard, B. & Pierrot-Deseilligny, C. Neurology of saccades and smooth pursuit. *Curr. Opin. Neurol*. **12**, 13-19 (1999).

35. Krauzlis, R.J. The control of voluntary eye movements: New Perspectives. *Neuroscientist* **11**, 124-137 (2005).

36. Stone, S. *et al.* The assessment of visuo-spatial neglect after acute stroke. *J. Neurol.* **54**, 345-350 (1991).

37. Malhotra, P., Coulthard, E. & Husain, M. Role of right posterior parietal cortex in maintaining attention to spatial locations over time. *Brain* **132**, 645-660 (2009).

38. Pisella, L. *et al.* Right-hemispheric dominance for visual remapping in humans. *Phil. Trans. R. Soc. B. Biol. Sci.* **366**, 572-585 (2011).

39. Gorbet, D., Staines, R. & Sergio, L. Brain mechanisms for preparing increasingly complex sensory to motor transformations. *Neuroimage* **23**, 1100-1111 (2004).

40. Fernandez-Ruiz, J., Goltz, H., DeSouza, J., Vilis, T. & Crawford, J. Human parietal "reach region" primarily encodes intrinsic visual direction, not extrinsic movement direction, in a visual-motor dissociation task. *Cereb. Cortex* **17**, 2283-2292 (2007).

41. Chen, Y., Monaco, S. & Crawford, J. Neural substrates for allocentric-to-egocentric conversion of remembered reach targets in humans. *Eur. J. Neurosci.* **47**, 901-917 (2018).

42. Galletti, C., Fattori, P., Battaglini, P.P., Shipp, S. & Zeki, S. Functional demarcation of a border between areas V6 and V6A in the superior parietal gyrus of the macaque monkey. *Eur. J. Neurosci*. **8**, 30-52 (1996).

43. Galletti, C., Fattori, P., Kutz, D.F. & Gamberini, M. Brain location and visual topography of cortical area V6A in the macaque monkey. *Eur. J. Neurosci.* **11**, 575-582 (1999).

44. Pitzalis, S., Fattori, P. & Galletti, C. The human cortical areas V6 and V6A. *Vis. Neurosci*. **32**, e007 (2015).

45. Schraa-Tam, C. *et al.* Cortical and cerebellar activation induced by reflexive and voluntary saccades. *Exp. Brain Res.* **192**, 157-187 (2009).

46. Tosoni, A. *et al*. Resting-state connectivity and functional specialization in human medial parieto-occipital cortex. *Brain Struct. Func.* **220**, 3307-3321 (2014).

47. Schott, B. *et al.* Gradual acquisition of visuospatial associative memory
representations via the dorsal precuneus. *Hum. Brain Mapp.* **40**, 1554-1570
(2018).
48. Cavanna, A. & Trimble, M. The precuneus: a review of its functional anatomy and
behavioural correlates. *Brain* **129**, 564-583 (2006).
49. Zhang, S. & Li, C.-S. Functional networks for cognitive control in a stop signal
task. *Hum. Brain Mapp.* **33**, 89-104 **(**2012).
50. Dean, H., Crowley, J. & Platt, M. Visual and saccade-related activity in macaque
posterior cingulate cortex. *J. Neurophysiol.* **92**, 3056-3068 **(**2004).
51. Pearson, J., Heilbronner, S., Barack, D., Hayden, B. & Platt, M. Posterior
cingulate cortex: adapting behavior to a changing world. *Trends Cogn. Sci.* **15**,
143-151 (2011).
52. Jonikaitis, D., Szinte, M., Rolfs, M. & Cavanagh, P. Allocation of attention across
saccades. *J. Neurophysiol*. **109**, 1425-1434 (2013).
53. Funahashi, S., Bruce, C. & Goldman-Rakic, P. Neuronal activity related to
saccadic eye movements in the monkey's dorsolateral prefrontal cortex. *J.*
Neurophysiol. **65**, 1464-1483 (1991).
54. Callicott, J. *et al.* Physiological characteristics of capacity constraints in working
memory as revealed by functional MRI. *Cereb. Cortex* **9**, 20-26 (1999).
55. DeSouza, J., Menon, R. & Everling, S. Preparatory set associated with pro-
saccades and anti-saccades in humans investigated with event-related fMRI. *J.*
Neurophysiol. **89**, 1016-1023 (2003).
56. Pierrot-Deseilligny, C., Müri, R., Nyffeler, T. & Milea, D. The role of the human
dorsolateral prefrontal cortex in ocular motor behavior. *Ann. N. Y. Acad. Sci.*
1039, 239-251 (2005).
57. Tanaka, L. *et al.* The effects of TMS over dorsolateral prefrontal cortex on trans-
saccadic memory of multiple objects. *Neuropsychologia (Oxford)* **63**, 185-193
(2014).
58. Hopfinger, J., Buonocore, M. & Mangun, G. The neural mechanisms of top-down
attentional control. *Nat. Neurosci*. **3**, 284-291 (2000).
59. Thiel, C., Zilles, K. & Zink, G. Cerebral correlates of alerting, orienting and
reorienting of visuospatial attention: an event-related fMRI study. *Neuroimage* **21**,
318-328 (2004).
60. Japee, S., Holiday, K., Satyshur, M., Mukai, I. & Ungerleider, L. A role of right
middle frontal gyrus in reorienting of attention: a case study. *Front. Syst.*
Neurosci. **9**, 1-15 (2015).
61. Vossel, S., Thiel, C. & Fink, G. Cue validity modulates the neural correlates of
covert endogenous orienting of attention in parietal and frontal cortex.
Neuroimage **32**, 1257-1264 (2006).
62. Ptak, R. The frontoparietal attention network of the human brain: Action, saliency,
and a priority map of the environment. *Neuroscientist* **18**, 502-515 (2012).
63. Petrides, M., Alivisatos, B., Meyer, E. & Evans, A.Fu nctional activation of the
human frontal cortex during the performance of verbal working memory tasks.
Proc. Natl. Acad. Sci. **90**, 878-882 (1993).
64. Cohen, J. *et al.* Temporal dynamics of brain activation during a working memory
task. *Nature* **386**, 604-608 (1997).

65. Treserras, S. *et al.* Transition from rest to movement: brain correlates revealed
by functional connectivity. *Neuroimage* **48**, 207-216 (2009).
66. O'Reilly, J., Woolrich, M., Behrens, T., Smith, S. & Johansen-Berg, H. Tools of
the trade: Psychophysiological interactions and functional connectivity. *Soc.*
Cogn. Affect. Neurosci. **7**, 604–609 (2012).
67. Tootell, R., Silverman, M., Hamilton, S., Switkes, E. & De Valois, R. Functional
anatomy of macaque striate cortex. V. Spatial frequency. *J. Neurosci.* **8**, 1610-
1624 (1988).
68. DeYoe, E. *et al.* Mapping straite and extrastriate visual areas in human cerebral
cortex. *Proc. Natl. Acad. Sci.* **93**, 2382-2386 (1996).
69. Tootell, R. *et al.* Functional analysis of V3A and related areas in human visual
cortex. *J. Neurosci.* **17**, 7060-7078 **(**1997).
70. Huntgeburth, S. & Petrides, M. Morphological patterns of the collateral sulcus in
the human brain. *Eur. J. Neurosci.* **35**, 1295-1311 (2012).
71. Nakamura, K. & Colby, C. Updating of the visual representation in monkey striate
and extrastriate cortex during saccades. *Proc. Natl. Acad. Sci*. **99**, 4026-4031
(2002).
72. Kauffman, L., Ramanoël, S., Guyader, N., Chauvin, A. & Peyrin, C. Spatial
frequency processing in scene-selective cortical regions. *Neuroimage* **112**, 86-95
(2015).
73. Tanji, J. & Evarts, E. Anticipatory activity of motor cortex neurons in relation to
direction of an intended movement. *J. Neurophysiol.* **39**, 1062-1068 (1976).
74. Donner, T., Siegel, M., Fries, P. & Engel. A. Buildup of choicie-predictive activity
in human motor cortex during perceptual decision making. *Curr. Biol.* **19**m, 1581-
1585 (2009).
75. Kaufman, M., Churchland, M. & Shenoy, K. The roles of monkey M1 neuron
classes in movement preparation and execution. *J. Neurophysiol* **110**, 817-825
(2013).
76. Curtis, C.&d D'Esposito, M. Persistent activity in the prefrontal cortex during
working memory. *Trends Cog. Sci.* **7**, 415-423 (2003).
77. Gaymard, B. *et al.* Cortical control of saccades. *Exp. Brain Res.* **123**, 159-163
(1998).
78. Duhamel, J., Colby, C. & Goldberg, M. The updating of the representation of
visual space in parietal cortex by intended eye movements. *Science* **255**, 90–92
(1992).
79. Oliva, A. & Torralba, A. Building the gist of a scene: The role of global image
features in recognition. *Prog. Brain Res.* **155**, 23-36 (2006).
80. Melcher, D. Predictive remapping of visual features precedes saccadic eye
movements*. Nat. Neurosci*. **10**, 903-907 (2007).
81. Hubel, D. & Wiesel, T. Receptive fields and functional architecture of monkey
striate cortex. *J. Physiol.* **195**, 215-243 (1968).
82. Nauhaus, I., Nielsen,K., Disney, A. & Callaway, E. Orthogonal micro-organization
of orientation and spatial frequency in primate primary visual cortex. *Nat.*
Neurosci. **15**, 1683-1690 (2012).

83. Monaco, S. *et al.* Functional magnetic resonance adaptation reveals the involvement of the dorsomedial stream in hand orientation for grasping. *J. Neurophysiol.* **106**, 2248-2263 (2011).

84. Valyear, K., Culham, J., Sharif, N., Westwood, D. & Goodale, M. A double dissociation between sensitivity to changes in object identity and object orientation in the ventral and dorsal visual streams: A human fMRI study. *Neuropsychologia* **44**, 218-228 (2006).

85. Melcher, D. & Morrone, C. Spatiotopic temporal integration of visual motion across saccadic eye movements. *Nat. Neurosci.* **6**, 877-881 (2003).

86. Ten Brink, A., Fabius, J., Weaver, N., Nijboer, T. & Van der Stighel, S. Trans-saccadic memory after right parietal brain damage. *Cortex* **120**, 284-297 (2019).

87. Faul, F., Erdfelder, E., Buchner, A. & Lang, A. Statistical power analyses using G*Power 3.1: Tests for correlation and regression analyses. *J. Behav. Res. Methods* **41**, 1149-1160 (2009).

88. Talairach, J. & Tournoux, P. Co-planar stereotaxic atlas of the human brain. New York: Thieme Medical. (1988).

89. Friston, K. et al. Psychophysiological and Modulatory Interactions in Neuroimaging. *Neuroimage* **6**, 218–229 (1997).

90. McLaren, D., Ries, M., Xy, G. & Johnson, S. A generalized form of context-dependent psychophysiological interactions (gPPI): a comparison to standard approaches. *Neuroimage* **61**, 1277-1286 (2012).

91. Sugiura, M., Shah, N., Zilles, K. & Fink, G. Cortical representations of personally familiar objects and places: functional organization of the human posterior cingulate cortex. *J. Cog. Neurosci.* **17**, 183-198 (2005).

92. Berman, R. *et al.* Cortical networks subserving pursuit and saccadic eye movements in humans: an fMRI study. *Hum. Brain Mapp.* **8**, 209-225 (1999).

93. Braver, T. *et al.* A parametric study of prefrontal cortex involvement in human working memory. *Neuroimage* **5**, 49-62 (1997).

94. Lim, J., Tan, J., Parimal, S., Dinges, D. & Chee, M. Sleep deprivation impairs object-selective attention: a view from the ventral visual cortex. *PloS one* **5**, e9087 (2010).

95. Ferri, F., Frassinetti, F., Ardizzi, M., Costantini, M. & Gallese, V. A sensorimotor network for the bodily self. *J. Cog. Neurosci.* **24**, 1584-1595 (2012).

96. Snow, J. *et al.* Bringing the real world into the fMRI scanner: Repetition effects for pictures versus real objects. *Sci. Rep.* **1**, 1-10 (2011).

97. Podrebarac, S., Goodale, M. & Snow, J. Are visual texture-selective areas recruited during haptic texture discrimination? *Neuroimage* **94**, 129-137 (2014).

Figure Legends

Figure 1. Experimental paradigm and eye movement traces. **A.** An example trial is

shown (0.7 cycles per degree, cpd, left fixation) with the four possible conditions:

Fixation Same, Fixation Different, Saccade Same, and Saccade Different. Each 22 s

trial had three major phases: 1) *Sensory/Motor,* for the first presentation of the stimulus

at one of the two possible spatial frequencies (0.7 or 1.1 cpd) while gaze could be to the

left or right; 2) *Visual/Oculomotor Updating*, for the second presentation of the stimulus

at the same spatial frequency (e.g., 0.7 cpd for first and second stimulus presentations;

Same condition) or different (e.g., 0.7 cpd for first presentation and 1.1 cpd for second

stimulus presentation, or vice versa; *Different* condition) while participants maintained

fixation on the same cross (*Fixate* condition) or made a directed saccade (*Saccade*

condition); and 3) *Response*, for the button press response period where an indication

of whether the spatial frequency across the two stimulus presentations was the same

('R') or different ('N'). **B.** An example eye position trace (º) for an example fixation and

saccade trial. In this figure, each of the two trials started with initial fixation on the right

(13.5º from center), then diverged after the initial presentation of the stimulus (1st visual

stimulus) and mask (black vertical bar), whereby gaze remained fixed for the fixation

trial or moved to the other fixation cross position (-13.5º from center) for the saccade

trial, where it remained after the second stimulus presentation (2nd visual stimulus), and

button press period.

Figure 2. Predictions used to test transsaccadic feature- and task-specificity. Prediction

1: For an area that shows saccade-related feature-specific modulations, there should be

a larger response 1) for a change in spatial frequency than a repetition, and 2) when a

saccade is produced than for fixation (i.e., (Saccade Different – Saccade Same) >

(Fixation Different – Fixation Same); SD, SS, FD, FS, respectively). Prediction 2: The

saccade-related effects in this region should also show task-specificity – i.e., there

should be larger modulations in the Saccade condition than the Fixation condition for

the experimental task compared with the independent saccade localizer (i.e., (Task,

Saccade – Task, Fixation) > (Saccade Localizer, Saccade – Saccade Localizer,

Fixation); TS, TF, S, F, respectively).

Figure 3. Testing Prediction 1: Saccade specificity in PPC during the *Visual/Oculomotor*

Updating phase. **A.** Upper panels: Voxelwise statistical map from an RFX GLM (n=15)

for Different > Same in the *Saccade* condition (orange) is overlaid on inflated cortical

surface renderings of right hemisphere of example participant (lateral view on left and

medial view on right). Regions in medial precuneus/posterior cingulate gyrus

(PCu/pCG) and frontal (MFG, middle frontal gyrus) cortex show saccade-related

feature-specific effects. **B.** Specific parietal site localization was performed, where the

site-of-interest is shown in white (i.e., ventral PCu, vPCu). The white dot corresponds to

location of peak voxel(s) of the site-of-interest. **C.** β-weights were extracted from vPCu,

analyzed, and plotted in bar graphs (n=12). In order to meet the first criterion (Prediction

1) to be considered a putative *transsaccadic integrator,* precuneus would need to show

a larger feature-related β-weight difference for the *Saccade* and *Fixation* conditions

(SD-SS: Saccade Different – Saccade Same; FD-FS: Fixation Different – Fixation

Same). As shown in the left bar graph, there is a (significantly) greater feature-related

difference for the *Saccade* than for the *Fixation* condition [(SD - SS) > (FD - FS)].

Values are mean differences ± SEM analyzed using one-tailed repeated measures t-

tests. (Please see Results for specific statistical values.)

Figure 4. Testing Prediction 2: Task specificity in PPC during the *Visual/Oculomotor*

Updating phase. **A.** Upper panels: Voxelwise statistical maps from an RFX GLM (n=15)

for Saccade > Fixation in the experimental task (orange; FDR (q < 0.05) and cluster

correction) versus the localizer (n=12; fuchsia; FDR (q < 0.05) and cluster correction)

are overlaid onto inflated cortical surface renderings of example participant (left

hemisphere on the left, right hemisphere on the right; upper panels showing the lateral

views, and lower panels showing medial views). There is substantial overlap in activity

in medial occipital regions for task and localizer. However, (experimental) task-specific

saccade modulations can be observed in particular occipital (middle occipital gyrus,

MOG), parietal (medial precuneus, mPCu; inferior parietal lobe, IPL; and superior

parietal lobule, SPL), and frontal saccade regions (medial superior frontal gyrus, mSFG

– likely pre-supplementary eye field; dorsal precentral sulcus, PCSd – likely frontal eye

field). **B.** The site-of-interest (i.e., ventral PCu, vPCu) is shown in white on the sagittal

view of the average brain of all the participants, which lies outside of the saccade-

specific activation for the task (orange) or localizer (fuchsia). The white dot corresponds

to location of peak voxel(s) of the site-of-interest. **C.** β-weights were extracted from

vPCu, analyzed, and plotted in bar graphs. To meet the second criterion (Prediction 2),

the observed saccade-related modulations for spatial frequency would have to be

greater for the task than for a separate saccade-only localizer. The β-weight difference

between *Saccade* and *Fixation* conditions is plotted for the task (TS-TF, Task, Saccade

and Task, Fixation) and localizer (S-F, Saccade Localizer, Saccade and Saccade

Localizer, Fixation). Precuneus also showed a (significant) saccade-related task-specific

effect [(TS - TF) > (S – F)]. Values are mean differences ± SEM analyzed using one-

tailed repeated measures t-tests. (Please see Results for specific statistical values.)

Figure 5. Functional connectivity network (A) and model (B) engaged in transsaccadic

updating of spatial frequency. **A.** Transverse slices through the average brain of all

participants are shown from most superior to inferior along the vertical axis. 'Z'

components of each region's Talairach coordinates are reported for reference. Shown in

cyan is the seed region, right precuneus (vPCu). Using a Saccade > Fixation contrast,

the psychophysiological interaction (PPI) analysis revealed statistically significant

communication between precuneus and right precentral gyrus (PCG, likely primary

motor cortex), as well as left lingual gyrus (LG) and medial occipitotemporal sulcus

(MOtS). This suggests that parietal precuneus communicates with frontal motor and

early/late object perception regions. **B.** A network model proposed to highlight the

potential interactions between right precuneus and saccade, memory, and object

perception regions. Expt. Region refers to a region that showed activation in our task;

Non-expt. Region refers to a region we did not find in our task (e.g., early visual cortex,

EVC); PPI connection refers to a connection derived from our PPI results; Hyp.

Connection refers to a hypothesized connection between regions in our model; Inhib.

Hyp. Connect. refers to the possible inhibitory connection between the medial PCu/pCG

region and MFG/dorsolateral prefrontal cortex (dlPFC).

Figure 1

872 **Figure 2**

Prediction 1:
Transsaccadic
Frequency Updating

Prediction 2:
Task-Specific
Saccade Effects

SD: Saccade Different

SS: Saccade Same

FD: Fixation Different

FS: Fixation Same

TS: Task, Saccade

TF: Task, Fixation

S: Saccade Localizer, Saccade

F: Saccade Localizer, Fixation

Figure 3

(**a**) Frequency Change Sensitivity Following Saccades

Lateral   **RH**   Medial

MFG

PCu/pCG →

········ Central Sulcus   = = · Intraparietal Sulcus

⬤ Sacc. Different > Sacc. Same   ⬤ Sacc. Same > Sacc. Different

(**b**) Parietal Site-of-Interest   (**c**) P1: Transsaccadic Feature Updating

RH
Ant.
vPCu

X = 14

SD-SS   FD-FS

*

Figure 4

(**a**) Saccade-Related Activity (Experiment vs. Localizer)
(Visual/Oculomotor Updating Phase)

LH   **RH**

Lateral

PCSd
SPL
IPL
MOG

········· Central Sulcus
= = Intraparietal Sulcus

Medial

mPCu
SPOC
Cu
LG

mSFG

mPCu
SOG
Cu
LG

🟠 Task (Sacc. > Fix.)
🟣 Localizer (Sacc. > Fix.)

(**b**) Site-of-Interest: vPCu

RH
Ant.

vPCu

X = 14

(**c**) P2: Task-Specific
Saccade Effects

β

TS-TF   S-F

0.06
0.04
0.02
0
-0.02
-0.04

*

910   **Figure 5**

Functional Connectivity with Right Ventral Precuneus
(Visual/Oculomotor Updating Phase)

923 **Table 1.** Names, Talairach coordinates, and sizes of regions of interest (ROIs) observed

924 during the *Visual/Oculomotor Updating* phase.

Region of Interest	Talairach coordinates						Active Voxels	References
	x	y	z	Std x	Std y	Std z	*n* voxels	
RH ventral Precuneus	14	-55	29	1.3	1.3	2.8	142	[48,91]
RH posterior Cingulate Gyrus	9	-49	14	2.5	2.8	2.1	641	[91,92]
RH Middle Frontal Gyrus	39	17	37	2.7	2.5	2.7	790	[93,94]
RH Precentral Gyrus	51	-8	34	2.8	2.7	2.8	876	[95]
LH Lingual Gyrus	-13	-65	-11	2.4	2.4	2.6	618	[96]
LH Medial Occipitotemporal Sulcus	-18	-77	-5	2.4	2.7	2.4	665	[97]