Local Gazetteers Reveal Contrasting Patterns of Historical Distribution Changes between Apex Predators and Mesopredators in Eastern China

Kaijin Hu[1], Chunhui Hao[2]

1. School of Sociology and Anthropology, Sun Yat-sen University, Guangzhou, China

2. School of Life Sciences, Sun Yat-sen University, Guangzhou, China

Correspondence author: Kaijin Hu, Email: hukaijin01@126.com

Abstract:

Background: Humans have been causing the sixth wave of mass extinction of biodiversity. The situation of predators, especially of carnivores, is a key indicator of biodiversity, and the mesopredator release is a typical phenomenon in ecosystem recess. Local gazetteers(地方志) are a rich resource for historical biodiversity research. But there are obvious biases in previous studies focusing in only presence records and neglecting the absence records. We recollected and analyze the records by fixed methods to research historical change of biodiversity.

Methods: Innovatively, this research uses both presence and absence records from local gazetteers to reconstruct the distribution of 8 kinds of mammalian predators (i.e. tiger, leopard, bear, wolf, fox, civet, dhole and mustelid) in eastern China from 1573 A.D. to 1949 A.D. (sorted into 4 periods). Then we analyze the distribution changes, the relation between animals and the influence from human.

Results: With the human population booming, the distribution of large/apex predators retreated overall, but the distribution of mesopredators expanded overall. Besides, the predator distribution relations formed different groups, not match the body size groups totally.

Conclusions: Based on our reconstructions, we provide direct proof of human disturbance on mammal distribution in China, and extend support for the mesopredator release hypothesis. And we tested the predator guild hypothesis by proving that dhole were successful to coexist with tiger and leopard in history.

Prospects: We have also collected records for other wild animals from local gazetteers in China. Based on the collection, we have built the Database of Wild Mammal Records in Chinese Local Gazetteers and are building the Database of Wild Bird Records in Chinese Local Gazetteers. We aim to continue relevant studies using these databases in the future.

Keywords: Local gazetteer; Historical distribution; Mesopredator release; Predator Guild

ORCID:
Kaijin Hu https://orcid.org/0000-0001-9639-3774

1.Introduction

It has been widely accepted that Earth is facing a biodiversity crisis (Rahbek & Colwell 2011). In the past 500 years, a wave of extinction and declines in local animal population have been recorded, with both the rate and magnitude of this recent period of extinction being comparable to the five previous periods of mass extinction (Pimm *et al.* 1995; He & Hubbell 2011; Dirzo *et al.* 2014). Unlike these five previous periods of mass extinction which were primarily driven by global changes in climate and atmospheric-ocean chemistry, volcanism, and bolide impacts (Barnosky *et al.* 2011), the modern loss of biodiversity is considered as the consequence of human socioeconomic activities (Dirzo & Raven 2003; Ceballos *et al.* 2015). Humans have been causing the sixth wave of extinction through changing global climate, overkilling species, co-opting resources, introducing exotic species, spreading infectious diseases, and destructing and modifying habitat (Leakey & Roger Lewin 1996; MacPhee *et al.* 1999; Dirzo & Raven 2003; Thomas *et al.* 2004; Martin 2005; Jackson 2008; Wake & Vredenburg 2008; Ceballos *et al.* 2015; Yap *et al.* 2015).

Although there is no doubt that the impact of the recent "sixth period of mass extinction" has extended across distinctive taxonomic groups, each species responds differently to climatic shifts, habitat fragmentation, and decline in species abundance caused by anthropogenic disturbance (Cardillo *et al.* 2008; Lorenzen *et al.* 2011). Moreover, the life history characteristics of different species are also to some extent correlated with their risk of being threatened by human activities (Davidson *et al.* 2009; Öckinger *et al.* 2010; Cardillo & Meijaard 2012). Despite this, such studies usually had poor predictive power to reliably link species' vulnerability with their life history traits (Jablonski 2001, 2008; Pocock 2011). Even relatively well-established patterns of correlation, such as large body size being known to be related strongly to extinction risk in mammals, still failed to accurately predict the extinction risk of a specific species affected by human activities. This is due to (*i*) multiple ecological factors and human activities interact to affect species that differ by orders of magnitude in life history traits (Davidson *et al.* 2009), and (*ii*) human disturbance to specific species is usually regulated by the local community structures and population dynamics in which the species is embedded (Kitahara & Fujii 1994; Gill 2007; Tylianakis *et al.* 2007). A phenomenon termed "mesopredator release" represents one of the best-referred examples. Large-bodied predators were persecuted by humans, but the abundance of mesopredators might undergo dramatic increases as a result of trophic cascading effects in communities (Prugh *et al.* 2009; Wallach *et al.* 2015). Consequently, in view of the complex synergy of both intrinsic species traits and human's role at different spatial-temporal and ecological scales, more specific quantitative assessments of the relationships between local extinction of endangered species and anthropogenic factors are urgently required (Fritz *et al.* 2009; Dirzo *et al.* 2014; Sandom *et al.* 2014; Wan *et al.* 2019; Teng *et al.* 2020).

In China, the distribution of wild animals has changed greatly over time, with many formerly widely distributed wild animals are now endangered or even extinct (Chatterjee et al. 2012; Qian *et al.* 2020). Meanwhile, the population in China has

increased significantly since the introduction of American crops in the 16th century,
which triggered prosperous socio-economic activities which may be responsible for
the extinction of wild animals in China (Li 1998; Cao 2001). China has a long history
of using local gazetteers to record momentous events and situations, including those
related to local wildlife, and previous researchers had published numerous collection
list of animal records (He 1993; 1997; Wen 2009; 2013; 2018). By exploring these
historical records in various ways, researchers have attempted to reconstruct the
spatio-temporal distribution and extinction of engendered animals in China, and then
disentangle the relative roles of anthropogenic and climate changes in shaping their
distribution and extinction (Jiang *et al.* 1995; Li *et al.* 2002, 2020; Zhou & Zhang
2013; Li *et al.* 2015; Turvey *et al.* 2015, 2017; Wan & Zhang 2017; Nüchel *et al.* 2018;
Zhao *et al.* 2018; Wan *et al.* 2019; Qian *et al.* 2020; Teng *et al.* 2020). However, such
studies have relied on biased species observation data which only focused on the
"presence" records of local animals but neglected the "absence" records. For each
species, they often defined the time of extirpation as the last observation (or called
last occurrence) time which is interpreted from the most recent record there. But the
collection of data from gazetteer-rich and gazetteer-poor regions/periods may cause
obvious biases in the estimation of the distribution of wild animals, for the
distribution of gazetteers is neither regular in space nor in time. For example, there are
few gazetteers in war-torn or sparsely populated areas, and there were only one or two
gazetteer from many regions throughout history. Therefore, to accurately evaluate the
spatio-temporal distribution and extinction of wild animals in China, we need to take
into account these biases in the data collected, and then do the re-collection job to
track changes in biodiversity and analyze how human activities would have affected
such changes (Boakes *et al.* 2010).

Here, we first recollected the presence and absence data of 8 kinds of wild
mammals in eastern China from 1573 A.D. to 1949 A.D. We chose this time period
and region to avoid potential biases, because (*i*) eastern China was the main activity
area of ancient Chinese people; (*ii*) during this period, China experienced a rapidly
growing human population; (*iii*) a greater quantity of records of wild animals in
regional gazetteers   The data were re-organized from scanned local gazetteers
(county-level or prefectural-level) in the Chinese Digital Gazetteer Database
(http://x.wenjinguan.com/) and *The Series of Rare Local Gazetteers in National*
Library（Fu *et al.* 2016）. All 8 of the mammals we studied are predators of distinctive
body sizes, are widely distributed across gazetteer records and are also easily
distinguishable in traditional folk taxonomy. Furthermore, by performing new
analytical techniques (*Methods*), we reconstruct the spatial and temporal distribution
patterns of 8 predators with both presence and absence records. The ultimate objective
of this study was to quantitatively estimate how specifically human socioeconomic
activities would have affected the distribution and extirpation of these 8 predators in
eastern China.

2.Methods:

Overview:

We chose the eastern part of China as research area, and the temporal span 1573-1949 as research period, for there are rich amount of local gazetteers preserved, and the human population experienced a huge increase. We collected both presence and absence data of each target mammal predator from local gazetteers, and sorted them into 4 periods. In each period, we used the tool Inverse Distance Weighed (IDW) to reconstruct the distribution probability layers by ArcGIS 10.3. Then we used 20km*20km grids to extract the continuous value in those layers and the human population density layers, to analyze the distribution changes, the relation between animals and the influence from human. (see *Fig. 1*)

(Insert Fig.1)

2.1 Data collection

We collected the data from the Chinese Digital Gazetteer Database (http://x.wenjinguan.com/) and *The Series of Rare Local Gazetteers in National Library*（Fu *et al.* 2016）. Previous studies used only presence points to reconstruct the presence and pseudo-absence(actually data missing) maps, then extracted the binary value in each grid to analyse with limited maths methods. Different to above way, we used both presence and absence points to reconstruct the probability maps. Then we extracted the continuous value in each grid to analyse.Previous studies collect the presence records of the target animal, but in this research, we collected the valid animal diversity records, and acquire the presence and absence records of the target animal. For example, there are two animal records in different local gazetteers(see row 1 in Fig.1), A recorded "Tiger, Leopard, Deer, Roe Deer .... Hedgehog, Vole, Ibex." (Wolf not existed), and B recorded "Cattle, Horse ... Wolf, Dog, Badger, Rabbit, Mouse" (Wolf existed). The previous studies would output "There were wolves in B" merely, but our study would output "There were wolves in B, and there were no wolves in A" (see row 2 in *Fig.1*).

There are two kinds of invalid records we excluded. The first kind recorded only livestock, and the second kind did not record comprehensively, but recorded only several large mammals roughly. To avoid above invalid records, we filtered the gazetteer records by the following rule: a valid record should record any wild mammal of which the body length is below 1 meter (including those are not analyzed in this research, like macaca monkey, gibbon or wild rabbit) (see *Table 1*).

(Insert Table.1)

We select 8 kinds of wide distributed predators, i.e. tiger, leopard, bear, wolf, fox, civet, dhole, mustelid, to research. And we match the folk animal kinds to the modern species in China(Smith & Xie 2009) (see *Table. S1*), the the folk toxonomy animal kind can be matched to one or several modern scientific taxonomy species. We sorted tiger, leopard, bear into the large/apex predator group, and the rest into the mesopredator group, according to the definition of megafauna, body mass>45kg or occupying the relevant ecological niche (Roberts *et al.* 2001; Barnosky 2008; Boulanger & Lyman 2014; Villavicencio *et al.* 2016; Moleon *et al.* 2020).

2.2 Spatial processing:

(Insert Fig.2)

We selected 22 province-level regions in eastern continent China as the research
/area (excluding Tibetan Plateau part in Sichuan and Yunnan) which roughly
corresponds with the eastern region of the Heihe-Tengchong Line (see *Fig.2*), ,
containing approximately 90% of the human population of China from ancient to
modern eras (Hu 1935). Although the names and areas of many counties have
changed since the Qing Dynasty, we were able to locate the data on modern maps by
using the current local government locations to identify the corresponding locations of
the local gazetteers.

We used the Inverse Distance Weighted (IDW) tool to reconstruct the
distribution based on the valid records using ArcGIS Desktop 10.3. This tool has been
widely applied to species distribution reconstruction (Roberts *et al.* 2004; Hijmans
2012; Takahashi *et al.* 2014; Barbosa 2015; Areias-Guerreiro *et al.* 2016; Palacio *et al.*
2020). The interpolation value in each grid was generated by the the nearest
12(default value) records, including presence records or absence records. The main
assumption is that species are more likely to be found closer to presence points and
farther from absences (i.e. spatial auto-correlation), and the local influence of points
(weighted average) diminishes with distance (Palacio *et al.* 2020）. The result value
means the probability of animal occurrence, in [0,1]. Besides the accuracy of data, the
smoothy probability value is continuous, so we can analyse the distributions much
subtler.

Finally, we use the fishnet tool to build 10819 20km*20km grids (in WGS 1984
World Mercator projected coordinate system), to extract the data from the layers.

2.3 Temporal processing:

To keep the balance between the diachronical-spatial dimensions, we selected
1573-1949 as the research span, and sorted the valid points into 4 periods according to
the reigning emperor, since many of local gazetteers are labeled under the reigning
emperor's title without specifying the year, and adopted the later one when there is
more than one record in the same location in one period.

Period 1 covers the span of 1573-1722, with 706 valid data points; Period 2
covers the span of 1723-1820, with 707 valid data points; Period 3 covers the span of
1821-1911 , with 787 valid data points; and Period 4 covers the span of 1912-1949,
with 451 valid data points. This layers are cross sections and based on both
presence/absence points, so it is appropriate that spans are different in periods. This
research has excluded the modern distribution of mammalian predators because their
distribution was greatly reduced as a consequence of a campaign by the Chinese
government to hunt the "harmful beasts" since the 1950s.

Since the America crops were introduced to China via Spanish Philippines in the
late 16[th] century and widely expanded in later centuries, the Chinese human

population(which was severely damaged in wars in early 17th century) recovered rapidly and surpassed the historic apex (Li 1998; Cao 2001). According to the HYDE 3.2 database(Goldewijk *et al.* 2017), the average human population density(HPD) increased from 34.9 persons/km² in 1720 to 145.3 persons/km² in 1950.

2.4 Relations of predator distribution：

We used the Kendall's tau-b coefficient to measure the relations between each predator in SPSS 22. This statistic is a measure of the strength and direction of associations between two Ordinal variables. Kendall's tau-b treats the variables symmetrically and ranges in [-1,1], with the ± symbol reflecting the direction of correlation and the absolute value indicating the strength of association.

$$\text{tau-b} = (N_c - N_d)/[(N_c + N_d + T_x)(N_c + N_d + T_y)]^{0.5}$$

N_c is the number of concordant pairs, N_d is the number of discordant pairs. T_x & T_y are the number of tied pairs on each variable.

2.5 Anthropogenic factor influence on predator distribution:

The official censuses in 1820, 1911, 1953 were widely recognized by academy and were the base of Chinese population reconstruction. We selected the nearest layers (1820, 1910 & 1950) in HYDE 3.2 database(Goldewijk *et al.* 2017) to match Period 2, 3 & 4, and selected layer in 1720 to match the breakpoint between Period 1 and Period 2.

We measured the influence of human population density(HPD) on predator distribution between each predator by Somers' d_{yx} statistic in software SPSS 22. This statistic is a measure of the influence strength of an ordinal variable on the other ordinal variable. The d_{yx} ranges in [-1,1], with the ± symbol reflecting the direction of influence and the absolute value reflects the strength (Somers 1962).

$$d_{yx} = (N_c - N_d)/(N_c + N_d + T_y)$$

N_c is the number of concordant pairs, N_d is the number of discordant pairs. T_y is the number of tied pairs on dependent variable.

3.Result：

3.1 The change of distribution area

We reconstructed the distribution of target predators and human population density (HPD) (see *Table S2*). In the large/apex predator group, distribution of two largest predators, tigers and bears retreated, while the smaller one, leopard, remained almost unchanged; by contrast, the distribution of predators in mesopredator group expanded in varying degrees (see *Fig.3, Fig.4 & Table S3*).

(Insert Fig.3, Fig.4)

3.2 Relations between each predator and others

(Insert Fig.5)

According to the relations of distributions between predators (see *Fig.5 & Table.*
S4), the predators in the large/apex predator group displayed strongly positive
correlations with each other in all periods. In the mesopredator group, wolves and
foxes were highly correlated, the correlation between wolves/foxes and large/apex
predators were negative overall; while for dholes and civets, they were positive
correlated with the large/apex predators and each other, but weakly negative with
wolves/foxes overall; for mustelids, the values with leopards and bears were negative
in earlier periods but turn positive later, and the value with other predators(including
tigers) were positive.

3.3 Anthropogenic influence on predator distribution

(Insert Fig.6)

The results (see *Fig.6 & Table. S5*) show that HPD exerted negative influence
upon the distributions of each large predator during each period, the influences on
tigers, leopards and bears kept are clearly negative throughout. But for mesopredators:
(1)wolves, foxes and civets were slightly positive in early periods, but became
negative in later periods; (2)dholes were negetive in all periods, like large predators;
and (3) mustelids were strongly positive in period 1 to period 3, but turn almost not
related and not significant in period 4.

3.4 The average probability in different HPD area in each period

We sorted the area into groups by HPD value to calculate the average probability
of predator distribution in these area groups (see *Fig.7*). In all periods, the average
probability of large/apex predators and dholes decreased with higher HPD in general.
By contrast, the trends of other mesopredators were increasing or inverse-U in earlier
periods, but also turn decreasing in the later periods.

(Insert Fig.7)

4.Discussion：

The reconstruction of distributions using both presence and absence data is more
accurate than previous studies (Zhou & Zhang 2013; Li *et al.* 2015; Turvey *et al.* 2015;
Wan *et al.* 2019; Teng *et al.* 2020) with have only used presence data (see *Fig.8*),
especially with the last observation method (see *Fig.9 & Fig.10*). It overcomes several
disadvantages of previous studies :

(1)Sampling bias: there are few gazetteers in the area/period with social
turbulence or few human population, causing the presence records were also few
there/then (*Fig.8.a,*). we fixed this bias by the IDW.

Besides, there are few local gazetteers which were written before 1573 A.D. and
still preserved today. That means the presence records (also absence records) are rare
in this periods. And the Chinese human population experienced several fluctuations in
history. It is inappropriate to infer the decline trend with the sample with these biases.
So we did not analyze this period in this research.

(2) New data: if there are new data, the reconstruction might change obviously
(e.g. there are much more new found records in Wen(2018)'s new research than his
previous research (Wen 2009) then the reconstruction changed obviously), for the last
observation method, it would lead to inverse interpolation. While in our methods, new
data will subtlize the reconstruction.

(3) Subjective: how large area (circle or square, 50km*50km or 100km*100km,
etc.) does a data point represent is determined by subjectivity of different researchers
(Zhou & Zhang 2013; Turvey *et al.* 2015; Wan *et al.* 2019; Teng *et al.* 2020), leading
to different result of distribution area, but in this research, it is replaced by objective
interpolation of IDW.

(4). Monotonic recession prejudged: Last observation method can not reflect the
expansion or recovery of animal distribution, but only monotonic recession, it is a
kind of prejudge. While in our method, the expansion or recovery can also be noted.

Above all, there are several biases in previous studies, and the methods in this
study mended them.

(Insert Fig.8, Fig.9 & Fig.10)

We exemplify the contrast of two methods by the distribution of wolf(see *Fig.9*
& Fig.10). The results in last observation method show a retracting trend, but the
result values are much different between 100km*100km square grids and
50km*50km square grids. In contrast, the result in IDW method shows a expanding
trend.

The distribution of wild animals is significantly shaped by anthropogenic
disturbances throughout the long history of China (Chatterjee *et al.* 2012). We found
that the distribution of large/apex predators retreated over successive periods of time,
supporting similar views of previous studies (He 1993; 1997; Wen 2009; 2013; 2018;
Wan *et al.* 2019; Teng *et al.* 2020) , and reflecting the recession of the eco-system
over the same time frame. The decline trend of bears was severe than it of tigers. but
the bears are still living in several protected area in research area, while tigers are
already extirpated. We infer that the reason is the different home range sizes. Tigers,
with large home range, would appear in wide sink-habitat out of source-habitat and be
exposed to anthropogenic dangers; while Bears' habitat were majorly source-habitat,
and they would survive when the small range habitats are preserved .

Not to be noted by previous studies, we find the distributions of mesopredator
expanded in eastern China, supporting the mesopredator release hypothesis (Prugh *et*
al. 2009), which said that the population of mesopredator would grow rapidly after
the constrains from large/apex predators were withdrawn.

The correlations among the three large/apex predators (tigers, leopards and bears)
were strongly positive during all periods, suggest that these species have similar
environmental requirements. The distributions of dholes and civets were also strongly
and positively correlated with the large/apex predators, even though its distribution

expanded slightly, suggest that they were adapted to coexisting with large/apex
predators.

The major relations between wolves/foxes/mustelids and large/apex predators
were negative firstly overall, but tend to turn slight positive overall. We infer that the
reasons are 1.there were many co-low-probability grids where the extremely high
HPD constrained all of the predators; 2.the mesopredators could be co-existent with
the large/apex predators, especially at the edges of both ranges (Newsome *et al*.
2017).

The expansion of wolves was mainly in low-mid HPD (<100 person/km^2) areas.
The body mass of wolves is transitive between apex predator and mesopredator.
Although wolves are attributed the role of a retreating apex predator in many studies
(Prugh *et al*. 2009; Newsome *et al*. 2017), our results are more consistent with
research conducted in Sikhote, Russia (Miquelle *et al*. 2005) where wolves are viewed
as mesopredators being constrained by tigers. This difference shows that the
classification between apex and mesopredator depends not only on species (Wallach
et al. 2015), but also on the local eco-system.

Although dholes are a kind of mesopredator with a middling body size and its
high-probability area expanded slightly, the distribution was strongly positively
correlated with the large/apex predators, and only slightly positive with the other
mesopredators. As a mesopredator dholes would be supposed to suffer from the
constraints of apex predators, but recent studies in South and South-east Asia revealed
that dholes can in general peacefully co-exist with tigers and leopards (Johnsingh
1992; Karanth & Sunquist 2000; Wang & Macdonald 2009; Jerks *et al*. 2012;
Hayward 2014; Rayan & Linkie 2016), and thus form a relationship as a "predator
guild". This study supports the view of such a predator guild, and supports it over a
longer temporal span than previous studies. The relationship between dholes and
bears is also strongly positively correlated, but we did not find any relevant predator
guild study. So we infer that the reason is that the environmental demands of bears are
similar to those of tigers and leopards, as the cause of such similar co-existing
relations. And civets are similar with dholes, but still not noticed by previous
researches. Mustelids were well adapted (most well one in mesopredator group) to the
environment with human disturbing, and they coexisted well with other mesopredator
and tiger.

The influence of HPD on large/apex predators is evidently negative, proving that
the retreat of large/apex predator is likely caused by human disturbance. For the
mesopredators, the increasingly negative influence of HPD, and the higher
distribution proportion in lower HPD area in the later period, also suggest that the
competition from large/apex predators constrained the distribution of mesopredators.
We propose that there was a trade-off phenomenon in which under a moderate HPD
the expulsion of larger predators may compensate the directly negative influence of
HPD on mesopredators.

5.Conclusion and Prospect:

Our study aimed to correct the existing methods for reconstruction of historical distribution based on local gazetteers by adopting the use of both presence and absence data. Based on our reconstructions, we provide direct proofs of human disturbance on mammal distribution in China, and extend support for the mesopredator release hypothesis and the predator guild hypothesis.

We have also collected records for other wild animals from local gazetteers in China. Based on the collection, we have built the Database of Wild Mammal Records in Chinese Local Gazetteers and are building the Database of Wild Bird Records in Chinese Local Gazetteers. We aim to continue relevant studies using these databases in the future.

Author Contributions:
Kaijin Hu designed research, collected data and wrote the paper; and Chunhui Hao modified the introduction.

ORCID:
Kaijin Hu https://orcid.org/0000-0001-9639-3774

Acknowledgements:
We thank Cuiyi He, Junming Liang, Jieyun He, Yao Zeng, Daishan Chen, Chenhui Shi and Lu Huang for assistance in collecting data, Zhu Deng for guidance in ArcGIS software, Pengfei Fan, Li Yang, Lu Zhang for useful advice, and Peter Tomlin for linguistic assistance.

Reference:

Areias Guerreiro, J., Mira, A., & Barbosa, A. M. (2016). How well can models predict changes in species distributions? A 13-year-old otter model revisited. *Hystrix*. https://doi.org/10.4404/hystrix-27.1-11867

Barbosa, A. M. (2015). fuzzySim: Applying fuzzy logic to binary similarity indices in ecology. *Methods in Ecology and Evolution*. https://doi.org/10.1111/2041-210X.12372

Barnosky, A. D. (2008). Megafauna biomass tradeoff as a driver of Quaternary and future extinctions. *Proceedings of the National Academy of Sciences of the United States of America*. https://doi.org/10.1073/pnas.0801918105

Barnosky, A. D., Matzke, N., Tomiya, S., Wogan, G. O. U., Swartz, B., Quental, T. B., Marshall, C., McGuire, J. L., Lindsey, E. L., Maguire, K. C., Mersey, B., & Ferrer, E. A. (2011). Has the Earth's sixth mass extinction already arrived? *Nature*, *471*(7336), 51–57. https://doi.org/10.1038/nature09678

Boakes, E. H., McGowan, P. J. K., Fuller, R. A., Chang-qing, D., Clark, N. E., O'Connor, K., & Mace,
G. M. (2010). Distorted Views of Biodiversity: Spatial and Temporal Bias in Species
Occurrence Data. *PLoS Biology*, *8*(6), e1000385.
https://doi.org/10.1371/journal.pbio.1000385

Boulanger, M. T., & Lyman, R. L. (2013). Northeastern North American Pleistocene megafauna
chronologically overlapped minimally with Paleoindians. *Quaternary Science Reviews*.
https://doi.org/10.1016/j.quascirev.2013.11.024

Cao, S. (2001). *History of Population in China, vol. V.* Fudan University Press.*(In Chinese)*

Cardillo, M., Mace, G. M., Gittleman, J. L., Jones, K. E., Bielby, J., & Purvis, A. (2008). The
predictability of extinction: biological and external correlates of decline in mammals.
Proceedings of the Royal Society B: Biological Sciences, *275*(1641), 1441–1448.
https://doi.org/10.1098/rspb.2008.0179

Cardillo, M., & Meijaard, E. (2012). Are comparative studies of extinction risk useful for
conservation? *Trends in Ecology & Evolution*, *27*(3), 167–171.
https://doi.org/10.1016/j.tree.2011.09.013

Ceballos, G., Ehrlich, P. R., Barnosky, A. D., García, A., Pringle, R. M., & Palmer, T. M. (2015).
Accelerated modern human-induced species losses: Entering the sixth mass extinction.
Science Advances, *1*(5), 9–13. https://doi.org/10.1126/sciadv.1400253

Chatterjee, H., Tse, J. S. Y., & Turvey, S. (2012). Using Ecological Niche Modelling to Predict
Spatial and Temporal Distribution Patterns in Chinese Gibbons: Lessons from the Present
and the Past. *Folia Primatologica*, *83*, 85–99. https://doi.org/10.1159/000342696

Davidson, A. D., Hamilton, M. J., Boyer, A. G., Brown, J. H., & Ceballos, G. (2009). Multiple
ecological pathways to extinction in mammals. *Proceedings of the National Academy of
Sciences*, *106*(26), 10702–10705. https://doi.org/10.1073/pnas.0901956106

Dirzo, R., & Raven, P. H. (2003). Global State of Biodiversity and Loss. *Annual Review of
Environment and Resources*, *28*(1), 137–167.
https://doi.org/10.1146/annurev.energy.28.050302.105532

Dirzo, R., Young, H. S., Galetti, M., Ceballos, G., Isaac, N. J. B., & Collen, B. (2014). Defaunation in
the Anthropocene. *Science*, *345*(6195), 401–406. https://doi.org/10.1126/science.1251817

Fritz, S. A., Bininda-Emonds, O. R. P., & Purvis, A. (2009). Geographical variation in predictors of
mammalian extinction risk: big is bad, but only in the tropics. *Ecology Letters*, *12*(6),
538–549. https://doi.org/10.1111/j.1461-0248.2009.01307.x

Fu, Y. Z. et al.(Edit) (2016) The Series of Rare Local Gazetteers in National Library(国家图书馆藏
地方志珍本丛刊). *Tianjin Ancient Books Press*, Tianjin*(in Chinese)*

Gill, J. A. (2007). Approaches to measuring the effects of human disturbance on birds. *Ibis*,
149(SUPPL. 1), 9–14. https://doi.org/10.1111/j.1474-919X.2007.00642.x

Goldewijk, K. K., Beusen, A., Doelman, J., & Stehfest, E. (2017). Anthropogenic land use estimates
for the Holocene - HYDE 3.2. *Earth System Science Data*.
https://doi.org/10.5194/essd-9-927-2017

Hayward, M. W., Lyngdoh, S., & Habib, B. (2014). Diet and prey preferences of dholes (Cuon
alpinus): dietary competition within Asia's apex predator guild. Journal of Zoology, 294(4),
255–266.

He, F., & Hubbell, S. P. (2011). Species-area relationships always overestimate extinction rates
from habitat loss. Nature, 473(7347), 368–371. https://doi.org/10.1038/nature09985

He, Y. (1997). The distributions of rare mammal (II) in China. Hunan Normal University Press.*(In*
Chinese)

He, Y. (1993). The distributions of rare mammal in China. Hunan Science & Technology Press.*(In*
Chinese)

Hijmans, R. J. (2012). Cross-validation of species distribution models: Removing spatial sorting
bias and calibration with a null model. *Ecology*. https://doi.org/10.1890/11-0826.1

Hu, H. (1935). Distribution of Population in China. Acta Geographica Sinica, 02, 33–74.*(In*
Chinese)

Jablonski, D. (2008). Extinction and the spatial dynamics of biodiversity. Proceedings of the
National Academy of Sciences, 105(Supplement 1), 11528–11535.
https://doi.org/10.1073/pnas.0801919105

Jablonski, D. (2001). Lessons from the past: Evolutionary impacts of mass extinctions.
Proceedings of the National Academy of Sciences, *98*(10), 5393–5398.
https://doi.org/10.1073/pnas.101092598

Jackson, J. B. C. (2008). Ecological extinction and evolution in the brave new ocean. *Proceedings*
of the National Academy of Sciences, *105*(Supplement 1), 11458–11465.
https://doi.org/10.1073/pnas.0802812105

Jenks, K. E., Kitamura, S., Lynam, A. J., Ngoprasert, D., Chutipong, W., Steinmetz, R., Sukmasuang,
R., Grassman, L. I., Cutter, P., Tantipisanuh, N., & others. (2012). Mapping the distribution
of dholes, Cuon alpinus (Canidae, Carnivora), in Thailand. *Mammalia*, *76*(2), 175–184.

Jiang, H., Lian, J., Wang, J., Liu, Z., & Feng, M. (1995). Influence of population growth stress on
distribution of macaque monkey in south China. *Chinese Journal of Applied Ecology*, *6*(2),
176–181.*(In Chinese)*

Johnsingh, A. J. T. (1992). Prey selection in three large sympatric carnivores in Bandipur.
Mammalia, *56*(4). https://doi.org/10.1515/mamm.1992.56.4.517

Karanth, K. U., & Sunquist, M. E. (2000). Behavioural correlates of predation by tiger (*Panthera*
tigris), leopard (*Panthera pardus*) and dhole (*Cuon alpinus*) in Nagarahole, India. *Journal of*
Zoology, *250*(2), 255–265.

Kitahara, M., & Fujii, K. (1994). Biodiversity and community structure of temperate butterfly
species within a gradient of human disturbance: An analysis based on the concept of
generalist vs. Specialist strategies. *Researches on Population Ecology*, *36*(2), 187–199.
https://doi.org/10.1007/BF02514935

Leakey, R. E., & Roger Lewin. (1996). *The sixth extinction: Patterns of life and the future of*
humankind. Anchor.

Li, B., He, G., Guo, S., Hou, R., Huang, K., Zhang, P., Zhang, H., Pan, R., & Chapman, C. A. (2020).
Macaques in China: Evolutionary dispersion and subsequent development. *American*
Journal of Primatology, *82*(7), e23142. https://doi.org/10.1002/ajp.23142

Li, B., Pan, R., & Oxnard, C. E. (2002). Extinction of snub-nosed monkeys in China during the past
400 years. *International Journal of Primatology*, *23*(6), 1227–1244.
https://doi.org/10.1023/A:1021122819845

Li, G. P. (1998). *Agriculture in Ancient China*. The Commercial Press.*(In Chinese)*

Li, X., Jiang, G., Tian, H., Xu, L., Yan, C., Wang, Z., Wei, F., & Zhang, Z. (2015). Human impact and
climate cooling caused range contraction of large mammals in China over the past two
millennia. *Ecography*, *38*(1), 74–82.

Lorenzen, E. D., Nogués-Bravo, D., Orlando, L., Weinstock, J., Binladen, J., Marske, K. A., Ugan, A.,
Borregaard, M. K., Gilbert, M. T. P., Nielsen, R., Ho, S. Y. W., Goebel, T., Graf, K. E., Byers, D.,
Stenderup, J. T., Rasmussen, M., Campos, P. F., Leonard, J. A., Koepfli, K. P., … Willerslev, E.
(2011). Species-specific responses of Late Quaternary megafauna to climate and humans.
Nature, *479*(7373), 359–364. https://doi.org/10.1038/nature10574

MacPhee, DE, R., & eds Hans-Dieter Sues. (1999). Extinctions in Near Time. In *Extinctions in Near*
Time. Springer Science & Business Media. https://doi.org/10.1007/978-1-4757-5202-1

Martin, P. S. (2005). *Twilight of the mammoths: Ice Age extinctions and the rewilding of America*.
Univ of California Press. https://doi.org/10.5860/choice.43-4679

Miquelle, D., Stephens, P., Smirnov, E. N., Goodrich, J., Zaumyslova, O. Y., & Myslenkov, A. I.
(2005). *Tigers and Wolves in the Russian Far East: Competitive Exclusion, Functional*
Redundancy and Conservation Implications (pp. 179–207).

Moleón, M., Sánchez-Zapata, J. A., Donázar, J. A., Revilla, E., Martín-López, B., Gutiérrez-Cánovas,
C., Getz, W. M., Morales-Reyes, Z., Campos-Arceiz, A., Crowder, L. B., Galetti, M.,
González-Suárez, M., He, F., Jordano, P., Lewison, R., Naidoo, R., Owen-Smith, N., Selva, N.,
Svenning, J. C., … Tockner, K. (2020). Rethinking megafauna. In *Proceedings of the Royal*
Society B: Biological Sciences. https://doi.org/10.1098/rspb.2019.2643

Newsome, T. M., Greenville, A. C., Ćirović, D., Dickman, C. R., Johnson, C. N., Krofel, M., Letnic,
M., Ripple, W. J., Ritchie, E. G., Stoyanov, S., & Wirsing, A. J. (2017). Top predators constrain
mesopredator distributions. *Nature Communications*, *8*(May), 1–7.
https://doi.org/10.1038/ncomms15469

Nüchel, J., Bøcher, P. K., Xiao, W., Zhu, A.-X., & Svenning, J.-C. (2018). Snub-nosed monkeys
(Rhinopithecus): potential distribution and its implication for conservation. *Biodiversity and*
Conservation, *27*(6), 1517–1538. https://doi.org/10.1007/s10531-018-1507-0

Öckinger, E., Schweiger, O., Crist, T. O., Debinski, D. M., Krauss, J., Kuussaari, M., Petersen, J. D.,
Pöyry, J., Settele, J., Summerville, K. S., & Bommarco, R. (2010). Life-history traits predict
species responses to habitat area and isolation: a cross-continental synthesis. *Ecology*
Letters, *13*(8), no--no. https://doi.org/10.1111/j.1461-0248.2010.01487.x

Palacio, R.D., et al., A data-driven geospatial workflow to improve mapping species distributions
and assessing extinction risk under the IUCN Red List. 2020.(Preprint in
BioRxiv)https://doi.org/10.1101/2020.04.27.064477

Pereira, H. M., Leadley, P. W., Proença, V., Alkemade, R., Scharlemann, J. P. W.,
Fernandez-Manjarrés, J. F., Araújo, M. B., Balvanera, P., Biggs, R., Cheung, W. W. L., Chini, L.,
Cooper, H. D., Gilman, E. L., Guénette, S., Hurtt, G. C., Huntington, H. P., Mace, G. M.,
Oberdorff, T., Revenga, C., … Walpole, M. (2010). Scenarios for global biodiversity in the
21st century. *Science*, *330*(6010), 1496–1501. https://doi.org/10.1126/science.1196624

Pimm, S. L., Russell, G. J., Gittleman, J. L., & Brooks, T. M. (1995). The Future of Biodiversity.
Science, *269*(5222), 347–350. https://doi.org/10.1126/science.269.5222.347

Pocock, M. J. O. (2011). Can traits predict species' vulnerability? A test with farmland passerines
in two continents. *Proceedings of the Royal Society B: Biological Sciences*, *278*(1711),
1532–1538. https://doi.org/10.1098/rspb.2010.1971

Prugh, L. R., Stoner, C. J., Epps, C. W., Bean, W. T., Ripple, W. J., Laliberte, A. S., & Brashares, J. S.
(2009). The Rise of the Mesopredator. *Bioscience*, *9*, 9.

Qian, T., Chi, Y., Xi, C., Li, Z., & Wang, J. (2020). Changes in the historical and current habitat
ranges of rare wild mammals in China: A case study of six taxa of small-to large-sized
mammals. *Sustainability (Switzerland)*, *12*(7), 1–19. https://doi.org/10.3390/su12072744

Rahbek, C., & Colwell, R. K. (2011). Species loss revisited. *Nature*, *473*(7347), 288–289. https://doi.org/10.1038/473288a

Rayan, D. M., & Linkie, M. (2016). Managing conservation flagship species in competition: Tiger, leopard and dhole in Malaysia. *Biological Conservation*, *204*, 360–366.

Roberts, R. G., Flannery, T. F., Ayliffe, L. K., Yoshida, H., Olley, J. M., Prideaux, G. J., Laslett, G. M., Baynes, A., Smith, M. A., Jones, R., & Smith, B. L. (2001). New ages for the last Australian megafauna: Continent-wide extinction about 46,000 years ago. Science. https://doi.org/10.1126/science.1060264

Roberts, E. A., Sheley, R. L., & Lawrence, R. L. (2004). Using sampling and inverse distance weighted modeling for mapping invasive plants. *Western North American Naturalist*.

Sala, O. E. (2000). Global Biodiversity Scenarios for the Year 2100. Science, 287(5459), 1770–1774. https://doi.org/10.1126/science.287.5459.1770

Sandom, C., Faurby, S., Sandel, B., & Svenning, J.-C. (2014). Global late Quaternary megafauna extinctions linked to humans, not climate change. *Proceedings of the Royal Society B: Biological Sciences*, *281*(1787), 20133254. https://doi.org/10.1098/rspb.2013.3254

Schipper, J., Chanson, J. S., Chiozza, F., Cox, N. A., Hoffmann, M., Katariya, V., Lamoreux, J., Rodrigues, A. S. L., Stuart, S. N., Temple, H. J., Baillie, J., Boitani, L., Lacher, T. E., Mittermeier, R. A., Smith, A. T., Absolon, D., Aguiar, J. M., Amori, G., Bakkour, N., … Young, B. E. (2008). The Status of the World's Land and Marine Mammals: Diversity, Threat, and Knowledge. *Science*, *322*(5899), 225–230. https://doi.org/10.1126/science.1165115

Somers, R. H. (1962). A New Asymmetric Measure of Association for Ordinal Variables. *American Sociological Review*, *27*(6), 799. https://doi.org/10.2307/2090408

Takahashi, M. K., Eastman, J. M., Griffin, D. A., Baumsteiger, J., Parris, M. J., & Storfer, A. (2014). A stable niche assumption-free test of ecological divergence. *Molecular Phylogenetics and Evolution*. https://doi.org/10.1016/j.ympev.2014.03.003

Teng, S. N., Xu, C., Teng, L., & Svenning, J. C. (2020). Long-term effects of cultural filtering on megafauna species distributions across China. *Proceedings of the National Academy of Sciences of the United States of America*, *117*(1), 486–493. https://doi.org/10.1073/pnas.1909896116

Thomas, C. D., Cameron, A., Green, R. E., Bakkenes, M., Beaumont, L. J., Collingham, Y. C., Erasmus, B. F. N., De Siqueira, M. F., Grainger, A., Hannah, L., Hughes, L., Huntley, B., Van Jaarsveld, A. S., Midgley, G. F., Miles, L., Ortega-Huerta, M. A., Peterson, A. T., Phillips, O. L., & Williams, S. E. (2004). Letter to nature: Extinction risk from climate change. *Nature*, *427*(145–148), 145–148.

Turvey, S. T., Crees, J. J., & Di Fonzo, M. M. I. (2015). Historical data as a baseline for conservation:
reconstructing long-term faunal extinction dynamics in Late Imperial–modern China.
Proceedings of the Royal Society B: Biological Sciences, *282*(1813), 20151299.
https://doi.org/10.1098/rspb.2015.1299

Turvey, S. T., Crees, J. J., Li, Z., Bielby, J., & Yuan, J. (2017). Long-term archives reveal shifting
extinction selectivity in China's postglacial mammal fauna. *Proceedings of the Royal Society*
B: Biological Sciences, *284*(1867), 20171979. https://doi.org/10.1098/rspb.2017.1979

Tylianakis, J. M., Tscharntke, T., & Lewis, O. T. (2007). Habitat modification alters the structure of
tropical host–parasitoid food webs. *Nature*, *445*(7124), 202–205.
https://doi.org/10.1038/nature05429

Villavicencio, N. A., Lindsey, E. L., Martin, F. M., Borrero, L. A., Moreno, P. I., Marshall, C. R., &
Barnosky, A. D. (2016). Combination of humans, climate, and vegetation change triggered
Late Quaternary megafauna extinction in the Última Esperanza region, southern Patagonia,
Chile. *Ecography*. https://doi.org/10.1111/ecog.01606

Wake, D. B., & Vredenburg, V. T. (2008). Are we in the midst of the sixth mass extinction? A view
from the world of amphibians. *Proceedings of the National Academy of Sciences*,
105(Supplement 1), 11466–11473. https://doi.org/10.1073/pnas.0801921105

Wallach, A. D., Izhaki, I., Toms, J. D., Ripple, W. J., & Shanas, U. (2015). What is an apex predator.
Oikos, *124*(11), 1453–1461.

Wan, X., Jiang, G., Yan, C., He, F., Wen, R., Gu, J., Li, X., Ma, J., Stenseth, N. C., & Zhang, Z. (2019).
Historical records reveal the distinctive associations of human disturbance and extreme
climate change with local extinction of mammals. *Proceedings of the National Academy of*
Sciences of the United States of America, *116*(38), 19001–19008.
https://doi.org/10.1073/pnas.1818019116

Wan, X., & Zhang, Z. (2017). Climate warming and humans played different roles in triggering
Late Quaternary extinctions in east and west Eurasia. *Proceedings of the Royal Society B:*
Biological Sciences, *284*(1851), 20162438. https://doi.org/10.1098/rspb.2016.2438

Wang, S. W., & Macdonald, D. W. (2009). Feeding habits and niche partitioning in a predator
guild composed of tigers, leopards and dholes in a temperate ecosystem in central Bhutan.
Journal of Zoology, *277*(4), 275–283.

Wen, R. S. (2013). *Geographical distribution of wild animals in ancient China*. Scientific and
Technological Press of Shandong.*(In Chinese)*

Wen, R. S. (2009). *The distribution and changes of rare wild animals in China*. Scientific and
Technological Press of Shandong.*(In Chinese)*

Wen, R. S. (2018). *The distribution and changes of rare wild animals in China(sequel)*. Scientific
and Technological Press of Shandong.*(In Chinese)*

Yap, T. A., Koo, M. S., Ambrose, R. F., Wake, D. B., & Vredenburg, V. T. (2015). Averting a North
American biodiversity crisis. *Science*, *349*(6247), 481–482.
https://doi.org/10.1126/science.aab1052

Zhao, X., Ren, B., Garber, P. A., Li, X., & Li, M. (2018). Impacts of human activity and climate
change on the distribution of snub-nosed monkeys in China during the past 2000 years.
Diversity and Distributions, *24*(1), 92–102. https://doi.org/10.1111/ddi.12657

Zhou, Y. H., & Zhang, P. (2013). Distribution and vicissitude of gibbons（*Hylobatidae*）in China
during the last 500 years. *Acta Theriologica Sinica*, *33*(3), 258–266.*(In Chinese)*

Philip, G. M., and D. F. Watson. "A Precise Method for Determining Contoured
Surfaces." Australian Petroleum Exploration Association Journal 22: 205–212. 1982.
612
613   Watson, D. F., and G. M. Philip. "A Refinement of Inverse Distance Weighted
614   Interpolation." Geoprocessing 2:315–327. 1985.

628 **Fig.1** Flow Chart

630
631
632

633 **Table.1** Qualifying the valid data.

Standards	Qualification
Only livestock	Invalid data
Only wild mammals with body length ≥1m	Invalid data
Wild mammals with body length ≥1m, and wild mammals with body length <1m	Valid data
Only wild mammals with body length <1m	Valid data

635 **Fig.2** Research area (blue) in eastern China and the Heihe-Tengchong Line (red).

654 **Fig.3** The change of average probability of predators from Period 1 to Period 4.

655
656

657 **Fig.4** The change from Period 1 to Period 4(Value=$(S_4-S_1)/S_1$)

673   **Fig.5** Relations between each predator and others (Kendall's tau-b) (N.S : not significant)

Fig.6 HPD Influence on predator distribution (Somers' dyx) (N.S. : not significant)

689   **Fig.7** Average probability in different HPD area in each period

693
694
695
696
Fig.8 Criticizing the presence-only method

a.The bear presence records(red points) in Period 2, both frames are almost empty; b. The bear presence

records(red points) and absence records(blue points) in Period 2, there are many absence records in the blue frame,

but few in the brown frame; c. The IDW reconstruction of bear distribution based on b., high-probability area(red)

and low-probability area(blue), the blue frame is fulfilled by low-probability area but the brown frame is fulfilled

by high-probability area.

704   a   b   c

715   **Fig.9** Criticizing the last observation method

722   **Fig.10** Different results of IDW, Last Observation(50km*50km) & Last Observation(50km*50km)

723
724
725
726
727   **Supplementary Information:**
728
729   **Table. S1** Folk name of predators in local gazetteers and relevant species in scientific taxonomy
730

Folk name in pre-modern China	Scientific name	English name
Tiger 虎	*Panthera tigris*	Tiger
Leopard 豹	*Panthera pardus*	Leopard
	Neofelis nebulosa	Clouded Leopard
	Uncia uncia	Snow Leopard
Bear[1.] 熊	*Helarctos malayanus*	Sun Bear
	Ursus arctos	Brown Bear
	Ursus thibetanus	Asian Black Bear
Wolf 狼	*Canis lupus*	Wolf
Fox 狐	*Vulpes corsac*	Corsac Fox
	Vulpes ferrilata	Tibetan Fox
	Vulpes vulpes	Red Fox
Civet (viverrids & small wild felids) 狸/香猫/野猫	*Vormela peregusna*	Marbled Polecat
	Catopuma temminckii	Asian Golden Cat
	Felis bieti	Chinese Mountain Cat
	Felis chaus	Jungle Cat
	Felis manul	Pallas' Cat

	Felis silvestris	Wildcat
	Pardofelis marmorata	Marbled Cat
	Prionailurus bengalensis	Leopard Cat
	Chrotogale owstoni	Owston's Palm Civet
	Arctictis binturong	Binturong
	Arctogalidia Trivirgata	Small-Toothed Palm Civet
	Paguma larvata	Masked Palm Civet
	Paradoxurus hermaphroditus	Common Palm Civet
	Prionodon pardicolor	Spotted Linsang
	Viverra megaspila	Large-Spotted Civet
	Viverra zibetha	Large Indian Civet
	Viverricula indica	Small Indian Civet
Dhole 豺/赤狗	*Cuon alpinus*	Dhole
Mustelid (Badger, Otter, etc.) 獭/鼬/獾/貂/龙狗/龙猪/山狗/猯	*Aonyx cinerea*	Asian Small-Clawed Otter
	Lutra lutra	Eurasian Otter
	Lutrogale perspicillata	Smooth-Coated Otter
	Arctonyx collaris	Hog Badger
	Martes flavigula	Yellow-Throated Marten
	Martes foina	Beech Marten
	Meles leucurus	Asian Badger
	Melogale moschata	Chinese Ferret Badger
	Melogale personata	Burmese Ferret Badger
	Mustela altaica	Mountain Weasel
	Mustela erminea	Ermine
	Mustela eversmanii	Steppe Polecat
	Mustela kathiah	Yellow-Bellied Weasel
	Mustela nivalis	Least Weasel
	Mustela sibirica	Siberian Weasel
	Mustela strigidorsa	Back-Striped Weasel

731   1. We didn't distinguish the brown bear, sun bear and Asian black bear in this research, because
732 the majority of local gazetteers only refer "熊(bear)", only a few of them distinguish "狗熊(Asian
733 black bear)", "马熊/罴(brown bear)" or "猪熊(suspected to be sun bear)". There are specific name
734 "貘"of panda, not included in this research.

742 **Table. S2** The distribution of predators and human population density(HPD)

746 **Table. S3** The change of average probability of predators from Period 1 to Period 4.

	Tiger	Leopard	Bear	Wolf	Fox	Civet	Dhole	Mustelid
Period 1	63.59%	59.25%	36.12%	41.50%	60.96%	47.06%	36.03%	59.69%
Period 2	61.15%	59.58%	38.30%	42.42%	65.42%	51.00%	35.16%	71.23%
Period 3	53.21%	54.18%	31.59%	46.51%	67.80%	61.92%	40.10%	76.47%
Period 4	49.13%	59.11%	24.00%	51.95%	69.64%	66.81%	40.93%	82.52%

749 **Table.S4** Relations between each predator and others (Kendall's tau-b coefficient)

Period 1	Tiger	Leopard	Bear	Wolf	Fox	Civet	Dhole	Mustelid
Tiger	1	.550**	.408**	−.076**	−.008 N.S.	.281**	.553**	.133**
Leopard	.550**	1	.511**	−.134**	−.120**	.130**	.475**	−.106**
Bear	.408**	.511**	1	−.121**	−.091**	.169**	.415**	−.089**
Wolf	−.076**	−.134**	−.121**	1	.293**	−.052**	−.026**	.081**
Fox	−.008 N.S.	−.120**	−.091**	.293**	1	−.042**	−.034**	.141**
Civet	.281**	.130**	.169**	−.052**	−.042**	1	.249**	.309**
Dhole	.553**	.475**	.415**	−.026**	−.034**	.249**	1	.116**
Mustelid	.133**	−.106**	−.089**	.081**	.141**	.309**	.116**	1

Period 2	Tiger	Leopard	Bear	Wolf	Fox	Civet	Dhole	Mustelid
Tiger	1	.610**	.395**	−.048**	−.082**	.190**	.476**	.067**
Leopard	.610**	1	.537**	−.137**	−.119**	.078**	.475**	−.086**
Bear	.395**	.537**	1	−.116**	−.146**	.002 N.S.	.379**	−.126**
Wolf	−.048**	−.137**	−.116**	1	.328**	−.024**	−.065**	.076**
Fox	−.082**	−.119**	−.146**	.328**	1	.014*	−.132**	.123**
Civet	.190**	.078**	.002 N.S.	−.024**	.014*	1	.177**	.330**
Dhole	.476**	.475**	.379**	−.065**	−.132**	.177**	1	.064**
Mustelid	.067**	−.086**	−.126**	.076**	.123**	.330**	.064**	1

Period 3	Tiger	Leopard	Bear	Wolf	Fox	Civet	Dhole	Mustelid
Tiger	1	.605**	.405**	−.050**	−.075**	.238**	.451**	.111**
Leopard	.605**	1	.478**	−.043**	.019**	.172**	.521**	.031**
Bear	.405**	.478**	1	−.085**	−.039**	.132**	.381**	−.031**
Wolf	−.050**	−.043**	−.085**	1	.298**	−.083**	−.045**	.097**
Fox	−.075**	.019**	−.039**	.298**	1	−.025**	−.013*	.155**
Civet	.238**	.172**	.132**	−.083**	−.025**	1	.276**	.210**
Dhole	.451**	.521**	.381**	−.045**	−.013*	.276**	1	.050**
Mustelid	.111**	.031**	−.031**	.097**	.155**	.210**	.050**	1

Period 4	Tiger	Leopard	Bear	Wolf	Fox	Civet	Dhole	Mustelid
Tiger	1	.599**	.480**	.015*	−.172**	.231**	.479**	.061**
Leopard	.599**	1	.560**	.112**	.001 N.S.	.214**	.522**	.071**
Bear	.480**	.560**	1	.028**	−.138**	.193**	.382**	.052**
Wolf	.015*	.112**	.028**	1	.368**	−.012 N.S.	.052**	.074**
Fox	−.172**	.001 N.S.	−.138**	.368**	1	−.047**	−.041**	.179**
Civet	.231**	.214**	.193**	−.012 N.S.	−.047**	1	.208**	.157**
Dhole	.479**	.522**	.382**	.052**	−.041**	.208**	1	.105**
Mustelid	.061**	.071**	.052**	.074**	.179**	.157**	.105**	1

751 **sig< .01;*sig< .05; N.S.: not significant.

753 **Table.S5** HPD Influence on predator distribution (Somers' dyx)

	Period 1	sig.	Period 2	sig.	Period 3	sig.	Period 4	sig.
Tiger	−.114	.000	−.149	.000	−.174	.000	−.241	.000
Leopard	−.297	.000	−.283	.000	−.243	.000	−.322	.000
Bear	−.292	.000	−.301	.000	−.302	.000	−.281	.000
Wolf	.030	.000	−.006	.368 N.S.	−.048	.000	−.200	.000
Fox	.042	.000	.034	.000	−.034	.000	−.098	.000
Civet	.027	.000	.076	.000	.003	.635 N.S.	−.045	.000
Dhole	−.193	.000	−.178	.000	−.202	.000	−.221	.000
Mustelid	.195	.000	.195	.000	.151	.000	.001	.918 N.S.

754 N.S. : not significant

30