Local Gazetteers Reveal Contrasting Patterns of Historical Distribution Changes between Apex Predators and Mesopredators in Eastern China

Kaijin Hu[1]

1. School of Sociology and Anthropology, Sun Yat-sen University, Guangzhou, China

Abstract:

Background: Humans have been causing the sixth wave of mass extinction of biodiversity. The situation of predators, especially of carnivores, is a key indicator of biodiversity, and the mesopredator release is a typical phenomenon in ecosystem recess. Local gazetteers are a rich resource for historical biodiversity research. However there are obvious biases in previous studies focusing only on presence records and neglecting the absence records. I recollected and analyzed the records by fixed methods to research historical biodiversity change.

Methods: Innovatively, this research used both presence and absence records from local gazetteers to reconstruct the distribution of 8 kinds of mammalian predators (i.e. large carnivores: tigers, leopards, and bears; medium-sized carnivores: wolves, foxes, civets, dholes, and mustelids) in eastern China from 1573C.E.(Common Era) to 1949C.E. (sorted into 4 periods). Then by utilizing statistics methods, e.g. Partial Least Squares Structural Equation Modeling(PLS-SEM), I analyzed the distribution changes(overall and sorted into 5 altitude groups), and the relation between carnivores and the influence from humans.

Results & Conclusions: I reconstructed the specific distribution changes of the carnivores and found the changes match the mesopredator release phenomenon. Furthermore, I found the specific relation(e.g. trade-off effect) between carnivores and the influence of humans. Besides, I found that bears and civets may be potential members of the "Tigers - Leopards - Dholes" predator guild.

Prospects: I have also collected records for other wild animals from local gazetteers in China. Based on this collection, I have built the Database of Wild Mammal Records in Chinese Local Gazetteers and am building the Database of Wild Bird Records in Chinese Local Gazetteers. I aim to continue relevant studies using these databases in the future.

Keywords: Local gazetteer; Historical distribution; Mesopredator release; Predator Guild; PLS-SEM

ORCID:
Kaijin Hu https://orcid.org/0000-0001-9639-3774

1.Introduction

It has been widely accepted that Earth is facing a biodiversity crisis (Rahbek & Colwell 2011). In the past 500 years, a wave of extinction and declines in local animal population have been recorded, with both the rate and magnitude of this recent period of extinction being comparable to the five previous periods of mass extinction (Pimm *et al.* 1995; He & Hubbell 2011; Dirzo *et al.* 2014). Unlike these five previous periods of mass extinction which were primarily driven by global changes in climate and atmospheric-ocean chemistry, volcanism, and bolide impacts (Barnosky *et al.* 2011), the modern loss of biodiversity is considered as the consequence of human socioeconomic activities (Dirzo & Raven 2003; Ceballos *et al.* 2015). Humans have been causing the sixth wave of extinction through changing global climate, overkilling species, overexploitation resources, introducing exotic species, spreading infectious diseases, and destructing and modifying habitats (Leakey & Roger Lewin 1996; MacPhee *et al.* 1999; Dirzo & Raven 2003; Thomas *et al.* 2004; Martin 2005; Jackson 2008; Wake & Vredenburg 2008; Ceballos *et al.* 2015; Yap *et al.* 2015).

Although there is no doubt that the impact of the recent "sixth period of mass extinction" has extended across distinctive taxonomic groups, each species responds differently to climatic shifts, habitat fragmentation, and decline in species abundance caused by anthropogenic disturbance (Cardillo *et al.* 2008; Lorenzen *et al.* 2011). Moreover, the life history characteristics of different species are also to some extent correlated with their risk of being threatened by human activities (Davidson *et al.* 2009; Öckinger *et al.* 2010; Cardillo & Meijaard 2012). Despite this, such studies usually had poor predictive power to reliably link species' vulnerability with their life history traits (Jablonski 2001, 2008; Pocock 2011). Even relatively well-established patterns of correlation, such as large body size being known to be related strongly to extinction risk in mammals, still failed to accurately predict the extinction risk of a specific species affected by human activities. This is due to (*i*) multiple ecological factors and human activities interact to affect species that differ by orders of magnitude in life history traits (Davidson *et al.* 2009), and (*ii*) human disturbance to specific species is usually regulated by the local community structures and population dynamics in which the species is embedded (Kitahara & Fujii 1994; Gill 2007; Tylianakis *et al.* 2007). A phenomenon termed "mesopredator release" represents one of the best-referred examples. Large-bodied predators were persecuted by humans, but the abundance of mesopredators might undergo dramatic increases as a result of trophic cascading effects in communities (Prugh *et al.* 2009; Wallach *et al.* 2015). Consequently, in view of the complex synergy of both intrinsic species traits and human's role at different spatial-temporal and ecological scales, more specific quantitative assessments of the relationships between local extinction of endangered species and anthropogenic factors are urgently required (Fritz *et al.* 2009; Dirzo *et al.* 2014; Sandom *et al.* 2014; Wan *et al.* 2019; Teng *et al.* 2020).

In China, the distribution of wild animals has changed greatly over time, with many formerly widely distributed wild animals now endangered or even extinct (Chatterjee et al. 2012; Qian *et al.* 2020). Meanwhile, the population in China has

increased significantly since the introduction of American crops in the 16th century,
which triggered prosperous socio-economic activities that may be responsible for the
extinction of wild animals in China (Li 1998; Cao 2001). China has a long history of
using local gazetteers to record momentous events and situations, including those
related to local wildlife, and previous researchers have published numerous collection
lists of animal records (He 1993; 1997; Wen 2009; 2013; 2018). By exploring these
historical records in various ways, researchers have attempted to reconstruct the
spatio-temporal distribution and extinction of engendered animals in China, and then
disentangle the relative roles of anthropogenic and climate changes in shaping their
distribution and extinction (Jiang *et al*. 1995; Li *et al*. 2002, 2020; Zhou & Zhang
2013; Li *et al*. 2015; Turvey *et al*. 2015, 2017; Wan & Zhang 2017; Nüchel *et al*. 2018;
Zhao *et al*. 2018; Wan *et al*. 2019; Qian *et al*. 2020; Teng *et al*. 2020). However, such
studies have relied on biased species observation data which only focused on the
"presence" records of local animals but neglected the "absence" records. For each
species, they often defined the time of extirpation as the last observation (or called
last occurrence) time which is interpreted from the most recent record there.

The previous studies neglected the absence records and thus led to biases. The
distribution of local gazetteers is not uniform in either space or time. If only
considering the presence (1) records, it will greatly confuse the absence (0) areas and
unknown(null) areas. The collection of data from gazetteer-rich and gazetteer-poor
regions/periods may cause obvious biases in estimating the distribution of wild
animals(Boakes *et al*. 2010). For example, there are few gazetteers in war-torn or
sparsely populated areas, and there were only one or two gazetteers from many
regions throughout history. Therefore, to accurately evaluate the spatio-temporal
distribution and extinction of wild animals in China, it is necessary to take into
account these biases in the data collected, and then do the re-collection job to track
changes in biodiversity and analyze how human activities would have affected such
changes.

In this research, I recollected the presence and absence data of 8 kinds of wild
mammals in eastern China from 1573C.E.(Common Era) to 1949C.E. I chose this
time period and region to avoid potential biases, because (*i*) eastern China was the
main activity area of ancient Chinese people; (*ii*) during this period, China
experienced a rapidly growing human population; (*iii*) a greater quantity of records of
wild animals in regional gazetteers. The data were re-organized from scanned local
gazetteers (county-level or prefectural-level) in the Chinese Digital Gazetteer
Database (http://x.wenjinguan.com/) and *The Series of Rare Local Gazetteers in the*
National Library（Fu *et al.* 2016）. All 8 kinds of the mammals I studied are predators
of distinctive body sizes, are widely distributed across gazetteer records, and are also
easily distinguishable in traditional folk taxonomy. Furthermore, by performing new
analytical techniques (see *Methods*), I reconstructed the spatial and temporal
distribution patterns of 8 kinds of predators with both presence and absence records.
The ultimate objective of this study was to quantitatively and specifically estimate
how human socioeconomic activities have affected the distribution and extirpation of

these 8 predators in eastern China.

2.Methods:

Overview:

I chose the eastern part of China as the research area, and the temporal span
1573C.E.-1949C.E. as the research period, for there are rich amount of local
gazetteers preserved, and the human population experienced a huge increase. I
collected both the presence and absence data of each target mammal predator from
local gazetteers and sorted them into 4 periods. In each period, I used the tool Inverse
Distance Weighted (IDW) to reconstruct the distribution probability layers by ArcGIS
10.3, I named this procedure as "Interpolation of Presence & Absence Records in
Same Periods". Then I used 20km*20km grids to extract the continuous value in those
layers and the human population density layers, to analyze the distribution changes,
the relation between animals, and the influence of humans. (see *Fig. S1*)

2.1 Data collection

I collected the data including both presence records and absence records from the
Chinese Digital Gazetteer Database(http://x.wenjinguan.com/) and *The Series of Rare
Local Gazetteers in the National Library*(Fu *et al.* 2016). Previous studies used only
presence points to reconstruct the presence and pseudo-absence(actually confused the
real absence(0) with data missing(null) maps, then extracted the binary value in each
grid to analyze with limited maths methods. Different from the above way, I used both
presence and absence points to reconstruct the probability maps. Then I extracted the
continuous value in each grid to analyze. Previous studies collect the presence records
of the target animal, but in this research, I collected the valid animal diversity records,
and acquired the presence and absence records of the target animal. For example,
there are two animal records in different local gazetteers(see row 1 in *Fig.S1*), A
recorded "Tiger, Leopard, Deer, Roe Deer .... Hedgehog, Vole, Ibex." (Wolf not
existed), and B recorded "Cattle, Horse ... Wolf, Dog, Badger, Rabbit, Mouse" (Wolf
existed). The previous studies would output "There were wolves in B" merely, but
my study would output "There were wolves in B, and there were no wolves in A"(see
row 2 in *Fig.S1*).

There are two kinds of invalid records I excluded. The first kind recorded only
livestock, and the second kind did not record comprehensively but recorded only
several large mammals roughly. To avoid the above invalid records, I filtered the
gazetteer records by the following rule: a valid record should record any wild
mammal of which the body weight is below 45kg(including those not analyzed in this
research, like macaca monkeys, gibbons, or wild rabbit) (see *Table.S1*).

I select 8 kinds of wide-distributed predators, i.e. tigers, leopards, bears, wolves,
foxes, civets, dholes, and mustelids, to research. And I match the folk animal kinds to
the modern species in China(Smith & Xie 2008) (see *Table.S2*), the folk taxonomy
animal kind can be matched to one or several modern scientific taxonomy species. I
sorted tigers, leopards, and bears into the large/apex predator group, and the rest into

the mesopredator group, according to the definition of megafauna, body mass ≥45kg or occupying the relevant ecological niche (Roberts *et al.* 2001; Barnosky 2008; Boulanger & Lyman 2013; Villavicencio *et al.* 2016; Moleon *et al.* 2020).

2.2 Spatial processing:

(Insert Fig.1)

I selected 22 province-level regions in the eastern continent of China as the research area (excluding the Tibetan Plateau part in Sichuan and Yunnan) which roughly corresponds with the eastern region of the Heihe-Tengchong Line (see *Fig.1*), containing approximately 90% of the human population of China from ancient to modern eras (Hu 1935). Although the names and areas of many administrative divisions have changed, it is able to locate the data on modern maps by using the current local government locations to identify the corresponding locations of the local gazetteers.

I used the Inverse Distance Weighted (IDW) tool to reconstruct the distribution based on the valid records using ArcGIS Desktop 10.3. This tool has been widely applied to species distribution reconstruction (Roberts *et al.* 2004; Hijmans 2012; Takahashi *et al.* 2014; Barbosa 2015; Areias-Guerreiro *et al.* 2016; Palacio *et al.* 2021). The interpolation value in each grid was generated by the nearest 12(default value) records, including presence records or absence records. The main assumption is that species are more likely to be found closer to presence points and farther from absences (i.e. spatial auto-correlation), and the local influence of points (weighted average) diminishes with distance (Palacio *et al.* 2021）．The result value means the probability of animal occurrence, in [0,1]. Besides the accuracy of data, the smoothy probability value is continuous, so I can analyze the distributions much subtler.

2.3 Temporal processing:

To keep the balance between the diachronical-spatial dimensions, I selected 1573-1949 as the research span, and sorted the valid points into 4 periods according to the reigning emperor, since many local gazetteers are labeled under the reigning emperor's title without specifying the year, and adopted the later one when there is more than one record in the same location in one period. Period 1 covers the span of 1573C.E.-1722C.E., with 694 valid data points; Period 2 covers the span of 1723C.E.-1820C.E., with 706 valid data points; Period 3 covers the span of 1821C.E.-1911C.E., with 786 valid data points; and Period 4 covers the span of 1912C.E.-1949C.E., with 451 valid data points. These layers are cross-sections and based on both presence/absence points, so it is appropriate that spans are different in periods.

This research has excluded the modern distribution of mammalian predators because their distribution was greatly reduced as a consequence of a campaign by the Chinese government to hunt the "harmful beasts" since the 1950s.

There are few local gazetteers that were written before 1573C.E. and are still
preserved today. That means the presence records (also absence records) are rare in
this period. Besides, the Chinese human population experienced several fluctuations
in history. It is inappropriate to infer the decline trend with the sample with these
biases. So I did not analyze this period in this research.

2.4 Statistics

2.4.1 Fishnet grids for overall surveys

I used the fishnet tool to build 10819 20km*20km grids (in WGS 1984 World
Mercator projected coordinate system), to extract the data from the layers.

The significance test is not necessary for this research, because the statistics are
not sampling surveys but overall surveys.

2.4.2 Relations of predator distribution：

I used Kendall's tau-b coefficient to measure the relations between each predator
in SPSS 22. This statistic is a measure of the strength and direction of associations
between two Ordinal variables. Kendall's tau-b treats the variables symmetrically and
ranges in [-1,1], with the $\pm$ symbol reflecting the direction of correlation and the
absolute value suggesting the strength of association.

$\text{tau-b} = (N_c - N_d)/[(N_c + N_d + T_x)(N_c + N_d + T_y)]^{0.5}$

N_c is the number of concordant pairs, N_d is the number of discordant pairs. T_x & T_y are the
number of tied pairs on each variable.

2.4.3 Influence of anthropogenic factor on predator distribution:

I added the human population density layers to make comparisons. I interpolated
human population data for the last three periods based on historical demographic
research. The population data in this study is referenced from Cao's(2001) "History of
the Chinese Population, Volume Five, Qing Dynasty" for historical population
statistics from the Qing Dynasty to 1953C.E. The 1820C.E. data corresponds to the
second period (1723C.E.-1820C.E.); the 1910C.E. data corresponds to the third period
(1821C.E.-1911C.E.), and the 1953C.E. data corresponds to the fourth period
(1912C.E.-1949C.E.) (Cao 2001). I imported the human population data into ArcGIS
Desktop 10.3 and created a kernel density map using a 200km search radius. For the
first period, where precise population statistics were unavailable, the study used the
population density layer from the HYDE 3.2 database for the year 1720C.E. as a
substitute (Goldewijk et al. 2017).

I measured the influence of human population density(HPD) on predator
distribution between each predator by Somers' d_{yx} statistic in the software SPSS 22.
This statistic is a measure of the influence strength of an ordinal variable on the other
ordinal variable. The d_{yx} ranges in [-1,1], with the $\pm$ symbol reflecting the direction

of influence and the absolute value reflecting the strength (Somers 1962).

$$d_{yx}=(N_c - N_d)/(N_c + N_d + T_y)$$

N_c is the number of concordant pairs, N_d is the number of discordant pairs. T_y is the number of

tied pairs on the dependent variable.

2.4.4 Terrain factors

This research incorporated the elevation layer and slope layer (both are from
Loess Plateau SubCenter, National Earth System Science Data Center, National
Science & Technology Infrastructure of China (http://loess.geodata.cn)). The entire
grids were approximately divided into five groups based on elevation (in meters):
alt1([-5, 58], n=2169 grids), alt2((58, 249], n=2157 grids), alt3((249, 670], n=2162
grids), alt4((670, 1241], n=2166 grids), and alt5((1241, 4414], n=2165 grids).
Subsequently, I used charts to display the variations in different animal species and
human population within each elevation range.

2.4.5The structural equation models analyzing the complex causal relationships
among large/apex carnivores, medium-sized carnivores, and human population
density.

I used PLS-SEM to analyze the complex causal relationships among the human
population, large carnivores, and medium-sized carnivores. In the first step, I
measured the latent variable of the overall large carnivores(reflecting the
completeness of the food chain) using the three kinds of large carnivores as reflective
indicators. In the second step, I conducted statistical analyses with human population
density as the independent variable, the overall large carnivores as the mediating
variable, and each kind of medium-sized carnivores as the dependent variable.
Because the large carnivores have predatory and competitive relationships with the
medium-sized carnivores in the food chain, and humans hunted both large and
medium-sized carnivores, I analyzed whether the large carnivores mediated the
impact of population density on medium-sized carnivores and the resulted masking
effect.

I chose PLS-SEM instead of the traditional CB-SEM, for the latter requires a
normal distribution of data. Compared to the traditional covariance-based structural
equation modeling (CB-SEM), partial least squares structural equation modeling
(PLS-SEM) does not require the data to follow a normal distribution as it employs a
non-parametric estimation method. In this study, the IDW data did not exhibit a
normal distribution, and there was a large amount of grids with values of 0 or 1
(representing extreme cases of 0% or 100% probability distribution). These grids
made it unsuitable to transform the data into a normal distribution through
standardization (which requires a larger sample size in the middle range and fewer
extreme values at both ends).

(Insert *Fig.2*)

3.Result：

3.1 The change of distribution area

I reconstructed the distribution of **target predators and human population density** (HPD) (see *Table.1*). In the large/apex predator group, the distribution of the two largest predators, tigers and bears retreated, while the smaller one, leopard, remained almost unchanged; by contrast, the distribution of predators in the mesopredator group expanded in varying degrees (see *Fig.3 & Table.S3*).

(Insert Fig.3)

Based on the analyses of elevation and slope, the average distribution of various carnivorous mammals overall shows an increasing trend in both elevation and slope (see *Fig.4*). Among them, the four periods of the large carnivores and dholes are all higher than the overall average elevation and slope of the entire research area, and they exhibit an increasing trend. In contrast, the average elevation and slope of the other medium-sized carnivores were mostly lower than the overall average values initially but increased to above the average later.

The grouped data on altitude suggests that: 1) generally, in lower altitude regions, the distribution of large carnivores decreased obviously, while the distributions of medium-sized carnivores increased obviously(civets), kept almost stable(foxes, dholes, and mustelids), or decreased obviously(wolves); 2) generally, in higher altitude regions, the distributions of large carnivores kept almost stable, while the distributions of medium-sized carnivores increased obviously(see *Fig.S2, Fig.5, Table.S4)*.

(Insert Fig.4, Fig.S2, &Fig.5)

3.2 Relations between each predator and others

(Insert Fig.S3)

According to the relations of distributions between predators (see *Fig.S3 & Table. S5*), the predators in the large/apex predator group displayed strongly positive correlations with each other in all periods**, suggesting that these species have similar environmental requirements**. In the mesopredator group, wolves and foxes were highly correlated, and the correlations between wolves/foxes and large/apex predators were negative overall; while for dholes and civets, they were positively correlated with the large/apex predators and each other, but weakly negative with wolves/foxes overall, suggest that they were adapted to coexisting with large/apex predators; for mustelids, the values with leopards and bears were negative in earlier periods but turn positive later, and the value with other predators(including tigers) were positive.

3.3 Anthropogenic influence on predator distributions

(Insert Fig.6)

The results (see *Fig.6 & Table. S6*) show that HPD exerted a negative influence upon the distributions of each large predator during each period, the influences on tigers, leopards, and bears kept are clearly negative throughout. But for mesopredators: (1)wolves, foxes, and civets were slightly positive in early periods, but became negative in later periods; (2)dholes were negative in all periods, like large predators; and (3) mustelids were strongly positive in period 1 to period 3, but turn slightly negative in period 4.

3.4 Complex causual relationships in PLS-SEM

（insert *Table.2*）：

According to the analysis(see *Table.2*), in the triangular relationship of "anthropogenic influence(human population density) - the overall Apex predators(large carnivores) - each of Meso-predators(medium-sized carnivores)" across four periods, human population density has consistently exerted a strong negative impact on the overall large carnivores (a).

For wolves and foxes, the direct impact of human population density (c) was consistently negative, and it became stronger (i.e. the absolute values increased) as the HPD increased in different periods. The impact of large carnivores was also consistently negative (b). In summary, the overall impact of HPD (s) initially was positive or nearly neutral, but later turned negative and became increasingly stronger.

For civets and dholes, the direct impact of HPD (c) was consistently weakly positive or nearly neutral, and the impact of large carnivores was strongly positive(b). In summary, the overall impact of HPD (s) fluctuates between weakly positive and weakly negative effects for civets, while it has consistently been negative for dholes.

For mustelids, the direct impact from HPD (c) was positive in the first three periods but became weakly negative in the fourth period. The impact of large carnivores (b) was weakly positive in the first period, weakly negative in the second period, and strongly positive in the third and fourth periods. The overall impact of HPD (s) was initially strongly positive but turned weakly negative in the fourth stage.

4.Discussion：

4.1 Discussion on procedure

In my procedure "Interpolation of Presence & Absence Records in Same Periods", the reconstruction of distributions using both presence and absence data is more accurate than previous studies (Zhou & Zhang 2013; Li *et al.* 2015; Turvey *et al.* 2015; Wan *et al.* 2019; Teng *et al.* 2020) with have only used presence data (see *Fig.7*), especially with the last observation procedure (see *Table.S7 & Table.3*). It

overcomes several disadvantages of previous studies:

(1)Sampling bias: The distribution of local gazetteers is not uniform in either space or time. If only considering the presence (1) records, it will greatly confuse the absence (0) areas and unknown(null) areas. There are few gazetteers in the area/period with social turbulence or few human population, causing the presence records to be few there/then (*Fig.7.a,*). I fixed this bias and interpolated the unknown areas by the IDW.

(2) New data: if there are new data, the reconstruction might change obviously (e.g. there are many more newly found records in Wen(2018)'s new research than his previous research (Wen 2009) then the reconstruction changed obviously), for the last observation precedure, it would lead to inverse interpolation. While in my procedure, new data will subtlize the reconstruction.

(3) Subjective: how large an area (circle or square, 50km*50km or 100km*100km, etc.) a data point represents is determined by the subjectivity of different researchers (Zhou & Zhang 2013; Turvey *et al.* 2015; Wan *et al.* 2019; Teng *et al.* 2020), leading to different result of distribution area, but in this research, it is replaced by objective interpolation of IDW.

(4) Monotonic recession prejudged: Last observation precedure can not reflect the expansion or recovery of animal distribution, but only monotonic recession, it is a kind of prejudge. In contrast, the expansion or recovery can also be noted in my procedure.

Above all, there are several biases in previous studies, and the procedure in this study mended them.

(Insert Fig.7, Table.S7 & Table.3)

I exemplify the contrast of the two procedures by the distribution of wolves(see *Fig.S2 & Table.3*). The results in the last observation precedure show a retracting trend, but the result values are much different between 100km*100km square grids and 50km*50km square grids. In contrast, the result in the IDW shows an expanding trend.

4.2 Discussion on results

4.2.1 Distribution of carnivores

The distribution of wild animals is significantly shaped by anthropogenic disturbances throughout the long history of China (Chatterjee *et al.* 2012).

The distribution of large carnivores decreased or kept almost stable. I found that the distribution of the two largest carnivores, i.e. tigers and bears, retreated over successive periods, supporting similar views of previous studies (He 1993; 1997; Wen 2009; 2013; 2018; Wan *et al.* 2019; Teng *et al.* 2020), and reflecting the recession of the eco-system over the same time frame. Leopards are the smallest one in large carnivores and also contain the larger medium-sized carnivores: cloud leopards. The

distribution of leopards kept almost stable, within the area in the lowest altitude group(alt1) decreased slightly but the area in the highest altitude group(alt5) increased slightly.

The distribution altitudes/slopes of large carnivores were consistently higher than the average in the research area and were increasing, suggesting a retraction towards higher altitudes (mountainous areas). Fisher and Shanee et al. have pointed out that for many animals currently concentrated in mountainous regions, the lower altitude areas are their most suitable habitats, however, human activities in these lower altitude areas compressed these animals' living spaces to the higher altitude, suboptimal habitats on the edges of their original suitable habitats. This study supports this viewpoint (Fisher 2011, Shanee et al. 2014).

The decline trend of bears was severe than that of tigers. but the bears are still living in several protected areas in the research area, while tigers are already extirpated. I infer that the reason is the different home range sizes. Tigers, with large home ranges, would appear in wide sink-habitat out of source-habitat and be exposed to anthropogenic dangers; while bears' habitats were majorly source-habitat, and they would survive when the small range habitats are preserved.

Not to be noted by previous studies, I find the distributions of mesopredator expanded in eastern China, matching the mesopredator release phenomenon, which suggests that the population of mesopredator would grow rapidly after the constraints from large/apex predators were withdrawn(Prugh *et al.* 2009). The distribution altitude/slope of medium-sized carnivores shifted from being lower than the average to approaching or higher than the average, suggesting an expansion of their distribution from lower altitude/slope areas to higher altitude/slope areas.

The results of mesopredators are not biased by data collection, because the distribution of other medium-sized mammals(2 kinds of primates and 3 kinds of medium-sized deer) decreased (Hu 2022).

I also provide a potential expression of the misunderstanding that wolves are not distributed in southern China. In numerous previous studies, wolves were once believed to be distributed only in the northern regions and the Qinghai-Tibet Plateau of China. However, Wang et al. clarified this misunderstanding by collating domestic specimen records and ecological research, confirming that wolves were also once widely distributed in various southern provinces of China (Wang et al. 2016). The evidence from this study supports Wang et al.'s viewpoint. It also indicates that the distribution of wolves in southern China undergoes a process of expansion, I speculate that it caused difficulties for early Western scholars to collect wolf specimens in southern China, thereby causing the misunderstanding.

4.2.2 The relationship among carnivores

4.2.2.1 Mesopredator release

The results match the changes of habitat overlaps in the process of mesopredator
release. Newsome et al (2017) proposed the existence of three zones/stages during the
process of mesopredator release: Zone A, the area monopolized by the large predators,
where large predators are present but mesopredators are absent; Zone B, the area
shared by large predators and mesopredators, where both present; Zone C, the area
monopolized by mesopredators, where large predators have disappeared but
mesopredators are still present. As HPD increased, the grouped altitude charts show
the transition from Zone A to Zone B and from Zone B to Zone C. This study also
identified a fourth zone/stage, Zone D, where mesopredators disappear as well when
human population density was very high, where neither large nor mesopredators are
present.

Exemplified by tigers and wolves, in the highest altitude group(alt5), the
distribution of tigers just decreased slightly while the distribution of wolves increased
obviously, corresponding to the transition from Zone A to Zone B. In the middle three
groups, the distribution of tigers decreased and the distribution of wolves increased,
corresponding to the transition from Zone B to Zone C. In the lowest altitude
group(alt1), as the population becomes highly dense, both tigers and wolves
decreased, corresponding to the transition from Zone C to Zone D.

It can explain why the relationships between large carnivores and wolves/
foxes/mustelids turned from negative to positive: the new co-exist area (Zone B
expanded and replaced Zone A) and the new co-non-exist area (Zone D expanded and
replaced Zone C) were larger than the new area where only large carnivores decreased
(Zone C expanded and replaced Zone B).

I provide a new case about the switchable role of wolves. The body mass of
wolves is transitive between apex predator and mesopredator. Although wolves are
attributed to the role of a retreating apex predator in many studies (Prugh *et al*. 2009;
Newsome *et al*. 2017), my results are more consistent with research conducted in
Sikhote, Russia (Miquelle *et al*. 2005) where wolves are viewed as mesopredators
being constrained by tigers. This difference shows that the classification between apex
and mesopredator depends not only on species (Wallach *et al*. 2015), but also on the
local eco-system.

4.2.2.2 Predator guild

I confirm the predator guild among tigers, leopards, and dholes existed in long
temporal span, and find out that bears (as a kind of large carnivore) and civets (as a
kind of medium-sized carnivore) are probably also in the guild. Although dholes are a
kind of mesopredator with a middling body size and its high-probability area
expanded slightly, the distribution was strongly positively correlated with the
large/apex predators, and only slightly positive with the other mesopredators. As a
kind of mesopredator, dholes would be supposed to suffer from the constraints of apex
predators, but recent studies in South Asia and Southeast Asia revealed that dholes can
in general peacefully co-exist with tigers and leopards (Johnsingh 1992; Karanth &
Sunquist 2000; Wang & Macdonald 2009; Jenks *et al*. 2012; Hayward 2014; Rayan &

Linkie 2016), and thus form a relationship as a "predator guild". This study supports
the view of such a predator guild and supports it over a longer temporal span than
previous studies. The relationship between dholes and bears is also strongly positively
correlated, but I did not find any relevant predator guild study. So I infer that the
reason is that the environmental demands of bears are similar to those of tigers and
leopards, as the cause of such similar co-existing relations. Besides, still not noticed
by previous research, civets were similar to dholes, but the positive relations with
large carnivores were weaker than those of dholes.

4.2.3 Influence of HPD

I quantified the specific negative impact of HPD on large carnivores,
demonstrating the decline in the distribution of large carnivores under human
interference. For medium-sized carnivores, the negative impact of HPD increased.
The changes indicate a trade-off effect: in regions with moderate HPD, the positive
impact of expelling large carnivores compensated for the direct negative impact of
humans' killing medium-sized carnivores.

The PLS-SEM analyzed the relation between human population expansion and
the mesopredator release. Wolves and foxes have been subject to sustained negative
impacts from large predators, and the disappearance of large predators has reduced the
survival pressure on wolves and foxes. Their distribution expansions were a typical
instance of the mesopredator release effect. The masking effect also confirms the
trade-off effect mentioned above. The trade-off effects of dholes, civets, and mustelids
were not typical, as they are influenced positively or fluctuated from positively to
negatively by large carnivores and HPD. Besides the predator guild effect in dholes
and civets, I speculate that these three kinds of animals also benefit directly from
human impact, such as providing more food sources like poultry and rodents.

In contemporary ecological research, red foxes (foxes in this research), wildcats
(civets in this research), and stone martens (mustelids in this research) are common
urban and suburban animals in Europe, and they are adapted to the human
environment (Recio et al. 2015). It is similar to the results of this study on civets and
mustelids. Dholes are not found in Europe, and the different situation of foxes in this
study may be due to the prohibition of fur hunting in the modern.

5.Conclusion and Prospect:

My study aimed to correct the existing procedures for reconstructions of
historical distribution based on local gazetteers by adopting the use of both presence
and absence data. Based on my reconstructions, I provide the direct proofs of human
disturbance on mammal distribution in China, and extend support for the
mesopredator release hypothesis and the predator guild hypothesis.

I have also collected records for other wild animals from local gazetteers in
China. Based on the collection, I have built the Database of Wild Mammal Records in

Chinese Local Gazetteers and are building the Database of Wild Bird Records in Chinese Local Gazetteers. I aim to continue relevant studies using these databases in the future.

ORCID:

Kaijin Hu https://orcid.org/0000-0001-9639-3774

Acknowledgements:

Ni, Ming provided advice to improve the figures and the tables. Hao, Chunhui provided advice to improve the introduction. Liu, Hanlun introduced the concept "mediating effect" to the author Hu, Kaijin.

Reference:

Areias Guerreiro, J., Mira, A., & Barbosa, A. M. (2016). How well can models predict changes in
species distributions? A 13-year-old otter model revisited. *Hystrix*.
https://doi.org/10.4404/hystrix-27.1-11867

Barbosa, A. M. (2015). fuzzySim: Applying fuzzy logic to binary similarity indices in ecology.
Methods in Ecology and Evolution. https://doi.org/10.1111/2041-210X.12372

Barnosky, A. D. (2008). Megafauna biomass tradeoff as a driver of Quaternary and future
extinctions. *Proceedings of the National Academy of Sciences of the United States of*
America. https://doi.org/10.1073/pnas.0801918105

Barnosky, A. D., Matzke, N., Tomiya, S., Wogan, G. O. U., Swartz, B., Quental, T. B., Marshall, C.,
McGuire, J. L., Lindsey, E. L., Maguire, K. C., Mersey, B., & Ferrer, E. A. (2011). Has the
Earth's sixth mass extinction already arrived? *Nature*, *471*(7336), 51–57.
https://doi.org/10.1038/nature09678

Boakes, E. H., McGowan, P. J. K., Fuller, R. A., Chang-qing, D., Clark, N. E., O'Connor, K., & Mace,
G. M. (2010). Distorted Views of Biodiversity: Spatial and Temporal Bias in Species
Occurrence Data. *PLoS Biology*, *8*(6), e1000385.
https://doi.org/10.1371/journal.pbio.1000385

Boulanger, M. T., & Lyman, R. L. (2013). Northeastern North American Pleistocene megafauna
chronologically overlapped minimally with Paleoindians. *Quaternary Science Reviews*.
https://doi.org/10.1016/j.quascirev.2013.11.024

Cao, S. (2001). *History of Population in China, vol. V.* Fudan University Press.*(In Chinese)*

Cardillo, M., Mace, G. M., Gittleman, J. L., Jones, K. E., Bielby, J., & Purvis, A. (2008). The
predictability of extinction: biological and external correlates of decline in mammals.
Proceedings of the Royal Society B: Biological Sciences, *275*(1641), 1441–1448.
https://doi.org/10.1098/rspb.2008.0179

Cardillo, M., & Meijaard, E. (2012). Are comparative studies of extinction risk useful for
conservation? *Trends in Ecology & Evolution*, *27*(3), 167–171.
https://doi.org/10.1016/j.tree.2011.09.013

Ceballos, G., Ehrlich, P. R., Barnosky, A. D., García, A., Pringle, R. M., & Palmer, T. M. (2015).
Accelerated modern human-induced species losses: Entering the sixth mass extinction.
Science Advances, *1*(5), 9–13. https://doi.org/10.1126/sciadv.1400253

Chatterjee, H., Tse, J. S. Y., & Turvey, S. (2012). Using Ecological Niche Modelling to Predict
Spatial and Temporal Distribution Patterns in Chinese Gibbons: Lessons from the Present
and the Past. *Folia Primatologica*, *83*, 85–99. https://doi.org/10.1159/000342696

Davidson, A. D., Hamilton, M. J., Boyer, A. G., Brown, J. H., & Ceballos, G. (2009). Multiple
ecological pathways to extinction in mammals. *Proceedings of the National Academy of*
Sciences, *106*(26), 10702–10705. https://doi.org/10.1073/pnas.0901956106

Dirzo, R., & Raven, P. H. (2003). Global State of Biodiversity and Loss. *Annual Review of*
Environment and Resources, *28*(1), 137–167.
https://doi.org/10.1146/annurev.energy.28.050302.105532

Dirzo, R., Young, H. S., Galetti, M., Ceballos, G., Isaac, N. J. B., & Collen, B. (2014). Defaunation in
the Anthropocene. *Science*, *345*(6195), 401–406. https://doi.org/10.1126/science.1251817

Fisher, D . (2011). Trajectories From Extinction: Where are Missing Mammals
Rediscovered? .Global Ecology & Biogeography, 20 (3): 415~425.
http://doi.org/10.1111/j.1466-8238.2010.00624.x

Fritz, S. A., Bininda-Emonds, O. R. P., & Purvis, A. (2009). Geographical variation in predictors of
mammalian extinction risk: big is bad, but only in the tropics. *Ecology Letters*, *12*(6),
538–549. https://doi.org/10.1111/j.1461-0248.2009.01307.x

Fu, Y. Z. et al.(Edit) (2016) The Series of Rare Local Gazetteers in the National Library. *Tianjin*
Ancient Books Press, Tianjin*(in Chinese)*

Gill, J. A. (2007). Approaches to measuring the effects of human disturbance on birds. *Ibis*,
149(SUPPL. 1), 9–14. https://doi.org/10.1111/j.1474-919X.2007.00642.x

Goldewijk, K. K., Beusen, A., Doelman, J., & Stehfest, E. (2017). Anthropogenic land use estimates
for the Holocene - HYDE 3.2. *Earth System Science Data*.
https://doi.org/10.5194/essd-9-927-2017

Hayward, M. W., Lyngdoh, S., & Habib, B. (2014). Diet and prey preferences of dholes (Cuon
alpinus): dietary competition within Asia's apex predator guild. Journal of Zoology, 294(4),
255–266.

He, F., & Hubbell, S. P. (2011). Species-area relationships always overestimate extinction rates
from habitat loss. Nature, 473(7347), 368–371. https://doi.org/10.1038/nature09985

He, Y. (1993). The distributions of rare mammal in China. Hunan Science & Technology Press.*(In*
Chinese)

He, Y. (1997). The distributions of rare mammal (II) in China. Hunan Normal University Press.*(In*
Chinese)

Hijmans, R. J. (2012). Cross-validation of species distribution models: Removing spatial sorting
bias and calibration with a null model. *Ecology*. https://doi.org/10.1890/11-0826.1

Hu, H. (1935). Distribution of Population in China. Acta Geographica Sinica, 02, 33–74.*(In*
Chinese)

Hu, K.(2022). The Distribution Changes of Primates and Other Common Mammals in Eastern
China During 1573~1949 — Quantitative Analysis Based on Local Gazetteers and GIS
Methods. Doctoral Thesis in Sun Yat-sen University.
http://doi.org/10.13140/RG.2.2.16959.28321

Jablonski, D. (2008). Extinction and the spatial dynamics of biodiversity. Proceedings of the
National Academy of Sciences, 105(Supplement 1), 11528–11535.
https://doi.org/10.1073/pnas.0801919105

Jablonski, D. (2001). Lessons from the past: Evolutionary impacts of mass extinctions.
Proceedings of the National Academy of Sciences, *98*(10), 5393–5398.
https://doi.org/10.1073/pnas.101092598

Jackson, J. B. C. (2008). Ecological extinction and evolution in the brave new ocean. *Proceedings*
of the National Academy of Sciences, *105*(Supplement 1), 11458–11465.
https://doi.org/10.1073/pnas.0802812105

Jenks, K. E., Kitamura, S., Lynam, A. J., Ngoprasert, D., Chutipong, W., Steinmetz, R., Sukmasuang,
R., Grassman, L. I., Cutter, P., Tantipisanuh, N., & others. (2012). Mapping the distribution
of dholes, Cuon alpinus (Canidae, Carnivora), in Thailand. *Mammalia*, *76*(2), 175–184.

Jiang, H., Lian, J., Wang, J., Liu, Z., & Feng, M. (1995). Influence of population growth stress on
distribution of macaque monkey in south China. *Chinese Journal of Applied Ecology*, *6*(2),
176–181.*(In Chinese)*

Johnsingh, A. J. T. (1992). Prey selection in three large sympatric carnivores in Bandipur.
Mammalia, *56*(4). https://doi.org/10.1515/mamm.1992.56.4.517

Karanth, K. U., & Sunquist, M. E. (2000). Behavioural correlates of predation by tiger (*Panthera*
tigris), leopard (*Panthera pardus*) and dhole (*Cuon alpinus*) in Nagarahole, India. *Journal of*
Zoology, *250*(2), 255–265.

Kitahara, M., & Fujii, K. (1994). Biodiversity and community structure of temperate butterfly
species within a gradient of human disturbance: An analysis based on the concept of
generalist vs. Specialist strategies. *Researches on Population Ecology*, *36*(2), 187–199.
https://doi.org/10.1007/BF02514935

Leakey, R. E., & Roger Lewin. (1996). *The sixth extinction: Patterns of life and the future of*
humankind. Anchor.

Li, B., He, G., Guo, S., Hou, R., Huang, K., Zhang, P., Zhang, H., Pan, R., & Chapman, C. A. (2020).
Macaques in China: Evolutionary dispersion and subsequent development. *American*
Journal of Primatology, *82*(7), e23142. https://doi.org/10.1002/ajp.23142

Li, B., Pan, R., & Oxnard, C. E. (2002). Extinction of snub-nosed monkeys in China during the past
400 years. *International Journal of Primatology*, *23*(6), 1227–1244.
https://doi.org/10.1023/A:1021122819845

Li, G. P. (1998). *Agriculture in Ancient China*. The Commercial Press.*(In Chinese)*

Li, X., Jiang, G., Tian, H., Xu, L., Yan, C., Wang, Z., Wei, F., & Zhang, Z. (2015). Human impact and
climate cooling caused range contraction of large mammals in China over the past two
millennia. *Ecography*, *38*(1), 74–82.

Lorenzen, E. D., Nogués-Bravo, D., Orlando, L., Weinstock, J., Binladen, J., Marske, K. A., Ugan, A.,
Borregaard, M. K., Gilbert, M. T. P., Nielsen, R., Ho, S. Y. W., Goebel, T., Graf, K. E., Byers, D.,
Stenderup, J. T., Rasmussen, M., Campos, P. F., Leonard, J. A., Koepfli, K. P., … Willerslev, E.
(2011). Species-specific responses of Late Quaternary megafauna to climate and humans.
Nature, *479*(7373), 359–364. https://doi.org/10.1038/nature10574

MacPhee, DE, R., & eds Hans-Dieter Sues. (1999). Extinctions in Near Time. In *Extinctions in Near*
Time. Springer Science & Business Media. https://doi.org/10.1007/978-1-4757-5202-1

Martin, P. S. (2005). *Twilight of the mammoths: Ice Age extinctions and the rewilding of America*.
Univ of California Press. https://doi.org/10.5860/choice.43-4679

Miquelle, D., Stephens, P., Smirnov, E. N., Goodrich, J., Zaumyslova, O. Y., & Myslenkov, A. I.
(2005). *Tigers and Wolves in the Russian Far East: Competitive Exclusion, Functional*
Redundancy and Conservation Implications (pp. 179–207).

Moleón, M., Sánchez-Zapata, J. A., Donázar, J. A., Revilla, E., Martín-López, B., Gutiérrez-Cánovas,
C., Getz, W. M., Morales-Reyes, Z., Campos-Arceiz, A., Crowder, L. B., Galetti, M.,
González-Suárez, M., He, F., Jordano, P., Lewison, R., Naidoo, R., Owen-Smith, N., Selva, N.,
Svenning, J. C., … Tockner, K. (2020). Rethinking megafauna. In *Proceedings of the Royal*
Society B: Biological Sciences. https://doi.org/10.1098/rspb.2019.2643

Newsome, T. M., Greenville, A. C., Ćirović, D., Dickman, C. R., Johnson, C. N., Krofel, M., Letnic,
M., Ripple, W. J., Ritchie, E. G., Stoyanov, S., & Wirsing, A. J. (2017). Top predators constrain
mesopredator distributions. *Nature Communications*, *8*(May), 1–7.
https://doi.org/10.1038/ncomms15469

Nüchel, J., Bøcher, P. K., Xiao, W., Zhu, A.-X., & Svenning, J.-C. (2018). Snub-nosed monkeys
(Rhinopithecus): potential distribution and its implication for conservation. *Biodiversity and*
Conservation, *27*(6), 1517–1538. https://doi.org/10.1007/s10531-018-1507-0

Öckinger, E., Schweiger, O., Crist, T. O., Debinski, D. M., Krauss, J., Kuussaari, M., Petersen, J. D.,
Pöyry, J., Settele, J., Summerville, K. S., & Bommarco, R. (2010). Life-history traits predict
species responses to habitat area and isolation: a cross-continental synthesis. *Ecology*
Letters, *13*(8), no--no. https://doi.org/10.1111/j.1461-0248.2010.01487.x

Palacio, R.D., et al. (2021). A data-driven geospatial workflow to map species distributions for
conservation assessments. *Diversity and Distributions*, 27(1):2559-2570.
http://dx.doi.org/10.1111/ddi.13424

Pimm, S. L., Russell, G. J., Gittleman, J. L., & Brooks, T. M. (1995). The Future of Biodiversity.
Science, *269*(5222), 347–350. https://doi.org/10.1126/science.269.5222.347

Pocock, M. J. O. (2011). Can traits predict species' vulnerability? A test with farmland passerines
in two continents. *Proceedings of the Royal Society B: Biological Sciences*, *278*(1711),
1532–1538. https://doi.org/10.1098/rspb.2010.1971

Prugh, L. R., Stoner, C. J., Epps, C. W., Bean, W. T., Ripple, W. J., Laliberte, A. S., & Brashares, J. S.
(2009). The Rise of the Mesopredator. *Bioscience*, *9*, 9.

Qian, T., Chi, Y., Xi, C., Li, Z., & Wang, J. (2020). Changes in the historical and current habitat
ranges of rare wild mammals in China: A case study of six taxa of small-to large-sized
mammals. *Sustainability (Switzerland)*, *12*(7), 1–19. https://doi.org/10.3390/su12072744

Rahbek, C., & Colwell, R. K. (2011). Species loss revisited. *Nature*, *473*(7347), 288–289.
https://doi.org/10.1038/473288a

Rayan, D. M., & Linkie, M. (2016). Managing conservation flagship species in competition: Tiger,
leopard and dhole in Malaysia. *Biological Conservation*, *204*, 360–366.

Recio, M.R., et al. (2015). Changes in Mediterranean mesocarnivore communities along urban and
ex-urban gradients. Current Zoology, 2015(5): p. 5. http://doi.org/10.1093/czoolo/61.5.793

Roberts, R. G., Flannery, T. F., Ayliffe, L. K., Yoshida, H., Olley, J. M., Prideaux, G. J., Laslett, G. M.,
Baynes, A., Smith, M. A., Jones, R., & Smith, B. L. (2001). New ages for the last Australian
megafauna: Continent-wide extinction about 46,000 years ago. Science.
https://doi.org/10.1126/science.1060264

Roberts, E. A., Sheley, R. L., & Lawrence, R. L. (2004). Using sampling and inverse distance
weighted modeling for mapping invasive plants. *Western North American Naturalist*.

Sandom, C., Faurby, S., Sandel, B., & Svenning, J.-C. (2014). Global late Quaternary megafauna
extinctions linked to humans, not climate change. *Proceedings of the Royal Society B:*
Biological Sciences, *281*(1787), 20133254. https://doi.org/10.1098/rspb.2013.3254

Shanee S, Shanee N, Campbell N, et al. (2014.) Biogeography and Conservation of Andean
Primates in Peru. In: Grow N., Gursky-Doyen S., Krzton A. (eds) High Altitude Primates.
Developments in Primatology: Progress and Prospects. Springer.

Smith, A. & Xie, Y. (2008) A Guide to the Mammals of China, *Princeton University Press*.

Somers, R. H. (1962). A New Asymmetric Measure of Association for Ordinal Variables. *American
Sociological Review*, *27*(6), 799. https://doi.org/10.2307/2090408

Takahashi, M. K., Eastman, J. M., Griffin, D. A., Baumsteiger, J., Parris, M. J., & Storfer, A. (2014).
A stable niche assumption-free test of ecological divergence. *Molecular Phylogenetics and
Evolution*. https://doi.org/10.1016/j.ympev.2014.03.003

Teng, S. N., Xu, C., Teng, L., & Svenning, J. C. (2020). Long-term effects of cultural filtering on
megafauna species distributions across China. *Proceedings of the National Academy of
Sciences of the United States of America*, *117*(1), 486–493.
https://doi.org/10.1073/pnas.1909896116

Thomas, C. D., Cameron, A., Green, R. E., Bakkenes, M., Beaumont, L. J., Collingham, Y. C.,
Erasmus, B. F. N., De Siqueira, M. F., Grainger, A., Hannah, L., Hughes, L., Huntley, B., Van
Jaarsveld, A. S., Midgley, G. F., Miles, L., Ortega-Huerta, M. A., Peterson, A. T., Phillips, O. L.,
& Williams, S. E. (2004). Letter to nature: Extinction risk from climate change. *Nature*,
427(145–148), 145–148.

Turvey, S. T., Crees, J. J., & Di Fonzo, M. M. I. (2015). Historical data as a baseline for conservation:
reconstructing long-term faunal extinction dynamics in Late Imperial–modern China.
Proceedings of the Royal Society B: Biological Sciences, *282*(1813), 20151299.
https://doi.org/10.1098/rspb.2015.1299

Turvey, S. T., Crees, J. J., Li, Z., Bielby, J., & Yuan, J. (2017). Long-term archives reveal shifting
extinction selectivity in China's postglacial mammal fauna. *Proceedings of the Royal Society
B: Biological Sciences*, *284*(1867), 20171979. https://doi.org/10.1098/rspb.2017.1979

Tylianakis, J. M., Tscharntke, T., & Lewis, O. T. (2007). Habitat modification alters the structure of
tropical host–parasitoid food webs. *Nature*, *445*(7124), 202–205.
https://doi.org/10.1038/nature05429

Villavicencio, N. A., Lindsey, E. L., Martin, F. M., Borrero, L. A., Moreno, P. I., Marshall, C. R., &
Barnosky, A. D. (2016). Combination of humans, climate, and vegetation change triggered
Late Quaternary megafauna extinction in the Última Esperanza region, southern Patagonia,
Chile. *Ecography*. https://doi.org/10.1111/ecog.01606

Wake, D. B., & Vredenburg, V. T. (2008). Are we in the midst of the sixth mass extinction? A view
from the world of amphibians. *Proceedings of the National Academy of Sciences*,
105(Supplement 1), 11466–11473. https://doi.org/10.1073/pnas.0801921105

Wallach, A. D., Izhaki, I., Toms, J. D., Ripple, W. J., & Shanas, U. (2015). What is an apex predator.
Oikos, *124*(11), 1453–1461.

Wan, X., Jiang, G., Yan, C., He, F., Wen, R., Gu, J., Li, X., Ma, J., Stenseth, N. C., & Zhang, Z. (2019).
Historical records reveal the distinctive associations of human disturbance and extreme
climate change with local extinction of mammals. *Proceedings of the National Academy of*
Sciences of the United States of America, *116*(38), 19001–19008.
https://doi.org/10.1073/pnas.1818019116

Wan, X., & Zhang, Z. (2017). Climate warming and humans played different roles in triggering
Late Quaternary extinctions in east and west Eurasia. *Proceedings of the Royal Society B:*
Biological Sciences, *284*(1851), 20162438. https://doi.org/10.1098/rspb.2016.2438

Wang, S. W., & Macdonald, D. W. (2009). Feeding habits and niche partitioning in a predator
guild composed of tigers, leopards and dholes in a temperate ecosystem in central Bhutan.
Journal of Zoology, *277*(4), 275–283.

Wang, L., et al.(2016.) The geographical distribution of grey wolves (*Canis lupus*) in China: a
systematic review. Zoological Research.
http://doi.org/10.13918/j.issn.2095-8137.2016.6.315

Wen, R. S. (2013). *Geographical distribution of wild animals in ancient China*. Scientific and
Technological Press of Shandong.*(In Chinese)*

Wen, R. S. (2009). *The distribution and changes of rare wild animals in China*. Scientific and
Technological Press of Shandong.*(In Chinese)*

Wen, R. S. (2018). *The distribution and changes of rare wild animals in China(sequel)*. Scientific
and Technological Press of Shandong.*(In Chinese)*

Yap, T. A., Koo, M. S., Ambrose, R. F., Wake, D. B., & Vredenburg, V. T. (2015). Averting a North
American biodiversity crisis. *Science*, *349*(6247), 481–482.
https://doi.org/10.1126/science.aab1052

Zhao, X., Ren, B., Garber, P. A., Li, X., & Li, M. (2018). Impacts of human activity and climate
change on the distribution of snub-nosed monkeys in China during the past 2000 years.
Diversity and Distributions, *24*(1), 92–102. https://doi.org/10.1111/ddi.12657

Zhou, Y. H., & Zhang, P. (2013). Distribution and vicissitude of gibbons（*Hylobatidae*）in China
during the last 500 years. Acta Theriologica Sinica, 33(3), 258–266.(In Chinese)

Fig.1 Research area (blue) in eastern China and the Heihe-Tengchong Line (red).

I selected 22 province-level regions in eastern continent China as the research area (excluding Tibetan Plateau

part in Sichuan and Yunnan) which roughly corresponds with the eastern region of the Heihe-Tengchong Line,

containing approximately 90% of the human population of China from ancient to modern eras (Hu 1935).

Fig.2 The PLS-SEM models.

In the triangle, the arrow "a" is the Anthropogenic influence on Apex predators; the arrow "b" is the Apex

predators' influence on Meso-predators; the arrow "c" is the direct Anthropogenic influence on Meso-predators.

The overall Anthropogenic influence on Meso-predators is: s=a*b+c. Exemplified by wolves in Period 1 and

Period 4: (i)HPD exerted a slight negative direct influence: -0.030. HPD also exert a large positive indirect

805   influence by kill the apex predators:(-.466)*(-0.244)≈0.114. The total HPD influence is 0.084, positive.HPD

806   exerted a large negative direct influence:-0.509. (ii)HPD also exert a smaller positive indirect influence by kill the

807   apex predators:(-.531)*(-0.169)≈0.090. The total HPD influence is -0.419, negative.

810 **Fig.3** The change of average probability of predators from Period 1 to Period 4, after the carnivores' name are the

811 total change proportion from Period 1 to Period 4: $(S_{Period\ 4}-S_{Period\ 1})/S_{Period\ 1}$.

826   **Fig.4** The average distribution altitude and average distribution slope of each kind of carnivores in each period.

828   a.Average altitude(meter),the green line is average altitude of research area(682m).

830   b.Average slope(°),the green line is average slope of research area(3.61°).

837   **Fig.5** The carnivore distribution change proportions from Period 1 to Period 4 in each altitude group, i.e.:

838    $(S_{Period\ 4}-S_{Period\ 1})/S_{Period\ 1}$.

839
840

841   **Fig.6** HPD Influence on predator distribution (Somers' dyx)

842
843

Fig.7 Criticizing the presence-only method

a.The bear presence records(red points) in Period 2, both frames are almost empty; b. The bear presence records(red points) and absence records(blue points) in Period 2, there are many absence records in the blue frame, but few in the brown frame; c. The IDW reconstruction of bear distribution based on b., high-probability area(red) and low-probability area(blue), the blue frame is fulfilled by low-probability area but the brown frame is fulfilled by high-probability area.

a   b   c

 Table.1 The distribution of carnivores, human population density(HPD), and Altitude/Slope

 Table.2 The results of PLS-SEM

		a.HPD→Apex	b.Apex→Meso	c.HPD→Meso (directly)	s.HPD→Meso (total)
Wolves	Period 1	-0.466	-0.244	-0.030	0.084
	Period 2	-0.477	-0.298	-0.218	-0.076
	Period 3	-0.481	-0.211	-0.213	-0.112
	Period 4	-0.531	-0.169	-0.509	-0.419
Foxes	Period 1	-0.466	-0.143	-0.067	0.000
	Period 2	-0.476	-0.232	-0.151	-0.040
	Period 3	-0.481	-0.107	-0.208	-0.156
	Period 4	-0.529	-0.278	-0.348	-0.201
Civets	Period 1	-0.460	0.390	0.155	-0.025
	Period 2	-0.472	0.250	0.256	0.138
	Period 3	-0.476	0.361	0.173	0.002
	Period 4	-0.529	0.326	0.065	-0.108
Dholes	Period 1	-0.460	0.678	0.032	-0.280
	Period 2	-0.467	0.686	0.140	-0.181
	Period 3	-0.474	0.648	-0.001	-0.309
	Period 4	-0.529	0.683	0.032	-0.329
Mustelids	Period 1	-0.467	0.032	0.225	0.210
	Period 2	-0.478	-0.039	0.223	0.238
	Period 3	-0.479	0.232	0.340	0.229
	Period 4	-0.530	0.075	-0.032	-0.071

894   **Table.3** Different results of IDW, Last Observation(100km*100km) & Last Observation(50km*50km)

	IDW	Last Observation 100km*100km	Last Observation 50km*50km
Period 1	41.4%	73.6%	31.8%
Period 2	42.4%	67.1%	28.3%
Period 3	46.5%	61.9%	24.2%
Period 4	51.9%	34.4%	11.3%

Supplementary Information:

934   **Fig.S1** Flow Chart of the procedure "Interpolation of Presence & Absence Records in Same"

938   **Fig.S2** The carnivore distribution in each altitude group in each period.

939　　　　　　　　　　HPD(Person(s)/km^2)

 FIg.S3 Relations between each predator and others (Kendall's tau-b)

961   **Table.S1** Qualifying the valid data.

Standards	Qualification
Only livestock	Invalid data
Only wild mammals with body mass ≥45kg	Invalid data
Wild mammals with body mass ≥ 45kg, and wild mammals with body mass <45kg	Valid data
Only wild mammals with body mass <45kg	Valid data

962
963
964

965   **Table.S2** Folk name of predators in local gazetteers and relevant species in scientific taxonomy
966

Folk name in pre-modern China	Scientific name	English name
Tiger	*Panthera tigris*	Tiger
Leopard	*Panthera pardus*	Leopard
	Neofelis nebulosa	Clouded Leopard
	Uncia uncia	Snow Leopard
Bear[1.]	*Helarctos malayanus*	Sun Bear
	Ursus arctos	Brown Bear
	Ursus thibetanus	Asian Black Bear
Wolf	*Canis lupus*	Wolf
Fox	*Vulpes corsac*	Corsac Fox
	Vulpes ferrilata	Tibetan Fox
	Vulpes vulpes	Red Fox
Civet (viverrids & small wild felids)	*Vormela peregusna*	Marbled Polecat
	Catopuma temminckii	Asian Golden Cat
	Felis bieti	Chinese Mountain Cat
	Felis chaus	Jungle Cat
	Felis manul	Pallas' Cat
	Felis silvestris	Wildcat
	Pardofelis marmorata	Marbled Cat
	Prionailurus bengalensis	Leopard Cat
	Chrotogale owstoni	Owston's Palm Civet
	Arctictis binturong	Binturong
	Arctogalidia Trivirgata	Small-Toothed Palm Civet
	Paguma larvata	Masked Palm Civet

	Paradoxurus hermaphroditus	Common Palm Civet
	Prionodon pardicolor	Spotted Linsang
	Viverra megaspila	Large-Spotted Civet
	Viverra zibetha	Large Indian Civet
	Viverricula indica	Small Indian Civet
Dhole	*Cuon alpinus*	Dhole
Mustelid (Badger, Otter, etc.)	*Aonyx cinerea*	Asian Small-Clawed Otter
	Lutra lutra	Eurasian Otter
	Lutrogale perspicillata	Smooth-Coated Otter
	Arctonyx collaris	Hog Badger
	Martes flavigula	Yellow-Throated Marten
	Martes foina	Beech Marten
	Meles leucurus	Asian Badger
	Melogale moschata	Chinese Ferret Badger
	Melogale personata	Burmese Ferret Badger
	Mustela altaica	Mountain Weasel
	Mustela erminea	Ermine
	Mustela eversmanii	Steppe Polecat
	Mustela kathiah	Yellow-Bellied Weasel
	Mustela nivalis	Least Weasel
	Mustela sibirica	Siberian Weasel
	Mustela strigidorsa	Back-Striped Weasel

1. I did not distinguish the brown bear, sun bear and Asian black bear in this research, because the majority of local gazetteers only refer "熊(bear)", only a few of them distinguish "Asian black bear", "brown bear" or "sun bear(suspected)". There are specific name of panda, not included in this research.

Table.S3 The change of average probability of carnivores from Period 1 to Period 4.

	Tigers	Leopards	Bears	Wolves	Foxes	Civets	Dholes	Mustelids
Period 1	63.8%	59.3%	36.0%	41.4%	60.9%	47.2%	36.1%	60.9%
Period 2	61.1%	59.5%	38.2%	42.4%	65.5%	51.1%	35.2%	71.8%
Period 3	53.4%	54.2%	31.6%	46.5%	67.8%	61.9%	40.1%	76.7%
Period 4	49.3%	59.3%	24.0%	51.9%	69.6%	67.3%	40.9%	83.1%

Table.S4 The average distribution probability in each altitude group in each period.

alt1

	P1	P2	P3	P4
Tigers	37.2%	29.6%	28.4%	21.5%
Leopards	25.0%	22.6%	22.3%	20.4%
Bears	10.8%	13.0%	11.3%	6.5%
Wolves	45.6%	43.4%	43.8%	33.3%
Foxes	61.5%	66.2%	64.3%	56.7%
Civets	42.5%	49.8%	55.2%	58.7%
Dholes	16.4%	18.0%	17.2%	17.7%
Mustelids	72.7%	86.1%	87.5%	80.0%

alt2

	P1	P2	P3	P4
Tigers	70.4%	67.4%	62.0%	57.6%
Leopards	59.5%	61.2%	54.3%	56.4%
Bears	31.0%	32.8%	26.8%	16.7%
Wolves	44.3%	39.9%	43.5%	46.3%
Foxes	59.8%	62.7%	60.0%	62.0%
Civets	53.3%	64.4%	73.8%	74.5%
Dholes	46.4%	45.5%	46.5%	40.7%
Mustelids	70.3%	79.2%	81.0%	84.0%

alt3

	P1	P2	P3	P4
Tigers	79.9%	72.3%	64.1%	59.2%
Leopards	69.3%	67.3%	61.8%	69.6%
Bears	42.7%	43.0%	35.0%	27.7%
Wolves	34.7%	33.9%	37.9%	48.7%
Foxes	60.1%	62.3%	61.0%	65.4%
Civets	53.4%	60.5%	66.0%	77.6%
Dholes	46.4%	44.6%	44.7%	45.6%
Mustelids	63.7%	73.0%	75.6%	83.1%

	alt4			
	P1	P2	P3	P4
Tigers	73.5%	72.1%	59.8%	53.7%
Leopards	70.0%	69.8%	64.1%	70.6%
Bears	45.4%	45.2%	34.8%	27.0%
Wolves	47.3%	48.9%	56.8%	64.7%
Foxes	66.8%	68.7%	73.3%	77.7%
Civets	47.9%	47.5%	57.6%	65.3%
Dholes	38.5%	34.6%	44.9%	43.6%
Mustelids	55.8%	65.7%	73.7%	82.8%

	alt5			
	P1	P2	P3	P4
Tigers	57.9%	64.2%	52.8%	54.4%
Leopards	72.6%	76.8%	68.7%	79.5%
Bears	49.9%	57.2%	50.1%	42.0%
Wolves	35.2%	46.2%	50.5%	66.7%
Foxes	56.6%	67.6%	80.4%	86.3%
Civets	38.7%	33.2%	57.1%	60.6%
Dholes	32.9%	33.4%	47.4%	57.1%
Mustelids	41.9%	55.0%	65.4%	85.7%

994 **Table.S5** Relations between each kind of carnivores and others (Kendall's tau-b coefficient).

Tigers

	Leopards	Bears	Wolves	Foxes	Civets	Dholes	Mustelids
Period 1	0.551	0.410	-0.076	-0.009	0.288	0.554	0.139
Period 2	0.613	0.396	-0.047	-0.079	0.191	0.480	0.068
Period 3	0.613	0.400	-0.046	-0.070	0.240	0.452	0.113
Period 4	0.603	0.479	0.017	-0.172	0.232	0.481	0.054

Leopards

	Tigers	Bears	Wolves	Foxes	Civets	Dholes	Mustelids
Period 1	0.551	0.513	-0.135	-0.124	0.133	0.477	-0.102
Period 2	0.613	0.541	-0.138	-0.118	0.079	0.480	-0.087
Period 3	0.613	0.480	-0.045	0.019	0.174	0.525	0.028
Period 4	0.603	0.561	0.115	0.003	0.212	0.525	0.065

Bears

	Tigers	Leopards	Wolves	Foxes	Civets	Dholes	Mustelids
Period 1	0.410	0.513	-0.125	-0.096	0.166	0.418	-0.075
Period 2	0.396	0.541	-0.117	-0.146	0.001	0.381	-0.128
Period 3	0.400	0.480	-0.088	-0.041	0.132	0.382	-0.039
Period 4	0.479	0.561	0.030	-0.139	0.197	0.385	0.030

Wolves

	Tigers	Leopards	Bears	Foxes	Civets	Dholes	Mustelids
Period 1	-0.076	-0.135	-0.125	0.295	-0.056	-0.023	0.076
Period 2	-0.047	-0.138	-0.117	0.332	-0.027	-0.065	0.071
Period 3	-0.046	-0.045	-0.088	0.303	-0.084	-0.047	0.101
Period 4	0.017	0.115	0.030	0.373	-0.013	0.052	0.100

Foxes

	Tigers	Leopards	Bears	Wolves	Civets	Dholes	Mustelids
Period 1	-0.009	-0.124	-0.096	0.295	-0.044	-0.034	0.151
Period 2	-0.079	-0.118	-0.146	0.332	0.017	-0.132	0.121
Period 3	-0.070	0.019	-0.041	0.303	-0.025	-0.014	0.163
Period 4	-0.172	0.003	-0.139	0.373	-0.059	-0.042	0.208

Civets

	Tigers	Leopards	Bears	Wolves	Foxes	Dholes	Mustelids
Period 1	0.288	0.133	0.166	-0.056	-0.044	0.256	0.340
Period 2	0.191	0.079	0.001	-0.027	0.017	0.176	0.336
Period 3	0.240	0.174	0.132	-0.084	-0.025	0.281	0.214
Period 4	0.232	0.212	0.197	-0.013	-0.059	0.216	0.171

Dholes

	Tigers	Leopards	Bears	Wolves	Foxes	Civets	Mustelids
Period 1	0.554	0.477	0.418	-0.023	-0.034	0.256	0.124
Period 2	0.480	0.480	0.381	-0.065	-0.132	0.176	0.056
Period 3	0.452	0.525	0.382	-0.047	-0.014	0.281	0.048
Period 4	0.481	0.525	0.385	0.052	-0.042	0.216	0.092

Mustelids

	Tigers	Leopards	Bears	Wolves	Foxes	Civets	Dholes
Period 1	0.139	-0.102	-0.075	0.076	0.151	0.340	0.124
Period 2	0.068	-0.087	-0.128	0.071	0.121	0.336	0.056
Period 3	0.113	0.028	-0.039	0.101	0.163	0.214	0.048
Period 4	0.054	0.065	0.030	0.100	0.208	0.171	0.092

997 **Table.S6**   HPD Influence on carnivore distribution (Somers' dyx)

	Tigers	Leopards	Bears	Wolves	Foxes	Civets	Dholes	Mustelids
Period 1	-0.112	-0.296	-0.293	0.030	0.042	0.028	-0.190	0.181
Period 2	-0.161	-0.304	-0.351	-0.053	0.013	0.150	-0.129	0.215
Period 3	-0.134	-0.264	-0.290	-0.100	-0.113	0.054	-0.188	0.179
Period 4	-0.254	-0.381	-0.289	-0.294	-0.173	-0.059	-0.222	-0.034

Table.S7 Criticizing the last observation precedure

	P1	P2	P3	P4	
Data points of Last Observation					Red points: Presence records
Last Observation 100km*100km					Black grids: Presence
Last Observation 50km*50km					Black grids: Presence
Data points of this research					Red points: presence records Blue points: Absence records
IDW					Red: High probability Blue: Low probability