Title: What's left in the tank? Identification of non-ascribed aquarium's coral collections with DNA barcodes

as part of an integrated diagnostic approach

Authors: Luigi Colin[1]; Daniel Abed-Navandi[2]; Dalia A. Conde[3,4]; Jamie Craggs[5]; Rita da Silva[3];

Max Janse[6]; Björn Källström[7]; Alexander Pearce-Kelly[8]; Chris yesson[1]

Institutions:

1) Institute of Zoology, Zoological Society of London, Regent's Park, London NW1 4RY, UK

2) Haus des Meeres - Aqua Terra Zoo, Fritz Gruenbaum Platz 1, 1060 Vienna, Austria

3) Interdisciplinary Centre on Population Dynamics (CPop), University of Southern Denmark,

Biology Department, Syddansk Universitet, Campusvej 55, DD-5230 Odense M, Denmark

4) Conservation Science Alliance, Species360, 7900 International Drive, Suite 1040 Bloomington,

MN, 55425 USA

5) Horniman Museum & Gardens, 100 London Road, Forest Hill, London SE23 3PQ, UK

6) Royal Burgers' Zoo, Antoon van Hooffplein 1, 6816 SH Arnhem, the Netherlands

7) Maritime Museum & Aquarium, Karl Johansgatan 1-3, SE-41459, Göteborg, Sweden

8) Zoological Society of London, Regent's Park, London NW1 4RY, UK

Communicating author: luigi.colin@ioz.ac.uk

Keywords: Coral, DNA Barcoding, Aquarium, Identification, Blast match, Species

Abstract

The unprecedented threats to coral reef ecosystems from global climate change (GCC) require an urgent response from the aquarium community, which is becoming an increasingly vital coral conservation resource. Unfortunately, many hermatypic corals in aquaria are not identified to species level, which hinders assessment of their conservation significance. Traditional methods of species identification using morphology can be challenging, especially to non-taxonomists. DNA barcoding is an option for species identification of Scleractinian corals, especially when used in concert with morphology-based assessment. This study uses DNA barcodes to try to identify aquarium specimens of the diverse reef-forming genus *Acropora* from 127 samples. We identified to our best current knowledge, to species name 44% of the analysed samples and provided provisional identification for 80% of them (101/127, in the form of a list of species names with associate confidence values). We highlighted a sampling bias in public nucleotide sequences repertories (e.g.: GenBank) towards more charismatic and more studied species, even inside a well-studied genus like *Acropora*. In addition, we showed a potential "single observer" effect with over a quarter of the reference sequences used for these identifications coming from the same study. We propose the use of barcoding and query matching as an additional tool for taxonomic experts and general aquarists, as an additional tool to increase their chances of making high confidence species-level identifications. We produce a standardised and easily repeatable methodology to increase the capacity of aquariums and other facilities to assess non-ascribed species, emphasising the value of integrating this approach with morphological identification optimising usage of authoritative identification guides and expert opinion.

Introduction

It is widely acknowledged that coral reef ecosystems are facing unprecedented threats from global climate change (GCC) e.g.(Pratchett et al. 2013; Hoegh-Guldberg et al. 2017; Cowburn et al. 2018;

Hughes et al. 2018; Masson-Delmotte et al. 2018; Brondizio et al. 2019; Sheppard et al. 2020).

Warm water reef-building corals create the most biodiverse ecosystems on the planet and have been

estimated to support 830'000 species of multi-cellular plants and animals worldwide (Bellwood and

Hughes 2001; Mora et al. 2008, 2011; Fisher et al. 2015), providing a variety of habitats for fish,

invertebrates and other taxa in shallow tropical seas (Bellwood and Hughes 2001). *Acropora* corals

play a key functional ecosystem role as they are the major reef builders in the majority of warm

water reefs ecosystems (Fukami et al. 2000).

Corals in aquaria

Ex situ conservation can play an important role in ensuring the survival of many species (Turley

1999; Hutchins 2003; Raven 2004; Blanco et al. 2009; Fa et al. 2011; Leus et al. 2011; Lacy 2013;

Skibins and Powell 2013). Given the severity of the species and ecosystem level climate change

threats to corals, and the alarming rates of recent losses (Masson-Delmotte et al. 2018), coral

collection can potentially play a valuable role in future conservation efforts. Aquaria institutions

therefore have a vitally important conservation remit to support coral restoration strategies allowing

for both sexual and asexual coral recruitment, biobanking and a wide range of research, education

and thus contribution towards *in situ* conservation activities. Early attempts in the 1980s at keeping

corals in aquaria faced difficulties (Borneman 2008; Brunner 2012) it was until 1980 that *Acropora*

was the first genus to be successfully cultured by Stüber (Borneman 2008). From the 1990s

onwards coral husbandry boomed, reaching a level where virtually all families of zooxanthellate

corals were not only being maintained for many years, but were being propagated and traded

between private and public aquaria (Borneman 2008). Specimens conserved in Aquaria thus,

became a highly valuable resource for restoration strategies (Rinkevich 1995).

Identification of corals

Coral taxonomy is a traditionally difficult discipline, subject to all the challenges of modern taxonomy (Godfray 2002). The genus *Acropora* is among the most diverse and geographically widespread reef building corals (Wallace 1999). Research on *Acropora* suggests that species limits may be sometimes narrower, sometimes broader, than generally perceived (Wallace and Willis 1994). Somewhat blurry species boundaries are confounded by hybridisation (Willis 1997; Hatta et al. 1999; Van Oppen et al. 2002; Vollmer and Palumbi 2002; Wolstenholme 2004; Willis et al. 2006; Palumbi et al. 2012; Wei et al. 2012; Isomura et al. 2013), often worsened by the possibility of synchronised spawning events (Harrison et al. 1984; Babcock et al. 1986; Márquez et al. 2002; Van Oppen et al. 2002). Currently the number of officially recognised *Acropora* species are between 135 (WORMS (Horton et al. 2020)) and 163 (Corals of the World (Veron et al. 2020)), reaching 186 if we include the "*taxon inquirendum*" on WORMS (Horton et al. 2020). There have been different attempts at grouping these species within the genus *Acropora* (Wallace 1999). These attempts lead to the creation of the concept of species group (aka syngameon). The concept of species groups, first defined for convenience of identification and without implying taxonomic relation, has been revised to reflect the evolution and phylogeny of the genus (Wallace 1999; Wallace et al. 2012). The use of genetic tools for species identification has provided another line of evidence for species delimitation, further adding to the blurred boundaries between species groups, species, and sub-species.

Traditionally, identification of coral has relied on morphological skeletal features, with the great majority of taxa originally described following typological species concepts defined around a century ago (Best et al. 1984; Todd 2008). The relationship between morphological variation within-species and the environment has been a source of contention since the late 1800s (Todd 2008). Whether this variability is due to different underlying genotypes or plastic phenotypes is unclear. Evidence supports both scenarios (difference in phenotype e.g.: (Willis 1985; Ayre and

Willis 1988; Todd et al. 2004)– Plastic phenotypes e.g.: (Foster 1979; Miller 1994; Bruno and

Edmunds 1997; Muko et al. 2000; Todd et al. 2004)) and these scenarios are not necessary mutually

exclusive but probable to operate simultaneously (Foster 1979; Amaral 1994; Todd 2008). This has

particular relevance in the face of continuous GCC, that is creating an increasingly unpredictable

wild reef environment.

Notwithstanding the taxonomic difficulties, the identification of coral species can be challenging

even in optimal 'wild' conditions. However, identification requirements are far more than just for

well documented field collections, but also cover aquaria collections, that in some cases are held in

conditions dramatically different from the wild. Furthermore, customs officials rely in proper

taxonomic identification even at the genera level for threatened taxa by the international wildlife

trade (AC25 Doc. 23 CITES 2011), although currently *Acropora* are covered by the blanket listing

of all Scleractinia under CITES appendix II. A status of these identification difficulties in aquaria

can be demonstrated by an analysis of Species360 Zoological Information Management Software

(ZIMS), adopted by more than 95 aquariums in 24 countries. Assuming correct identification ZIMS

indicates c. 42.9% of corals of the orders Scleractinia, Alcyonacea, Helioporacea, Antipatharia,

Corallimorpharia, Pennatulacea, and Zoanthidea in aquaria are identified to genus or higher

taxonomic level (ZIMS list of species holdings – 4th June 2020). Most of the records correspond to

the order Scleractinia, which is also the order with more species represented in aquariums.

Corallimorpharia is, however, the order with the highest number of reported individuals in

aquariums members of the Species360 network (Fig. 1). In addition to this the non-species

associated genera (e.g.: *Acropora* sp.) is usually recorded once in the ZIMS database and most other

aquarium inventory systems, which possibly significantly underestimates the actual number of non-

identified specimens. Moreover, there are currently no confidence values ascribed to any taxonomic

level a collection allocates to a specimen which means that there is likely to mean that a significant

number of species ascribed specimens are actually at the lower end of the confidence spectrum and

would benefit from being reassessed.

In many aquariums', corals come from three sources: fragments from other aquaria, via ornamental

trade, or from confiscated and seized shipments (which aquariums are usually asked to

accommodate). All three sources present identification challenges as often exact knowledge from

their wild origin is absent. This specially a challenge for confiscated specimens since they have

none or relatively poor source records, sample integrity, and any other additional background

information that might be of assistance for the identification process. This, together with the

difficulties caused by environmental phenotypic plasticity makes coral identification in aquariums a

particularly challenging task. Data from the Convention on the International Trade of Endangered

Species of Flora and Fauna (CITES) shows a total of 4557 instances of coral confiscation from

1982 to 2018 worldwide, for a total of 27027 individual items and 5461 kg per year (CITES data -

https://trade.cites.org 02/07/2020). Of these, less than 20% are reported at species level, meaning

that for most cases we know only the genus, family or even order of the specimens confiscated.

Given these challenges CITES currently requires identification only to the order level.

DNA Barcoding

An alternative or complementary diagnostic option to morphological identification is genetic

analysis. DNA Barcodes have been proposed by Hebert et al. (Hebert et al. 2003) as a way to

circum-vent the "limitations inherent in morphology-based identification systems". The core

concept of DNA barcoding is the use of a standard sequence corresponding to a single homologous

region that can be amplified with universal primers that is sufficiently variable to distinguish

between species – usually represented by the mitochondrially encoded cytochrome c oxidase I (MT-

CO1 or CO1 or COX1) (Hebert et al. 2003).

Beside the inherent limitations (Shearer et al. 2002; Ferguson 2002; Moritz and Cicero 2004;

DeSalle et al. 2005; Meyer and Paulay 2005; Prendini 2005; Cognato 2006; Hickerson et al. 2006;

Meier et al. 2006, 2008; Little and Stevenson 2007), DNA barcoding is especially attractive as an

option for species identification of scleractinian corals, as this method bypasses the problems

caused by phenotypic plasticity or life stage (larva, juvenile or adult). Unfortunately, the proposed

'universal' Barcoding region CO1 is at least 10-20 times slower in Anthozoa than the standard

vertebrate mtDNA mutation rate, and 2-5 times slower than the average Anthozoan nuclear

sequence(Van Oppen et al. 1999; Shearer et al. 2002; Hellberg 2006; Huang et al. 2008; Chen et al.

2009; Ortman et al. 2010) with zero interspecific divergence for many species (Shearer and

Coffroth 2008). This has led to the use of a number of different barcoding regions for corals

(Shearer et al. 2002).

The diverse, reef forming Acropora group are widely grown in aquaria and have been a target for

barcoding work. Potential barcodes were reviewed by Shearer et al. (Shearer et al. 2002) examining

both nuclear and mitochondrial regions. Substitution rates of mitochondrial genes in *Acropora* tend

to be lower relative to the nuclear counterparts mentioned by Shearer (i.e. Pax-C intron; internal

transcribed spacer 1 and 2 (ITS) (Shearer et al. 2002). Although, Van Oppen et al. (van Oppen et al.

1999) and Vollmer et al. (Vollmer 2002) have highlighted the mitochondrial putative control region

(mtCR) is more variable than the other mitochondrial regions. Subsequently Shearer et al. (Shearer

et al. 2002) compared the genetic divergence of mitochondrial and nuclear regions in Scleractinia,

highlighting how the mtCR region has a similar range of divergence to nuclear coding regions like

the Pax-C intron, but is much slower than the hyper-variable ITS1-2. Notwithstanding all the above

mentioned work, *Acropora*'s molecular taxonomy remains to be resolved, and to do so it would

require extensive taxonomic work and genome-wide markers (Cowman et al. 2020).

This study seeks to build on these studies by investigating the utility of DNA barcoding as a method

for provisional species identification of *Acropora* collections in aquaria, as an aid to morphological

taxonomy rather than an alternative, and as part of an integrated diagnostic approach.

Material and Methods

Study Area and Collection

As part of the "idCoral" project we collected 224 samples across seven institutions (Aquarium de Paris (France), Chester's Zoo (UK), Haus des Meeres-Vienna (Austria), Horniman Museum and Gardens (UK), Royal Burgers' Zoo (Netherlands), Tierpark Hagenbeck Aquarium (Germany), ZSL London Zoo (UK)). Branch tip fragments of 2 cm were collected and stored in single ethanol (+95%) vials labelled with a unique identifier. During the sample collection the colonies were tagged and photographed with collection dates and tank locations were recorded. The photographs and tags were also recorded in the "idCoral" database (database - idcoral.org (2020)). After collection samples were stored at -20°C until extraction, with an initial change of ethanol after 1-2 days aimed at reducing degradation. We analysed 127 *Acropora* samples from these collections before COVID-19 restrictions stopped further progress on this project.

DNA Extraction, Amplification and Sequencing

The DNA extractions were undertaken using the DNeasy PowerSoil Pro Kit from Qiagen© with a modified version of the manufacturer's protocol. An incubation step with proteinase K was added and the physical breaking of the cell structure with a chemical dissolution was removed from the protocol as it resulted in excessive shearing of the DNA (Online Resource 1). Extracted DNA was then preserved at -20°C, and the DNA concentration and quality tested with both Nanodrop assay and gel electrophoresis. The DNA was diluted to standardise the concentration to between 5 to 20 µg/ml.

Two genes, PaxC and mtCR were selected as barcodes based on an assessment of their variability and availability as reference sequences (Van Oppen et al. 2001; Shearer et al. 2002). The ITS1-2 region was excluded due to the lower number of taxa available as reference sequences. PCR

amplification was conducted with Q5® High- Fidelity DNA Polymerase mixed following

manufacturer's indication. 3μl of DNA template and 0.5 μl of Bovine Serum Albumin (BSA)1μg

were added, reaching a final volume of 25μl with RNA free water. For the PaxC gene the primer

used were PaxC FP1 5'-TCC AGA GCA TTA GAG ATG CTG G-3' and PaxC RP1 5'-GGC GAT

TTG AGA ACC AAA CCT GTA-3' (Van Oppen et al. 2001) with a protocol of 98°C for 30s,

followed by 31 cycles at 98°C for 10 s, 62°C for 30 s and 72°C for 60 s, ending with a final phase

of 72°C for 2min. For the mtCR the primer used were RNS2: 5'-CAG AGT AAG TCG TAA CAT

AG-3' and GR: 5'-AAT TCC GGT GTG TGT TCT CT-3' (Suzuki et al. 2008) with a protocol of

98°C for 30s, followed by 40 cycles at 98°C for 10 s, 62°C for 15 s and 72°C for 60 s, ending with

a final phase of 72°C for 5min.

Successful PCR amplifications (assessed by gel electrophoresis) were sent to Eurofins Genomics

for PCR cleaning and custom Sanger sequencing on both the forward and the reverse primers using

cycle sequencing technology (dideoxy chain termination / cycle sequencing) on ABI 3730XL

sequencing machines. The returned sequences were manually trimmed to remove poor quality

sections and assembled using Geneious Prime® (2019.1.2.).

Identification

Taxonomy

Both the barcode gap and specimen identifications were initially performed at the species level, but

were also performed at the level of species group, based on the revision of Acropora by Wallace et

al. (Wallace et al. 2012), to provide an additional layer of information in the final integrated

approach.

Barcoding Gap

Reference sequences for the two targeted regions were collated. FASTA files were created with the records matching the following query on GenBank: mtCR=""Acroporidae"[Organism] AND mitochondrial "complete genome" OR "control region" OR "putative control region" OR mitochondrial control region"; PaxC=""Acroporidae"[Organism] AND PaxC[gene] OR pax-c[gene]". Two multiple alignments were performed, one for each gene, using the Geneious$^{®}$ algorithm (progressive pairwise aligner) and trimmed. For both the produced references alignments distance matrices were calculated using the function dist.dna (R package apex (Jombart et al. 2017)) and with different models (Raw ; JC69; K80; K81; F84; BH87; paralin; indel Y; indelblock). Successively the frequency distribution of intra/inter-specific distances was plotted. The quantile function (R package "stats") was used to define the 99; 50; 1% quantile confidence interval used in the blast match analysis. Note: the two alignments were constructed and used only for the specific purposes of examining the barcoding gap for the targeted region and determining potential confidence interval for the blast match analysis.

Phylogenetic tree

Separate unrooted Maximum-likelihood (ML) phylogenetic trees were created, one for each targeted region. The references obtained from the previously mentioned query were cleaned for duplicated sequences and subsequently aligned. The two phylogenetic analyses were performed using the program IQ-TREE, with default settings and automatic model selection (Minh et al. 2013; Nguyen et al. 2015) with analysis conducted on the usegalaxy.eu public server (Afgan et al. 2018).

Blast match

Query based specimen identification with blast match (Benson 2000) was determined using a custom Blastn database. All records from the query ""Acroporidae"[Organism]" on GenBank were

downloaded and included in the local database built with the packages rBLAST (Hahsler and Nagar

2019) and rentrez (Winter 2017). A local Blast match (Camacho et al. 2009) was ultimately

performed. To filter for species match with identical scores a decision tree (Fig. 2) was designed

relying on Bit score, Percentage identity, and number of mismatches. Matches shorter than 2/3 of

the original sample sequence query were removed. The decision tree prioritised 100% pairwise

identity matches where possible, then to the highest bit score, then lowest number of base

mismatches. 100% pairwise identity matches to a species group rather than a single species name

were prioritised over single species name matches with lower pairwise identity with higher bit

score, both possible IDs were retained. We confronted the match from the two targeted regions to

obtain a single most probable ID. To account for conflicting matches between the two region we

relied on the 99; 50; 1% quantile distribution of the intraspecific distances obtained from the

barcoding gap analysis. Using pairwise identity threshold as confidence intervals to discern which

region offers the better match for the ID (R-scripts as Online Resource 3 and 4).

Results

258   From the 127 treated samples all were successfully extracted and, 107 mtCR and 100 PaxC had a

259   successful DNA amplification, respectively 84% and 79% with 76 samples overlap between the two

260   regions (Table 1). 348 reads were successfully sequenced, 84 mtCR sample and 90 PaxC. The

261   median sequence lengths were 734 post assembly for the mtCR region and 649 for the PaxC region

262   (Table 1).

Barcoding

Our GenBank query produced a total of 1348 mtCR sequences and 427 PaxC sequences. The mtCR

references query, once all sequence duplicates with the same species name were removed, produced

a 579-sequence alignment of length 914bp. The PaxC references query, after similar cleaning

produced a 352-sequence alignment of length 770 bp.

The mtCR alignment of 579 sequences resulted in a matrix of 335 241 pairwise genetic distances,

grouped by 57 species names. 6.4% of these (21 471) are intraspecific distances, while 93.6% (313

770) are interspecific distances (Fig. 3). The PaxC alignment of 352 sequences resulted in a matrix

of 123 904 genetic distances, grouped by 55 species names. 3% of these (3 726) are intraspecific

distances, while 97% (120 178) are interspecific distances (Fig. 3). Between the two groups there is

an overlap of 49 species names. Table 3 and 4 with the sources of the sequences used in these

alignments.

Fig. 3 highlights in blue the intraspecific variation (sequences with the same name on GenBank)

and in the light green the interspecific distances (sequences with different names on GenBank). In

both Fig. 3A) and 3B), and both barcoding regions, we see no discernable barcoding gap.

The quantile within species distribution produce the following distance threshold for our confidence

interval: mtCR 0.027 (99%); 0.0068 (50%); 0 (1%) and PaxC 0.17 (99%); 0.014 (50%); 0 (1%).

In Fig. 4, for the mtCR and the PaxC respectively, we can see the above-mentioned uneven

distribution of the GenBank records towards specific species and species groups. With example like

species group *humilis* showing 4 times the frequency of less represented species group like *loripes*.

Phylogeny

Both the mtCR (Online Resource 5) and the PaxC (Online Resource 6) phylogenies show clear sign

of polyphyly. Looking at our best matches (i.e.: 100% match to a reference), most of our sample

sequences match to single species – 63% of mtCR 100% matches and all of the PaxC 100%

matches. Focusing on just these sequences with 100% matches to a reference, we see that 0/19

mtCR sequences and 1/4 PaxC match to single-species clades (a monophyletic group composed of

more than one sample of a single taxon name). However, some of these match to species that a

polyphyletic within the tree (All of 100% mtCR matches tie to polyphyletic taxa and 75% for

PaxC).

Blast match

The local database for blast matching contained 510826 sequences. In Table 2 we see that 23 of our

76 samples (with both regions sequenced) were uniquely identified to a single species, while 13

samples matched to conflicting names. 20 of our samples had a unique species name match for the

PaxC region but mtCR region matched to 2 or more names with all but 6 of these including the

matching PaxC name. Very few samples matched to multiple species for both regions (4).

We have 47 % of samples matching to a single species name with both regions, 5% with the same

identical match. While we have 42% of the matches to a single species name that are in conflict,

25% of these conflicts are resolvable (i.e.: one of the two region with a match better fitting than the

other one, based on the defined confidence thresholds of each region, rather than directly on the

pairwise identity). Similar results for the other sample matching for two or more in at least one of

the two regions. Out of the 76 samples, in 15 cases we observe the second region adding

information to the first ones (Table 5 as Online Resource 2). Mainly from a mtCR multi name ID to

a single PaxC name (i.e.: sample P269 has 7 mtCR identical matches to 7 different species names,

and the PaxC matches at a lower rate to only one of them). In addition, the second region functions

as backup option whenever one of the two falls too short (5/76 sample have too low-quality match

to be trusted independently).

The complete list of our current best species name match, with species group information and

detailed genetic info is provided as Online Resource (Table 5 – Online Resource 2).

Discussion

Barcoding gap

Using DNA barcodes to identify unknown samples is only achievable if a well- studied, well-

sampled reference sequence database is available (Meyer and Paulay 2005), and even though

databases such as GenBank (which comprises the DNA DataBank of Japan (DDBJ), the European

Nucleotide Archive (ENA), and GenBank at NCBI - http://www.ncbi.nlm.nih.gov/genbank) and

BOLD (Barcode Of Life Data System - http://www.boldsystems.org) contain an ever increasing

number of species-level identified sequences (i.e.: 216 531 829 total sequences on GenBank as

April 2020), these databases are far from complete. The open nature of GenBank is destined to

produce a bias towards species with a higher research interest (i.e.: Human sequences constitute

56% of the total sequences on GenBank (Benson 2000)). The genus focus of our study, *Acropora*, is

one of the more commonly studied and widespread hard corals in the ocean (Wallace 1999; Fukami

et al. 2000) making it one of the most covered coral genera in GenBank. The reality is that even for

a widely studied group such as *Acropora* we can observe a disparity in the number of sequences

available for each species name (Fig. 4). The best example of this is the delta between two species

group: group *humilis* being the most sampled (251 mtCR and 61 PaxC references sequences from

our query) and group *robusta* one of the least sampled (11 mtCR and 13 PaxC references sequences

from our query). Note that both species group are formed by of the same number of species (8

species each (Wallace et al. 2012). However, while counting the number of sequences available for

a group may be an indication of the level of research that group receives, it does not mean that

groups with the most sequences have the best reference collection for a DNA barcoding project.

The most important quality for a reference sequence collection is it's representation of the

underlying species diversity, ideally covering a wide geographical and morphological range

(Vellend and Geber 2005; Rozenfeld et al. 2007; Ma et al. 2017), all-the-while using a marker that

shows sufficient variability to permit robust identifications. Sampling bias is a problem for all

specimen collections and this must always be kept in mind when interpreting results (Leray et al.

2019).

From our analysis we could not identify a clear barcoding gap in both our targeted regions (Fig. 3),

and furthermore we can observe a total overlap of the intra-specific distances by the inter-specific

distances. For both our targeted regions, even if more evident in the mtCR (Fig. 3A), we see that the

inter-specific distances follow a distribution reminiscent of a more common and more expected

barcoding distribution (Fig. 3B, mtCR). We performed the same barcoding gap analysis at the

higher taxonomic level using the species group information without obtaining a noticeable

difference. With the lack of a clear Barcoding gap, and accounting for the extensive criticism

present in the literature: For the use of distances in taxonomy and systematics (Ferguson 2002; Lee

2004; Moritz and Cicero 2004; DeSalle et al. 2005; Prendini 2005; Knapp et al. 2005; Meyer and

Paulay 2005; Cognato 2006; Hickerson et al. 2006; Little and Stevenson 2007; Meier et al. 2008);

for the drawbacks of using GenBank data (Meier et al. 2006); for the evidence of different rates of

evolution in Scleractinia (Van Oppen et al. 1999; Shearer et al. 2002; Kitahara et al. 2010). We

cannot establish a simple distance base threshold to use for our identification purposes, leading us

to depend more on the blast match.

Another potential issue with the use of these methods of identification is the source of the reference

sequences. With databases like GenBank, with little or no review on the uploaded information, the

question is the confidence that the initial identification of the now available reference sequences is

indeed correct (James Harris 2003). In our specific case after the processing we performed for the

mtCR reference alignment, out of the total 579 included 42% (242 sequences) come from a single

doctoral thesis project, while 5.4% (33 sequences) are from unpublished papers (Table 3). For the

PaxC 16% (58 sequences) are from unpublished papers, and another 39% (138 sequences) from a

single paper (Richards et al. 2008) (Table 4). Highlighting clearly the need for more study on the

subject to increase the number of potential sources and alleviate the possible bias of a "single

observer".

Phylogeny

While the construction of phylogenetic trees is a more refined and, by some, recommended form of

species identification, it is more demanding in terms of time and expertise, and has stages such as

sequence alignment and phylogenetic interpretation that often require significant manual

interpretation and intervention (Nixon and Wheeler 1990; Minh et al. 2013; Nguyen et al. 2015).

We observed high levels of polyphyly across all of our highest confidence matches, with only 1 of

evident case in which the 100% match was to a monophyletic species (*Acropora humilis* PaxC

region). Our preference is to employ a standardised, quickly repeatable, objective process, while

keeping in mind that the results are imperfect and designed to inform but not dictate identification

efforts.

Blast match

The use of query match with blast is far from being a complete solution, with 26 out of 76 samples

without a single species or species group resolved ID (Table 5), we have proposed a theoretical

decision tree that standardise the choice when there is a need of sorting through the output of a

multi-species name blast match (Fig. 2). In addition we used two different regions, a mitochondrial

one and a nuclear one to increase the confidence in our results (Suzuki et al. 2008). Our proposed

species ID are provisional and subject to revision when more species and sequences are available,

hence the value of a repeatable process for matching and re-matching.

All methods of identification that rely on DNA barcodes (e.g.: Blast matching, Barcoding Gap

analysis, etc.) use genetic divergence thresholds to assign individuals to correct species. This based

on two fundamental assumptions 1) monophyly of species with respect to the molecular marker

used, and 2) intraspecific genetic divergence is much smaller than interspecific genetic differences

(Toffoli et al. 2008). The first assumption is more dependent on the experimental choices made (i.e.

sequence region selected). The second assumption is where most of the criticism of barcodes

methods lies, on not comparing sister species or geographical distribution thus, underestimating

intraspecific genetic variability (Moritz and Cicero 2004; Toffoli et al. 2008). Fundamentally, the

issue of using thresholds lies in the fact that species naturally embodies an evolutionary process,

being subject to demographic and selective processes that will act on the genetic diversity (e.g.

(Avise et al. 2000; Hey 2001; Coyne and Orr 2004; Toffoli et al. 2008)). Since species are real

evolutionary groups and not categories which are created as a direct function of perceived

distinction (Hey 2001), the use of thresholds in "discovering" new species is overly simplistic, and

in some cases even misleading (Toffoli et al. 2008). We argue against simply substituting

morphology based taxonomy with DNA barcoding based taxonomy, as each system is more adapt

to answer different questions (Toffoli et al. 2008). Correct assignment is thus only possible by

complementing DNA barcoding with other data types, such as morphological, and ecological

characters as part of an integrated diagnostic approach (Toffoli et al. 2008).

While the simple blast match is inheritably less complex than other traditional DNA barcodes

identification methods (i.e.: the above mentioned barcoding gap analysis and, the potentially more

direct Automatic Barcoding Gap Discovery (ABGD) (Puillandre et al. 2012)). The argument we

make is that given the shared limitation and the common need for additional data in an integrated

approach, a faster and simpler blast match provides a relatively similar additional amount of

information for the species identification while being more easily repeatable, especially considering

the value of standardized data among aquariums (da Silva et al. 2019) and in light of the ever

increasing quantity of reference materials being added to GenBank and BOLD.

Available sequences libraries are not exhaustive, and probably never will be. That should not stop

us from attempting identifications based on the best available knowledge. All identifications must

fit within the context of our evolving understanding of taxonomy, where new descriptions

invalidate existing ones and often we do not have unambiguous taxonomic descriptions for every

species (Simpson 1951; Wiley 1978; Mayr 1982, 2000; Nixon and Wheeler 1990; Medlin et al.

1995). The reality is that barcoding as an independent tool of identification has its clear limitation

(Van Oppen et al. 1999; Shearer et al. 2002; Ferguson 2002; Moritz and Cicero 2004; Lee 2004;

DeSalle et al. 2005; Meyer and Paulay 2005; Prendini 2005; Knapp et al. 2005; Cognato 2006;

Hickerson et al. 2006; Meier et al. 2006, 2008; Little and Stevenson 2007; Kitahara et al. 2010), and

coral barcoding is no exception. With some of the concern on the use of genetic repertories in

common of all the methods explored in this study, we strongly recommend an integrated diagnostic

approach, combining morphological and genetic means of identification, in agreement with C.

Moritz in "DNA Barcoding: Promise and Pitfalls" (Moritz and Cicero 2004). Proposing the use of

barcoding and query matching as an additional tool for identification, increasing confidence of

experts and both confidence and capacity of non-experts' taxonomist. Producing this way reliably

identified sequences that can potentially become new reference, adding to the repertories already

freely available. Considering the specific needs of the aquarium community and their critical role to

support corals conservation (i.e., by hosting at least 24% of the corals assessed as highly vulnerable

to climate change (da Silva et al. 2019)), we emphasise the value of integrating standardised

barcoding analysis, and specimen morphology photographs to optimise usage of authoritative

identification guides and expert opinion.

Acknowledgments

Genetic analysis costs were partly covered by a FAITAG genetics fund grant an initiative from the

EAZA (European Association of Zoos and Aquaria) and EUAC (European Union of Aquarium

Curators) Fish and Aquatic Invertebrate Taxon Advisory Group (FAITAG).

Conflict of interest

On behalf of all authors, the corresponding author states that there is no conflict of interest.

Fig. 1: Number of records from member institutions of the Species360 network at genus or higher

level in light blue (i.e. order, family and genus) and at species or lower level in purple (i.e. species

and subspecies). This information is based on the rank field from the holdings information reported

by each institution through the ZIMS software. Information downloaded from ZIMS (Species360,

2020) on 4th June 2020.

Fig. 2: Decision tree for conflict solving based on Pairwise percentage identity, Bit score and

number of mismatches.

Fig. 3: Barcoding gap (raw distance matrix frequency distribution) – mtCR max distance = 0.0877

minimum distance = 0 ; PaxC max distance = 0.5895 minimum distance = 0. (Bin size=0.002) A)

Minimum interspecific distance and maximum intraspecific distance B) all distances

Fig. 4: Number of mtCR and PaxC references sequences with the same species name by species

group

Bibliography

Afgan E, Baker D, Batut B, Van Den Beek M, Bouvier D, Ech M, Chilton J, Clements D, Coraor N, Grüning BA, Guerler A, Hillman-Jackson J, Hiltemann S, Jalili V, Rasche H, Soranzo N, Goecks J, Taylor J, Nekrutenko A, Blankenberg D (2018) The Galaxy platform for accessible, reproducible and collaborative biomedical analyses: 2018 update. Nucleic Acids Res 46:W537–W544

Amaral FD (1994) Morphological variation in the reef coral Montastrea cavernosa in Brazil. Coral Reefs 13:113–117

Avise JC, Baker RJ, Porter CA, Patton JC, Van Den Bussche RA, Ballard JW, Chemoff B, James AC (2000) Phylogeography: the history and formation of species. Harvard university press,

Ayre DJ, Willis BL (1988) Population structure in the coral Pavona cactus: clonal genotypes show little phenotypic plasticity. Mar Biol 99:495–505

Babcock RC, Bull GD, Harrison PL, Heyward AJ, Oliver JK, Wallace CC, Willis BL (1986) Synchronous spawnings of 105 scleractinian coral species on the Great Barrier Reef. Mar Biol 90:379–394

Bellwood DR, Hughes TP (2001) Regional-scale assembly rules and biodiversity of coral reefs. Science (80-) 292:1532–1534

Benson DA (2000) GenBank. Nucleic Acids Res 28:15–18

Best, Borel M, G. J. Boekschoten, Oosterbaan A (1984) Species concept and ecomorph variation in living and fossil Scleractinia. Palaeontogr Am 54 70–79

Blanco JM, Wildt DE, Höfle U, Voelker W, Donoghue AM (2009) Implementing artificial insemination as an effective tool for ex situ conservation of endangered avian species. Theriogenology 71:200–213

Borneman E (2008) Introduction to the husbandry of corals in aquariums : A review. Adv coral Husb public aquariums, Leewis, RJ Janse, M 2:3–14

Brondizio ES, Settele J, Dìaz S, Others (2019) IPBES. 2019 Global assessment report on biodiversity and ecosystem services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services. Bonn, Ger

Brunner B (2012) The Ocean at Home: an illustrated history of the aquarium. Reaktion Books,

Bruno JF, Edmunds PJ (1997) Clonal variation for phenotypic plasticity in the coral Madracis mirabilis. Ecology 78:2177–2190

Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL (2009) BLAST+: Architecture and applications. BMC Bioinformatics 10:1–9

Chen I-P, Tang C-Y, Chiou C-Y, Hsu J-H, Wei NV, Wallace CC, Muir P, Wu H, Chen CA (2009) Comparative Analyses of Coding and Noncoding DNA Regions Indicate that Acropora (Anthozoa: Scleractina) Possesses a Similar Evolutionary Tempo of Nuclear vs. Mitochondrial Genomes as in Plants. Mar Biotechnol 11:141–152

Cognato AI (2006) Standard Percent DNA Sequence Difference for Insects Does Not Predict Species Boundaries. J Econ Entomol 99:1037–1045

Cowburn B, Samoilys MA, Obura D (2018) The current status of coral reefs and their vulnerability to climate change and multiple human stresses in the Comoros Archipelago, Western Indian Ocean. Mar Pollut Bull 133:956–969

Cowman PF, Quattrini AM, Bridge TCL, Watkins-Colwell GJ, Fadli N, Grinblat M, Roberts TE,
McFadden CS, Miller DJ, Baird AH (2020) An enhanced target-enrichment bait set for
Hexacorallia provides phylogenomic resolution of the staghorn corals (Acroporidae) and close
relatives. Mol Phylogenet Evol 153:106944

Coyne JA, Orr HA (2004) Speciation Sinauer Associates. Sunderland, MA 276:281

DeSalle R, Egan MG, Siddall M (2005) The unholy trinity: Taxonomy, species delimitation and
DNA barcoding. Philos Trans R Soc B Biol Sci 360:1905–1916

Fa JE, Funk SM, O'Connell D (2011) Zoo conservation biology. Cambridge University Press,

Ferguson JWH (2002) On the use of genetic divergence for identifying species. Biol J Linn Soc
75:509–516

Fisher R, O'Leary RA, Low-Choy S, Mengersen K, Knowlton N, Brainard RE, Caley MJ (2015)
Species richness on coral reefs and the pursuit of convergent global estimates. Curr Biol
25:500–505

Foster AB (1979) Phenotypic plasticity in the reef corals Montastraea annularis (Ellis & Solander)
and Siderastrea siderea (Ellis & Solander). J Exp Mar Bio Ecol 39:25–54

Fukami H, Omori M, Hatta M (2000) Phylogenetic Relationships in the Coral Family Acroporidae,
Reassessed by Inference from Mitochondrial Genes. Zoolog Sci 17:689–696

Godfray HCJ (2002) Challenges for taxonomy. Nature 417:17–19

Hahsler M, Nagar A (2019) rBLAST:Interface to the Basic Local Alignment Search Tool (BLAST).

Harrison PL, Babcock RC, Bull GD, Oliver JK, Wallace CC, Willis BL (1984) Mass Spawning in
Tropical Reef Corals. Science (80-) 223:1186–1189

Hatta M, Fukami H, Wang W, Omori M, Shimoike K, Hayashibara T, Ina Y, Sugiyama T (1999)
Reproductive and genetic evidence for a reticulate evolutionary history of mass-spawning
corals. Mol Biol Evol 16:1607–1613

Hebert PDN, Cywinska A, Ball SL, DeWaard JR (2003) Biological identifications through DNA
barcodes. Proc R Soc B Biol Sci 270:313–321

Hellberg ME (2006) No variation and low synonymous substitution rates in coral mtDNA despite
high nuclear variation. BMC Evol Biol 6:1–8

Hey J (2001) Genes, categories, and species: The evolutionary and cognitive cause of the species
problem. Oxford University Press,

Hickerson MJ, Meyer CP, Moritz C (2006) DNA barcoding will often fail to discover new animal
species over broad parameter space. Syst Biol 55:729–739

Hoegh-Guldberg O, Poloczanska ES, Skirving W, Dove S (2017) Coral Reef Ecosystems under
Climate Change and Ocean Acidification. Front Mar Sci 4:158

Horton T, Kroh A, Ahyong S, Bailly N, Boyko CB, Brandão SN, Gofas S, Hooper JNA, Hernandez
F, Holovachov O, Mees J, Molodtsova TN, Paulay G, Decock W, Dekeyzer S, Lanssens T,
Vandepitte L, Vanhoorne B, Adlard R, Adriaens P, Agatha S, Ahn KJ, Akkari N, Alvarez B,
Anderberg A, Anderson G, Angel M V, Antic D, Arango C, Artois T, Atkinson S, Baldwin
BG, Bank R, Barber A, Barbosa JP, Bartsch I, Bellan-Santini D, Bergh N, Bernot J, Berta A,
Bezerra TN, Bieler R, Blanco S, Blasco-Costa I, Blazewicz M, Bock P, de León M, Böttger-
Schnack R, Bouchet P, Boury-Esnault N, Boxshall G, Bray R, Bruce NL, Cairns S, Calvo
Casas J, Carballo JL, Cárdenas P, Carstens E, Chan BK, Chan TY, Cheng L, Churchill M,
Coleman CO, Collins AG, Collins GE, Corbari L, Cordeiro R, Cornils A, Coste M, Costello
MJ, Crandall KA, Cremonte F, Cribb T, Cutmore S, Dahdouh-Guebas F, Daly M, Daneliya M,

Dauvin JC, Davie P, De Broyer C, De Grave S, de Mazancourt V, de Voogd NJ, Decker P, Decraemer W, Defaye D, d'Hondt JL, Dippenaar S, Dohrmann M, Dolan J, Domning D, Downey R, Ector L, Eisendle-Flöckner U, Eitel M, Encarnação SC d., Enghoff H, Epler J, Ewers-Saucedo C, Faber M, Feist S, Figueroa D, Finn J, Fišer C, Fordyce E, Foster W, Frank JH, Fransen C, Freire S, Furuya H, Galea H, Gao T, Garcia-Alvarez O, Garcia-Jacas N, Garic R, Garnett S, Gasca R, Gaviria-Melo S, Gerken S, Gibson D, Gibson R, Gil J, Gittenberger A, Glasby C, Glover A, Gómez-Noguera SE, González-Solís D, Gordon D, Gostel M, Grabowski M, Gravili C, Guerra-García JM, Guidetti R, Guiry MD, Gutierrez D, Hadfield KA, Hajdu E, Hallermann J, Hayward BW, Heiden G, Hendrycks E, Herbert D, Herrera Bachiller A, Ho J s., Hodda M, Høeg J, Hoeksema B, Houart R, Hughes L, Hyžný M, Iniesta LFM, Iseto T, Ivanenko S, Iwataki M, Janssen R, Jarms G, Jaume D, Jazdzewski K, Jersabek CD, Jóźwiak P, Kabat A, Kantor Y, Karanovic I, Karthick B, Katinas L, Kim YH, King R, Kirk PM, Klautau M, Kociolek JP, Köhler F, Kolb J, Kotov A, Kremenetskaia A, Kristensen RM, Kulikovskiy M, Kullander S, Lambert G, Lazarus D, Le Coze F, LeCroy S, Leduc D, Lefkowitz EJ, Lemaitre R, Lichter-Marck IH, Liu Y, Loeuille B, Lörz AN, Lowry J, Ludwig T, Lundholm N, Macpherson E, Madin L, Mah C, Mamo B, Mamos T, Manconi R, Mapstone G, Marek PE, Marshall B, Marshall DJ, Martin P, Mast R, McFadden C, McInnes SJ, Meidla T, Meland K, Merrin KL, Messing C, Miljutin D, Mills C, Moestrup Ø, Mokievsky V, Monniot F, Mooi R, Morandini AC, da Rocha R, Morrow C, Mortelmans J, Mortimer J, Musco L, Nesom G, Neubauer TA, Neubert E, Neuhaus B, Ng P, Nguyen AD, Nielsen C, Nishikawa T, Norenburg J, O'Hara T, Opresko D, Osawa M, Osigus HJ, Ota Y, Páll-Gergely B, Panero JL, Pasini E, Patterson D, Paxton H, Pelser P, Peña-Santiago R, Perrier V, Perrin W, Petrescu I, Picton B, Pilger JF, Pisera AB, Polhemus D, Poore GC, Potapova M, Pugh P, Read G, Reich M, Reimer JD, Reip H, Reuscher M, Reynolds JW, Richling I, Rimet F, Ríos P, Rius M, Rogers DC, Roque N, Rosenberg G, Rützler K, Sabbe K, Saiz-Salinas J, Sala S, Santagata S, Santos S, Sar E, Satoh A, Saucède T, Schatz H, Schierwater B, Schilling E, Schmidt-Rhaesa A, Schneider S, Schönberg C, Schuchert P, Senna AR, Serejo C, Shaik S, Shamsi S, Sharma J, Shear WA, Shenkar N, Shinn A, Short M, Sicinski J, Sierwald P, Simmons E, Sinniger F, Sivell D, Sket B, Smit H, Smit N, Smol N, Souza-Filho JF, Spelda J, Sterrer W, Stienen E, Stoev P, Stöhr S, Strand M, Suárez-Morales E, Summers M, Suppan L, Susanna A, Suttle C, Swalla BJ, Taiti S, Tanaka M, Tandberg AH, Tang D, Tasker M, Taylor J, Taylor J, Tchesunov A, ten Hove H, ter Poorten JJ, Thomas JD, Thuesen E V, Thurston M, Thuy B, Timi JT, Timm T, Todaro A, Turon X, Tyler S, Uetz P, Uribe-Palomino J, Urtubey E, Utevsky S, Vacelet J, Vachard D, Vader W, Väinölä R, Valls Domedel G, de Vijver B, van der Meij SE, van Haaren T, van Soest RW, Vanreusel A, Venekey V, Vinarski M, Vonk R, Vos C, Walker-Smith G, Walter TC, Watling L, Wayland M, Wesener T, Wetzel CE, Whipps C, White K, Wieneke U, Williams DM, Williams G, Wilson R, Witkowski A, Witkowski J, Wyatt N, Wylezich C, Xu K, Zanol J, Zeidler W, Zhao Z (2020) World Register of Marine Species (WoRMS).

Huang D, Meier R, Todd PA, Chou LM (2008) Slow Mitochondrial COI Sequence Evolution at the Base of the Metazoan Tree and Its Implications for DNA Barcoding. J Mol Evol 66:167–174

Hughes TP, Connolly SR, Eakin CM, Dietzel A, McWilliam MJ, Hoey AS, Pears RJ, Liu G, Skirving WJ, Heron SF, Pratchett MS, Baird AH, Stella JS, Hoogenboom MO, Torda G, Kerry JT (2018) Global warming transforms coral reef assemblages. Nature 556:492–496

Hutchins M (2003) Zoo and aquarium animal management and conservation: current trends and future challenges. Int Zoo Yearb 38:14–28

Isomura N, Iwao K, Fukami H (2013) Possible Natural Hybridization of Two Morphologically Distinct Species of Acropora (Cnidaria, Scleractinia) in the Pacific: Fertilization and Larval Survival Rates. PLoS One 8:1–8

James Harris D (2003) Can you bank on GenBank? Trends Ecol Evol 18:317–319

Jombart T, Kamvar ZN, Schliep K, Archer E, Harris R (2017) apex: Phylogenetic Methods for Multiple Gene Data.

Kitahara M V., Cairns SD, Stolarski J, Blair D, Miller DJ (2010) A comprehensive phylogenetic analysis of the scleractinia (Cnidaria, Anthozoa) based on mitochondrial CO1 sequence data. PLoS One 5:

Knapp S, Lughadha EN, Paton A, Mallet J, Isaac NJB, MacE GM, James Harris D, Froufe E (2005) Taxonomic inflation, species concepts and global species lists (multiple letters). Trends Ecol Evol 20:7–8

Lacy RC (2013) Achieving True Sustainability of Zoo Populations. Zoo Biol 32:19–26

Lee MSY (2004) The molecularisation of taxonomy. Invertebr Syst 18:1–6

Leray M, Knowlton N, Ho SL, Nguyen BN, Machida RJ (2019) GenBank is a reliable resource for 21st century biodiversity research. Proc Natl Acad Sci U S A 116:22651–22656

Leus K, Traylor-Holzer K, Lacy RC (2011) Genetic and demographic population management in zoos and aquariums: Recent developments, future challenges and opportunities for scientific research. Int Zoo Yearb 45:213–225

Little DP, Stevenson DW (2007) A comparison of algorithms for the identification of specimens using DNA barcodes: Examples from gymnosperms. Cladistics 23:1–21

Ma B, Liao L, Peng Q, Fang T, Zhou H, Korban SS, Han Y (2017) Reduced representation genome sequencing reveals patterns of genetic diversity and selection in apple. J Integr Plant Biol 59:190–204

Márquez LM, Van Oppen MJHH, Willis BL, Reyes A, Miller DJ (2002) The highly cross-fertile coral species, Acropora hyacinthus and Acropora cytherea, constitute statistically distinguishable lineages. Mol Ecol 11:1339–1349

Masson-Delmotte V, Zhai P, Pörtner H-O, Roberts D, Skea J, Shukla PR, Pirani A, Moufouma-Okia W, Péan C, Pidcock R, Connors S, Matthews JBR, Chen Y, Zhou X, Gomis MI, Lonnoy E, Maycock T, Tignor M, Waterfield T (2018) IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change, sustainable development, an. Sustain Dev Efforts to Eradicate Poverty https//www ipcc.ch/sr15/ 1:

Mayr E (1982) The growth of biological thought: Diversity, evolution, and inheritance. Harvard University Press.,

Mayr E (2000) The biological species concept. Species concepts phylogenetic theory a debate Columbia Univ Press New York 17–29

Medlin LK, Lange M, Barker GLA, Hayes PK (1995) Can Molecular Techniques Change Our Ideas About the Species Concept? Molecular Ecology of Aquatic Microbes. Springer Berlin Heidelberg, Berlin, Heidelberg, pp 133–152

Meier R, Shiyang K, Vaidya G, Ng PKL (2006) DNA barcoding and taxonomy in diptera: A tale of high intraspecific variability and low identification success. Syst Biol 55:715–728

Meier R, Zhang G, Ali F (2008) The use of mean instead of smallest interspecific distances exaggerates the size of the "barcoding gap" and leads to misidentification. Syst Biol 57:809–813

Meyer CP, Paulay G (2005) DNA barcoding: Error rates based on comprehensive sampling. PLoS Biol 3:e422

Miller KJ (1994) Morphological variation in the coral genus Platygyra - Environmental influences and taxonomic implications. Mar Ecol Prog Ser 110:19–28

Minh BQ, Nguyen MAT, von Haeseler A (2013) Ultrafast Approximation for Phylogenetic Bootstrap. Mol Biol Evol 30:1188–1195

Mora C, Tittensor DP, Adl S, Simpson AGB, Worm B (2011) How many species are there on earth and in the ocean? PLoS Biol 9:1–8

Mora C, Tittensor DP, Myers RA (2008) The completeness of taxonomic inventories for describing the global diversity and distribution of marine fishes. Proc R Soc B Biol Sci 275:149–155

Moritz C, Cicero C (2004) DNA Barcoding: Promise and Pitfalls. PLoS Biol 2:e354

Muko S, Kawasaki K, Sakai K, Takasu F, Shigesada N (2000) Morphological plasticity in the coral Porites sillimaniani and its adaptive significance. Bull Mar Sci 66:225–239

Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ (2015) IQ-TREE: A Fast and Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. Mol Biol Evol 32:268–274

Nixon KC, Wheeler QD (1990) An amplification of the phylogenetic species concept. Cladistics 6:211–223

van Oppen MJH, Hislop NR, Hagerman PJ, Miller DJ (1999) Gene Content and Organization in a Segment of the Mitochondrial Genome of the Anthozoan Subclass Zoantharia. Mol Biol Evol 16:1812–1815

Van Oppen MJH, Willis BL, Van Rheede T, Miller DJ (2002) Spawning times, reproductive compatibilities and genetic structuring in the Acropora aspera group: Evidence for natural hybridization and semi-permeable species boundaries in corals. Mol Ecol 11:1363–1376

Van Oppen MJHH, McDonald BJ, Willis B, Miller DJ (2001) The evolutionary history of the coral genus acropora (Scleractinia, Cnidaria) based on a mitochondrial and a nuclear marker: Reticulation, incomplete lineage sorting, or morphological convergence? Mol Biol Evol 18:1315–1329

Van Oppen MJHH, Willis BL, Miller DJ (1999) Atypically low rate of cytochrome b evolution in the scleractinian coral genus Acropora. Proc R Soc London Ser B Biol Sci 266:179–183

Ortman BD, Bucklin A, Pagès F, Youngbluth M (2010) DNA Barcoding the Medusozoa using mtCOI. Deep Sea Res Part II Top Stud Oceanogr 57:2148–2156

Palumbi SR, Vollmer S, Romano S, Oliver T, Ladner J (2012) The role of genes in understanding the evolutionary ecology of reef building corals. Evol Ecol 26:317–335

Pratchett MS, Pisapia C, Sheppard CRC (2013) Background mortality rates for recovering populations of Acropora cytherea in the Chagos Archipelago, central Indian Ocean. Mar Environ Res 86:29–34

Prendini L (2005) Comment on "Identifying spiders through DNA barcodes." Can J Zool 83:498–504

Puillandre N, Lambert A, Brouillet S, Achaz G (2012) ABGD, Automatic Barcode Gap Discovery for primary species delimitation. Mol Ecol 21:1864–1877

Raven PH (2004) Ex situ plant conservation: supporting species survival in the wild. Island Press,

Richards ZT, van Oppen MJH, Wallace CC, Willis BL, Miller DJ (2008) Some rare Indo-Pacific coral species are probable hybrids. PLoS One 3:1–8

Rinkevich B (1995) Restoration Strategies for Coral Reefs Damaged by Recreational Activities:

691   The Use of Sexual and Asexual Recruits. Restor Ecol 3:241–251

692   Rozenfeld AF, Arnaud-Haond S, Hernández-García E, Eguíluz VM, Matías MA, Serrão E, Duarte
693   CM (2007) Spectrum of genetic diversity and networks of clonal organisms. J R Soc Interface
694   4:1093–1102

695   Shearer TL, Coffroth MA (2008) DNA BARCODING: Barcoding corals: limited by interspecific
696   divergence, not intraspecific variation. Mol Ecol Resour 8:247–255

697   Shearer TL, van Oppen MJH, Romano SL, Wörheide G (2002) Slow mitochondrial DNA sequence
698   evolution in the Anthozoa (Cnidaria). Mol Ecol 11:2475–2487

699   Sheppard C, Sheppard A, Fenner D (2020) Coral mass mortalities in the Chagos Archipelago over
700   40 years: Regional species and assemblage extinctions and indications of positive feedbacks.
701   Mar Pollut Bull 154:111075

702   da Silva R, Pearce-Kelly P, Zimmerman B, Knott M, Foden W, Conde DA (2019) Assessing the
703   conservation potential of fish and corals in aquariums globally. J Nat Conserv 48:1–11

704   Simpson GG (1951) The Species Concept. Evolution (N Y) 5:285

705   Skibins JC, Powell RB (2013) Conservation caring: Measuring the influence of zoo visitors'
706   connection to wildlife on pro-conservation behaviors. Zoo Biol 32:528–540

707   Suzuki G, Hayashibara T, Shirayama Y, Fukami H (2008) Evidence of species-specific habitat
708   selectivity of Acropora corals based on identification of new recruits by two molecular
709   markers . Mar Ecol Prog Ser 355:149–159

710   Todd PA (2008) Morphological plasticity in scleractinian corals. Biol Rev 83:315–337

711   Todd PA, Sidle RC, Lewin-Koh NJI (2004) An aquarium experiment for identifying the physical
712   factors inducing morphological change in two massive scleractinian corals. J Exp Mar Bio
713   Ecol 299:97–113

714   Toffoli D, Hrbek T, de Araújo MLG, de Almeida MP, Charvet-Almeida P, Farias IP (2008) A test
715   of the utility of DNA barcoding in the radiation of the freshwater stingray genus Potamotrygon
716   (Potamotrygonidae, Myliobatiformes). Genet Mol Biol 31:324–336

717   Turley SK (1999) Conservation and tourism in the traditional UK zoo. J Tour Stud 10:2

718   Vellend M, Geber MA (2005) Connections between species diversity and genetic diversity. Ecol
719   Lett 8:767–781

720   Veron JEN, Stafford-Smith MG, Turak E, L.M. D (2020) Corals of the World.
721   http://www.coralsoftheworld.com

722   Vollmer S V. (2002) Hybridization and the Evolution of Reef Coral Diversity. Science (80-)
723   296:2023–2025

724   Vollmer S V, Palumbi SR (2002) Hybridization and the Evolution of Reef Coral Diversity. Science
725   (80-) 296:2023 LP – 2025

726   Wallace CC (1999) Staghorn corals of the world: a revision of the coral genus Acropora
727   (Scleractinia; Astrocoeniina; Acroporidae) worldwide, with emphasis on morphology,
728   phylogeny and biogeography. CSIRO publishing,

729   Wallace CC, Done BJ, Muir PR (2012) Revision and catalogue of worldwide staghorn corals
730   acropora and isopora (scleractinia: Acroporidae) in the museum of tropical Queensland. Mem
731   Queensl Museum 57:1–257

732   Wallace CC, Willis BL (1994) Systematics of the coral genus Acropora: implications of new
733   biological findings for species concepts. Annu Rev Ecol Syst 25:237–262

Wei NV, Hsieh HJ, Dai C, Wallace CC, Baird AH, Chen CA (2012) Acroporidae) in a Marginal
Coral Assemblage. Mar Biol Res 51:85–92

Wiley EO (1978) The Evolutionary Species Concept Reconsidered. Syst Zool 27:17

Willis BL (1985) Phenotypic plasticity versus phenotypic stability in the reef corals Turbinaria
mesenterina and Pavona cactus. Proc Fifth Int Coral Reef Symp Vol. 4:

Willis BL (1997) Hybridization and breeding incompatibilities within the mating systems of mass
spawning reef corals. 1:81–90

Willis BL, van Oppen MJH, Miller DJ, Vollmer S V., Ayre DJ (2006) The Role of Hybridization in
the Evolution of Reef Corals. Annu Rev Ecol Evol Syst 37:489–517

Winter DJ (2017) rentrez: An R package for the NCBI eUtils API. R J 9:520–526

Wolstenholme JK (2004) Temporal reproductive isolation and gametic compatibility are
evolutionary mechanisms in the Acropora humilis species group (Cnidaria; Scleractinia). Mar
Biol 144:567–582

(2020) idCoral.org. https://idcoral.org

Figure 1: Number of records from member institutions of the Species360 network at genus or higher level in light blue(i.e. order, family and genus) and at species or lower level in purple (i.e. species and subspecies). This information is based on the rank field from the holdings information reported by each institution through the ZIMS software. Information downloaded from ZIMS (Species360, 2020) on 4th June 2020.

Figure 2: Decision tree for conflict solving based on Pairwise percentage identity, Bit score and number of mismatches.

Figure 3: Barcoding gap (raw distance matrix frequency distribution) – mtCR max distance = 0.0877 minimum distance = 0 ; PaxC max distance = 0.5895 minimum distance = 0. (Bin size=0.002) A) Minimum interspecific distance and maximum intraspecific distance B) all distances

Figure 4: Number of mtCR and PaxC references sequences with the same species name by species group

Table 1: Success rate across genes

N samples	N samples overlapping	Gene	N Sequences	Minimum sequence length	Maximum sequence length	Median sequence length
127	76	*mtCR*	84	563	904	734
		PaxC	90	429	869	649

Table 2: Number of sample matching to a single or more (2+) species name on GenBank. Above the diagonal line match that agree between the two region. Below the diagonal line matches that disagree between the two regions

Table 3:Sources of mtCR references sequences by Paper title

Journal	Authors	Title	N	%
Doctroal Thesis (2003) James Cook University, Townsville, Queensland, Australia	Wolstenholme JK	Species boundaries in scleractinian corals: a case study of the Acropora humilis species group	242	41.8
PLoS ONE 3 (9), E3240 (2008)	Z. T. Richards, M. J. H. van Oppen, C. C. Wallace, B. L. Willis, and D. J. Miller	Some rare Indo-Pacific coral species are probable hybrids	79	13.6
Conserv. Genet. 18 (4) 825–835 (2017)	M. D. Waterhouse, C. Blair, K. W. Larsen, and M. A. Russello	Genetic Variation and Fine-Scale Population Structure in the Threatened Acropora palmata and Acropora cervicornis around Puerto Rico	48	8.3
Science 296 (5575), 2023-2025 (2002)	S. V. Vollmer	Hybridization and the evolution of reef coral diversity	42	7.3
Mol. Biol. Evol. 18 (7), 1315-1329 (2001)	M. J. H. van Oppen, B. J. McDonald, B. Willis, and D. J. Miller	The evolutionary history of the coral genus Acropora (Scleractinia, Cnidaria) based on a mitochondrial and a nuclear marker: reticulation, incomplete lineage sorting, or morphological convergence?	41	7.1
PLoS One 5 (1) e8652 (2010)	E. M. Hemond and S. V Vollmer	Genetic Diversity and Connectivity in the Threatened Staghorn Coral (Acropora cervicornis) in Florida	32	5.5
Mol. Ecol. 11 (8), 1339-1349 (2002)	L. M. Márquez, M. J. H. Van Oppen, B. L. Willis, A. Reyes, and D. J. Miller	The highly cross-fertile coral species, Acropora hyacinthus and Acropora cytherea, constitute statistically distinguishable lineages	21	3.6
Zool. Sci. 29 (2), 134-140 (2012)	G. Suzuki and H. Fukami	Evidence of genetic and reproductive isolation between two morphs of subtropical-dominant coral Acropora solitaryensis in the non-reef region of Japan	13	2.2
Mol. Ecol. 24 (19), 5006-5019 (2015)	N. L. Rosser	Asynchronous spawning in sympatric populations of a hard coral reveals cryptic species and ancient genetic lineages	8	1.4
PeerJ 7, e6429 (2019)	H. Fukami, K. Iwao, N. H. Kumagai, M. Morita, and N. Isomura	Maternal inheritance of F1 hybrid morphology and colony shape in the coral genus Acropora	6	1.0
Syst. Biodivers. 8 (2), 281–288 (2010)	Z. T. Richards, C. C. Wallace, and D. J. Miller	Archetypal 'elkhorn' coral discovered in the Pacific Ocean	6	1.0
Mol. Phylogenet. Evol. 63 (2), 527-531 (2012)	Y. Nakajima, A. Nishikawa, A. Iguchi, and K. Sakai	The population genetic approach delineates the species boundary of reproductively isolated corymbose acroporid corals	4	0.7
Mar. Ecol. Prog. Ser. 355, 149–159 (2008)	G. Suzuki, T. Hayashibara, Y. Shirayama, and H. Fukami	Evidence of species-specific habitat selectivity of Acropora corals based on identification of new recruits by two molecular markers	3	0.5
J. Mol. Evol. 55 (1), 1-13 (2002)	M. J. H. Van Oppen, J. Catmull, B. J. McDonald, N. R. Hislop, P. J. Hagerman, and D. J. Miller	The mitochondrial genome of Acropora tenuis (Cnidaria; Scleractinia) contains a large group I intron and a candidate control region	1	0.2
Mitochondrial DNA, 1-2 (2015) In press	Y. Zhang, X. Yu, Z. Zhou, and B. Huang	The complete mitochondrial genome of Acropora aculeus (cnidaria, scleractinia, acroporidae)	1	0.2
Proc. R. Soc. Lond., B, Biol. Sci. 266 (1415), 179-183 (1999)	M. J. H. Van Oppen, B. L. Willis, and D. J. Miller	Atypically low rate of cytochrome b evolution in the scleractinian coral genus Acropora	1	0.2
Unpublished	Santacruz-Castro,A. and Dai,C.F.	Latitudinal cline in the reproductive traits and local adaptation of the Acropora hyacinthus species complex	20	3.5
Unpublished	Chan,C.L. and Chen,C.A.	Multiplex next generation sequencing of scleractinian mitochondrial genomes	11	1.9

Table 4:Sources of PaxC references sequences by Paper title

Journal	Authors	Title	N	%
PLoS ONE 3 (9), E3240 (2008)	Z. T. Richards, M. J. H. van Oppen, C. C. Wallace, B. L. Willis, and D. J. Miller	Some rare Indo-Pacific coral species are probable hybrids	138	39.2
Mol. Biol. Evol. 18 (7), 1315-1329 (2001)	M. J. H. van Oppen, B. J. McDonald, B. Willis, and D. J. Miller	The evolutionary history of the coral genus Acropora (Scleractinia, Cnidaria) based on a mitochondrial and a nuclear marker: reticulation, incomplete lineage sorting, or morphological convergence?	87	24.7
Mol. Ecol. 11 (8), 1339-1349 (2002)	L. M. Márquez, M. J. H. Van Oppen, B. L. Willis, A. Reyes, and D. J. Miller	The highly cross-fertile coral species, Acropora hyacinthus and Acropora cytherea, constitute statistically distinguishable lineages	34	9.7
Mol. Ecol. 24 (19), 5006-5019 (2015)	N. L. Rosser	Asynchronous spawning in sympatric populations of a hard coral reveals cryptic species and ancient genetic lineages	16	4.5
Syst. Biodivers. 8 (2), 281–288 (2010)	Z. T. Richards, C. C. Wallace, and D. J. Miller	Archetypal 'elkhorn' coral discovered in the Pacific Ocean	10	2.8
Evol. Ecol. (2011) In press	S. R. Palumbi, S. Vollmer, S. Romano, T. Oliver, and J. Ladner	The role of genes in understanding the evolutionary ecology of reef building corals	3	0.9
Galaxea, J. Coral Reef Stud. 10 (2), 91 (2008)	M. Hatta and K. Matsushima	Presumed natural hybrids between Acropora donei and Acropora tenuis	3	0.9
Mol. Ecol. 9 (9), 1363-1373 (2000)	M. J. H. Van Oppen, B. L. Willis, H. W. J. A. Van Vugt, and D. J. Miller	Examination of species boundaries in the Acropora cervicornis group (Scleractinia, cnidaria) using nuclear DNA sequence analyses	3	0.9
Unpublished	Wei,N.V., Tang,P.-C. and Chen,A.C.	Hybridization does not occur among common Acropora species in a marginal coral assemblage	29	8.2
Unpublished	Ohki,S., Kowalski,R.K., Kitanobou,S. and Morita,M.	Different timing of the spawning is related to reproductive isolation and gamete species recognition in the broadcast spawning coral Acropora	27	7.7
Unpublished	van Oppen,M.J.H. and Miller,D.J.	A new single-locus nuclear DNA marker for evolutionary studies in the scleractinian coral genus Acropora	2	0.6