Toward a Monte Carlo approach
to selecting climate variables in MaxEnt:
A case study using Cassin's Sparrow (*Peucaea cassinii*)

John L. Schnase[*], Mark L. Carroll, Roger L. Gill, Glenn S. Tamkin,

Jian Li, Savannah L. Strong, Thomas P. Maxwell, and Mary E. Aronne

Office of Computational and Information Sciences and Technology,

NASA Goddard Space Flight Center, Greenbelt, Maryland, 20708, USA

*Corresponding author

Email: john.l.schnase@nasa.gov

Author contributions

Abstract

MaxEnt is an important aid in understanding the influence of climate change on species distributions and abundance. There is growing interest in using IPCC-class global climate model outputs as environmental predictors in this work. These models provide realistic, global representations of the climate system, projections for hundreds of variables (including Essential Climate Variables), and combine observations from an array of satellite, airborne, and *in-situ* sensors. Unfortunately, direct use of this important class of data in MaxEnt modeling has been limited by the large size of climate model output collections and the fact that MaxEnt can only operate on a relatively small set of predictors stored in a computer's main memory. In this study, we demonstrate the feasibility of a Monte Carlo method that overcomes this limitation by finding a useful subset of predictors in a larger, externally-stored collection of environmental variables in a reasonable amount of time. Our proposed solution takes an ensemble approach wherein many MaxEnt runs, each drawing on a small random subset of variables, converges on a global estimate of the top contributing subset of variables in the larger collection. In preliminary tests, the Monte Carlo approach selected a consistent set of top six variables within 540 runs, with the four most contributory variables of the top six accounting for approximately 93% of overall permutation importance in the final model. These results suggest that a Monte Carlo approach could offer a viable means of selecting environmental predictors for MaxEnt models that is amenable to parallelization and scalable to very large data sets. This point to the possibility of near-real-time multiprocessor implementations that could enable broader and more exploratory use of global climate model outputs in environmental niche modeling and aid in the discovery of viable predictors.

Introduction

MaxEnt is one of the most popular software packages in use today by the ecological research community [1–3]. Based on a machine learning approach to maximum entropy modeling, MaxEnt allows researchers to construct ecological niche models (ENMs) that estimate the habitat suitability of a species using occurrence data and a set of environmental variables [1,2,4–6]. An abundant literature points to MaxEnt's effectiveness across a wide range of applications in fields as diverse as biogeography and phylogeny [7], conservation biology and epidemiology [8,9], invasion biology [10–12], and archaeology [13]. Its merits compared to alternative approaches have been the subject of numerous statistical and methodological analyses, many of which have led to software improvements and refinements to the way MaxEnt is used [14–21]. In this paper, we contribute to this ongoing dialog by describing our efforts to overcome a specific technical limitation of the MaxEnt software that makes the tool difficult to use with large predictor data sets.

In recent years, MaxEnt has become a particularly important aid in understanding the influence of climate change on species distributions and abundance [22–27]. The need for reliable climate projections in this work is leading to greater use of global climate model (GCM) outputs as predictors [23]. While creating important new opportunities for research, this trend is also creating a "Big Data" challenge for the MaxEnt community [16]. The largest and most sophisticated GCMs — sometimes referred to as "IPCC-class" models because of the critical role they play in the work of the Intergovernmental Panel on Climate Change (IPCC) — produce petabyte-scale data sets comprising hundreds of variables, a volume that vastly exceeds what is generally used in bioclimatic modeling today [28–30]. Moreover, the direct outputs of these systems are being transformed into derived climate data

products on an unprecedented scale [24,31]. As a result, model tuning and variable selection,

which are crucial aspects of any species distribution modeling effort, are becoming more

complicated issues [32].

Part of the problem lies in the fact that MaxEnt, like many machine learning systems,

acts on its inputs as a piece: predictors and observations must be memory-resident for the

program to work [33]. This results in run-times and space requirements that scale linearly

with the size of a model's inputs. In most cases, these scaling properties pose few difficulties.

But when the number of predictors under consideration becomes large, compute times can

become impractically long, models can become overly complex, and efforts to understand

any particular variable's contribution to model formation, either as an aspect of model

analysis or as a way of selecting subsets of variables for further model refinement, can

become challenging [17,32,34–36]. Clearly, an effective way of dealing with large

environmental data sets that preserves the many advantages of MaxEnt while overcoming its

current limitations would benefit the MaxEnt community.

In this study, we investigated the potential of a Monte Carlo method to help

accomplish such an outcome. Monte Carlo optimizations are a common way of finding

approximate answers to problems that are solvable in principle but lack a practical means of

solution [37]. Our objective was to find a useful subset of predictors in a larger collection of

environmental variables in a reasonable amount of time. Our proposed solution takes an

ensemble approach wherein many MaxEnt runs, each drawing on a small random subset of

variables, converges on a global estimate of the top contributing subset of variables in the

larger collection.

Preliminary results suggest that the method reliably selects a subset of the original

predictors that is capable of producing a well-tuned, parsimonious model of high quality.

Since each model run is independent and uses a set number of variables, the method is totally

parallelizable, independent of the scaling properties of MaxEnt, and amenable to

implementation as an external memory algorithm. If proved to be effective, such an approach

could provide a practical way of constructing MaxEnt models when there is a need to select a

small set of predictors in a pool comprising a potentially very large number of predictors.

This could lead to greater use of climate model outputs by the ecological research community

and aid the search for viable predictors when variable selection through ecological reasoning

is not apparent.

Materials and Methods

Cassin's Sparrow (*Peucaea cassinii* Woodhouse, 1852) is an elusive resident of arid

shrub grasslands of Middle America and the Southwestern United States [38]. Desert-adapted

birds, such as Cassin's Sparrow, appear to be especially vulnerable to climate change

[39,40]. We chose Cassin's Sparrow as a target for our development efforts as an example of

a species whose study could benefit from the technical advances described here. Occurrence

data was obtained from the Global Biodiversity Information Facility (GBIF) for the year

2016 [41]. After removing replicates, a total of 1865 records were acquired. To limit spatial

extent and avoid pseudo-replication, we thinned the points to a radius of 16 km, which

resulted in a total of 609 observations. For predictors, we used Worldclim's standard 19

Bioclimatic (bioclim) environmental variables at a resolution of 2.5 arc-minutes throughout

(Table 1) [42,43]. These predictor layers were clipped to the coverage area of our

observational data, reprojected, and formatted for use with MaxEnt using the Geospatial Data

Abstraction Library (GDAL) software package [44] following the guidelines of Hijmans et

al. [45]. We did not attempt to minimize collinearity by removing variables, because the

current study focuses on an assessment of stochastic down-selection from a full variable set,

and because MaxEnt has a demonstrated ability to account well for redundant variables [46].

We used MaxEnt Version 3.4.1 [47], R Version 4.0.1 [48], the ENMEval Version

0.3.0 R package [49], RStudio Version 1.2.5033 [50], and ENMTools Version 1.4.4 [51]

running on a 2.8 GHz Intel Core i7 MacBook Pro with 16 GB of memory in the study. First,

we developed a baseline model using the stand-alone MaxEnt program operated through its

graphical user interface (GUI). MaxEnt users can apply various combinations of five

mathematical transformations ('feature classes' or FCs) to predictor variables to enable more

complex fits to the observational data. The available feature types for continuous variables

are linear (L), quadratic (Q), hinge (H), product (P), and threshold (T) [4]. Users can also

adjust a regularization multiplier (RM) to maximize predictive accuracy and offset the

Table 1. Worldclim Bioclimatic Variables.

bio01	Annual Mean Temperature
bio02	Mean Diurnal Range (Mean of monthly (max temp - min temp))
bio03	Isothermality (BIO2/BIO7) (×100)
bio04	Temperature Seasonality (standard deviation ×100)
bio05	Max Temperature of Warmest Month
bio06	Min Temperature of Coldest Month
bio07	Temperature Annual Range (BIO5-BIO6)
bio08	Mean Temperature of Wettest Quarter
bio09	Mean Temperature of Driest Quarter
bio10	Mean Temperature of Warmest Quarter
bio11	Mean Temperature of Coldest Quarter
bio12	Annual Precipitation
bio13	Precipitation of Wettest Month
bio14	Precipitation of Driest Month
bio15	Precipitation Seasonality (Coefficient of Variation)
bio16	Precipitation of Wettest Quarter
bio17	Precipitation of Driest Quarter
bio18	Precipitation of Warmest Quarter
bio19	Precipitation of Coldest Quarter

overfitting that FC adjustments can introduce. We applied MaxEnt's default FC and RM

settings (i.e. the "Auto features" setting) with 10 replicate cross-validation and jackknife

evaluation of variable importance. By default, MaxEnt uses all feature classes and a

regularization multiplier of 1.0 when there are more than 80 training samples, which was the

case here [47]. Ten thousand background points were selected from across the study area

following the recommendations of Phillips et al. [52] and Fourcade et al. [53]. We

determined the average permutation importance for each variable in three replicated runs.

The top six predictors in the three-run ensemble were used to develop the final MaxEnt

baseline model.

We then developed an alternative method to select the top six variables that is based

on random sampling. We implemented our Monte Carlo approach as an R script that invokes

MaxEnt through ENMEval, which provides convenient control over model settings, built-in

evaluation metrics, and improved performance [35,49]. To reduce variability and isolate

outcomes as much as possible to the effects of the sampling process, we adopted a feature

class setting of LQHP and a regularization multiplier setting of 1.0 as fixed parameters in all

the Monte Carlo runs. We defined ensemble, in this case, to mean a collection of 100 sprints,

where each sprint consisted of ten model runs. A tally table was used to maintain a count of

the number of times a variable was used in a model run along with a cumulative sum of the

variable's permutation importance. The tally table thus provided the information needed to

determine the average permutation importance of a predictor at any point along the way.

To process a sprint, we initialized each of its ten model runs with a random subset of

environmental variables read from the filesystem. Random integers drawn from a uniform

distribution ranging 1–19 corresponding to the 19 bioclim predictors were used to make the

selection. At the conclusion of each model run, the tally table was updated appropriately. At

the conclusion of each sprint, we computed a MaxEnt model using the six predictors in the

original starting set having the highest average permutation importance values at that point.

This process was repeated 100 times to produce a complete ensemble. We assessed the

algorithm's performance in two ensembles. In the first, we chose two random variables for

each sprint run; in the second, six random variables were used for each run. This resulted in

an overall total of 2000 model runs.

The predictive distribution maps produced by the models were judged for

reasonableness based on first-hand knowledge of the species, its habitat preferences, and

known range [54]. We further compared model predictions to observational records from

Cornell Lab's eBird citizen-scientist database [55]. We used the area under the operating

curve (AUC) [56] as an indication of a model's classification accuracy (higher values

indicating greater accuracy) and the Akaike information criterion corrected for small sample

size (AICc) [57] as a measure of relative explanatory power (lower values indicating less

information loss). Model similarity was compared with Warren's I-statistic [58] and

Schoener's D statistic [59] (higher values in both indicating greater similarity) using

ENMTools. Single-processor run times were recorded to aid our understanding of algorithm

performance and help identify opportunities for multiprocessor parallelization.

Results

On the basis of permutation importance, 13 of the 19 original bioclim variables were

among the top ten most contributory predictors across all three replicated runs of the MaxEnt

baseline: bio02, bio03, bio05, bio06, bio08–bio12, bio14, bio15, bio17, and bio18 (Table 2).

Of those, bio02, bio05, and bio14 appeared in only one run each at 10th place. Bio18 showed

strong dominance throughout. When performance was averaged across all three runs, the top

six contributory variables in the ensemble collectively accounted for 65% of overall

permutation importance (ensemble average). In descending order of importance, the top six

predictors included bio18, bio03, bio10, bio15, bio11, and bio06. When these six top-

contributing variables were used in a final MaxEnt run, the model's four most contributory

variables (bio18, bio03, bio10, and bio15) accounted for approximately 86% of overall

permutation importance, and its predicted habitat suitability distribution corresponded well

with what is known about the natural history of the species and observational records for

Cassin's Sparrow for the year 2016 (Fig 1) [55].

A distinct pattern of progression toward a stable subset of key variables was observed

in the Monte Carlo ensembles (Figs 2,3). In both cases, the top three contributory variables

among the top six were selected early in the sprint runs, and AICc values fluctuated within a

narrow range around an average that changed little over the course of the selection process.

Greater variability in the composition of the top six subset was seen in Ensemble #1 where

two random variables at a time were selected for each sprint run (Table 2, Fig 2). In

Ensemble #2, where six random variables at a time were selected for the MaxEnt runs, the

top six variables were identified by the 25th sprint and had settled into their final rank order

by sprint 54 (Fig 3). Ensemble #2 appeared to produce the best overall results and shared

four variables in common with the top six selected by the MaxEnt baseline (bio03, bio06,

bio11, and bio18) (Table 2). Ensemble #2's final model had the lowest overall AICc, and its

four most contributory variables accounted for approximately 93% of overall permutation

importance, the highest attained overall.

Ensemble #1 had only one variable in common with the top six selected by both the

baseline run and Ensemble #2. What accounts for this difference is not immediately apparent;

however, we speculate that the random pair-wise comparisons occurring in Ensemble #1 may

alter the relative global influence of the collinearities known to exist in the bioclim variables

[62–64]. The average number of times a variable was sampled appeared to have a marginal,

positive influence on resulting model quality once an adequate minimum was attained.

Ensemble #2 results suggest that at least 80 uniformly distributed samples per starting-set

variable are needed to identify a reasonable top six set of variables; the best overall model

resulted from over 300 samples per variable (Table 2).

Table 2. Results of Maxent Baseline and Monte Carlo Selection Trials.

MODELS	bio01	bio02	bio03	bio04	bio05	bio06	bio07	bio08	bio09	bio10	bio11	bio12	bio13	bio14	bio15	bio16	bio17	bio18	bio19	AUC	AICc	% Top 3	% Top 4	Total Run Count	Total Run Time Mins	Hrs	Avg Random Samples/Variable
Maxent Baseline																											
Run #1	1.0	4.3	5.6	2.8	0.6	0.9	0.9	5.2	4.2	8.4	4.9	4.6	1.0	2.7	8.3	0.0	2.2	30.6	1.7					1	151	2.5	–
Run #2	1.7	2.9	6.4	1.9	3.5	9.9	0.5	3.5	3.9	5.4	7.7	4.8	1.0	3.5	8.3	0.8	0.5	30.1	3.6					1	120	2.0	–
Run #3	0.8	3.4	6.1	1.3	1.9	10.4	1.0	2.5	4.5	7.6	8.3	4.6	2.3	3.4	7.1	0.1	3.4	28.9	2.5					1	100	1.7	–
Ensemble avg	1.2	3.5	6.0	2.0	2.0	7.1	0.8	3.7	4.2	7.1	7.0	4.7	1.4	3.2	7.9	0.3	2.0	29.9	2.6								
Final model			14.1			6.6				13.8	7.3				10.9			47.4		0.818	12,222	75.3	86.2	1	18.0	0.3	–
Monte Carlo Selection																											
Ensemble #1	*(two random variables per sprint run)*																										
Sprint 025					17.6			10.8	9.3				22.2			14.3		25.9		0.801	12,229	65.7	80.0	250	181	3.0	26
Sprint 050					15.1			11.7	11.2				21.2			15.2		25.5		0.802	12,231	61.9	77.0	500	355	5.9	53
Sprint 100				24.1	18.2			6.2					7.6			4.1		39.7		0.806	12,252	82.0	89.6	1000	710	11.8	105
Ensemble #2	*(six random variables per sprint run)*																										
Sprint 025			25.4			3.6					15.6		7.7			3.8		43.9		0.801	12,376	84.9	92.6	250	438	7.3	79
Sprint 050			26.6			4.9					13.5		3.8			3.7		47.7		0.807	12,168	87.8	92.7	500	856	14.3	158
Sprint 100			26.5			5.5					18.9		4.1			2.6		42.4		0.805	12,152	87.8	93.3	1000	1793	29.9	316

Fig 1. Cassin's Sparrow distribution maps. Range map depicting the seasonally-averaged estimated relative abundance of Cassin's Sparrow (A) compared to the species' predicted habitat suitability distributions obtained from the MaxEnt baseline (B), Monte Carlo Ensemble #1 (C), and Monte Carlo Ensemble #2 (D). Image (A) provided by eBird (www.ebird.org) and created 28 July 2020; images (B)–(D) created by the authors using MaxEnt Version 3.4.1 [61].

Sprint	Runs	Time (min)	AICc	1st	2nd	3rd	4th	5th	6th
1	10	7.58	12379.01	bio05	bio08	bio14	bio04	bio17	bio03
2	20	8.04	12200.28	bio15	bio08	bio16	bio12	bio01	bio14
3	30	6.99	12167.31	bio08	bio16	bio18	bio14	bio05	bio09
4	40	6.87	12232.75	bio18	bio16	bio08	bio05	bio09	bio12
5	50	7.31	12170.46	bio18	bio16	bio08	bio05	bio13	bio09
6	60	7.17	12149.38	bio18	bio16	bio08	bio05	bio13	bio09
7	70	6.69	12226.38	bio18	bio16	bio13	bio08	bio05	bio09
8	80	7.73	12222.55	bio18	bio16	bio13	bio05	bio08	bio09
9	90	6.40	12157.66	bio18	bio16	bio13	bio08	bio05	bio09
10	100	7.89	12195.88	bio18	bio16	bio13	bio08	bio05	bio15
11	110	8.48	12221.87	bio18	bio16	bio13	bio08	bio05	bio09
12	120	7.11	12265.25	bio18	bio16	bio13	bio05	bio08	bio09
13	130	5.60	12251.47	bio18	bio16	bio13	bio05	bio08	bio11
14	140	8.10	12218.51	bio18	bio16	bio13	bio08	bio05	bio09
15	150	7.16	12198.43	bio18	bio16	bio13	bio08	bio05	bio09
16	160	7.32	12222.59	bio18	bio16	bio13	bio08	bio05	bio09
17	170	6.99	12216.19	bio18	bio16	bio13	bio08	bio05	bio03
18	180	6.95	12199.30	bio18	bio16	bio13	bio08	bio05	bio12
19	190	6.20	12211.87	bio18	bio16	bio13	bio08	bio05	bio03
20	200	8.30	12257.88	bio18	bio16	bio13	bio08	bio05	bio03
21	210	7.08	12220.48	bio18	bio16	bio13	bio08	bio05	bio09
22	220	6.94	12235.24	bio18	bio16	bio13	bio08	bio05	bio09
23	230	7.53	12209.08	bio18	bio16	bio13	bio08	bio05	bio09
24	240	8.17	12229.47	bio18	bio16	bio13	bio08	bio05	bio09
25	250	6.61	12229.17	bio18	bio16	bio13	bio08	bio05	bio09
26	260	7.67	12227.33	bio18	bio16	bio13	bio08	bio05	bio09
27	270	7.68	12240.54	bio18	bio16	bio13	bio08	bio05	bio09
28	280	5.29	12253.73	bio18	bio16	bio13	bio08	bio05	bio09
29	290	7.69	12245.23	bio18	bio16	bio13	bio08	bio05	bio09
30	300	8.49	12277.86	bio18	bio16	bio13	bio08	bio05	bio09
31	310	6.38	12245.44	bio18	bio16	bio13	bio08	bio05	bio09
32	320	7.53	12189.91	bio18	bio16	bio13	bio08	bio05	bio09
33	330	7.08	12233.77	bio18	bio16	bio13	bio08	bio05	bio09
34	340	6.79	12249.50	bio18	bio16	bio13	bio08	bio05	bio09
35	350	6.52	12213.96	bio18	bio16	bio13	bio08	bio05	bio09
36	360	6.80	12243.56	bio18	bio16	bio13	bio08	bio05	bio09
37	370	7.71	12245.80	bio18	bio16	bio13	bio08	bio05	bio09
38	380	7.96	12225.07	bio18	bio16	bio13	bio08	bio05	bio09
39	390	6.42	12249.01	bio18	bio16	bio13	bio08	bio05	bio09
40	400	5.70	12253.59	bio18	bio16	bio13	bio08	bio05	bio09
41	410	7.08	12206.76	bio18	bio16	bio13	bio08	bio05	bio09
42	420	7.06	12259.68	bio18	bio16	bio13	bio08	bio05	bio09
43	430	6.41	12264.71	bio18	bio16	bio13	bio08	bio05	bio09
44	440	6.70	12246.71	bio18	bio16	bio13	bio08	bio05	bio09
45	450	6.06	12297.09	bio18	bio16	bio13	bio08	bio05	bio09
46	460	6.93	12214.90	bio18	bio16	bio13	bio08	bio05	bio09
47	470	6.81	12250.17	bio18	bio16	bio13	bio08	bio05	bio09
48	480	6.71	12212.33	bio18	bio16	bio13	bio08	bio05	bio09
49	490	6.95	12230.75	bio18	bio16	bio13	bio08	bio05	bio09
50	500	7.06	12230.62	bio18	bio16	bio13	bio08	bio05	bio09
51	510	7.78	12233.56	bio18	bio16	bio13	bio08	bio05	bio09
52	520	8.06	12259.47	bio18	bio16	bio13	bio08	bio05	bio09
53	530	6.91	12241.89	bio18	bio16	bio13	bio08	bio05	bio09
54	540	8.19	12247.57	bio18	bio16	bio13	bio08	bio05	bio09
55	550	7.69	12226.76	bio18	bio16	bio13	bio08	bio05	bio09
56	560	6.49	12210.05	bio18	bio16	bio13	bio08	bio05	bio09
57	570	7.90	12205.59	bio18	bio16	bio13	bio08	bio05	bio09
58	580	5.90	12254.37	bio18	bio16	bio13	bio08	bio05	bio09
59	590	6.75	12273.07	bio18	bio16	bio13	bio08	bio05	bio09
60	600	6.82	12236.51	bio18	bio16	bio13	bio08	bio05	bio09
61	610	7.22	12243.03	bio18	bio16	bio13	bio08	bio05	bio09
62	620	7.73	12198.01	bio18	bio16	bio13	bio08	bio05	bio09
63	630	8.41	12338.98	bio18	bio16	bio13	bio08	bio05	bio09
64	640	6.79	12245.66	bio18	bio16	bio13	bio08	bio05	bio09
65	650	6.98	12251.55	bio18	bio16	bio13	bio08	bio05	bio09
66	660	7.02	12348.13	bio18	bio16	bio13	bio08	bio05	bio09
67	670	7.28	12239.33	bio18	bio16	bio13	bio08	bio05	bio09
68	680	7.92	12343.54	bio18	bio16	bio13	bio08	bio05	bio09
69	690	7.40	12237.15	bio18	bio16	bio13	bio08	bio05	bio09
70	700	6.83	12222.24	bio18	bio16	bio13	bio08	bio05	bio09
71	710	7.89	12280.63	bio18	bio16	bio13	bio08	bio05	bio09
72	720	6.65	12234.48	bio18	bio16	bio13	bio08	bio05	bio09
73	730	6.97	12256.76	bio18	bio16	bio13	bio08	bio05	bio09
74	740	7.58	12279.34	bio18	bio16	bio13	bio08	bio05	bio09
75	750	7.08	12210.23	bio18	bio16	bio13	bio08	bio05	bio09
76	760	6.97	12232.46	bio18	bio16	bio13	bio08	bio05	bio09
77	770	7.14	12298.37	bio18	bio16	bio13	bio08	bio05	bio09
78	780	6.80	12220.16	bio18	bio16	bio13	bio08	bio05	bio09
79	790	7.62	12214.21	bio18	bio16	bio13	bio08	bio05	bio09
80	800	7.41	12227.38	bio18	bio16	bio13	bio08	bio05	bio09
81	810	7.18	12250.13	bio18	bio16	bio13	bio08	bio05	bio09
82	820	7.65	12264.45	bio18	bio16	bio13	bio08	bio05	bio03
83	830	6.20	12245.57	bio18	bio16	bio13	bio08	bio05	bio03
84	840	6.33	12216.66	bio18	bio16	bio13	bio08	bio05	bio03
85	850	8.21	12242.54	bio18	bio16	bio13	bio08	bio05	bio03
86	860	6.49	12282.52	bio18	bio16	bio13	bio08	bio05	bio03
87	870	6.87	12256.19	bio18	bio16	bio13	bio08	bio05	bio03
88	880	7.06	12251.20	bio18	bio16	bio13	bio08	bio05	bio03
89	890	6.78	12237.27	bio18	bio16	bio13	bio08	bio05	bio03
90	900	8.21	12278.73	bio18	bio16	bio13	bio08	bio05	bio09
91	910	6.06	12307.16	bio18	bio16	bio13	bio08	bio05	bio09
92	920	5.29	12210.69	bio18	bio16	bio13	bio08	bio05	bio09
93	930	7.93	12245.61	bio18	bio16	bio13	bio08	bio05	bio03
94	940	5.88	12229.42	bio18	bio16	bio13	bio08	bio05	bio03
95	950	8.25	12236.33	bio18	bio16	bio13	bio08	bio05	bio03
96	960	6.15	12280.85	bio18	bio16	bio13	bio08	bio05	bio03
97	970	6.42	12273.77	bio18	bio16	bio13	bio08	bio05	bio03
98	980	6.38	12247.16	bio18	bio16	bio13	bio08	bio05	bio03
99	990	7.06	12214.92	bio18	bio16	bio13	bio08	bio05	bio03
100	1000	7.17	12252.44	bio18	bio16	bio13	bio08	bio05	bio03

Fig 2. Monte Carlo Ensemble #1 results. Two random variables at a time were chosen for each MaxEnt sprint run. The sprint log on top shows the progressive selection of a stable set of top six variables in yellow. The graph on the bottom shows the narrow range of fluctuating AICc values over the course of the ensemble runs. Maximum and minimum AICc values are shown in red.

Sprint	Runs	Time (min)	AICc	1st	2nd	3rd	4th	5th	6th
1	10	18.07	12185.35	bio18	bio06	bio12	bio16	bio03	bio11
2	20	18.32	12182.82	bio18	bio16	bio06	bio12	bio03	bio14
3	30	17.60	12232.96	bio18	bio16	bio13	bio06	bio03	bio15
4	40	18.92	12180.38	bio18	bio16	bio13	bio03	bio12	bio14
5	50	16.99	12207.91	bio18	bio16	bio13	bio03	bio06	bio04
6	60	16.67	12191.40	bio18	bio16	bio13	bio03	bio06	bio11
7	70	15.97	12167.45	bio18	bio16	bio13	bio06	bio03	bio11
8	80	17.16	12188.02	bio18	bio16	bio13	bio06	bio03	bio11
9	90	18.62	12161.62	bio18	bio16	bio13	bio06	bio03	bio11
10	100	17.70	12179.81	bio18	bio16	bio13	bio06	bio11	bio03
11	110	18.57	12163.00	bio18	bio16	bio13	bio06	bio11	bio03
12	120	17.67	12161.86	bio18	bio16	bio13	bio06	bio11	bio03
13	130	18.03	12195.77	bio18	bio16	bio13	bio06	bio11	bio03
14	140	17.45	12162.09	bio18	bio16	bio13	bio06	bio11	bio03
15	150	16.41	12153.21	bio18	bio16	bio13	bio06	bio11	bio03
16	160	17.04	12169.11	bio18	bio16	bio13	bio06	bio11	bio03
17	170	16.71	12173.60	bio18	bio16	bio13	bio06	bio11	bio03
18	180	16.99	12160.81	bio18	bio16	bio13	bio06	bio11	bio03
19	190	19.12	12165.07	bio18	bio16	bio13	bio06	bio11	bio03
20	200	17.50	12160.62	bio18	bio16	bio13	bio06	bio11	bio03
21	210	18.51	12151.23	bio18	bio16	bio13	bio06	bio11	bio03
22	220	17.35	12281.16	bio18	bio16	bio13	bio06	bio11	bio03
23	230	16.64	12200.43	bio18	bio16	bio13	bio06	bio11	bio03
24	240	16.82	12172.95	bio18	bio16	bio13	bio06	bio11	bio03
25	250	17.11	12176.06	bio18	bio16	bio13	bio06	bio11	bio03
26	260	17.25	12166.65	bio18	bio16	bio13	bio06	bio11	bio03
27	270	16.91	12203.32	bio18	bio16	bio13	bio06	bio11	bio03
28	280	15.79	12182.27	bio18	bio16	bio13	bio06	bio11	bio03
29	290	17.31	12186.27	bio18	bio16	bio13	bio06	bio11	bio03
30	300	18.19	12238.20	bio18	bio16	bio13	bio06	bio11	bio03
31	310	16.43	12177.40	bio18	bio16	bio13	bio06	bio11	bio03
32	320	17.14	12159.92	bio18	bio16	bio13	bio06	bio11	bio03
33	330	18.26	12149.59	bio18	bio16	bio13	bio06	bio11	bio03
34	340	16.17	12178.16	bio18	bio16	bio13	bio06	bio11	bio03
35	350	16.78	12153.80	bio18	bio16	bio13	bio11	bio06	bio03
36	360	18.70	12158.82	bio18	bio16	bio13	bio06	bio11	bio03
37	370	18.12	12179.92	bio18	bio16	bio13	bio11	bio06	bio03
38	380	17.30	12187.95	bio18	bio16	bio13	bio11	bio06	bio03
39	390	17.48	12208.89	bio18	bio16	bio13	bio11	bio06	bio03
40	400	17.13	12203.25	bio18	bio16	bio13	bio11	bio06	bio03
41	410	18.86	12167.30	bio18	bio16	bio13	bio11	bio06	bio03
42	420	18.89	12166.73	bio18	bio16	bio13	bio11	bio06	bio03
43	430	17.32	12170.09	bio18	bio16	bio13	bio11	bio06	bio03
44	440	17.30	12181.02	bio18	bio16	bio13	bio11	bio06	bio03
45	450	16.42	12194.59	bio18	bio16	bio13	bio11	bio06	bio03
46	460	18.32	12148.92	bio18	bio16	bio13	bio11	bio06	bio03
47	470	17.03	12196.84	bio18	bio16	bio13	bio11	bio06	bio03
48	480	18.00	12178.66	bio18	bio16	bio13	bio11	bio06	bio03
49	490	17.28	12168.32	bio18	bio16	bio13	bio11	bio06	bio03
50	500	16.30	12198.48	bio18	bio16	bio13	bio11	bio06	bio03

Sprint	Runs	Time (min)	AICc	1st	2nd	3rd	4th	5th	6th
51	510	18.53	12175.76	bio18	bio16	bio13	bio11	bio06	bio03
52	520	20.15	12198.96	bio18	bio16	bio13	bio11	bio06	bio03
53	530	20.48	12168.05	bio18	bio16	bio13	bio06	bio11	bio03
54	540	17.22	12163.76	bio18	bio16	bio13	bio11	bio06	bio03
55	550	17.28	12165.65	bio18	bio16	bio13	bio06	bio11	bio03
56	560	20.14	12189.90	bio18	bio16	bio13	bio06	bio11	bio03
57	570	20.86	12161.10	bio18	bio16	bio13	bio06	bio11	bio03
58	580	19.10	12182.45	bio18	bio16	bio13	bio06	bio11	bio03
59	590	17.63	12179.96	bio18	bio16	bio13	bio06	bio11	bio03
60	600	19.88	12182.09	bio18	bio16	bio13	bio06	bio11	bio03
61	610	18.92	12166.12	bio18	bio16	bio13	bio06	bio11	bio03
62	620	19.17	12163.61	bio18	bio16	bio13	bio06	bio11	bio03
63	630	18.74	12152.40	bio18	bio16	bio13	bio06	bio11	bio03
64	640	20.34	12201.92	bio18	bio16	bio13	bio06	bio11	bio03
65	650	19.29	12202.01	bio18	bio16	bio13	bio06	bio11	bio03
66	660	19.49	12185.66	bio18	bio16	bio13	bio06	bio11	bio03
67	670	19.05	12177.32	bio18	bio16	bio13	bio06	bio11	bio03
68	680	18.96	12192.10	bio18	bio16	bio13	bio06	bio11	bio03
69	690	18.21	12164.35	bio18	bio16	bio13	bio06	bio11	bio03
70	700	20.04	12183.12	bio18	bio16	bio13	bio06	bio11	bio03
71	710	18.20	12200.74	bio18	bio16	bio13	bio06	bio11	bio03
72	720	19.34	12173.30	bio18	bio16	bio13	bio06	bio11	bio03
73	730	17.71	12160.29	bio18	bio16	bio13	bio06	bio11	bio03
74	740	19.39	12187.62	bio18	bio16	bio13	bio06	bio11	bio03
75	750	17.44	12164.62	bio18	bio16	bio13	bio06	bio11	bio03
76	760	17.59	12197.39	bio18	bio16	bio13	bio06	bio11	bio03
77	770	17.78	12191.93	bio18	bio16	bio13	bio06	bio11	bio03
78	780	19.23	12180.93	bio18	bio16	bio13	bio06	bio11	bio03
79	790	18.43	12175.62	bio18	bio16	bio13	bio06	bio11	bio03
80	800	18.13	12154.68	bio18	bio16	bio13	bio06	bio11	bio03
81	810	18.11	12164.86	bio18	bio16	bio13	bio06	bio11	bio03
82	820	20.02	12161.18	bio18	bio16	bio13	bio06	bio11	bio03
83	830	19.08	12176.15	bio18	bio16	bio13	bio06	bio11	bio03
84	840	19.17	12178.22	bio18	bio16	bio13	bio06	bio11	bio03
85	850	20.01	12187.92	bio18	bio16	bio13	bio06	bio11	bio03
86	860	17.05	12165.70	bio18	bio16	bio13	bio06	bio11	bio03
87	870	17.78	12164.13	bio18	bio16	bio13	bio06	bio11	bio03
88	880	18.00	12155.70	bio18	bio16	bio13	bio06	bio11	bio03
89	890	16.87	12250.99	bio18	bio16	bio13	bio06	bio11	bio03
90	900	17.98	12186.50	bio18	bio16	bio13	bio06	bio11	bio03
91	910	17.64	12179.19	bio18	bio16	bio13	bio06	bio11	bio03
92	920	17.89	12175.84	bio18	bio16	bio13	bio06	bio11	bio03
93	930	15.97	12186.11	bio18	bio16	bio13	bio06	bio11	bio03
94	940	17.25	12170.26	bio18	bio16	bio13	bio06	bio11	bio03
95	950	16.83	12199.93	bio18	bio16	bio13	bio06	bio11	bio03
96	960	16.43	12176.31	bio18	bio16	bio13	bio06	bio11	bio03
97	970	18.13	12221.18	bio18	bio16	bio13	bio06	bio11	bio03
98	980	17.58	12169.45	bio18	bio16	bio13	bio06	bio11	bio03
99	990	15.28	12197.56	bio18	bio16	bio13	bio06	bio11	bio03
100	1000	16.84	12151.75	bio18	bio16	bio13	bio06	bio11	bio03

216 **Fig 3. Monte Carlo Ensemble #2 results.** Six random variables at a time were chosen for

217 each MaxEnt sprint run. The sprint log on top shows the progressive selection of a stable set

218 of top six variables in yellow. The graph on the bottom shows the narrow range of fluctuating

219 AICc values over the course of the ensemble runs. Maximum and minimum AICc values are

220 shown in red.

Discussion

The most striking outcome of the study is the similarity in results. Final models in the MaxEnt baseline and the Monte Carlo ensembles produced predicted habitat suitability distributions that are nearly indistinguishable from one another (Fig 1). Internal metrics likewise reveal little difference in outcomes, with AUC values ranging only from 0.801 to 0.818 and AICc ranging from 12,152 to 12,222, suggesting that the traditional MaxEnt runs and the Monte Carlo approach both produced reasonable models (Table 2). The two approaches also each identified four variables that collectively contributed more than 80% to the formulation of their respective models. Across the board, models showed a high degree of similarity in Schoener's D and the I-statistic (Table 3).

Table 3. Model Similarity Metrics.

Schoener's D Statistic

MODELS ↓→	Maxent-Run1	Maxent-Run2	Maxent-Run3	Maxent-Final	MC-E1-025	MC-E1-050	MC-E1-100	MC-E2-025	MC-E2-050	MC-E2-100
Maxent-Run1	1	0.9712	0.9738	0.9355	0.8629	0.8630	0.9044	0.8882	0.8903	0.8906
Maxent-Run2	x	1	0.9793	0.9397	0.8639	0.8648	0.9078	0.8915	0.8949	0.8947
Maxent-Run3	x	x	1	0.9354	0.8579	0.8586	0.9039	0.8871	0.8903	0.8902
Maxent-Final	x	x	x	1	0.8667	0.8673	0.9214	0.9104	0.9138	0.9142
MC-E1-025	x	x	x	x	1	0.9880	0.9006	0.8810	0.8801	0.8785
MC-E1-050	x	x	x	x	x	1	0.9026	0.8813	0.8804	0.8786
MC-E1-100	x	x	x	x	x	x	1	0.9393	0.9389	0.9365
MC-E2-025	x	x	x	x	x	x	x	1	0.9815	0.9810
MC-E2-050	x	x	x	x	x	x	x	x	1	0.9844
MC-E2-100	x	x	x	x	x	x	x	x	x	1

Warren's I Statistic

MODELS ↓→	Maxent-Run1	Maxent-Run2	Maxent-Run3	Maxent-Final	MC-E1-025	MC-E1-050	MC-E1-100	MC-E2-050	MC-E2-100	MC-E2-100
Maxent-Run1	1	0.9991	0.9994	0.9948	0.9760	0.9758	0.9874	0.9828	0.9833	0.9834
Maxent-Run2	x	1	0.9995	0.9949	0.9760	0.9760	0.9878	0.9831	0.9838	0.9839
Maxent-Run3	x	x	1	0.9945	0.9748	0.9748	0.9869	0.9823	0.9830	0.9831
Maxent-Final	x	x	x	1	0.9755	0.9753	0.9894	0.9871	0.9878	0.9880
MC-E1-025	x	x	x	x	1	0.9998	0.9887	0.9798	0.9797	0.9784
MC-E1-050	x	x	x	x	x	1	0.9889	0.9795	0.9794	0.9781
MC-E1-100	x	x	x	x	x	x	1	0.9910	0.9912	0.9906
MC-E2-025	x	x	x	x	x	x	x	1	0.9996	0.9994
MC-E2-050	x	x	x	x	x	x	x	x	1	0.9996
MC-E2-100	x	x	x	x	x	x	x	x	x	1

The most significant drawback identified in the study was the long run times.

MaxEnt's linear scaling behavior can be challenging in a single-processor environment. In

the baseline runs, producing a single model through MaxEnt's GUI using our selected

settings involved writing many files to disk and took from 18 minutes (with six variables) to

over two hours (with all 19 variables). MaxEnt in the R environment outputs memory-

resident objects, which results in faster run times. Still, with its repeated invocations of

MaxEnt, Ensemble #2 took nearly 30 hours to complete (Table 2).

The Monte Carlo approach also exhibits linear scaling properties; however, its scaling

behavior is not determined by the MaxEnt program, since each of the MaxEnt runs in the

Monte Carlo method operates on a set number of predictors. The method's linear scaling

property is determined, instead, by the need to adequately sample the starting set of

environmental variables in order to obtain a good result. What makes a practical

implementation of the Monte Carlo method possible is that each of its MaxEnt runs is

entirely independent from all other runs in the ensemble. This high level of subtask

independence is sometimes referred to as an "embarrassingly parallel" workload, which

makes it relatively straightforward to implement in a cluster computing environment. If 1000

processors were recruited into service — which is becoming increasingly convenient with the

proliferation of multiprocessor, high performance cloud computing — a 1000-run ensemble

could conceivably take about as long as a single MaxEnt run.

The potential significance of this advantage becomes apparent when one considers

the method's use with large collections of environmental data. The Monte Carlo approach

described here provides an approximate solution to the problem of finding a useful k-size

subset of an n-size collection of variables. In principal, there are $n! \, / \, [k!(n-k)!]$ variable

combinations to consider in such an evaluation, a staggering 27,000-plus six-variable subsets

with the 19 bioclim variables alone. Algorithms that accomplish variable selection through

stepwise removal or are otherwise bound to the linear scaling properties of underlying

software components are inherently unable to exhaustively explore this combinatorial space.

A Monte Carlo method makes such a search possible by randomly sampling the universe of

possible combinations and returning approximate solutions in practical amounts of time,

particularly if implemented as a high-performance cloud service.

The use of IPCC-class climate model outputs in efforts to assess the impacts of

climate change on biodiversity and other ecosystem processes is growing. Exploring the

potential of these massive data sets, expanded use of ensemble modeling, and the actual work

of fitting models for the thousands of species scientists wish to study will require hundreds to

thousands of projections [16,23]. An improved capacity to use large environmental data sets

in MaxEnt modeling would benefit this work. We are encouraged to think that innovative use

of Monte Carlo techniques might provide a helpful means of meeting this challenge.

Conclusions

This small-scale, proof-of-concept study leaves many practical and theoretical

questions unanswered. Preliminary results, however, suggest that a Monte Carlo method

could offer a viable means of selecting environmental predictors for MaxEnt models that is

amenable to parallelization and scalable to large data sets, including externally-stored

collections. This points to the possibility of near-real-time multiprocessor implementations

that could enable broader and more exploratory use of global climate model outputs in

environmental niche modeling and aid in the discovery of viable predictors. Next steps will

focus on implementing this capability in NASA's Advanced Data Analytics Platform

(ADAPT) science cloud, evaluating the method's behavior using products generated by the

Goddard Earth Observing System, Version 5 (GEOS-5) modeling system, extending

stochasticity to feature class and regularization multiplier selection, developing automatic

stopping rules, and evaluating the method's effectiveness in addressing research questions

relating to climate change influences on Cassin's Sparrow abundance and distribution.

282　References

1.　Elith J, Phillips SJ, Hastie T, Dudík M, Chee YE, Yates CJ. A statistical explanation of

MaxEnt for ecologists. Diversity and distributions. 2011;17: 43–57.

2.　Phillips SJ, Anderson RP, Dudík M, Schapire RE, Blair ME. Opening the black box: An

open-source release of Maxent. Ecography. 2017;40: 887–893.

3.　Phillips SJ. A brief tutorial on Maxent. AT&T Research. 2005;190: 231–259.

4.　Phillips SJ, Anderson RP, Schapire RE. Maximum Entropy Modeling of Species

Geographic Distributions. Ecological Modelling. 2006;190: 231–259.

doi:10.1016/j.ecolmodel.2005.03.026.

5.　Kalinski CE. Building Better Species Distribution Models with Machine Learning:

Assessing the Role of Covariate Scale and Tuning in Maxent Models. 2019; 129.

6.　Merow C, Smith MJ, Silander JA. A practical guide to MaxEnt for modeling species'

distributions: what it does, and why inputs and settings matter. Ecography. 2013;36:

1058–1069. doi:10.1111/j.1600-0587.2013.07872.x

7.   Schmidt-Lebuhn AN, Knerr NJ, Miller JT, Mishler BD. Phylogenetic diversity and

endemism of Australian daisies (Asteraceae). Journal of Biogeography. 2015;42: 1114–

1122. doi:10.1111/jbi.12488

8.   Warren DL, Wright AN, Seifert SN, Shaffer HB. Incorporating model complexity and

spatial sampling bias into ecological niche models of climate change risks faced by 90

California vertebrate species of concern. Diversity and distributions. 2014;20: 334–343.

9.   Cardoso-Leite R, Vilarinho AC, Novaes MC, Tonetto AF, Vilardi GC, Guillermo-

Ferreira R. Recent and future environmental suitability to dengue fever in Brazil using

species distribution model. Transactions of The Royal Society of Tropical Medicine and

Hygiene. 2014;108: 99–104. doi:10.1093/trstmh/trt115

10.   Morisette JT, Jarnevich CS, Ullah A, Cai W, Pedelty JA, Gentle JE, et al. A Tamarisk

Habitat Suitability Map for the Continental United States. Frontiers in Ecology and the

Environment. 2006;4: 11–17. doi:10.1890/1540-9295(2006)004

11.   Stohlgren TJ, Schnase JL. Risk Analysis for Biological Hazards: What We Need to

Know about Invasive Species. Risk Analysis. 2006;26: 163–73. doi:10.1111/j.1539-

6924.2006.00707.x.

12.   Beauchamp VB, Koontz SM, Suss C, Hawkins C, Kyde KL, Schnase JL. An

Introduction to Oplismenus Undulatifolius (Ard.) Roem. & Schult (Wavyleaf

Basketgrass), a Recent Invader in Mid-Atlantic Forest Understories 1,2. The Journal of

the Torrey Botanical Society. 2013;140: 391–413. doi:10.3159/torrey-d-13-00033.1.

13.   alwi Muttaqin luthfi, Heru Murti S, Susilo B. MaxEnt (Maximum Entropy) model for

predicting prehistoric cave sites in Karst area of Gunung Sewu, Gunung Kidul,

Yogyakarta. In: Wibowo SB, Rimba AB, A. Aziz A, Phinn S, Sri Sumantyo JT,

Widyasamratri H, et al., editors. Sixth Geoinformation Science Symposium.

Yogyakarta, Indonesia: SPIE; 2019. p. 3. doi:10.1117/12.2543522

14.   Feng X, Park DS, Walker C, Peterson AT, Merow C, Papeş M. A checklist for

maximizing reproducibility of ecological niche models. Nature Ecology & Evolution.

2019;3: 1382–1395. doi:10.1038/s41559-019-0972-5

15.   Morales NS, Fernández IC, Baca-González V. MaxEnt's parameter configuration and

small samples: are we paying attention to recommendations? A systematic review.

PeerJ. 2017;5: e3093. doi:10.7717/peerj.3093

16.   Araújo M, New M. Ensemble forecasting of species distributions. Trends in Ecology &

Evolution. 2007;22: 42–47. doi:10.1016/j.tree.2006.09.010

17.   Zeng Y, Low BW, Yeo DCJ. Novel methods to select environmental variables in

MaxEnt: A case study using invasive crayfish. Ecological Modelling. 2016;341: 5–13.

doi:https://doi.org/10.1016/j.ecolmodel.2016.09.019

18.   Araújo MB, Anderson RP, Barbosa AM, Beale CM, Dormann CF, Early R, et al.

Standards for distribution models in biodiversity assessments. Science Advances.

2019;5: eaat4858.

19.   Qiao H, Soberón J, Peterson AT. No silver bullets in correlative ecological niche

modelling: insights from testing among many potential algorithms for niche estimation.

Kriticos D, editor. Methods in Ecology and Evolution. 2015;6: 1126–1136.

doi:10.1111/2041-210X.12397

20. Ashraf U, Peterson AT, Chaudhry MN, Ashraf I, Saqib Z, Rashid Ahmad S, et al.

Ecological niche model comparison under different climate scenarios: a case study of

Olea spp. in Asia. Ecosphere. 2017;8: e01825. doi:10.1002/ecs2.1825

21. Guisan A, Zimmermann NE. Predictive habitat distribution models in ecology.

Ecological modelling. 2000;135: 147–186.

22. Li Y, Li M, Li C, Liu Z. Optimized Maxent Model Predictions of Climate Change

Impacts on the Suitable Distribution of Cunninghamia lanceolata in China. Forests.

2020;11: 302. doi:10.3390/f11030302

23. Cavanagh RD, Murphy EJ, Bracegirdle TJ, Turner J, Knowland CA, Corney SP, et al. A

Synergistic Approach for Evaluating Climate Model Output for Ecological

Applications. Frontiers in Marine Science. 2017;4: 308. doi:10.3389/fmars.2017.00308

24. Harris RMB, Grose MR, Lee G, Bindoff NL, Porfirio LL, Fox-Hughes P. Climate

projections for ecologists: Climate projections for ecologists. Wiley Interdisciplinary

Reviews: Climate Change. 2014;5: 621–637. doi:10.1002/wcc.291

25. Stock CA, Alexander MA, Bond NA, Brander KM, Cheung WW, Curchitser EN, et al.

On the use of IPCC-class models to assess the impact of climate on living marine

resources. Progress in Oceanography. 2011;88: 1–27.

26.   Bojinski S, Verstraete M, Peterson TC, Richter C, Simmons A, Zemp M. The Concept

357   of Essential Climate Variables in Support of Climate Research, Applications, and

358   Policy. Bulletin of the American Meteorological Society. 2014;95: 1431–1443.

359   doi:10.1175/BAMS-D-13-00047.1

360   27.   Braunisch V, Coppes J, Arlettaz R, Suchant R, Schmid H, Bollmann K. Selecting from

361   correlated climate variables: a major source of uncertainty for predicting species

362   distributions under climate change. Ecography. 2013;36: 971–983. doi:10.1111/j.1600-

363   0587.2013.00138.x

364   28.   Schnase JL. Climate Analytics as a Service. Cloud Computing in Ocean and

365   Atmospheric Sciences. 2016. pp. 187–219. doi:10.1016/b978-0-12-803192-6.00011-6

366   29.   Edwards PN. A Vast Machine: Computer Models, Climate Data, and the Politics of

367   Global Warming. Cambridge, MA: MIT Press; 2010.

368   30.   IPCC — Intergovernmental Panel on Climate Change. 2020 [cited 14 Mar 2020].

369   Available: https://www.ipcc.ch/

370   31.   Responding to the Challenge of Climate and Environmental Change: NASA's Plan for a

371   Climate-Centric Architecture for Earth Observations and Applications from Space.

372   National Aeronautics and Space Administration; 2010. Available:

373   https://gmao.gsfc.nasa.gov/reanalysis/MERRA/.

374   32.   Araújo MB, Guisan A. Five (or so) challenges for species distribution modelling.

375   Journal of Biogeography. 2006;33: 1677–1688. doi:10.1111/j.1365-2699.2006.01584.x

376   33.   Duan Y, Edwards JS, Dwivedi YK. Artificial intelligence for decision making in the era

377   of Big Data – evolution, challenges and research agenda. International Journal of

378   Information Management. 2019;48: 63–71. doi:10.1016/j.ijinfomgt.2019.01.021

379   34.   Galante PJ, Alade B, Muscarella R, Jansa SA, Goodman SM, Anderson RP. The

380   challenge of modeling niches and distributions for data-poor species: a comprehensive

381   approach to model complexity. Ecography. 2018;41: 726–736.

382   35.   Muscarella R, Galante PJ, Soley-Guardia M, Boria RA, Kass JM, Uriarte M, et al.

383   ENMeval: An R package for conducting spatially independent evaluations and

384   estimating optimal model complexity for Maxent ecological niche models. Methods in

385   Ecology and Evolution. 2014;5: 1198–1205. doi:10.1111/2041-210X.12261

386   36.   Radosavljevic A, Anderson RP. Making better MaxEnt models of species distributions:

387   complexity, overfitting and evaluation. Araújo M, editor. Journal of Biogeography.

388   2014;41: 629–643. doi:10.1111/jbi.12227

389   37.   Kroese DP, Brereton T, Taimre T, Botev ZI. Why the Monte Carlo method is so

390   important today: Why the MCM is so important today. Wiley Interdisciplinary

391   Reviews: Computational Statistics. 2014;6: 386–392. doi:10.1002/wics.1314

392   38.   Dunning, Jr. JB, Bowers, Jr. RK, Suter SJ, Bock CE. Cassin's Sparrow (Peucaea

393   cassinii), Version 1.0. In: Birds of the World (P. G. Rodewald, Editor) [Internet]. 2020

394   [cited 22 May 2020]. Available: https://doi.org/10.2173/bow.casspa.01

39. Iknayan KJ, Beissinger SR. Collapse of a desert bird community over the past century

driven by climate change. Proc Natl Acad Sci USA. 2018;115: 8597.

doi:10.1073/pnas.1805123115

40. Radchuk V, Reed T, Teplitsky C, van de Pol M, Charmantier A, Hassall C, et al.

Adaptive responses of animals to climate change are most likely insufficient. Nature

Communications. 2019;10: 3109. doi:10.1038/s41467-019-10924-4

41. GBIF.org (21 February 2019) GBIF Occurrence Download

https://doi.org/10.15468/dl.0s8yak.

42. Fick SE, Hijmans RJ. WorldClim 2: new 1-km spatial resolution climate surfaces for

global land areas. International Journal of Climatology. 2017;37: 4302–4315.

doi:10.1002/joc.5086

43. Worldclim bioclimatic variables. 2020 [cited 22 May 2020]. Available:

https://worldclim.org/data/worldclim21.html

44. GDAL/OGR Geospatial Data Abstraction Software Library. Open Source Geospatial

Foundation; 2020. Available: https://gdal.org/

45. Hijmans RJ, Phillips S, Elith J, Leathwick J. dismo: Species Distribution Modeling.

2017. Available: https://CRAN.R-project.org/package=dismo

46. Feng X, Park DS, Liang Y, Pandey R, Papeş M. Collinearity in ecological niche

modeling: Confusions and challenges. Ecology and Evolution. 2019;9: 10365–10376.

doi:10.1002/ece3.5555

47.   Maxent Version 3.4.1 Download Site. [cited 22 May 2020]. Available:

https://biodiversityinformatics.amnh.org/open_source/maxent/

48.   R: The R Project for Statistical Computing. [cited 22 May 2020]. Available:

https://www.r-project.org/

49.   Muscarella R, Galante PJ, Soley-Guardia M, Boria RA, Kass JM, Anderson MU and

RP. ENMeval: Automated Runs and Evaluations of Ecological Niche Models. 2018.

Available: https://CRAN.R-project.org/package=ENMeval

50.   RStudio | Open source & professional software for data science teams. [cited 27 May

2020]. Available: https://rstudio.com/

51.   Warren DL, Glor RE, Turelli M. ENMTools: a toolbox for comparative studies of

environmental niche models. Ecography. 2010 [cited 27 Mar 2020]. doi:10.1111/j.1600-

0587.2009.06142.x

52.   Phillips SJ, Dudík M, Elith J, Graham CH, Lehmann A, Leathwick J, et al. Sample

selection bias and presence-only distribution models: implications for background and

pseudo-absence data. Ecological applications. 2009;19: 181–197.

53.   Fourcade Y, Engler JO, Rödder D, Secondi J. Mapping Species Distributions with

MAXENT Using a Geographically Biased Sample of Presence Data: A Performance

Assessment of Methods for Correcting Sampling Bias. Valentine JF, editor. PLoS ONE.

2014;9: e97122. doi:10.1371/journal.pone.0097122

54.   Schnase JL, Grant WE, Maxwell TC, Leggett JJ. Time and energy budgets of Cassin's

sparrow (Aimophila cassinii) during the breeding season: evaluation through modelling.

Ecological Modelling. 1991;55: 285–319.

55.   (Peucaea cassinii) - Species Map - eBird. 2020 [cited 31 May 2020]. Available:

https://ebird.org/map/casspa

56.   Fielding AH, Bell JF. A review of methods for the assessment of prediction errors in

conservation presence/absence models. Environmental Conservation. 1997;24: 38–49.

doi:10.1017/S0376892997000088

57.   H. Akaike. A new look at the statistical model identification. IEEE Transactions on

Automatic Control. 1974;19: 716–723. doi:10.1109/TAC.1974.1100705

58.   Warren DL, Glor RE, Turelli M. Environmental niche equivalency versus conservatism:

quantitative approaches to niche evolution. Evolution. 2008;62: 2868–2883.

doi:10.1111/j.1558-5646.2008.00482.x

59.   Schoener TW. The Anolis Lizards of Bimini: Resource Partitioning in a Complex

Fauna. Ecology. 1968;49: 704–726. doi:10.2307/1935534

60.   Fink D, Auer T, Johnston A, Strimas-Mackey M, Robinson O, Ligocki S, et al. Cassin's

Sparrow - Abundance map - eBird Status and Trends. In: eBird Status and Trends, Data

Version: 2018; Released: 2020 [Internet]. 2020 [cited 5 Oct 2020]. Available:

https://ebird.org/ebird/science/status-and-trends/casspa/abundance-map

61.   Phillips SJ, Dudík M, Schapire RE. Maxent software for modeling species niches and

distributions (Version 3.4.1). 2020. Available: url:

http://biodiversityinformatics.amnh.org/open_source/maxent/

62.   Tang Y, Winkler JA, Viña A, Liu J, Zhang Y, Zhang X, et al. Pearson pairwise

correlation matrix between the bioclimatic variables. 2018.

doi:10.1371/journal.pone.0189496.g003

63.   O'Donnell MS, Ignizio D a. Bioclimatic Predictors for Supporting Ecological

Applications in the Conterminous United States. Reston, VA: US Geological Survey;

2012 p. 10. Report No.: 691. Available: https://pubs.usgs.gov/ds/691/

64.   Dormann CF, Elith J, Bacher S, Buchmann C, Carl G, Carré G, et al. Collinearity: a

review of methods to deal with it and a simulation study evaluating their performance.

Ecography. 2013;36: 27–46. doi:10.1111/j.1600-0587.2012.07348.x