Influence of local landscape and time of year on bat-road collision risks

Charlotte Roemer[1, 2, 3], Aurélie Coulon[1, 3] Thierry Disca[2] & Yves Bas[1, 3]

[1] Centre d'Ecologie et des Sciences de la Conservation (CESCO), Muséum national d'Histoire

naturelle, Centre National de la Recherche Scientifique, Sorbonne Université, CP 135, 57 rue Cuvier

75005 Paris, France

[2] Biotope, Mèze, France

[3] CEFE, Univ Montpellier, CNRS, Univ Paul Valéry Montpellier 3, EPHE, IRD, Montpellier, France

Corresponding author:

Charlotte Roemer

22 Boulevard Maréchal Foch, BP 58 - 34140 Mèze, France

+33 (0)4 67 18 46 20

charlotte.roemer@lilo.org

Abstract

Roads impact bat populations through habitat loss and collisions. High quality habitats particularly

increase bat mortalities at roads, yet many questions remain concerning how local landscape features

may influence bat behaviour and lead to high collision risks. Moreover, no study provides an

understanding of the extent to which mortalities result from bat density on one hand and from bat

flight behaviour on the other hand, which is required when designing risk mitigation measures.

In this study, we recorded bat activity and characterised flight behaviour with three variables: position

at collision risk, vehicle avoidance, and flight path orientation, using acoustic flight path tracking at 66

study sites in the Mediterranean region during two full nights. We modelled the effect of the local

landscape, i.e. in a radius of 30 m around the road (vegetation height, distance, density and

orientation), road features (road width, traffic volume) and time of the year on eleven species or

species groups. We built models for each conditional probability of the road collision risk (i.e. species

density, presence in the zone at risk, vehicle avoidance) and multiplied their estimates to calculate the

overall collision risk.

Our results show that the local landscape had different effects on bat density and presence at collision

risk. Increasing distance to trees and decreasing tree height were associated with a decrease in bat

density at roads. Forests were the local landscapes where bats flew more often in the zone at collision

risk. The overall collision risk was higher either at forests or at tree rows perpendicular to the road

depending on species. Contrary to common preconceptions, mid-range echolocators seemed to be

generally more at collision risk than short-range or long-range echolocators. In addition, collision risk

was maximal in summer or autumn for most species. Finally, bats mainly followed the road axis even

when trees were present on both sides or absent.

Our results contribute to a better understanding of bat movements in different local environments at

the scale where they directly sense their surroundings with echolocation calls. Disentangling bat

density from flight behaviour allowed us to better understand the temporal and spatial contributors of

roadkills, and to provide guidance for road impact assessment studies.

Introduction

Highways and main or secondary roads cover important surfaces of industrialised countries worldwide

while road construction and traffic density rise continuously (Ibisch et al., 2016; van der Ree et al.,

2015a). Both networks lead to concerning impacts on wildlife, namely death by collision, loss of

habitat amount and quality, population fragmentation, which in turn lead to negative impacts on

population survival in numerous taxa (Rytwinski and Fahrig, 2015).

To explain the direct ecological impact of roads, i.e. mortality by collision, several studies have

investigated the role of road and land features on roadkill occurrence. They showed for example that

increasing road width, traffic and/or speed limit enhances collisions in large mammals (Nelli et al.,

2018; Neumann et al., 2012; Seiler, 2005; Valero et al., 2015) but the results concerning the role of

traffic or speed limit on roadkills were contrasted for other vertebrate taxa (Clevenger et al., 2003;

D'Amico et al., 2015; Mazerolle, 2004). Studies on a variety of animal groups also found that habitats

favourable for species foraging or movement, described at the home-range scale (e.g. presence or

absence of woodland, cropland, wetland …), are more often associated with the occurrence of road-

kills (Grilo et al., 2016; Gunson et al., 2011; Malo et al., 2004).

Very few studies have investigated the effect of local landscape features (i.e. within the few meters on

either side of the road) on collisions. However, when mitigation measures are recommended, they

often deal with vegetation structure at this local scale (van der Ree et al., 2015b). Indeed, it is likely

that the landscape in the immediate vicinity of roads affects animal movement trajectories – and, as a

result, collisions risks. In ungulates, Meisingset et al. (2014) found that increasing road edge clearance

decreased the rate of collisions. However, as the authors suggest, this is probably partly a driver effect

since drivers benefitting from a better visibility will in all likelihood have more time to avoid

collisions. Driver avoidance of collision is probably less important for smaller animals. Large animals

that may be avoided by drivers represent only a very small percentage of the species impacted by

collisions (D'Amico et al., 2015; Rytwinski and Fahrig, 2015) and the effects of local landscape are

likely to be species dependent, but knowledge is very scarce at the species level.

The movement of aerial animals is expected to be particularly conditioned by height, density and

spatial arrangement of three-dimensional structures (Brigham et al., 1997; Norberg, 1994, 1986). For

example in birds, gaps in vegetation are an important factor of road collisions (Lin, 2016; Orłowski,

2008). Among aerial animals impacted by road collisions, bats are long-lived mammals with a low

reproductive rate, having one offspring – exceptionally two – per year (Dietz et al., 2009).

Additionally, temperate bats have suffered from an important decline of their populations in the

second half of the twentieth century, which translates into a poor conservation status today (Van der

Meij et al., 2015), and North-American bats have experienced dramatic declines due to the white-nose

syndrome, a fungal disease (Langwig et al., 2015). For these reasons, even moderate increases in

mortality rates may represent a serious threat to their survival. As a result, all European bats are now

under strict protection (*Council Directive 92/43/EEC on the conservation of natural habitats and of*

wild fauna and flora, 1992). A good understanding of the mechanisms leading to collisions between

road vehicles and bats is therefore necessary to efficiently mitigate them (Fensome and Mathews,

2016).

At the home range scale, several studies showed that bat collisions are concentrated in habitats

qualified as favourable for foraging and commuting (e.g. water bodies, forests and riparian habitats)

(Gaisler et al., 2009; Lesiński, 2007; Medinas et al., 2019, 2013). At the local scale, it is suspected that

the orientation of tree lines and vegetation structure (i.e. height, density and distance from road edge)

direct bat movement, and are consequently major factors of road collision risks (Fensome and

Mathews, 2016). The influence of vegetation structure on bat activity has been relatively well studied

in the literature (Kelm et al., 2014; Pourshoushtari et al., 2018; Toffoli, 2016; Verboom and Spoelstra,

1999). However, the influence of local landscape on bat movement has been almost exclusively

described from visual observations (Arthur and Lemaire, 2015; Dietz et al., 2009). Some studies at

roads suggest that increasing distance to surrounding trees decreases bat crossing frequency, and that

increasing tree height elevates bat crossing height (hence reducing the risk of collisions with vehicles)

(Abbott, 2012; Bennett and Zurcher, 2013; Russell et al., 2009). But small sample sizes and poor

taxonomic resolution limit the generalisation of these results. Moreover, Bennett and Zurcher (2013)

considered bat trajectories initially directed perpendicular to the road and determined that vegetation

structure and vehicle presence influenced bat decision to cross the road or to fly away. But they did

not take into account bats flying parallel to the road axis, although this behaviour may be a

determinant factor of collisions, because in this situation bats spend much more time in the zone at

collision risk.

Road collision risks in a species depend on (1) its local density, (2) the proportion of time spent in the

zone at collision risk and (3) the simultaneous presence of bats and vehicles in the zone at collision

risk (Jaeger et al., 2005; Zimmermann Teixeira et al., 2017). It is therefore necessary to take each of

these variables into account when investigating road collisions. Indeed, when comparing two different

road locations within different landscape features, a higher bat acoustic activity (used as a proxy of bat

density) does not necessarily lead to a higher proportion of flights at collision risk for all species (see

Abbott et al., 2012). A measure of per capita mortality risk is essential to avoid wrong

recommendations for the siting of mitigation measures (Zimmermann Teixeira et al., 2017).

Our study aimed at (1) assessing the effects of the characteristics of the local landscape on bat activity

and movement behaviour and consequently on road collision risks and (2) disentangling the roles of

density and movement behaviour in collision risks. In order to provide species-specific answers, our

analyses were mostly performed at the species level, using the guild level only for species with small

sample sizes. We used acoustic monitoring to detect bat passes and car passes, and acoustic flight path

tracking (AFPT) to locate bat echolocation calls in three dimensions. This method allows

reconstructing three-dimensional flight paths, and then model separately: (1) bat species density, (2) a

probability of flight at collision risk, (3) a probability of car avoidance and (4) a probability to fly

parallel or perpendicular to the road.

Material and Methods

Study sites and description of the local landscape

The study took place in 2016 and 2017 in the French Mediterranean lowland region. This area is

composed of a mosaic of garrigues, cultivated areas (often vines), young forests of oaks and pines, and

urban areas consisting of small traditional villages and large cities with extensive conurbation. The

national and departmental road network is built around one main highway going from the east to the

west and following the French southern coastline (Figure 1). Four other highways link this main

highway to the inner lands through large natural valleys.

Bat behaviour was recorded at 66 sites (Figure 1, supplementary table 1) at national or departmental

roads, during two to five nights per site (mean = 2.6 +/- SD 0.9). The local landscape was described

within a radius of 30 m, equivalent to the sonar range of mid-range echolocating bats (Holderied and

von Helversen, 2003). This scale was chosen under the assumption that bats adapt their flight

movements according to the environment perceived acoustically. At most study sites, landscape

description would have been similar with a 100 m radius, which corresponds to the sonar range of

long-range echolocators (Holderied and von Helversen, 2003). Study sites were chosen so as to reach

a balanced representation of six major types of road landscapes in the study area: simple parallel tree

rows (SPT), double parallel tree rows (DPT), perpendicular tree rows (PT), forests (F), forest edges

(FE) and no vegetation taller than 1.5 m (NV) (Table 1, Figure 1, Figure A 1). Tree species were very

often associated with a type of landscape: 70 % of SPT and DPT were plane trees (*Platanus sp.*), and

more rarely olive trees (*Olea sp.*), Celtis (*Celtis sp.*), Aleppo pines (*Pinus halepensis*) or mulberries

(*Morus sp.*). F and FE consisted in 80 % oaks (*Quercus ilex*, *Q. pubescens*, *Q. coccifera* and/or *Q.*

suber), and 20 % Aleppo pines (*Pinus halepensis*). PT were a mix of Mediterranean riparian species

(mostly *Fraxinus* sp., *Populus* sp., *Quercus* sp. and/or *Arundo donax*) typically associated with

temporary watercourses. The category NV consisted in land either occupied by vines, wheat, recently

ploughed or left uncultivated. Pastureland was almost non-existent. Parallel tree rows had gaps of

about 10 to 20 m between trees while the other types of vegetated landscapes had little or no gaps.

Figure 1: Geographical distribution of the 66 sampling sites in the secondary road network of the
French Mediterranean lowland region. The symbols of the sampling sites represent the type of local
landscape they belong to. Top right: French biogeographical regions (white: Mediterranean; light grey:
Atlantic; dark grey: Continental; black: Alpine). (Road network source: ROUTE500 from the Institut
National de l'Information Géographique et Forestière, 2017).

Table 1: Description of variables used to model bat density, bat position at collision risk, and flight path
orientation. Percentage of night, time lag and time lag direction were only used in the last two models.
154

Variable		Description
Landscape type	*Simple parallel tree row (SPT)*	One row of trees arranged linearly and parallel to the road.
	Double parallel tree row (DPT)	Two rows of trees arranged linearly at each side and parallel to the road.
	Perpendicular tree row (PT)	One row of trees at each side and perpendicular to the road. Associated with small seasonal streams where water was either absent at the time of monitoring, or likely not accessible for bats due to dense tree cover.
	Forest (F)	Dense tree patch at each side of the road.
	Forest edge (FE)	Dense tree patch at one side of the road.
	No vegetation (NV)	No vegetation taller than 1.5 m at each side of the road.
Road width		Distance between both outer edge lines of the roadway.
Traffic volume		Mean number of vehicles per night.
Distance to tree foliage		Mean distance between road outer edge line and tree foliage over all present trees. If foliage runs over the road, distance is negative.
Tree height		Mean tree height from ground to canopy over all present trees.

All sites were situated in lowlands, on two-lane asphalt roads wide of 4 to 8 m, on straight portions,

where vehicles were allowed to drive up to 90 km/h. Several features were avoided: (1) artificial street

lights and urban areas (the smallest distance to lit streets and urban areas was 300 m) (2) important

three-dimensional structures, such as electric poles (3) highways (the smallest distance to a highway

was 1.1 km) (4) water bodies or wetlands other than the small streams sampled in the PT category (the

shortest distance to water was 100 m) and (5) sparse trees within the landscape matrix. The minimum

distance between study sites was 500 m. Monitoring was performed exclusively during nights with

optimal weather conditions for bat activity (mean temperature = 20.6 +/- 6.5 °C; mean wind speed =

7.5 +/- 8 km/h; mean cumulated rain per night = 0.2 +/- 1.3 mm) but the percentage of visible moon

was not a criterion we used (mean = 49.2 +/- 35.8 %).

Four secondary landscape characteristics likely to affect flight behaviour were measured at each study

site: road width, traffic volume, distance between road and tree foliage, and tree height (Table 1).

Measurements were made with a laser telemeter. Traffic volume was calculated using the

TADARIDA-L software (Bas et al., 2017) to identify and count vehicle passes. Sound event detection

was done in the low frequency mode, and any acoustic sequence of 5 s or less that contained an

uninterrupted sound event with a duration superior to 1.2 s was counted as one vehicle pass, even if

several vehicles followed each other very closely. This threshold was chosen based on a verification of

false and true positives and negatives on 100 random sound sequences from different study sites

stratified by sound duration (unpublished data).

Bat acoustic monitoring

On each site, two pairs of microphones (either SMX-US or SMX-U1 (Wildlife Acoustics, USA), or

BMX-US (Biotope, France)) were plugged into two SM2BATs or SM3BATs (Wildlife Acoustics,

USA), each connected to a GPS unit used to timely synchronise recorders. Microphones were either

mounted on wooden poles (at a maximum height of 4 m) or attached to vegetation (at a minimum

height of 20 cm) (Figure A 1). Microphone pairs were installed on each side of the road (0.5 – 4 m

distance from the road edge) in arbitrarily shaped non-coplanar microphone arrays. Depending on the

study site, minimum distance between microphones was 8.1 (+/- SD 1.6) m and maximum distance

was 16.8 (+/- SD 2.7) m. Recorders were programmed to start each day 30 min before sunset and to

stop 30 min after sunrise. Gain was set at 36 dB for SMX-US and BMX-US microphones, or at 0 dB

for SMX-U1 microphones. Sampling rate was set at 192 kHz, trigger at 6 dB above background noise

and trigger window at 2.5 sec.

Species identification was performed based on echolocation calls, which carry enough information to

allow the identification of the majority of European bat species, depending on the quality and the

context of the recordings (Barataud, 2015). We used the SonoChiro software (Biotope/MNHN,

France) to automatically sort sequences by species, and then verified most of sequences manually on

Syrinx (John Burt, USA) (except for sequences classified as *Pipistrellus* which are too numerous for a

detailed verification, and because SonoChiro has a very low error rate for *P. pipistrellus* and *P.*

pygmaeus in the Mediterranean region according to our experience). *Plecotus* species were grouped in

Plecotus sp., *Myotis blythii* and *M. myotis* were grouped in *M. blythii/myotis*, and *Pipistrellus kuhlii*

and *P. nathusii* were grouped in *Pipistrellus kuhlii/nathusii*. Acoustic sequences that could not be

identified to the species level or groups were left unidentified (0.15 % of all bat passes). From our

knowledge of bat assemblages of France Mediterranean lowlands (unpublished mist-netting data), we

expect the last group to contain a very large majority of *P. kuhlii*, and the *Plecotus* group to contain a

very large majority of *P. austriacus*.

Three-dimensional positioning of bat calls

Bat three-dimensional flight paths were generated from the three-dimensional source location of

echolocation calls recorded on the four microphones. After species acoustic identification, call

location was achieved by (1) measuring time of arrival differences (TOAD) of each call between pairs

of microphones and (2) deducing the coordinates of the sound source by comparing those field TOAD

($TOAD_F$) with theoretical TOAD ($TOAD_T$). Indeed, since the speed of sound in the air is known (here

we approximated it to 340 m/s), TOADs of a sound source recorded by at least four microphones can

be used to calculate the location of the source (see Koblitz, 2018).

$TOAD_F$ were calculated by measuring the starting time of bat calls using the SonoChiro software

(Biotope/MNHN, France). Call association between pairs of microphones was achieved using the R (R

Core Team, 2014) function find.matches of the Hmisc package (Harrell, 2018). Because there are four

microphones, six $TOAD_F$ per call are calculated. $TOAD_T$ were calculated for each simulated point of a

matrix of 40 x 40 x 40 m with a one-meter resolution and centred around the centroid of the 4

microphones, inputting the same microphone configurations as the ones used in the field. The

dimensions of this matrix were chosen according to the spatial range of our equipment (i.e. maximal

distance of detection of a bat position) for the location of middle-range echolocators (e.g. *Pipistrellus*

pipistrellus). This range is dependent on the acoustic range of the individual recorded and the position

of the individual in relation to the microphones, i.e. accuracy is maximal at the centre of the device

and minimal at the far edges.

The position of the bat was deduced from the comparison of the differences between the six $TOAD_F$

and the six $TOAD_T$ using the R (R Core Team, 2014) function find.matches of the Hmisc package

(Harrell, 2018). The closest TOAD match was selected as a candidate bat position. During test

calibrations of our setting with different microphone configurations, we calculated that $TOAD_F$

resulting in a position more than 10 m away from the centroid of the microphones had a difference

with the real position larger than one meter. Imprecise positions were systematically reconstructed

away from the centroid, which means for example that a bat flying in reality at 15 m from the centroid

could be located with our device at 17 m (away from the centroid), but never at 13 m (toward the

centroid). We therefore rejected any field position found at more than 10 m from the centroid of the

microphones and did not use them for further analyses (Table **A 1**).

Grouping of calls in individual flight trajectories

Calls were then attributed a flight trajectory ID using successive filters. During the first round, a same

temporary ID was first given to all calls separated by less than two seconds and the flight speed

between the preceding and the actual call was calculated in the X and Y dimensions. Several rounds

were then run successively. At each round, to keep their temporary ID, calls had to (1) have a peak

frequency differing by less than 5 kHz from the median peak frequency of all calls within the same ID

(2) be separated by less than 2 seconds from the preceding call (3) be preceded and followed by

positions conferring a speed lower than 20 m/s (i.e. the maximum possible speed of flying bats

(Holderied and Jones, 2009; Popa-Lisseanu, 2007). Else, calls were attributed a new (unique)

temporary ID and went through a new round of filtering. Successive rounds were applied until all IDs

were stabilized. Flight trajectories with less than three calls were not considered as a full flight path

and were classified as non-located bat passes (Figure 2). R scripts and tables are available at

https://github.com/Charlotte-Roemer/bat-road-collision-risks.

Definition of collision risk and calculation of flight path orientation

Each successfully located bat position above the road and at vehicle height (< 5 m) (Berthinussen and

Altringham, 2012) was classified as 'unsafe' (Figure 2). All other successfully located positions were

classified as 'safe'. Bat calls which were not recorded by all four microphones at once – and that could

therefore not be precisely located – were assumed to be far from the microphones' centroid and thus

probably far from the road and hence also classified as 'safe'. For the same reason, positions

potentially not successfully located (> 10 m from microphones centroid) were disregarded to avoid

location errors. Since this error rate is similar across landscape types, we do not expect any resulting

bias. If any of the bat positions within a flight trajectory was unsafe, the complete flight trajectory was

classified as unsafe, else it was classified as safe. Flight trajectories with less than three calls were

assumed to be far from the microphones' centroid and thus probably far from the road and hence also

classified as 'safe' (Figure 2).

Figure 2: Process of classification of flight trajectory at collision risk. NA: non-available data. The area
where calls are accurate is 10 m around the centroid of microphones. Unsafe call: located at vehicle
height and above the road.

We then calculated the angle between the road axis and the axis of the vector linking the first to the

last position of each flight paths. Trajectories were classified in two categories: 0-45° = parallel; 45-

90° = perpendicular to the road.

Response variables

We tested how the local landscape affects the different determinants of collision risks, building one

model for each of them: (1) local species density (2) the proportion of flights in the zone at risk (3)

vehicle avoidance and (4) flight path orientations (Figure 3). To summarise results, the estimates of

the first three models (i.e. quantitative models) were multiplied to obtain as a product (5) the number

of bat passes at collision risk per night, depending on the characteristics of the local landscape.

Figure 3: Successive steps in modelling of bat density and flight behaviour in function of landscape
variables. If the flight path is at risk, position at risk = yes. If the time lag between the bat pass and the
vehicle pass is greater than 10 s, vehicle avoidance = yes.
272

(1) Local bat density

The density of the most common species in our dataset (i.e. occurrence per night larger than 30 % and

occurrence per site larger than 50 %) was modelled as a negative binomial distribution. The response

variable was the median number (among the four microphones) of five second intervals per night

within which a species was identified. This acoustic activity was then used as a proxy of bat density

(number of acoustic sequences per night within the acoustic range of the setting) (Froidevaux et al.,

2017).

(2) Probability of bats to fly through the zone at risk

The probability of trajectories to be in the zone at risk (i.e. at vehicle height) was modelled using the

risk status of each trajectory as the binomial response variable (0 = safe; 1 = unsafe) (Figure 2).

(3) Bat avoidance of vehicles

The probability of bats to fly through the zone at risk gives a spatial evaluation of risk. To make a

more precise risk assessment, the vehicle avoidance by bats (i.e. temporal evaluation of risk) was also

modelled. For bat flight trajectories at risk only, the probability of bats to avoid vehicles was modelled

using bat avoidance behaviour as the binomial response variable (1 = vehicle avoidance; 0 = no

vehicle avoidance). To do this, the time lag between an acoustic sequence containing a bat and the

closest sequence containing a vehicle pass was calculated using the function find.matches of the

Hmisc package (Harrell, 2018). If the time lag was lower than 10 s, we considered that there was no

vehicle avoidance (0). If the time lag was higher than 10 s, we considered that there was vehicle

avoidance (1). In fact, it might be argued that very long time lags between a vehicle pass and a bat

pass cannot be considered as true vehicle avoidance, but rather coincidental bat absence on the road in

the simultaneous absence of a vehicle. However, a relatively higher proportion of such bat passes

could be interpreted as a higher probability that bats avoid vehicles, thus we used this metric as a

proxy for vehicle avoidance.

(4) Flight path orientation

For all bat flight trajectories, the proportion of flight paths parallel to the road axis was modelled using

flight orientation as a binomial response variable (0 = perpendicular; 1 = parallel).

(5) Number of bat passes at collision risk per night

If an explanatory variable was selected in several models, then each of those models gives a partial

evaluation of bat collision risks at roads. In fact, all quantitative models succeeding the density model

can be interpreted as conditional probabilities that an individual is at collision risk. Thus, a proxy for

fatality rate at a road section can be computed by the multiplication of all outputs of the quantitative

models:

Eqn 1: $Expected\ value\ of\ collision\ risk = E_1 \times E_2 \times E_3,$

where:

E_1 = Prediction of the number of bats present on site per night

E_2 = Prediction of the probability that a detected bat flies in the volume at collision risk

E_3 = Prediction of the probability that a bat in the volume at risk does not avoid vehicles

To estimate the confidence intervals of this product, we needed a large number of responses for each

value along the gradient of each predictor for each response variable. For this, we first simulated

responses according to model estimates and their standard error using the rnorm function (R Core

Team, 2014): 20,000 replicates for each of 60 values along a gradient for a given predictor. For a

given predictor, if one E_1, E_2 or E_3 was missing, meaning that the predictor was not selected in one of

the models, we used the mean value of the original observations instead. If two or all three of E_1, E_2 or

E_3 were missing for a given predictor, we did not compute their product.

Model selection

We used the R (R Core Team, 2014) package glmmTMB (Brooks et al., 2017) to model each response

variable in generalised linear mixed models (GLMM). When sample size of a given species was too

small, we did not model the species response. In addition, three bat guilds were created based on the

adaptation of species to clutter of the environment, which is strongly linked to sonar features (Aldridge

and Rautenbach, 1987; Denzinger et al., 2018). Species were thereby split into the guilds "short-range

echolocator" (SRE), "mid-range echolocator" (MRE) or "long-range echolocator" (LRE) according to

the definition of Frey-Ehrenbold et al. (2013) (see Table A 1 for complete list).

All descriptive variables were normalised if necessary and scaled to follow a normal distribution and

to compare their effects. Variables considered for fixed effects were landscape type, road width, traffic

volume, distance to tree foliage and tree height (Table 1). We first calculated the correlation

coefficients between predictors using the corrplot function of the stats package in the R program (R

Core Team, 2014). Tree height and distance to tree foliage were correlated ($r = -0.57$), as well as road

width and traffic volume ($r = 0.64$). We excluded road width for further analysis and created the

possibility to select either tree height or distance to tree foliage (but not both) during stepwise model

selection (see next paragraph). Candidate predictors were also included in simple interactions with

each other. In addition, Julian day was included as a fixed quadratic effect to account for seasonal

variations in bat density and flight behaviour. Site ID was introduced as a random effect.

An upward stepwise model selection was performed to select the relevant variables (except for Julian

day which was part of the null model). We operated an upward model selection because the full model

led to overfitting for species with a low occurrence. At each step of model selection, the VIF (Variance

inflation factor), which quantifies the degree of multicollinearity in least square regression analyses,

was calculated. If any of the selected variables had a $VIF > 3$ (Heiberger and Holland, 2004; Zuur et

al., 2010), the model was not considered as a candidate model. At each step of model selection, the

model with the smallest Akaike's information criterion for small sample sizes (AICc) was considered.

This model was retained and selected if its AICc was at least inferior by two points to the AICc of the

best model of the previous step (supporting that the newly added parameter is informative) (Arnold,

2010).

For each retained model, we checked the uniformity of the residuals using the DHARMa package

(Hartig, 2018). Goodness of fit, autocorrelation, overdispersion and zero-inflation (for density data

only) were checked and revealed no problematic situation.

Results

In total, 122,294 bat passes were recorded and identified at the group or species level, from which

30,954 successful flight trajectories could be located (Table A 1). Because of technical problems on

two study sites (one of the two recorders was once destroyed by a rotary flail and once displaced by

someone), flight path tracking could not be performed and these sites were used for modelling bat

density only. The density of nine species and three species groups (*Pipistrellus kuhlii/nathusii*,

Plecotus sp. and *Myotis blythii/myotis*) could be modelled, but their flight behaviour (i.e. presence at

risk, vehicle avoidance and flight path orientation) could not be modelled for all of them, due to the

lack of data.

Models showed no convergence problem during selection, except for *E. serotinus* (model position in

the zone at risk, for interactions), *M. daubentonii* (model vehicle avoidance, for landscape type), *H.*

savii (model trajectory orientation, for landscape type) and *M.myotis/blythii* (model vehicle avoidance,

for landscape type; model trajectory orientation, for landscape type).

Figure 4: Predicted effects of landscape type on density (left), proportion of trajectories at risk (middle) and predicted proportion of vehicle avoidance (trajectories at risk and at more than 10 s from a vehicle pass) (right). 95% confidence intervals are shown. Only the effects present in the final models are shown. Bottom figures represent landscape type viewed from the top (road in light grey and trees in black). LRE: long-range echolocators. MRE: mid-range echolocators.

 Model 1 – Bat density

Landscape type had an important influence on bat density (Table 2). It was selected in the model of

four species. Density was much higher at PT for *Pipistrellus* species and was higher at forested

landscapes for *H. savii* (Figure 4). Increasing distances to tree foliage were associated with a decrease

in bat density for five species (*M. daubentonii, P. pipistrellus, P. pygmaeus, M. schreibersii* and *N.*

leisleri) and for the MRE guild, while it was associated with an increase for *Plecotus* species (Figure

5). Increasing tree height was associated with an increase in the density of *M. blythii/myotis* and of the

LRE guild (Table 2). With an increasing traffic volume, the density of *Plecotus sp.* and of the SRE

guild decreased (Table 2). Throughout the year, species density showed a typical peak in mid-summer,

except for *P. pygmaeus, M. schreibersii, Plecotus* sp. and *N. leisleri*, that were more active in the

autumn (Figure A 2).

Distance to tree foliage (m)

376
377   Figure 5: Predicted effect of distance to foliage on bat density for a selection of four species. 95%
378   confidence intervals are shown. Ticks in x axis represent sampled values. Negative values mean that
379   foliage was running over the road.
380

Model 2 – Bat presence in the zone at risk

Landscape type also greatly influenced the proportion of bat positions in the zone at risk. It was

selected in five of the ten species-specific models, and in two of the guild models. The proportion of

positions at risk was generally higher in forests and lower without trees (Figure 4 and Table 3).

An increasing distance to tree foliage was associated with an increase in the presence at risk for *E.*

serotinus (Table 3). Increasing traffic density was associated with a decrease in the proportion of

flights in the zone at risk for *P. kuhlii/nathusii* (Table 3). Throughout the year, the different species

displayed quite different patterns in presence at risk, but all three guilds showed a tendency for a

higher proportion of flights in the zone at risk toward the end of the year (Figure A 2).

Model 3 – Bat avoidance of vehicles for trajectories in the zone at risk

Landscape type was only selected in the model for the LRE (Table 4). For this guild, simple parallel

tree rows were associated with higher vehicle avoidance than double parallel tree rows (Figure 4).

Increasing tree height was associated with a decrease in vehicle avoidance in *M. daubentonii, P.*

pipistrellus, P. pygmaeus and the MRE guild (Table 4). An increasing traffic density was associated to

less vehicle avoidance for all *Pipistrellus* species and for *M. myotis/blythii* (Table 4). The SRE guild

had a higher avoidance rate than the MRE guild (Figure 6). Season had different effects on the

avoidance of vehicles according to species and guilds (Table 4, Figure A 2).

399
400   Figure 6: Predicted effect of traffic volume on proportion of vehicles avoided (trajectories positioned at
401   risk and at more than 10 s from a vehicle pass). 95% confidence intervals are shown. Ticks in x axis
402   represent sampled values (bottom = MRE; top = SRE). SRE: short-range echolocators. MRE: mid-
403   range echolocators.
404

Model 4 – Orientation of flight trajectories

The large majority of flight paths followed the road axis in all landscape types (results not shown).

Landscape type, distance to tree foliage, and tree height were not selected to explain trajectory

orientation (Table 5). Nonetheless, an increasing traffic volume was associated to a larger proportion

of trajectories parallel to the road axis in *P. pipistrellus* (Table 5). Season had a very weak, or even no

effect on the proportion of trajectories parallel to the road (Figure A 2).

Product: number of bat passes at collision risk per night

There was only a small selection of species for which a same variable had an effect on at least two

aspects of the collision risk (i.e. on bat density, bat presence in the zone a risk, or bat avoidance of

vehicles). These cases are all described in this section. An increasing traffic was associated with a

tendency of an increasing number of bat passes at collision risk for *P. kuhlii/nathusii* and the SRE

guild (Figure A 3). The number of bat passes at collision risk was higher at perpendicular tree rows for

P. pipistrellus but higher at forests and forest edges for *H. savii* (Figure A 3). The number of bat

passes at collision risk was higher in summer for *E. serotinus*, *H. savii*, *P. kuhlii/nathusii* and *P.*

pipistrellus, while it was higher in autumn for *M. schreibersii*, *M. daubentonii*, *P. pygmaeus* and

Plecotus sp, and higher in spring for *M. myotis/blythii* (Figure A 2).

Table 2: Summarised statistical results of the negative binomial distributed GLMM for the density of each species. $\hat{\beta}$ = estimate. SE = standard error. p = significance of p value. Species names are given with the three first letters of the species and genera. Dist.tree = distance to tree foliage. Tree.H = tree height. DPT = double parallel tree rows. F = forest. FE = forest edge. PT = perpendicular tree rows. SPT = simple parallel tree rows. Intercept is for NV (no vegetation) landscape type. LRE: long-range echolocators. MRE: mid-range echolocators. SRE: short-range echolocators. There is a total of 163 nights of observations.

Variable	Eptser			Hypsav			Minsch			Myodau			Myoema			Myobly/myo			Nyclei			Pipkuh/nat			Pippip			Pippyg			Plesp		
Type	$\hat{\beta}$	SE	p	$\hat{\beta}$	SE	p	$\hat{\beta}$	SE	p	$\hat{\beta}$	SE	p	$\hat{\beta}$	SE	p	$\hat{\beta}$	SE	p	$\hat{\beta}$	SE	p	$\hat{\beta}$	SE	p	$\hat{\beta}$	SE	p	$\hat{\beta}$	SE	p	$\hat{\beta}$	SE	p
Intercept	0.64	0.48		2.64	0.50	***	-0.17	0.56		-0.76	0.41	.	0.00	0.28		0.32	0.39		2.24	0.22	***	5.50	0.19	***	4.67	0.29	***	3.03	0.44	***	0.26	0.23	
Dist.tree							-0.58	0.23	*	-0.92	0.29	**							-0.91	0.17	***				-0.29	0.13	*	-0.80	0.20	***	0.42	0.16	*
Tree.H																1.21	0.33	***															
Traffic																															-0.44	0.17	*
DPT				-0.20	0.62		1.04	0.72																	0.44	0.39		0.62	0.59				
F				0.63	0.64		3.34	0.71	***																0.82	0.38	*	0.06	0.58				
FE				0.76	0.63		3.34	0.72	***																0.67	0.39	.	0.14	0.59				
PT				-1.42	0.67	*	3.26	0.72	***																2.22	0.39	***	1.87	0.59	**			
SPT				-1.35	0.67	*	2.15	0.75	**																-0.16	0.40		-0.11	0.61				
Julian Day	-0.19	0.30		-0.21	0.18		0.42	0.15	**	1.17	0.32	***	-0.30	0.18		-0.85	0.28	**	0.40	0.14	**	-0.09	0.13		0.40	0.10	***	0.74	0.14	***	0.53	0.15	***
Julian Day²	-0.96	0.30	**	-0.99	0.19	***	0.36	0.14	**	-0.58	0.26	*	-0.44	0.19	*	-0.75	0.23	**	0.28	0.14	*	-0.50	0.12	***	-0.58	0.09	***	0.07	0.13		0.41	0.14	**

P < 0.1 = .
P < 0.05 = *
P < 0.01 = **
P < 0.001 = ***

Table 2 (continued)

Variable	SRE			MRE			LRE		
Type	$\hat{\beta}$	SE	p	$\hat{\beta}$	SE	p	$\hat{\beta}$	SE	p
Intercept	3.09	0.16	***	5.56	0.26	***	3.17	0.19	***
Dist.tree				-0.31	0.12	**			
Tree.H							0.71	0.15	***
Traffic	-0.36	0.14	**						
DPT				0.62	0.35	.			
F				1.35	0.34	***			
FE				1.01	0.35	**			
PT				1.72	0.35	***			

Variable	SRE			MRE			LRE		
Type	$\hat{\beta}$	SE	p	$\hat{\beta}$	SE	p	$\hat{\beta}$	SE	p
SPT				0.53	0.36				
Julian Day	0.19	0.10	.	0.18	0.09	*	0.30	0.13	*
Julian Day²	-0.29	0.10	**	-0.30	0.08	***	-0.05	0.12	

433 Table 3: Summarised statistical results of the binomial distributed GLMM for the proportion of trajectories in the zone at risk for each species. $\hat{\beta}$ = estimate. SE = standard
434 error. p = significance of p value. Species names are given with the three first letters of the species and genera. Dist.tree = distance to tree foliage. Tree.H = tree height. DPT
435 = double parallel tree rows. NV = no vegetation. FE = forest Edge. PT = perpendicular tree rows. SPT = simple parallel tree rows. Intercept is for F (forest) landscape type.
436 LRE: long-range echolocators. MRE: mid-range echolocators. SRE: short-range echolocators. Interactions are shown with a colon.
437

Variable	Eptser			Hypsav			Minsch			Myodau			Myobly/myo			Nyclei			Pipkuh/nat			Pippip			Pippyg			Plesp			SRE			MRE			LRE		
N observations	876			4509			10280			401			1629			8401			45475			40279			19446			743			5744			119991			10025		
Type	$\hat{\beta}$	SE	p	$\hat{\beta}$	SE	p	$\hat{\beta}$	SE	p	$\hat{\beta}$	SE	p	$\hat{\beta}$	SE	p	$\hat{\beta}$	SE	p	$\hat{\beta}$	SE	p	$\hat{\beta}$	SE	p	$\hat{\beta}$	SE	p	$\hat{\beta}$	SE	p	$\hat{\beta}$	SE	p	$\hat{\beta}$	SE	p	$\hat{\beta}$	SE	p
Intercept	-4.77	1.5	**	-2.53	0.46	***	-3.6	0.31	***	-1.95	0.18	***	-3.7	0.42	***	-3.21	0.46	***	-1.2	0.29	***	-1.7	0.28	***	-2.91	0.17	***	-3.69	0.54	***	-3.51	0.36	***	-1.55	0.32	***	-3.48	0.73	***
Dist.tree	1.09	0.31	***																																				
Tree.H																																							
Traffic																			-0.31	0.14	*																		
DPT	-0.4	1.31		-2.57	0.79	**										-1.18	0.55	*	-1.92	0.4	***	-1.24	0.37	***										-1.66	0.36	***	-1.01	0.52	.
FE	-1.31	1.43		0.38	0.59											-1.05	0.66		-1.35	0.41	***	-0.9	0.39	*										-1.07	0.37	**	-0.75	0.59	
NV	0.18	1.46		-0.51	0.67											-2.47	0.79	**	-1.83	0.43	***	-1.5	0.42	***										-1.85	0.38	***	-2.11	0.71	**
PT	1.46	1.4		-0.49	1.28											-0.74	0.82		-1.64	0.41	***	-0.65	0.38	.										-1.24	0.37	***	0.02	0.61	
SPT	1.29	1.59		-1.09	0.84											-0.14	0.54		-1.38	0.4	***	-1.04	0.39	**										-1.4	0.36	***	-0.07	0.52	
Julian Day	0.03	0.34		-1.39	0.53	**	0.43	0.21	*	0.25	0.26		0.59	0.28	*	0.38	0.2	.	0.01	0.1		-0.01	0.08		0.23	0.13	.	0.48	0.33		0.38	0.15	*	0.11	0.06	*	0.24	0.17	
Julian Day²	1.61	0.48	***	-1	0.65		-0.37	0.15	*	-0.14	0.21		0.45	0.24	.	0	0.19		0.04	0.09		-0.09	0.09		-0.04	0.12		-0.11	0.3		-0.1	0.14		-0.05	0.06		0.13	0.17	

442 Table 4: Summarised statistical results of the binomial distributed generalised linear mixed effect models (GLMM) for the vehicle avoidance for each species. $\hat{\beta}$ = estimate. SE
443 = standard error. p = significance of p value. Species names are given with the three first letters of the species and genera. Dist.tree = distance to tree foliage. Tree.H = tree
444 height. DPT = double parallel tree rows. NV = no vegetation. FE = forest Edge. PT = perpendicular tree rows. SPT = simple parallel tree rows. Intercept is for F (Forest)
445 landscape type. LRE: long-range echolocators. MRE: mid-range echolocators. SRE: short-range echolocators.

Variable	Eptser (β̂ SE p)	Hypsav	Minsch	Myodau	Myobly/myo	Nyclei	Pipkuh/nat	Pippip	Pippyg	SRE	MRE	LRE
N observations	35	869	661	46	60	432	5155	5291	1356	213	13332	480
Intercept	-0.23 0.63	-0.25 0.27	1.32 0.30 ***	1.83 0.60 **	0.63 0.81	0.41 0.34	0.66 0.09 ***	0.69 0.12 ***	0.89 0.14 ***	1.56 0.25 ***	0.51 0.19 **	0.43 0.44
Dist.tree												-0.57 0.21 **
Tree.H				-0.72 0.31 *				-0.18 0.09 *	-0.32 0.09 ***		-0.15 0.07 *	
Traffic					-2.79 1.81		-0.79 0.09 ***	-0.60 0.10 ***	-0.53 0.12 ***	-0.89 0.25 ***	-0.65 0.08 ***	
DPT												-1.12 0.52 *
FE												0.18 0.80
NV												-2.03 1.18 .
PT												0.27 0.72
SPT												0.16 0.55
Julian Day	-0.74 0.53	0.40 0.59	-0.49 0.19 **	0.46 0.58	1.00 0.48 *	0.45 0.30	0.01 0.10	-0.20 0.10 .	-0.05 0.12	0.45 0.21 *	-0.09 0.08	0.16 0.25
Julian Day²	-0.10 0.74	1.00 0.58 .	-0.37 0.17 *	-0.49 0.45	0.10 0.38	0.16 0.24	-0.07 0.09	0.11 0.10	-0.08 0.11	-0.23 0.18	-0.02 0.07	0.10 0.19

p < 0.1 = .
p < 0.05 = *
p < 0.01 = **
p < 0.001 = ***

Table 5: Summarised statistical results of the binomial distributed generalised linear mixed effect models (GLMM) for the flight path orientation for each species. β̂ = estimate. SE = standard error. p = significance of p value. Species names are given with the three first letters of the species and genera. Dist.tree = distance to tree foliage. LRE: long-range echolocators. MRE: mid-range echolocators. SRE: short-range echolocators.

Variable	Eptser (β̂ SE p)	Hypsav	Minsch	Myobly/myo	Nyclei	Pipkuh/nat	Pippip	Pippyg	Plesp	SRE	MRE	LRE
N observations	97	1956	1309	196	657	10787	10440	2834	81	625	27236	625
Intercept	2.32 0.48 ***	1.00 0.38 **	1.08 0.21 ***	1.53 0.33 ***	1.19 0.34 ***	1.42 0.11 ***	1.38 0.12 ***	1.56 0.17 ***	1.41 0.65 *	1.77 0.29 ***	1.51 0.12 ***	1.55 0.30 ***
Traffic						0.24 0.10 *						
Julian Day	0.68 0.47	0.65 0.77	0.22 0.14	0.06 0.18	-0.01 0.26	0.02 0.10	-0.17 0.09 .	0.22 0.12 .	0.31 0.32	0.36 0.20 .	0.06 0.07	0.20 0.23
Julian Day²	-0.66 0.51	1.38 1.09	0.02 0.13	-0.28 0.20	0.02 0.22	-0.12 0.09	-0.05 0.09	-0.06 0.12	-0.27 0.35	-0.35 0.20 .	-0.14 0.07 *	-0.18 0.21

P < 0.1 = .
P < 0.05 = *
P < 0.01 = **
P < 0.001 = ***

Discussion

This study aimed at disentangling the different mechanisms that influence collision risk: (1) density of

individuals recorded from the road edge, (2) position in the zone at risk (low flight height over the

road), (3) vehicle avoidance, and (4) risk duration (approximated by flight path orientation).

Disentangling those mechanisms was possible thanks to three-dimensional acoustic flight path

tracking (AFPT) of bat behaviour at roads at a large scale (French Mediterranean region) on a large

amount of data (122,294 bat passes). Moreover, contrary to the classic method of collecting bat

carcasses, the results of AFPT are not biased by predation or observer efficiency, and AFPT may be

applied to study roads after as well as before they are in service, if necessary.

Local landscape type

The effect of local landscape type on density was heterogeneous according to species. For example,

perpendicular tree rows led to a higher density of *P. pygmaeus* compared to simple parallel tree rows,

and forest edges led to a higher density of *H. savii* compared to simple parallel tree rows. It is rather

unlikely that the structural composition (density and orientation of linear vegetation) is the only

explanation for these differences, because other confounding effects may very well increase bat

density also, such as the different tree essences that were often associated with a type of landscape.

Interestingly, landscape type did not produce similar effects on bat density and on bat flight behaviour.

Indeed, in the case of *P. pipistrellus* for instance, bat density was the highest in perpendicular tree

rows formed by small streams. However, local landscapes eliciting the highest proportion of flights at

collision risk for this species were forests. This type of landscape was in fact a very high factor of

presence at risk for most species and guilds. Vegetation density may play an important role here.

Indeed, parallel tree rows consisted in individual planted trees with gaps of 10 to 20 m while forests

had dense shrub layers most of the time. Tree rows probably allow bats to benefit from the edge effect

(i.e. easy access to flying insects abundant in or near tree foliage) (Brigham et al., 1997; Verboom and

Spoelstra, 1999) without the need of flying directly above the road, contrarily to forest landscapes with

hard edges, which act as conduits (Kalcounis-Rueppell et al., 2013). In addition, for the LRE guild,

results show that vehicle avoidance was higher at simple parallel tree rows compared to double

parallel tree rows. It should be noted that we recorded a particularly high social activity of *N. leisleri*

in most double parallel tree rows of hollow plane trees that provide roost for this species. If bats are

less aware of the danger of vehicles when engaging in social activity, or if social activity is a proxy for

the presence of maternity roosts with naïve juveniles, it could explain the fact that bats were less

cautious in avoiding vehicles in double parallel tree rows.

Contrary to our expectations, landscape types were not selected to explain the orientation of flight

trajectories. Our results show that even in the presence of a perpendicular tree row or in the absence of

trees, bats fly most of the time parallel to the road axis. This supports the idea, not often enough

emphasised, that bats may use roads as corridors, because road verges may offer foraging

opportunities by attracting more insects than adjacent habitats (Medinas et al., 2019; Villemey et al.,

2018), and because of the verge effect when trees are present (Brigham et al., 1997; Kalcounis-

Rueppell et al., 2013; Verboom and Spoelstra, 1999).

Tree height and distance to tree foliage

As said in material and methods, distance to tree foliage was correlated to tree height in our study. In

all species except *Plecotus sp.*, density was negatively affected by an increasing distance to tree

foliage, when selected. Conversely, taller trees led to a higher density of *M. myotis/blythii* and of the

LRE guild. This effect of distance to trees was shown in several studies out of road context (Heim et

al., 2015; Kelm et al., 2014; Verboom and Spoelstra, 1999), and once at hedgerows crossing roads

(Abbott, 2012), but never at road study sites with different landscape structures and for several

distinctly identified species. In our study, the proportion of flights in the zone at collision risk was

rarely influenced by tree height or distance to trees; nonetheless, increasing distance to trees was

associated with higher proportions of flights at risk for *E. serotinus*. In addition, in several species (*M.*

daubentonii, P. pipistrellus and *P. pygmaeus*), taller trees (that generally had they foliage over the

road) led to less vehicle avoidance.

Traffic

Traffic volume did not affect bat density or position in the zone at risk for most species, contrary to

our expectations. Nonetheless, *P. kuhlii/nathusii* avoided more the zone at collision risk when traffic

increased, probably because they recognise the danger triggered by vehicles. Moreover, *P. pipistrellus*

flew more often parallel and over the external sides of the road with increasing traffic. These results

show that bats spatially avoid the vicinity of vehicles, completing the observations of Zurcher et al.

(2010). The latter did not distinguish between species, but found that 60% of approaching individuals

reversed their course in the presence of a vehicle. In addition, it was shown that flying insect

abundance decreases with increasing traffic (Martin et al., 2018). Reduced foraging opportunities

might discourage bats from foraging over the road, concomitantly with a spatial avoidance of vehicles.

Our results trivially showed that the higher the traffic volume, the lower the temporal vehicle

avoidance, probably because bats have no choice but crossing the road closely in time to vehicle

passes when traffic is high. In addition, our study showed that SRE are less likely to fly in the zone at

collision risk when a vehicle is present compared to MRE. It could be that because SRE fly lower and

are more at risk when crossing, they are more reluctant to approach vehicles, but roads could also be

foraging grounds for MRE and not for SRE, and MRE might take more risks when foraging.

Therefore, even if SRE are known to fly lower than MRE and thus at heights more similar to those of

vehicles (Berthinussen and Altringham, 2012; Roemer et al., 2019), their vehicle avoidance behaviour

should partially mitigate their susceptibility to collisions. This result emphasises the importance of

accounting for the different aspects of species behaviour when evaluating their susceptibility to

collisions (Chamberlain et al., 2006).

Time of the year

Our results show typical activity patterns throughout the year with peak density in summer or autumn,

but it is the first time to our knowledge that it is demonstrated that flight proportion in the zone at risk

at roads increases in autumn (for most species and all guilds). An increased flight proportion in the

zone at risk in autumn could partly be attributed to the naïve behaviour of juveniles, not aware of the

danger of road vehicles. Juveniles are indeed more vulnerable to road collisions than adults (Fensome

and Mathews, 2016). This result could also be explained by increasing foraging opportunities if insects

are more attracted to the warm asphalt during colder times. Bats may also be less cautious in this time

of the year where they are sexually active (i.e. increases in sexual hormones may decrease awareness

of potential danger (Alcock, 2009)), and where they need to fatten before hibernation.

Species differences

Our study provides detailed information at the species level except for species with small sample sizes,

for which readers are referred to the guild level. Models for guilds also inform on the extent of

generalisation of results because variables selected at the guild level are assumed to exert a significant

influence on several species.

Forests clearly stood out as a landscape type with a higher proportion of trajectories in the zone at risk

for MRE, *P. pipistrellus* and *P. kuhlii/nathusii*. For *H. savii* and *E. serotinus*, double parallel tree rows

elicited the smallest proportion of trajectories in the zone at risk. In most species, locations without

trees generated a relatively low proportion of trajectories in the zone at risk.

MRE seem to be generally more at risk of collision with vehicles than other species because they

showed a lower proportion of vehicle avoidance than SRE and because they had a higher proportion of

trajectories in the zone at collision risk than SRE or LRE. *Rhinolophus* species are assumed to be very

susceptible to road collisions because they fly very close to ground level (Fensome and Mathews,

2016; Jones and Rayner, 1989; Roemer et al., 2017). However *Rhinolophus* species, because of their

very high sonar frequencies (Kingston et al., 2000), are very difficult to detect and to record, and this

is why we could not study their flight behaviour with our method. AFPT with only two microphones

would allow a study of *Rhinolophus* collision risks, although with simpler metrics (Claireau et al.,

2018). One could speculate on *Rhinolophus* collision risks on roads from the SRE guild. However,

since these bats have very particular foraging strategies (Denzinger et al., 2018), they could

demonstrate quite different responses from other SRE.

Consistency (or discrepancy) between bat collision risk and measures of activity

All quantitative models succeeding the density model were interpreted as conditional probabilities that

an individual is at collision risk, and their predicted probabilities were multiplied to obtain the overall

bat collision risk if a variable was selected in several of them. The product of all quantitative models

showed that *H. savii* was more at collision risk at forests and forests edges (and to a lesser extend at

roads without trees), while *P. pipistrellus* was more at collision risk at perpendicular tree rows. These

products match the patterns of bat density in function of landscape type. The product of quantitative

models also showed that the yearly patterns of collision risks matched the ones of bat density.

Collision risks are more numerous in summer or autumn according to species, and explain the

mortality patterns found in Fensome and Mathews (2016).

However, while increasing traffic density was associated with a decrease in SRE density, it was

associated with an increase in the overall collision risk (the product of quantitative models). This

demonstrates that the measure of the number of bat passes can be a good proxy of bat collision risks in

certain contexts, but that it is necessary to also measure bat behaviour to assess collision risks with

certainty in all contexts.

Recommendations for road siting and management

Our first group of recommendations applies to habitat selection during road planning to avoid

situations with enhanced collision risks. In a trivial way, 'quality habitats' – depending on species

ecology – should generally be avoided to ensure that roads will avoid habitats with high bat density.

However, bat activity is highly dependent on distance to roost and may be under- or overrepresented at

certain habitats according to the distance to roosts. Therefore, measuring species activity at different

seasons on site will always provide more insights on the potential risks. Moreover, since it is assumed

that species do not have comparable susceptibilities to road collisions (Fensome and Mathews, 2016a),

possessing information on species presence and density is highly relevant. The present study also

allows us to emit recommendations for road siting based on the behavioural reactions to landscape

features that we measured. Forested areas should be avoided because they elicit high proportions of

flights at risk. Areas without trees should be prioritised because they almost always led to very low

activity levels and low proportions of flights at risk. However, to explain position at risk, landscape

types were only selected in models for species belonging to MRE and LRE and we cannot conclude on

their effect on SRE.

Our second recommendation applies to the management of roadside vegetation during construction

work and during the operational phase, to reduce collision risks. A gap of five meters between the road

edge and tree foliage significantly decreased the activity levels of most species. Our appreciation of

study sites suggests that this effect could be due to higher primary productivity when vegetation is

higher and closer to the road. If less primary biomass is available to insects, foraging opportunities for

bats decrease, and so does their density (Threlfall et al., 2012). It is however controversial to

recommend cutting trees at road sides, because this decision will engender habitat loss in numerous

taxa, especially in large-scale impacted areas such as linear transport infrastructures. Opening habitat

at road edges also creates suitable foraging grounds for birds of prey for instance (Morelli et al., 2014),

and will increase their collision probability. It is possible to make these open verges less attractive by

converting them to gravel surface (Kociolek et al., 2015), but this will eliminate plant habitats.

Another possibility is to only cut a certain number of trees and clear shrub layers periodically (a

frequent practice in French Mediterranean forests to prevent fires) to reduce primary production and to

allow bats to navigate between trees rather than above the asphalt. The local management will thus

depend on the biodiversity stakes of the area. In areas of high stakes, reducing vehicle speed limit

could be an efficient solution, but this was not tested on bats to our knowledge.

Limits of the study and perspectives

Our recommendations can only apply to landscapes and bat communities similar to the ones that we

sampled. Therefore, complementary studies should be conducted in other biogeographical areas (e.g.

Continental or Atlantic areas) to make sure that bats react consistently to the same road landscape

features. However, we expect this endeavour to be quite difficult because of the local particularities in

landscape management. Since we expect bats to be more active at prolific foraging grounds, it would

be interesting to see if the measure of primary production - for example using satellite imaging - can

be a more universal descriptor of bat activity than the description of the local landscape.

Several questions remain unanswered, such as the role of tree essence, topography at a medium scale

(i.e. slope of the terrain), and topography at a small scale (i.e. road embankments) in bat collision risks

at roads. The nearby presence of a bat roost is also expected to be an important factor of collisions. At

last, it was reported that juveniles and males are more prone to road collisions (Fensome and Mathews,

2016). It would be interesting to study the behaviour of bats of different age and sex to explain this

finding.

623   Acknowledgments

624   This study was a collaboration between Biotope and the Muséum national d'Histoire naturelle (Paris,

625   France) in the form of a PhD thesis funded by Biotope and the Association Nationale de la Recherche

626   et de la Technologie. We thank Fiona Mathews for her comments, which helped to improve the

627   manuscript. We would like to thank Bruno Sanchez, Dominique Guicheteau (RNN de la Plaine des

628   Maures), Ugo Schumpp and Julien Penvern for their precious help during field work.

630   Conflict of interest

631   Biotope is an environmental consultancy involved in road impact assessment studies. Two of the

632   authors, Charlotte Roemer and Thierry Disca, were employees at Biotope during the time of the study.

633   Authors thus declare a conflict of interest. However, authors take complete responsibility for the

634   integrity of the data and the accuracy of their analysis.

636   References

Abbott, I.M., 2012. Assessment of the effectiveness of mitigation measures employed on Irish national road schemes for the conservation of bats (PhD Thesis). Ph. D. thesis, University College Cork, Ireland.

Abbott, I.M., Butler, F., Harrison, S., 2012. When flyways meet highways – The relative permeability of different motorway crossing sites to functionally diverse bat species. Landsc. Urban Plan. 106, 293–302. https://doi.org/10.1016/j.landurbplan.2012.03.015

Alcock, J., 2009. Animal behavior, ninth edition. ed. Sinauer Associates, Sunderland, MA, USA.

Aldridge, H.D.J.N., Rautenbach, I.L., 1987. Morphology, Echolocation and Resource Partitioning in Insectivorous Bats. J. Anim. Ecol. 56, 763. https://doi.org/10.2307/4947

Arnold, T.W., 2010. Uninformative Parameters and Model Selection Using Akaike's Information Criterion. J. Wildl. Manag. 74, 1175–1178. https://doi.org/10.2193/2009-367

Arthur, L., Lemaire, M., 2015. Les chauves-souris de France, Belgique, Luxembourg et Suisse. Biotope Editions, Mèze; Paris.

Barataud, M., 2015. Acoustic ecology of European bats: species identification, study of their habitats and foraging behaviour. Biotope éditions.

Bas, Y., Bas, D., Julien, J.-F., 2017. Tadarida: A Toolbox for Animal Detection on Acoustic Recordings. J. Open Res. Softw. 5. https://doi.org/10.5334/jors.154

Bennett, V.J., Zurcher, A.A., 2013. When corridors collide: Road-related disturbance in commuting bats. J. Wildl. Manag. 77, 93–101. https://doi.org/10.1002/jwmg.467

Berthinussen, A., Altringham, J., 2012. Do bat gantries and underpasses help bats cross roads safely? PloS One 7, e38775.

Brigham, R.M., Grindal, S.D., Firman, M.C., Morissette, J.L., 1997. The influence of structural clutter on activity patterns of insectivorous bats. Can. J. Zool. 75, 131–136.

Brooks, M.E., Kristensen, K., van Benthem, K.J., Magnusson, A., Berg, C.W., Nielsen, A., Skaug, H.J., Maechler, M., Bolker, B.M., 2017. Modeling Zero-Inflated Count Data With glmmTMB. https://doi.org/10.1101/132753

Chamberlain, D.E., Rehfisch, M.R., Fox, A.D., Desholm, M., Anthony, S.J., 2006. The effect of avoidance rates on bird mortality predictions made by wind turbine collision risk models. Ibis 148, 198–202.

Claireau, F., Bas, Y., Puechmaille, S.J., Julien, J.-F., Allegrini, B., Kerbiriou, C., 2018. Bat overpasses: An insufficient solution to restore habitat connectivity across roads. J. Appl. Ecol. 56, 573–584. https://doi.org/10.1111/1365-2664.13288

Clevenger, A.P., Chruszcz, B., Gunson, K.E., 2003. Spatial patterns and factors influencing small vertebrate fauna road-kill aggregations. Biol. Conserv. 109, 15–26.

Council Directive 92/43/EEC on the conservation of natural habitats and of wild fauna and flora, 1992. , 206.

D'Amico, M., Román, J., de los Reyes, L., Revilla, E., 2015. Vertebrate road-kill patterns in Mediterranean habitats: Who, when and where. Biol. Conserv. 191, 234–242. https://doi.org/10.1016/j.biocon.2015.06.010

Denzinger, A., Tschapka, M., Schnitzler, H.-U., 2018. The role of echolocation strategies for niche differentiation in bats. Can. J. Zool. 96, 171–181. https://doi.org/10.1139/cjz-2017-0161

Dietz, C., Nill, D., Helversen, O. von, 2009. Bats of Britain, Europe and Northwest Africa. A & C Black.

Fensome, A.G., Mathews, F., 2016. Roads and bats: a meta-analysis and review of the evidence on vehicle collisions and barrier effects. Mammal Rev. 46, 311–323. https://doi.org/10.1111/mam.12072

Frey-Ehrenbold, A., Bontadina, F., Arlettaz, R., Obrist, M.K., 2013. Landscape connectivity,
habitat structure and activity of bat guilds in farmland-dominated matrices. J. Appl.
Ecol. 50, 252–261. https://doi.org/10.1111/1365-2664.12034

Froidevaux, J.S.P., Louboutin, B., Jones, G., 2017. Does organic farming enhance
biodiversity in Mediterranean vineyards? A case study with bats and arachnids. Agric.
Ecosyst. Environ. 249, 112–122. https://doi.org/10.1016/j.agee.2017.08.012

Gaisler, J., Řehák, Z., Bartonička, T., 2009. Bat casualties by road traffic (Brno-Vienna). Acta
Theriol. (Warsz.) 54, 147–155.

Grilo, C., de Resende Cardoso, T., Solar, R., Bager, A., 2016. Do the size and shape of spatial
units jeopardize the road mortality-risk factors estimates? Nat. Conserv. 14, 8–13.
https://doi.org/10.1016/j.ncon.2016.01.001

Gunson, K.E., Mountrakis, G., Quackenbush, L.J., 2011. Spatial wildlife-vehicle collision
models: A review of current work and its application to transportation mitigation
projects. J. Environ. Manage. 92, 1074–1082.
https://doi.org/10.1016/j.jenvman.2010.11.027

Harrell, F.E., 2018. Hmisc: Harrell Miscellaneous.

Hartig, F., 2018. DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed)
Regression Models. Theoretical Ecology, University of Regensburg, Regensburg,
Germany.

Heiberger, R.M., Holland, B., 2004. Multiple Comparisons, in: Statistical Analysis and Data
Display, Springer Texts in Statistics. Springer, New York, NY, pp. 155–185.
https://doi.org/10.1007/978-1-4757-4284-8_7

Heim, O., Treitler, J.T., Tschapka, M., Knörnschild, M., Jung, K., 2015. The Importance of
Landscape Elements for Bat Activity and Species Richness in Agricultural Areas.
PLOS ONE 10, e0134443. https://doi.org/10.1371/journal.pone.0134443

Holderied, M.W., Jones, G., 2009. Flight Dynamics. Ecol. Behav. Methods Study Bats 459–
475.

Holderied, M.W., von Helversen, O., 2003. Echolocation range and wingbeat period match in
aerial-hawking bats. Proc. R. Soc. B Biol. Sci. 270, 2293–2299.
https://doi.org/10.1098/rspb.2003.2487

Ibisch, P.L., Hoffmann, M.T., Kreft, S., Pe'er, G., Kati, V., Biber-Freudenberger, L.,
DellaSala, D.A., Vale, M.M., Hobson, P.R., Selva, N., 2016. A global map of roadless
areas and their conservation status. Science 354, 1423–1427.
https://doi.org/10.1126/science.aaf7166

Institut National de l'Information Géographique et Forestière, 2017. BD TOPO version 2.2.

Jaeger, J.A., Bowman, J., Brennan, J., Fahrig, L., Bert, D., Bouchard, J., Charbonneau, N.,
Frank, K., Gruber, B., von Toschanowitz, K.T., 2005. Predicting when animal
populations are at risk from roads: an interactive model of road avoidance behavior.
Ecol. Model. 185, 329–348.

Jones, G., Rayner, J.M., 1989. Foraging behavior and echolocation of wild horseshoe bats
Rhinolophus ferrumequinum and R. hipposideros (Chiroptera, Rhinolophidae). Behav.
Ecol. Sociobiol. 25, 183–191.

Kalcounis-Rueppell, M.C., Briones, K.M., Homyack, J.A., Petric, R., Marshall, M.M., Miller,
D.A., 2013. Hard forest edges act as conduits, not filters, for bats. Wildl. Soc. Bull. 37,
571–576.

Kelm, D.H., Lenski, J., Kelm, V., Toelch, U., Dziock, F., 2014. Seasonal Bat Activity in
Relation to Distance to Hedgerows in an Agricultural Landscape in Central Europe
and Implications for Wind Energy Development. Acta Chiropterologica 16, 65–73.
https://doi.org/10.3161/150811014X683273

Kingston, T., Jones, G., Zubaid, A., Kunz, T.H., 2000. Resource partitioning in rhinolophoid bats revisited. Oecologia 124, 332–342. https://doi.org/10.1007/PL00008866

Koblitz, J.C., 2018. Arrayvolution-Using microphone arrays to study bats in the field. Can. J. Zool.

Kociolek, A., Grilo, C., Jacobson, S., 2015. Flight doesn't solve everything: mitigation of road impacts on birds. Handb. Road Ecol. 1.

Langwig, K.E., Hoyt, J.R., Parise, K.L., Kath, J., Kirk, D., Frick, W.F., Foster, J.T., Kilpatrick, A.M., 2015. Invasion Dynamics of White-Nose Syndrome Fungus, Midwestern United States, 2012–2014. Emerg. Infect. Dis. 21, 1023–1026. https://doi.org/10.3201/eid2106.150123

Lesiński, G., 2007. Bat road casualties and factors determining their number. mammalia 71. https://doi.org/10.1515/MAMM.2007.020

Lin, S.-C., 2016. Landscape and traffic factors affecting animal road mortality. J. Environ. Eng. Landsc. Manag. 24, 10–20. https://doi.org/10.3846/16486897.2015.1098652

Malo, J.E., Suárez, F., Díez, A., 2004. Can we mitigate animal–vehicle accidents using predictive models? J. Appl. Ecol. 41, 701–710. https://doi.org/10.1111/j.0021-8901.2004.00929.x

Martin, A.E., Graham, S.L., Henry, M., Pervin, E., Fahrig, L., 2018. Flying insect abundance declines with increasing road traffic. Insect Conserv. Divers. https://doi.org/10.1111/icad.12300

Mazerolle, M.J., 2004. AMPHIBIAN ROAD MORTALITY IN RESPONSE TO NIGHTLY VARIATIONS IN TRAFFIC INTENSITY. Herpetologica 60, 45–53. https://doi.org/10.1655/02-109

Medinas, D., Marques, J.T., Mira, A., 2013. Assessing road effects on bats: the role of landscape, road features, and bat activity on road-kills. Ecol. Res. 28, 227–237. https://doi.org/10.1007/s11284-012-1009-6

Medinas, D., Ribeiro, V., Marques, J.T., Silva, B., Barbosa, A.M., Rebelo, H., Mira, A., 2019. Road effects on bat activity depend on surrounding habitat type. Sci. Total Environ. 660, 340–347.

Meisingset, E.L., Loe, L.E., Brekkum, Ø., Mysterud, A., 2014. Targeting mitigation efforts: The role of speed limit and road edge clearance for deer–vehicle collisions. J. Wildl. Manag. 78, 679–688. https://doi.org/10.1002/jwmg.712

Morelli, F., Beim, M., Jerzak, L., Jones, D., Tryjanowski, P., 2014. Can roads, railways and related structures have positive effects on birds?–A review. Transp. Res. Part Transp. Environ. 30, 21–31.

Nelli, L., Langbein, J., Watson, P., Putman, R., 2018. Mapping risk: Quantifying and predicting the risk of deer-vehicle collisions on major roads in England. Mamm. Biol. 91, 71–78. https://doi.org/10.1016/j.mambio.2018.03.013

Neumann, W., Ericsson, G., Dettki, H., Bunnefeld, N., Keuler, N.S., Helmers, D.P., Radeloff, V.C., 2012. Difference in spatiotemporal patterns of wildlife road-crossings and wildlife-vehicle collisions. Biol. Conserv. 145, 70–78. https://doi.org/10.1016/j.biocon.2011.10.011

Norberg, U.M., 1994. Wing design, flight performance, and habitat use in bats. Ecol. Morphol. Integr. Org. Biol. 205–239.

Norberg, U.M., 1986. Evolutionary convergence in foraging niche and flight morphology in insectivorous aerial-hawking birds and bats. Ornis Scand. 253–260.

Orłowski, G., 2008. Roadside hedgerows and trees as factors increasing road mortality of birds: Implications for management of roadside vegetation in rural landscapes. Landsc. Urban Plan. 86, 153–161. https://doi.org/10.1016/j.landurbplan.2008.02.003

Popa-Lisseanu, A.G., 2007. Roosting Behaviour, Foraging Ecology & the Enigmatic Dietary Habits of the Aerial-Hawking Bat Nyctalus lasiopterus. Universidad de Sevilla, Sevilla, Spain.

Pourshoushtari, R.D., Pauli, B.P., Zollner, P.A., Haulton, G.S., 2018. Road and Habitat Interact to Influence Selection and Avoidance Behavior of Bats in Indiana. Northeast. Nat. 25, 236–247. https://doi.org/10.1656/045.025.0206

R Core Team, 2014. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.

Roemer, C., Coulon, A., Disca, T., Bas, Y., 2019. Bat sonar and wing morphology predict species vertical niche. J. Acoust. Soc. Am. 145, 3242–3251.

Roemer, C., Disca, T., Coulon, A., Bas, Y., 2017. Bat flight height monitored from wind masts predicts mortality risk at wind farms. Biol. Conserv. 215, 116–122. https://doi.org/10.1016/j.biocon.2017.09.002

Russell, A., Butchkoski, C., Saidak, L., McCracken, G., 2009. Road-killed bats, highway design, and the commuting ecology of bats. Endanger. Species Res. 8, 49–60. https://doi.org/10.3354/esr00121

Rytwinski, T., Fahrig, L., 2015. The Impacts of Roads and Traffic on Terrestrial Animal Populations. https://doi.org/10.1002/9781118568170.ch28

Seiler, A., 2005. Predicting locations of moose–vehicle collisions in Sweden. J. Appl. Ecol. 42, 371–382. https://doi.org/10.1111/j.1365-2664.2005.01013.x

Threlfall, C.G., Law, B., Banks, P.B., 2012. Influence of landscape structure and human modifications on insect biomass and bat foraging activity in an urban landscape. PLoS One 7, e38800.

Toffoli, R., 2016. The Importance of Linear Landscape Elements for Bats in a Farmland Area: The Influence of Height on Activity. J. Landsc. Ecol. 9. https://doi.org/10.1515/jlecol-2016-0004

Valero, E., Picos, J., Álvarez, X., 2015. Road and traffic factors correlated to wildlife–vehicle collisions in Galicia (Spain). Wildl. Res. 42, 25. https://doi.org/10.1071/WR14060

Van der Meij, T., Van Strien, A.J., Haysom, K.A., Dekker, J., Russ, J., Biala, K., Bihari, Z., Jansen, E., Langton, S., Kurali, A., Limpens, H., Meschede, A., Petersons, G., Presetnik, P., Prüger, J., Reiter, G., Rodrigues, L., Schorcht, W., Uhrin, M., Vintulis, V., 2015. Return of the bats? A prototype indicator of trends in European bat populations in underground hibernacula. Mamm. Biol. - Z. Für Säugetierkd. 80, 170–177. https://doi.org/10.1016/j.mambio.2014.09.004

van der Ree, R., Smith, D.J., Grilo, C., 2015a. The ecological effects of linear infrastructure and traffic. Handb. Road Ecol. 1–9.

van der Ree, R., Smith, D.J., Grilo, C., 2015b. Handbook of Road Ecology. John Wiley & Sons.

Verboom, B., Spoelstra, K., 1999. Effects of food abundance and wind on the use of tree lines by an insectivorous bat, Pipistrellus pipistrellus. https://doi.org/10.1139/cjz-77-9-1393

Villemey, A., Jeusset, A., Vargac, M., Bertheau, Y., Coulon, A., Touroult, J., Vanpeene, S., Castagneyrol, B., Jactel, H., Witte, I., Deniaud, N., Flamerie De Lachapelle, F., Jaslier, E., Roy, V., Guinard, E., Le Mitouard, E., Rauel, V., Sordello, R., 2018. Can linear transportation infrastructure verges constitute a habitat and/or a corridor for insects in temperate landscapes? A systematic review. Environ. Evid. 7. https://doi.org/10.1186/s13750-018-0117-3

Zimmermann Teixeira, F., Kindel, A., Hartz, S.M., Mitchell, S., Fahrig, L., 2017. When road-kill hotspots do not indicate the best sites for road-kill mitigation. J. Appl. Ecol. https://doi.org/10.1111/1365-2664.12870

834 Zurcher, A.A., Sparks, D.W., Bennett, V.J., 2010. Why the Bat Did Not Cross the Road? Acta
835   Chiropterologica 12, 337–340. https://doi.org/10.3161/150811010X537918
836 Zuur, A., Ieno, E., Elphick, C., 2010. A protocol for data exploration to avoid common
837   statistical problems. Methods Ecol. Evol. 1, 3–14.

841 **Supplementary Material**

843
844 Figure A 1: Landscape types. Microphones are shown on poles but when possible, they were
845 attached to vegetation instead. The box between two microphones is the recorder.

846
847
848
849
Table A 1: Summary table for the total amount of data in each category and for each species. SRE:
Short-range echolocators. MRE: Mid-range echolocators. LRE: Long-rang echolocators. N bat passes:
rounded median number of bat passes among all four microphones. N flight trajectories: acoustic
recordings with more than 3 consecutive calls recorded by all 4 microphones. N successful flight
trajectories: flight trajectories with at least one call in the precision zone (<10 m from microphones
centroid).

Species	Guild	N bat passes	N flight trajectories	N successful flight trajectories
B. barbastellus	SRE	1157	311	97

Species	Category	Col1	Col2	Col3
M. alcathoe	SRE	2	1	1
M. bechsteinii	SRE	32	11	2
M. capaccinii	SRE	98	40	24
M. daubentonii	SRE	368	287	150
M. emarginatus	SRE	287	29	17
M. blythii/myotis	SRE	1477	583	201
M. nattereri	SRE	345	75	37
Myotis sp.	SRE	405	101	40
Plecotus sp.	SRE	908	281	89
R. euryale	SRE	6	0	0
R. ferrumequinum	SRE	68	2	2
R. hipposideros	SRE	129	3	1
H. savii	MRE	4634	3912	2053
M. schreibersii	MRE	9226	2268	1417
P. kuhlii/nathusii	MRE	45152	28601	11685
P. pipistrellus	MRE	32383	19830	11261
P. pygmaeus	MRE	14410	6136	3033
Eptesicus/ Nyctalus/ Vespertilio	LRE	503	141	31
E. serotinus	LRE	1161	475	107
N. lasiopterus	LRE	13	2	0
N. leisleri	LRE	9306	3506	698
N. noctula	LRE	32	16	5
T. teniotis	LRE	192	53	3
TOTAL		122294	66664	30954

Myotis emarginatus

Myotis myotis/blythii

Nyctalus leisleri

Pipistrellus kuhlii/nathusii

Pipistrellus pipistrellus

Pipistrellus pygmaeus

Plecotus sp.

860
861 Figure A 2: Predicted values for each model and each species or guild in function of time of the year (in month). 95% confidence intervals are shown. The last
862 column displays the product of the first three models (density x in zone at risk x vehicles not avoided) to obtain the number of bat passes at risk per night
863 (logarithmic scale).
864
865

889
890 Figure A 3: Product of the first three models (density x in zone at risk x vehicles not avoided) to obtain
891 the number of bat passes at risk per night (logarithmic scale) for variables selected in at least two
892 models for the same species. 95% confidence intervals are shown. Figures represent landscape type
893 viewed from the top (road in light grey and trees in black). SRE: short-range echolocators. LRE: long-
894 range echolocators. MRE: mid-range echolocators.
895

899