Systematic Engineering of Artificial Metalloenzymes for New-to-Nature Reactions

Tobias Vornholt,[1,3] Fadri Christoffel,[2,3,§] Michela M. Pellizzoni,[2,3,§] Sven Panke,[1,3] Thomas R. Ward[2,3] and Markus Jeschek[1,3,*]

[1]Department of Biosystems Science and Engineering, ETH Zurich, Basel CH-4058, Switzerland

[2]Department of Chemistry, University of Basel, Mattenstrasse 24a, BPR 1096, CH-4002 Basel Switzerland

[3]National Centre of Competence in Research (NCCR) Molecular Systems Engineering, Switzerland; Web: www.nccr-mse.ch

[§]Both authors contributed equally

[*]Correspondence to: markus.jeschek@bsse.ethz.ch

Abstract

Artificial metalloenzymes (ArMs) catalyze new-to-nature reactions under mild conditions and could therefore play an important role in the transition to a sustainable, circular economy. While ArMs have been created for a variety of reactions, their activity for most biorthogonal transformations has remained modest and attempts at optimizing them by means of enzyme engineering have been case-specific and unsystematic. To realize the great potential of ArMs for biocatalysis and synthetic biology, there is a need for methods that enable the rapid discovery of highly active ArM variants for any reaction of interest. Here, we present a broadly applicable, automation-compatible ArM engineering platform. It relies on periplasmic compartmentalization of the ArM in *Escherichia coli* to rapidly and reliably identify improved variants based on the biotin-streptavidin technology. We assess 400 sequence-verified ArM mutants for five bio-orthogonal transformations involving different metal cofactors, reaction mechanisms and substrate-product pairs, including novel ArMs for gold-catalyzed hydroamination and hydroarylation. The achieved activity enhancements of six- to fifteen-fold highlight the potential of the proposed systematic approach to ArM engineering. We further capitalize on the sequence-activity data to suggest and validate smart strategies for future screening campaigns. This systematic, multi-reaction study has important implications for the development of highly active ArMs for novel applications in biocatalysis and synthetic biology.

Introduction

Artificial metalloenzymes (ArMs) combine the broad reaction scope of organometallic catalysis with the selectivity and mild reaction conditions of enzymes[1,2]. Therefore, they have a great potential to enable more sustainable synthetic routes to various compounds of interest[3,4] and, if functional in a cellular environment, open up new possibilities for metabolic engineering and synthetic biology[5]. ArMs have been generated by repurposing natural metalloenzymes[6,7], designing binding sites for metal ions[8–11] and incorporating organometallic cofactors into protein scaffolds[12–16]. Among these, one of the most versatile approaches relies on the biotin-streptavidin technology. This strategy employs the high affinity of the homotetrameric protein streptavidin (Sav) for the vitamin D-biotin to non-covalently anchor biotinylated metal complexes (referred to as cofactor hereafter) within the Sav protein. Using this approach (and modifications thereof), artificial enzymes have been created for multiple reactions, including hydrogenation, sulfoxidation, C−H activation and olefin metathesis[17–19]. While wild-type Sav may impart some selectivity or rate acceleration on the reaction, protein engineering is usually required in order to obtain proficient biocatalysts[20–23]. Initial studies in this direction relied on screening (semi-)purified Sav mutants[24,25], but more recently a trend towards whole-cell screening with *Escherichia coli* has emerged for Sav-based ArMs[26,27] as well as other artificial enzymes[8,28,29]. This allows significantly increased throughput and is the method of choice if variants for *in vivo* synthetic biology applications are desired.

However, a number of challenges arise for cell-based ArM assays, most notably insufficient cofactor uptake into the cell as well as cofactor poisoning by cellular components such as reduced glutathione[30]. In order to circumvent these challenges, Sav has been exported to the periplasmic space[26] or the cell surface[27]. While these studies have established the possibility of engineering ArMs using whole-cell screenings, they did so using case-specific engineering strategies and with a focus on reactions affording fluorescent products. The general applicability of these methods across various reactions remains to be demonstrated. To generalize the development of ArMs, however, broadly applicable strategies for ArM engineering are required, which enable rapid identification of active variants for any reaction of interest. This requires a robust screening protocol that is compatible with various reaction conditions and read-outs. Moreover, it imposes a demand for Sav mutant libraries that embody a high potential to contain highly active variants for different reactions while maintaining a comparably small (and thus screenable) library size. Such a combination of a broadly applicable screening method with a "concise" library could serve as a universal starting point for various ArM engineering campaigns and would render tedious case-by-case method and library development obsolete. Herein, we present a screening platform that meets the aforementioned requirements as a first systematic approach to ArM engineering. We establish a well-plate based screening protocol that can be easily adapted to new reaction conditions or analytical methods and show that it is amenable to lab automation as an important prerequisite to generalizable ArM engineering approaches. In addition, we create a full-factorial, sequence-defined Sav library by simultaneously diversifying two crucial amino acid positions in close vicinity to the catalytic metal. To demonstrate the versatility of this platform, we selected

five bio-orthogonal reactions requiring different reaction conditions and analytical readouts. This platform

enabled the identification of substantially improved ArMs for all tested reactions with fast turnaround times.

Moreover, the systematic characterization of the sequence-activity landscape enabled us to identify smart

library designs to further streamline the future development of ArMs. The efforts presented herein represent

the first systematic assessment of ArMs based on the biotin-streptavidin technology and constitute an

important step towards a streamlined optimization of these novel biocatalysts.

Results

Construction and characterization of a sequence-defined Sav library

In order to establish a generalizable first engineering step for ArMs, we sought to generate a library of Sav

variants that offers a high likelihood of identifying active variants at a minimized screening effort. Based on

previous experience with Sav expression and whole-cell ArM catalysis[26,27], we selected a periplasmic

compartmentalization strategy for this library. Secretion to the periplasm and subsequent ArM assembly

represents a good trade-off between accessibility to the cofactor, expression levels and compatibility of the

reaction conditions[31]. In order to facilitate periplasmic export in *E. coli*, the signal peptide of the outer

membrane protein A (OmpA) was N-terminally fused to T7-tagged mature Sav (referred to as wild type

hereafter) as previously reported[26]. The holoenzyme (i.e. full ArM with cofactor) can then be assembled in

the periplasm by incubating cells in a buffer containing a biotinylated metal cofactor (Figure 1a). We selected

the amino acid residues 112 and 121 in Sav as randomization targets, which correspond to serine and

lysine in wild-type Sav. These residues are in close proximity to the biotinylated cofactor[24] (Figure 1b) and

have repeatedly been found to have a substantial impact on the activity of ArMs for diverse cofactors and

reactions[25–27]. More specifically, we created a combinatorial Sav 112X 121X library (X representing all 20

canonical amino acids) consisting of all 400 possible amino acid combinations for these two positions. Such

a full-factorial library makes it possible to identify improved mutants that are the result of synergistic

interactions between the two positions. We applied a three-step cloning and sequencing strategy (see

Methods and Supplementary Figs. 1-3) to produce an arrayed, sequence-verified set of all 400 possible

Sav 112X 121X mutants. This minimizes the screening effort by avoiding the requirement for oversampling.

Figure 1 | Platform for the systematic screening of artificial metalloenzymes (ArMs) in *E. coli*'s periplasm. a, Sav is secreted
to *E. coli*'s periplasm by fusion to the signal peptide of the outer membrane protein A (OmpA), where it binds to an externally-added
biotinylated cofactor (containing a catalytically active metal M and its ligands L and L') to afford an ArM. **b**, Left: Close-up view of a
bound biotinylated metathesis cofactor within the biotin-binding vestibule of homotetrameric Sav (PDB 5IRA). The symmetry-related
amino acids S112 & S'112 (blue) and K121 and K'121 (purple) in Sav are located in the immediate vicinity of the cofactor and were
mutated in a combinatorial manner (right), affording an Sav library with 400 amino acid combinations in these two positions. **c**,
Expression level of all 400 Sav mutants as determined in cell lysate using a fluorescence quenching assay[32] (Methods). The values
displayed are the mean of biological triplicates (n=3) and were normalized to the OD_{600} of the cultures. Refer to Supplementary
Figure 4 for the standard deviation between replicates. **d**, Percentage of Sav mutants with unoccupied biotin-binding sites as a
function of the concentration of periplasmic cofactor (given in terms of the volume of the cell suspension). When adding 10 µM
cofactor to the cell suspension, it can be expected that less than half of the cofactor enters the periplasm. This means that an
excess of binding sites is present for more than 90 % of Sav variants (dashed lines). **e**, Overview of the screening workflow
(Methods). Following periplasmic Sav expression in deep-well plates, cells are incubated with the respective biotinylated cofactor
and subsequently washed once prior to the addition of substrate (circular arrows represent centrifugation steps for buffer exchange).
Depending on the reaction, the product concentration is determined via absorbance or fluorescence measurements or by LC-MS.

Even single mutations can substantially alter the expression level of proteins[33], which complicates the
identification of variants with increased specific activity. For this reason, we aimed to screen at cofactor
concentrations that do not saturate the available biotin-binding sites even for low-expressing variants.

Consequently, we analyzed the expression level of all 400 Sav mutants from the S112X K121X library,

relying on a quenching assay with biotinylated fluorescent probes[32] (Figure 1c, Methods). The vast majority

of variants showed good expression levels ranging from 17 to 536 mg L^{-1}, or 1 and 33 µM biotin-binding

sites. When normalized by the density of the culture, these values ranged from 0.2 to 3.2 µM binding sites

in a culture with an optical density at 600 nm (OD$_{600}$) of one. Importantly, while the expression of variants

harboring a cysteine at position 112 or a tryptophan at either of the two positions appeared to be reduced,

87 % of Sav mutants showed an expression level greater than 1 µM binding sites per OD$_{600}$. Based on

these measurements, we determined a maximum permitted cofactor concentration (Fig. 1d). This critical

experimental parameter should on the one hand be high enough to result in well-detectable product

concentrations and on the other hand remain below the concentration of available biotin-binding sites for

the majority of the library. The latter requirement is important in order to keep the concentration of

assembled ArMs constant, irrespective of the expression level of Sav. In light of previous studies[26] and our

own observations (Supplementary Fig. 5), we conservatively assumed that less than half of externally-

added cofactor enters the periplasm. As a consequence, we set the maximum cofactor concentration at the

incubation step to 10 µM, which is expected to result in a concentration of periplasmic cofactor of less than

5 µM. This means that an excess of binding sites is maintained for more than 90 % of variants (Figure 1d),

thus largely eliminating the expression level dependence of the screening results.

Systematic screening of ArMs for bio-orthogonal reactions

In order to facilitate screening of improved ArMs for diverse reactions, we relied on a 96-well-plate based

assay (Methods)[26]. In brief, periplasmic expression cultures are spun down and cells are resuspended in

an incubation buffer containing the cofactor of interest to assemble the ArM in the periplasm. After

incubation, cells are washed once to remove unbound cofactor and resuspended in a reaction-specific

buffer containing the substrate. Following overnight incubation, the product concentration is determined

using a suitable analytical method (Figure 1e). This screening procedure is compatible with various reaction

conditions and analytical methods, and consequently not restricted to model reactions, which, for example,

produce a fluorescent product. Furthermore, it is amenable to lab automation. We thus automated the steps

required for incubation with cofactor, washing and substrate addition and applied this automated protocol

to two reactions. This substantially reduces the manual labor required for a screening, as well as the

consumption of consumables. The robotic platform can handle up to eight 96-well plates concurrently.

Accordingly, the 400-member library can be processed in one day and an entire screening can be performed

within one week.

Relying on this protocol, we sought to systematically test the full-factorial 400-mutant Sav library for various

ArM reactions of interest (Figure 2a). To this end, we selected three reactions based on previously reported

ArMs: A ring-closing metathesis (RCM, **I**) of diallyl-sulfonamide **1** to 2,5-dihydro-pyrrole **2**[26] and two

deallylation reactions (**II**, **III**) of allyl carbamate-protected substrates **3** and **5,** affording amino coumarin **4**

and indole **6**, respectively. The corresponding cofactors for these reactions are either a biotinylated second-

generation Hoveyda-Grubbs catalyst **(Biot-NHC)Ru** (**I**) or a biotinylated ruthenium cyclopentadienyl complex **(Biot-HQ)CpRu**[34,35] (**II**, **III**), respectively (Figure 2b). Further, with the aim of extending the scope of ArM-catalyzed reactions towards biocompatible nucleophilic cyclizations of alkynes, we developed two novel gold-containing ArM cofactors, (**Biot-NHC)Au1** and (**Biot-NHC')Au2** (Fig. 2b), to create ArMs for hydroamination (**IV**) and hydroarylation[36] (**V**) reactions. Importantly, none of these reactions are known to be catalyzed by natural enzymes, and therefore they have potential for applications ranging from the design of novel metabolic pathways to prodrug activation and *in vivo* labelling[5,37].

While preliminary experiments on the RCM **I** and deallylations **II** and **III** confirmed the compatibility of these reactions with the periplasmic screening platform, reactions **IV** and **V** showed only little conversion in the whole-cell assay. Based on previous experience with ArMs[26,30], we hypothesized that cellular thiols might inhibit these gold-catalyzed reactions, which we could confirm by *in vitro* experiments with purified Sav. These revealed a marked inhibitory effect of glutathione and cysteine (Supplementary Fig. 6a). Previously, we had reported that thiol inhibition can be overcome *in vitro* by adding the oxidizing agent diamide[30]. Gratifyingly, we observed that diamide also neutralizes the detrimental effect of thiols in whole-cell experiments, rendering periplasmic gold catalysis feasible (Supplementary Fig. 6b).

With a functional periplasmic screening assay for the five reactions at hand, we tested all 400 mutants for each reaction at least in biological duplicates relying on the aforementioned workflow in 96-deep well plates. This allowed us to determine a local sequence-activity landscape for each ArM (Figure 2c). The activity patterns varied substantially between reactions, indicating that the observed effects are indeed the result of specific interactions between protein, cofactor and substrate, as opposed to unspecific effects such as the expression level. In line with this, we observed no correlation between expression level and activity for any of the reactions (Supplementary Fig. 7). Hence, the concentration of periplasmic cofactor, and not the number of available biotin-binding sites is limiting, which confirms the validity of the determined maximum permitted cofactor concentration (see above). Similar activity patterns were only found between the two deallylation reactions (**II** and **III**), suggesting that in these cases either the substrates are sufficiently similar or that the observed activity is mainly the result of interactions between protein and cofactor.

For all reactions, the wild-type variant (S112 K121) had a comparably low activity that was only slightly above the background observed for cells lacking Sav. Note that in this case the residual activity results from residual cofactor that is unspecifically bound and thus removed in the washing step. Importantly, we identified significantly more active mutants for all five ArM reactions (Figure 2d). In order to validate these results, we measured the activity of the most promising variants of each ArM again in eight replicates (Fig. 2e), which confirmed the observed enhancements. The best mutants reached fold-improvements over the wild type varying between six- and fifteen-fold. Importantly, these activity improvements compare favorably to previous efforts of optimizing Sav-based ArMs[26,27]. Similarly, the "hits" appear to be more active than variants we identified previously for the same reactions. For instance, several studies had previously described allylic deallylases that uncage allylcarbamate-protected substrates. The most extensive

engineering study of these was a screening of 80 surface-displayed variants with mutations at positions
112 and 121, yielding the double mutants 112M 121A and 112Y 121S as most active variants for the
uncaging of allylcarbamate-protected amino coumarin **3** (**II**)[27]. While both of these mutants were highly
active in our periplasmic screen, several others displayed even higher activities, reaching an improvement
over the wild type of up to approximately fifteen-fold, compared to nine- and ten-fold for the previously
identified variants.

Similarly, we identifed substantially improved variants of the novel, gold-based ArMs for hydroamination
(**IV**) and hydroarylation (**V**) reactions. For the latter reaction, we observed comparably low product
concentrations for most variants, which was likely due to the poor solubility of substrate and product. As a
result of this, fluorescence measurements were close to the background of cellular autofluorescence for
many variants. Nonetheless, significantly improved variants could be identified reliably using the established
screening workflow.

Figure 2 | Systematic screening for ArMs catalyzing diverse reactions. a, ArM-catalyzed reactions: **I**, Ring-closing metathesis (RCM) with a diallyl-sulfonamide **1** to afford a 2,5-dihydro-pyrrole **2**. **II**, Deallylation of allyl carbamate-protected coumarin **3** to the corresponding amino coumarin **4**. **III**, Deallylation of allyl carbamate indole **5** to indole **6**. **IV**, Hydroamination of 2-ethynylaniline **7** to indole **6**. **V**, hydroarylation of profluorophore **8** to afford amino coumarin **9**. **b**, Structures of the biotinylated cofactors employed in this study. Biot refers to D-biotin as depicted in Figure 1a. **c,** Cell-specific activity of 400 ArMs mutated at positions 112 and 121 in Sav normalized to the activity of wild-type Sav (S112 K121). The displayed activities correspond to the product concentrations after 20 h of reaction and are the mean of biological duplicate reactions. The corresponding standard deviations between replicates can be found in Supplementary Figure 8. **d**, Activity distribution in the Sav mutant library for the five different ArM reactions. Violins comprise all 400 double mutants with the ten most active ArMs depicted as circles. **e,** Validation of "hits" from the 400-mutant screens. Bars correspond to the mean activity of eight biological replicates with standard deviation shown as error bars and individual replicates (circles). Mutants are designated by the amino acids at position 112 and 121.

While the screening was designed towards increased ArM activity in a cellular environment, we sought to investigate whether the identified variants also display an increased activity *in vitro*. We therefore purified the three most active mutants for each reaction and performed ArM reactions *in vitro* (Figure 3). Notably, all mutants significantly outperformed the wild-type ArM as well as the corresponding free cofactor, as reflected by markedly higher turnover numbers (TONs). Depending on the reaction, the best variants reached between five- and twenty-fold higher TONs than the free cofactor, demonstrating the benefit of embedding these cofactors within an engineered protein scaffold. It should be noted that in some cases, the relative ranking of variants changed compared to the whole-cell biotransformations, which is likely due to the different reaction conditions and the absence of cellular components *in vitro* (Methods). Nevertheless, the *in vitro* experiments further corroborated the potential of the periplasmic screening platform for the rapid and straightforward discovery of different ArMs with improved catalytic properties and confirmed that the identified variants indeed have a significantly increased specific activity.

Figure 3 | *In vitro* turnover number of ArM variants identified in the periplasmic screening. For each reaction, the three most active variants identified in the whole-cell screening were purified and their TON was determined. Reactions were carried out at 37 °C and 200 rpm for 20 h. Bars represent mean TONs of three technical triplicate reactions with standard deviation as error bars and individual replicates as circles. For comparison, the free cofactor and wild-type Sav variant (wt) were included.

Analysis of sequence-activity landscapes

The recording of sequence-activity data as performed herein provides the basis for additional insight beyond the mere identification of active mutants. To this end, we first sought to understand which biophysical properties result in the observed activity changes. Therefore, we analyzed the effect of individual amino acids on the activity for the individual ArM reactions (Fig. 4a): First, we scaled the activity measurements (Fig. 2c) such that the average activity across the 400 mutant library equals zero and the corresponding standard deviation equals one. Next, we averaged the scaled activities of all 20 variants that carry the same amino acid in one position. This was performed for all 20 canonical amino acids and either of the two diversified positions. As a result, amino acid- and position-specific averages greater or smaller than zero represent a positive or negative effect on ArM catalysis, respectively. While effects of amino acids varied between positions and reactions (Fig. 4a), an overall positive effect of hydrophobic amino acids on the activity of the corresponding ArMs could be observed, whereas charged amino acids as well as proline tend towards a negative impact. Additionally, we used hierarchical clustering to analyze which amino acids have similar effects on activity (Methods). In line with the previous observation, hydrophobic amino acids clustered clearly separated from hydrophilic ones, indicating that this property is crucial in terms of catalytic activity (Figure 4b). Further, amino acids with similar chemical properties showed a strong tendency to cluster closely together (note for instance L-I-V, D-E, H-K and S-T clusters). This points to an anticipated large potential of library strategies that employ reduced amino acid alphabets to maintain a large chemical diversity while significantly reducing the screening effort (see below).

Next, we compared the full-factorial screening approached as pursued herein to other common enzyme engineering strategies. To this end, we used our screening results to analyze which variants would have been identified using such heuristics, and how their activity compares to the most active variant identified in our exhaustive screening. As highlighted in Figure 4c, screening only single mutants (S112X K121 or S112 K121X) would have led to "hits" with activities ranging from 26 to 65 % relative to the most active double mutant for the respective ArM reaction. Subsequent combination of the most beneficial single mutations would have led to an activity increase in two cases, but a decrease in three cases. This suggests that there are interactions between the two amino acid positions 112 and 121 that lead to non-additive effects on the overall activity[38]. Indeed, further analysis revealed that between 24 and 37 % of the observed variance can be attributed to interactions between the two positions (Supplementary Table 1). Notably, the activity of the best mutants identified in our screening can only be explained by considering non-additive effects (Supplementary Fig. 9).

Iterative saturation mutagenesis (ISM) relies on the sequential randomization of positions, while using the best variant from the previous round as a starting point for the next[39]. In this way, it can leverage non-additive interactions to some extent while keeping the experimental effort limited. Indeed, this strategy would have been more successful than the simple combination of single mutations and would have led to the discovery of mutants at or near the local activity optimum for two out of five reactions. Nonetheless, this

demonstrates that a full-factorial library enables the discovery of active mutants that are the result of non-
trivial interactions, which would likely be missed using other strategies. In addition, note that while sequential
strategies require the analysis of fewer variants, they do not provide universally applicable libraries, which
would be a substantial disadvantage for a general ArM screening platform as proposed herein.

Unfortunately, full-factorial screening quickly becomes intractable if more positions are to be diversified. An
established strategy to decrease library size is to use reduced amino acid sets, such as the twelve amino
acids encoded by NDT codons, which still covers all major classes of amino acid chemistries and lacks
redundant or stop codons[40]. Our analysis shows that this strategy would have been more successful than
those discussed above, reaching consistently high activities for all ArM reactions. This observation is in line
with the fact that these twelve amino acids cover the clusters identified in our previous analysis well (Figure
4b). For this reason, it seems promising to screen combinatorial libraries of Sav mutants based on such a
reduced amino acid set. Further reductions of the amino acid set are possible (for example eight amino
acids encoded by NRT), but at the cost of a reduced probability of finding highly active mutants.

Figure 4 | Analysis of amino acid effects and implications for directed evolution. a, Average effect of amino acids on ArM activity by reaction and position. Values outside the dark grey circle indicate a positive mean effect on activity, whereas values inside of it denote a negative effect. Values were standardized by subtracting the mean of the relative activity of all 400 mutants for the respective reaction and dividing by the corresponding standard deviation. The mean across all 20 variants harboring the respective amino acid at the indicated position is shown. **b**, Hierarchical clustering of amino acids based on the activity of variants

containing these residues at position 112 or 121 across all reactions (Methods). Amino acids resulting in similar activity patterns
are grouped at lower heights than more dissimilar ones. Amino acids that are encoded by the degenerate codon NDT are highlighted
in blue. **c**, Comparison of enzyme engineering strategies based on the results presented in this study. The success of various
strategies was compared to the full-factorial approach applied here (from left to right): Screening single mutants for both positions
individually, combining the best mutations of the two individual positions, iterative saturation mutagenesis (ISM) as well as screening
combinatorial libraries based on reduced amino acid sets encoded by NDT (see 4b) and NRT (C, D, G, H, N, R, S, Y) codons. The
activity of the best mutant identified using the respective strategy is displayed relative to the best double mutant among all 400
tested for each ArM reaction. Grey bars represent the mean across all "hits" for the five reactions as identified by the different
strategies.

Discussion

The results presented in this study highlight that Sav-based ArMs can be tailored to catalyze various new-
to-nature reactions. Further, all reactions tested in the context of this study were found to be compatible
with whole-cell biotransformations under close-to-physiological conditions, which is an important property
on the way towards advanced applications in the context of synthetic biology[5]. Our results show that the
activity landscapes of ArMs are distinct for each reaction, underscoring the need for flexible enzyme
engineering strategies that can be easily adapted to new reactions. The Sav 112X 121X library along with
the automation-compatible, periplasmic screening platform represents an important step in this direction.

Based on the analyses presented in this study, it seems reasonable to assume that differences in the
expression level of Sav mutants do not have a major influence on the screening results using our platform,
as an excess of biotin binding sites is consistently ensured. At the same time, the results also suggest that
there is substantial potential for increasing the catalytic capacity of each cell by adding higher
concentrations of cofactor and increasing the uptake capacity for applications in biocatalysis.

Our results highlight that combinatorial libraries focused on crucial residues are a powerful tool for the
engineering of ArMs, particularly when multiple catalytic activities are being pursued originating from a
common starting point. The combination of such libraries with the screening platform presented herein will
enable the rapid discovery of active ArMs for any new reaction as an initial step for more extensive
engineering campaigns. For the latter, the results from this study provide valuable implications such as the
indicated high potential of reduced amino acid sets in the context of ArMs. Combined with lab automation,
such amino acid sets render larger screening campaigns involving more amino acid positions feasible, while
maintaining the advantages of a universal library that can be used for any reaction of interest. For much
larger numbers of targeted positions in Sav, ISM, potentially in combination with reduced amino acid
alphabets, appears to be a good alternative since it offers a good trade-off between experimental effort and
the probability of finding highly active mutants. Lastly, as an alternative to the exhaustive search pursued
in this study, the screening platform presented herein may be used to perform undersampling of libraries
with a much larger theoretical diversity. In combination with powerful machine learning techniques, which
have been ascribed a high potential for enzyme engineering in general[41], this may enable smartly guided
and efficient directed evolution approaches for ArMs.

Acknowledgments

TRW thanks the ERC for the award of an advanced grant (DrEAM, Advanced Grant 694424), the NCCR "Molecular Systems Engineering" and the SNF (Grant 200020_182046) for generous support. MJ thanks ETH Zurich for generous support. TV acknowledges support by the SNF (Grant 31003A_179521) and by the NCCR "Molecular Systems Engineering". The authors kindly thank Valerio Sabatino, Fabian Schwizer, Yoann Cotelle and Jaicy Vallapurackal for their help with substrate and cofactor synthesis, and Gregor Schmidt for technical support regarding lab automation.

Author Contributions

MJ and TRW conceived and supervised the study. SP advised the project. TV carried out biological and screening experiments and analyzed the data. FC and MP developed the gold-catalyzed reactions *in vitro* and synthesized the corresponding substrates and cofactors. TV, MJ and TRW discussed the data. TV and MJ wrote the manuscript with input from all authors.

Materials and Methods

Chemicals and reagents. Unless stated otherwise, chemicals and primers were obtained from Sigma-Aldrich, Acros or Fluorochem. Enzymes for molecular cloning were obtained from New England Biolabs.

Cloning of Sav library. The Sav library was created based on a previously described expression plasmid that contains a T7-tagged Sav gene with an N-terminal *ompA* signal peptide for export to the periplasm under control of the T7 promoter in a pET30b vector (Addgene #138589)[26]. Positions 112 and 121 were randomized using the codon set described by Tang et al[42]. The plasmid was amplified in two fragments using either primer 1 and a mix of primers 2-5 (molar ratio of 12:6:1:1) or primer 10 and a mix of primers 6-9 (molar ratio of 12:6:1:1, see Supplementary Tab. 2 for primer sequences). PCRs were carried out using Q5 polymerase. Template plasmid was digested using DpnI and the PCR products were purified using a PCR purification kit (Sigma-Aldrich). Subsequently, the fragments were joined by Gibson assembly and used to transform chemically competent BL21-Gold(DE3) cells (Agilent Technologies). Individual clones were sequence-verified by Sanger sequencing (Microsynth AG). Having identified 250 out of 400 possible distinct double mutants this way, a second pool containing only the 150 missing variants was cloned. To this end, sequence-verified plasmids from the first library were used as PCR templates and 40 fragments, each containing a single amino acid exchange at position 112 or 121, were obtained by amplification with primers 1 and 12 or 10 and 11 for positions 112 and 121, respectively. Following DpnI digest and purification, these fragments were added to Gibson assembly reactions in combinations suitable for obtaining pools of missing variants. More specifically, this was achieved by adding one fragment with a desired mutation at position 112 per reaction, along with multiple fragments for position 121. Again, individual clones were sequenced after transformation. Eventually, 36 remaining variants were cloned

individually by assembling the corresponding fragments. Refer to Supplementary table 1 for an overview of
the library generation.

Sav expression in 96 well plates. 96 deep-well plates were filled with 500 µL LB (+ 50 mg L^{-1} kanamycin)
per well. Cultures were inoculated from glycerol stocks and grown overnight at 37 °C and 300 rpm in a
Kuhner LT-X shaker (50 mm shaking diameter). 20 µL per culture were used to inoculate expression
cultures in 1 mL LB with kanamycin. These cultures were grown at 37 °C and 300 rpm for 1.5 h. At this
point, the plates were placed at room temperature for 20 min and subsequently Sav expression was induced
by addition of 50 µM isopropyl β-D-1-thiogalactopyranoside (IPTG). Expression was carried out at 20 °C
and 300 rpm for an additional 20 h.

Quantification of biotin-binding sites. For measurement of Sav expression levels, the OD$_{600}$ of the
cultures was determined in a plate reader (Infinite M1000 PRO, Tecan Group AG) using 50 µL samples
diluted with an equal volume of PBS. The remaining cultures were then centrifuged (3,220 rcf, 20 °C, 10
min) and the pellets were resuspended in 250 µL lysis buffer (50 mM Tris, 150 mM NaCl, 5 mM EDTA,
1 g L^{-1} lysozyme, pH 7.4). After 30 min incubation at room temperature, the cell suspensions were subjected
to three freeze-thaw cycles. Afterwards, 150 µL of DNaseI buffer (50 mM Tris, 150 mM NaCl, 1 µL mL^{-1}
DNaseI, pH 7.4) were added and plates were incubated at 37 °C for 45 min before centrifugation (4800 rcf,
20 °C, 20 min). Subsequently, the concentration of biotin binding sites in the supernatant was determined
using a modified version of the assay described by Kada *et al.*[32], which relies on the quenching of the
fluorescence of a biotinylated fluorophore upon binding to Sav. Specifically, 190 µL binding site buffer (1 µM
biotin-4-fluorescein, 0.1 g L^{-1} bovine serum albumin in PBS) were mixed with 10 µL supernatant or purified
Sav standard. After incubation at room temperature for 90 min, the fluorescence intensity was measured
(excitation at 485 nm, emission at 525 nm) and a calibration curve produced with purified Sav was used to
calculate the concentration of biotin binding sites in each sample.

Synthesis of cofactors and substrates. Substrates **1**[23], **3**[27] and cofactors **(Biot-NHC)Ru**[43] and **(Biot-**
HQ)CpRu[27] were prepared according to reported procedures. Substrate **7** was obtained from Sigma-
Aldrich. A detailed description of the synthesis of **(Biot-NHC)Au1**, **(Biot-NHC')Au2**, substrate **8** and product
9 is available in the Supplementary Notes.

Whole-cell screening. Following the expression of Sav mutants in deep-well plates, the OD$_{600}$ of the
cultures was determined using 50 µL samples diluted with an equal volume of PBS. Afterwards, the plates
were centrifuged (3,220 rcf, 15 °C, 10 min), the supernatant was discarded and the pellets were
resuspended in 400 µL incubation buffer containing the respective reaction-specific cofactor. The
composition of the incubation buffer varied for metathesis (2 µM **(Biot-NHC)Ru** in 50 mM Tris, 0.9 % (w/v)
NaCl, pH 7.4), deallylation (5 µM **(Biot-HQ)CpRu** in 50 mM MES, 0.9 % NaCl, pH 6.1), hydroamination
(10 µM **(Biot-NHC)Au1** in 50 mM MES, 0.9 % NaCl, 5 mM diamide, pH 6.1) and hydroarylation (10 µM
(Biot-NHC')Au2 in 50 mM MES, 0.9 % NaCl, 5 mM diamide, pH 5) reactions. The compositions of different
buffers are summarized in Supplementary Table 3. Cells were incubated with the cofactor for 1 h at 15 °C

and 300 rpm. Afterwards, plates were centrifuged (2,000 rcf, 15 °C, 10 min), the supernatant was discarded and the pellets were resuspended in 500 µL of the respective incubation buffer lacking cofactor to remove unbound cofactor. Following another centrifugation step, cell pellets were resuspended in 200 µL reaction buffer containing the respective substrate. The composition of this reaction buffer varied for metathesis (5 mM **1** in 100 mM sodium acetate, 0.5 M MgCl$_2$, pH 4), deallylation (500 µM **3** or **5** in 50 mM MES, 0.9 % NaCl, pH 6.1), hydroamination (5 mM **7** in 50 mM MES, 0.9 % NaCl, 5 mM diamide, pH 6.1) and hydroarylation (5 mM **9** in 50 mM MES, 0.9 % NaCl, 5 mM diamide, pH 5) reactions (Supplementary Tab. 3). Reactions were performed at 37 °C and 300 rpm for 20 h before determining the product concentration. Each 96-well plate contained 80 mutants to be tested along with four replicates of cells expressing wild-type Sav, Sav SL, Sav MA and cells lacking Sav as controls. To account for plate-to-plate variations, the product concentrations were normalized to the mean of the product concentrations measured for the Sav SL (metathesis, hydroamination and hydroarylation) or Sav MA (deallylation) controls on the respective plate. These mutants had been identified as active in preliminary experiments. All variants were tested at least in biological duplicates.

Lab automation. The steps required for incubation of cells with cofactor, washing, substrate addition and OD measurement were implemented on two connected Tecan Freedom EVO 200 Robots, equipped with a Kuhner deep-well plate shaker for incubations, a Rotanta 46 RSC centrifuge and a Tecan Infinite M200 PRO plate reader for OD measurement. The automated protocol was used for the deallylation with allyl carbamate coumarin and the gold-catalyzed hydroarylation, while the screenings for the other reactions were performed manually.

Product quantification by UPLC-MS. To quantify the metathesis product **2**, an extraction was performed by adding 775 µL methanol and 25 µL internal standard (100 µM in methanol) to each 200 µL sample. The samples were incubated for one hour at 20 °C and 300 rpm, followed by centrifugation at 3,220 rcf and 20 °C for 10 min. Subsequently, 50 µL of the supernatant were mixed with 200 µL water and analyzed by UPLC-MS. UPLC analysis was performed using a Waters H-Class Bio using a BEH C18 1.7 µM column and a flow rate of 0.6 ml min^{-1} (eluent A, 0.1 % formic acid in water; eluent B, 0.1% formic acid in acetonitrile; gradient at 0 min: 90 % A, 10 % B; at 0.5 min: 90 % A, 10 % B; at 2.5 min: 10 % A, 90 % B; at 3.5 min: 90 % A, 10 % B; at 4.5 min: 90 % A, 10 % B). Peak integration for SIR (single ion recording) were used for quantification, and concentrations of the metathesis product (retention time of 1.0 min ± 0.25 min) were determined on the basis of a standard curve with the ring-closed product in the presence of a fixed concentration of the nonadeuteraded product (retention time of 1.0 min ± 0.25 min).

Fluorescence measurements. The fluorescent product **4** was quantified by measuring the fluorescence intensity at an excitation of 394 nm and emission of 460 nm. Product **9** was excited at 390 nm and measured at 488 nm. Measurements were carried out in black 96 well plates in an Infinite M1000 PRO plate reader (Tecan Group AG).

Kovac's assay. Indole was quantified using the photometric Kovac's assay (adapted from Piñero-Fernandez *et al.*[44]). For measurements in culture supernatant, plates were centrifuged (3,220 rcf, 20 °C, 10 min) and 110 µL supernatant was mixed with 165 µL Kovac's reagent (50 g L^{-1} 4-(dimethylamino)benzaldehyde, 710 g L^{-1} isoamylic alcohol, 240 g L^{-1} hydrochloric acid) in a separate plate. After 5 min incubation, these plates were centrifuged (3,220 rcf, 20 °C, 10 min). Subsequently, 75 µL of the upper phase were transferred to a new transparent plate and the absorbance at 540 nm was measured in a plate reader (Infinite M1000 PRO, Tecan Group AG). The same procedure (omitting the first centrifugation step) was applied to measure indole after *in vitro* experiments. A standard curve was then used to determine indole concentrations.

Sav expression for purification. A single colony of BL21-Gold(DE3) harboring a plasmid from the previously described library for periplasmic expression of the desired Sav variant was used to inoculate a starter culture (4 mL LB with 50 mg L^{-1} kanamycin), which was grown overnight at 37 °C and 200 rpm. On the following day, 100 mL LB with kanamycin in a 500 mL flask were inoculated to an OD$_{600}$ of 0.01. The culture was grown at 37 °C and 200 rpm until it reached an OD$_{600}$ of 0.5. At this point, the flask was placed at room temperature for 20 min and 50 µM IPTG were added to induce Sav expression. Expression was performed at 20 °C and 200 rpm overnight, and cells were harvested by centrifugation (3,220 rcf, 4 °C, 15 min). Pellets were stored at -20 °C until purification.

Sav purification. Cell pellets were resuspended in lysis buffer (50 mM Tris, 150 mM NaCl, 1 g L^{-1} lysozyme, pH 7.4) added to an amount so as to reach a final OD$_{600}$ of 10. After 30 min of incubation at room temperature, cell suspensions were subjected to three freeze-thaw cycles. Subsequently, nucleic acids were digested by addition of 1 µL benzonase (Merck KGaA) and 10 µL 1 M MgSO$_4$ followed by incubation at room temperature for 10 min. After centrifugation, the supernatant was transferred to a new tube and mixed with carbonate buffer (50 mM ammonium bicarbonate, 500 mM NaCl, pH 11) in a ratio of 2:3. 500 µL iminobiotin beads were added and the tubes were incubated at room temperature with shaking (120 rpm) for 1 h. Afterwards, the beads were washed twice with 15 mL carbonate buffer and resuspended in 2 mL acetate buffer (50 mM ammonium acetate, 500 mM NaCl, pH 4). After 20 min of incubation at room temperature with shaking, the tubes were centrifuged and the supernatant was dialyzed (SnakeSkin dialysis tubing with 3.5 kDa molecular weight cut-off, Thermo Fisher Scientific) against 1 L of the desired buffer (Supplementary Table 4) for 24 h at room temperature with one buffer exchange.

***In vitro* catalysis.** *In vitro* reactions were performed with 2.5 µM purified Sav (tetrameric; corresponding to 10 µM biotin binding sites) in a volume of 200 µL in glass vials. Cofactor and substrate concentrations as well as buffer conditions varied between reactions and are listed in Supplementary Table 4. Reactions were carried out at 37 °C and 200 rpm for 20 h.

Data analysis. Data were analyzed using R 3.6.1[45]. For clustering and calculating amino acid effects, activity values were standardized by subtracting the mean and dividing by the standard deviation of all activity values for the respective reaction. Hierarchical clustering was done by calculating the Euclidian

distances between all standardized activity values of variants harboring the respective amino acid at position 112 or 121 (20 values per position and reaction, amounting to 200 values in total) and clustering these using the hclust function[45] with complete linkage. The activity of all 400 mutants was further analyzed by two-way analysis of variance (ANOVA). To this end, position 112 and 121 were treated as explanatory variables (with 20 levels corresponding to the canonical amino acids). For analyzing the variance explained by individual factors, an interaction term was included, whereas it was omitted in order to generate a purely additive model.

Data and code availability. All data and code supporting the findings of this study are available from the authors upon request.

References

1. Lewis, J. C. Artificial metalloenzymes and metallopeptide catalysts for organic synthesis. *ACS Catal.* **3**, 2954–2975 (2013).

2. Rosati, F. & Roelfes, G. Artificial metalloenzymes. *ChemCatChem* **2**, 916–927 (2010).

3. Bornscheuer, U. T. The fourth wave of biocatalysis is approaching. *Philos. Trans. R. Soc. A Math. Phys. Eng. Sci.* **376**, 20170063 (2018).

4. Hammer, S. C., Knight, A. M. & Arnold, F. H. Design and evolution of enzymes for non-natural chemistry. *Curr. Opin. Green Sustain. Chem.* **7**, 23–30 (2017).

5. Vornholt, T. & Jeschek, M. The quest for xenobiotic enzymes - from new enzymes for chemistry to a novel chemistry of life. *ChemBioChem* (2020). doi:10.1002/cbic.202000121

6. Kan, S. B. J., Lewis, R. D., Chen, K. & Arnold, F. H. Directed evolution of cytochrome c for carbon–silicon bond formation: Bringing silicon to life. *Science* **354**, 1048–1051 (2016).

7. Tinoco, A., Steck, V., Tyagi, V. & Fasan, R. Highly diastereo- and enantioselective synthesis of trifluoromethyl-substituted cyclopropanes via myoglobin-catalyzed transfer of trifluoromethylcarbene. *J. Am. Chem. Soc.* **139**, 5293–5296 (2017).

8. Song, W. J. & Tezcan, F. A. A designed supramolecular protein assembly with in vivo enzymatic activity. *Science* **346**, 1525–1528 (2014).

9. Studer, S. *et al.* Evolution of a highly active and enantiospecific metalloenzyme from short peptides. *Science* **362**, 1285–1288 (2018).

10. Drienovská, I. *et al.* Design of an enantioselective artificial metallo-hydratase enzyme containing an unnatural metal-binding amino acid. *Chem. Sci.* **8**, 7228–7235 (2017).

11. Cangelosi, V. M., Deb, A., Penner-Hahn, J. E. & Pecoraro, V. L. A de novo designed metalloenzyme for the hydration of CO2. *Angew. Chem. Int. Ed.* **53**, 7900–7903 (2014).

12. Key, H. M., Dydio, P., Clark, D. S. & Hartwig, J. F. Abiological catalysis by artificial haem proteins containing noble metals in place of iron. *Nature* **534**, 534–537 (2016).

13. Yang, H. *et al.* Evolving artificial metalloenzymes via random mutagenesis. *Nat. Chem.* **10**, 318–324 (2018).

14. Mirts, E. N., Petrik, I. D., Hosseinzadeh, P., Nilges, M. J. & Lu, Y. A designed heme-[4Fe-4S] metalloenzyme catalyzes sulfite reduction like the native enzyme. *Science* **361**, 1098–1101 (2018).

15. Raines, D. J. *et al.* Redox-switchable siderophore anchor enables reversible artificial metalloenzyme assembly. *Nat. Catal.* **1**, 680–688 (2018).

16.   Eda, S. *et al.* Biocompatibility and therapeutic potential of glycosylated albumin artificial
metalloenzymes. *Nat. Catal.* **2**, (2019).

17.   Heinisch, T. & Ward, T. R. Artificial metalloenzymes based on the biotin-streptavidin technology:
challenges and opportunities. *Acc. Chem. Res.* **49**, 1711–1721 (2016).

18.   Nödling, A. R. *et al.* Reactivity and selectivity of iminium organocatalysis improved by a protein
host. *Angew. Chem. Int. Ed.* **57**, 12478–12482 (2018).

19.   Hassan, I. S. *et al.* Asymmetric δ-lactam synthesis with a monomeric streptavidin artificial
metalloenzyme. *J. Am. Chem. Soc.* **141**, 4815–4819 (2019).

20.   Zeymer, C. & Hilvert, D. Directed evolution of protein catalysts. *Annu. Rev. Biochem.* **87**, 131–157
(2018).

21.   Zhang, R. K., Romney, D. K., Kan, S. B. J. & Arnold, F. H. Directed evolution of artificial
metalloenzymes: bridging synthetic chemistry and biology. in *Artificial metalloenzymes and*
metalloDNAzymes in catalysis 137–170 (Wiley-VCH Verlag GmbH & Co. KGaA, 2018).
doi:10.1002/9783527804085.ch5

22.   Reetz, M. T. Directed evolution of artificial metalloenzymes: a universal means to tune the
selectivity of transition metal catalysts? *Acc. Chem. Res.* (2019).
doi:10.1021/acs.accounts.8b00582

23.   Markel, U., Sauer, D. F., Schiffels, J., Okuda, J. & Schwaneberg, U. Towards the evolution of
artificial metalloenzymes—a protein engineer's perspective. *Angew. Chem. Int. Ed.* **58**, 4454–4464
(2019).

24.   Creus, M. *et al.* X-ray structure and designed evolution of an artificial transfer hydrogenase. *Angew.*
Chem. Int. Ed. **47**, 1400–1404 (2008).

25.   Hyster, T. K., Knorr, L., Ward, T. R. & Rovis, T. Biotinylated Rh(III) complexes in engineered
streptavidin for accelerated asymmetric C-H activation. *Science* **338**, 500–503 (2012).

26.   Jeschek, M. *et al.* Directed evolution of artificial metalloenzymes for in vivo metathesis. *Nature* **537**,
661–665 (2016).

27.   Heinisch, T. *et al.* E. coli surface display of streptavidin for directed evolution of an allylic deallylase.
Chem. Sci. **9**, 5383–5388 (2018).

28.   Grimm, A. R. *et al.* A whole cell E. coli display platform for artificial metalloenzymes:
poly(phenylacetylene) production with a rhodium-nitrobindin metalloprotein. *ACS Catal.* **8**, 2611–
2614 (2018).

29.   Donnelly, A. E., Murphy, G. S., Digianantonio, K. M. & Hecht, M. H. A de novo enzyme catalyzes a
life-sustaining reaction in Escherichia coli. *Nat. Chem. Biol.* **14**, 253–255 (2018).

30.   Wilson, Y. M., Dürrenberger, M., Nogueira, E. S. & Ward, T. R. Neutralizing the detrimental effect
of glutathione on precious metal catalysts. *J. Am. Chem. Soc.* **136**, 8928–8932 (2014).

31.   Jeschek, M., Panke, S. & Ward, T. R. Periplasmic screening for artificial metalloenzymes. in
Methods in Enzymology **580**, 539–556 (Elsevier Inc., 2016).

32.   Kada, G., Kaiser, K., Falk, H. & Gruber, H. J. Rapid estimation of avidin and streptavidin by
fluorescence quenching or fluorescence polarization. *Biochim. Biophys. Acta* **1427**, 44–8 (1999).

33.   Tokuriki, N. & Tawfik, D. S. Stability effects of mutations and protein evolvability. *Curr. Opin. Struct.*
Biol. **19**, 596–604 (2009).

34.   Völker, T., Dempwolff, F., Graumann, P. L. & Meggers, E. Progress towards bioorthogonal
catalysis with organometallic compounds. *Angew. Chem. Int. Ed.* **53**, 10536–10540 (2014).

35.   Völker, T. & Meggers, E. Chemical activation in blood serum and human cell culture: improved

ruthenium complex for catalytic uncaging of alloc-protected amines. *ChemBioChem* **18**, 1083–1086
(2017).

36. Do, J. H., Kim, H. N., Yoon, J., Kim, J. S. & Kim, H. J. A rationally designed fluorescence turn-on
probe for the gold(III) ion. *Org. Lett.* **12**, 932–934 (2010).

37. Sasmal, P. K., Streu, C. N. & Meggers, E. Metal complex catalysis in living biological systems.
Chem. Commun. **49**, 1581–1587 (2013).

38. Reetz, M. T. The importance of additive and non-additive mutational effects in protein engineering.
Angew. Chem. Int. Ed. **52**, 2658–2666 (2013).

39. Reetz, M. T. & Carballeira, J. D. Iterative saturation mutagenesis (ISM) for rapid directed evolution
of functional enzymes. *Nat. Protoc.* **2**, 891–903 (2007).

40. Reetz, M. T., Kahakeaw, D. & Lohmer, R. Addressing the numbers problem in directed evolution.
ChemBioChem **9**, 1797–1804 (2008).

41. Yang, K. K., Wu, Z. & Arnold, F. H. Machine-learning-guided directed evolution for protein
engineering. *Nat. Methods* **16**, 687–694 (2019).

42. Tang, L. *et al.* Construction of 'small-intelligent' focused mutagenesis libraries using well-designed
combinatorial degenerate primers. *Biotechniques* **52**, 149–158 (2012).

43. Kajetanowicz, A., Chatterjee, A., Reuter, R. & Ward, T. R. Biotinylated metathesis catalysts:
synthesis and performance in ring closing metathesis. *Catal. Letters* **144**, 373–379 (2014).

44. Piñero-Fernandez, S., Chimerel, C., Keyser, U. F. & Summers, D. K. Indole transport across
Escherichia coli membranes. *J. Bacteriol.* **193**, 1793–1798 (2011).

45. R Core Team. R: A language and environment for statistical computing. *R Foundation for Statistical*
Computing (2019). Available at: https://www.r-project.org/.