Note: This is a pre-print that has not undergone peer-review.

Variability in infants' functional brain network connectivity is associated with differences in affect and behavior

Caroline M. Kelsey[1,2], Katrina Farris[2,3] & Tobias Grossmann[2]

[1] Department of Pediatrics, Division of Developmental Medicine, Boston Children's Hospital,

Boston, MA, USA

[2] Department of Psychology, University of Virginia, Charlottesville, VA, USA

[3] Department of Psychology, Georgia State University, Atlanta, GA, USA

Correspondence:

Tobias Grossmann

Department of Psychology

PO BOX 400400

University of Virginia

Charlottesville, VA 22904

Tg3ny@virginia.edu

Keywords: fNIRS, Functional connectivity, Default Mode Network, Fronto-parietal Network, Infancy, Temperament.

Abstract

Variability in functional brain network connectivity has been linked to individual differences in cognitive, affective, and behavioral traits in adults. However, little is known about the developmental origins of such brain-behavior correlations. The current study examined functional brain network connectivity and its link to behavioral temperament in newborn and 1-month-old infants (*M* [age] = 25 days; N = 75) using functional near-infrared spectroscopy (fNIRS). Specifically, we measured long-range connectivity between cortical regions approximating fronto-parietal, default mode, and homologous-interhemispheric networks. Our results show that connectivity in these functional brain networks varies across infants and maps onto individual differences in behavioral temperament. Specifically, connectivity in the fronto-parietal network was positively associated with regulation and orienting behaviors, whereas connectivity in the default mode network showed the opposite effect on these behaviors. Our analysis also revealed a significant positive association between the homologous-interhemispheric network and infants' negative affect. The current results suggest that variability in long-range intra-hemispheric and cross-hemispheric functional connectivity between frontal, parietal and temporal cortex is associated with individual differences in affect and behavior. These findings shed new light on the brain origins of individual differences in early-emerging behavioral traits and thus represent a viable novel approach for investigating developmental trajectories in typical and atypical neurodevelopment.

44   **Variability in infants' functional brain network connectivity is associated with differences**

45   **in affect and behavior**

47   Spontaneous brain activity is characterized by intrinsic dynamics of synchronized low-

frequency fluctuations within structurally and functionally connected brain networks

(Damoiseaux et al., 2006; Kaiser, Andrews-Hanna, Wager, & Pizzagalli, 2015). Much research

has been focused on mapping the human connectome and delineating its anatomical and

functional properties (Smith et al., 2013). Individual variability in functional connectivity

profiles can accurately identify specific individuals likened to a fingerprint (Finn et al., 2015)

and is linked to individual differences in cognitive, affective and behavioral traits in adults

(Jauniaux, Khatibi, Rainville, & Jackson, 2019; Kaiser, Andrews-Hanna, Wager, et al., 2015;

Picó-Pérez et al., 2019). More generally, the study of functional brain network connectivity has

been argued to be one of the most promising and effective ways in bridging between brain and

behavior (Friston, 2011).

58   From a developmental perspective, functional connectivity within brain networks can be

detected from very early in human brain development. A host of studies employing resting-state

functional magnetic resonance imaging (rs-fMRI) have mapped and identified functional

networks in newborn infants (Fransson et al., 2009; Gao, Alcauter, Smith, Gilmore, & Lin, 2015;

Schöpf, Kasprian, Brugger, & Prayer, 2012; Thomason et al., 2019; Zhang, Shen, & Lin, 2019).

In fact, a body of work relying on progress in fetal rs-fMRI suggests that the basic organization

and architecture of the functional connectome emerges during the late second trimester of

pregnancy (Schöpf et al., 2012; Thomason et al., 2013; Thomason et al., 2019; van den Heuvel &

Thomason, 2016). In addition, already within the first week of life early prenatal and postnatal

experiences, such as premature birth, have begun to shape the development of these network

(Rogers, Lean, Wheelock, & Smyser, 2018; Smyser, Wheelock, Limbrick, & Neil, 2019). The

existing research with fetuses and infants points to a developmental progression whereby

functional connectivity in primary short-range sensory-motor and homologous-interhemispheric

networks are already in place at birth, whereas functional connectivity in higher-order cortical

networks across longer ranges involving frontal, temporal and parietal cortex shows more

protracted development during infancy (Gao et al., 2015; van den Heuvel & Thomason, 2016).

For example, there is evidence to suggest that higher-order networks such as the default mode

network (DMN) exist in a rudimentary form even in the fetus and newborn (Gao, Lin, Grewen,

& Gilmore, 2017; van den Heuvel & Thomason, 2016), but functional network integration and

synchronization continues to develop during infancy and beyond (Gilmore, Knickmeyer, & Gao,

2018). Other higher-order networks, such as the fronto-parietal network, show an even more

protracted development as it is still considered immature by the end of the first year of postnatal

life (see Gao et al., 2017, for a review). Moreover, it is these networks with the prolonged

development, such as the fronto-parietal network, that go on to have the greatest inter-person

variability and are considered to be unique identifiers of individuals (Finn et al., 2015). Taken

together, much progress has been made in mapping the functional connectome in early human

brain development, however, to date, little is known about whether and how functional brain

network connectivity in these networks is linked to early affective, cognitive, and behavioral

traits. This is a particularly important question considering that many mental health disorders are:

(a) accompanied by alterations in functional connectivity (Hu et al., 2017; Kaiser, Andrews-

Hanna, Wager, et al., 2015; Steward, Menchon, Jiménez-Murcia, Soriano-Mas, & Fernandez-

Aranda, 2018) and (b) are argued to have deep developmental origins (Insel, 2010).

Depression as one main neuropsychiatric disorder affecting about 20.6% of the US

population across their lifetime has been linked to specific alterations in functional brain network

connectivity (Hasin et al., 2018). Specifically, a host of studies show that depression in adults is

characterized by: (1) hypoconnectivity within the fronto-parietal network (FPN; composed of

regions in the anterior cingulate cortex, dorsolateral prefrontal cortex, and parietal cortex)

implicated in the cognitive control of attention and emotion (2) hyperconnectivity within the

default mode network (DMN; composed of regions in the medial prefrontal cortex (mPFC), the

precuneus, the posterior and anterior cingulate cortex, the inferior parietal cortex, and the lateral

temporal cortex) involved in internally-oriented thought, mind-wandering, social cognition and

(3) hypoconnectivity within the homologous-interhemispheric network (HIN; examining cross-

hemispheric connections between frontal, temporal, and parietal lobes) involved in emotion

regulation (Banich & Karol, 1992; Kaiser, Andrews-Hanna, Wager, et al., 2015; Patashov,

Goldstein, & Balberg, 2019; Wang et al., 2013). Considering these alterations in functional

connectivity seen in depression and the possibility that they might have their origins in early

human brain development, it is important to examine variability in these brain networks and how

it links to individual differences in affective and behavioral traits.

Therefore, the current study followed two major goals. First, we aimed to identify and

map individual variability in the three functional brain networks (FPN, DMN, HIN) in young

infants using functional near-infrared spectroscopy (fNIRS). FNIRS is a non-invasive, portable,

and safe, optical neuroimaging technique for assessing functional connectivity in cortical brain

networks during infancy (see the following papers for other examples of functional connectivity

analysis using fNIRS with infants Bulgarelli et al., 2019; Bulgarelli et al., 2020; Homae et al.,

2010). To capture functional connectivity patterns in young infants using fNIRS we pre-defined

the following three long-range brain networks including available channels in specific frontal,

temporal and parietal regions: 1) the FPN was created by measuring functional connectivity

between the dorsolateral prefrontal cortex and inferior parietal cortex, 2) the DMN was created

by measuring functional connectivity between the lateral temporal cortex and medial prefrontal

cortex (note that our probe layout did not allow us to measure activity from superior parietal

cortical regions including the precuneus, which is typically included in the DMN), and 3) the

HIN was created by measuring functional connectivity between homologous cross-hemispheric

connections in frontal, temporal, parietal cortex. In addition, based previous work measuring

functional connectivity using fNIRS in adults (Sasai, Homae, Watanabe, & Taga, 2011), we

created a so-called control network, computing functional connectivity between left frontal

cortex and right temporal cortex and right frontal cortex and left temporal cortex. This served as

a non-functional control network, because it is not known to have any functional associations

and shows much lower levels of functional connectivity than established functional brain

networks (Sasai et al., 2011).We hypothesized that functional connectivity within the three

functional brain networks (FPN, DMN and HIN) will be significantly greater than in the control

network, attesting to the existence of these long-range cortical networks in young infants. In this

context, it is important to mention that, to our knowledge, there is no prior work demonstrating

long-range functional connectivity in FPN and DMN in newborns and 1-month-old infants (Gao

et al., 2017; van den Heuvel & Thomason, 2016), whereas functional connectivity in HIN has

been shown to exist in the fetal brain (Thomason et al., 2013).

Critically, we also examined whether and how variability in functional brain network

connectivity maps onto individual differences in infant temperament, which can be readily and

reliably assed through parental report (Rothbart, 2007). Infant and child temperament is

considered to reflect robust, biologically-based individual differences in affective and behavioral

traits (Rothbart, 2007) linked to long-term developmental outcomes in prospective longitudinal

studies, including adult personality, depression and anxiety (Asendorpf, Denissen, & van Aken,

2008; Caspi et al., 2003; Caspi, Moffitt, Newman, & Silva, 1996; Tang et al., 2020). In addition,

brain networks assessed during the first few weeks of life has been linked to behavioral

temperament 6 months later (Graham et al., 2016; Thomas et al., 2019). However, no work to the

authors knowledge have examined the concurrent relation temperament in infants younger than 6

months. Specifically, we focused our investigation on three critical dimensions of infant

temperament (regulation/orienting, negative emotionality, positive emotionality/surgency),

which have been previously identified in a factor analysis (Gartstein & Rothbart, 2003). Based

on prior work with adults linking functional connectivity in FPN to cognitive control of attention

and behavior (Kaiser, Andrews-Hanna, Spielberg, et al., 2015; Kaiser, Andrews-Hanna, Wager,

et al., 2015), we hypothesized that infants' regulation/orienting behaviors will be associated with

functional connectivity in the FPN with greater connectivity in this network being linked to

enhanced regulation and orienting. In contrast, we predicted the opposite pattern of association in

the DMN, lower connectivity being linked to enhanced regulation and orienting, as it has

previously been linked to self-referential, stimulus-independent thought, and mind-wandering in

adults (Kaiser, Andrews-Hanna, Spielberg, et al., 2015; Kaiser, Andrews-Hanna, Wager, et al.,

2015). Moreover, we expected that infants' reduced functional connectivity in the HIN will be

linked to higher levels of negative emotionality, based on previous findings linking reduced

cross-hemispheric connectivity to depression in adults (Patashov et al., 2019; Wang et al., 2013).

Critically, in our analysis, we expected to see these predicted associations only for the specific

functional networks but not for the (non-functional) control network. Finally, considering that

there is no prior work informing how surgency/positive emotionality is linked to network

connectivity, we did not have a specific hypothesis regarding this trait, but still included it in our

analysis because surgency/positive emotionality has been identified as an important factor in

previous work (Gartstein & Rothbart, 2003). Together, the current study presents a systematic

examination of functional connectivity in long-range brain networks and its links to behavioral

temperament in young infants.

Materials and Methods

Seventy-five newborn and one-month-old infants (*M* [age] = 25 days; *Median* [age] = 24

days; ranging from 9 days to 56 days; 32 females; 43 males) were included in the final sample

used for the present analyses. Participants were recruited from a local hospital. The diverse

sample of infants were representative of the surrounding Mid-Atlantic college town area such

that the majority of infants were Caucasian (*n* = 49 Caucasian; *n* = 14 Black; *n* = 3 South Asian;

n = 3 Pacific Islander; *n* = 2 Asian; *n* = 4 Other), from highly-educated parents (*n* = 31 obtained a

Graduate Degree; *n* = 19 Bachelor's Degree; *n* = 12 some College/Associates Degree; *n* = 11

High School Diploma/GED; *n* = 2 some High School), and low to medium-income families (*n* =

21 $15-45,000; *n* = 18 $75-110,000; *n* = 11 $45-75,000; *n* = 11 $110-175,000; *n* = 8 $175,000+;

n = 5 less than $15,000; *n* = 1 did not respond). All participants were born at term, with normal

birth weight (>2,500g), and did not have any hearing or visual impairments. Thirty-three

additional infants were tested but were excluded from the present analyses for the following

reasons: *n* = 25 were excluded because they failed to reach our pre-determined inclusion criterion

of having at least 100 seconds of continuous data with non-disruptive behaviors (see below); *n* =

4 were excluded because of inaccurate placement of the cap; *n* = 4 were excluded because more

than 33% of the measured fNIRS channels had poor light intensity readings, more specifically, a

signal-to-noise ratio of less than 1.5 (Bulgarelli et al., 2019; Xu et al., 2015). Note that the

current attrition rate (30%) is lower than in previous infant fNIRS studies (Cristia et al., 2013).

Moreover, temperament profiles (negative emotionality, regulation/orienting, and

surgency/positive emotionality) were compared using independent samples t-tests between

infants that were included and excluded from the present analyses and no significant differences

were found between the two groups (all *p-values* > .29). All parents gave informed consent for

their infants to participate in accordance with the Declaration of Helsinki and families received a

payment for their participation. All procedures were approved by and carried out in accordance

with The University of Virginia Institutional Review Board for Health Sciences (Protocol

number 20381).

Infant Temperament

Infant temperament was assed using parental reports of the 91-item Infant Behavior

Questionnaire Revised Short Form (IBQ-R; Gartstein & Rothbart, 2003). Parents filled out the

questionnaire online using Qualtrics survey platform prior to their appointment. This measure

has been widely used and shown to be reliable and valid at the newborn time point (see the

following papers for examples of prior work using this measure with newborns Rigato, Stets,

Bonneville-Roussy, & Holmboe, 2018; Stifter & Fox, 1990; Worobey & Blajda, 1989). The

questionnaire asks parents to report their infant's behavior during the previous two weeks and

rate the occurrence/frequency of the behavior on a 1 (Never) to 7 (Always) scale. Based on prior

work using factor analysis (Gartstein & Rothbart, 2003), three general temperament dimensions

were computed summarizing information from various sub-scales: (1) negative emotionality

(contributing sub-scales: fear, distress to limitations, falling reactivity, sadness), (2)

regulation/orienting (contributing sub-scales: low intensity pleasure, cuddliness, duration of

orienting, soothability), and (3) surgency/positive emotionality (contributing sub-scales: activity

level, smiling and laughing, high intensity pleasure, perceptual sensitivity, approach, vocal

reactivity) (Gartstein & Rothbart, 2003). If parents reported the behavior was not applicable at

the current time then this item was given a value of 0. Chronbach's alpha coefficients were

calculated to determine reliability of the temperament measures and all values were in acceptable

ranges for each of the three dimensions: surgency/positive emotionality α = .78,

regulation/orienting α = .78, and negative emotionality α = .91. Finally, correlation analyses

between Edinburgh Postnatal Depression Scale scores (assessed at the same time as behavioral

temperament; Cox, Holden, & Sagovsky, 1987) and behavioral temperament scores were

conducted in order to statistically account for any variance in maternal-reported behavioral

temperament that may be related to maternal mental health. Here, we did not find any significant

associations (all *p-values* > .24). Therefore, maternal depression was not used as a covariate in

later analyses.

Procedure

The resting state fNIRS task took place in a quiet, dimly-lit testing area. Infants were

seated on their parents' lap and placed approximately 60 cm from the screen (23-inch monitor).

The infants were fitted with a fNIRS fabric cap (EasyCap, Germany) which was secured in place

using infant overalls and outside netting. The experimental paradigm was presented using the

Presentation software package (Neurobehavioral Systems, USA). A non-social stimulus was

created by selecting non-social clips from a popular infant video (Baby Einstein - Kids2 Inc.)

that featured videos of toys, stuffed animals, and still images of everyday objects, which was

accompanied by classical music (Lordier et al., 2019). Similar screen-saver-like videos have

been used in prior work examining functional connectivity using fNIRS (see Bulgarelli et al.,

2020). This video was played for a total of seven minutes while fNIRS data were being recorded.

The clips were segmented into 30 second intervals and the order of presentation was randomized

for each infant. Parents were asked to remain quiet throughout the fNIRS recording session.

Sessions were video-recorded using a camera mounted above the screen. This allowed for later

offline coding of infants' alertness and cap placement.

Data acquisition

Infants' fNIRS data were recorded using a NIRx Nirscout system and NirStar acquisition

software. The fNIRS method quantifies concentration changes of oxygenated hemoglobin

(oxyHb) and deoxygenated hemoglobin (deoxyHb) in the cerebral cortex through shining

specific frequencies of light that are selectively absorbed by these chromophores (for more

information regarding this technique see Lloyd-Fox, Széplaki-Köllőd, Yin, & Csibra, 2015). The

fNIRS system used contains 16 source-detector pairs (approximately 2.0 cm apart) resulting in a

total of 49 channels positioned over frontal and temporal-parietal regions (see Altvater-

Mackensen & Grossmann, 2016; Grossmann, Missana, & Krol, 2018; Kelsey, Krol, Kret, &

Grossmann, 2019; Krol, Puglia, Morris, Connelly, & Grossmann, 2019 for infant work using the

identical channel positioning/layout). The system emits two wavelengths of light in the Near-

Infrared spectrum, 760 nm and 850 nm, and captures both deoxyHb and oxyHb. The diodes have

a power of 25 mW/wavelength and data were recorded at a preset default sampling rate of 3.91

Hz.

Behavioral Coding

Infants' behavior during the fNIRS recording session was coded by a trained research

assistant using video recordings of the experimental session. Specifically, coders identified

timepoints where the infants were crying, excessively moving, looking at the parents, or parents

were talking in order to remove these periods from the analysis. To assess the reliability of the

attentional coding done by the primary coder, an additional trained coder also coded infant

behavior from selected subsample of infants (25.3%; $n = 19$). This analysis showed that inter-

rater reliability amount of data included was excellent (Cronbach's $\alpha = 0.94$). In line with

previous studies, infants were only included in the present analysis if they had at least 100

seconds of disruption-free (see aforementioned behaviors) data (Bulgarelli et al., 2019).

Moreover, as it takes a minimum of 8 seconds for the Hemodynamic response function to return

to baseline after a stimulus-evoked event, the onset of useable data was delayed for 8 seconds

(Bulgarelli et al., 2019; Bulgarelli et al., 2020). However, unlike Bulgarelli et al. (2019; 2020),

the time series of fNIRS obtained in the current study was continuous. On average, infants

contributed 317.59 seconds of data ($SD = 115.46$ seconds; range = 100-420 seconds).

Furthermore, the amount of data included in the current analysis is comparable to other

functional connectivity work using fNIRS with older infants (Bulgarelli et al., 2020). In addition,

we coded infants' state of alertness on a 1 (Deep Sleep) to 6 (Crying) scale. On average infants

were rated as being in a Active Light Sleep to Drowsy State ($M = 2.68$, $SD = 1.27$). Finally, we

assessed how infants' behavior throughout the session, specifically the amount of useable data

related to functional network connectivity in each of the networks (see supplemental materials).

Data Analysis

The fNIRS data were analyzed using the functional connectivity program, FC-NIRS (Xu

et al., 2015). First, channels were assessed for light intensity quality and channels were removed

if the signal-to-noise ratio was less than 1.5 (Xu et al., 2015). In order to be included in the

present analyses, infants needed to have at least 70% of their channels passing this threshold

(Bulgarelli et al., 2019). Next, data were band-pass filtered (using a .08 Hz low-pass filter, to

remove fast fluctuations related to heart rate, and a high-pass filter of .01 Hz, to remove changes

that were too slow and related to drift; Bulgarelli et al., 2019; Lu et al., 2009). This range of .01

to .08 Hz was chosen on the basis of prior work (Bulgarelli et al., 2019; Sasai et al., 2011). This

range was also selected because it falls well below the reported range for cardiac fluctuations

(greater than 1 Hz), providing us with greater confidence that the measured changes reflect

hemodynamic events tied to cortical activity rather than (systemic) cardiovascular system

activity (e.g., heart rate Elwell, Springett, Hillman, & Delpy, 1999; Obrig et al., 2000). Finally,

concentration changes were calculated using the modified Beer-Lambert law [partial path length

factor: 6.0] (Villringer & Chance, 1997).

For each infant, we obtained a 49 by 49 correlation matrix corresponding to all of the

relations between all of the channels measured. Considering that negative values are difficult to

interpret in terms of their neurobiological basis, and based on prior work, we first checked data

to see if there were any negative values and found out there were none (Fox, Zhang, Snyder, &

Raichle, 2009; Murphy, Birn, Handwerker, Jones, & Bandettini, 2009). In order to standardize

the values, Fisher Z-transformations were performed on all correlation matrices. Networks of

interest were created by selecting channels that corresponded to specific regions of interest.

Brain networks were composed based on the anatomical information available in Kabdebon et al.

(2014), a meta-analysis of resting state fMRI (Kaiser, Andrews-Hanna, Wager, et al., 2015), a

large resting state fMRI functional connectivity analysis of newborn infants (Eyre et al., 2020),

and prior work infant and adult work using rs-fNIRS (Bulgarelli et al., 2020; Imai et al., 2014;

Patashov et al., 2019; Sasai et al., 2011). Based on this information four networks were created:

(1) The FPN was created by averaging all correlations between three channels in the dorsolateral

prefrontal cortex (corresponding with the F3, F4, F5, F6 electrodes) and two channels in the

parietal area (corresponding with CP3 and CP4 electrodes); (2) The DMN was created by

averaging all correlations between three channels in the medial prefrontal cortex (corresponding

with the Fpz electrode) and four channels in the lateral temporal cortex (corresponding with FT7,

T7, FT8, T8 electrodes); (3) The HIN was created by averaging all correlations between the 21

channels in the left hemisphere (including frontal, temporal and parietal cortical regions) with

their corresponding (homologous) channels in the right hemisphere; and, (4) a (non-functional)

control network was created by averaging all correlations between three channels in the left

frontal area (corresponding with the F7 electrode) with three channels in the right temporal area

(corresponding with the T8 electrode) and three channels in the right frontal area (corresponding

with F8 electrode) with three channels in the left temporal area (corresponding with the T7

electrode; see Figure 1 for schematic of network configurations). Cortical projections onto a

standard MNI newborn (0-2 months old) atlas (Fonov et al., 2011) were created using NIRSite

(Nirx) by using 10-20 system references from the cap layout.

All analyses were conducted for both oxyHb and deoxyHb (for deoxyHb results please

see supplemental materials). Moreover, statistical outliers – values that were more than 3 SD

above the mean – were removed for the subsequent analyses (FPN $n = 2$, negative emotionality n

$= 1$).

Figure 1. Shows the configurations for each of the network patterns in both a 2-dimensional 10-20 system layout and estimated projections onto cortical space of a 0-2 month-old Atlas (Fonov et al., 2011). Note, each network consists of the average of all of the connections between red and blue channels of the same letter. In addition, the orange dots represent relevant 10-20 landmarks.

Figure 2. Shows the channels that are significantly different than zero for each of the networks. Please note, channels in red, blue, and black represent significant changes for oxyHb, deoxyHb, and both oxy and deoxyHb respectively.

Results

      A series of Spearman's rho correlations were used to identify significant associations between variables of interest and potential socio-demographic factors (for a schematic representation for all associations see supplementary Figure 1). Any demographic variables found to be significantly associated with a study variable of interest were then included in the subsequent models assessing differences in said study variable as a covariate. Negative Emotionality was significantly associated with both infant age (Spearman's rho correlation $r_s =$ .47, $p < $ .001) and family income (Spearman's rho correlation $r_s=$ .29, $p = $ .011).

Regulation/orienting was significantly associated with Education (Spearman's rho correlation $r_s=$

$-.30$, $p = .009$). However, there were no significant associations found between any of the

functional connectivity measures and any of the covariates.

Functional connectivity across networks

As a first step, a series of one-sample t-tests were conducted to assess whether Fisher-

transformed correlation between individual channels within the pre-defined networks of interest

differed from zero. As shown in Figure 2, this analysis identified significant functional

connectivity between individual channels within the pre-defined networks of interest (see

supplementary Table 1). Next, we conducted a series of one-sample t-tests to assess connectivity

at the network level (combining across all channels of interest). Here, all networks (FPN, DMN,

HIN, control) were found to be greater than zero (all p's < .001; see Figure 3). To analyze

differences in overall connectivity levels across networks an omnibus repeated measures

ANOVA with network type (FPN, DMN, HIN, control) as a within-subjects factor was

conducted. This analysis revealed a significant within-subjects effect across network types, $F(3,$

$216) = 18.78$, $p < .001$, $\eta^2 = .207$. Post-hoc analyses with Bonferroni adjustments for multiple

comparisons were conducted to assess which networks significantly differed from one another.

Importantly, all functional networks of interest had significantly higher connectivity than the

(non-functional) control network ($M = .05$; $SD = .12$; range: -.20–.44), all p's < .001. In addition,

we found that there was significantly greater connectivity in the FPN ($M = .21$; $SD = .20$; range:

-.16–.72) compared to both the HIN ($M = .13$; $SD = .12$; range: -.12–.48), $p = .003$, and the DMN

($M = .13$; $SD = .16$; range: -.28–.73), $p = .01$. However, there was no significant difference found

between the level of connectivity for the HIN from the DMN, $p = 1.00$ (see Figure 3).

Figure 3. Shows the average levels of functional connectivity (oxyHb) and range of variability

for each network. The boxplot horizontal lines from bottom to top reflect values for the lower

quartile, median, and upper quartile respectively. Note, * $p < .05$.

Functional connectivity and behavioral temperament

In order to assess how functional connectivity patterns differentially predicted

temperament characteristics, three separate regressions with all four network types (FPN, DMN,

HIN, control) predicting each of the three domains of temperament (negative emotionality,

regulation/orienting, surgency/positive emotionality) were conducted.

Regulation/Orienting. A linear regression was conducted with the four network types

(FPN, DMN, HIN, control) predicting regulation/orienting using the entry method. The

regression model significantly predicted regulation/orienting, $F(5, 72) = 4.84$, $p = .001$, $R^2 = .27$.

More specifically, connectivity in the DMN was negatively associated with regulation/orienting

(β = -1.02, *SE* = .42, *p* = .018); whereas, connectivity in the FPN was positively associated with

regulation/orienting (β = .71, *SE* = .35, *p* = .049; see Figure 4). Neither the HIN nor the Control

network were found to be related to regulation/orienting, all *p-values* > .24.

Negative Emotionality. A multiple linear regression was conducted with the socio-

demographic covariates (age, income) and four network types (FPN, DMN, HIN, control)

predicting negative emotionality using the entry method. The model significantly predicted

negative emotionality $F(6, 64)$ = 5.50, $p < .001$, R^2 = .34. More specifically, we found a

significant positive relation between HIN connectivity and negative emotionality, (β = 1.60, *SE* =

.75, *p* = .038; See Figure 5). However, none of the other networks (functional nor control) were

found to be related to Negative Emotionality, all *p-values* > .57.

Surgency/Positive Emotionality. A linear regression was conducted with the four network

types (FPN, DMN, HIN, control) predicting surgency/positive emotionality using the entry

method. Here, the regression model did not significantly predict surgency/positive emotionality,

p = .39. Moreover, none of the network types were significantly associated with

surgency/positive emotionality (all *p's* > .14).

Figure 4. Shows the unadjusted (solid line) and adjusted relation (covariates not shown: HIN,

Control) between functional connectivity (oxyHb) Z-score and regulation/orienting. Here, we

found that connectivity in the FPN was positively associated with regulation/orienting (p = .049)

whereas, connectivity in the DMN was negatively associated with regulation/orienting (p =

.018). Note, shaded regions represent 90% confidence intervals for the raw (unadjusted) data.

Figure 5. Shows the unadjusted (solid line) and adjusted relation (covariates: FPN, DMN,

Control, Income, age) between HIN functional connectivity (oxyHb) Z-score and negative

emotionality. Here, we found a significant positive relation between the HIN and negative

emotionality (p = .038). Note, shaded regions represent 90% confidence interval for the raw

(unadjusted) data.

Discussion

The current study examined functional connectivity in brain networks using fNIRS and

behavioral temperament using parental report in young infants. Our results show that functional

connectivity in long-range cortical brain networks (FPN, DMN and HIN) can be identified in

very young infants and that functional connectivity in these networks varied considerably among

infants. This supports the suitability of fNIRS in assessing functional connectivity and its

variability in newborn infants. Importantly, our results also show that such variability in

functional brain network connectivity systematically maps onto individual differences in infant

behavioral temperament. Overall, the current findings provide novel insights into the brain

origins of individual differences in affect and behavior, pointing to the early perinatal foundation

of human temperament.

In line with our hypothesis, functional connectivity within the three brain networks (FPN,

DMN, and HIN) was significantly greater than in the control network and significantly greater

than a zero-value, indicating the existence of these long-range cortical brain networks in young

infants. This provides further evidence that functional brain networks exist from early in

ontogeny and are detectable in young infants (Gao et al., 2017; Gao et al., 2009; Zhang et al.,

2019). To our knowledge, this is the first study to demonstrate long-range functional

connectivity in FPN and DMN in young infants, suggesting a remarkably early emergence of

long-range connectivity in higher-order brain networks linked to cognitive control and self-

referential processes, respectively. The current findings are noteworthy also in regard to the fact

that both networks involve regions in prefrontal cortex, providing new evidence from newborns

and 1-month-old infants supporting the view that prefrontal cortex plays a critical role in human

brain function from very early in development (Grossmann, 2013a, 2013b; Grossmann, 2015;

Powell, Kosakowski, & Saxe, 2018; van den Heuvel & Thomason, 2016).

In addition to the general difference in connectivity between the functional and the (non-

functional) control network, we also found that activity in the FPN was significantly greater than

in the DMN and HIN (whereas there was no difference in connectivity levels found between the

DMN and HIN). One possible interpretation of this finding is that functional connectivity in the

FPN might have been enhanced when compared to the other functional networks, because like

other resting-state studies with infants, the participants were presented with a video accompanied

by music during the fNIRS measurement (Bulgarelli et al., 2019). In other words, the FPN might

have been more engaged because infants were attending to external audio-visual stimuli (note

that all infants were exposed to the same video [audio-visual] stimulus). Here it is important to

mention that prior work with adults using fMRI shows that functional connectivity in higher-

order cortical resting-state networks can be reliably acquired during the presentation of videos

and corresponds to functional connectivity acquired in the absence of any stimulus (Finn et al.,

2015; Vanderwal, Kelly, Eilbott, Mayes, & Castellanos, 2015). Nonetheless, based on recent

work showing that preterm infants display enhanced functional connectivity in higher-order

cognitive networks in response to music (Lordier et al., 2019), we speculate that enhanced

functional connectivity in FPN might at least be partly explained by having newborn infants

listen to music in the current study. Clearly, future research with infants, which systematically

compares between stimulation protocols, is needed to examine whether and how functional

connectivity is influenced by the measurement context and the stimulation protocol used.

Overall, our functional connectivity analysis supports the notion that intrinsic functional

connectivity in cortical brain networks and its variability can be effectively mapped in newborn

infants using fNIRS.

Having established functional connectivity in these brain networks as variable and

distinct from a (non-functional) control network then allowed for the examination of specific

associations between brain network connectivity and infant behavioral temperament. Our results

confirmed our hypothesis and showed that infants' regulation/orienting behaviors were

associated with functional connectivity in the FPN with greater connectivity in this network

being associated with enhanced regulation and orienting. This result is in line with prior work

linking functional connectivity in FPN to cognitive control of attention and behavior in adults

(Kaiser, Andrews-Hanna, Spielberg, et al., 2015; Kaiser, Andrews-Hanna, Wager, et al., 2015).

The current results further showed the opposite pattern of association for functional connectivity

in DMN, with greater connectivity associated with reduced regulation and orienting, which is in

agreement with our hypothesis based on the DMN previously being linked to self-referential,

stimulus-independent thought and mind-wandering in adults (Kaiser, Andrews-Hanna, Spielberg,

et al., 2015; Kaiser, Andrews-Hanna, Wager, et al., 2015). To obtain such opposing effects of

functional connectivity in FPN and DMN is reminiscent of seminal findings supporting the

existence of anti-correlated brain networks in adults (Fox et al., 2005) and may suggest that

similar organizational principles are at play in newborn infants. However, it should be

emphasized that, functional connectivity in FPN and DMN in the current study was not

anticorrelated as such, but rather had opposing effects on infants' behavioral and attentional

regulation.

Assuming that depression represents variability at more extreme ends of a spectrum in

FPN and DMN function, our results concerning behavioral and attentional regulation and their

functional connectivity correlates in infants, are principally in line with prior research showing

hyperconnectivity in the DMN and hypoconnectivity in the FPN in adults with depression

(Kaiser, Andrews-Hanna, Wager, et al., 2015). Moreover, our data show that infants' functional

connectivity in the HIN was associated with negative emotionality. Contrary to prior work with

adults indicating that hypoconnectivity is associated with depression (Patashov et al., 2019;

Wang et al., 2013), the current infant data show that greater connectivity between homologous

brain regions in both hemispheres was associated with greater negative affect. It is unclear why

the direction of the association (positive versus negative) would differ as a function of age

(newborn infants in the current study and adults in previous work), but it is worth noting that the

experience and display of negative affect only gradually emerges during infancy and may thus

not be fully present in newborn infants (Stifter & Fox, 1990). The current approach of measuring

functional connectivity in brain networks during early development previously shown to be

associated with depression therefore affords the possibility in designing protocols to assess risk

for developing depression. Future research employing prospective longitudinal designs is needed

to systematically investigate these brain-behavior correlations with implications for our

understanding and treatment of depression.

482   Taken together, the current findings demonstrate specific associations between functional

brain network connectivity and behavioral temperament in newborn infants. This suggests a

remarkably early emergence of functional networks with behavioral relevance and highlights the

importance of evaluating individual differences reflected in intrinsic brain connectivity.

Although there are many advantages in the current approach of using fNIRS to examine

functional brain connectivity, including its cost-effective and non-confining application, there are

some limitations that need to be mentioned. First, because fNIRS is limited in monitoring

activity from (superficial) cortical structures (Lloyd-Fox et al., 2010), our approach did not allow

us to measure activity from deeper cortical and subcortical regions and include those in our

network analyses. Second, from a developmental perspective, it should be noted that our analysis

is limited to only one age group and comprised of very young infants. It is thus critically

important to further assess the development of variability in these brain networks and their

associations with behavioral temperament over developmental time to determine its long-term

effects and the robustness of these associations (Imai et al., 2014).

496   In conclusion, the current study provides novel insights into the use of fNIRS in

identifying neural endophenotypes -variability in functional brain network connectivity- linked

to behavioral temperament traits in early human development. The present findings support the

notion that functionally distinct neural networks are implicated in regulatory and emotional

behaviors already in newborn infants, adding a critical developmental component to efforts

directed at mapping how the individual functional connectome links to affective, cognitive and

behavioral traits. The current findings shed light on the brain origins of individual differences in

early-emerging behavioral traits and provide the basis for future research examining the genetic

and environmental factors contributing to and the long-term developmental consequences of this

brain-behavior correlation. More generally, the current study provides early ontogenetic evidence

for the idea that studying functional brain network connectivity is an effective way in helping

bridge the gap between brain and behavior.

Acknowledgements

We are grateful to all families who participated in this study as well as Sarah Thomas, Christina
Marlow, Kate Haynes, Carolynn McElroy, Julia Larsen, Heath Yancey, Sujal Sigdel, and
Shefalika Prasad for assistance with infant data collection at the University of Virginia. This
research was supported by *Danone North America, Gut Microbiome, Yogurt and Probiotics*
Fellowship Grant, Jefferson Scholars Foundation and *UVA Data Science Fellowship* (to C.M.K*)*
and National Science Foundation #2017229 and *UVA Brain Institute Seed fund* (to T.G*).*

Conflicts of interest

The authors have no conflicts of interest to declare.

Preprint

This   manuscript   is   available   as   a   pre-print   on   BioRxiv   (link:

https://www.biorxiv.org/content/10.1101/2020.07.15.204271v1)

Data Availability

Data are available upon request to the authors.

References

Altvater-Mackensen, N., & Grossmann, T. (2016). The role of left inferior frontal cortex during audiovisual speech perception in infants. *Neuroimage, 133*, 14-20. doi:10.1016/j.neuroimage.2016.02.061

Asendorpf, J. B., Denissen, J. J., & van Aken, M. A. (2008). Inhibited and aggressive preschool children at 23 years of age: personality and social transitions into adulthood. *Developmental psychology, 44*(4), 997-1011. doi:10.1037/0012-1649.44.4.997

Banich, M. T., & Karol, D. L. (1992). The sum of the parts does not equal the whole: Evidence from bihemispheric processing. *Journal of Experimental Psychology: Human Perception and Performance, 18*(3), 763. doi:10.1037/0096-1523.18.3.763

Bulgarelli, C., Blasi, A., de Klerk, C. C., Richards, J. E., Hamilton, A., & Southgate, V. (2019). Fronto-temporoparietal connectivity and self-awareness in 18-month-olds: A resting state fNIRS study. *Developmental cognitive neuroscience*, 100676. doi:10.1016/j.dcn.2019.100676

Bulgarelli, C., de Klerk, C., Richards, J. E., Southgate, V., Hamilton, A., & Blasi, A. (2020). The developmental trajectory of fronto-temporoparietal connectivity as a proxy of the default mode network: a longitudinal fNIRS investigation. *Hum Brain Mapp, 41*(10), 2717-2740. doi:10.1002/hbm.24974

Caspi, A., Harrington, H., Milne, B., Amell, J. W., Theodore, R. F., & Moffitt, T. E. (2003). Children's behavioral styles at age 3 are linked to their adult personality traits at age 26. *J Pers, 71*(4), 495-513. doi:10.1111/1467-6494.7104001

Caspi, A., Moffitt, T. E., Newman, D. L., & Silva, P. A. (1996). Behavioral observations at age 3

years predict adult psychiatric disorders. Longitudinal evidence from a birth cohort. *Arch*

Gen Psychiatry, 53(11), 1033-1039. doi:10.1001/archpsyc.1996.01830110071009

Cox, J. L., Holden, J. M., & Sagovsky, R. (1987). Detection of postnatal depression.

Development of the 10-item Edinburgh Postnatal Depression Scale. *The British journal*

of psychiatry, 150(6), 782-786.

Cristia, A., Dupoux, E., Hakuno, Y., Lloyd-Fox, S., Schuetze, M., Kivits, J., . . . Minagawa-

Kawai, Y. (2013). An online database of infant functional near infrared spectroscopy

studies: a community-augmented systematic review. *PLoS One, 8*(3), e58906.

doi:10.1371/journal.pone.0058906

Damoiseaux, J. S., Rombouts, S. A., Barkhof, F., Scheltens, P., Stam, C. J., Smith, S. M., &

Beckmann, C. F. (2006). Consistent resting-state networks across healthy subjects.

Proceedings of the National Academy of Sciences of the United States of America,

103(37), 13848-13853. doi:10.1073/pnas.0601417103

Elwell, C., Springett, R., Hillman, E., & Delpy, D. T. (1999). Oscillations in cerebral

haemodynamics. In *Oxygen transport to tissue XXI* (pp. 57-65): Springer.

Eyre, M., Fitzgibbon, S. P., Ciarrusta, J., Cordero-Grande, L., Price, A. N., Poppe, T., . . .

Edwards, A. D. (2020). The Developing Human Connectome Project: typical and

disrupted perinatal functional connectivity. *bioRxiv*, 2020.2001.2020.912881.

doi:10.1101/2020.01.20.912881

Finn, E. S., Shen, X., Scheinost, D., Rosenberg, M. D., Huang, J., Chun, M. M., . . . Constable,

R. T. (2015). Functional connectome fingerprinting: identifying individuals using

patterns of brain connectivity. *Nat Neurosci, 18*(11), 1664-1671. doi:10.1038/nn.4135

Fonov, V., Evans, A. C., Botteron, K., Almli, C. R., McKinstry, R. C., & Collins, D. L. (2011).

Unbiased average age-appropriate atlases for pediatric studies. *Neuroimage, 54*(1), 313-

327. doi:10.1016/j.neuroimage.2010.07.033

Fox, M. D., Snyder, A. Z., Vincent, J. L., Corbetta, M., Van Essen, D. C., & Raichle, M. E.

(2005). The human brain is intrinsically organized into dynamic, anticorrelated functional

networks. *Proceedings of the National Academy of Sciences of the United States of*

America, 102(27), 9673-9678. doi:10.1073/pnas.0504136102

Fox, M. D., Zhang, D., Snyder, A. Z., & Raichle, M. E. (2009). The global signal and observed

anticorrelated resting state brain networks. *Journal of neurophysiology, 101*(6), 3270-

3283. doi:10.1152/jn.90777.2008

Fransson, P., Skiöld, B., Engström, M., Hallberg, B., Mosskin, M., Aden, U., . . . Blennow, M.

(2009). Spontaneous brain activity in the newborn brain during natural sleep--an fMRI

study in infants born at full term. *Pediatr Res, 66*(3), 301-305.

doi:10.1203/PDR.0b013e3181b1bd84

Friston, K. J. (2011). Functional and effective connectivity: a review. *Brain Connect, 1*(1), 13-

36. doi:10.1089/brain.2011.0008

Gao, W., Alcauter, S., Smith, J. K., Gilmore, J. H., & Lin, W. (2015). Development of human

brain cortical network architecture during infancy. *Brain Struct Funct, 220*(2), 1173-

1186. doi:10.1007/s00429-014-0710-3

Gao, W., Lin, W., Grewen, K., & Gilmore, J. H. (2017). Functional Connectivity of the Infant

Human Brain: Plastic and Modifiable. *Neuroscientist, 23*(2), 169-184.

doi:10.1177/1073858416635986

Gao, W., Zhu, H., Giovanello, K. S., Smith, J. K., Shen, D., Gilmore, J. H., & Lin, W. (2009).

Evidence on the emergence of the brain's default network from 2-week-old to 2-year-old

healthy pediatric subjects. *Proceedings of the National Academy of Sciences of the*

United States of America, 106(16), 6790-6795. doi:10.1073/pnas.0811221106

Gartstein, M. A., & Rothbart, M. K. (2003). Studying infant temperament via the revised infant

behavior questionnaire. *Infant Behavior and Development, 26*(1), 64-86.

doi:10.1016/S0163-6383(02)00169-8

Gilmore, J. H., Knickmeyer, R. C., & Gao, W. (2018). Imaging structural and functional brain

development in early childhood. *Nature reviews neuroscience, 19*(3), 123.

Graham, A. M., Buss, C., Rasmussen, J. M., Rudolph, M. D., Demeter, D. V., Gilmore, J. H., . . .

Fair, D. A. (2016). Implications of newborn amygdala connectivity for fear and cognitive

development at 6-months-of-age. *Developmental cognitive neuroscience, 18*, 12-25.

doi:10.1016/j.dcn.2015.09.006

Grossmann, T. (2013a). Mapping prefrontal cortex functions in human infancy. *Infancy, 18*(3),

303-324. doi:10.1111/infa.12016

Grossmann, T. (2013b). The role of medial prefrontal cortex in early social cognition. *Frontiers*

in human neuroscience, 7. doi:10.3389/fnhum.2013.00340

Grossmann, T. (2015). The early development of social brain functions in infancy. *Psychological*

bulletin, 141, 1266-1297.

Grossmann, T., Missana, M., & Krol, K. M. (2018). The neurodevelopmental precursors of

altruistic behavior in infancy. *PLoS Biol, 16*(9), e2005281.

doi:10.1371/journal.pbio.2005281

Hasin, D. S., Sarvet, A. L., Meyers, J. L., Saha, T. D., Ruan, W. J., Stohl, M., & Grant, B. F.

(2018). Epidemiology of Adult DSM-5 Major Depressive Disorder and Its Specifiers in

the   United   States.   *JAMA   Psychiatry,   75*(4),   336-346.

doi:10.1001/jamapsychiatry.2017.4602

Homae, F., Watanabe, H., Otobe, T., Nakano, T., Go, T., Konishi, Y., & Taga, G. (2010).

Development of global cortical networks in early infancy. *J Neurosci, 30*(14), 4877-4882.

doi:10.1523/jneurosci.5618-09.2010

Hu, M. L., Zong, X. F., Mann, J. J., Zheng, J. J., Liao, Y. H., Li, Z. C., . . . Tang, J. S. (2017). A

Review of the Functional and Anatomical Default Mode Network in Schizophrenia.

Neurosci Bull, 33(1), 73-84. doi:10.1007/s12264-016-0090-1

Imai, M., Watanabe, H., Yasui, K., Kimura, Y., Shitara, Y., Tsuchida, S., . . . Taga, G. (2014).

Functional connectivity of the cortex of term and preterm infants and infants with Down's

syndrome. *Neuroimage, 85*, 272-278. doi:10.1016/j.neuroimage.2013.04.080

Insel,   T.   R.   (2010).   Rethinking   schizophrenia.   *Nature,   468*(7321),   187-193.

doi:10.1038/nature09552

Jauniaux, J., Khatibi, A., Rainville, P., & Jackson, P. L. (2019). A meta-analysis of

neuroimaging studies on pain empathy: investigating the role of visual information and

observers'   perspective.   *Soc   Cogn   Affect   Neurosci,   14*(8),   789-813.

doi:10.1093/scan/nsz055

Kabdebon, C., Leroy, F., Simmonet, H., Perrot, M., Dubois, J., & Dehaene-Lambertz, G. (2014).

Anatomical correlations of the international 10–20 sensor placement system in infants.

Neuroimage, 99, 342-356. doi:10.1016/j.neuroimage.2014.05.046

Kaiser, R. H., Andrews-Hanna, J. R., Spielberg, J. M., Warren, S. L., Sutton, B. P., Miller, G. A.,

645   . . . Banich, M. T. (2015). Distracted and down: neural mechanisms of affective

646   interference in subclinical depression. *Soc Cogn Affect Neurosci, 10*(5), 654-663.

647   doi:10.1093/scan/nsu100

648   Kaiser, R. H., Andrews-Hanna, J. R., Wager, T. D., & Pizzagalli, D. A. (2015). Large-Scale

649   Network Dysfunction in Major Depressive Disorder: A Meta-analysis of Resting-State

650   Functional   Connectivity.   *JAMA   Psychiatry,   72*(6),   603-611.

651   doi:10.1001/jamapsychiatry.2015.0071

652   Kelsey, C. M., Krol, K. M., Kret, M. E., & Grossmann, T. (2019). Infants' brain responses to

653   pupillary changes in others are affected by race. *Sci Rep, 9*(1), 4317. doi:10.1038/s41598-

654   019-40661-z

655   Krol, K. M., Puglia, M. H., Morris, J. P., Connelly, J. J., & Grossmann, T. (2019). Epigenetic

656   modification of the oxytocin receptor gene is associated with emotion processing in the

657   infant brain. *Dev Cogn Neurosci, 37*, 100648. doi:10.1016/j.dcn.2019.100648

658   Lloyd-Fox, S., Blasi, A., & Elwell, C. (2010). Illuminating the developing brain: the past, present

659   and future of functional near infrared spectroscopy. *Neuroscience & Biobehavioral

660   Reviews, 34*(3), 269-284. doi:10.1016/j.neubiorev.2009.07.008

661   Lloyd-Fox, S., Széplaki-Köllőd, B., Yin, J., & Csibra, G. (2015). Are you talking to me? Neural

662   activations in 6-month-old infants in response to being addressed during natural

663   interactions. *Cortex, 70*, 35-48. doi:10.1016/j.cortex.2015.02.005

664   Lordier, L., Meskaldji, D.-E., Grouiller, F., Pittet, M. P., Vollenweider, A., Vasung, L., . . .

665   Hüppi, P. S. (2019). Music in premature infants enhances high-level cognitive brain

networks. *Proceedings of the National Academy of Sciences of the United States of America, 116*(24), 12103-12108. doi:10.1073/pnas.1817536116

Lu, C., Zhang, Y.-J., Biswal, B., Zang, Y.-F., Peng, D., & Zhu, C.-Z. (2009). Use of fNIRS to assess resting state functional connectivity. *Journal of neuroscience methods, 186*, 242-249. doi:10.1016/j.jneumeth.2009.11.010

Murphy, K., Birn, R. M., Handwerker, D. A., Jones, T. B., & Bandettini, P. A. (2009). The impact of global signal regression on resting state correlations: are anti-correlated networks introduced? *Neuroimage, 44*(3), 893-905. doi:10.1016/j.neuroimage.2008.09.036

Obrig, H., Neufang, M., Wenzel, R., Kohl, M., Steinbrink, J., Einhäupl, K., & Villringer, A. (2000). Spontaneous low frequency oscillations of cerebral hemodynamics and metabolism in human adults. *Neuroimage, 12*(6), 623-639. doi:10.1006/nimg.2000.0657

Patashov, D., Goldstein, D., & Balberg, M. (2019). *Homologous Connectivity Maps Can Discriminate Diseased from Healthy Brains.* Paper presented at the Optics and the Brain.

Picó-Pérez, M., Alemany-Navarro, M., Dunsmoor, J. E., Radua, J., Albajes-Eizagirre, A., Vervliet, B., . . . Fullana, M. A. (2019). Common and distinct neural correlates of fear extinction and cognitive reappraisal: A meta-analysis of fMRI studies. *Neurosci Biobehav Rev, 104*, 102-115. doi:10.1016/j.neubiorev.2019.06.029

Powell, L. J., Kosakowski, H. L., & Saxe, R. (2018). Social Origins of Cortical Face Areas. *Trends Cogn Sci, 22*(9), 752-763. doi:10.1016/j.tics.2018.06.009

Rigato, S., Stets, M., Bonneville-Roussy, A., & Holmboe, K. (2018). Impact of maternal depressive symptoms on the development of infant temperament: Cascading effects during the first year of life.

Rogers, C. E., Lean, R. E., Wheelock, M. D., & Smyser, C. D. (2018). Aberrant structural and

functional connectivity and neurodevelopmental impairment in preterm children. *J*

Neurodev Disord, 10(1), 38. doi:10.1186/s11689-018-9253-x

Rothbart, M. K. (2007). Temperament, development, and personality. *Current directions in*

psychological science, 16(4), 207-212. doi:10.1111/j.1467-8721.2007.00505.x

Sasai, S., Homae, F., Watanabe, H., & Taga, G. (2011). Frequency-specific functional

connectivity in the brain during resting state revealed by NIRS. *Neuroimage, 56*(1), 252-

257. doi:10.1016/j.neuroimage.2010.12.075

Schöpf, V., Kasprian, G., Brugger, P. C., & Prayer, D. (2012). Watching the fetal brain at 'rest'.

Int J Dev Neurosci, 30(1), 11-17. doi:10.1016/j.ijdevneu.2011.10.006

Smith, S. M., Beckmann, C. F., Andersson, J., Auerbach, E. J., Bijsterbosch, J., Douaud, G., . . .

Harms, M. P. (2013). Resting-state fMRI in the human connectome project. *Neuroimage,*

80, 144-168.

Smyser, C. D., Wheelock, M. D., Limbrick, D. D., Jr., & Neil, J. J. (2019). Neonatal brain injury

and   aberrant   connectivity.   *Neuroimage,*   *185*,   609-623.

doi:10.1016/j.neuroimage.2018.07.057

Steward, T., Menchon, J. M., Jiménez-Murcia, S., Soriano-Mas, C., & Fernandez-Aranda, F.

(2018). Neural Network Alterations Across Eating Disorders: A Narrative Review of

fMRI   Studies.   *Curr*   *Neuropharmacol,*   *16*(8),   1150-1163.

doi:10.2174/1570159x15666171017111532

Stifter, C. A., & Fox, N. A. (1990). Infant reactivity: Physiological correlates of newborn and 5-

month temperament. *Developmental psychology, 26*(4), 582. doi:10.1037/0012-

1649.26.4.582

Tang, A., Crawford, H., Morales, S., Degnan, K. A., Pine, D. S., & Fox, N. A. (2020). Infant

behavioral inhibition predicts personality and social outcomes three decades later.

Proceedings of the National Academy of Sciences of the United States of America,

117(18), 9800-9807. doi:10.1073/pnas.1917376117

Thomas, E., Buss, C., Rasmussen, J. M., Entringer, S., Ramirez, J. S., Marr, M., . . . Wadhwa, P.

D. (2019). Newborn amygdala connectivity and early emerging fear. *Developmental*

cognitive neuroscience, 37, 100604. doi:10.1016/j.dcn.2018.12.002

Thomason, M. E., Dassanayake, M. T., Shen, S., Katkuri, Y., Alexis, M., Anderson, A. L., . . .

Romero, R. (2013). Cross-hemispheric functional connectivity in the human fetal brain.

Sci Transl Med, 5(173), 173ra124. doi:10.1126/scitranslmed.3004978

Thomason, M. E., Hect, J. L., Rauh, V. A., Trentacosta, C., Wheelock, M. D., Eggebrecht, A. T.,

. . . Burt, S. A. (2019). Prenatal lead exposure impacts cross-hemispheric and long-range

connectivity in the human fetal brain. *Neuroimage, 191*, 186-192.

doi:10.1016/j.neuroimage.2019.02.017

van den Heuvel, M. I., & Thomason, M. E. (2016). Functional Connectivity of the Human Brain

in Utero. *Trends Cogn Sci, 20*(12), 931-939. doi:10.1016/j.tics.2016.10.001

Vanderwal, T., Kelly, C., Eilbott, J., Mayes, L. C., & Castellanos, F. X. (2015). Inscapes: A

movie paradigm to improve compliance in functional magnetic resonance imaging.

Neuroimage, 122, 222-232. doi:10.1016/j.neuroimage.2015.07.069

Villringer, A., & Chance, B. (1997). Non-invasive optical spectroscopy and imaging of human

brain function. *Trends Neurosci, 20*(10), 435-442. doi:10.1016/s0166-2236(97)01132-6

Wang, L., Li, K., Zhang, Q.-E., Zeng, Y.-W., Jin, Z., Dai, W.-J., . . . Yu, X. (2013).

Interhemispheric functional connectivity and its relationships with clinical characteristics

in major depressive disorder: a resting state fMRI study. *PLoS One, 8*(3), e60191.

doi:10.1371/journal.pone.0060191

Worobey, J., & Blajda, V. M. (1989). Temperament ratings at 2 weeks, 2 months, and 1 year:

Differential stablity of activity and emotionality. *Developmental psychology, 25*(2), 257.

doi:10.1037/0012-1649.25.2.257

Xu, J., Liu, X., Zhang, J., Li, Z., Wang, X., Fang, F., & Niu, H. (2015). FC-NIRS: a functional

connectivity analysis tool for near-infrared spectroscopy data. *BioMed research*

international, 2015. doi:10.1155/2015/248724

Zhang, H., Shen, D., & Lin, W. (2019). Resting-state functional MRI studies on infant brains: A

decade of gap-filling efforts. *Neuroimage, 185*, 664-684.

doi:10.1016/j.neuroimage.2018.07.004

755