Interleukin-10 suppression enhances T-cell antitumor immunity

and responses to checkpoint blockade in chronic lymphocytic

leukemia

J.R. Rivas[1,2], Y. Liu[3,4], S.S. Alhakeem[1,2], J.M. Eckenrode[3], F. Marti[5], J.P. Collard[1,2], Y.

Zhang[3,4], K.A. Shaaban[3,4], N. Muthusamy[6], G.C. Hildebrandt[1,2,7], R.A. Fleischman[2,7], L.

Chen[2,8], J.S. Thorson[2,3,4], M. Leggas[2,3,4], S. Bondada[1,2,#]

[1]*Dept. of Microbiology, Immunology and Molecular Genetics, University of Kentucky,*

Lexington, KY 40536

[2]*Markey Cancer Center, University of Kentucky, Lexington, KY 40536*

[3]*Dept. of Pharmaceutical Sciences, University of Kentucky, Lexington, KY 40536*

[4]*Center for Pharmaceutical Research and Innovation, University of Kentucky, Lexington,*

KY 40536

[5]*Dept. of Surgery, University of Kentucky, Lexington, KY 40536*

[6]*Dept. of Internal Medicine and Comprehensive Cancer Center, Ohio State University,*

Columbus, OH 43210

[7]*Division of Hematology and Blood and Marrow Transplant, University of Kentucky,*

Lexington, KY 40536

[8]*Dept. of Biostatistics, University of Kentucky, Lexington, KY 40536*

Running title: Suppressing interleukin-10 improves immune checkpoint blockade

#Correspondence: Subbarao Bondada, University of Kentucky, 800 Rose St., 212

Combs Research Building, Lexington, KY 40536; e-mail: bondada@email.uky.edu

Phone number: 859-323-4705; Fax number: 859-257-9608

Manuscript word count: 3990   **Abstract word count:** 196

Number of Figures: 7   **Number of references:** 60

Keywords: CLL, immune checkpoint blockade, IL-10, antitumor immunity

Abstract:

T-cell dysfunction is a hallmark of B-cell Chronic Lymphocytic Leukemia (CLL), where CLL cells downregulate T-cell responses through regulatory molecules including programmed death ligand-1 (PD-L1) and Interleukin-10 (IL-10). Immune checkpoint blockade (ICB) aims to restore T-cell function by preventing the ligation of inhibitory receptors like PD-1. However, most CLL patients do not respond well to this therapy. Thus, we investigated whether IL-10 suppression could enhance antitumor T-cell activity and responses to ICB. Since CLL IL-10 expression depends on Sp1, we utilized a novel, better tolerated analogue of the Sp1 inhibitor mithramycin ($MTM_{ox}32E$) to suppress CLL IL-10. $MTM_{ox}32E$ treatment inhibited mouse and human CLL IL-10 production and maintained T-cell effector function *in vitro*. In the Eµ-Tcl1 mouse model, treatment reduced plasma IL-10 and CLL burden and increased $CD8^+$ T-cell proliferation, effector and memory cell prevalence, and interferon-γ production. When combined with ICB, suppression of IL-10 improved responses to anti-PD-L1 as shown by a 4.5-fold decrease in CLL cell burden compared to anti-PD-L1 alone. Combination therapy also produced more interferon-$γ^+$, cytotoxic effector $KLRG1^+$, and memory $CD8^+$ T-cells, and fewer exhausted T-cells. Since current therapies for CLL do not target IL-10, this provides a novel strategy to improve immunotherapies.

Introduction:

B-cell Chronic Lymphocytic Leukemia (CLL) is a malignancy of CD5$^+$CD19$^+$ B-lymphocytes that leads to severe immune dysfunction. T-cells play multiple roles in the CLL microenvironment and disease progression, from aiding CLL growth to lacking functionality which impedes immunity. CLL cells decrease T-cell antitumor immunity by expressing regulatory molecules including checkpoint ligands and anti-inflammatory cytokines like interleukin-10 (IL-10)(1-5). Patient T-cells then proliferate less after stimulation, increase inhibitory receptor expression, and experience impaired immune synapse formation(1, 6, 7). These functional defects skew T-cell subset distributions, alter T-cell metabolism, and create pseudo-exhausted T-cells(1, 8). Ultimately this results in prevalent infections and secondary cancers, causing much of CLL patient morbidity and mortality(9). Since CLL is rarely curable, there is a need for new approaches to restore immunity.

After repeated antigen stimulation, T-cells upregulate checkpoint receptors such as programmed death-1 (PD-1) that decrease T-cell receptor sensitivity. Ligation of these receptors decreases CD8$^+$ T-cell antitumor immunity via T-cell exhaustion(1). Antibody-based immune checkpoint blockade (ICB) can increase T-cell activity by preventing this interaction. Pembrolizumab (anti-PD-1) ICB was tested to reestablish T-cell functionality in high risk CLL patients with limited success. Only those with Richter's transformation responded(10), suggesting additional mechanisms downregulate T-cell antitumor immunity. Even adoptively transferred anti-CD19 chimeric antigen receptor (CAR) T-cells upregulate apoptosis and exhaustion genes in most CLL patients, resulting in surprisingly low response rates(11). Hence several studies are exploring combination therapies to improve responses to T-cell immunotherapy in CLL(12-14).

Since IL-10 is a well-known anti-inflammatory cytokine, we reasoned that CLL-derived IL-10 may suppress host antitumor T-cell responses(3, 4). Elevated IL-10 levels

correlate with aggressive disease in CLL and other cancers(15, 16), and CLL patients

with elevated plasma IL-10 have a worse 3-year survival rate than those with low

levels(15). Furthermore, reduced DNA methylation at the IL-10 locus or polymorphisms

that increase IL-10 production are correlated with decreased survival in CLL(17, 18).

Despite the importance of IL-10 in the microenvironment and improved antitumor

responses with IL-10 blockade in other cancer types(19-21), no small molecules

blocking IL-10 are currently available(22, 23). Previously, we showed that IL-10R

deletion improved CD8[+] T-cell-efficacy to control CLL growth in the Eμ-TCL1 mouse CLL

model(4). Eμ-TCL1 mice express the oncogene T-cell leukemia 1 (TCL1) under the μ-

enhancer and the VH promoter leading to an aggressive CLL-like disease in 9 to 11

months(24). Like their human counterparts, murine CLL cells secrete IL-10 in a BCR

dependent manner, which can be interrupted by small molecules that inhibit BCR

signaling(3-5).

We hypothesized that reducing IL-10 would restore host anti-tumor responses

and improve the efficacy of T-cell-based immunotherapy. We suppressed CLL-derived

IL-10 with anti-IL-10 or by inhibiting Sp1, a transcription factor required for CLL IL-10

production(4), with a novel analog of mithramycin (MTM$_{ox}$32E, structure in

supplementary methods). MTM$_{ox}$32E exhibits reduced toxicity and increased potency

over the parent molecule in other cancer models(25, 26). This treatment significantly

decreased CLL burden in the blood, spleen and bone marrow, and when combined with

anti-PD-L1, suppressing IL-10 reduced tumor burden and enhanced T-cell functionality

over ICB alone. Thus, suppressing IL-10 could potentially improve antitumor T-cell

function in human CLL and other immunosuppressive cancers.

Methods:

Human samples

Patient blood and healthy donor leukopaks were obtained with informed consent

in accordance with the regulations of the Institutional Review Board of the University of

Kentucky Research Foundation. Diagnosis of CLL was assigned by board-certified

hematologists (Table S1). Peripheral blood mononuclear cells (PBMCs) were isolated by

Ficoll-Paque density centrifugation, then used fresh for healthy donor cells and thawed

from frozen CLL samples. $CD45^+CD5^+CD19^+$ CLL cell expression of CD38, Zap70 and

CD49d were evaluated by flow cytometry (antibodies in Table S2) on an LSR II flow

cytometer (BD Biosciences, Franklin Lakes, NJ). Mutational status of the expressed

BCR was determined by PCR from PBMC RNA, as previously described(4).

Mice

Eµ-TCL1 (C57BL/6J background), NOD.SCID IL-2Rγ$^{-/-}$ (NSG), and C57BL/6J

mice were bred in house. Experimental groups were randomized by including 8-12

week-old mice of both genders (numbers in figure legends). All animal studies were

approved by the University of Kentucky Institutional Animal Care and Use Committee

(IACUC). Three to four million $CD19^+$ bead purified Eµ-TCL1 cells were adoptively

transferred into NSG mice with or without primed $CD8^+$ T-cells at 32:1 (CLL:T-cell). Cell

sorts were performed with MojoSort Mouse Pan B Isolation Kit II (BioLegend, San Diego,

CA) for CD19+ cells and with Miltenyi's magnetic-activated cell sorting (MACS) T-cell

subset Isolation Kits (Miltenyi Biotec, Bergisch Gladbach, Germany) for CD4+ and CD8+

cells according to the manufacturers' protocols. To prime $CD8^+$ T-cells, ten million Ficoll-

Paque purified Eµ-TCL1 splenocytes (>90% $CD5^+CD19^+$) were adoptively transferred

into C57BL/6J mice and ten days later T-cells were purified from the spleen. Disease

burden was monitored in the blood by $CD45^+CD5^+CD19^+$ frequency on an LSR II flow

cytometer (BD Biosciences) and CD45$^+$ cell count on a MACSQuant VYB Flow

Cytometer (Miltenyi Biotec). Mice were treated with 12mg/kg MTM$_{ox}$32E or vehicle (5%

Kolliphor EL, 1% DMSO) by tail vein injection, 150µg/mouse anti-IL-10 (clone JES5-2A5,

BioXCell, West Lebanon, NH), or 10mg/kg anti-PD-L1 or isotype control by

intraperitoneal injection (clone B7-H1, BioXCell). Investigators were not blinded. When

CD5$^+$CD19$^+$ cells comprised more than 70% of the peripheral blood, mice were

euthanized. For $in vivo$ proliferation, 1mg of BrdU (Sigma Aldrich, St. Louis, MO) in

100µL PBS was injected intraperitoneally three hours before euthanasia.

Cell culture

Eµ-TCL1 splenocytes, human CLL PBMCs (hCLL), and sorted T-cell subsets

from mice and humans were cultured at 1x10^6/mL for 24 hours in 96 well plates.

Concentrations of inhibitors and stimulants are listed in figure legends. Anti-IgM

(Jackson ImmunoResearch, West Grove, PA), anti-CD3, anti-CD28 (BioLegend) and

MTM (Enzo Life Sciences, Farmingdale, NY) are commercially sourced, while

MTM$_{ox}$32E was synthesized in house(26). Secreted cytokines were measured with

ELISA MAX Standard Set Mouse or Human kits according to the manufacturer's protocol

(BioLegend). Cellular viability was determined by resuspending cells in cRPMI with

0.1mM resazurin (Sigma Aldrich) I, incubating for four hours at 37°C, and measuring

fluorescence (ex:560nm, em:590nm) on a SpectraMax M5 plate reader (Molecular

Devices, Sunnyvale, CA)(27).

Proliferation assays

CD8$^+$ cells were sorted as above from primed WT mice, healthy donor leukopaks

or hCLL PBMCs. Mouse T-cells were cultured at 0.75x10^6/mL for three days with soluble

10µg/mL anti-CD3 and/or Ficoll-Paque purified Eµ-TCL1 splenocytes. Human T-cells

were cultured at 1×10^6/mL for five days with plate-bound 10µg/mL anti-CD3 and soluble

1µg/mL anti-CD28. Cells were then pulsed with 1µCi ^{3}H-thymidine (Perkin Elmer,

Waltham, MA) or 40µM BrdU for four hours at 37°C. Counts from ^{3}H-thymidine

incorporation were measured with a TopCount NXT Microplate Scintillation &

Luminescence Counter (Perkin Elmer) and BrdU incorporation was measured on an

LSR II or Symphony PRO flow cytometer (BD Biosciences).

Statistics

A priori power analysis determined the number of mice used in experiments. GraphPad

Prism 8 software was used for statistical tests (GraphPad Software, San Diego, CA), as

listed in each figure legend. One-way ANOVA was used for three or more groups, two-

way ANOVA was used for two or more groups over time points or concentrations, and

two-tailed student's t-test was used between two groups. Variance was comparable

between groups in the same test.

Additional information is in the Supplementary Materials and Methods.

Results:

Blocking CLL-derived IL-10 restores T-cell function

Previously we observed that removing CD8$^+$ T-cell IL-10 signaling improves control of CLL growth in the Eμ-TCL1 adoptive transfer model(4). Considering IL-10 is a potent regulator of T-cell antitumor immunity(23), we hypothesized that CLL-derived IL-10 inhibits CLL-primed CD8$^+$ T-cell responses to antigenic stimuli. We primed C57BL/6J CD8$^+$ T-cells with Eμ-TCL1 splenocytes *in vivo* and tested their responses to re-challenge with CLL splenocytes *ex vivo* with and without anti-IL-10. Anti-CD3 induced T-cell proliferation was not changed by anti-IL-10 antibody in the absence of CLL (Fig. 1A). In CLL-T-cell cocultures (3:1), IL-10 levels were only moderately increased with few CLL present, and anti-IL-10 did not change T-cell proliferation (Fig. 1A, 1B). In contrast, proliferation increased four-fold when IL-10 was blocked in cultures with higher CLL:CD8$^+$ ratios (Fig. 1A). This is not explained by CLL cell activity, as blocking IL-10 does not enhance Eμ-TCL1 proliferation under similar culture conditions(4). In CLL:T-cell co-cultures IFN-γ levels were very low, which were substantially increased by neutralizing IL-10 (Fig. 1C).

In earlier studies we showed CLL IL-10 production depends on Sp1 activity(4). Sp1 binds GC rich regions of DNA to regulate transcription, which can be competitively inhibited by MTM or MTM analogues(28). Inhibiting CLL IL-10 with the MTM analogue MTM$_{ox}$32E also increased T-cell functionality in our CD8$^+$:CLL co-cultures. Proliferation of primed CD8+ T-cells was rescued by MTM$_{ox}$32E treatment in the 1:1 co-cultures (Fig. 1D). MTM$_{ox}$32E suppressed IL-10 production in all conditions, even in 3:1 co-cultures, where low IL-10 levels only had a small effect on T-cell proliferation (Fig. 1D-E). Drug treatment did not affect anti-CD3 driven proliferation of primed CD8$^+$ T-cells in isolation (Fig. 1D). Like co-cultures blocked with anti-IL-10, decreasing IL-10 with MTM$_{ox}$32E (Fig. 1E) also increased IFN-γ secretion (Fig. 1F). Notably, 1μM MTM$_{ox}$32E treatment did not

alter IFN-γ secretion from anti-CD3 stimulated, primed CD8$^+$ T-cells in the absence of

CLL (Fig. 1F). Lastly, sorted human CD8$^+$ T-cells both from healthy donors (Fig. 1G, 1I)

and CLL patients (Fig. 1H, 1J) also show susceptibility to IL-10. When stimulated with

plate-bound anti-CD3 and soluble anti-CD28, exogenous IL-10 treatment decreased

human CD8$^+$ T-cell proliferation (Fig. 1G-H), IFN-γ expression (Fig. 1I-J), and CD69

expression (Fig. S1A-B) in a dose-dependent manner. 1µM MTM$_{ox}$32E treatment did not

dampen healthy donor nor CLL patient human CD8$^+$ T-cell responses to stimuli *ex vivo*

(Fig. S1C-F), suggesting this analogue could suppress CLL IL-10 without decreasing T-

cell functionality.

MTM$_{ox}$32E suppresses IL-10 and preserves T-cell function

Dose response studies showed that MTM$_{ox}$32E decreased Eµ-TCL1 IL-10

production with an IC50 of 70nM (Fig. 2A). In contrast, MTM$_{ox}$32E treatment did not

significantly inhibit cytokine production by isolated murine CD4$^+$ or CD8$^+$ T-cells in 24-

hour cultures (Fig. 2A), or CD8$^+$ T-cell proliferation in 48-hour cultures (Fig. 2B).

MTM$_{ox}$32E decreased hCLL PBMC IL-10 production in 93% of patients tested (Fig. 2C,

Table S3) and suppressed IL-10 production by the human CLL cell line, Mec-1 (Fig.

S1G). Chromatin immunoprecipitation with Sp1 in Mec-1 cells showed MTM$_{ox}$32E

reduced Sp1 occupancy of two GC rich sites in the IL-10 promoter (Fig. 2D) and reduced

IL-10 transcription (Fig. S1H), consistent with Sp1 displacement. Notably, the mRNA

levels of Sp1 target genes involved in cell cycle and survival were less affected by

MTM$_{ox}$32E than MTM in hCLL cells (Fig. S1I), leading to less inhibition of cellular viability

(Fig. S1-J-L). Relative to the parent molecule MTM, MTM$_{ox}$32E is similarly potent at

suppressing Eµ-TCL1 IL-10 (Fig. S2A) but is significantly less inhibitory to murine T-cell

cytokine production (Fig. S2B) and viability (Fig. S2C-F). MTM$_{ox}$32E only mildly affected

the viability of healthy human PBMCs after five days in culture (Fig. S3A) and did not

decrease human CD4$^+$ and CD8$^+$ T-cell cytokine production in mixed cultures after 5

days at the dose (100nM) that blocks IL-10 production (Fig. S3B-C), though there was

some inhibition at higher doses (data not shown). Taken together, these results indicate

MTM$_{ox}$32E reduces CLL-derived IL-10 production while preserving T-cell functionality.

Inhibiting CLL-derived IL-10 production enhances T-cell antitumor immunity

Since anti-IL-10 and MTM$_{ox}$32E similarly suppressed CLL IL-10 and enhanced

primed CD8+ T-cell activity in co-culture, these treatments were tested in the adoptive

transfer model of Eμ-TCL1 in NSG mice(4). Disease burden was monitored by the

appearance of CD5$^+$CD19$^+$ CLL cells in the blood and mice were euthanized when blood

CLL reached 70-80% (Fig. S4A, S5A). Without T-cells, CLL grew to 70-80% of

peripheral blood cells in 15-16 days (Fig. 3A-B). Transferring primed CD8+ T-cells at

1:32 ratio (CD8$^+$:CLL) significantly delayed the development of disease, and suppressing

IL-10 further delayed disease development (Fig. 3A-B). When mice were treated with

anti-IL-10, leukemic burden is reduced in blood (Fig. 3A), spleen and bone marrow (Fig.

S4B-C). Similarly, 12mg/kg MTM$_{ox}$32E slowed disease progression in the blood, with an

earlier reduction in CLL burden (statistically significant decrease at day 14 for MTM$_{ox}$32E

but not anti-IL-10, Fig. 3A-B) and less variability in the spleen and bone marrow (Fig.

S4B-C, S5B-D). Both treatments reduced plasma IL-10 levels (Fig. 3C-D),

demonstrating that MTM$_{ox}$32E can suppress CLL-derived IL-10 *in vivo*, leading to

decreased CLL burden.

Spleens from NSG mice also showed decreased CLL burden when MTM$_{ox}$32E

suppressed IL-10. Groups without T-cells had to be euthanized earlier due to faster CLL

progression (day 17 vs 22), so comparisons are made within T-cell and no T-cell groups.

MTM$_{ox}$32E treatment reduced CLL cell frequency in the spleen and bone marrow (Fig.

S5B-D). However, splenic CLL burden (count) was only reduced in NSG mice given

primed CD8$^+$ T-cells and MTM$_{ox}$32E treatment (Fig. 3E). The number of IL-10+ splenic

CLL cells (Fig. 3F, Fig. S5E) and proliferative CLL cells (Fig. 3G) were also only reduced

by IL-10 suppression when T-cells were present, indicating a requirement for T-cells.

The lack of significance without T-cells may be because IL-10 does not affect CLL

directly but inhibits the anti-CLL T-cell response.

In vivo T-cell activation increases with IL-10 suppression

When IL-10 was suppressed with MTM$_{ox}$32E, CD8$^+$ T-cell functionality improved

in Eμ-TCL1 recipient NSG mice. CD8$^+$ T-cells persisted throughout the experiment, with

an increased count in the blood and a trend towards increase in the spleen after

MTM$_{ox}$32E treatment (Fig. 4A, Fig. S5F-G). Like anti-IL-10 (Fig. S4D-E), MTM$_{ox}$32E

treatment increased CD8$^+$ T-cell proliferation, shown by increased number and

frequency of splenic BrdU$^+$CD5$^+$CD19$^-$ T-cells (Fig. 4B). On average, the number and

frequency of IFN-γ$^+$CD8$^+$ (Fig. 4C) and Granzyme-B$^+$CD8$^+$ (GzB$^+$CD8$^+$) splenic T-cells

(Fig. 4D) increased with MTM$_{ox}$32E treatment. Some mice upregulated cytokine

production more than others, but mice with more proliferative T-cells also had more

cytokine competent T-cells. With IL-10 suppression the frequency of CD8$^+$ cells

expressing the effector T-cell marker, killer cell lectin-like receptor G1 (KLRG1), and the

memory marker CD27 (Fig. 4E) increased. KLRG1+ effector T-cells are more cytotoxic

and enhance protective immunity, and may be more active in control of CLL[29]. Despite

these changes, T-cells increased PD-1 expression with time, a marker of activation and

eventual exhaustion, regardless of drug treatment (Fig. 4F). Therefore, we hypothesized

anti-CLL immunity could be improved by combining IL-10 suppression with ICB.

1 **Inhibiting CLL-derived IL-10 production improves responses to anti-PD-L1**

2 **checkpoint blockade**

3   Previous reports described ICB reducing CLL burden in Eµ-TCL1 mice,

4 especially in combination with other therapies(13, 30). Our results were similar, where

5 anti-PD-L1 ICB alone modestly reduced CLL prevalence in blood and bone marrow but

6 not spleen (Fig. S6A-D). Although there were more proliferating CD8$^+$ T-cells in spleen

7 (Fig S6E), ICB did not alter T-cell abundance, and yielded fewer IFN-γ$^+$ and more PD-

8 1$^+$CD8$^+$ T-cells (Fig. S6F-G). We tested if IL-10 suppression could enhance the efficacy

9 of anti-PD-L1 ICB in NSG mice given purified CD19$^+$ Eµ-TCL1 CLL cells and primed

10 CD8$^+$ T-cells.

11   With anti-PD-L1 ICB and MTM$_{ox}$32E-mediated IL-10 suppression combined (Fig.

12 S7A), antitumor CD8$^+$ T-cells were more effective at controlling Eµ-TCL1 CLL growth.

13 Combination therapy slowed CLL growth in the blood compared to anti-PD-L1 alone or

14 isotype plus vehicle treated control mice (Fig. 5A). By the end of the experiment, CLL

15 frequency (Fig. 5B) and burden (Fig. 5C) in the spleen were on average 4.5-fold lower in

16 combination therapy mice versus anti-PD-L1 alone (68.9% vs 15.4%; 94.9x10^6 vs

17 21.1x10^6 cells). In the spleen, the frequency and number of CD8$^+$ T-cells more than

18 doubled (combination vs ICB: 6.97% vs 15.4%, 9.81x10^6 vs 22.14x10^6 cells, Fig. 5D).

19 Blood CD8$^+$ T-cell levels were also elevated with combination therapy (Fig. 5E, Fig.

20 S7B). Spleen size was not different between the therapy groups, which could be due to

21 the expansion of T-cells with combination therapy (Fig. S7C-D). Combination therapy

22 reduced CLL spread to the bone marrow (combination vs ICB: 1.58% vs 0.28%, 1.6x10^5

23 vs 0.2x10^5, Fig. 5F), and dramatically altered the severity of disease, as only one mouse

24 receiving combination therapy developed more than 30% CLL in the blood by the end of

25 the experiment (Fig. 5G).

Adding IL-10 suppression to ICB altered the proliferative capacity of both CD8+

T-cells and CLL cells in NSG mice. As expected, PD-L1$^+$ CLL cells were decreased with

ICB treatment (Fig. S7E). Plasma IL-10 levels were lowest in mice receiving combination

therapy, with intermediate IL-10 levels for anti-PD-L1 alone (Fig. S7F). The number of

BrdU$^+$CD8$^+$ T-cells were elevated in both groups (Fig. 6A), but the number of dividing

CLL cells only decreased with combination therapy (Fig. 6B). This resulted in a greatly

elevated ratio of total and proliferative CD8$^+$ to CLL cells with combination therapy (Fig.

S7G). Interestingly, plasma IL-10 levels inversely correlated with the number of total and

dividing CD8$^+$ T-cells, suggesting IL-10 plays a role in decreasing T-cell persistence and

proliferation (Fig. S7H).

When combining IL-10 suppression and ICB therapy, the outcome of T-cell

activation was further altered. MTM$_{ox}$32E treatment increased the proportion of highly

cytotoxic effector T-cells (KLRG1$^+$) and memory T-cells (CD27$^+$) over ICB alone (Fig.

6C). There were also higher numbers of splenic IFN-γ$^+$CD8$^+$ and GzB$^+$CD8$^+$ T-cells (Fig.

6D-E), and CD8$^+$ T-cells expressing CD107a, a marker of degranulation, in treated mice

(Fig. 6F). Still, similar levels of PD-1$^+$CD8$^+$ T-cells were detected in the blood throughout

the experiment (Fig. S8A). PD-1 is upregulated after T-cell activation, and truly

exhausted T-cells often express other exhaustion markers in addition to PD-1 (Fig. S8B).

Though there was a higher frequency of splenic PD-1$^+$CD8$^+$ T-cells in combination

therapy mice (Fig. S8C), very few CD8$^+$ T-cells expressed other markers of exhaustion,

such as CTLA4, Lag3 or Tim-3 (Fig. S8D). The total frequency of exhausted cells (PD-

1$^+$CTLA4$^+$, PD-1$^+$Lag3$^+$, or PD-1$^+$Tim-3$^+$) decreased for both treatment groups (Fig.

S8E), suggesting that most of the PD-1+ T-cells were not yet exhausted.

The increase in T-cell functionality with IL-10 suppression and ICB combination

therapy shows clearly enhanced antitumor immunity, so lastly, we compared CLL burden

within the previous experiment. With this comparison, the additive effect of IL-10

suppression and ICB became clear. CLL frequency is markedly lower in the spleen (Fig.

6G) and blood (Fig. 6H) with the combination therapy compared to either control or

monotherapy. Furthermore, when mice are treated with this combination, median

survival increases significantly from 41 to 49.5 days (Fig. S8F). Taken together, these

data show improved CD8[+] T-cell antitumor immunity in CLL with combined ICB and IL-

10 suppression.

Discussion:

IL-10 is a potent anti-inflammatory cytokine that downregulates CD8$^+$ antitumor immunity(4). In our studies the therapeutic suppression of IL-10 enhanced CD8$^+$ T-cell control of CLL. IL-10 suppression was most effective at enhancing anti-CLL CD8$^+$ T-cell activity when combined with anti-PD-L1 ICB. This combination reduced disease burden compared to ICB alone and increased the number of CD8$^+$ T-cells, suggesting improved persistence. The outcome of T-cell activation changed as the numbers of CD8$^+$ T-cells expressing functional markers like IFN-γ, GzB, CD27, and KLRG1 increased and exhausted CD8$^+$ T-cells decreased. These data show IL-10 suppression can improve CD8$^+$ T-cell functionality, antitumor immunity, and responses to ICB. They point to IL-10 suppression as a novel therapeutic strategy that may improve responses to T-cell-based immunotherapy in human CLL.

In CLL and several solid tumors, IL-10 reduced the inflammatory state of immune responses including CD8$^+$ T-cells(23, 31-33), although there are examples where IL-10 supported antitumor T-cell function(34-36). The sources of IL-10 are diverse ranging from T-reg cells, macrophages, plasma cells, B-reg cells and cancer cells, as in the present case. Indeed, the role of IL-10 in cancer may be context dependent. In liver and prostate cancers, IL-10 inhibits cytotoxic T-cell function and anti-cancer immunity(37, 38). PEGylated IL-10 treatment increased STAT1 and STAT3 activation in CD8$^+$ T-cells, enhancing control of PDV6 squamous carcinoma tumors(35). The dose and duration of IL-10 exposure is critical to its regulatory effects on CD8$^+$ T-cell maturation into functional memory cells(39). Our data suggest that IL-10 suppression in aggressive CLL enhances memory and effector cell development from CLL-primed CD8$^+$ T-cells, which may differ from T-cells that are mostly tumor-antigen naïve or exhausted. The IL-10 pathway also plays a role in T-cell exhaustion during chronic viral infection, and can directly downregulate CD8$^+$ T-cell function(40, 41). Blocking IL-10 and PD-1 during

chronic viral infection synergistically enhanced viral control and reversed CD8[+] T-cell

exhaustion(42). Furthermore, we previously showed IL-10 inhibits macrophage pro-

inflammatory cytokine secretion in the aged, which may be important as CLL mainly

affects older individuals(43). CD138[+]PD-L1[+] plasma cell derived IL-10 also affects

immune responses to infectious agents in part by inhibiting innate immune cells(44).

These innate cells and stromal cells are part of the microenvironment which supports

CLL cell growth. We find that coculture of CLL cells with stromal cells dramatically

enhances IL-10 production by CLL cells (Collard *et al*, manuscript in preparation).

We describe $MTM_{ox}32E$, a novel small molecule inhibitor of CLL cell IL-10

production, which does not inhibit effector T-cell function, and enhances anti-CLL CD8[+]

T-cell activity. $MTM_{ox}32E$ was more effective than anti-IL-10 in inducing a more

consistent reduction in CLL growth. A small molecule may have better access than

antibodies to microenvironments where CLL cells interact with other cell types.

$MTM_{ox}32E$ treatment reduced Sp1 occupancy of the IL-10 promoter and CLL IL-10

secretion but is uniquely less toxic than MTM. Like MTM, $MTM_{ox}32E$ interacts with GC

rich regions of DNA. $MTM_{ox}32E$ had better pharmacokinetics, tolerability, and selectivity

than MTM and other analogues(26). $MTM_{ox}32E$ showed some selectivity for inhibiting

Sp1 binding on the IL-10 promoter in cell culture. This may be due to interactions

between the analogue and additional proteins in the transcription complex promoting

CLL IL-10 production, which requires further investigation. $MTM_{ox}32E$ also has a small

inhibitory effect on CLL growth *in vivo* but less so when used as single agent *in vitro*.

This could be due to its effect on the microenvironment, which requires further

investigation. Sp1 positively regulates murine T-BET (T-box expressed in T-cells), which

controls natural killer and T-cell IFN-γ expression, and MTM treatment reduced both T-

BET and IFN-γ production in natural killer cells(45). Sp1 expression and transcriptional

activity was shown to be upregulated in activated and proliferating CD4[+] T-cells(46, 47),

but not in activated CD8$^+$ T-cell subsets(48, 49). Our studies focused on CD8$^+$ T-cells,

where we see no detrimental effect of MTM$_{ox}$32E treatment. Additional studies are

required to elucidate the differential effects of MTM and MTM$_{ox}$32E on naïve, effector

and memory T-cells.

Expression of IL-10 by CLL cells is upregulated by B-cell receptor (BCR)

signaling and may be reduced by inhibitors of this pathway. Many therapeutics for CLL

inhibit kinases in BCR signaling(50, 51), and have yielded great clinical outcomes for

CLL patients. The ability of Bruton's tyrosine kinase (Btk) inhibitors (i.e. ibrutinib) and

phosphoinoside-3 kinase (PI3K) inhibitors (i.e. idelalisib) to inhibit CLL growth are well

studied, and may also enhance immune responses(52). However, some patients can

develop resistance mutations or have worse disease after treatment cessation(53, 54).

Ibrutinib treatment also affects T-cells, enhancing their function in CLL patients, which

may be explained by reduced CLL-derived IL-10 in addition to its direct effects(55). CD8$^+$

T-cells in hCLL exist in a pseudo-exhausted state(8), and T-cell exhaustion may occur

within certain T-cell populations in patients(56). Reversing this exhaustion may be

possible with the right therapeutic strategy(57). In mouse studies blocking the PD-1/PD-

L1 pathway was promising, but was more effective with additional blockade of Lag-3 or

Tim-3(30, 58). Anti-PD-1 ICB primarily affects CD8+ T-cells and can rescue exhausted

T-cells, in contrast with anti-CTLA4(59). Nivolumab (anti-PD-1) is currently in clinical

trials for human CLL in combination with the Btk inhibitor ibrutinib (NCT02420912) since

previous reports suggested this combination would be more effective than

monotherapy(12-14). Our findings suggest combining PD-1/PD-L1 blockade with IL-10

suppression could also improve antitumor T-cell responses (Fig. 7) and may still be

effective for patients who have developed resistance to Btk inhibitors.

With no known cure and increasing CLL incidence(60), there is a need for novel

therapies for CLL. Decreasing IL-10 to improve antitumor immunity provides an

opportunity to overcome CLL-induced immunosuppression and enhance current CLL

immunotherapies. Immune evasion through T-cell suppression is common to many types

of cancer, so this strategy may also be applicable to other tumors(16, 19, 20, 22, 32).

Further studies will reveal whether suppressing IL-10 will be effective in treating human

cancer patients.

Acknowledgements:

We would like to thank the patients and their families for participating in our research

and Dr. Eric Durbin with the Kentucky Cancer Registry. We also thank Dr. Siva

Gandhapudi and Dylan Rivas for their helpful discussion. Jacqueline R. Rivas is a Fellow

of the Leukemia & Lymphoma Society and was partially supported by the NCI Training

Grant T32CA165990.

This project was supported in part by R01CA165469 (SB), R01CA217934 (SB),

R01CA217255 (JST), and R01GM115261 (JST). We also acknowledge the support from

the National Center for Advancing Translational Sciences (UL1TR001998 and

UL1TR000117), the Center of Biomedical Research Excellence (COBRE) in

Pharmaceutical Research and Innovation (CPRI, P20GM130456), the University of

Kentucky College of Pharmacy PharmNMR Center, the University of Kentucky Markey

Cancer Center and Markey Cancer Center's Flow Cytometry and Immune Monitoring

Shared Resource Facility (P30CA177558). The content is solely the responsibility of the

authors and does not necessarily represent the official views of the NIH.

Competing Interests:

J.S.T. is a co-founder of Centrose (Madision, WI, USA). No other authors have

competing financial interests.

References:

1.	Brusa D, Serra S, Coscia M, Rossi D, D'Arena G, Laurenti L, et al. The PD-1/PD-L1 axis contributes to T-cell dysfunction in chronic lymphocytic leukemia. Haematologica. 2013;98(6):953-63.

2.	McClanahan F, Riches JC, Miller S, Day WP, Kotsiou E, Neuberg D, et al. Mechanisms of PD-L1/PD-1-mediated CD8 T-cell dysfunction in the context of aging-related immune defects in the Emicro-TCL1 CLL mouse model. Blood. 2015;126(2):212-21.

3.	DiLillo DJ, Weinberg JB, Yoshizaki A, Horikawa M, Bryant JM, Iwata Y, et al. Chronic lymphocytic leukemia and regulatory B cells share IL-10 competence and immunosuppressive function. Leukemia. 2013;27(1):170-82.

4.	Alhakeem SS, McKenna MK, Oben KZ, Noothi SK, Rivas JR, Hildebrandt GC, et al. Chronic Lymphocytic Leukemia-Derived IL-10 Suppresses Antitumor Immunity. J Immunol. 2018;200(12):4180-9.

5.	Alhakeem SS, Sindhava VJ, McKenna MK, Gachuki BW, Byrd JC, Muthusamy N, et al. Role of B cell receptor signaling in IL-10 production by normal and malignant B-1 cells. Ann N Y Acad Sci. 2015;1362:239-49.

6.	Ramsay AG, Johnson AJ, Lee AM, Gorgun G, Le Dieu R, Blum W, et al. Chronic lymphocytic leukemia T cells show impaired immunological synapse formation that can be reversed with an immunomodulating drug. J Clin Invest. 2008;118(7):2427-37.

7.	Palma M, Gentilcore G, Heimersson K, Mozaffari F, Nasman-Glaser B, Young E, et al. T cells in chronic lymphocytic leukemia display dysregulated expression of immune checkpoints and activation markers. Haematologica. 2017;102(3):562-72.

8.	Riches JC, Davies JK, McClanahan F, Fatah R, Iqbal S, Agrawal S, et al. T cells from CLL patients exhibit features of T-cell exhaustion but retain capacity for cytokine production. Blood. 2013;121(9):1612-21.

9.	Forconi F, Moss P. Perturbation of the normal immune system in patients with CLL. Blood. 2015;126(5):573-81.

10.	Ding W, LaPlant BR, Call TG, Parikh SA, Leis JF, He R, et al. Pembrolizumab in patients with CLL and Richter transformation or with relapsed CLL. Blood. 2017;129(26):3419-27.

11.	Fraietta JA, Lacey SF, Orlando EJ, Pruteanu-Malinici I, Gohil M, Lundh S, et al. Determinants of response and resistance to CD19 chimeric antigen receptor (CAR) T cell therapy of chronic lymphocytic leukemia. Nat Med. 2018;24(5):563-71.

12.	Freeman CL, Gribben JG. Immunotherapy in Chronic Lymphocytic Leukaemia (CLL). Curr Hematol Malig Rep. 2016;11(1):29-36.

13.	Sagiv-Barfi I, Kohrt HE, Czerwinski DK, Ng PP, Chang BY, Levy R. Therapeutic antitumor immunity by checkpoint blockade is enhanced by ibrutinib, an inhibitor of both BTK and ITK. Proc Natl Acad Sci U S A. 2015;112(9):E966-72.

14.	Fraietta JA, Beckwith KA, Patel PR, Ruella M, Zheng Z, Barrett DM, et al. Ibrutinib enhances chimeric antigen receptor T-cell engraftment and efficacy in leukemia. Blood. 2016;127(9):1117-27.

15.	Fayad L, Keating MJ, Reuben JM, O'Brien S, Lee BN, Lerner S, et al. Interleukin-6 and interleukin-10 levels in chronic lymphocytic leukemia: correlation with phenotypic characteristics and outcome. Blood. 2001;97(1):256-63.

16.	Zhao S, Wu D, Wu P, Wang Z, Huang J. Serum IL-10 Predicts Worse Outcome in Cancer Patients: A Meta-Analysis. PLoS One. 2015;10(10):e0139598.

17.	Dai ZM, He AL, Zhang WG, Liu J, Cao XM, Chen YX, et al. Association of the four common polymorphisms in interleukin-10 (rs1800890, rs1800896, rs1800871, and rs1800872) with non-Hodgkin's lymphoma risk: a meta-analysis. Int J Clin Exp Med. 2014;7(12):4720-33.

18.	Drennan S, D'Avola A, Gao Y, Weigel C, Chrysostomou E, Steele AJ, et al. IL-10 production by CLL cells is enhanced in the anergic IGHV mutated subset and associates with reduced DNA methylation of the IL10 locus. Leukemia. 2017;8:1686-94.

19.	Lamichhane P, Karyampudi L, Shreeder B, Krempski J, Bahr D, Daum J, et al. IL10 Release upon PD-1 Blockade Sustains Immunosuppression in Ovarian Cancer. Cancer Res. 2017;77(23):6667-78.

20.	Gordy JT, Luo K, Francica B, Drake C, Markham RB. Anti-IL-10-mediated Enhancement of Antitumor Efficacy of a Dendritic Cell-targeting MIP3alpha-gp100 Vaccine in the B16F10 Mouse Melanoma Model Is Dependent on Type I Interferons. J Immunother. 2018;41(4):181-9.

21.	Wu X, Hsu DK, Wang KH, Huang Y, Mendoza L, Zhou Y, et al. IL-10 is overexpressed in human cutaneous T-cell lymphoma and is required for maximal tumor growth in a mouse model. Leuk Lymphoma. 2019;60(5):1244-52.

22.	Sato T, Terai M, Tamura Y, Alexeev V, Mastrangelo MJ, Selvan SR. Interleukin 10 in the tumor microenvironment: a target for anticancer immunotherapy. Immunol Res. 2011;51(2-3):170-82.

23.	Oft M. IL-10: master switch from tumor-promoting inflammation to antitumor immunity. Cancer Immunol Res. 2014;2(3):194-9.

24.	Bichi R, Shinton SA, Martin ES, Koval A, Calin GA, Cesari R, et al. Human chronic lymphocytic leukemia modeled in mouse by targeted TCL1 expression. Proc Natl Acad Sci U S A. 2002;99(10):6955-60.

25.	Hou C, Weidenbach S, Cano KE, Wang Z, Mitra P, Ivanov DN, et al. Structures of mithramycin analogues bound to DNA and implications for targeting transcription factor FLI1. Nucleic Acids Res. 2016;44(18):8990-9004.

26.	Liu Y, Eckenrode JM, Zhang Y, Zhang J, Hayden RC, Kyomuhangi A, et al. Mithramycin 2'-Oximes with Improved Selectivity, Pharmacokinetics, and Ewing Sarcoma Antitumor Efficacy. J Med Chem. 2020;63(22):14067-86.

27.	Mitra P, Eckenrode JM, Mandal A, Jha AK, Salem SM, Leggas M, et al. Development of Mithramycin Analogues with Increased Selectivity toward ETS Transcription Factor Expressing Cancers. J Med Chem. 2018;61(17):8001-16.

28.	Choi ES, Nam JS, Jung JY, Cho NP, Cho SD. Modulation of specificity protein 1 by mithramycin A as a novel therapeutic strategy for cervical cancer. Sci Rep. 2014;4:7162.

29.	Herndler-Brandstetter D, Ishigame H, Shinnakasu R, Plajer V, Stecher C, Zhao J, et al. KLRG1(+) Effector CD8(+) T Cells Lose KLRG1, Differentiate into All Memory T Cell Lineages, and Convey Enhanced Protective Immunity. Immunity. 2018;48(4):716-29 e8.

30.	Wierz M, Pierson S, Guyonnet L, Viry E, Lequeux A, Oudin A, et al. Dual PD1/LAG3 immune checkpoint blockade limits tumor development in a murine model of chronic lymphocytic leukemia. Blood. 2018;131(14):1617-21.

31.	Ruffell B, Chang-Strachan D, Chan V, Rosenbusch A, Ho CM, Pryer N, et al. Macrophage IL-10 blocks CD8+ T cell-dependent responses to chemotherapy by suppressing IL-12 expression in intratumoral dendritic cells. Cancer Cell. 2014;26(5):623-37.

32.	Mahipal A, Terai M, Berd D, Chervoneva I, Patel K, Mastrangelo MJ, et al. Tumor-derived interleukin-10 as a prognostic factor in stage III patients undergoing adjuvant treatment with an autologous melanoma cell vaccine. Cancer Immunol Immunother. 2011;60(7):1039-45.

33.   Sawant DV, Yano H, Chikina M, Zhang Q, Liao M, Liu C, et al. Adaptive plasticity of IL-10(+) and IL-35(+) Treg cells cooperatively promotes tumor T cell exhaustion. Nat Immunol. 2019;20(6):724-35.

34.   Gu T, De Jesus M, Gallagher HC, Burris TP, Egilmez NK. Oral IL-10 suppresses colon carcinogenesis via elimination of pathogenicCD4(+) T-cells and induction of antitumor CD8(+) T-cell activity. Oncoimmunology. 2017;6(6):e1319027.

35.   Emmerich J, Mumm JB, Chan IH, LaFace D, Truong H, McClanahan T, et al. IL-10 directly activates and expands tumor-resident CD8(+) T cells without de novo infiltration from secondary lymphoid organs. Cancer Res. 2012;72(14):3570-81.

36.   Santin AD, Hermonat PL, Ravaggi A, Bellone S, Pecorelli S, Roman JJ, et al. Interleukin-10 increases Th1 cytokine production and cytotoxic potential in human papillomavirus-specific CD8(+) cytotoxic T lymphocytes. J Virol. 2000;74(10):4729-37.

37.   Shalapour S, Lin XJ, Bastian IN, Brain J, Burt AD, Aksenov AA, et al. Inflammation-induced IgA+ cells dismantle anti-liver cancer immunity. Nature. 2017;551(7680):340-5.

38.   Shalapour S, Font-Burgada J, Di Caro G, Zhong Z, Sanchez-Lopez E, Dhar D, et al. Immunosuppressive plasma cells impede T-cell-dependent immunogenic chemotherapy. Nature. 2015;521(7550):94-8.

39.   Cui W, Liu Y, Weinstein JS, Craft J, Kaech SM. An interleukin-21-interleukin-10-STAT3 pathway is critical for functional maturation of memory CD8+ T cells. Immunity. 2011;35(5):792-805.

40.   Brooks DG, Trifilo MJ, Edelmann KH, Teyton L, McGavern DB, Oldstone MB. Interleukin-10 determines viral clearance or persistence in vivo. Nat Med. 2006;12(11):1301-9.

41.   Smith LK, Boukhaled GM, Condotta SA, Mazouz S, Guthmiller JJ, Vijay R, et al. Interleukin-10 Directly Inhibits CD8(+) T Cell Function by Enhancing N-Glycan Branching to Decrease Antigen Sensitivity. Immunity. 2018;48(2):299-312 e5.

42.   Brooks DG, Ha SJ, Elsaesser H, Sharpe AH, Freeman GJ, Oldstone MB. IL-10 and PD-L1 operate through distinct pathways to suppress T-cell activity during persistent viral infection. Proc Natl Acad Sci U S A. 2008;105(51):20428-33.

43.   Chelvarajan RL, Liu Y, Popa D, Getchell ML, Getchell TV, Stromberg AJ, et al. Molecular basis of age-associated cytokine dysregulation in LPS-stimulated macrophages. J Leukoc Biol. 2006;79(6):1314-27.

44.   Lino AC, Dang VD, Lampropoulou V, Welle A, Joedicke J, Pohar J, et al. LAG-3 Inhibitory Receptor Expression Identifies Immunosuppressive Natural Regulatory Plasma Cells. Immunity. 2018;49(1):120-33 e9.

45.   Yu J, Wei M, Boyd Z, Lehmann EB, Trotta R, Mao H, et al. Transcriptional control of human T-BET expression: the role of Sp1. Eur J Immunol. 2007;37(9):2549-61.

46.   Lacroix I, Lipcey C, Imbert J, Kahn-Perles B. Sp1 transcriptional activity is up-regulated by phosphatase 2A in dividing T lymphocytes. J Biol Chem. 2002;277(11):9598-605.

47.   Juang YT, Rauen T, Wang Y, Ichinose K, Benedyk K, Tenbrock K, et al. Transcriptional activation of the cAMP-responsive modulator promoter in human T cells is regulated by protein phosphatase 2A-mediated dephosphorylation of SP-1 and reflects disease activity in patients with systemic lupus erythematosus. J Biol Chem. 2011;286(3):1795-801.

48.   Moskowitz DM, Zhang DW, Hu B, Le Saux S, Yanes RE, Ye Z, et al. Epigenomics of human CD8 T cell differentiation and aging. Sci Immunol. 2017;2(8).

49.   He B, Xing S, Chen C, Gao P, Teng L, Shan Q, et al. CD8(+) T Cells Utilize Highly Dynamic Enhancer Repertoires and Regulatory Circuitry in Response to Infections. Immunity. 2016;45(6):1341-54.

50.   Roberts AW, Davids MS, Pagel JM, Kahl BS, Puvvada SD, Gerecitano JF, et al. Targeting
BCL2 with Venetoclax in Relapsed Chronic Lymphocytic Leukemia. N Engl J Med.
2016;374(4):311-22.
51.   Jain N, O'Brien S. Targeted therapies for CLL: Practical issues with the changing
treatment paradigm. Blood Rev. 2016;30(3):233-44.
52.   Woyach JA, Johnson AJ, Byrd JC. The B-cell receptor signaling pathway as a therapeutic
target in CLL. Blood. 2012;120(6):1175-84.
53.   Jain P, Thompson PA, Keating M, Estrov Z, Ferrajoli A, Jain N, et al. Long-term outcomes
for patients with chronic lymphocytic leukemia who discontinue ibrutinib. Cancer.
2017;123(12):2268-73.
54.   Ahn IE, Underbayev C, Albitar A, Herman SE, Tian X, Maric I, et al. Clonal evolution
leading to ibrutinib resistance in chronic lymphocytic leukemia. Blood. 2017;129(11):1469-79.
55.   Long M, Beckwith K, Do P, Mundy BL, Gordon A, Lehman AM, et al. Ibrutinib treatment
improves T cell number and function in CLL patients. J Clin Invest. 2017;127(8):3052-64.
56.   Hanna BS, Roessner PM, Yazdanparast H, Colomer D, Campo E, Kugler S, et al. Control of
chronic lymphocytic leukemia development by clonally-expanded CD8(+) T-cells that undergo
functional exhaustion in secondary lymphoid tissues. Leukemia. 2019;33(3):625-37.
57.   Bachireddy P, Hainz U, Rooney M, Pozdnyakova O, Aldridge J, Zhang W, et al. Reversal of
in situ T-cell exhaustion during effective human antileukemia responses to donor lymphocyte
infusion. Blood. 2014;123(9):1412-21.
58.   Ngiow SF, von Scheidt B, Akiba H, Yagita H, Teng MW, Smyth MJ. Anti-TIM3 antibody
promotes T cell IFN-gamma-mediated antitumor immunity and suppresses established tumors.
Cancer Res. 2011;71(10):3540-51.
59.   Wei SC, Anang NAS, Sharma R, Andrews MC, Reuben A, Levine JH, et al. Combination
anti-CTLA-4 plus anti-PD-1 checkpoint blockade utilizes cellular mechanisms partially distinct
from monotherapies. Proc Natl Acad Sci U S A. 2019;116(45):22699-709.
60.   Chen Q, Jain N, Ayer T, Wierda WG, Flowers CR, O'Brien SM, et al. Economic Burden of
Chronic Lymphocytic Leukemia in the Era of Oral Targeted Therapies in the United States. J Clin
Oncol. 2017;35(2):166-74.

Figure Legends:

Figure 1: CLL IL-10 suppresses T-cell responses. (A) Proliferation of CLL-primed CD8$^+$ T-cells in response to anti-CD3 or varying numbers of Eμ-TCL1 splenocytes after 72 hours of stimulation, with or without 10μg/mL anti-IL-10. Secreted IL-10 (B) and IFN-γ (C) from cultures in panel A (Eμ-TCL1 spleen cells alone produced 986pg/mL IL-10 and 80pg/mL IFN-γ). (D) Proliferation of CLL-primed CD8$^+$ T-cells stimulated with anti-CD3 or varying numbers of Eμ-TCL1 splenocytes after 72 hours with or without 1μM MTM$_{ox}$32E. Secreted IL-10 (E) and IFN-γ (F) from cultures in panel D (Eμ-TCL1 spleen cells alone produced 948pg/mL IL-10 and 78pg/mL IFN-γ). (G-H) Normalized proliferation (BrdU incorporation) of healthy donor (G) or CLL patient (H) human CD8$^+$ T-cells in response to a 5-day stimulation with plate bound anti-CD3 with soluble anti-CD28 and recombinant human IL-10. (I-J) Normalized intracellular IFN-γ production from cells in (G-H), measured by flow cytometry. Frequency of responding cells in G-J were normalized to responses of CD8$^+$ T-cells from same donor/patient cultured without IL-10. Two-way ANOVA between control and antibody or inhibitor treatment was used to calculate statistical significance. **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$

Figure 2: MTM$_{ox}$32E suppresses CLL IL-10 without dampening *in vitro* T-cell responses. (A) Secreted IL-10 from Eμ-TCL1 splenocytes, IL-2 from murine C57BL/6 CD4$^+$ T-cells, and IFN-γ from murine C57BL/6 CD8$^+$ T-cells cultured for 24 hours with MTM$_{ox}$32E, normalized to vehicle control (660pg/mL, 398pg/mL, 106pg/mL, respectively). T-cells were stimulated with 10μg/mL soluble anti-CD3. (B) Proliferation (BrdU incorporation) of C57BL/6 CD8$^+$ T-cells cultured for 72 hours with 10μg/mL anti-CD3 $\pm$ MTM$_{ox}$32E (error bars are too small to be seen). (C) Secreted IL-10 from human CLL PBMCs stimulated with 25μg/mL anti-IgM for 24 hours with MTM$_{ox}$32E, normalized to vehicle control (17-656pg/mL). (D) Promoter occupancy of Sp1 GC-rich sites on the

human IL-10 promoter in Mec1 cells treated with 1000nM $MTM_{ox}32E$. Statistical

comparisons made by two-way ANOVA in A and one-way ANOVA in D. *p<0.05,

**p<0.01, ***p<0.001, ****p<0.0001

Figure 3: $MTM_{ox}32E$ enhances anti-CLL activity of CD8$^+$ T-cells and suppresses

CLL IL-10 production *in vivo*. (A-B) Frequency of Eµ-TCL1 CLL cells in the blood of

NSG mice (mean + SEM). Six groups of eight NSG mice were injected with Eµ-TCL1

CLL, with or without Eµ-TCL1 primed CD8$^+$ T-cells (CLL to T-cell ratio of 32:1) and

received 150µg anti-IL-10 or isotype every two days (A) or 12mg/kg $MTM_{ox}32E$ or

vehicle every three days (B). Stars indicate statistical significance of differences between

control and anti-IL-10 (A) or $MTM_{ox}32E$ (B) treated group. (C-D) Plasma IL-10 levels in

NSG mice (mean + SEM). Stars indicate significance between control and anti-IL-10 (C)

or $MTM_{ox}32E$ (D) treated group. (E) Burden of Eµ-TCL1 CLL cells in the spleens of NSG

mice without CD8$^+$ T-cells (left panel, euthanized day 17) and with CD8$^+$ T-cells (right

panel, euthanized day 22) in the $MTM_{ox}32E$ group. (F) Count of IL-10$^+$ Eµ-TCL1 CLL

cells in the spleens of NSG mice without CD8$^+$ T-cells (left, day 17) and with CD8$^+$ T-

cells (right, day 22). (G) Count of proliferative (BrdU$^+$) Eµ-TCL1 CLL cells in the spleens

of NSG mice without CD8$^+$ T-cells (left, day 17) and with CD8$^+$ T-cells (right, day 22).

Statistical comparisons made by two-way ANOVA in A-D, and one-way ANOVA in E-G.

*p<0.05, **p<0.01, ***p<0.001, ****p<0.0001, VC=vehicle control, x=mice euthanized at

indicated earlier timepoint

Figure 4: Inhibiting CLL IL-10 with $MTM_{ox}32E$ increases the numbers of T- effector

cells. Mice were treated as in Figure 3B. (A) The number of CD8$^+$ T-cells in the blood of

NSG mice. (B) (Left) Representative histograms of BrdU incorporation by splenic

CD5$^+$CD19$^-$ T-cells (>99% CD8$^+$). Frequency (middle) and number (right) of BrdU$^+$

splenic CD8$^+$ T-cells. (C) (Left) Representative histograms of IFN-γ production by splenic

CD8$^+$ T-cells. Frequency (middle) and number (right) of splenic IFN-γ$^+$ CD8$^+$ T-cells. (D)

Frequency (left) and number (right) of GzB$^+$ CD8$^+$ T-cells in the spleen. (E) Frequency of

KLRG1$^+$ highly cytotoxic effector and CD27$^+$ memory CD8$^+$ T-cells in the spleen (left)

and pie chart representations of CD8$^+$ T-cell proportions. (F) Frequency of PD-1

expressing CD8$^+$ T-cells in the blood. Statistical comparisons in A+F were obtained by

two-way ANOVA comparing vehicle to MTM$_{ox}$32*E* treated mice, and in B-E by one-way

ANOVA between vehicle to MTM$_{ox}$32*E*. *p<0.05

Figure 5: Anti-PD-L1 checkpoint blockade is more effective when combined with

IL-10 suppression by MTM$_{ox}$32*E*. Three groups of thirteen NSG mice were injected

with Eμ-TCL1 and Eμ-TCL1 primed CD8$^+$ T-cells at a ratio of one T-cell to 32 CLL cells.

Mice received 12mg/kg MTM$_{ox}$32*E* or vehicle every two to three days plus 10mg/kg anti-

PD-L1 or isotype control every three days. (A) Frequency of Eμ-TCL1 CLL cells in the

blood of NSG mice (mean + SEM). (B-C) Frequency (B) and count (C) of Eμ-TCL1 CLL

cells in the spleen. (D) Frequency (left) and count (right) of CD8$^+$ T-cells in the spleen of

recipient NSG mice. (E) Frequency of CD8$^+$ T-cells in the blood of NSG recipients (mean

+ SEM). (F) Frequency (left) and count (right) of Eμ-TCL1 CLL cells in the bone marrow.

(G) Percent of mice with less than 30% CLL in the blood over time. Statistical

significance in A, E, G was calculated by two-way ANOVA comparing vehicle to treated

mice, and by one-way ANOVA comparing vehicle to treated mice in remaining panels.

*p<0.05, **p<0.01, ***p<0.001, ****p<0.0001 indicate statistical significance of

*difference between combination and ICB alone, #difference between combination and

control, &difference between ICB alone and control

Figure 6: T-cells from Eμ-TCL1 mice are more functional when treated with both anti-PD-L1 checkpoint blockade and IL-10 blockade. Mice were treated as in Figure 5. (A) Count of BrdU$^+$ splenic CD8$^+$ T-cells. (B) Count of BrdU$^+$ Eμ-TCL1 CLL cells in the spleen. (C) Frequency of KLRG1$^+$ highly cytotoxic effector and CD27$^+$ memory CD8$^+$ T-cells in the spleen (left) and pie chart representations of CD8$^+$ T-cell proportions. (D-E) Count of IFN-γ$^+$ (D) and GzB$^+$ (E) splenic CD8$^+$ T-cells. (F) Frequency of CD107a$^+$ splenic CD8$^+$ T-cells. (G) Comparison of CLL frequency in spleen from Fig. 5B and Fig. S5B. (H) Comparison of CLL frequency in the blood from Figures 3B and 5A. Statistical comparisons done by one-way ANOVA comparing vehicle to treated mice, except in G+H where all groups were compared to each other. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001

Figure 7: Cooperation of IL-10 suppression and immune checkpoint blockade. Created with BioRender.com. Model shows combining IL-10 suppression with anti-PD-L1 ICB enhances antitumor CD8$^+$ T-cell activity by counteracting multiple methods of CLL immune suppression.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7